# Transcriptional modulation during photomorphogenesis in rice seedlings

Parul Gupta, Pankaj Jaiswal[#]

**Affiliations:**

Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR, USA

**# Correspondence:**

Prof. Pankaj Jaiswal

Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA

Email: Jaiswal.pankaj@oregonstate.edu

**Keywords:** Rice, photomorphogenesis, RNA-Seq, transcriptome, alternative splicing, lncRNA, MVA and MEP pathways, circadian clock

**ABSTRACT**

Light is one of the most important factors regulating plant gene expression patterns, metabolism,

physiology, growth, and development. To explore how light may induce or alter transcript

splicing, we conducted RNA-Seq-based transcriptome analyses by comparing the samples

harvested as etiolated seedlings grown under continuous dark conditions vs. the light-treated

green seedlings. We identified 14,766 differentially expressed genes, of which 4369 genes

showed alternative splicing. We observed that genes mapped to the plastid-localized methyl-

erythritol-phosphate (MEP) pathway were upregulated in light compared to the cytosolic

mevalonate (MVA) pathway genes. Many of these genes also undergo splicing. These pathways

provide crucial metabolite precursors for the biosynthesis of secondary metabolic compounds

needed for chloroplast biogenesis, the establishment of successful photosynthetic apparatus, and

photomorphogenesis. In the chromosome-wide survey of the light-induced transcriptome, we

observed intron retention as the most predominant splicing event. In addition, we identified 1709

novel lncRNA transcripts in our transcriptome data.

## INTRODUCTION

Light is an essential growth factor for sustaining autotrophic plant life. The quality, quantity, direction, and length of light exposure affect plant growth and development [1,2]. The natural light-dark cycle maintains the carbon and nitrogen metabolism of rice plants [3]. Light induces transcriptional reprogramming in various plant species [3–10] via an array of photoreceptors [11,12]. The photoreceptors phytochrome (red light) and cryptochrome (blue light) are involved in light signaling and photomorphogenesis [13,14]. Genes encoding phytochrome interacting factors (PIFs) promote skotomorphogenesis and development in the dark [15–17]. Light-mediated gene expression modulation is also triggered by translational enhancement of preexisting mRNA pool instead of an enhanced rate of transcription [18,19].

Intron splicing in multiexonic mRNA is a post-transcriptional regulatory process that often produces different mRNA isoforms transcribed from a single gene locus, thus contributing to proteome plasticity. Light-induced alternative splicing (AS) was observed in ~10% of the protein-coding genes of *Arabidopsis* and *Physcomitrella patens* [20,21]. A flash of light applied in the middle of the dark or nighttime is sufficient to induce splicing [22]. In contrast to animals, where the most common AS event is exon skipping (ES), intron retention (IR) is the most common AS event in rice [23], Arabidopsis [23,24], and poplar [25]. It is now well established that transcriptome modulation via AS is vital for plant growth, development, and response to stress [26–28].

Compared to the protein-coding genes, long noncoding RNAs (lncRNAs) are transcripts >200 bp in length that do not have protein-coding potential. lncRNAs are classified into categories: (i) sense lncRNAs, (ii) antisense lncRNAs, (iii) intergenic RNAs (lincRNAs), (iv) intronic RNAs, and (v) bidirectional lncRNAs [29]. In addition to the numerous studies coupling gene expression and AS studies on protein-coding genes, it is now known that lncRNAs also play a role in regulating gene expression through transcriptional, post-transcriptional, and chromatin remodeling mechanisms [29–32]. In rice, lncRNAs regulate biological processes, such as ovule development and female gametophyte abortion [33], sexual reproduction [34], and stress response [35,36]. However, the role of light in regulating lncRNAs is not well studied. *Arabidopsis* noncoding RNA *HID1* is a known positive regulator of photomorphogenesis in continuous red light [37].

Rice is a global staple crop and is a model for studying crop genomics. The transition from

skotomorphogenesis under dark conditions to photomorphogenesis under light exposure is

critical for seedling survival and requires precise control of gene expression by different

regulatory mechanisms. Results from our rice study show that exposure to light alters the

expression and splicing of a wide array of protein-coding genes but not so much for the

noncoding lncRNA genes.

## MATERIALS & METHODS

### Plant material, growth conditions, and treatment

Seeds of rice (*Oryza sativa spp. japonica* cv. Nipponbare) were grown and processed for the

experiment by following the growth conditions and sampling described previously [38]. Seeds were

sown in the dark and germinated on day 2. After sowing, these germinated seedlings grew in the

dark for 8 days (8DD). At the end of day 8, three biological replicates of the dark-grown

etiolated shoots were harvested. The remaining dark-grown seedlings were exposed to

continuous white light at 120 µmol/m2/sec (measured at the soil surface) for 48 hours or 2 days

(days 9 and 10 after sowing). The shoots of three biological replicates of light-treated green-

colored seedling samples (8DD-2LL) were harvested at the end of day 10. Harvested samples

were frozen using liquid nitrogen and stored at -80°C until further processing. Throughout this

report, 8DD-treated plants are called dark samples, and 8DD-2LL light-treated plants are called

light samples. Data analysis is described for light regulation.

### Sample preparation and sequencing

Total RNA from frozen tissue samples was extracted using RNA Plant reagent (Invitrogen Inc.,

USA) and RNeasy kits (Qiagen Inc., USA) and treated with RNase-free DNase (Life

Technologies Inc., USA) according to the manufacturer's protocol. The total RNA quality and

concentration were determined using an ND-1000 spectrophotometer (Thermo Fisher Scientific

Inc., USA) and Bioanalyzer 2100 (Agilent Technologies Inc., USA). PolyA-enriched mRNA

libraries were prepared from three biological replicates of dark and light samples using the

TruSeqTM RNA Sample Preparation Kits (v2) and sequenced as 51-bp single-end reads using

the Illumina HiSeq 2000 instrument (Illumina Inc., USA) according to the manufacturer's

protocol at the Center for Genomic Research and Biocomputing, Oregon State University. The

strand-specific sequencing reads and metadata were deposited at EMBL-EBI ArrayExpress

(accession number E-MTAB-5689).

**RNA-seq data analysis**

The generation of FASTQ files from the RNA-Seq sequences was performed by CASAVA

software v1.8.2 (Illumina Inc.). Sequence reads were filtered and trimmed for low quality at a

score of 20 using Sickle v1.33 [39]. Clean, high-quality reads from each sample and replicates were

aligned to the *Oryza sativa japonica* cv Nipponbare reference genome (IRGSP-1.0.31) using

TopHat v2.1.1 [40]. Mapped reads were assembled using Cufflinks, and the reference-guided

assembled transcripts from each replicate were merged using Cuffmerge [41]. Assembled

transcripts were compared to the reference genome annotation using Cuffcompare. The RSEM

software package estimated normalized baseline expression from the aligned sequence reads [42].

For differential gene expression analysis, read count data obtained from RSEM were used in

EBSeq [43]. Differentially expressed (DE) genes were filtered based on the false discovery rate

corrected P value ≤0.05.

**Functional annotation**

105 We carried out the Gene Ontology (GO) enrichment analysis tool provided by the GO

106 consortium [44] to determine the biological roles played by the enriched gene set. Plant pathway

107 enrichment analysis was done by mapping the DE genes using the Plant Reactome analysis tool

108 [46] (http://plantreactome.gramene.org/PathwayBrowser/#TOOL=AT).

**Alternative splicing analysis**

Splicing events in the transcripts from the samples were identified by the SpliceGrapher v0.2.5

pipeline [47]. Sequence reads from each sample, and replicates were aligned to the reference rice

genome. Splice site-specific classifiers were built using build_classifiers.py script using

canonical (GT) and noncanonical (GC) donor sites and acceptor site (AG) for *Oryza sativa*

genome annotation version 31 (Oryza_sativa.IRGSP-1.0.31). Read alignments in SAM format

from each replicate were used as input for SpliceGrapher's sam_filter.py script to filter out false-

positive sites. SpliceGrapher Python scripts were used for the generation of depth files

(sam_to_depths.py), splice graph prediction (predict_graphs.py), generating statistics

(splicegraph_statistics.py) from a set of splice graphs, gene-by-gene summary

(genewise_statistics.py) of splicing events and splice graph visualization (plotter.py). The

Realignment pipeline was used to construct putative transcripts from unresolved exons with

sufficient coverage from the alignments [47]. In the following steps, IsoLasso [48], an extension of

the SpliceGrapher workflow, was used to predict novel splicing events.

**Prediction of long noncoding RNA (lncRNA)**

All transcripts annotated as intergenic transcripts, intron transcripts, antisense exon transcripts

overlapping the reference exons, and antisense intron transcripts overlapping the reference

introns were considered potential lncRNA candidates. Transcript sequences of length <=200

nucleotides were filtered out, and the gffread function of Cufflinks was used to extract fasta

sequences of potential lncRNA transcripts from the gtf file. CPC2 [49] was used to predict the

coding potential of transcripts. Predicted lncRNAs were scanned by InterProScan [50] to ensure the

absence of protein-coding domains. To identify novel lncRNAs, a BLASTn [51] search was

performed against a custom BLAST database generated using rice lncRNAs downloaded from

the CANTATAdb2.0 (http://cantata.amu.edu.pl), PNRD [52], GreeNC [53], and RiceLNCPedia [54]

databases. Secondary structures for the lncRNA molecules were predicted by the RNAfold

software [55,56].

**RESULTS**

**Light-mediated differential gene expression during photomorphogenesis**

To explore the transcriptome modulation in rice in response to light, we sequenced the strand-

specific poly-A enriched RNA fraction isolated from three biological replicates of rice plants

grown under dark and light exposure conditions (see methods). A total of ~38 million and ~42

million high-quality reads were generated from the dark and light-treated samples, respectively.

More than 92% of reads from each sample aligned to the rice reference genome (Supplementary

Table S1). We found 38,642 genes showed baseline normalized expression in the samples, of

which 33,943 genes expressed in the dark vs 35,772 genes that expressed under light,

respectively. Differential expression analysis identified 14,766 light-regulated genes (Figure 1,

Supplementary Figure S1, Supplementary Table S2).

**Figure 1:** Summary of differential gene expression and transcript splicing observed in rice
seedling shoots undergoing photomorphogenesis.

149
150

## Gene function and pathway enrichment analyses

Gene Ontology-based functional annotation analysis of the light-regulated differentially
expressed gene set (Supplementary Figure S2) revealed enrichment for biological processes
(BP), such as chlorophyll and cell wall biosynthesis, besides photosystem light reaction
pathways and ribosome assembly. Dominant molecular functions (MF) were rRNA binding,
cytoskeletal motor activity, oxidoreductase and metallopeptidase activities, translation
elongation, and chaperone binding. As expected, chloroplast, cytoskeleton, ribosomes,
peroxisomes, and nucleus were the major cellular component sites of activity.

Pathway enrichment analysis using the Plant Reactome pathway analysis tool [57] mapped 475
light-upregulated genes to 243 pathways (Supplementary Table S3). A total of 164 pathways
overlapped between light-up and light-downregulated genes. Most pathways showed higher
mappings to light-upregulated genes, except hormone auxin and brassinosteroid signaling and
reproductive structure development (seed). Pathways unique to light-upregulated genes include
those for the biosynthesis of photosynthesis components chlorophyll, carotenoid, and

166 phylloquinone, hormones like gibberellin, auxin, abscisic acid, *etc*. The pathways with unique

167 mapping to light-downregulated genes include polar auxin transport, mevalonate (MVA)

168 pathway, circadian clock, salicylic acid metabolism and signaling, reproductive plant part

169 development, root-specific gene network of NAC10 transcription factor (Supplementary Figure

170 S3).

**Identifying light-regulated transcription factors**

To identify the light-regulated transcription factors (TF), a list of rice TFs was downloaded from

the Plant Transcription Factor Database [58] and searched against the DE genes. We found 429

light-upregulated and 498 light-downregulated TFs (Supplementary Table S4). We found

WRKY, NAC, and orphans were the most abundantly expressed TFs in light, compared to many

light-downregulated bHLH, bZIP, and C3H gene family members (Table 1). Highly upregulated

(fold change $\geq$10) TFs belong to the MYB, AP2-EREBP, WRKY, Orphans, NAC, MADS, and

bHLH gene families; however, down upregulated TFs belong to the AP2-EREBP, C2C2-CO-

like, C3H, HB, and NAC gene families (Supplementary Table S4). To investigate whether TFs

targeted the MVA and MEP pathway genes, we surveyed the list of TFs and their potential

targets identified by the Plant Transcription Factor Database. We identified 23 TFs that

potentially bind to the promoter region of 6 MVA pathway genes. Two MVA pathway genes,

hydroxymethylglutaryl-CoA synthase (*HMGS*; Os08g0544900) and 3-hydroxy-3-methylglutaryl

coenzyme A reductase (*HMGR*; Os08g0512700), were targeted by bZIP and AP2-EREBP

factors, respectively, whereas the mevalonate 5-diphosphate decarboxylase (*MDD*;

Os02g0109100) gene was a target of by 13 AP2-EREBPs, one C3H protein, and one C2H2

protein (Supplementary Table S5). None of the TFs we found bind to the promoter of MEP

pathway genes.

195 **Table 1:** Light-regulated transcription factor gene families and their members.

| Gene family | Upregulated | Downregulated | Gene family | Upregulated | Downregulated |
|---|---|---|---|---|---|
| ABI3VP1 | 6 | 4 | GRF | 1 | 2 |
| Alfin-like | 1 | 4 | HB | 21 | 24 |
| AP2-EREBP | 23 | 32 | HSF | 3 | 10 |
| ARF | 8 | 8 | LIM | 2 | 1 |
| ARR-B | 1 | 4 | LOB | 2 | 1 |
| BBR-BPC | 0 | 1 | MADS | 6 | 5 |
| BES1 | 0 | 3 | mTERF | 9 | 4 |
| bHLH | 19 | 27 | MYB | 28 | 14 |
| BSD | 1 | 6 | MYB-related | 16 | 25 |
| bZIP | 10 | 27 | NAC | 21 | 19 |
| C2C2-CO-like | 4 | 7 | OFP | 2 | 1 |
| C2C2-Dof | 5 | 4 | Orphans | 20 | 10 |
| C2C2-GATA | 7 | 4 | PBF-2-like | 1 | 0 |
| C2C2-YABBY | 3 | 1 | PLATZ | 1 | 4 |
| C2H2 | 14 | 19 | RWP-RK | 4 | 2 |
| C3H | 16 | 24 | S1Fa-like | 1 | 1 |
| CAMTA | 0 | 3 | SBP | 3 | 5 |
| CCAAT | 9 | 11 | Sigma 70-like | 4 | 0 |
| CPP | 1 | 4 | SRS | 1 | 0 |
| CSD | 0 | 1 | TAZ | 1 | 1 |
| DBP | 2 | 3 | TCP | 4 | 6 |
| E2F/DP | 2 | 1 | Tify | 6 | 2 |
| EIL | 1 | 1 | Trihelix | 3 | 9 |
| FAR1 | 5 | 10 | TUB | 2 | 5 |
| FHA | 7 | 6 | ULT | 1 | 0 |
| G2-like | 9 | 5 | VOZ | 0 | 2 |
| GeBP | 0 | 3 | WRKY | 22 | 10 |
| GRAS | 8 | 7 | zf-HD | 5 | 3 |
| **Other transcriptional regulators** | | | | | |
| Gene family | Upregulated | Downregulated | Gene family | Upregulated | Downregulated |
| ARID | 2 | 1 | PHD | 12 | 22 |
| AUX/IAA | 3 | 19 | Pseudo ARR-B | 2 | 1 |
| Coactivator p15 | 0 | 1 | RB | 0 | 1 |
| DDT | 3 | 0 | Rcd1-like | 0 | 2 |
| GNAT | 12 | 8 | SET | 11 | 6 |
| HMG | 4 | 5 | SNF2 | 11 | 9 |
| Jumonji | 4 | 5 | SOH1 | 0 | 2 |
| LUG | 0 | 2 | SWI/SNF-BAF60b | 4 | 4 |
| MBF1 | 0 | 1 | SWI/SNF-SWI3 | 2 | 1 |
| MED6 | 0 | 1 | TRAF | 7 | 11 |

196
197

**Transcript splicing during photomorphogenesis**

To study how light exposure modulates transcript splicing (AS) events in the rice seedling shoots, we analyzed the transcriptome data by using SpliceGrapher [47]. We created highly accurate splice site classifiers for rice with canonical (GT-AG) and semi-canonical (GC-AG) splice sites to filter the splice junctions in the sequence reads aligned to the reference genome. After removing the false positive splice sites, we predicted chromosome-wise splice graphs from the two transcriptome sample datasets. In the dark-treated dataset, 63.9% were true positive junctions, with 5.5% novel splice sites. In the light-treated dataset, 64.3% of splice junctions were true positives, of which 5.8% were predicted as novel sites. We observed 6214 spliced genes with 9685 splicing events in the dark samples compared to the 6809 spliced genes with 10432 splicing events in the light samples (Figure 1, Table 2). The highest number of spliced genes and events were on chromosome 1, and the lowest was on chromosome 10 (Table 2). Intron retention (IR) was the most prevalent type of splicing event in both samples, followed by exon skipping (ES). Alternative 3' splicing (Alt.3') was the least common event. To resolve ambiguous combinations of donor and acceptor splice sites, we realigned the sequenced reads from the two samples to the putative transcripts and resolved the novel exons to generate splice graphs. Light-induced splicing in 2162 unique genes compared to 1567 unique genes that undergo splicing in the dark. Of the 4647 spliced genes shared between the two samples, 165 genes displayed differential splicing events.

**Table 2:** Summary of spliced genes and events under dark and light conditions detected by the SpliceGrapher and the IsoLasso workflows. IR: Intron retention; ES: Exon Skipping; Alt5': alternative 5′ splice site; Alt3': alternative 3′ splice site. Color scales: Red (Large counts) to Blue (Lower counts).

| Chromosome # | Treatment | SpliceGrapher Prediction | | | | | IsoLasso | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Genes | IR | ES | Alt5' | Alt3' | Genes | IR | ES | Alt5' | Alt3' |
| 1 | Dark | 924 | 897 | 214 | 163 | 97 | 939 | 925 | 214 | 181 | 99 |
| | Light | 1028 | 994 | 264 | 184 | 109 | 1048 | 1021 | 264 | 191 | 112 |
| 2 | Dark | 747 | 728 | 224 | 167 | 109 | 771 | 781 | 227 | 189 | 117 |
| | Light | 802 | 777 | 219 | 153 | 99 | 822 | 812 | 222 | 165 | 103 |
| 3 | Dark | 825 | 797 | 216 | 136 | 94 | 858 | 868 | 216 | 162 | 120 |
| | Light | 945 | 893 | 219 | 143 | 98 | 960 | 920 | 219 | 153 | 118 |
| 4 | Dark | 565 | 551 | 182 | 112 | 64 | 581 | 576 | 183 | 117 | 67 |
| | Light | 596 | 572 | 193 | 104 | 61 | 611 | 592 | 194 | 109 | 66 |
| 5 | Dark | 519 | 503 | 139 | 139 | 70 | 538 | 534 | 140 | 114 | 75 |
| | Light | 552 | 537 | 144 | 103 | 80 | 562 | 550 | 144 | 109 | 92 |
| 6 | Dark | 509 | 459 | 186 | 128 | 62 | 527 | 493 | 186 | 135 | 64 |
| | Light | 531 | 501 | 169 | 103 | 63 | 554 | 535 | 169 | 109 | 67 |
| 7 | Dark | 466 | 432 | 166 | 77 | 48 | 482 | 468 | 166 | 85 | 51 |
| | Light | 481 | 473 | 129 | 72 | 52 | 498 | 498 | 129 | 82 | 56 |
| 8 | Dark | 400 | 390 | 97 | 63 | 62 | 415 | 416 | 98 | 68 | 64 |
| | Light | 463 | 433 | 134 | 62 | 84 | 475 | 457 | 134 | 67 | 87 |
| 9 | Dark | 341 | 342 | 79 | 53 | 32 | 349 | 359 | 79 | 64 | 37 |
| | Light | 375 | 361 | 118 | 49 | 38 | 382 | 377 | 118 | 53 | 39 |
| 10 | Dark | 290 | 275 | 94 | 43 | 29 | 295 | 288 | 94 | 49 | 36 |
| | Light | 308 | 295 | 102 | 61 | 32 | 314 | 307 | 102 | 65 | 35 |
| 11 | Dark | 304 | 266 | 113 | 49 | 35 | 313 | 278 | 113 | 53 | 36 |
| | Light | 363 | 306 | 150 | 51 | 35 | 374 | 325 | 150 | 57 | 37 |
| 12 | Dark | 324 | 310 | 117 | 69 | 45 | 329 | 315 | 117 | 74 | 46 |
| | Light | 365 | 347 | 150 | 63 | 53 | 372 | 359 | 155 | 68 | 61 |

In our differentially expressed and spliced genes (DES), we found that approximately 30%
(4369) differentially expressed (DE) genes also undergo splicing (AS). Approximately 10%
(2440) of the non-DE genes undergo splicing (Figure 1). Light-upregulated DES genes show
enrichment for various molecular functions like oxidoreductase, hydrolase, isomerase activity,
and RNA binding, which play roles in photosynthesis, transmembrane transport, small molecule
metabolic process, and located in the cellular components, chloroplast, and membrane-bound
organelles. In contrast, the light-downregulated DES genes enriched for RNA processing,
response to stress, protein binding, and macromolecule metabolic process, and localized in
cellular components cytoplasm and nucleus (Supplementary Figure S4).

We also surveyed the DES aspects of the 57 known spliced genes in rice [23,59] and observed
splicing in 41 genes (Table 3). Of these, seven spliced under dark conditions, three under light
conditions, and 31 genes spliced under both dark and light conditions. Of the known genes, we
also observed that 18 genes were light-upregulated and 24 downregulated. Only 32 genes
showed a DES profile. Three light-upregulated genes (Os02g0130600, Os05g0348100,
Os12g0567300) show complete splicing in light compared to intron retention in the dark. In
contrast, the two light-downregulated genes (Os02g0666200 and Os04g0656100) show intron
retention.

**Table 3:** Expression and splicing profile of genes in our data that are known to undergo splicing from previous reports

| Known spliced genes | Spliced in | Regulation in light | Gene description |
|---|---|---|---|
| Os01g0619000 | Dark, Light | Down | RNA recognition motif domain-containing protein |
| Os01g0649900 | Dark, Light | - | GDSL esterase/lipase protein 20 |
| Os01g0764000 | Dark, Light | Up | PHI glutathione s-transferase 2 |
| Os01g0834400 | Dark | - | HAP3A subunit of CCAAT-box binding complex |
| Os02g0122800 | Dark, Light | Down | Similar to Arginine/serine-rich splicing factor |
| Os02g0130600 | Dark | Up | Conserved hypothetical protein |
| Os02g0161900 | Dark, Light | Down | Rice ubiquitin2 |
| Os02g0197900 | Dark, Light | Down | OsSTA53 |
| Os02g0274900 | Dark, Light | Up | Similar to metabolite transport protein CSBC |
| Os02g0577100 | Dark, Light | Down | Zinc finger, RING/PHD-type domain containing protein |
| Os02g0666200 | Light | Down | Aquaporin |
| Os02g0834000 | Dark, Light | Up | Rac-like GTP-binding protein 5 |
| Os02g0291000 | - | Down | Calcineurin B-like protein 7 |
| Os02g0291400 | - | - | Calcineurin B-like protein 8 |
| Os02g0823100 | - | Up | Plasma membrane intrinsic protein 1;3 |
| Os03g0265600 | Dark, Light | Down | Transformer-2-like protein |
| Os03g0314100 | Dark, Light | Down | DEAD-like helicase |
| Os03g0395900 | Dark, Light | Down | Splicing factor |
| Os03g0670700 | Dark, Light | Down | Glycine-rich rna-binding protein 3 |
| Os03g0698500 | Dark, Light | Down | Similar to Yippee-like protein 3 |
| Os03g0745000 | Dark, Light | - | Heat stress transcription factor A-2a |
| Os03g0717600 | - | Down | Zinc finger, C2H2-type |
| Os04g0115400 | Dark, Light | Down | D111/G-patch domain containing protein |
| Os04g0649100 | Dark, Light | - | Shattering abortion 1 |
| Os04g0656100 | Light | Down | $H^+$-ATPase |
| Os04g0665800 | Dark, Light | Up | Similar to H1005F08.12 protein |
| Os04g0402300 | - | - | Cysteine-type peptidase |
| Os04g0479200 | - | Up | Similar to NAD-dependent isocitrate dehydrogenase c;1 |

| | | | |
|---|---|---|---|
| Os05g0348100 | Dark | Up | Similar to CRR23 (chlororespiratory reduction 23) |
| Os05g0463800 | Dark, Light | Down | HAP3C subunit of CCAAT-box binding complex |
| Os05g0554400 | Dark, Light | Down | Phosphatidyl serine synthase family protein |
| Os05g0574700 | Dark, Light | - | Similar to cDNA clone:002-182-C01 |
| Os05g0534400 | - | Up | Calcineurin B-like protein 4 |
| Os05g0548900 | - | Up | Phosphoethanolamine N-Methyltransferase 2 |
| Os06g0172800 | Light | - | Similar to alkaline alpha galactosidase 2 |
| Os06g0651600 | Dark, Light | Down | Protein phosphatase 2C58 |
| Os06g0727200 | Dark | Down | Catalase B |
| Os06g0128500 | - | Up | Ribosomal protein L47, mitochondrial family protein |
| Os06g0133000 | - | - | Glutinous Endosperm, waxy |
| Os06g0506600 | - | - | Ubiquitin-conjugating enzyme 17 |
| Os07g0490400 | Dark, Light | Up | FK506 binding protein 20-2 |
| Os07g0574800 | - | Up | Tubulin alpha-1 chain |
| Os07g0613300 | - | - | Similar to PAUSED |
| Os08g0191600 | Dark, Light | Up | Autophagy associated gene 8C |
| Os08g0530400 | Dark, Light | Up | Moco containing protein, Similar to sulfite oxidase |
| Os08g0436200 | - | - | Zinc finger, RING/PHD-type domain containing protein |
| Os10g0115600 | Dark, Light | Down | U1 snRNP 70K |
| Os10g0535800 | Dark, Light | - | Cys-rich domain containing protein |
| Os10g0564900 | Dark | Down | Similar to protein kinase CK2 regulatory subunit CK2B2 |
| Os10g0567400 | Dark, Light | Up | Chlorophyll a oxygenase 1 |
| Os10g0577900 | Dark, Light | - | Glycerol-3-phosphate acyltransferase |
| Os10g0411500 | - | Down | Q calmodulin-binding region domain containing protein |
| Os11g0157100 | Dark, Light | Down | Cyclin-T1-4 |
| Os11g0700500 | Dark | - | MybAS1 |
| Os11g0600700 | - | Up | Zinc finger, RING domain protein |
| Os12g0567300 | Dark | Up | MybAS2, R2R3-Myb |
| Os12g0632000 | Dark, Light | Down | Glycine-rich Protein GRP162 |

**Light-mediated shift in gene expression and splicing events**

*Circadian clock pathway genes*

The circadian clock is entrained by light, and the diurnal photoperiod regulated clock genes [60,61].
We investigated the light-regulated DES profile of clock genes. Most of the circadian clock
genes spliced under either or both conditions. Four genes, *Casein kinase alpha subunit* (*CK2α-
1/Hd6*, Os03g0762000), *Casein kinase beta subunit* (*CK2β-2*, Os10g0564900), *Phytochrome B*
(*PhyB*, Os03g0309200;), and *Timing of CAB Expression 1* (*TOC1*, Os02g0618200), showed
differential splicing (only reference splicing events or complete splicing and no novel isoforms)
upon light treatment compared to the presence of unspliced intron in the absence of light
(Supplementary Figure S5). In contrast, only one gene, *Phytochrome Interacting Factor 3* (*PIF3*,
Os01g0286100), showed intron retention under light. All five genes were upregulated in light
(Supplementary Figure S5). Seven genes, *Casein kinase alpha subunit* (*CK2α-2*,
Os03g0763000), *Casein kinase beta subunit* (*CK2β-1*, Os07g0495100), blue light receptor
*Cryptochrome 2* (*CRY2*, Os02g0625000), *Gigantea* (*GI*, Os01g0182600), *Pseudo-Response
Regulator 37* or *Heading date 2* (*PRR37*, *Hd2*, Os07g0695100), and *Pseudo-Response Regulator
73* (*PRR73*, Os03g0284100), showed differential splicing under both conditions (Supplementary
Figure S5, S6). Exon skipping and alternative 5' (Alt.5') splicing events in the *CK2α-2*
(Os03g0763000) gene were induced in light. Both *CK2β-1* (Os07g0495100) and *GI*
(Os01g0182600) display incomplete splicing of intron (IR) in the dark. *PRR37* showed three
types of splicing events (Alt.3', ES, and IR) under light, whereas the second IR event observed
toward the 3' end under dark was spliced in light. Conversely, *PRR73* showed three types of
events (Alt.3', ES, and IR) under dark conditions, whereas under light conditions, two new Alt.5'
events were identified. Expression of both PRR genes (*PRR37* and *PRR73*) was light
downregulated (Supplementary Figure S5). Light-upregulated *PRR95* shows an additional
unspliced intron in light.

*MVA and MEP pathway genes*

Light regulates the biosynthesis and accumulation of secondary plant products like isoprenoid-
derived metabolites required for plant growth and development. Isoprenoid precursors are
synthesized via cytosolic MVA and plastid-localized MEP pathways, and their fluxes are

regulated by light [62,63], and their products are essential for the successful development and

function of chloroplast. We observed that MEP pathway genes were light-upregulated compared

to the downregulated MVA pathway genes (Figure 2). Light positively regulates MEP pathway

activity, whereas MVA pathway activity appears negatively regulated by light [64].

We investigated the light-induced differential splicing events in the MEP and MVA pathway

genes. Four MVA pathway genes (*acetyl-CoA acetyltransferase: AACT, hydroxymethyl glutaryl*

*CoA synthase: HMGS, mevalonate kinase: MK, phosphomevalonate kinase: PMK*) and 3 MEP

pathway genes (*1-deoxy-D-xylulose-5-phosphate synthase: DXS, 1-deoxy-D-xylulose-5-*

*phosphate reductoisomerase: DXR, 2-C-methyl-D-erythritol2,4-cyclodiphosphate synthase:*

*MECPS*) showed intron retention events in dark (Figure 2). Gene *MK* showed a novel light-

induced exon-skipping event.

**Figure 2:** Gene expression and splicing of cytosolic mevalonate (MVA) and plastid-localized

methyl erythritol 4-phosphate (MEP) pathway genes. *AACT*, acetyl-CoA acetyltransferase;

*HMGS*, hydroxymethyl glutaryl CoA synthase; *HMGR*, 3-hydroxy-3-methylglutaryl-coenzyme

A reductase; *MK*, mevalonate kinase; *PMK*, phosphomevalonate kinase; *MDD*, mevalonate

diphosphosphate decarboxylase; *DXS*, 1-deoxy-D-xylulose-5-phosphate synthase; *DXR*, 1-

deoxy-D-xylulose-5-phosphate reductoisomerase; *CDPMES*, 2-C-methyl-D-erythritol4-

phosphate cytidylyl transferase; *CDPMEK*, 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase;

*MECPS*, 2-C-methyl-D-erythritol2,4-cyclodiphosphate synthase; *HDS*, 4-hydroxy-3-methylbut-

2-enyldiphosphate synthase; *HDR*, 4-hydroxy-3-methylbut-2-enyldiphosphate reductase. Blue:

intron retention, Green: exon skipping, Orange: Alt3' splicing, Purple: Alt5' splicing

**lncRNA discovery and splicing during photomorphogenesis**

lncRNAs are known to regulate diverse biological processes in plants, including

photomorphogenesis [65], and as described above, are categorized into four major types based on

their location in the gene space and on the DNA strand. We used our transcriptome data to

characterize light-induced rice lncRNAs (Supplementary Figure S7). We identified 1485 and

1407 lncRNA transcripts in the dark and light samples, of which 309 were common. Many

lncRNAs from intergenic and intronic regions were identified (Supplementary Figure S7). To

discover novel lncRNAs, the transcripts from our datasets were queried against the custom

BLAST database of rice lncRNAs. About 44% of the total lncRNA transcripts (2583) found a

match in the BLAST database (Supplementary Table S6). 173 lncRNAs spliced under dark and

142 spliced under light conditions, most located in the intergenic regions (Supplementary Figure

S7). Interestingly, all the spliced lncRNA genes from both datasets were from Chr1, Chr10,

Chr11, and Chr12. The maximum number of spliced lncRNA genes was from Chr1, followed by

Chr11, 12, and 10. Among all the splicing events, IR was predominant under both conditions.

We found lncRNA transcripts from light (TCONS_00012152) and dark (TCONS_00011794)

datasets that were present on the antisense strand overlapping an exon region of the *MADS27*

(Os02g0579600) gene. The transcript from the light-treated sample appears longer and showed

two alternative 3' splice sites and one exon skipping event. The TCONS_00011794 may

compete with MADS27 to form a complex with miRNA osa-miR444a known to play multiple

roles in the nitrate-dependent development pathway [66]. Among the list of lncRNAs present on

the antisense strand overlapping annotated genes, we identified a light-induced lncRNA

(TCONS_00001515) overlapping *ERF99* transcription factor (Os01g0868000) exon that may

play a role in its silencing. *ERF99* plays a central role in mediating abiotic stress responses [67].

Similarly, lncRNA transcript (TCONS_00015825) is present exclusively in light is transcribed

from the intronic region of the antisense strand of a B-type response regulator gene

(Os03g0224200), known for its involvement in cytokinin signaling, meristem maintenance, and

stress response [68,69]. The lncRNA transcript (TCONS_00026617) present exclusively in the dark

overlaps the intronic region of a phytochrome-interacting factor gene *PIF14* (Os07g0143200).

*PIF14* is known to bind the active form of phytochrome B and plays a crucial role in cross-talk

between light and stress signaling [70].

## Discussion

Light is known to induce changes in the transcriptome, metabolome, and proteome of the plants

[71], which not only regulates the development, function, and physiology of the chloroplast but

also provides signals for modulating the plant's morphological, developmental, and

physiological adaptions in response to growth environment [7,38,72–74]. Though plants experience

constantly changing light conditions under the natural environment daily during their lifetime,

early development from germination to the seedling stage and acquiring the full autotrophic

capability is very much programmed by exposure to light. Therefore, we investigated

transcriptome modulation in rice seedlings during photomorphogenesis. Compared to earlier

studies in rice and *Arabidopsis*, where ~20% of the genes were reported differentially expressed

in dark-grown etiolated seedlings compared to light-exposed green seedlings [8,75,76], we observed

that the transition from skotomorphogenesis to photomorphogenesis alters differential expression

of ~38% of the rice genes.

Functional annotation of the differentially expressed genes enriched for roles in secondary

metabolism, chloroplast-related biosynthetic pathways, hormone biosynthesis and signaling,

amino acid biosynthesis, fatty acid metabolism, *etc*. (Supplementary Figure S3, Supplementary

Table S3). This result was expected based on the earlier reports on light regulation of such events

[22,77].

The plastids develop into chloroplasts by following an essential step of developing the thylakoid

membrane system and recruiting and assembling components of light and dark reactions to

establish a functional photosynthetic process. Many terpenoid compounds are essential

participants in plants' light-harvesting function and provide protection against damage from

reactive oxygen species (ROS). The basic isoprenoid units for terpenoid biosynthesis, such as

tocopherols, plastoquinones, carotenoids, chlorophylls, and precursors of the growth hormones

gibberellins and abscisic acid, are synthesized by the plastid-localized MEP pathway [78,79]. At the

same time, isoprenoid units synthesized by the MVA pathway contribute to the synthesis of

triterpenes, phytosterols, and phytohormones [80]. MEP and MVA pathways complement and

contribute to the formation of chlorophylls and carotenoids required for plastid development [81].

MEP pathway genes were light-upregulated in our transcriptome data compared to the light-

downregulated MVA pathway genes (Figure 2). These results were consistent with earlier

reports [82,83] and confirm that light acts as a critical regulator in modulating the availability of

isoprenoid precursors during photomorphogenesis [84]. Arabidopsis circadian clock genes (*LHY*,

*PRR9*, *CCA1*, *TOC1*) regulate MVA and MEP pathways [84]. In our dataset, we did not observe

any significant change in the expression of *LHY/CCA1*(Os08g0157600), *PRR95* (Os09g0532400,

a homolog of *AtPRR9*), and *TOC1* (Os02g0618200). However, all three genes spliced

differently. Under the light, *TOC1* showed no splicing but retained two introns in the dark. We

observed one exon-skipping event in light for the *LHY/CCA1* gene, whereas *PRR95* showed one

additional intron retention (Supplementary Figure S5).

Previous transcriptome studies showed that ~42% of *Arabidopsis*, ~63% of *soybean*, and ~56%

of maize genes undergo alternative splicing events [85–87]. Studies on rice by Wang and Brendel [23]

reported IR for 69% of 6568 AS genes, and Campbell *et al.* [88] reported 44.7% of 8772 AS genes.

In our dataset, 4369 genes undergo differential expression and alternative splicing, called DES

genes (Figure 1A). About 80% (5395) of the spliced genes (6809) showed IR events, suggesting

that IR is the major splicing event. The more significant number of splicing events in the light-

treated samples suggests that light-mediated gene expression and post-transcriptional mRNA

splicing play an important role in photomorphogenesis. The splicing events in rice circadian

clock genes suggest that IR regulates the expression of PRR genes (Supplementary Figure S5).

Many of the light-regulated genes of the MEP and MVA pathways show IR events, suggesting

that light plays a positive role in completing the splicing and provides a hypothesis that exposure

to light is an adaptive feature and even if the gene is expressed in dark, it stays unspliced until it

encounters light. Our data showed the presence of all three states, partially spliced in the dark

(*GI*, Os01g0182600), fully spliced in light (*GI*, Os01g0182600), and both states in light (*CK2β-*

*1*, Os07g0495100) (Supplementary Figure S6).

lncRNAs play regulatory functions in essential biological processes such as vernalization,

photomorphogenesis, and stress regulation [37,89,90] and display splicing [91]. 6480 long intergenic

noncoding RNAs (lincRNAs) were identified from 200 *Arabidopsis* transcriptome datasets [92];

however, we identified 827 lincRNAs in the dark and 727 lincRNAs in the light datasets.

Corona-Gomez and coworkers [93] characterized the splicing conservation of lncRNAs in

*Brassicaceae*, which revealed that ~18% of lincRNAs display splicing conservation in at least

one exon. In our dataset, ~10% of lncRNA transcripts undergo splicing, and the majority were of

long intergenic noncoding RNA type (Supplementary Figure S7). Komiya and colleagues

reported [94] that phased small interfering RNAs (phasiRNAs) are generated from over 700

lincRNAs, and these phasiRNAs bind to rice argonaute protein MEL1. MEL1 has a specific

function in developing pre-meiotic germ cells and the progression of meiosis. In our dataset,

many lincRNAs transcribed from MEL1-phasiRNA clusters appeared.

Identification of light-induced lncRNAs antisense to *ERF99*, B-type response regulator genic

region may play a role in its transcriptional regulation. *ERF99* is known to modulate root

architecture and downregulated in crown root primordia [95]. *MADS27* gene targeted by lncRNAs

present in both dark and light conditions. However, the dark lncRNA (TCONS_00011794) is

longer and, based on its predicted RNA fold structure and lower free energy, is likely more stable

than its transcript TCONS_00012152 in light. For the MADS27 gene, we also observed an

additional IR event in light. MADS27-miR444 complex is known to play a role in plant

development in a nitrate-dependent manner [96–98]. Therefore, we investigated the nitrogen

assimilation pathway genes. Nitrogen assimilation is necessary for sustaining plant growth and

development. Various nitrogen assimilatory enzymes are known to show isoform and cellular

component-specific responses under light and dark conditions [99,100]. We observed light

upregulation for the nitrate transporter, nitrate reductase, and nitrite reductase genes. We also

observed differential splicing patterns for these genes (Supplementary Figure S8).

**Conclusion**

This study suggests that light is a significant regulatory factor controlling genome-wide gene

expression through an alternative splicing mechanism in rice. All spliced genes did not

necessarily produce novel isoforms, which indicates the coupling of AS and nonsense-mediated

decay (NMD). NMD prevents the translation of mutant mRNAs harboring potential premature

stop codons by targeting them for degradation. In *Arabidopsis*, 77.2% of light-regulated AS

events exhibit NMD features within a splicing isoform [101]. We conclude that light induces a

significant number of splicing events in rice protein-coding and noncoding lncRNA genes. This

photomorphogenesis transcriptome study is a valuable resource for lncRNA research in rice and

provides insights into the portion of the genome regulated at the level of alternative splicing in

response to light. We expect the condition-dependent novel splice events discovered in this study

will help improve reference rice genome annotation.

**Acknowledgments**

The authors acknowledge Dr. Mamatha Hanumappa and Dr. Samuel Fox for helping with the

experimental setup and preparation of libraries for sequencing. We recognize the Center for

Genome Research and Biocomputing (CGRB) at Oregon State University for providing

sequencing services and support and maintaining Jaiswal Laboratory's high-performance

computing infrastructure for data analysis. We also acknowledge Prof. ASN Reddy for critical

suggestions on the earlier draft and Justin Preece for technical and English editing.

**Author's Contributions**

PJ conceptualized the project. PG performed the data analysis. PG and PJ wrote the manuscript.

**Funding**

This work was supported by the startup funds provided to PJ by the Department of Botany and

Plant Pathology in the College of Agricultural Sciences at Oregon State University. Authors are

partially supported by the National Science Foundation awards #1127112, #1340112, #2029854.

Funding agencies had no role in project design and data analytics.

**Conflict of interest**

The authors declare no conflict of interest.

**Availability of data**

The sequence reads and metadata were deposited at EMBL-EBI ArrayExpress (accession

number E-MTAB-5689). https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-5689/#.

**Figure legends**

**Figure 1:** Summary of differential gene expression and transcript splicing observed in rice

seedling shoots undergoing photomorphogenesis.

**Figure 2:** Gene expression and splicing of cytosolic mevalonate (MVA) and plastid-localized

methyl erythritol 4-phosphate (MEP) pathway genes. *AACT*, acetyl-CoA acetyltransferase;

*HMGS*, hydroxymethyl glutaryl CoA synthase; *HMGR*, 3-hydroxy-3-methylglutaryl-coenzyme

A reductase; *MK*, mevalonate kinase; *PMK*, phosphomevalonate kinase; *MDD*, mevalonate

diphosphosphate decarboxylase; *DXS*, 1-deoxy-D-xylulose-5-phosphate synthase; *DXR*, 1-

deoxy-D-xylulose-5-phosphate reductoisomerase; *CDPMES*, 2-C-methyl-D-erythritol4-

phosphate cytidylyl transferase; *CDPMEK*, 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase;

*MECPS*, 2-C-methyl-D-erythritol2,4-cyclodiphosphate synthase; *HDS*, 4-hydroxy-3-methylbut-

2-enyldiphosphate synthase; *HDR*, 4-hydroxy-3-methylbut-2-enyldiphosphate reductase. Blue:

intron retention, Green: exon skipping, Orange: Alt3' splicing, Purple: Alt5' splicing

**Table legends**

**Table 1:** Light-regulated transcription factor gene families and their members.

**Table 2:** Summary of spliced genes and events under dark and light conditions detected by the

SpliceGrapher and the IsoLasso workflows. IR: Intron retention; ES: Exon Skipping; Alt5':

alternative 5′ splice site; Alt3': alternative 3′ splice site. Color scales: Red (Large counts) to Blue

(Lower counts).

**Table 3:** Expression and splicing profile of genes in our data known to undergo splicing from

previous reports

**Supplementary material**

**Supplementary Figure S1:**

Expression pattern of differentially expressed genes (14,766) with target FDR controlled at 5%

**Supplementary Figure S2:**

Gene Ontology enrichment analysis of differentially expressed genes.

**Supplementary Figure S3:**

Pathway enrichment analysis using the Plant Reactome. **(A)** Plant Reactome pathway enrichment

analysis plots, **(B)** Unique and shared pathways enriched for the light upregulated and

downregulated gene sets; **(C)** Counts of genes mapped to some of the common Plant Reactome

pathways. Light upregulated (green) and downregulated (grey).

**Supplementary Figure S4**:

Bar plots of most significantly enriched GO terms for DES genes with -log10 transformed FDR

values. Light-upregulated (green), light-downregulated (grey).

**Supplementary Figure S5**:

The summary of transcript splicing events and expression pattern of circadian clock genes during

morphogenesis.

**Supplementary Figure S6**:

Transcript graphs showing splicing of *Casein kinase beta subunit:CK2β-1* (Os07g0495100),

*Gigantea:GI* (Os01g0182600) genes during photomorphogenesis.

**Supplementary Figure S7**:

Classification of lncRNAs identified in dark and light.

**Supplementary Figure S8:**

An overviw of splicing and expression pattern of nitrogen assimilation cycle genes during

photomorphogenesis. NRT: Nitrate transporter; NR: Nitrate reductase; NiR: Nitrate reductase.

**Supplementary Table S1**:

Summary of the RNA-Seq read mapping to the reference rice genome.

**Supplementary Table S2:**

Differentially expressed genes

**Supplementary Table S3:**

Plant Reactome Pathway enrichment analysis for light-regulated genes

**Supplementary Table S4:**

Transcription factor genes regulated in light

**Supplementary Table S5:**

Transcription factors regulating MVA pathway genes

**Supplementary Table S6:**

lncRNA annotation using BLAST

**Reference**

1.  Kami, C., Lorrain, S., Hornitschek, P. & Fankhauser, C. Chapter Two - Light-Regulated Plant Growth and Development. in *Current Topics in Developmental Biology* (ed. Timmermans, M. C. P.) vol. 91 29–66 (Academic Press, 2010).

2.  Poolman, M. G., Kundu, S., Shaw, R. & Fell, D. A. Responses to Light Intensity in a Genome-Scale Model of Rice Metabolism. *Plant Physiology* **162**, 1060–1072 (2013).

3.  Li, H. *et al.* A Natural Light/Dark Cycle Regulation of Carbon-Nitrogen Metabolism and Gene Expression in Rice Shoots. *Front. Plant Sci.* **7**, (2016).

4.  Bai, S. *et al.* BBX16, a B-box protein, positively regulates light-induced anthocyanin accumulation by activating MYB10 in red pear. *Plant Biotechnology Journal* **0**,.

5.  Tobin, E. M. & Silverthorne, J. Light Regulation of Gene Expression in Higher Plants. *Annual Review of Plant Physiology* **36**, 569–593 (1985).

6.  Ma, L. *et al.* Light Control of Arabidopsis Development Entails Coordinated Regulation of Genome Expression and Cellular Pathways. *Plant Cell* **13**, 2589–2608 (2001).

7.  Casal, J. J. & Yanovsky, M. J. Regulation of gene expression by light. *Int. J. Dev. Biol.* **49**, 501–511 (2004).

8.  Jiao, Y., Ma, L., Strickland, E. & Deng, X. W. Conservation and Divergence of Light-Regulated Genome Expression Patterns during Seedling Development in Rice and Arabidopsis. *The Plant Cell* **17**, 3239–3256 (2005).

9.  Majláth, I. *et al.* Effect of light on the gene expression and hormonal status of winter and spring wheat plants during cold hardening. *Physiologia Plantarum* **145**, 296–314 (2012).

10. Petrillo, E., Herz, M. A. G., Barta, A., Kalyna, M. & Kornblihtt, A. R. Let there be light: Regulation of gene expression in plants. *RNA Biology* **11**, 1215–1220 (2014).

11. Quail, P. H. Phytochrome photosensory signalling networks. *Nature Reviews Molecular Cell Biology* **3**, 85–93 (2002).

12. Lin, C. & Shalitin, D. Cryptochrome Structure and Signal Transduction. *Annual Review of Plant Biology* **54**, 469–496 (2003).

13. Huang, X., Ouyang, X. & Deng, X. W. Beyond repression of photomorphogenesis: role switching of COP/DET/FUS in light signaling. *Current Opinion in Plant Biology* **21**, 96–103 (2014).

14. Sheerin, D. J. *et al.* Light-Activated Phytochrome A and B Interact with Members of the
SPA Family to Promote Photomorphogenesis in Arabidopsis by Reorganizing the
COP1/SPA Complex. *Plant Cell* **27**, 189–201 (2015).

15. Leivar, P. *et al.* Multiple phytochrome-interacting bHLH transcription factors repress
premature photomorphogenesis during early seedling development in darkness. *Curr Biol*
**18**, 1815–1823 (2008).

16. Shin, J. *et al.* Phytochromes promote seedling light responses by inhibiting four negatively-
acting phytochrome-interacting factors. *Proc Natl Acad Sci U S A* **106**, 7660–7665 (2009).

17. Xie, C., Zhang, G., An, L., Chen, X. & Fang, R. Phytochrome-interacting factor-like
protein OsPIL15 integrates light and gravitropism to regulate tiller angle in rice. *Planta*
(2019) doi:10.1007/s00425-019-03149-8.

18. Liu, M.-J., Wu, S.-H., Chen, H.-M. & Wu, S.-H. Widespread translational control
contributes to the regulation of Arabidopsis photomorphogenesis. *Molecular Systems*
*Biology* **8**, 566 (2012).

19. Kianianmomeni, A. More light behind gene expression. *Trends in Plant Science* **19**, 488–
490 (2014).

20. Wu, H.-P. *et al.* Genome-wide analysis of light-regulated alternative splicing mediated by
photoreceptors in Physcomitrella patens. *Genome Biology* **15**, R10 (2014).

21. Shikata, H. *et al.* Phytochrome controls alternative splicing to mediate light responses in
Arabidopsis. *PNAS* **111**, 18781–18786 (2014).

22. Mancini, E. *et al.* Acute Effects of Light on Alternative Splicing in Light-Grown Plants.
*Photochem Photobiol* **92**, 126–133 (2016).

23. Wang, B.-B. & Brendel, V. Genomewide comparative analysis of alternative splicing in
plants. *Proc Natl Acad Sci U S A* **103**, 7175–7180 (2006).

24. Ner-Gaon, H. *et al.* Intron retention is a major phenomenon in alternative splicing in
Arabidopsis. *The Plant Journal* **39**, 877–885 (2004).

25. Filichkin, S. A. *et al.* Abiotic Stresses Modulate Landscape of Poplar Transcriptome via
Alternative Splicing, Differential Intron Retention, and Isoform Ratio Switching. *Front.*
*Plant Sci.* **9**, (2018).

26. Chaudhary, S. *et al.* Alternative Splicing and Protein Diversity: Plants Versus Animals.
*Front Plant Sci* **10**, 708 (2019).

27.   Ganie, S. A. & Reddy, A. S. N. Stress-Induced Changes in Alternative Splicing Landscape
in Rice: Functional Significance of Splice Isoforms in Stress Tolerance. *Biology (Basel)* **10**,
309 (2021).

28.   Gao, P. *et al.* Alternative splicing dynamics and evolutionary divergence during
embryogenesis in wheat species. *Plant Biotechnol J* **19**, 1624–1643 (2021).

29.   Mercer, T. R., Dinger, M. E. & Mattick, J. S. Long noncoding RNAs: insights into
functions. *Nature Reviews Genetics* **10**, 155–159 (2009).

30.   Liu, J., Wang, H. & Chua, N.-H. Long noncoding RNA transcriptome of plants. *Plant*
*Biotechnol J* **13**, 319–328 (2015).

31.   Palos, K., Yu, L., Railey, C. E., Nelson Dittrich, A. C. & Nelson, A. D. L. Linking
discoveries, mechanisms, and technologies to develop a clearer perspective on plant long
noncoding RNAs. *Plant Cell* **35**, 1762–1786 (2023).

32.   Wang, X., Fan, H., Wang, B. & Yuan, F. Research progress on the roles of lncRNAs in
plant development and stress responses. *Front Plant Sci* **14**, 1138901 (2023).

33.   Liu, H., Wang, R., Mao, B., Zhao, B. & Wang, J. Identification of lncRNAs involved in rice
ovule development and female gametophyte abortion by genome-wide screening and
functional analysis. *BMC Genomics* **20**, 90 (2019).

34.   Zhang, Y.-C. *et al.* Genome-wide screening and functional analysis identify a large number
of long noncoding RNAs involved in the sexual reproduction of rice. *Genome Biol* **15**,
(2014).

35.   Chen, L. *et al.* Genome-wide analysis of long noncoding RNAs affecting roots development
at an early stage in the rice response to cadmium stress. *BMC Genomics* **19**, 460 (2018).

36.   Li, W. Q. *et al.* Genome-wide identification and characterization of long noncoding RNAs
responsive to Dickeya zeae in rice. *RSC Adv.* **8**, 34408–34417 (2018).

37.   Wang, Y. *et al.* Arabidopsis noncoding RNA mediates control of photomorphogenesis by
red light. *PNAS* **111**, 10359–10364 (2014).

38.   Fox, S. E. *et al.* De Novo Transcriptome Assembly and Analyses of Gene Expression
during Photomorphogenesis in Diploid Wheat Triticum monococcum. *PLoS One* **9**, (2014).

39.   najoshi. najoshi/sickle. (2020).

40.   Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-
Seq. *Bioinformatics* **25**, 1105–1111 (2009).

41.   Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562–578 (2012).

42.   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

43.   Leng, N. *et al.* EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq
experiments. *Bioinformatics* **29**, 1035–1043 (2013).

44.   The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing
strong. *Nucleic Acids Res* **47**, D330–D338 (2019).

45.   Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C. & Kanehisa, M. KAAS: an automatic
genome annotation and pathway reconstruction server. *Nucleic Acids Res* **35**, W182–W185
(2007).

46.   Naithani, S. *et al.* Plant Reactome: a resource for plant pathways and comparative analysis.
*Nucleic Acids Res* **45**, D1029–D1039 (2017).

47.   Rogers, M. F., Thomas, J., Reddy, A. S. & Ben-Hur, A. SpliceGrapher: detecting patterns
of alternative splicing from RNA-Seq data in the context of gene models and EST data.
*Genome Biol* **13**, R4 (2012).

48.   Li, W., Feng, J. & Jiang, T. IsoLasso: A LASSO Regression Approach to RNA-Seq Based
Transcriptome Assembly. *J Comput Biol* **18**, 1693–1707 (2011).

49.   Kang, Y.-J. *et al.* CPC2: a fast and accurate coding potential calculator based on sequence
intrinsic features. *Nucleic Acids Res* **45**, W12–W16 (2017).

50.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*
**30**, 1236–1240 (2014).

51.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *Journal of Molecular Biology* **215**, 403–410 (1990).

52.   Yi, X., Zhang, Z., Ling, Y., Xu, W. & Su, Z. PNRD: a plant noncoding RNA database.
*Nucleic Acids Res* **43**, D982–D989 (2015).

53.   Paytuví Gallart, A., Hermoso Pulido, A., Anzar Martínez de Lagrán, I., Sanseverino, W. &
Aiese Cigliano, R. GREENC: a Wiki-based database of plant lncRNAs. *Nucleic Acids Res*
**44**, D1161–D1166 (2016).

54.   Zhang, Z., Xu, Y., Yang, F., Xiao, B. & Li, G. RiceLncPedia: a comprehensive database of
rice long noncoding RNAs. *Plant Biotechnol J* **19**, 1492–1494 (2021).

55. Hofacker, I. L. Vienna RNA secondary structure server. *Nucleic Acids Res* **31**, 3429–3431 (2003).

56. Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R. & Hofacker, I. L. The Vienna RNA Websuite. *Nucleic Acids Res* **36**, W70–W74 (2008).

57. Naithani, S. *et al.* Plant Reactome: a knowledgebase and resource for comparative pathway analysis. *Nucleic Acids Res* **48**, D1093–D1103 (2020).

58. Jin, J. *et al.* PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Res* **45**, D1040–D1045 (2017).

59. Li, J. *et al.* A subgroup of MYB transcription factor genes undergoes highly conserved alternative splicing in Arabidopsis and rice. *J Exp Bot* **57**, 1263–1273 (2006).

60. Sugiyama, N., Izawa, T., Oikawa, T. & Shimamoto, K. Light regulation of circadian clock-controlled gene expression in rice. *The Plant Journal* **26**, 607–615 (2001).

61. Matsuzaki, J., Kawahara, Y. & Izawa, T. Punctual Transcriptional Regulation by the Rice Circadian Clock under Fluctuating Field Conditions[OPEN]. *Plant Cell* **27**, 633–648 (2015).

62. Cordoba, E., Salmi, M. & León, P. Unravelling the regulatory mechanisms that modulate the MEP pathway in higher plants. *J Exp Bot* **60**, 2933–2943 (2009).

63. Vranová, E., Coman, D. & Gruissem, W. Network Analysis of the MVA and MEP Pathways for Isoprenoid Synthesis. *Annual Review of Plant Biology* **64**, 665–700 (2013).

64. Vranová, E., Coman, D. & Gruissem, W. Structure and Dynamics of the Isoprenoid Pathway Network. *Molecular Plant* **5**, 318–333 (2012).

65. Liu, X., Hao, L., Li, D., Zhu, L. & Hu, S. Long Noncoding RNAs and Their Biological Roles in Plants. *Genomics Proteomics Bioinformatics* **13**, 137–147 (2015).

66. Zhou, R. *et al.* Analysis of Rice Transcriptome Reveals the LncRNA/CircRNA Regulation in Tissue Development. *Rice (N Y)* **14**, 14 (2021).

67. Serra, T. S. *et al.* OsRMC, a negative regulator of salt stress response in rice, is regulated by two AP2/ERF transcription factors. *Plant Mol Biol* **82**, 439–455 (2013).

68. Argueso, C. T., Ferreira, F. J. & Kieber, J. J. Environmental perception avenues: the interaction of cytokinin and environmental response pathways. *Plant, Cell & Environment* **32**, 1147–1160 (2009).

69.  Worthen, J. M. *et al.* Type-B response regulators of rice play key roles in growth,

development and cytokinin signaling. *Development* **146**, dev174870 (2019).

70.  Cordeiro, A. M. *et al.* Rice phytochrome-interacting factor protein OsPIF14 represses

OsDREB1B gene expression through an extended N-box and interacts preferentially with

the active form of Phytochrome B. *Biochim Biophys Acta* **1859**, 393–404 (2016).

71.  Liang, C. *et al.* Transcriptomic, proteomic and metabolic changes in Arabidopsis thaliana

leaves after the onset of illumination. *BMC Plant Biology* **16**, 43 (2016).

72.  Ding, J. *et al.* Comparative Transcriptome Analysis of Gene Expression Patterns in Tomato

Under Dynamic Light Conditions. *Genes (Basel)* **10**, 662 (2019).

73.  Jin, X. *et al.* The Coordinated Upregulated Expression of Genes Involved in MEP,

Chlorophyll, Carotenoid and Tocopherol Pathways, Mirrored the Corresponding Metabolite

Contents in Rice Leaves during De-Etiolation. *Plants (Basel)* **10**, 1456 (2021).

74.  Bian, Z. *et al.* A Transcriptome Analysis Revealing the New Insight of Green Light on

Tomato Plant Growth and Drought Stress Tolerance. *Front Plant Sci* **12**, 649283 (2021).

75.  Ma, L. *et al.* Light Control of Arabidopsis Development Entails Coordinated Regulation of

Genome Expression and Cellular Pathways. *Plant Cell* **13**, 2589–2608 (2001).

76.  Jiao, Y., Lau, O. S. & Deng, X. W. Light-regulated transcriptional networks in higher

plants. *Nature Reviews Genetics* **8**, 217–230 (2007).

77.  de Wit, M., Galvão, V. C. & Fankhauser, C. Light-Mediated Hormonal Regulation of Plant

Growth and Development. *Annual Review of Plant Biology* **67**, 513–537 (2016).

78.  Hunter, W. N. The Non-mevalonate Pathway of Isoprenoid Precursor Biosynthesis. *J. Biol.*

*Chem.* **282**, 21573–21577 (2007).

79.  Gupta, P. *et al.* Cloning and characterization of 2-C-methyl-d-erythritol-4-phosphate

pathway genes for isoprenoid biosynthesis from Indian ginseng, Withania somnifera.

*Protoplasma* **250**, 285–295 (2013).

80.  Pulido, P., Perello, C. & Rodriguez-Concepcion, M. New Insights into Plant Isoprenoid

Metabolism. *Molecular Plant* **5**, 964–967 (2012).

81.  Nagata, N., Suzuki, M., Yoshida, S. & Muranaka, T. Mevalonic acid partially restores

chloroplast and etioplast development in Arabidopsis lacking the non-mevalonate pathway.

*Planta* **216**, 345–350 (2002).

82.    Rodríguez-Concepción, M. Early Steps in Isoprenoid Biosynthesis: Multilevel Regulation
720           of the Supply of Common Precursors in Plant Cells. *Phytochem Rev* **5**, 1–15 (2006).

721    83.    Ghassemian, M. *et al.* Integrative analysis of transcript and metabolite profiling data sets to
722           evaluate the regulation of biochemical pathways during photomorphogenesis. *Archives of*
723           *Biochemistry and Biophysics* **448**, 45–59 (2006).

724    84.    Vranová, E., Coman, D. & Gruissem, W. Structure and Dynamics of the Isoprenoid
725           Pathway Network. *Molecular Plant* **5**, 318–333 (2012).

726    85.    Filichkin, S. A. *et al.* Genome-wide mapping of alternative splicing in Arabidopsis thaliana.
727           *Genome Res* **20**, 45–58 (2010).

728    86.    Li, P. *et al.* The developmental dynamics of the maize leaf transcriptome. *Nature Genetics*
729           **42**, 1060–1067 (2010).

730    87.    Shen, Y. *et al.* Global Dissection of Alternative Splicing in Paleopolyploid Soybean. *The*
731           *Plant Cell* **26**, 996–1008 (2014).

732    88.    Campbell, M. A., Haas, B. J., Hamilton, J. P., Mount, S. M. & Buell, C. R. Comprehensive
733           analysis of alternative splicing in rice and comparative analyses with Arabidopsis. *BMC*
734           *Genomics* **7**, 327 (2006).

735    89.    Csorba, T., Questa, J. I., Sun, Q. & Dean, C. Antisense COOLAIR mediates the coordinated
736           switching of chromatin states at FLC during vernalization. *PNAS* **111**, 16160–16165
737           (2014).

738    90.    Filichkin, S. A. *et al.* Abiotic Stresses Modulate Landscape of Poplar Transcriptome via
739           Alternative Splicing, Differential Intron Retention, and Isoform Ratio Switching. *Front.*
740           *Plant Sci.* **9**, (2018).

741    91.    Deng, P., Liu, S., Nie, X., Weining, S. & Wu, L. Conservation analysis of long noncoding
742           RNAs in plants. *Sci. China Life Sci.* **61**, 190–198 (2018).

743    92.    Liu, J. *et al.* Genome-Wide Analysis Uncovers Regulation of Long Intergenic Noncoding
744           RNAs in Arabidopsis[C][W]. *Plant Cell* **24**, 4333–4345 (2012).

745    93.    Corona-Gomez, J. A., Garcia-Lopez, I. J., Stadler, P. F. & Fernandez-Valverde, S. L.
746           Splicing conservation signals in plant long noncoding RNAs. *bioRxiv* 588954 (2019)
747           doi:10.1101/588954.

748    94.    Komiya, R. *et al.* Rice germline-specific Argonaute MEL1 protein binds to phasiRNAs
749           generated from more than 700 lincRNAs. *The Plant Journal* **78**, 385–397 (2014).

95. Lavarenne, J. *et al.* Transcriptome profiling of laser-captured crown root primordia reveals new pathways activated during early stages of crown root formation in rice. *PLoS One* **15**, e0238736 (2020).

96. Yan, Y., Wang, H., Hamera, S., Chen, X. & Fang, R. miR444a has multiple functions in the rice nitrate-signaling pathway. *The Plant Journal* **78**, 44–55 (2014).

97. Chen, H. *et al.* OsMADS27 regulates the root development in a NO3−—Dependent manner and modulates the salt tolerance in rice (Oryza sativa L.). *Plant Science* **277**, 20–32 (2018).

98. Pachamuthu, K. *et al.* Nitrate-dependent regulation of miR444-OsMADS27 signalling cascade controls root development in rice. *Journal of Experimental Botany* **73**, 3511–3530 (2022).

99. Pathak, R. R. *et al.* Transcriptomic and network analyses reveal distinct nitrate responses in light and dark in rice leaves (Oryza sativa Indica var. Panvel1). *Sci Rep* **10**, 12228 (2020).

100. Yoneyama, T. & Suzuki, A. Light-Independent Nitrogen Assimilation in Plant Leaves: Nitrate Incorporation into Glutamine, Glutamate, Aspartate, and Asparagine Traced by 15N. *Plants (Basel)* **9**, 1303 (2020).

101. Hartmann, L. *et al.* Alternative Splicing Substantially Diversifies the Transcriptome during Early Photomorphogenesis and Correlates with the Energy Availability in Arabidopsis. *The Plant Cell* **28**, 2715–2734 (2016).

**Figure-1**

# Figure-2

| Pathway | Gene | Dark | Light | |
|---|---|---|---|---|
| | | IR | IR | ES |
| MVA | OS01G0110400 (AACT) | 2 | 1 | - |
| | OS08G0544900 (HMGS) | 1 | - | - |
| | OS10G0329300 (MK) | 1 | 1 | 1 |
| | OS03G0253100 (PMK) | 1 | - | - |
| MEP | OS05G0408900 (DXS) | - | 1 | - |
| | OS01G0106900 (DXR) | - | 2 | - |
| | OS01G0887100 (MECPS) | - | 1 | - |

■ Intron Retention (IR)