Transcriptional modulation during photomorphogenesis in rice seedlings

Parul Gupta, Pankaj Jaiswal[#]

Affiliations:

Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR, USA

 # Correspondence:

Prof. Pankaj Jaiswal

Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA

Email: Jaiswal.pankaj@oregonstate.edu

Keywords: Rice, photomorphogenesis, RNA-Seq, transcriptome, alternative splicing, lncRNA, MVA and MEP pathways, circadian clock

ABSTRACT

Light is one of the most important factors regulating plant gene expression patterns, metabolism, physiology, growth, and development. To explore how light may induce or alter transcript splicing, we conducted RNA-Seq-based transcriptome analyses by comparing the samples harvested as etiolated seedlings grown under continuous dark conditions vs. the light-treated green seedlings. We identified 14,766 differentially expressed genes, of which 4369 genes showed alternative splicing. We observed that genes mapped to the plastid-localized methyl-erythritol-phosphate (MEP) pathway were light-upregulated compared to the cytosolic mevalonate (MVA) pathway genes. Many of these genes also undergo splicing. These pathways provide crucial metabolite precursors for the biosynthesis of secondary metabolic compounds needed for chloroplast biogenesis, the establishment of successful photosynthetic apparatus, and photomorphogenesis. In the chromosome-wide survey of the light-induced transcriptome, we observed intron retention as the most predominant splicing event. In addition, we identified 1709 novel lncRNA transcripts in our transcriptome data.

INTRODUCTION

Light is an essential growth factor for sustaining autotrophic plant life. The quality, quantity, direction, and length of light exposure affect plant growth and development [1,2,3]. The natural light-dark cycle maintains the carbon and nitrogen metabolism of rice plants [4]. Light induces transcriptional reprogramming in various plant species [4–12] via an array of photoreceptors [13,14]. The photoreceptors phytochrome (red light) and cryptochrome (blue light) are involved in light signaling and photomorphogenesis [15,16]. Genes encoding phytochrome interacting factors (PIFs) promote skotomorphogenesis and development in the dark [17–19]. Light-mediated gene expression modulation is also triggered by translational enhancement of the preexisting mRNA pool instead of an enhanced transcription rate [20,21].

Intron splicing in multiexonic mRNA is a post-transcriptional regulatory process that often produces different mRNA isoforms transcribed from a single gene locus, thus contributing to proteome plasticity. Light-induced alternative splicing (AS) was observed in ~10% of the protein-coding genes of *Arabidopsis* and *Physcomitrella patens* [22,23]. A flash of light applied in the middle of the dark or nighttime is sufficient to induce splicing [24]. In contrast to animals, where the most common AS event is exon skipping (ES), intron retention (IR) is the most common AS event in rice [25], Arabidopsis [25,26], and Poplar [27]. It is now well established that transcriptome modulation via AS is vital for plant growth, development, and stress response [28–30].

Compared to the protein-coding genes, long non-coding RNAs (lncRNAs) are transcripts >200 bp in length that do not have protein-coding potential. lncRNAs are classified into categories: (i) sense lncRNAs, (ii) antisense lncRNAs, (iii) intergenic RNAs (lincRNAs), (iv) intronic RNAs, and (v) bidirectional lncRNAs [31]. In addition to the numerous studies coupling gene expression and AS studies on protein-coding genes, it is now known that lncRNAs also play a role in regulating gene expression through transcriptional, post-transcriptional, and chromatin remodeling mechanisms [31–34]. In rice, lncRNAs regulate biological processes, such as ovule development and female gametophyte abortion [35], sexual reproduction [36], and stress response [37,38]. However, the role of light in regulating lncRNAs is not well studied. *Arabidopsis* non-coding RNA *HID1* is a known positive regulator of photomorphogenesis in continuous red light [39].

Rice is a global staple crop and is a model for studying crop genomics. The transition from
skotomorphogenesis under dark conditions to photomorphogenesis under light exposure is
critical for seedling survival and requires precise control of gene expression by different
regulatory mechanisms. Results from our rice study show that exposure to light alters the
expression and splicing of a wide array of protein-coding genes but not so much for the non-
coding lncRNA genes.

MATERIALS & METHODS

Plant material, growth conditions, and treatment

Seeds of rice (*Oryza sativa spp. japonica* cv. Nipponbare) were grown and processed for the
experiment by following the growth conditions and sampling described previously [40]. Seeds were
sown in the dark and germinated on day 2. After sowing, these germinated seedlings grew in the
dark for 8 days (8DD). At the end of day 8, three biological replicates of the dark-grown
etiolated shoots were harvested. The remaining dark-grown seedlings were exposed to
continuous white light at 120 µmol/m2/sec (measured at the soil surface) for 48 hours or 2 days
(days 9 and 10 after sowing). The shoots of three biological replicates of light-treated green-
colored seedling samples (8DD-2LL) were harvested at the end of day 10. Harvested samples
were frozen using liquid nitrogen and stored at -80°C until further processing. Throughout this
report, 8DD-treated plants are called dark samples, and 8DD-2LL light-treated plants are called
light samples. Data analysis is described for light regulation.

Sample preparation and sequencing

Total RNA from frozen tissue samples was extracted using RNA Plant reagent (Invitrogen Inc.,
USA) and RNeasy kits (Qiagen Inc., USA) and treated with RNase-free DNase (Life
Technologies Inc., USA) according to the manufacturer's protocol. The total RNA quality and
concentration were determined using an ND-1000 spectrophotometer (Thermo Fisher Scientific
Inc., USA) and Bioanalyzer 2100 (Agilent Technologies Inc., USA). PolyA-enriched mRNA
libraries were prepared from three biological replicates of dark and light samples using the
TruSeqTM RNA Sample Preparation Kits (v2) and sequenced as 51-bp single-end reads using
the Illumina HiSeq 2000 instrument (Illumina Inc., USA) according to the manufacturer's
protocol at the Center for Genomic Research and Biocomputing, Oregon State University. The

strand-specific sequencing reads and metadata were deposited at EMBL-EBI ArrayExpress

(accession number E-MTAB-5689).

RNA-seq data analysis

The generation of FASTQ files from the RNA-Seq sequences was performed by CASAVA

software v1.8.2 (Illumina Inc.). Sequence reads were filtered and trimmed for low quality at a

score of 20 using Sickle v1.33 [41]. Clean, high-quality reads from each sample and replicates were

aligned to the *Oryza sativa japonica* cv Nipponbare reference genome (IRGSP-1.0.31) using

TopHat v2.1.1 [42]. Mapped reads were assembled using Cufflinks, and the reference-guided

assembled transcripts from each replicate were merged using Cuffmerge [43]. Assembled

transcripts were compared to the reference genome annotation using Cuffcompare. The RSEM

software package estimated normalized baseline expression from the aligned sequence reads [44].

For differential gene expression analysis, read count data obtained from RSEM were used in

EBSeq [45]. Differentially expressed (DE) genes were filtered based on the false discovery rate

corrected P value ≤0.05.

Functional annotation

We carried out the Gene Ontology (GO) enrichment analysis tool provided by the GO

consortium [46] to determine the biological roles played by the enriched gene set. Plant pathway

enrichment analysis was done by mapping the DE genes using the Plant Reactome analysis tool

[47,48,49] (http://plantreactome.gramene.org/PathwayBrowser/#TOOL=AT).

Alternative splicing analysis

Splicing events in the transcripts from the samples were identified by the SpliceGrapher v0.2.5

pipeline [50]. Sequence reads from each sample, and replicates were aligned to the reference rice

genome. Splice site-specific classifiers were built using build_classifiers.py script using

canonical (GT) and noncanonical (GC) donor sites and acceptor site (AG) for *Oryza sativa*

genome annotation version 31 (Oryza_sativa.IRGSP-1.0.31). Read alignments in SAM format

from each replicate were used as input for SpliceGrapher's sam_filter.py script to filter out false-

positive sites. SpliceGrapher Python scripts were used for the generation of depth files

(sam_to_depths.py), splice graph prediction (predict_graphs.py), generating statistics

(splicegraph_statistics.py) from a set of splice graphs, gene-by-gene summary

(genewise_statistics.py) of splicing events and splice graph visualization (plotter.py). The

Realignment pipeline was used to construct putative transcripts from unresolved exons with

sufficient coverage from the alignments [50]. In the following steps, IsoLasso [51], an extension of

the SpliceGrapher workflow, was used to predict novel splicing events.

Prediction of long non-coding RNA (lncRNA)

All transcripts annotated as intergenic transcripts, intron transcripts, antisense exon transcripts

overlapping the reference exons, and antisense intron transcripts overlapping the reference

introns were considered potential lncRNA candidates. Transcript sequences of length <=200

nucleotides were filtered out, and the gffread function of Cufflinks was used to extract fasta

sequences of potential lncRNA transcripts from the gtf file. CPC2 [52] was used to predict the

coding potential of transcripts. Predicted lncRNAs were scanned by InterProScan [53] to ensure the

absence of protein-coding domains. To identify novel lncRNAs, a BLASTn [54] search was

performed against a custom BLAST database generated using rice lncRNAs downloaded from

the CANTATAdb2.0 (http://cantata.amu.edu.pl), PNRD [55], GreeNC [56], and RiceLNCPedia [57]

databases. Secondary structures for the lncRNA molecules were predicted by the RNAfold

software [58,59].

RESULTS

Light-mediated differential gene expression during photomorphogenesis

To explore the transcriptome modulation in rice in response to light, we sequenced the strand-

specific poly-A enriched RNA fraction isolated from three biological replicates of rice plants

grown under dark and light exposure conditions (see methods). A total of ~38 million and ~42

million high-quality reads were generated from the dark and light-treated samples, respectively.

More than 92% of reads from each sample aligned to the rice reference genome (Supplementary

Table S1). We found 38,642 genes showed baseline normalized expression in the samples, of

which 33,943 genes expressed in the dark vs 35,772 genes that expressed under light,

respectively. Differential expression analysis identified 14,766 light-regulated genes (Figure 1,

Supplementary Figure S1, Supplementary Table S2).

Figure 1: Summary of differential gene expression and transcript splicing observed in rice

seedling shoots undergoing photomorphogenesis.

Gene function and pathway enrichment analyses

Gene Ontology-based functional annotation analysis of the light-regulated differentially

expressed gene set (Supplementary Figure S2) revealed enrichment for biological processes

(BP), such as chlorophyll and cell wall biosynthesis, besides photosystem light reaction

pathways and ribosome assembly. Dominant molecular functions (MF) were rRNA binding,

cytoskeletal motor activity, oxidoreductase and metallopeptidase activities, translation

elongation, and chaperone binding. As expected, chloroplast, cytoskeleton, ribosomes,

peroxisomes, and nucleus were the major cellular component sites of activity.

Pathway enrichment analysis using the Plant Reactome pathway analysis tool [60] mapped 475

light-upregulated genes to 243 pathways (Supplementary Table S3). A total of 164 pathways

overlapped between light-up and light-downregulated genes. Most pathways showed higher

mappings to light-upregulated genes, except hormone auxin and brassinosteroid signaling and

reproductive structure development (seed). Pathways unique to light-upregulated genes include

those for the biosynthesis of photosynthesis components chlorophyll, carotenoid, and

phylloquinone, hormones like gibberellin, auxin, abscisic acid, *etc*. The pathways with unique

mapping to light-downregulated genes include polar auxin transport, mevalonate (MVA)

pathway, circadian clock, salicylic acid metabolism and signaling, reproductive plant part

development, root-specific gene network of NAC10 transcription factor (Supplementary Figure

S3). To our surprise, we observed enrichment of the

Identifying light-regulated transcription factors

To identify the light-regulated transcription factors (TF), a list of rice TFs was downloaded from

the Plant Transcription Factor Database [61] and searched against the DE genes. We found 429

light-upregulated and 498 light-downregulated TFs (Supplementary Table S4). We found

WRKY, NAC, and orphans were the most abundantly expressed TFs in light, compared to many

light-downregulated bHLH, bZIP, and C3H gene family members (Table 1). Highly upregulated

(fold change ≥10) TFs belong to the MYB, AP2-EREBP, WRKY, Orphans, NAC, MADS, and

bHLH gene families; however, down upregulated TFs belong to the AP2-EREBP, C2C2-CO-

like, C3H, HB, and NAC gene families (Supplementary Table S4). To investigate whether TFs

targeted the MVA and MEP pathway genes, we surveyed the list of TFs and their potential

targets identified by the Plant Transcription Factor Database. We identified 23 TFs that

potentially bind to the promoter region of 6 MVA pathway genes. Two MVA pathway genes,

hydroxymethylglutaryl-CoA synthase (*HMGS*; Os08g0544900) and 3-hydroxy-3-methylglutaryl

coenzyme A reductase (*HMGR*; Os08g0512700), were targeted by bZIP and AP2-EREBP

factors, respectively, whereas the mevalonate 5-diphosphate decarboxylase (*MDD*;

Os02g0109100) gene was a target of by 13 AP2-EREBPs, one C3H protein, and one C2H2

protein (Supplementary Table S5). None of the TFs we found bind to the promoter of MEP

pathway genes.

Table 1: Light-regulated transcription factor gene families and their members.

Gene family	Upregulated	Downregulated	Gene family	Upregulated	Downregulated
ABI3VP1	6	4	GRF	1	2
Alfin-like	1	4	HB	21	24
AP2-EREBP	23	32	HSF	3	10
ARF	8	8	LIM	2	1
ARR-B	1	4	LOB	2	1
BBR-BPC	0	1	MADS	6	5
BES1	0	3	mTERF	9	4
bHLH	19	27	MYB	28	14
BSD	1	6	MYB-related	16	25
bZIP	10	27	NAC	21	19
C2C2-CO-like	4	7	OFP	2	1
C2C2-Dof	5	4	Orphans	20	10
C2C2-GATA	7	4	PBF-2-like	1	0
C2C2-YABBY	3	1	PLATZ	1	4
C2H2	14	19	RWP-RK	4	2
C3H	16	24	S1Fa-like	1	1
CAMTA	0	3	SBP	3	5
CCAAT	9	11	Sigma 70-like	4	0
CPP	1	4	SRS	1	0
CSD	0	1	TAZ	1	1
DBP	2	3	TCP	4	6
E2F/DP	2	1	Tify	6	2
EIL	1	1	Trihelix	3	9
FAR1	5	10	TUB	2	5
FHA	7	6	ULT	1	0
G2-like	9	5	VOZ	0	2
GeBP	0	3	WRKY	22	10
GRAS	8	7	zf-HD	5	3
Other transcriptional regulators					
Gene family	**Upregulated**	**Downregulated**	**Gene family**	**Upregulated**	**Downregulated**
ARID	2	1	PHD	12	22
AUX/IAA	3	19	Pseudo ARR-B	2	1
Coactivator p15	0	1	RB	0	1
DDT	3	0	Rcd1-like	0	2
GNAT	12	8	SET	11	6
HMG	4	5	SNF2	11	9
Jumonji	4	5	SOH1	0	2
LUG	0	2	SWI/SNF-BAF60b	4	4
MBF1	0	1	SWI/SNF-SWI3	2	1
MED6	0	1	TRAF	7	11

Transcript splicing during photomorphogenesis

To study how light exposure modulates transcript splicing (AS) events in the rice seedling

shoots, we analyzed the transcriptome data by using SpliceGrapher [50]. We created highly

accurate splice site classifiers for rice with canonical (GT-AG) and semi-canonical (GC-AG)

splice sites to filter the splice junctions in the sequence reads aligned to the reference genome.

After removing the false positive splice sites, we predicted chromosome-wise splice graphs from

the two transcriptome sample datasets. In the dark-treated dataset, 63.9% were true positive

junctions, with 5.5% novel splice sites. In the light-treated dataset, 64.3% of splice junctions

were true positives, of which 5.8% were predicted as novel sites. We observed 6214 spliced

genes with 9685 splicing events in the dark samples compared to the 6809 spliced genes with

10432 splicing events in the light samples (Figure 1, Table 2). The highest number of spliced

genes and events were on chromosome 1, and the lowest was on chromosome 10 (Table 2).

Intron retention (IR) was the most prevalent type of splicing event in both samples, followed by

exon skipping (ES). Alternative 3' splicing (Alt.3') was the least common event. To resolve

ambiguous combinations of donor and acceptor splice sites, we realigned the sequenced reads

from the two samples to the putative transcripts and resolved the novel exons to generate splice

graphs. Light-induced splicing in 2162 unique genes compared to 1567 unique genes that

undergo splicing in the dark. Of the 4647 spliced genes shared between the two samples, 165

genes displayed differential splicing events.

Table 2: Summary of spliced genes and events under dark and light conditions detected by the SpliceGrapher and the IsoLasso workflows. IR: Intron retention; ES: Exon Skipping; Alt5': alternative 5′ splice site; Alt3': alternative 3′ splice site. Color scales: Red (Large counts) to Blue (Lower counts).

Chromosome #	Treatment	SpliceGrapher Prediction					IsoLasso				
		Genes	IR	ES	Alt5'	Alt3'	Genes	IR	ES	Alt5'	Alt3'
1	Dark	924	897	214	163	97	939	925	214	181	99
	Light	1028	994	264	184	109	1048	1021	264	191	112
2	Dark	747	728	224	167	109	771	781	227	189	117
	Light	802	777	219	153	99	822	812	222	165	103
3	Dark	825	797	216	136	94	858	868	216	162	120
	Light	945	893	219	143	98	960	920	219	153	118
4	Dark	565	551	182	112	64	581	576	183	117	67
	Light	596	572	193	104	61	611	592	194	109	66
5	Dark	519	503	139	139	70	538	534	140	114	75
	Light	552	537	144	103	80	562	550	144	109	92
6	Dark	509	459	186	128	62	527	493	186	135	64
	Light	531	501	169	103	63	554	535	169	109	67
7	Dark	466	432	166	77	48	482	468	166	85	51
	Light	481	473	129	72	52	498	498	129	82	56
8	Dark	400	390	97	63	62	415	416	98	68	64
	Light	463	433	134	62	84	475	457	134	67	87
9	Dark	341	342	79	53	32	349	359	79	64	37
	Light	375	361	118	49	38	382	377	118	53	39
10	Dark	290	275	94	43	29	295	288	94	49	36
	Light	308	295	102	61	32	314	307	102	65	35
11	Dark	304	266	113	49	35	313	278	113	53	36
	Light	363	306	150	51	35	374	325	150	57	37
12	Dark	324	310	117	69	45	329	315	117	74	46
	Light	365	347	150	63	53	372	359	155	68	61

In our differentially expressed and spliced genes (DES), we found that approximately 30% (4369) differentially expressed (DE) genes also undergo splicing (AS). Approximately 10% (2440) of the non-DE genes undergo splicing (Figure 1). Light-upregulated DES genes show enrichment for various molecular functions like oxidoreductase, hydrolase, isomerase activity, and RNA binding, which play roles in photosynthesis, transmembrane transport, small molecule metabolic process, and located in the cellular components, chloroplast, and membrane-bound organelles. In contrast, the light-downregulated DES genes enriched for RNA processing, response to stress, protein binding, and macromolecule metabolic process, and localized in cellular components cytoplasm and nucleus (Supplementary Figure S4).

We also surveyed the DES aspects of the 57 known spliced genes in rice [25,62] and observed splicing in 41 genes (Table 3). Of these, seven spliced under dark conditions, three under light conditions, and 31 genes spliced under both dark and light conditions. Of the known genes, we also observed that 18 genes were light-upregulated and 24 downregulated. Only 32 genes showed a DES profile. Three light-upregulated genes (Os02g0130600, Os05g0348100, Os12g0567300) show complete splicing in light compared to intron retention in the dark. In contrast, the two light-downregulated genes (Os02g0666200 and Os04g0656100) show intron retention.

235 **Table 3:** Expression and splicing profile of genes in our data that are known to undergo splicing
236 from previous reports

Known spliced genes	Spliced in	Regulation in light	Gene description
Os01g0619000	Dark, Light	Down	RNA recognition motif domain-containing protein
Os01g0649900	Dark, Light	-	GDSL esterase/lipase protein 20
Os01g0764000	Dark, Light	Up	PHI glutathione s-transferase 2
Os01g0834400	Dark	-	HAP3A subunit of CCAAT-box binding complex
Os02g0122800	Dark, Light	Down	Similar to Arginine/serine-rich splicing factor
Os02g0130600	Dark	Up	Conserved hypothetical protein
Os02g0161900	Dark, Light	Down	Rice ubiquitin2
Os02g0197900	Dark, Light	Down	OsSTA53
Os02g0274900	Dark, Light	Up	Similar to metabolite transport protein CSBC
Os02g0577100	Dark, Light	Down	Zinc finger, RING/PHD-type domain containing protein
Os02g0666200	Light	Down	Aquaporin
Os02g0834000	Dark, Light	Up	Rac-like GTP-binding protein 5
Os02g0291000	-	Down	Calcineurin B-like protein 7
Os02g0291400	-	-	Calcineurin B-like protein 8
Os02g0823100	-	Up	Plasma membrane intrinsic protein 1;3
Os03g0265600	Dark, Light	Down	Transformer-2-like protein
Os03g0314100	Dark, Light	Down	DEAD-like helicase
Os03g0395900	Dark, Light	Down	Splicing factor
Os03g0670700	Dark, Light	Down	Glycine-rich rna-binding protein 3
Os03g0698500	Dark, Light	Down	Similar to Yippee-like protein 3
Os03g0745000	Dark, Light	-	Heat stress transcription factor A-2a
Os03g0717600	-	Down	Zinc finger, C2H2-type
Os04g0115400	Dark, Light	Down	D111/G-patch domain containing protein
Os04g0649100	Dark, Light	-	Shattering abortion 1
Os04g0656100	Light	Down	H^+-ATPase
Os04g0665800	Dark, Light	Up	Similar to H1005F08.12 protein
Os04g0402300	-	-	Cysteine-type peptidase
Os04g0479200	-	Up	Similar to NAD-dependent isocitrate dehydrogenase c;1

Os05g0348100	Dark	Up	Similar to CRR23 (chlororespiratory reduction 23)
Os05g0463800	Dark, Light	Down	HAP3C subunit of CCAAT-box binding complex
Os05g0554400	Dark, Light	Down	Phosphatidyl serine synthase family protein
Os05g0574700	Dark, Light	-	Similar to cDNA clone:002-182-C01
Os05g0534400	-	Up	Calcineurin B-like protein 4
Os05g0548900	-	Up	Phosphoethanolamine N-Methyltransferase 2
Os06g0172800	Light	-	Similar to alkaline alpha galactosidase 2
Os06g0651600	Dark, Light	Down	Protein phosphatase 2C58
Os06g0727200	Dark	Down	Catalase B
Os06g0128500	-	Up	Ribosomal protein L47, mitochondrial family protein
Os06g0133000	-	-	Glutinous Endosperm, waxy
Os06g0506600	-	-	Ubiquitin-conjugating enzyme 17
Os07g0490400	Dark, Light	Up	FK506 binding protein 20-2
Os07g0574800	-	Up	Tubulin alpha-1 chain
Os07g0613300	-	-	Similar to PAUSED
Os08g0191600	Dark, Light	Up	Autophagy associated gene 8C
Os08g0530400	Dark, Light	Up	Moco containing protein, Similar to sulfite oxidase
Os08g0436200	-	-	Zinc finger, RING/PHD-type domain containing protein
Os10g0115600	Dark, Light	Down	U1 snRNP 70K
Os10g0535800	Dark, Light	-	Cys-rich domain containing protein
Os10g0564900	Dark	Down	Similar to protein kinase CK2 regulatory subunit CK2B2
Os10g0567400	Dark, Light	Up	Chlorophyll a oxygenase 1
Os10g0577900	Dark, Light	-	Glycerol-3-phosphate acyltransferase
Os10g0411500	-	Down	Q calmodulin-binding region domain containing protein
Os11g0157100	Dark, Light	Down	Cyclin-T1-4
Os11g0700500	Dark	-	MybAS1
Os11g0600700	-	Up	Zinc finger, RING domain protein
Os12g0567300	Dark	Up	MybAS2, R2R3-Myb
Os12g0632000	Dark, Light	Down	Glycine-rich Protein GRP162

Light-mediated shift in gene expression and splicing events

Circadian clock pathway genes

The circadian clock is entrained by light, and the diurnal photoperiod regulated clock genes [63,64]. We investigated the light-regulated DES profile of clock genes. Most of the circadian clock genes spliced under either or both conditions. Four genes, *Casein kinase alpha subunit* (*CK2α-1/Hd6*, Os03g0762000), *Casein kinase beta subunit* (*CK2β-2*, Os10g0564900), *Phytochrome B* (*PhyB*, Os03g0309200;), and *Timing of CAB Expression 1* (*TOC1*, Os02g0618200), showed differential splicing (only reference splicing events or complete splicing and no novel isoforms) upon light treatment compared to the presence of unspliced intron in the absence of light (Supplementary Figure S5). In contrast, only one gene, *Phytochrome Interacting Factor 3* (*PIF3*, Os01g0286100), showed intron retention under light. All five genes were upregulated in light (Supplementary Figure S5). Seven genes, *Casein kinase alpha subunit* (*CK2α-2*, Os03g0763000), *Casein kinase beta subunit* (*CK2β-1*, Os07g0495100), blue light receptor *Cryptochrome 2* (*CRY2*, Os02g0625000), *Gigantea* (*GI*, Os01g0182600), *Pseudo-Response Regulator 37* or *Heading date 2* (*PRR37*, *Hd2*, Os07g0695100), and *Pseudo-Response Regulator 73* (*PRR73*, Os03g0284100), showed differential splicing under both conditions (Supplementary Figure S5, S6). Exon skipping and alternative 5' (Alt.5') splicing events in the *CK2α-2* (Os03g0763000) gene were induced in light. Both *CK2β-1* (Os07g0495100) and *GI* (Os01g0182600) display incomplete splicing of intron (IR) in the dark. *PRR37* showed three types of splicing events (Alt.3', ES, and IR) under light, whereas the second IR event observed toward the 3' end under dark was spliced in light. Conversely, *PRR73* showed three types of events (Alt.3', ES, and IR) under dark conditions, whereas under light conditions, two new Alt.5' events were identified. Expression of both PRR genes (*PRR37* and *PRR73*) was light downregulated (Supplementary Figure S5). Light-upregulated *PRR95* shows an additional unspliced intron in light.

MVA and MEP pathway genes

Light regulates the biosynthesis and accumulation of secondary plant products like isoprenoid-derived metabolites required for plant growth and development. Isoprenoid precursors are synthesized via cytosolic MVA and plastid-localized MEP pathways, and their fluxes are

regulated by light [65,66], and their products are essential for the successful development and
function of chloroplast. We observed that MEP pathway genes were light-upregulated compared
to the downregulated MVA pathway genes (Figure 2). Light positively regulates MEP pathway
activity, whereas MVA pathway activity appears negatively regulated by light [67].

We investigated the light-induced differential splicing events in the MEP and MVA pathway
genes. Four MVA pathway genes (*acetyl-CoA acetyltransferase: AACT, hydroxymethyl glutaryl*
CoA synthase: HMGS, mevalonate kinase: MK, phosphomevalonate kinase: PMK) and 3 MEP
pathway genes (*1-deoxy-D-xylulose-5-phosphate synthase: DXS, 1-deoxy-D-xylulose-5-*
phosphate reductoisomerase: DXR, 2-C-methyl-D-erythritol2,4-cyclodiphosphate synthase:
MECPS) showed intron retention events in dark (Figure 2). Gene *MK* showed a novel light-
induced exon-skipping event.

Figure 2: Gene expression and splicing of cytosolic mevalonate (MVA) and plastid-localized
methyl erythritol 4-phosphate (MEP) pathway genes. *AACT*, acetyl-CoA acetyltransferase;
HMGS, hydroxymethyl glutaryl CoA synthase; *HMGR*, 3-hydroxy-3-methylglutaryl-coenzyme
A reductase; *MK*, mevalonate kinase; *PMK*, phosphomevalonate kinase; *MDD*, mevalonate
diphosphosphate decarboxylase; *DXS*, 1-deoxy-D-xylulose-5-phosphate synthase; *DXR*, 1-
deoxy-D-xylulose-5-phosphate reductoisomerase; *CDPMES*, 2-C-methyl-D-erythritol4-
phosphate cytidylyl transferase; *CDPMEK*, 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase;
MECPS, 2-C-methyl-D-erythritol2,4-cyclodiphosphate synthase; *HDS*, 4-hydroxy-3-methylbut-
2-enyldiphosphate synthase; *HDR*, 4-hydroxy-3-methylbut-2-enyldiphosphate reductase. Blue:
intron retention, Green: exon skipping, Orange: Alt3' splicing, Purple: Alt5' splicing.

lncRNA discovery and splicing during photomorphogenesis

lncRNAs are known to regulate diverse biological processes in plants, including
photomorphogenesis [68], and as described above, are categorized into four major types based on
their location in the gene space and on the DNA strand. We used our transcriptome data to
characterize light-induced rice lncRNAs (Supplementary Figure S7). We identified 1485 and
1407 lncRNA transcripts in the dark and light samples, of which 309 were common. Many
lncRNAs from intergenic and intronic regions were identified (Supplementary Figure S7). To
discover novel lncRNAs, the transcripts from our datasets were queried against the custom
BLAST database of rice lncRNAs. About 44% of the total lncRNA transcripts (2583) found a
match in the BLAST database (Supplementary Table S6). 173 lncRNAs spliced under dark and
142 spliced under light conditions, most located in the intergenic regions (Supplementary Figure
S7). Interestingly, all the spliced lncRNA genes from both datasets were from Chr1, Chr10,
Chr11, and Chr12. The maximum number of spliced lncRNA genes was from Chr1, followed by
Chr11, 12, and 10. Among all the splicing events, IR was predominant under both conditions.

We found lncRNA transcripts from light (TCONS_00012152) and dark (TCONS_00011794)
datasets that were present on the antisense strand overlapping an exon region of the *MADS27*
(Os02g0579600) gene. The transcript from the light-treated sample appears longer and showed
two alternative 3' splice sites and one exon skipping event. The TCONS_00011794 may
compete with MADS27 to form a complex with miRNA osa-miR444a known to play multiple
roles in the nitrate-dependent development pathway [69]. Among the list of lncRNAs present on
the antisense strand overlapping annotated genes, we identified a light-induced lncRNA
(TCONS_00001515) overlapping *ERF99* transcription factor (Os01g0868000) exon that may
play a role in its silencing. *ERF99* plays a central role in mediating abiotic stress responses [70].
Similarly, lncRNA transcript (TCONS_00015825) is present exclusively in light is transcribed
from the intronic region of the antisense strand of a B-type response regulator gene
(Os03g0224200), known for its involvement in cytokinin signaling, meristem maintenance, and
stress response [71,72]. The lncRNA transcript (TCONS_00026617) present exclusively in the dark
overlaps the intronic region of a phytochrome-interacting factor gene *PIF14* (Os07g0143200).

319 *PIF14* is known to bind the active form of phytochrome B and plays a crucial role in cross-talk

320 between light and stress signaling [73].

Discussion

Light is known to induce changes in the transcriptome, metabolome, and proteome of the plants

[74], which not only regulates the development, function, and physiology of the chloroplast but

also provides signals for modulating the plant's morphological, developmental, and

physiological adaptions in response to growth environment [8,40,75–77]. Though plants experience

constantly changing light conditions under the natural environment daily during their lifetime,

early development from germination to the seedling stage and acquiring the full autotrophic

capability is very much programmed by exposure to light. Therefore, we investigated

transcriptome modulation in rice seedlings during photomorphogenesis. Compared to earlier

studies in rice and *Arabidopsis*, where ~20% of the genes were reported differentially expressed

in dark-grown etiolated seedlings compared to light-exposed green seedlings [9,78,79], we observed

that the transition from skotomorphogenesis to photomorphogenesis alters differential expression

of ~38% of the rice genes.

Functional annotation of the differentially expressed genes enriched for roles in secondary

metabolism, chloroplast-related biosynthetic pathways, hormone biosynthesis and signaling,

amino acid biosynthesis, fatty acid metabolism, *etc*. (Supplementary Figure S3, Supplementary

Table S3). This result was expected based on the earlier reports on light regulation of such events

[24,80].

The plastids develop into chloroplasts by following an essential step of developing the thylakoid

membrane system and recruiting and assembling components of light and dark reactions to

establish a functional photosynthetic process. Many terpenoid compounds are essential to plants'

light-harvesting function and protect against damage from reactive oxygen species (ROS). The

basic isoprenoid units for terpenoid biosynthesis, such as tocopherols, plastoquinones,

carotenoids, chlorophylls, and precursors of the growth hormones gibberellins and abscisic acid,

are synthesized by the plastid-localized MEP pathway [81,82]. At the same time, isoprenoid units

synthesized by the MVA pathway contribute to the synthesis of triterpenes, phytosterols, and

phytohormones [83]. MEP and MVA pathways complement and contribute to the biosynthesis of

chlorophylls and carotenoids required for plastid development [84]. MEP pathway genes were

light-upregulated in our transcriptome data compared to the light-downregulated MVA pathway

genes (Figure 2). These results were consistent with earlier reports [85,86] and confirm that light

acts as a critical regulator in modulating the availability of isoprenoid precursors during

photomorphogenesis [87]. Arabidopsis circadian clock genes (*LHY*, *PRR9*, *CCA1*, *TOC1*) regulate

MVA and MEP pathways [87]. *AtTOC1* is also known to regulate the G1-to-S phase transition in

the mitotic cell cycle in early leaf development [88]. In our dataset, we did not observe any

significant change in the expression of *LHY/CCA1*(Os08g0157600), *PRR95* (Os09g0532400, a

homolog of *AtPRR9*), and *TOC1* (Os02g0618200). However, all three genes spliced differently.

Under the light, *TOC1* showed no splicing but retained two introns in the dark. We observed one

exon-skipping event in light of the *LHY/CCA1* gene, whereas *PRR95* showed one additional

intron retention (Supplementary Figure S5). Similar to *Arabidopsis* [88], the functional enrichment

analysis also identified the *TOC1*-regulated DNA biosynthesis pathway of the G1-to-S transition

of mitotic cell cycle (Figure 3) along with a greater coverage of reaction events in the light-

upregulated set. To our knowledge this is the first report of the light regulation of mitotic cell-

cycle pathway in rice.

Figure 3: Plant Reactome pathway enrichment showing light-regulated genes and reaction event coverage for the DNA biosynthesis pathway of the G1-to-S phase mitotic cell cycle.

Previous transcriptome studies showed that ~42% of *Arabidopsis*, ~63% of *soybean*, and ~56% of maize genes undergo alternative splicing events [89–91]. Studies on rice by Wang and Brendel [25] reported IR for 69% of 6568 AS genes, and Campbell *et al.* [92] reported 44.7% of 8772 AS genes. In our dataset, 4369 genes undergo differential expression and alternative splicing, called DES genes (Figure 1A). About 80% (5395) of the spliced genes (6809) showed IR events, suggesting that IR is the major splicing event. The more significant number of splicing events in the light-treated samples suggests that light-mediated gene expression and post-transcriptional mRNA splicing play an important role in photomorphogenesis. The splicing events in rice circadian clock genes suggest that IR regulates the expression of PRR genes (Supplementary Figure S5). Many of the light-regulated genes of the MEP and MVA pathways show IR events, suggesting that light plays a positive role in completing the splicing and provides a hypothesis that exposure to light is an adaptive feature and even if the gene is expressed in dark, it stays unspliced until it encounters light. Our data showed the presence of all three states, partially spliced in the dark

(*GI*, Os01g0182600), fully spliced in light (*GI*, Os01g0182600), and both states in light (*CK2β-*

1, Os07g0495100) (Supplementary Figure S6).

lncRNAs play regulatory functions in essential biological processes such as vernalization,

photomorphogenesis, and stress regulation [39,93,94] and display splicing [95]. 6480 long intergenic

non-coding RNAs (lincRNAs) were identified from 200 *Arabidopsis* transcriptome datasets [96];

however, we identified 827 lincRNAs in the dark and 727 lincRNAs in the light datasets.

Corona-Gomez and coworkers [97] characterized the splicing conservation of lncRNAs in

Brassicaceae, which revealed that ~18% of lincRNAs display splicing conservation in at least

one exon. In our dataset, ~10% of lncRNA transcripts undergo splicing, and the majority were of

long intergenic non-coding RNA type (Supplementary Figure S7). Komiya and colleagues

reported [98] that phased small interfering RNAs (phasiRNAs) are generated from over 700

lincRNAs, and these phasiRNAs bind to rice argonaute protein MEL1. MEL1 has a specific

function in developing pre-meiotic germ cells and the progression of meiosis. In our dataset,

many lincRNAs transcribed from MEL1-phasiRNA clusters appeared.

Identification of light-induced lncRNAs antisense to *ERF99*, B-type response regulator genic

region may play a role in transcriptional regulation. *ERF99* is known to modulate root

architecture and downregulated in crown root primordia [99]. *MADS27* gene targeted by lncRNAs

present in both dark and light conditions. However, the dark lncRNA (TCONS_00011794) is

longer and, based on its predicted RNA fold structure and lower free energy, is likely more stable

than its transcript TCONS_00012152 in light. For the MADS27 gene, we also observed an

additional IR event in light. MADS27-miR444 complex is known to play a role in plant

development in a nitrate-dependent manner [100–102]. Therefore, we investigated the nitrogen

assimilation pathway genes. Nitrogen assimilation is necessary for sustaining plant growth and

development. Various nitrogen assimilatory enzymes are known to show isoform and cellular

component-specific responses under light and dark conditions [103,104]. We observed light

upregulation for the nitrate transporter, nitrate reductase, and nitrite reductase genes. We also

observed differential splicing patterns for these genes (Supplementary Figure S8). We also found

light regulation of the recently reported rice abiotic and biotic stress response genes [105,106].

Conclusion

This study suggests that light is a significant regulatory factor controlling genome-wide gene expression through an alternative splicing mechanism in rice. All spliced genes did not necessarily produce novel isoforms, which indicates the coupling of AS and nonsense-mediated decay (NMD). NMD prevents the translation of mutant mRNAs harboring potential premature stop codons by targeting them for degradation. In *Arabidopsis*, 77.2% of light-regulated AS events exhibit NMD features within a splicing isoform [107]. We conclude that light induces a significant number of splicing events in rice protein-coding and non-coding lncRNA genes. This photomorphogenesis transcriptome study is a valuable resource for lncRNA research in rice and provides insights into the portion of the genome regulated at the level of alternative splicing in response to light. We expect the condition-dependent novel splice events discovered in this study will help improve the annotation of the rice reference genome and pan-genomes [49,108]

Conflict of interest

The authors declare no conflict of interest.

Availability of data

The sequence reads and metadata were deposited at EMBL-EBI ArrayExpress (accession number E-MTAB-5689).

Figure legends

Figure 1: Summary of differential gene expression and transcript splicing observed in rice

seedling shoots undergoing photomorphogenesis.

Figure 2: Gene expression and splicing of cytosolic mevalonate (MVA) and plastid-localized

methyl erythritol 4-phosphate (MEP) pathway genes. *AACT*, acetyl-CoA acetyltransferase;

HMGS, hydroxymethyl glutaryl CoA synthase; *HMGR*, 3-hydroxy-3-methylglutaryl-coenzyme

A reductase; *MK*, mevalonate kinase; *PMK*, phosphomevalonate kinase; *MDD*, mevalonate

diphosphosphate decarboxylase; *DXS*, 1-deoxy-D-xylulose-5-phosphate synthase; *DXR*, 1-

deoxy-D-xylulose-5-phosphate reductoisomerase; *CDPMES*, 2-C-methyl-D-erythritol4-

phosphate cytidylyl transferase; *CDPMEK*, 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase;

MECPS, 2-C-methyl-D-erythritol2,4-cyclodiphosphate synthase; *HDS*, 4-hydroxy-3-methylbut-

2-enyldiphosphate synthase; *HDR*, 4-hydroxy-3-methylbut-2-enyldiphosphate reductase. Blue:

intron retention, Green: exon skipping, Orange: Alt3' splicing, Purple: Alt5' splicing

Figure 3: Plant Reactome pathway enrichment showing light regulated genes and reaction event

coverage for the DNA biosynthesis pathway of the G1-to-S phase mitotic cell cycle.

 Table legends

 Table 1: Light-regulated transcription factor gene families and their members.

 Table 2: Summary of spliced genes and events under dark and light conditions detected by the

 SpliceGrapher and the IsoLasso workflows. IR: Intron retention; ES: Exon Skipping; Alt5':

 alternative 5′ splice site; Alt3': alternative 3′ splice site. Color scales: Red (Large counts) to Blue

 (Lower counts).

 Table 3: Expression and splicing profile of genes in our data known to undergo splicing from

 previous reports

Supplementary material

Supplementary Figure S1:

Expression pattern of differentially expressed genes (14,766) with target FDR controlled at 5%

Supplementary Figure S2:

Gene Ontology enrichment analysis of differentially expressed genes.

Supplementary Figure S3:

Pathway enrichment analysis using the Plant Reactome. **(A)** Plant Reactome pathway enrichment

analysis plots, **(B)** Unique and shared pathways enriched for the light upregulated and

downregulated gene sets; **(C)** Counts of genes mapped to some of the common Plant Reactome

pathways. Light upregulated (green) and downregulated (grey).

Supplementary Figure S4:

Bar plots of most significantly enriched GO terms for DES genes with -log10 transformed FDR

values. Light-upregulated (green), light-downregulated (grey).

Supplementary Figure S5:

The summary of transcript splicing events and expression pattern of circadian clock genes during

morphogenesis.

Supplementary Figure S6:

Transcript graphs showing splicing of *Casein kinase beta subunit:CK2β-1* (Os07g0495100),

Gigantea:GI (Os01g0182600) genes during photomorphogenesis.

Supplementary Figure S7:

Classification of lncRNAs identified in dark and light.

Supplementary Figure S8:

An overviw of splicing and expression pattern of nitrogen assimilation cycle genes during

photomorphogenesis. NRT: Nitrate transporter; NR: Nitrate reductase; NiR: Nitrate reductase.

Supplementary Table S1:

Summary of the RNA-Seq read mapping to the reference rice genome.

Supplementary Table S2:

Differentially expressed genes

Supplementary Table S3:

Plant Reactome Pathway enrichment analysis for light-regulated genes

Supplementary Table S4:

Transcription factor genes regulated in light

Supplementary Table S5:

Transcription factors regulating MVA pathway genes

Supplementary Table S6:

lncRNA annotation using BLAST

Reference

1. Kami, C., Lorrain, S., Hornitschek, P. & Fankhauser, C. Chapter Two - Light-Regulated Plant Growth

and Development. in *Current Topics in Developmental Biology* (ed. Timmermans, M. C. P.) vol. 91

29–66 (Academic Press, 2010).

2. Poolman, M. G., Kundu, S., Shaw, R. & Fell, D. A. Responses to Light Intensity in a Genome-Scale

Model of Rice Metabolism. *Plant Physiology* **162**, 1060–1072 (2013).

3. Naithani, S., Nonogaki, H. & Jaiswal, P. Exploring Crossroads Between Seed Development and

Stress Response. in *Mechanism of Plant Hormone Signaling under Stress* 415–454 (John Wiley &

Sons, Ltd, 2017). doi:10.1002/9781118889022.ch32.

4. Li, H. *et al.* A Natural Light/Dark Cycle Regulation of Carbon-Nitrogen Metabolism and Gene

Expression in Rice Shoots. *Front. Plant Sci.* **7**, (2016).

5. Bai, S. *et al.* BBX16, a B-box protein, positively regulates light-induced anthocyanin accumulation

by activating MYB10 in red pear. *Plant Biotechnology Journal* **0**,.

6. Tobin, E. M. & Silverthorne, J. Light Regulation of Gene Expression in Higher Plants. *Annual Review*

of Plant Physiology **36**, 569–593 (1985).

7. Ma, L. *et al.* Light Control of Arabidopsis Development Entails Coordinated Regulation of Genome

Expression and Cellular Pathways. *Plant Cell* **13**, 2589–2608 (2001).

8. Casal, J. J. & Yanovsky, M. J. Regulation of gene expression by light. *Int. J. Dev. Biol.* **49**, 501–511

(2004).

9. Jiao, Y., Ma, L., Strickland, E. & Deng, X. W. Conservation and Divergence of Light-Regulated

Genome Expression Patterns during Seedling Development in Rice and Arabidopsis. *The Plant Cell*

17, 3239–3256 (2005).

10. Majláth, I. *et al.* Effect of light on the gene expression and hormonal status of winter and spring

wheat plants during cold hardening. *Physiologia Plantarum* **145**, 296–314 (2012).

11.   Petrillo, E., Herz, M. A. G., Barta, A., Kalyna, M. & Kornblihtt, A. R. Let there be light: Regulation of

536   gene expression in plants. *RNA Biology* **11**, 1215–1220 (2014).

537   12.   Godoy Herz, M. A. *et al.* Light Regulates Plant Alternative Splicing through the Control of

538   Transcriptional Elongation. *Molecular Cell* (2019) doi:10.1016/j.molcel.2018.12.005.

539   13.   Quail, P. H. Phytochrome photosensory signalling networks. *Nature Reviews Molecular Cell Biology*

540   **3**, 85–93 (2002).

541   14.   Lin, C. & Shalitin, D. Cryptochrome Structure and Signal Transduction. *Annual Review of Plant*

542   *Biology* **54**, 469–496 (2003).

543   15.   Huang, X., Ouyang, X. & Deng, X. W. Beyond repression of photomorphogenesis: role switching of

544   COP/DET/FUS in light signaling. *Current Opinion in Plant Biology* **21**, 96–103 (2014).

545   16.   Sheerin, D. J. *et al.* Light-Activated Phytochrome A and B Interact with Members of the SPA Family

546   to Promote Photomorphogenesis in Arabidopsis by Reorganizing the COP1/SPA Complex. *Plant Cell*

547   **27**, 189–201 (2015).

548   17.   Leivar, P. *et al.* Multiple phytochrome-interacting bHLH transcription factors repress premature

549   photomorphogenesis during early seedling development in darkness. *Curr Biol* **18**, 1815–1823

550   (2008).

551   18.   Shin, J. *et al.* Phytochromes promote seedling light responses by inhibiting four negatively-acting

552   phytochrome-interacting factors. *Proc Natl Acad Sci U S A* **106**, 7660–7665 (2009).

553   19.   Xie, C., Zhang, G., An, L., Chen, X. & Fang, R. Phytochrome-interacting factor-like protein OsPIL15

554   integrates light and gravitropism to regulate tiller angle in rice. *Planta* (2019) doi:10.1007/s00425-

555   019-03149-8.

556   20.   Liu, M.-J., Wu, S.-H., Chen, H.-M. & Wu, S.-H. Widespread translational control contributes to the

557   regulation of Arabidopsis photomorphogenesis. *Molecular Systems Biology* **8**, 566 (2012).

558   21.   Kianianmomeni, A. More light behind gene expression. *Trends in Plant Science* **19**, 488–490 (2014).

559   22.   Wu, H.-P. *et al.* Genome-wide analysis of light-regulated alternative splicing mediated by

560   photoreceptors in Physcomitrella patens. *Genome Biology* **15**, R10 (2014).

561   23.   Shikata, H. *et al.* Phytochrome controls alternative splicing to mediate light responses in

562   Arabidopsis. *PNAS* **111**, 18781–18786 (2014).

563   24.   Mancini, E. *et al.* Acute Effects of Light on Alternative Splicing in Light-Grown Plants. *Photochem

564   Photobiol* **92**, 126–133 (2016).

565   25.   Wang, B.-B. & Brendel, V. Genomewide comparative analysis of alternative splicing in plants. *Proc

566   Natl Acad Sci U S A* **103**, 7175–7180 (2006).

567   26.   Ner-Gaon, H. *et al.* Intron retention is a major phenomenon in alternative splicing in Arabidopsis.

568   *The Plant Journal* **39**, 877–885 (2004).

569   27.   Filichkin, S. A. *et al.* Abiotic Stresses Modulate Landscape of Poplar Transcriptome via Alternative

570   Splicing, Differential Intron Retention, and Isoform Ratio Switching. *Front. Plant Sci.* **9**, (2018).

571   28.   Chaudhary, S. *et al.* Alternative Splicing and Protein Diversity: Plants Versus Animals. *Front Plant

572   Sci* **10**, 708 (2019).

573   29.   Ganie, S. A. & Reddy, A. S. N. Stress-Induced Changes in Alternative Splicing Landscape in Rice:

574   Functional Significance of Splice Isoforms in Stress Tolerance. *Biology (Basel)* **10**, 309 (2021).

575   30.   Gao, P. *et al.* Alternative splicing dynamics and evolutionary divergence during embryogenesis in

576   wheat species. *Plant Biotechnol J* **19**, 1624–1643 (2021).

577   31.   Mercer, T. R., Dinger, M. E. & Mattick, J. S. Long non-coding RNAs: insights into functions. *Nature

578   Reviews Genetics* **10**, 155–159 (2009).

579   32.   Liu, J., Wang, H. & Chua, N.-H. Long noncoding RNA transcriptome of plants. *Plant Biotechnol J* **13**,

580   319–328 (2015).

581   33.   Palos, K., Yu, L., Railey, C. E., Nelson Dittrich, A. C. & Nelson, A. D. L. Linking discoveries,

582   mechanisms, and technologies to develop a clearer perspective on plant long noncoding RNAs.

583   *Plant Cell* **35**, 1762–1786 (2023).

584   34.   Wang, X., Fan, H., Wang, B. & Yuan, F. Research progress on the roles of lncRNAs in plant

585   development and stress responses. *Front Plant Sci* **14**, 1138901 (2023).

586   35.   Liu, H., Wang, R., Mao, B., Zhao, B. & Wang, J. Identification of lncRNAs involved in rice ovule

587   development and female gametophyte abortion by genome-wide screening and functional

588   analysis. *BMC Genomics* **20**, 90 (2019).

589   36.   Zhang, Y.-C. *et al.* Genome-wide screening and functional analysis identify a large number of long

590   noncoding RNAs involved in the sexual reproduction of rice. *Genome Biol* **15**, (2014).

591   37.   Chen, L. *et al.* Genome-wide analysis of long non-coding RNAs affecting roots development at an

592   early stage in the rice response to cadmium stress. *BMC Genomics* **19**, 460 (2018).

593   38.   Li, W. Q. *et al.* Genome-wide identification and characterization of long non-coding RNAs

594   responsive to Dickeya zeae in rice. *RSC Adv.* **8**, 34408–34417 (2018).

595   39.   Wang, Y. *et al.* Arabidopsis noncoding RNA mediates control of photomorphogenesis by red light.

596   *PNAS* **111**, 10359–10364 (2014).

597   40.   Fox, S. E. *et al.* De Novo Transcriptome Assembly and Analyses of Gene Expression during

598   Photomorphogenesis in Diploid Wheat Triticum monococcum. *PLoS One* **9**, (2014).

599   41.   najoshi. najoshi/sickle. (2020).

600   42.   Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-Seq.

601   *Bioinformatics* **25**, 1105–1111 (2009).

602   43.   Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq experiments with

603   TopHat and Cufflinks. *Nat Protoc* **7**, 562–578 (2012).

44.   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a

reference genome. *BMC Bioinformatics* **12**, 323 (2011).

45.   Leng, N. *et al.* EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments.

Bioinformatics **29**, 1035–1043 (2013).

46.   The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing strong.

Nucleic Acids Res **47**, D330–D338 (2019).

47.   Naithani, S. *et al.* Plant Reactome: a resource for plant pathways and comparative analysis. *Nucleic*

Acids Res **45**, D1029–D1039 (2017).

48.   Gupta, P. *et al.* Plant Reactome and PubChem: The Plant Pathway and (Bio)Chemical Entity

Knowledgebases. *Methods Mol Biol* **2443**, 511–525 (2022).

49.   Tello-Ruiz, M. K. *et al.* Gramene 2021: harnessing the power of comparative genomics and

pathways for plant research. *Nucleic Acids Res* **49**, D1452–D1463 (2021).

50.   Rogers, M. F., Thomas, J., Reddy, A. S. & Ben-Hur, A. SpliceGrapher: detecting patterns of

alternative splicing from RNA-Seq data in the context of gene models and EST data. *Genome Biol*

13, R4 (2012).

51.   Li, W., Feng, J. & Jiang, T. IsoLasso: A LASSO Regression Approach to RNA-Seq Based Transcriptome

Assembly. *J Comput Biol* **18**, 1693–1707 (2011).

52.   Kang, Y.-J. *et al.* CPC2: a fast and accurate coding potential calculator based on sequence intrinsic

features. *Nucleic Acids Res* **45**, W12–W16 (2017).

53.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**,

1236–1240 (2014).

54.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool.

Journal of Molecular Biology **215**, 403–410 (1990).

55.   Yi, X., Zhang, Z., Ling, Y., Xu, W. & Su, Z. PNRD: a plant non-coding RNA database. *Nucleic Acids Res*

43, D982–D989 (2015).

56.   Paytuví Gallart, A., Hermoso Pulido, A., Anzar Martínez de Lagrán, I., Sanseverino, W. & Aiese

Cigliano, R. GREENC: a Wiki-based database of plant lncRNAs. *Nucleic Acids Res* **44**, D1161–D1166

(2016).

57.   Zhang, Z., Xu, Y., Yang, F., Xiao, B. & Li, G. RiceLncPedia: a comprehensive database of rice long

non-coding RNAs. *Plant Biotechnol J* **19**, 1492–1494 (2021).

58.   Hofacker, I. L. Vienna RNA secondary structure server. *Nucleic Acids Res* **31**, 3429–3431 (2003).

59.   Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R. & Hofacker, I. L. The Vienna RNA Websuite.

Nucleic Acids Res **36**, W70–W74 (2008).

60.   Naithani, S. *et al.* Plant Reactome: a knowledgebase and resource for comparative pathway

analysis. *Nucleic Acids Res* **48**, D1093–D1103 (2020).

61.   Jin, J. *et al.* PlantTFDB 4.0: toward a central hub for transcription factors and regulatory

interactions in plants. *Nucleic Acids Res* **45**, D1040–D1045 (2017).

62.   Li, J. *et al.* A subgroup of MYB transcription factor genes undergoes highly conserved alternative

splicing in Arabidopsis and rice. *J Exp Bot* **57**, 1263–1273 (2006).

63.   Sugiyama, N., Izawa, T., Oikawa, T. & Shimamoto, K. Light regulation of circadian clock-controlled

gene expression in rice. *The Plant Journal* **26**, 607–615 (2001).

64.   Matsuzaki, J., Kawahara, Y. & Izawa, T. Punctual Transcriptional Regulation by the Rice Circadian

Clock under Fluctuating Field Conditions[OPEN]. *Plant Cell* **27**, 633–648 (2015).

65.   Cordoba, E., Salmi, M. & León, P. Unravelling the regulatory mechanisms that modulate the MEP

pathway in higher plants. *J Exp Bot* **60**, 2933–2943 (2009).

66.   Vranová, E., Coman, D. & Gruissem, W. Network Analysis of the MVA and MEP Pathways for

Isoprenoid Synthesis. *Annual Review of Plant Biology* **64**, 665–700 (2013).

67.   Vranová, E., Coman, D. & Gruissem, W. Structure and Dynamics of the Isoprenoid Pathway

Network. *Molecular Plant* **5**, 318–333 (2012).

68.   Liu, X., Hao, L., Li, D., Zhu, L. & Hu, S. Long Non-coding RNAs and Their Biological Roles in Plants.

Genomics Proteomics Bioinformatics **13**, 137–147 (2015).

69.   Zhou, R. *et al.* Analysis of Rice Transcriptome Reveals the LncRNA/CircRNA Regulation in Tissue

Development. *Rice (N Y)* **14**, 14 (2021).

70.   Serra, T. S. *et al.* OsRMC, a negative regulator of salt stress response in rice, is regulated by two

AP2/ERF transcription factors. *Plant Mol Biol* **82**, 439–455 (2013).

71.   Argueso, C. T., Ferreira, F. J. & Kieber, J. J. Environmental perception avenues: the interaction of

cytokinin and environmental response pathways. *Plant, Cell & Environment* **32**, 1147–1160 (2009).

72.   Worthen, J. M. *et al.* Type-B response regulators of rice play key roles in growth, development and

cytokinin signaling. *Development* **146**, dev174870 (2019).

73.   Cordeiro, A. M. *et al.* Rice phytochrome-interacting factor protein OsPIF14 represses OsDREB1B

gene expression through an extended N-box and interacts preferentially with the active form of

Phytochrome B. *Biochim Biophys Acta* **1859**, 393–404 (2016).

74.   Liang, C. *et al.* Transcriptomic, proteomic and metabolic changes in Arabidopsis thaliana leaves

after the onset of illumination. *BMC Plant Biology* **16**, 43 (2016).

75.   Ding, J. *et al.* Comparative Transcriptome Analysis of Gene Expression Patterns in Tomato Under

Dynamic Light Conditions. *Genes (Basel)* **10**, 662 (2019).

76.   Jin, X. *et al.* The Coordinated Upregulated Expression of Genes Involved in MEP, Chlorophyll,

Carotenoid and Tocopherol Pathways, Mirrored the Corresponding Metabolite Contents in Rice

Leaves during De-Etiolation. *Plants (Basel)* **10**, 1456 (2021).

77.   Bian, Z. *et al.* A Transcriptome Analysis Revealing the New Insight of Green Light on Tomato Plant

Growth and Drought Stress Tolerance. *Front Plant Sci* **12**, 649283 (2021).

78.   Ma, L. *et al.* Light Control of Arabidopsis Development Entails Coordinated Regulation of Genome

Expression and Cellular Pathways. *Plant Cell* **13**, 2589–2608 (2001).

79.   Jiao, Y., Lau, O. S. & Deng, X. W. Light-regulated transcriptional networks in higher plants. *Nature*

Reviews Genetics **8**, 217–230 (2007).

80.   de Wit, M., Galvão, V. C. & Fankhauser, C. Light-Mediated Hormonal Regulation of Plant Growth

and Development. *Annual Review of Plant Biology* **67**, 513–537 (2016).

81.   Hunter, W. N. The Non-mevalonate Pathway of Isoprenoid Precursor Biosynthesis. *J. Biol. Chem.*

282, 21573–21577 (2007).

82.   Gupta, P. *et al.* Cloning and characterization of 2-C-methyl-d-erythritol-4-phosphate pathway

genes for isoprenoid biosynthesis from Indian ginseng, Withania somnifera. *Protoplasma* **250**,

285–295 (2013).

83.   Pulido, P., Perello, C. & Rodriguez-Concepcion, M. New Insights into Plant Isoprenoid Metabolism.

Molecular Plant **5**, 964–967 (2012).

84.   Nagata, N., Suzuki, M., Yoshida, S. & Muranaka, T. Mevalonic acid partially restores chloroplast and

etioplast development in Arabidopsis lacking the non-mevalonate pathway. *Planta* **216**, 345–350

(2002).

85.   Rodríguez-Concepción, M. Early Steps in Isoprenoid Biosynthesis: Multilevel Regulation of the

Supply of Common Precursors in Plant Cells. *Phytochem Rev* **5**, 1–15 (2006).

86.   Ghassemian, M. *et al.* Integrative analysis of transcript and metabolite profiling data sets to

evaluate the regulation of biochemical pathways during photomorphogenesis. *Archives of*

Biochemistry and Biophysics **448**, 45–59 (2006).

87.   Vranová, E., Coman, D. & Gruissem, W. Structure and Dynamics of the Isoprenoid Pathway

Network. *Molecular Plant* **5**, 318–333 (2012).

88.   Fung-Uceda, J. *et al.* The Circadian Clock Sets the Time of DNA Replication Licensing to Regulate

Growth in Arabidopsis. *Developmental Cell* **45**, 101-113.e4 (2018).

89.   Filichkin, S. A. *et al.* Genome-wide mapping of alternative splicing in Arabidopsis thaliana. *Genome*

Res **20**, 45–58 (2010).

90.   Li, P. *et al.* The developmental dynamics of the maize leaf transcriptome. *Nature Genetics* **42**,

1060–1067 (2010).

91.   Shen, Y. *et al.* Global Dissection of Alternative Splicing in Paleopolyploid Soybean. *The Plant Cell* **26**,

996–1008 (2014).

92.   Campbell, M. A., Haas, B. J., Hamilton, J. P., Mount, S. M. & Buell, C. R. Comprehensive analysis of

alternative splicing in rice and comparative analyses with Arabidopsis. *BMC Genomics* **7**, 327

(2006).

93.   Csorba, T., Questa, J. I., Sun, Q. & Dean, C. Antisense COOLAIR mediates the coordinated switching

of chromatin states at FLC during vernalization. *PNAS* **111**, 16160–16165 (2014).

94.   Filichkin, S. A. *et al.* Abiotic Stresses Modulate Landscape of Poplar Transcriptome via Alternative

Splicing, Differential Intron Retention, and Isoform Ratio Switching. *Front. Plant Sci.* **9**, (2018).

95.   Deng, P., Liu, S., Nie, X., Weining, S. & Wu, L. Conservation analysis of long non-coding RNAs in

plants. *Sci. China Life Sci.* **61**, 190–198 (2018).

96.   Liu, J. *et al.* Genome-Wide Analysis Uncovers Regulation of Long Intergenic Noncoding RNAs in

Arabidopsis[C][W]. *Plant Cell* **24**, 4333–4345 (2012).

97.   Corona-Gomez, J. A., Garcia-Lopez, I. J., Stadler, P. F. & Fernandez-Valverde, S. L. Splicing

conservation signals in plant long non-coding RNAs. *bioRxiv* 588954 (2019) doi:10.1101/588954.

98.   Komiya, R. *et al.* Rice germline-specific Argonaute MEL1 protein binds to phasiRNAs generated

from more than 700 lincRNAs. *The Plant Journal* **78**, 385–397 (2014).

99.   Lavarenne, J. *et al.* Transcriptome profiling of laser-captured crown root primordia reveals new

722   pathways activated during early stages of crown root formation in rice. *PLoS One* **15**, e0238736

723   (2020).

724   100.   Yan, Y., Wang, H., Hamera, S., Chen, X. & Fang, R. miR444a has multiple functions in the rice

725   nitrate-signaling pathway. *The Plant Journal* **78**, 44–55 (2014).

726   101.   Chen, H. *et al.* OsMADS27 regulates the root development in a NO3−—Dependent manner and

727   modulates the salt tolerance in rice (Oryza sativa L.). *Plant Science* **277**, 20–32 (2018).

728   102.   Pachamuthu, K. *et al.* Nitrate-dependent regulation of miR444-OsMADS27 signalling cascade

729   controls root development in rice. *Journal of Experimental Botany* **73**, 3511–3530 (2022).

730   103.   Pathak, R. R. *et al.* Transcriptomic and network analyses reveal distinct nitrate responses in light

731   and dark in rice leaves (Oryza sativa Indica var. Panvel1). *Sci Rep* **10**, 12228 (2020).

732   104.   Yoneyama, T. & Suzuki, A. Light-Independent Nitrogen Assimilation in Plant Leaves: Nitrate

733   Incorporation into Glutamine, Glutamate, Aspartate, and Asparagine Traced by 15N. *Plants (Basel)*

734   **9**, 1303 (2020).

735   105.   Geniza, M. *et al.* Comparative Analysis of Salinity Response Transcriptomes in Salt-Tolerant Pokkali

736   and Susceptible IR29 Rice. 2023.08.20.551368 Preprint at

737   https://doi.org/10.1101/2023.08.20.551368 (2023).

738   106.   Al-Bader, N. *et al.* Loss of a Premature Stop Codon in the Rice Wall-Associated Kinase 91 (WAK91)

739   Gene Is a Candidate for Improving Leaf Sheath Blight Disease Resistance. *Genes* **14**, 1673 (2023).

740   107.   Hartmann, L. *et al.* Alternative Splicing Substantially Diversifies the Transcriptome during Early

741   Photomorphogenesis and Correlates with the Energy Availability in Arabidopsis. *The Plant Cell* **28**,

742   2715–2734 (2016).

743   108.   Naithani, S., Deng, C. H., Sahu, S. K. & Jaiswal, P. Exploring Pan-Genomes of Crops: Genomic

744   Resources and Tools for Unraveling Gene Evolution, Function, and Adaptation. Preprint at

745   https://doi.org/10.20944/preprints202308.1641.v1 (2023).

Figure-1

Figure-2

MVA Pathway

	Dark	Light
AACT_OS01G0110400		
HMGS_OS08G0544900		
HMGS_OS09G0521400		
HMGR_OS08G0512700		
HMGR_OS09G0492700		
MK_OS10G0329300		
PMK_OS03G0253100		
MDD_OS02G0107200		
MDD_OS02G0109100		
IPI1_OS05G0413400		
IPI1_OS07G0546000		

Pathway diagram:

MVA Pathway

Acetyl-CoA → AACT (2.3.1.9) → Acetoacetyl-CoA → HMGS (2.3.3.10) → HMG-CoA → HMGR (1.1.1.34) → MVA → MK (2.7.1.36) → MVA-5-P → PMK (2.7.4.2) → MVAPP → MDD (4.1.1.33) → DMAPP ↔ IPP

MEP Pathway (Plastid)

Pyruvate + Gly-3-P → DXS (2.2.1.7) → DOXP → DXR (1.1.1.267) → MEP → CDP-MES (2.7.7.60) → CDP-ME → CDP-MEK (2.7.1.148) → CDP-MEP → MECPS (4.6.1.12) → MECP → HDS (1.17.7.3) → HMBPP → HDR (1.17.7.4) → DMAPP ↔ IPP

MEP Pathway

	Dark	Light
DXS_OS05G0408900		
DXS_OS07G0190000		
DXR_OS01G0106900		
MECPS_OS01G0887100		
CDPMEK_OS01G0802100		
CDPMES_OS02G0680600		
HDS_OS02G0603800		
HDR_OS03G0731900		

Color scale: 1, 0.5, 0, -0.5, -1

Pathway	Gene	Dark IR	Light IR	Light ES
MVA	OS01G0110400 (AACT)	2	1	-
	OS08G0544900 (HMGS)	1	-	-
	OS10G0329300 (MK)	1	1	1
	OS03G0253100 (PMK)	1	-	-
MEP	OS05G0408900 (DXS)	-	1	-
	OS01G0106900 (DXR)	-	2	-
	OS01G0887100 (MECPS)	-	1	-

DXS (Light) — OS05G0408900_Light; OS05G0408900 Read Coverage; 19963120 – 19957171

DXR (Light) — OS01G0106900_Light; OS01G0106900 Read Coverage; 380713 – 374728

HMGS (Dark) — OS08G0544900_Dark; OS08G0544900 Read Coverage; 27302249 – 27306827

PMK (Dark) — OS03G0253100_Dark; OS03G0253100 Read Coverage; 8071826 – 8067630

■ Intron Retention (IR)

Figure-3