No single PCR test is sufficient to determine parvovirus

IHHNV presence in or impact on farmed shrimp production

Kallaya Sritunyalucksana[1,#], Piyachat Sanguanrut[1,#], Jiraporn Srisala[1],

Jumroensri Thawonsuwan[2], Nattakan Saleetid[3], Rapeepun Vanichviriyakit[4,5],

Charoonroj Chotwiwatthanakun[5,6], Timothy W. Flegel[5,7],

Suparat Taengchaiyaphum[1,*]

[1]Aquatic Animal Health Research Team, Integrative Aquaculture Biotechnology Research

Group, National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science

and Technology Development Agency (NSTDA), Yothi office, Rama VI Rd., Bangkok, 10400,

Thailand

[2]Songkhla Aquatic Animal Health Research and Development Center, Aquatic Animal Health

Research and Development Division, Department of Fisheries, 130/2 Pawong, Amphur Muang,

Songkhla, 90100, Thailand

[3]Coastal Aquaculture Research and Development Division, Department of Fisheries, 50

Phahonyothin Rd., Lat Yao, Chatuchak, Bangkok 10900, Thailand

[4]Department of Anatomy, Faculty of Science, Mahidol University, Rama VI rd., Bangkok,

10400, Thailand

[5]Center of Excellence for Shrimp Molecular Biology and Biotechnology (Centex Shrimp),

Faculty of Science, Mahidol University, Rama VI Rd., Bangkok, 10400, Thailand

[6]Mahidol University, Nakhonsawan Campus, Nakhonsawan, 60000, Thailand

[7]National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and

Technology Development Agency (NSTDA), Klong Luang, Pathumthani, 12120, Thailand

#These authors with equally contribution to the work

*corresponding author

e-mail address: suparat.tae@biotec.or.th

ABSTRACT

The main purpose of this report is to provide hard evidence that the shrimp parvovirus IHHNV has not resulted "in significant consequences e.g. production losses, morbidity or mortality at a zone or country level" in Thailand since at least 2010. It also reveals that no single PCR test is sufficient to identify IHHNV-infected shrimp. It presents historical evidence and new evidence from 11 commercial ponds cultivating the giant tiger shrimp *Penaeus monodon* in Thailand. These ponds were selected because they were the ponds that gave positive PCR test results for IHHNV using 2 methods recommended for IHHNV diagnosis by WOAH. However, an additional in-house "IHHNV Long-Amp method" (IHHNV-LA) was also used to amplify 90% of the 4 kb IHHNV genome sequence, and it also gave false-positive test results with 2 of the 11 ponds. Further tests using normal histopathological analysis for the presence of pathognomonic Cowdry A type inclusions (CAI), *in situ* hybridization (ISH) and immunohistochemistry (IHC) could confirm IHHNV infections in only 2 of the 4 ponds PCR-positive using all 3 PCR methods. In addition, positive detection of CAI alone was equivalent to ISH or IHC in confirming IHHNV infection after a positive test with any of the PCR methods used. In summary, the recommended WOAH PCR methods gave false positive test results for IHHNV infection with 9/11 ponds (82%). All 11 ponds gave profitable harvests despite the confirmation of IHHNV in 2 ponds, where it was accompanied by various additional pathogens. Unfortunately, according to current practice, positive PCR test results with the WOAH methods alone sometimes leads to rejection of traded shrimp products without assurance that the test results are not false positive results that may arise from endogenous viral elements (EVE).

Keywords: Infectious hypodermal and hematopoietic necrosis virus (IHHNV), *Penaeus monodon*, OIE IHHNV detection method, False positive, Long-amp IHHNV detection

1. INTRODUCTION

1.1. Identity of IHHNV

The official scientific name of infectious hypodermal and hematopoietic necrosis virus (IHHNV) is now *Penstylhamaparvovirus* 1 in the family *Parvoviridae* and sub-family *Hamaparvovirinae* (Pénzes et al., 2020). Acute, catastrophic disease with mass mortality was originally reported in IHHNV-infected *Penaeus stylirostris* (Lightner et al., 1983). It was

discovered that the virus originated from grossly normal stocks of captured wild *Penaeus monodon* that were imported for experimental culture in the Americas in facilities where *P. stylirostris* was also under experimental culture. Subsequent IHHNV disease outbreaks in *Penaeus vannamei* in the Americas resulted not in mortality, but instead retarded growth accompanied by anatomical deformities (e.g., bent rostra and stunted growth) called runt deformity syndrome (RDS) (Bell and Lightner, 1984). The coefficient of variation (CV) in size for cultivated populations with RDS was typically 35-90%, with the mean shifted towards unmarketable small sizes that dominated crops and made them unprofitable. This compared to normal IHHNV-free populations with CVs of 10-30% (Bray et al., 1994; Brock and Main, 1994; Carr et al., 1996; Lightner, Donald V, 1996; Primavera and Quinitio, 2000; Pruder, 1995). The effect of IHHNV on growth of *P. monodon* is controversial, but confirmed studies (Flegel et al., 2004; Withyachumnarnkul et al., 2006) indicate that it does not cause RDS or significant negative effects on growth similar to those that occur in *P. vannamei*.

IHHNV was among the 5 shrimp viruses listed in the first edition of the WOAH Aquatic Animal Health Code (AAHC) in 1995 and it is still listed in the current edition (WOAH, 2019a). Three distinct genotypes of IHHNV were listed in the 2019 online edition of the OIE Aquatic Manual of Diagnostic Tests for Aquatic Animals (WOAH, 2019b). These are Types 1 and 2 that are infectious viruses, while Type 3 was called non-infectious IHHNV. All are detectable by conventional PCR tests. The primer pair 309F/R can detect all the known genotypes of infectious IHHNV while primer pair 389F/R can also detect non-infectious Type 3 (Tang et al., 2000; Tang et al., 2007) which comprises IHHNV-related DNA sequences present in the genome of *P. monodon* (Saksmerprome et al., 2011; Tang and 84 Lightner, 2006; Tang et al., 2007). Type 3 has recently been revealed to be an assembly artifact that arose from scrambled sequences that sometimes occur in disjunctive clusters within the genome of *P. monodon* and constitute remnants of an ancient form of IHHNV (Huerlimann et al., 2022; Taengchaiyaphum et al., 2022). Such non-retroviral virus sequences in animal genomes are called endogenous viral elements (EVE) (Feschotte and Gilbert, 2012).

The target tissues for IHHNV infection in penaeid shrimp are of ectodermal origin (e.g., the gills and sub-cuticular epidermis), and mesodermal origin (e.g., haematopoietic tissue and antennal

glands). Histological characteristic IHHNV lesions obtained using hematoxylin and eosin (H&E)

staining are called Cowdry Type A inclusions (CAI) comprising a central, eosinophilic,

intranuclear inclusion separated by a small unstained space from marginated (basophilic)

chromatin (Bell and Lightner, 1984; Lightner, 1996). CAIs are an artifact of the Davidson's acid

fixative that is recommended to preserve shrimp tissue for histological analysis and they do not

occur when other fixatives such as neutral phosphate buffered fixative are used.

1.2. Decline in negative economic impact of IHHNV

The negative impact of IHHNV on cultured shrimp production in the USA dropped

dramatically after the development and adoption of domesticated, specific pathogen free (SPF)

shrimp stocks free of the virus (Wyban, 2007; Wyban et al., 1992). Work in Hawaii revealed that

stocking PL derived from such shrimp in cultivation ponds did not result in significant RDS or

other negative impact even if the shrimp became infected after stocking by residual virus from the

previous culture cycle. Consistent, subsequent stocking of the same pond with IHHNV-free PL

resulted in no negative impact and in a progressive decline in the number of IHHNV-positive

specimens. Similarly, upon the first introduction of non-SPF *P. vannamei* for cultivation in

Thailand in the late 1990's a few cases of RDS were seen, but after the mid 2000's in Thailand

upon the widespread use of IHHNV-free PL of *P. vannamei*, no cases of RDS have been reported

(Flegel, 2006).

In September 2020, a report from the OIE Aquatic Animal Health Standards Commission (AAHC)

responded to an official request from Ecuador to review the impact of IHHNV and to consider

removing it from its list of crustacean diseases [Report of the meeting of the OIE Aquatic Animal

Health Standards Commission (AAHC), Paris, 19–26 February 2020 in Section 4.6, Annex 9].

The AAHC upheld the continued listing of IHHNV solely because it concluded that IHHNV fit

Criterion 4b: "The disease has been shown to affect the health of cultured aquatic animals at the

level of a country or a zone resulting in significant consequences e.g. production losses, morbidity

or mortality at a zone or country level". If the AAHC had deemed that IHHNV did not satisfy this

criterion, it would have been required to delist it.

To support their decision that IHHNV did satisfy Criterion 4b, the AAHC report cited 22
references, 8 of which were published prior to 2010 while only 14 covered the period 2010 to 2020
during which the impact of IHHNV on shrimp production should have been assessed. In
Supplementary Information 1, these remaining 14 cited references are examined in detail and it
is shown that only 2 of them (Jagadeesan et al., 2019; Sellars et al., 2019) contained new data
showing production losses in ponds infected with IHHNV. All the other 12 publications from 2010
onward concerned studies of IHHNV itself in terms of biology and detection and none contained
original data regarding the impact of IHHNV on production.

The reasons why the remaining 2 publications are inappropriate to use to justify Criterion 4b are
described in the Supplementary Information 1. Both publications were based predominantly on
positive PCR results for detection of IHHNV with no additional histological examination (e.g., *in*
situ hybridization or immunohistochemistry) to confirm IHHNV infections. Whilst the data of
Sellars et al. 2019 was taken from a broader data set where additional PCR was carried out to
confirm samples were negative from all other endemic pathogens in the region, it is unknown if
those of Jagadeesan et al. 2019 had any additional PCR data to demonstrate the presence of other
pathogenic agents. Sellars et al. 2019 also based the financial cost calculations on an Australian
production model that is unique compared to the rest of the world in that they achieve very high 8
tons/ha averages whilst the average globally is around 6.0-7.5 tons/ha. Clearly, all the harvests
would have been profitable, but some more-so than others. Profits from farm to farm vary based
on multiple, complex factors including management. Thus, it is not rational to refer to the profits
of some farmers as losses because they are lower than those of another.

The study of Jagadeesan et al. 2019 was a survey of 350 *P*. *vannamei* ponds in India of which 30
were PCR positive for IHHNV. Of these 16 had problems while 14 did not. Since approximately
half of the farms positive for IHHNV had problems while half did not, it is our opinion that IHHNV
was not significantly associated with problems and that other causes for the problems should have
been further sought. The publications from the interval 2010 to 2020 that describe lack of
significant problems linked to IHHNV were not included in the AAHC review (e.g., Hernández-
Ruíz et al., 2020; Mendoza-Cano et al., 2014). Furthermore, a recent publication by Aranguren

Caro et al. (2022) showed lack of a significant impact of IHHNV on production, with IHHNV only causing small measurable reductions in growth in some ponds.

1.3. Study of arbitrarily selected Thai cultivation ponds positive for IHHNV by PCR

In this paper we present results from 11 recently harvested Thai shrimp ponds that were selected because they gave positive PCR test results for IHHNV using two WOAH PCR detection methods via active surveillance but did not exhibit significant negative effects on production that could be ascribed to IHHNV. The results also revealed significant problems might arise from false positive tests for IHHNV using currently recommended WOAH methods, if additional confirmatory tests are not carried out. In our experience, these should not be additional PCR tests, even with sequencing, but should be histopathological analysis for Cowdry A type inclusions (CAI) and/or *in situ* hybridization (ISH) or immunohistochemistry (IHC). We hope that this information will stir others to carry out similar studies and provide more evidence from production ponds to support the contention that IHHNV no longer has "significant consequences e.g. production losses, morbidity or mortality at a zone or country level."

1.4 A precautionary note

It is important to understand from the outset that we do not consider IHHNV to be innocuous. It should not be ignored and must remain included in the group of pathogens that should be excluded from specific pathogen free (SPF) shrimp used to supply shrimp farmers with postlarvae (PL) to stock their cultivation ponds. The situation is the same as that for the viruses MBV and BP that were formerly listed by OIE but were later removed from the list (WOAH, 2008). They were removed because full understanding of their biology allowed for their exclusion from hatcheries and from post-larvae supplied to shrimp farmers for stocking in their cultivation ponds. Hepatopancreatic parvovirus (HPV), now called decapod hepanhamaparvovirus (DHPV), causes much more severe growth retardation than IHHNV but has never been listed for similar reasons. However, producers that use wild, captured PL or broodstock to produce PL must be aware that these unlisted pathogens must be excluded by screening broodstock or PL for their absence. This is because their presence may decrease production efficiency and profitability even though they may not cause financial losses.

The same situation now applies for IHHNV. The current, high-quality PL produced from SPF

stocks of *P. vannamei* are not infected with IHHNV or other known pathogens and they produce

offspring that can achieve average daily weight gains of 0.3 to 0.4 g or more under ideal culture

conditions. In addition, they are unaffected by IHHNV infection. The weight gain is well beyond

that possible with offspring obtained from captured, wild broodstock. This achievement has

resulted from a combination of innovations in breeding, selection, feed and culture management.

Thus, no pathogen can simply be ignored and assumed to have no possible negative impact on

production. At the same time, no manageable shrimp pathogen should be considered as a perpetual

dire threat when the evidence clearly shows that it is not.

Rational aquaculture management requires that shrimp farmers take precautions to ensure

optimum culture conditions such as exclusion of known pathogens, including IHHNV. This can

be achieved either by testing before stocking or by using PL from a supplier with a certified long

history of providing PL free of known pathogens. For comparison, we point out from WHO data

from June 2024 that total global reports of Covid-19 infection were approximately 35,000 while

deaths were approximately 600. Thus, thousands more survive after suffering light to severe

illness. All these Covid-19 infections, lethal or not, involve negative impacts. Despite this, all

restrictions on movements of people worldwide have now been removed. This does not mean that

Covid-19 presents no threat so long as reasonable precautions are taken. In our opinion, the

situation is similar to that for the shrimp viruses MBV, BP and HPV and the work described herein

supports the argument that like them, IHHNV should not be a listed pathogen.

2.   MATERIALS AND METHODS

2.1. Shrimp specimens

Juvenile giant tiger shrimp (*P. monodon*) were obtained from 2 sets of shrimp cultivation

ponds. These ponds were selected based on being positive for IHHNV during an active

surveillance using WOAH-recommended PCR methods. The PL used to stock the ponds were not

tested for IHHNV, so the IHHNV source was unknown. The first, **Pond Set 1** consisted of 4 ponds,

1 from Chachoengsao province on the eastern coast of the Gulf of Thailand and 3 from Krabi

province on the southwestern Andaman Sea coast of Thailand. These were cultivated from

February to August, during the optimum time for shrimp production in Thailand. Another set of 7

ponds, **Pond Set 2** were obtained from the adjacent provinces Phuket and Phang Nga on the southwestern coast of Thailand on the Andaman Sea. These were cultivated from October to April, through the Thai winter (November to February) which is the riskiest time for shrimp cultivation in Thailand. However, this period attracts some farmers due to higher product prices.

Thirty shrimp were collected randomly from each pond at harvest to assess the range of shrimp sizes by fresh weight. For Pond Set 1, an additional 5 were collected from each pond for PCR and histological analysis. Pleopods from the 5 shrimp specimens were removed for PCR analysis, while the cephalothoraxes of the same specimens were processed using standard methods (Bell and Lightner, 1988) to confirm IHHNV presence by normal histological analysis plus ISH and IHC analysis. For Pond Set 2, the technician mistakenly prepared 6 shrimp specimens from each pond for PCR analysis and a separate set of 3 shrimp each for histological analysis. Thus, for Pond Set 2, results from the PCR tests and histological analysis could not be directly compared, but production results could be compared between ponds confirmed or not for the presence of infectious IHHNV.

2.2. PCR detection methods for IHHNV

Three PCR detection methods for IHHNV detection were used in this study (**Table 1**). Two comprised WOAH-recommended methods here called the IHHNV-309 method employing primer pairs 309F/R and the IHHNV-389 method employing primer pairs 389F/R (Tang et al., 2000; Tang et al., 2007). These methods were carried out following the WOAH manual (WOAH, 2019b). A third, long-amp IHHNV (IHHNV-LA) detection method consisted of a two-step nested PCR protocol that was established in our laboratory to cover 90% (~3.6 kb) of the whole genome sequence of infectious IHHNV (4.1 kb). The primers were designed from the GenBank database of IHHNV accession number AF218266. The sensitivity limit of the IHHNV-LA method (10 copies/ng total DNA) was determined using a DNA plasmid containing an insert of the 3,665 bp-product from the first amplification step as the template (Taengchaiyaphum et al., 2021).

Table 1. Primers used for detection of IHHNV by PCR in this study.

Method	Primer	Sequence (5'-3')	Product size	Reference
IHHNV-309	309F	TCCAACACTTAGTCAAAACCAA	309 bp	Tang et al., 2007
	309R	TGTCTGCTACGATGATTATCCA		
IHHNV-389	389F	CGGAACACAACCCGACTTTA	389 bp	Tang et al., 2000
	389R	GGCCAAGACCAAAATACGAA		
IHHNV-LA	First step PCR			
	3665F	CCCAGTTTCTAACTGACGAGTGAAGAGG	3,665 bp	Taengchaiyaphum et al., 2021
	3665R	CCTGACTCTAAATGACTGACTGACGATAGGG		
	Nested PCR			
	1044F	ACAGATGTCTACAATTCAAT	1,044 bp	Taengchaiyaphum et al., 2021
	1044R	AATAGAAGAAAGTGTCCC		

2.3. DNA extraction

Pleopods were collected from 5 individual shrimp from each pond and homogenized for separate analysis in DNA lysis buffer (50 mM Tris-base, 100 mM EDTA, 50 mM NaCl, 2% (w/v) SDS and 100 µg/ml proteinase K). The mixtures were incubated at 56 ℃ for 1 h before DNA was extracted using an QIAamp DNA Mini Kit (Qiagen, Germany) according to the manufacturer's protocol. Total DNA concentration was determined by Qubit 3.0 Fluorometer (Life Technologies, USA).

2.4. Histological analysis

Histological analysis was carried out by standard methods (Bell & Lightner, 1988). Briefly, the cephalothorax of each shrimp sample was fixed with Davidson's fixative for 24 h and then changed to 70% ethanol. The tissue was next dehydrated with a series of graded ethanol ending with xylene and embedded in paraffin. Sections were cut at 4 µm and placed onto glass slides for H&E staining to detect of Cowdry Type A inclusions (CAI) characteristic for IHHNV, for *in situ* hybridization (ISH) analysis and for immunohistochemistry (IHC) analysis (Lightner 1996). Positive control slides were prepared using archived tissue blocks of shrimp infected with IHHNV and showing typical IHHNV-CAI. We consider the detection of CAI as the gold standard for confirmation of IHHNV infection. The farm samples were also examined histologically for the

presence of lesions in the hepatopancreas (HP) caused by monodon baculovirus (MBV),
hepatopancreatic parvovirus (HPV) and other pathogens such as bacteria known to cause retarded
growth in *P. monodon* (Flegel, 2004; Flegel et al., 2006). The viruses MBV and HPV are now
called *Penaeus monodon* nudivirus (PmNV) (Yang et al., 2014) and *Penaeus monodon*
hepandhamaparvovirus (Cotmore et al., 2019), respectively. Analysis of the lesions for these 2
viruses was carried out as recommended by Lightner (1996). The HP of the specimens were also
screened for the presence of the microsporidian *Enterocytozoon hepatopenaei* (EHP) that is known
to infect the HP and to be associated with retarded shrimp growth (Chaijarasphong et al., 2021).

2.5. *In situ* hybridization (ISH)

The PCR products amplified from both IHHNV-309 and IHHNV-389 methods were used as
templates to prepare digoxigenin (DIG)-labeled IHHNV probes using a PCR DIG-labeling kit
(Roche, Germany). After preparation, probes were purified using a Gel/PCR cleanup kit (Geneaid,
Taiwan) using the protocol described by the manufacturer. The protocol for ISH has been
previously described (Sanguanrut et al., 2022). Briefly, slides containing adjacent tissue sections
to those used for histological analysis by H&E staining were de-paraffinized, rehydrated and TNE
buffer (500 mM Tris-Cl, 100 mM NaCl, l0 mM EDTA) before treatment with proteinase K (Sigma,
USA) and assay using 200 ng per slide of the DIG-labeled probes in hybridization buffer. A control
reaction without probe was included in a separate container. After incubation overnight at 42 ℃
the slides were washed, incubated with blocking buffer and further incubated with 1:500 anti-DIG-
AP antibody in the dark, washed and developed for signal using NBT/BCIP solution (Roche,
Germany) and counterstained with 0.5% Bismarck Brown Y (Sigma, USA) for microscopic
examination (Leica DM750, ICC50W model with a digital camera and LASV4.12 Software).

2.6. Immunohistochemistry (IHC) analysis

A polyclonal antibody to the capsid protein of IHHNV was produced in a rabbit using
recombinant IHHNV capsid protein and the concentration appropriate for IHC was determined.
The tissue sections were de-paraffinized, rehydrated with xylene and a series of graded alcohols
to water. They were then treated with 0.6% H_2O_2 in methanol for 15 min to quench endogenous
peroxidase activity followed by washing two times with 1X PBS for 5 min each. The slides were
incubated with blocking buffer (10% fetal bovine serum in 1X PBS pH 7.4) for 1 h, at RT and

replaced with rabbit anti-capsid IHHNV antibody (1:1000 in blocking buffer) (Kiatmetha et al., 2018) in a moist chamber at 37 °C for 1-2 h. The slides were washed three times to remove unbound antibody with 1X PBS for 5 min each followed by incubation with horseradish peroxidase (HRP)-conjugated goat anti rabbit antibody (1:500 in the blocking buffer) at 37 °C for 1 h in a moist chamber before washing three times with 1X PBS for 5 min each. The slides were incubated with a VECTOR®NovaRED™ substrate kit (Vector laboratories, USA) for 5-15 min and washed with water for 5 min before counterstaining with Mayer's hematoxylin, dehydrating in a series of graded alcohols to xylene and covering with a cover glass.

3. RESULTS

3.1. *History of official surveillance reports of IHHNV in Thailand*

For general interest, the percentage prevalence of IHHNV in samples of both *P. monodon* and *P. vannamei* received by the Thai Department of Fisheries (Thai DOF), the competent authority for shrimp diseases in Thailand from the years 2011 to 2021 is shown in **Fig. 1** together with National production. In 2013, production dropped precipitously due mostly to reduced cultivation because farmers feared acute hepatopancreatic necrosis disease (AHPND) (Sanguanrut et al., 2018). Subsequent recovery to pre-2013 levels has not occurred due to bankruptcy of the majority of Thai shrimp export companies during the AHPND crisis, and to concomitant loss of export market share to other shrimp producing countries. In any case, the graph clearly shows that the percentage of IHHNV positive samples was not significantly correlated with a negative effect on shrimp production. In fact, a Pearson correlation analysis (**Fig. 2**) showed a significant (p = 0.015) positive association between production quantity and the percentage of positive IHHNV detections in the total number of farmed shrimp specimens examined by active and passive surveillance by the Thai DOF. This result clearly reveals that simple data on detection of IHHNV prevalence cannot be used to predict shrimp production level.

Figure 1. Thailand shrimp production (blue bars and left Y axis) for both *P. Monodon* and *P. Vannamei* versus the percentage of IHHNV-PCR positive cases (dotted red line and right Y axis) from 2011 to 2021. The IHHNV detection from 2011-2017 was recorded through a Thai DOF active surveillance program, whereas those in 2018-2021 were recorded through a Thai DOF passive surveillance program. The dotted line indicates the percentage (right axis) of positive IHHNV-PCR test results in the total number of samples tested for any reason by Thai DOF in each individual year.

Figure 2. Correlation between percentage of IHHNV-PCR positive cases and shrimp production in Thailand from 2011 to 2021. The Pearson product-moment correlation coefficient (r) is 0.7 (p = 0.015) indicating a significant positive correlation between the yearly percentage of IHHNV-PCR positive samples and yearly shrimp production.

3.2. *IHHNV relationship to production in Pond Set 1*

3.2.1. All ponds in Pond Set 1 gave profitable harvests

Shrimp from all 4 culture ponds in Pond Set 1 (selected due to positive PCR test results for IHHNV) were profitably harvested with total culture periods between 123 and 185 days (**Table 2**). Cultivation was carried out beginning during the months of February to August which is the optimum period of year for shrimp cultivation in Thailand. The coefficients of variation for weight

were in the normal range from 16-25%. This contrasts with those reported for RDS at 35-90% (see introduction for references). In addition, the average daily weight gain for all the ponds was approximately 0.2 g, which is also normal for *P. monodon* in Thailand (Limsuwan and Chanratchakool, 2004).

Table 2. Shrimp growth performance and pond productivity in Pond Set 1.

Pond	Stocking density (shrimp/m^2)	Duration of culture (days)	CV (%)	Mean harvested weight (grams) (mean± SD)	Avg. Daily gain (g)
1	36	123	16	25.2±8.50	0.20
2	50	152	19	30.9±5.97	0.20
3	54	149	22	37.0±8.13	0.25
4	55	185	25	32.5±8.14	0.18

The size distribution for the shrimp in the 4 ponds is shown in **Fig 3**. Pond #1 (blue bars) was cultivated for the normal period of approximately 4 months and the mean weight of the shrimp was 25.2±8.50 g. The size-group distribution had a median of 29-35 g characterized by a tail towards larger sizes. This is the opposite to RDS where the median would be below the normal size and the distribution would be shifted sufficiently towards small sizes to make the whole crop unprofitable. That was not the situation with any of the 11 ponds in this study. The time of final harvest, and sometimes partial harvesting times, chosen by farmers are dependent on highly variable market conditions. Despite these variables, the overall ADGs, shrimp sizes and CVs for all the ponds (16 to 25) can be considered normal to better than normal for Thai conditions and not characteristic of RDS. RDS is characterized by a mean weight at harvest of non-profitable shrimp sizes and a CV of 35% or more shifted towards sizes of low value (Kalagayan et al., 1991).

Figure 3. Bar graph showing the size distribution of the shrimp in the 4 ponds studied. The medians for all the ponds were in the range of 22 to 35g.

3.2.2. Two ponds from Pond Set 1 were PCR-positive using the IHHNV-LA method

Using a plasmid carrying the target sequence for IHHNV, the detection sensitivity of the IHHNV-LA nested PCR method was found to be 10 copies (**Supplementary Figure S1**). Of the DNA extracts from 5 shrimp randomly collected from each of the 4 ponds in Pond Set 1, all gave positive PCR test results with both the OIE-IHHNV-309 and OIE-IHHNV-389 methods, while only 2 ponds (Pond #1 and #2) gave additional positive PCR test results with the IHHNV-LA method. Representative agarose gels of PCR amplicons from Pond Set 1 are shown in **Fig. 4** and the results are summarized for all 20 shrimp specimens in **Table 3**.

Figure 4. Example gels of PCR amplicons from shrimp specimens using 3 IHHNV-PCR detection methods. (A) Agarose gels showing positive PCR test results for all 3 PCR detection methods with the representative specimens from Pond #1. (B) Agarose gels from representative specimens from Pond #3 showing positive PCR test results with the 2 OIE methods only and negative results for the IHHNV-LA method.

Table 3. Summary of IHHNV-DNA detection by PCR and histological methods in juvenile shrimp from the 4 culture ponds. ND = not determined.

Pond #	PCR methods			Histological methods		
	IHHNV-309	IHHNV-389	IHHNV-LA	CAI	ISH	IHC
1	5/5	5/5	5/5	1/5	1/5	1/5
2	5/5	5/5	5/5	0/5	ND	ND
3	5/5	5/5	0/5	0/5	0/5	0/5
4	5/5	5/5	0/5	0/5	ND	ND

3.2.3. Histological failure to confirm IHHNV in 4/5 IHHNV-LA-positive shrimp

Shrimp from all 4 ponds (5 each, total 20) were subjected to histological examination. There were no cells of the hematopoietic tissue, the cuticular epidermis or any other IHHNV target tissues that showed Cowdry Type A inclusions (CAI), except for 1 specimen from Pond #1 that showed 1 CAI in the antennal gland (**Fig. 5c**) and 2 in the nerve cord (**Fig. 5d**). In other words, only 1 of 10 specimens positive for IHHNV with all three PCR methods showed CAI, and even then, at a very low prevalence. An example of IHHNV-CAI is shown in **Fig. 5a** from a slide of archived positive control tissue. In contrast, **Fig. 5b** shows an example of normal hematopoietic tissue seen in samples from the study ponds. Results for ISH and IHC for adjacent, serial sections from the same paraffin blocks used for histological analysis by H&E staining are given in the succeeding section.

Figure 5. Photomicrographs of tissues from IHHNV positive and negative shrimp. (a) Example photomicrograph from the positive-control specimen showing IHHNV characteristic lesions (Cowdry Type A inclusions or CAI) (black arrows), with the upper right inclusion magnified in the insert at the upper left. (b) Example photomicrograph of normal hematopoietic tissue (i.e., no CAI) from one of the study specimens even though positive for IHHNV using the IHHNV-LA method. (c, d) Photomicrographs of antennal gland and nerve cord showing Cowdry Type A inclusions or CAI, respectively, from a single specimen (Pm 1.3) from Pond #1.

3.2.4. Positive ISH results from only 1 CAI-positive specimen from Pond #1

For confirmatory ISH tests using probes based on amplicons from the two WOAH detection methods (IHHNV-309 and IHHNV-389), samples of 5 shrimp each were examined only from Pond #1 (i.e., 5/5 specimens positive using the IHHNV-LA method) and Pond #3 (all 5 specimens negative using the IHHNV-LA method). All 5 shrimp from Ponds #1 and #3 were tested and results of the ISH tests are shown in Table 2. However, photomicrographs of ISH results in **Fig. 6** show only representative photos from 2 shrimp specimens from Pond #1, one that showed CAI (Pm 1.3) and another (Pm 1.1) that did not. In **Fig. 6a-c**, the positive control specimen shows strong ISH reactions (dark staining) in the HPT while Pm 1.1 shows no positive ISH

reactions in either the HPT or subcuticular epithelium (Cep), despite giving positive PCR test results with all 3 methods used. In contrast, positive ISH reactions can be seen in both the HPT and Cep of these tissues for Pm 1.3 which gave not only positive PCR tests results with all three methods but also showed CAI (although few) by H&E staining.

Figure 6. Representative photomicrographs of *in situ* hybridization results using IHHNV-309 and IHHNV-389 DNA probes with adjacent tissue sections from 2 shrimp specimens from Pond #1 in which all 5 specimens were positive for all 3 PCR detection methods. The no-probe controls for all tests gave negative ISH results (Column 3c,f,i,l,o). (a, b) Photomicrographs of positive ISH results (dark staining) for IHHNV using both ISH probes with the archived, positive-control HPT infected with IHHNV. (d, e, g, h) Photomicrographs from specimen Pm 1.1 that gave negative ISH

results with both probes (no dark staining) in the HPT and Cep, respectively. (j, k, m, n) Representative photomicrographs from specimen Pm 1.3 showing positive ISH results (dark staining) for both probes in HPT and Cep.

In **Fig. 7**, negative ISH reactions are shown for both shrimp specimens from Pond #3 that were positive with both WOAH PCR methods but negative for both the IHHNV-LA method and for the presence of CAI. Although the proportional numbers of IHHNV positive cells in the HPT were much lower than in the Cep in Fig. 6, highly variable loads of IHHNV in different tissues of the same shrimp specimen and between shrimp specimens have been previously reported for IHHNV (Chayaburakul et al., 2005).

Figure 7. Representative photomicrographs of negative *in situ* hybridization results for adjacent tissue sections from 2 shrimp specimens (Pm 3.1 and 3.2) from Ponds #3 showing negative ISH reactions with both IHHNV-309 and -389 probes both in HPT and Cep tissues.

3.2.5. Immunohistochemistry (IHC) results paralleled the ISH results

The same 2 shrimp specimens each from Pond #1 (positive with all 3 PCR detection methods) and from Pond #3 (positive with both WOAH-PCR methods but negative for the IHHNV-LA method) were also subjected to IHC analysis using an anti-IHHNV capsid protein antibody. Results obtained using the IHC method exactly matched the ISH positive and negative results obtained using the same specimens (Table 3 above). Specifically, the positive (brown staining) and negative (no brown staining) IHC results for each of the specimens matched the positive and negative ISH results for the same shrimp specimens. Only one specimen, Pm 1.3 from Pond #1 gave positive IHC results, corresponding with positive results for all 3 of the PCR methods used. These results indicate that even the IHHNV-LA method may also give false positive test results for the presence of infectious IHHNV.

Representative photomicrographs of IHC results from the specimens Pm 1.1 and 1.3 from Pond #1 are shown in **Fig. 8**. The positive control result (dark staining) is shown in Fig. 8a while all the negative controls (no antibody) show no brown staining. Representative photomicrographs of negative IHC reactions for IHHNV are shown in the HPT (**Fig. 8c**) and Cep (**Fig. 8e**) from specimen Pm 1.1, and these matched the negative ISH results for the same specimen (Pm 1.1) in Fig. 6. Similarly, representative photomicrographs for specimen Pm 1.3 show positive IHC reactions (brown) in HPT (**Fig. 8g**) and Cep (**Fig. 8i**) that matched the positive ISH results for the same specimen (Pm 1.3) in Fig. 6.

Figure 8. Representative photomicrographs of immunohistochemistry (IHC) results for the same
2 shrimp specimens (Pm 1.1 and Pm 1.3) from Pond #1 used for ISH experiments. (a, b) Archived

positive control shrimp tissue infected with IHHNV and showing positive IHC staining (dark brown) in the HPT. (c, d) Negative IHC results for HPT of Pm 1.1. (e, f) Negative IHC results for Cep of Pm 1.1. (g, h) Positive IHC results for HPT of Pm 1.3. (i, j) Positive IHC results for Cep of Pm 1. 3.

Representative IHC results for specimens Pm 3.1 and 3.2 from Pond #3 are shown in **Fig. 9**. Aside from the positive control, none of the shrimp specimens gave positive IHC results (i.e., no dark staining) in either the HPT or Cep, and this matched the negative ISH results for the same specimens shown in **Fig. 7**.

628 **Figure 9**. Example photomicrographs of IHC results from the same two specimens (Pm 3.1 and
629 Pm 3.2) from Pond #3. (a, b) Archived positive control shrimp infected with IHHNV showing
630 positive IHC staining (dark brown) in the HPT. All other photomicrographs for both tissues show
631 negative IHC results (no brown staining).
632

3.2.6. Pathogens other than IHHNV were also present in Pond Set 1

In addition to microscopic examination for CAI, all 20 shrimp specimens in Pond Set 1were

examined for other histopathology that might be associated with retarded growth (e.g., MBV,

HPV, EHP and bacteria in the HP). No signs of MBV, HPV or EHP were seen in the HP of any of

the 20 specimens. However, some specimens showed the presence of bacterial granulomas, some

of which were melanized, indicating long presence. Other anomalies seen are summarized in **Table

4**.

Table 4. Other histopathological signs in tissues of the 20 specimens from the 4 study ponds.
Totals for each pond less than 5 indicate absence of the target tissue in the microscope slide. PMC
indicates postmortem change from poor tissue fixation of the target organ, which makes it difficult
to assess pathology. Example photomicrographs of the histopathologies are shown in
Supplementary Figure 2.

Histopathology	Pond 1 Chachoengsao	Pond 2 Krabi	Pond 3 Krabi	Pond 4 Krabi	Total	#Ponds
LO spheroids	5/5	4/4	5/5	5/5	19/19	4/4
HP bacterial lesions	0/5	PMC	3/5	0/1	3/11	1/3
Gill fouling	4/5	0/4	0/5	0/5	4/19	1/4
Gill pyknotic nuclei	2/5	3/4	4/5	0/5	9/19	3/4
ATM	4/5	PMC	0/5	0/1	4/11	1/3
Muscle lesions	0/5	4/5	1/5	5/5	10/20	¾
Total # of problems	4/6	3/6	4/6	2/6		

None of the anomalies (e.g., LO spheroids, HP lesions, etc.) listed in Table 4 gave ISH-positive or

IHC-positive signals using the IHHNV-specific probes. Thus, absence of ISH and IHC signals

indicated that the histological anomalies did not contain detectable IHHNV. Nor have such

anomalies been reported to be caused by IHHNV. Lymphoid organ (LO) spheroids develop in

response to foreign material accumulated by phagocytic hemocytes and also during viral infections

with Taura syndrome virus (TSV) (Hasson et al., 1999) and yellow head virus (YHV)

(Soowannayan et al., 2003). The spheroids caused by these pathogens are positive by ISH and IHC

specific for the respective viruses. Thus, we can assume that the spheroids in our specimens were
not formed in response to IHHNV but to other causes.

The other problems exhibited in the specimens were the presence of bacterial lesions in the HP,
pyknotic nuclei in gill tissue, muscle necrosis and gill fouling. According to the Lightner scale of
disease severity (Lightner, 1996), these other problems were present at the levels from "Trace"
(Signs of infection/infestation by internal pathogens, parasite or epicommensals present at just
above diagnostic procedure minimum detection limits) or "1" ("Signs of infection/infestation by
pathogen, parasite or epicommensal present, but at levels that may be below those needed for
clinical disease"). These levels were similar to those for IHHNV in the few specimens where
IHHNV-CAI were observed.

The presence of LO spheroids in specimens from all ponds indicated a substantial immune
response to an unknown factor in all ponds and this was compounded by additional sub-clinical
problems that varied from pond to pond, also from undetermined causes. In addition, gill fouling
is a sign of low pond water quality leading to reduced ability of shrimp to maintain clean gills, and
muscle necrosis is a sign of shrimp stress. It is important that even 1 positive specimen for any
pathogen in a random sample of 10 specimens is indicative of 26% or more prevalence with 95%
confidence in the source population (Cameron, 2002). Thus, even 1 positive sample in 6 or 5
specimens would indicate prevalences of 40 to 46% or more. As a result, the histological
information clearly indicates that conclusions about the effect of IHHNV on production from these
ponds based solely on PCR detection of IHHNV would be scientifically untenable.

3.2.7. Summary for Pond Set 1

In summary for Pond Set 1, all 4 ponds and all 5 sampled shrimp from each gave positive
PCR test results with the 2 WOAH-recommended PCR methods while only two ponds (#1 and #2)
gave additional positive tests using the IHHNV-LA method. However, only 1 of 5 specimens from
Pond #1 that gave positive PCR test results employing all three PCR methods showed the presence
of characteristic CAI lesions supported by matching positive ISH and IHC results, proving the
presence of infectious IHHNV. Thus, 4 out of 5 of these specimens gave false positive PCR test
results for infectious IHHNV using 2 WOAH-recommended detection methods and using an

IHHNV-LA detection method previously suggested as a possible PCR method to confirm the presence of infectious IHHNV (Taengchaiyaphum et al., 2022). The work herein reveals that the IHHNV-LA method can be used only as an additional control for the 2 WOAH-recommended methods, in that a negative PCR test result with the IHHNV-LA method after positive test results with one or both of the WOAH methods would indicate that the WOAH test results were false positive results. Thus, the best tests for confirmation of IHHNV presence would be to require that positive test results with either or both the WOAH methods or with the IHHNV-LA method be followed by detection of CAI or by positive ISH or IHC signals in IHHNV target tissues. Ideally, viral purification followed by bioassays would provide absolute proof. The only pond confirmed to contain IHHNV-infected shrimp gave a profitable harvest without signs of RDS.

3.3. *IHHNV relationship to production in Pond Set 2*

3.3.1. Al1 ponds studied in Pond Set 2 gave profitable harvests

Similar to Pond Set 1, all the ponds in Set 2 also gave profitable harvests (**Table 5**) including Ponds #1 and #7 that showed abnormally high CV (44% and 39%, respectively). However, the mean size for Pond #1 was 29±12 g and the median (**Fig. 10**, blue arrow) was 36 to 42g indicating that the high CV was due to a wide range of sizes shifted towards large sizes (a profile opposite to the RDS profile). On the other hand, Pond #7 did have the smallest mean size (17±7 g) and smallest median size (15-21 g) (**Fig. 10**, gray arrow), fitting the profile of an RDS pond. Pond #6 (**Fig. 10**, black bar) was intermediate, with a CV of 24%, a mean size 20±5g and a median size of 15-21g (**Fig. 10**, black arrow). Like Pond #1, this does not fit the RDS profile. In summary, only Pond #7 showed a CV and size profile that corresponded with the description of RDS in terms of size distribution (i.e., there was no shrimp deformity). Despite the differences among these ponds and despite the profile of Pond #7, profitable crops were obtained by all the farm operators. This was because the cultivations were carried out during the months of October to April which is the riskiest "Thai winter season" characterized by low temperatures conducive to outbreaks of white spot disease (WSD) caused by white spot syndrome virus (WSSV). However, during this period shrimp prices are high, attracting some farmers to attempt cultivation despite the risks.

Figure 10. Bar graph showing the size distribution of the shrimp in the 7 ponds studied in Pond Set 2. The median for 4 of the ponds (2 to 5) were in the range of 22 to 35 g, similar to Ponds #1 to #4 in Pond Set 1. Pond #7 (gray arrow) shows the smallest median size (6-14g), followed by Pond #6 (black arrow) (15-21g). Pond #1 is unusual with a median size of 36-42g and a mean of 29±12g with a CV of 44% but shifted towards large sizes (i.e., opposite to RDS).

Table 5. Shrimp growth performance and pond productivity in Pond Set 2.

Pond	Stocking density (shrimp/m²)	Duration of culture (days)	CV (%)	Mean harvested weight (g) (mean± SD)	Avg. daily weight gain (g)
1	22	160	44	29±12	0.18
2	63	164	22	33±7	0.20
3	326	166	31	32±10	0.19
4	152	166	24	28±7	0.17
5	47	120	19	32±6	0.27
6	200	174	24	20±5	0.11
7	87	132	39	17±7	0.13

3.3.2. False-positive WOAH-PCR test results for 5/7 ponds in Set 2

The 6 shrimp samples individually tested for IHHNV from each of 7 ponds in Set 2 gave positive test results with both WOAH-recommended methods (**Table 6**). However, with the IHHNV-LA method, only specimens from Pond #3 (1 in 6) and Pond #7 (3 in 6) gave positive test results. This indicated that 5 out of 7 ponds had given false positive PCR test results for infectious

743 IHHNV using the 2 WOAH methods. For Ponds #3 and #7, the results suggested that the
744 prevalence of IHHNV was higher in Pond #7 than Pond #3.

746 **Table 6.** Summary of IHHNV-DNA detection by PCR and normal histological methods with Pond
747 Set 2. Five ponds (grey background) that gave negative test results with the IHHNV-LA method
748 (gray background) had given false positive test results with the 2 WOAH methods.
749
750

Pond #	PCR methods for IHHNV		
	IHHNV-309	IHHNV-389	IHHNV-LA
1	6/6	6/6	0/6
2	6/6	6/6	0/6
3	6/6	6/6	1/6
4	6/6	6/6	0/6
5	6/6	6/6	0/6
6	6/6	6/6	0/6
7	6/6	6/6	3/6

3.3.3. IHHNV confirmation by histology in Pond Set 2

Out of the 7 ponds in Set 2, only Pond #7 gave positive test results for IHHNV by ISH, IHC

and presence of CAI in 2 of 3 specimens tested (**Table 7**). It is possible that if more samples had

been provided, that Pond #3 may have yielded some positive specimens. What can be concluded,

as with the results from Pond Set 1, is that positive test results using either ISH, IHC or CAI can

be used to confirm positive PCR tests using any of the 3 PCR methods, while a negative IHHNV-

LA test result may be useful in revealing some false positive WOAH-PCR test results by using it

as a 3rd PCR test.

769
770
771
772

773
774
Table 7. Summary of IHHNV-DNA detection by PCR and normal histological methods with Pond
Set 2. Unfortunately, due to communication problems, the results shown were from different
specimens than those whose results are shown in Table 6. Thus, the protocol differed from that of
the specimens whose results are shown for Pond Set 1 in Table 3 above where the PCR and
histology results came from the same specimens. Note that only Pond #7 (gray background) gave
positive test results for IHHNV with all three methods. ND = not determine.
781

Pond #	Histological methods for IHHNV		
	H&E CAI	ISH	IHC
1	0/3	0/3	0/3
2	0/3	0/3	0/3
3	0/3	0/3	0/3
4	0/3	ND	ND
5	0/3	ND	ND
6	0/3	ND	ND
7	2/3	2/3	2/3

3.3.4. Pathogens other than IHHNV were present in ponds from Set 2

The 3 specimens examined by PCR, ISH and IHC in Pond Set 2 (**Table 7**) were also examined

histologically in H&E-stained tissue sections for the presence of pathologies other than IHHNV

(**Table 8**). Because of the small number of shrimp sampled, even 1 positive specimen in 3 indicated

a very high prevalence (64% or more) in the source population. Similar to Pond Set 1, all samples

from every pond (total 3 x 7 = 21) showed the presence of lymphoid organ spheroids indicative of

an immune response by phagocytic cells (Hasson et al., 1999) but not a response to IHHNV

because of the negative ISH and IHC reactions in the LO to both types of IHHNV probes.

801
802

Table 8. Other histopathological signs in shrimp tissues from 7 ponds of Pond Set 2. Note that samples (grey background) from Ponds 1, 2 and 3 gave high estimated prevalence for HPV (gray background) and from Pond 7 gave a high estimated prevalence for MBV (gray background), both of which are known to be associated with retarded growth in *P. monodon*. Example photomicrographs of the histopathologies are shown in Supplementary Figures 3.

Histopathology	Pond 1 Phuket	Pond 2 Phuket	Pond 3 Phuket	Pond 4 Phuket	Pond 5 Phuket	Pond 6 Phuket	Pond 7 Phang Nga	Total	# Ponds
LO spheroids	3/3	3/3	3/3	3/3	3/3	3/3	3/3	19/21	7/7
HP bacterial lesions	0/3	0/3	0/3	1/3	0/3	1/3	1/3	3/21	3/7
HP hemocyte aggregation	0/3	0/3	1/3	1/3	3/3	1/3	0/3	6/21	4/7
HP sloughing cells	0/3	0/3	0/3	0/3	3/3	0/3	1/3	4/21	2/7
HPV	2/3	1/3	1/3	0/3	0/3	0/3	0/3	3/21	3/7
MBV	0/3	0/3	0/3	0/3	0/3	0/3	1/3	1/21	1/7
Total # problems	2/6	2/6	3/6	3/6	3/6	3/6	4/6		

Present in the HP were granulomas suggestive of bacterial infections (4/7 ponds), hemocyte aggregation (4/7 ponds) indicating an immune response, and cell sloughing (2/7 ponds) similar to that caused by acute hepatopancreatic necrosis disease (AHPND). In addition, Ponds #1, #2 and #3 gave high estimated prevalences for HPV and Pond #7 gave a high prevalence for MBV, both of which are known to be associated with retarded growth in *P. monodon* (Flegel et al., 2002; Flegel, 2006). Pond #6 also had a median of relatively small shrimp sizes, but was negative for IHHNV by all tests and also negative for HPV and MBV, clearly showing that retarded growth can arise from causes other than IHHNV. In contrast, Pond #3 positive for IHHNV by the IHHNV-LA method might have been confirmed for IHHNV by histology if more samples had been taken, but it showed a normal to better than normal size profile. Each pond exhibited from 2 to 4 of these 6 histological anomalies. Clearly, any proposal regarding the shrimp production in these study ponds based solely on IHHNV-PCR detection results and ignorant of the presence or absence of other possible pathologies would be scientifically unsupportable. This is especially important when one considers that HPV and MBV are both known to cause retarded growth in *P. monodon* (Flegel et al., 2002; Flegel, 2006). Overall, the histological analysis of the shrimp in Pond Set 2 was worse than that in Pond Set 1 probably due to off-season cultivation.

According to the Lightner scale of disease severity (Lightner, 1996), these other problems were present at the levels from "1" ("Signs of infection/ infestation by pathogen, parasite or epicommensal present, but at levels that may be below those needed for clinical disease") to "2" ("Moderate signs of infection/infestation as shown by low to moderate numbers of parasite or epicommensal, or by number and severity of pathogen caused lesions. Prognosis is for possible production losses and/or slight increases in mortality if no treatment (if treatable) or management change is applied."). In contrast, the IHHNV lesions (when present) were at the "Trace" level.

3.3.5. Summary for Pond Set 2

In summary for Pond Set 2, all 7 ponds and all 6 sampled shrimp from each gave positive PCR test results with the 2 WOAH-recommended PCR methods while only two ponds (#3 and #7) gave additional positive tests using the IHHNV-LA method. These results alone were sufficient to reveal that Ponds #1, #2, #4, #5 and #6 (5/7 = 71%) had given false positive PCR test results for IHHNV using the 2 WOAH methods. In addition, Pond #3 with only 1 in 6 specimens positive and Pond #7 with only 3 in 6 positive by the IHHNV-LA method revealed that many shrimp in each of the ponds were not infected with IHHNV. For the 3 additional specimens from each pond examined histologically, IHHNV infection could be confirmed in only 2 specimens from Pond #7, leaving the status of Pond #3 (1 in 6 positive with the IHHNV-LA test) open to question due to the low number of samples tested by histology.

4. COMBINED SUMMARY FOR BOTH POND SETS 1 and 2

For both Pond Sets 1 and 2, any specimen giving positive test results using all three PCR test methods was not sufficient to confirm an IHHNV infection. We now believe that no single PCR method or multiple PCR method can confirm IHHNV infection. However, a positive test with any one of the three PCR methods together with a positive test with any one of the three histological methods was sufficient to confirm the presence of IHHNV infection in a pond. In addition, a positive test result with either or both WOAH PCR methods but a negative result with the IHHNV-LA method served as a negative test result for the presence of IHHNV infection. In addition to the PCR results, the histological analysis in Pond Set 1 revealed less serious pathologies than those in Pond Set 2 where additional HPV and MBV infections plus sloughing of HP tubule epithelial cells characteristic of AHPND were observed. This was probably due to off-season cultivation in Pond

Set 2. Overall, the results for Pond Sets 1 and 2 were parallel with respect to IHHNV detection
and lack of direct impact on productivity.

5.   DISCUSSION

5.1. IHHNV false positive PCR results frequently arise

ISH reactions for IHHNV were negative for 3/5 specimens from Pond #1 that gave PCR
positive results for IHHNV using all 3 methods. It was possible that positive PCR results but
negative ISH and IHC results in those 3 specimens arose because the viral loads were too low to
give ISH and IHC signals. This was our former belief for shrimp positive for viral pathogens
detectable only with nested PCR and negative with histochemical methods. However, the results
in this study for both Pond Set 1 and Pond Set 2 revealed a very good correlation between positive
ISH and IHC results with results from normal histological analysis using H&E staining to detect
CAI characteristic for IHHNV. An absolute, final answer to the question of infectivity would be
obtained only by preparation of purified viral extracts followed by bioassays with SPF shrimp.
Lack of transmission would indicate the absence of infectious IHHNV in the extracts.

We believe it is most likely that the positive PCR results that could not be confirmed by positive
results for CAI, ISH and IHC arose from endogenous viral elements (EVE) of IHHNV that contain
the primer target sequences as has previously been reported (Saksmerprome et al., 2011). The
presence of EVE for both IHHNV and white spot syndrome virus (WSSV) in shrimp has been
confirmed (see Flegel, 2020 for a review).

EVE with high sequence identity to infectious IHHNV were recently found (Taengchaiyaphum et
al., 2022) in pseudochromosome 7 (PC7) of a published draft genome of *P. monodon*
(Uengwetwanit et al., 2021). Some of these EVE showed 100% sequence identity to infectious
IHHNV Type 1 with sequences that matched 100% with existing types of IHHNV at GenBank
and especially with the targets of the IHHNV-309 and -389 methods. PCR tests with archived
DNA from the genome project using the IHHNV-309 and -389 methods gave positive test results.
However, the same DNA was negative for IHHNV using the IHHNV-LA method. Additional PCR
testing confirmed that the targets for the methods were in the genomic material and not in

contaminating IHHNV DNA. Thus, a DNA extract from the specimen used for sequencing would

have given a positive PCR test result for infectious IHHNV, even though it was not infected.

Similar EVE in mosquitoes have been shown to be protective against their homologous viruses

(Suzuki et al., 2020). Thus, eliminating EVE in breeding stocks to avoid false positive PCR test

results and avoid trade restrictions might have a negative impact on farmed shrimp production if

the eliminated EVE are protective. The same arguments may apply to EVE for other shrimp viruses

such as TSV and WSSV that are also included in the WOAH list of controlled crustacean diseases.

These precautions regarding interpretation of PCR test results have been discussed previously

(Alday-Sanz et al., 2020; Saksmerprome et al., 2022).

In the end, we were forced to accept from our results herein that other than using sets of 7 to 10

overlapping primers pairs to cover the whole IHHNV genome (Chayaburakul et al., 2005;

Saksmerprome et al., 2011), it would be impossible to be certain using only 1 to 3 primer sets that

focus on short regions of the genome whether amplicons arose from IHHNV or EVE derived from

it. Instead, we now believe that obtaining a positive test result using any of the PCR methods

recommended by WOAH (or any equivalent) must be confirmed by a histological test (CAI, ISH

or IHC). One might propose that the IHHNV-LA method be used as a temporary check to screen

WOAH positive specimens if histological analysis is not possible. This might reveal about half of

the potential false positives and reduce unjust rejection or destruction of at least some farmed

shrimp products. On the other hand, of the 10 shrimp that gave positive PCR test results with the

IHHNV-LA method from Pond set 1, only 1/10 showed CAI, so there would still be a significant

risk of rejections based on false positive test results. On the other hand, the use of 7 to 10

overlapping primer pairs to cover the whole IHHNV genome (Saksmerprome et al., 2011) would

be an alternate choice.

5.2. Histopathology is required to confirm IHHNV detection by PCR

This study was carried out using shrimp (*P. monodon*) cultivation ponds that had been

selected solely on the basis that sampled shrimp specimens had given positive test results using

two WOAH-recommended PCR methods for detection of IHHNV infections. Such positive PCR

test results are often used to justify the rejection of imported shrimp products and result in serious

negative economic impacts. Here we have shown that rejections based solely on the use of either

one or both of the recommended WOAH-PCR methods would be unjustified. With 11 ponds tested

in the study, presence of IHHNV could be confirmed in only 2 ponds (2/11 = 18%) with 1 pond

open to question (3/11 = 27%). This suggests that economically important decisions might be made

based on false positive PCR test results 73 to 82% of the time.

As early as 2011 (Saksmerprome et al., 2011; 2022), it was revealed that fragments of the genome

of existing types of IHHNV occur commonly in the shrimp genome and may collectively cover

almost the whole genome (Taengchaiyaphum et al., 2022). Thus, individual specimens may

potentially give false positive PCR test results for the presence of IHHNV for any target sequence.

Since around 2012, such viral genome fragments have been referred to as endogenous viral

elements (EVE) (Feschotte & Gilvert 2012). Recent, analysis of the whole genome of *P. monodon*

has revealed clusters of scrambled EVE from IHHNV on pseudochromosome 7 that can give false

positive PCR test results for existing types of infectious IHHNV (Taengchaiyaphum et al., 2020).

EVE of extant types of white spot syndrome virus (WSSV) have also been reported in *P. monodon*

(Utari et al., 2017). Warnings regarding the propensity of EVE to give false negative PCR and RT-

PCR test results (Saksmerprome et al., 2011, 2022; Alday-Sanz et al., 2020) for viral infections

have been largely ignored.

Of the 3 PCR methods used in this study, none alone or in combination with the other PCR methods

were sufficient to confirm the presence of IHHNV infections. Due to the existence of highly

variable EVE profiles in individual shrimp specimens, confirmation required additional

histological examination using H&E-stained tissues, ISH or IHC. Thus, we recommend that the

WOAH diagnostic manual for shrimp viral diseases be modified to specify that no viral infection

can be confirmed in shrimp by one or more positive PCR test results without additional histological

analysis to demonstrate the detection of characteristic IHHNV lesions or of positive ISH or IHC

results. Based on the clusters of scrambled EVE from IHHNV on pseudochromosome 7 described

by Taengchaiyaphum et al. (2020), it is possible that even using a set of 7-10 overlapping primer

sets to cover the whole IHHNV genome might give a false positive test result.

5.3. Histopathology required to determine the true impact of IHHNV infection

In the introduction section, we reviewed the history of IHHNV infection in *P. monodon* and showed that valid reports reveal no significant negative impact on farm production (Flegel et al., 2004; Withyachumnarnkul et al., 2006). This was also the conclusion recently reported for cultivated *P. vannamei* in Ecuador and Peru (Aranguren et al., 2022). For Pond Set 1 in this study, only Pond #1 contained shrimp with confirmed IHHNV infections, and its production did not differ significantly from that of the other 3 ponds in Set 1 that were negative for IHHNV. None of the ponds exhibited gross signs associated with IHHNV (e.g., RDS).

In Pond Set 2, only Pond #7 was fully confirmed to contain shrimp infected with IHHNV and the shrimp size distribution for this pond was abnormal and growth resembled that of *P. vannamei* exhibiting RDS deformities, which the *P. monodon* did not exhibit. If this pond had been tested for IHHNV by PCR only and not subjected to histological analysis, an unwary examiner might have concluded that the abnormal size distribution was caused by IHHNV. However, this conclusion would not only be contrary to the history of IHHNV in *P. monodon* but also ignorant of the presence of LO spheroids, bacterial lesions, HP cell sloughing and the presence of MBV (associated with retarded growth in *P. monodon*). The co-occurrence of these 4 pathologies in addition to IHHNV indicated a very poor overall health status for the shrimp in Pond #7. Failure to consider such issues and failure to carry out a full health analysis but instead to focus simply on PCR test results can result in flawed publications claiming negative impacts on shrimp production caused by any pathogen. Misuse of PCR detection as the single criterion in assigning IHHNV as the cause for problems with cultivated shrimp has been particularly frequent. This is most important when shrimp mortality is part of the disease expression profile since death is not a characteristic of IHHNV in shrimp other than *P. stylirostris*. Thus, Pond #7 from Pond Set 2 is an excellent example of the perils involved in using PCR detection alone to implicate IHHNV as the cause of disease expression without backup histological analysis.

5.4. Seedstock impacts on production must also be considered

In this study, comparison of the harvest outcomes for the shrimp ponds studied in Pond Set 1 and Pond Set 2 could not be made without considering farm management issues including details related to cultivation season, farm location and infrastructure, source of seedstock (post larvae),

feed and water management, etc. Already addressed was the issue of poorer health of the shrimp
in the off-season farms, but other management details were not investigated in our study. However,
from Pond Set 2, the presence of HPV in Ponds #1, #2 and #3 and MBV in Pond #7 suggested that
viral infections might have occurred from the surrounding environment after pond stocking.
However, our previous experience has shown that these viruses are most often introduced by PL
used to stock the ponds. The farms that use genuine SPF stocks free of HPV and MBV do not
develop these infections in rearing ponds. Instead, unscreened PL derived from non-SPF, captured
broodstock or false-SPF-derived PL from unscrupulous PL suppliers are available in Thailand at
lower prices than genuine SPF-derived PL. These examples reinforce the importance for farmers
to be aware of the prime needs of using SPF-derived PL and recommended biosecurity measures
during cultivation.

5.5. The WOAH reporting for shrimp viruses needs to be modified

As stated in the Introduction, it is generally recognized in Thailand that IHHNV has had
no significant negative impact on the production of cultivated *P. monodon* or *P. vannamei* since
around 2010. Thus, it is not normally tested for or reported by farmers. On the other hand, samples
from shrimp farms are sometimes submitted to the Thai competent authority (passive surveillance)
to be tested for other reasons such as clearance before export. If such shrimp samples are tested
for IHHNV by PCR for any reason and give a positive result, Thailand's WTO obligations require
that WOAH be notified, because IHHNV is a listed crustacean disease. Clearly, a report to WOAH
based solely on PCR detection using the WOAH recommended method with the specimens
described here for Pond #3 in Pond Set 1 would have constituted an unintended false positive
report. In addition, the confirmed detection of IHHNV from Pond #1 would also be reported, but
the report would not be accompanied by a note indicating that detection was not associated with
any significant impact on production. This is important, because a key characteristic of a WOAH-
listed disease is that it has a significant, negative economic impact on production.

1006

1007 This is a very important issue, given the current knowledge showing that shrimp are capable of a
1008 specific, adaptive immune response based on nucleic acids that can give rise to EVE that are
1009 heritable if they occur in germ cells (Flegel, 2020; 2022). As a result, shrimp that survive a viral
1010 infection give rise to a vast variety of offspring, each with an individual EVE profile such that

protective EVE combinations become naturally selected in nature over time or can be purposefully selected in shrimp breeding programs (Alday-Sanz et al., 2020). The eventual result is that disease outbreaks caused by the relevant pathogens decline over time as a result of genetic tolerance and development of effective prevention protocols such that the negative economic impact may decline sufficiently to justify WOAH delisting. In order to monitor this decline, it would be necessary that obligatory country reports of listed shrimp pathogens be accompanied by an estimate of any production losses associated with each report. One might propose that listing does no harm, even if there is no negative economic impact. However, this is a naïve position. For example, the cost of unnecessary PCR testing for a listed pathogen can be high and the losses arising from tests that result in unjust rejection of products can also be very high.

6. CONCLUSIONS

Our research has shown that confirmation of IHHNV infection in shrimp can be confirmed by PCR detection only when combined with histopathological analysis by any 1 of 3 histological methods (i.e., H&E, ISH or IHC staining) or by H&E plus ISH or IHC. It cannot be confirmed by single or multiple PCR. There were no production losses in the sampled ponds positive for IHHNV. Only 2 of 11 ponds contained shrimp confirmed for IHHNV infection. These results correspond to the lack of reliable reports since 2010 on significant negative impacts of IHHNV on commercially farmed *P. monodon* or *P. vannamei* in Thailand. Similarly, there are no reports from the world capture fishery for these shrimp species, in areas where IHHNV is known to occur. Many reports claiming losses to IHHNV focus largely or solely on PCR detection results without confirmation by histological analysis or by bioassays and without investigation for other pathogens or without realistic evaluation of actual farm losses. Furthermore, recent reports that claim losses to IHHNV in hatcheries, experimental shrimp or on farms could have been avoided by use of IHHNV-negative broodstock and/or stocking of PL free of IHHNV combined with simple biosecurity. At the same time, we have found no reliable reports of significant losses to IHHNV from commercial cultivation ponds of the world's major shrimp producers since 2010. A thorough investigation regarding any significant negative impact on farmed shrimp production from IHHNV is needed to reassess whether it fits the criteria required to be listed by WOAH. It is our opinion that the massive expenditures required for needless PCR testing and the current losses suffered via

1041 trade restrictions currently justified by WOAH listing are now the only significant negative
1042 economic impacts of IHHNV.

1043

Acknowledgements

This work was funded by the National Research Council of Thailand (NRCT): High-Potential
Research Team Grant Program, grant no. N42A650869 (to K. Sritunyalucksana), Mahidol
University under the New Discovery and Frontier Research Grant 2023 (Grant no. FF-056/2566)
and the NSRF via the Program Management Unit for Human Resources & Institutional
Development, Research and Innovation (Grant No: B05F640137) and the joint Chinese Academy
of Science and National Science and Technology Development Agency (CAS-NSTDA) project
(Grant no. P2350044). We also thank to the Agricultural Research Development Agency (Public
Organization), contract no. PRP6605031130, (to J. Kongkumnerd).

1053

1054 **Conflicts of interest**

1055 None

1056

1057 **Ethics statement**

1058 This work followed Thailand's laws for ethical animal care under the Animal for Scientific
1059 83 Purposes ACT, B.E. 2558 (A.D. 2015) under project approval number BT-Animal document
1060 no. BT37/2563) from the National Center for Genetic Engineering and Biotechnology (BIOTEC),
1061 National Science and Technology Development Agency (NSTDA), Thailand.

1062

1063 **Author contributions**

1064 KS, PS, JS, JT, NS, ST Methodology, Investigation, Data curation, Writing-Original draft
1065 preparation, KS, RV, CC, TWF, ST Conceptualization, supervision, manuscript writing and final
1066 approval of the manuscript.

1067

REFERENCES

Alday-Sanz, V., Brock, J., Flegel, T.W., McIntosh, R., Bondad-Reantaso, M.G., Salazar, M.,
Subasinghe, R., 2020. Facts, truths and myths about SPF shrimp in Aquaculture. Rev.
Aquac. 12(1), 76-84. https://doi.org/10.1111/raq.12305

Aranguren Caro, L.F., Gomez-Sanchez, M.M., Piedrahita, Y., Mai, H.N., Cruz-Flores, R.,
Alenton, R.R.R., Dhar, A.K., 2022. Current status of infection with infectious hypodermal
and hematopoietic necrosis virus (IHHNV) in the Peruvian and Ecuadorian shrimp industry.
PLoS One. 17, e0272456.

Bell, T.A., Lightner, D.V., 1984. IHHN virus: Infectivity and pathogenicity studies in *Penaeus*
stylirostris and *Penaeus vannamei*. Aquaculture 38(3), 185-194. https://doi.org/10.1016/00
44-8486(84)90142-X

Bell, T.A., Lightner, D.V., 1988. A handbook of normal shrimp histology. World Aquaculture
Society, Baton Rouge, LA.

Bray, W., Lawrence, A., Leung-Trujillo, J., 1994. The effect of salinity on growth and survival of
Penaeus vannamei, with observations on the interaction of IHHN virus and salinity.
Aquaculture 122(2-3), 133-146.

Brock, J.A., Main, K.L., 1994. Guide to the Common Problems and Diseases of cultured *Penaeus*
vannamei. The Oceanic Institute, Makapuu Point, P.O. Box 25280, Honolulu, Hawaii, USA.

Cameron, A., 2002. Survey toolbox for aquatic animal diseases; a practical manual and software
package. Australian Centre for International Agricultural Research, Canberra.

Carr, W.H., Sweeney, J.N., Nunan, L., Lightner, D.V., Hirsch, H.H., Reddington, J.J., 1996. The
use of an infectious hypodermal and hematopoietic necrosis virus gene probe serodiagnostic
field kit for the screening of candidate specific pathogen-free *Penaeus vannamei*
broodstock. Aquaculture 147(1-2), 1-8.

Chaijarasphong, T., Munkongwongsiri, N., Stentiford, G.D., Aldama-Cano, D.J., Thansa, K.,
Flegel, T.W., Sritunyalucksana, K., Itsathitphaisarn, O., 2021. The shrimp microsporidian
Enterocytozoon hepatopenaei (EHP): Biology, pathology, diagnostics and control. J.
Invertebr. Pathol. 186, 107458. https://doi.org/10.1016/j.jip.2020.107458

Chayaburakul, K., Lightner, D.V., Sriurairattana, S., Nelson, K.T., Withyachumnarnkul, B., 2005.
Different responses to infectious hypodermal and hematopoietic necrosis virus (IHHNV) in
Penaeus monodon and *P. vannamei*. Dis. Aquat. Organ. 67(3), 191-200.

Cotmore, S.F., Agbandje-McKenna, M., Canuti, M., Chiorini, J.A., Eis-Hubinger, A.-M., Hughes,
J., Mietzsch, M., Modha, S., Ogliastro, M., Pénzes, J.J., Pintel, D.J., Qiu, J., Soderlund-
Venermo, M., Tattersall, P., Tijssen, P., Consortium, I.R., 2019. ICTV Virus Taxonomy
Profile: *Parvoviridae*. J. Gen. Virol. 100(3), 367-368.

Feschotte, C., Gilbert, C., 2012. Endogenous viruses: insights into viral evolution and impact on
host biology. Nat. Rev. Genet. 13(4), 283-296. https://doi.org/10.1038/nrg3199

Flegel, T.W., Nielsen, L., Thamavit, V., Kongtim, S., Pasharawipas, T., 2004. Presence of multiple
viruses in non-diseased, cultivated shrimp at harvest. Aquaculture. 240, 55-68.

Flegel, T.W., 2006. Detection of major penaeid shrimp viruses in Asia, a historical perspective
with emphasis on Thailand. Aquaculture 258(1), 1-33.

Flegel, T.W., 2020. Research progress on viral accommodation 2009 to 2019. Dev. Comp.
Immunol. 112, 103771. https://doi.org/10.1016/j.dci.2020.103771

Flegel, T.W., 2022. Viral accommodation: A New mechanism for disease control in shrimp
aquaculture, in: Alday-Sanz, V. (Ed.), The Shrimp Book II. 5M Books Ltd., United
Kingdom, pp. 427-445.

Flegel, T.W., Nielsen, L., Thamavit, V., Kongtim, S., Pasharawipas, T., 2004. Presence of multiple
viruses in non-diseased, cultivated shrimp at harvest. Aquaculture 240(1), 55-68.

Hasson, K.W., Lightner, D.V., Mohney, L.L., Redman, R.M., Poulos, B.T., White, B.M., 1999.
Taura syndrome virus (TSV) lesion development and the disease cycle in the Pacific white
shrimp *Penaeus vannamei*. Dis. Aquat. Organ. 36(2), 81-93.

Hernández-Ruíz, H., Montaldo, H.H., Bustos-Martínez, J., Campos-Montes, G.R., Castillo-Juárez,
H., 2020. Heritability and genetic correlations for infectious hypodermal and hematopoietic
necrosis virus load, body weight at harvest, and survival rate in Pacific white shrimp
(*Litopenaeus vannamei*). J. World Aquac. Soc. 51(1), 312-323. https://doi.org/
10.1111/jwas.12664

Jagadeesan, V., Praveena, P.E., Otta, S.K., Jithendran, K.P., 2019. Classical Runt Deformity
Syndrome Cases in Farmed *Penaeus vannamei* Along the East Coast of India. J. Coast. Res.
86(sp1), 107-111.

Jitrakorn, S., Arunrut, N., Sanguanrut, P., Flegel, T.W., Kiatpathomchai, W., Saksmerprome, V.,
2016. In situ DIG-labeling, loop-mediated DNA Amplification (ISDL) for highly sensitive
detection of infectious hypodermal and hematopoietic necrosis virus (IHHNV). Aquaculture
456, 36-43. https://doi.org/10.1016/j.aquaculture.2016.01.023

Kalagayan, G., Godin, D., Kanna, R., Hagino, G., Sweeney, J., Wyban, J., Brock, J., 1991. IHHN
virus as an etiological factor in runt-deformity syndrome of juvenile *Penaeus vannamei*
cultured in Hawaii. J. World Aquacul. Soc. 22, 235-243.

Kiatmetha, P., Chotwiwatthanakun, C., Jariyapong, P., Santimanawong, W., Ounjai, P., Weerachatyanukul, W., 2018. Nanocontainer designed from an infectious hypodermal and hematopoietic necrosis virus (IHHNV) has excellent physical stability and ability to deliver shrimp tissues. PeerJ 6, e6079. https://doi.org/10.7717/peerj.6079

Lightner, D.V., 1996. A handbook of shrimp pathology and diagnostic procedures for diseases of cultured penaeid shrimp, illustrated ed. World Aquaculture Society, Baton Rouge, LA (USA)

Lightner, D.V., Redman, R.M., Bell, T.A., 1983. Infectious hypodermal and hematopoietic necrosis, a newly recognized virus disease of penaeid shrimp. J. Invertebr. Pathol. 42(1), 62-70.

Limsuwan, C., Chanratchakool, P., 2004. Shrimp Culture industry in Thailand: Components of Thai shrimp industry, in: Shrimp Culture industry in Thailand. Faculty of Fisheries, Kasetsart University; Department of Fisheries; National Research Council of Thailand, pp. 1-6.

Marín-Riffo, M.C., Raadsma, H.W., Jerry, D.R., Coman, G.J., Khatkar, M.S., 2021. Bioeconomic modelling of hatchery, grow-out and combined business of Australian black tiger shrimp *Penaeus monodon* farming. Rev. Aquac. 13(3), 1695-1708. https://doi.org/10.1111/raq.12541

Mendoza-Cano, F., Enríquez-Espinoza, T., Encinas-García, T., Sánchez-Paz, A., 2014. Prevalence of the infectious hypodermal and hematopoietic necrosis virus in shrimp (*Penaeus vannamei*) broodstock in northwestern Mexico. Prev. Vet. Med. 117(1), 301-304. https://doi.org/10.1016/j.prevetmed.2014.09.006

Pénzes, J.J., Söderlund-Venermo, M., Canuti, M., Eis-Hübinger, A.M., Hughes, J., Cotmore, S.F., Harrach, B., 2020. Reorganizing the family Parvoviridae: a revised taxonomy independent of the canonical approach based on host association. Arch. Virol., 1-14.

Primavera, J.H., Quinitio, E.T., 2000. Runt-Deformity Syndrome in Cultured Giant Tiger Prawn *Penaeus monodon*. J. Crustac. Biol. 20(4), 796-802. https://doi.org/10.1163/20021975-99990101

Pruder, G.D., 1995. High health shrimp systems: seed supply-theory and practice, Swimming through troubled water, Proceedings of the special session on shrimp farming, Aquaculture'95. World Aquaculture Society, San Diego, California.

Saksmerprome, V., Jitrakorn, S., Chayaburakul, K., Laiphrom, S., Boonsua, K., Flegel, T.W.,
2011. Additional random, single to multiple genome fragments of *Penaeus stylirostris*
densovirus in the giant tiger shrimp genome have implications for viral disease diagnosis.
Virus Res. 160(1), 180-190. https://doi.org/10.1016/j.virusres.2011.06.010

Saksmerprome, V., Jitrakorn, S., Arunrut, N., Kiatpathomchai, W., Felgel, T.W., 2022.
Geologically recent shrimp endogenous viral elements and their implications for shrimp
disease control and trade. in: Alday-Sanz, V. (Ed.), The Shrimp Book II. 5m Books Ltd.,
United Kingdom, pp. 547-561.

Sanguanrut, P., Munkongwongsiri, N., Kongkumnerd, J., Thawonsuwan, J., Thitamadee, S.,
Boonyawiwat, V., Tanasomwang, V., Flegel, T.W., Sritunyalucksana, K., 2018. A cohort
study of 196 Thai shrimp ponds reveals a complex etiology for early mortality syndrome
(EMS). Aquaculture 493, 26-36. https://doi.org/10.1016/j.aquaculture.2018.04.033

Sanguanrut, P., Thaiue, D., Thawonsuwan, J., Aldama-Cano, D.J., Flegel, T.W., Sritunyalucksana,
K., 2022. The lymphoid organ (LO) is an additional, prime target for decapod iridescent
virus 1 (DIV1) in *Penaeus vannamei*. Aquaculture 547, 737482.

Sellars, M.J., Cowley, J.A., Musson, D., Rao, M., Menzies, M.L., Coman, G.J., Murphy, B.S.,
2019. Reduced growth performance of Black Tiger shrimp (*Penaeus monodon*) infected
with infectious hypodermal and hematopoietic necrosis virus. Aquaculture 499, 160-166.
https://doi.org/10.1016/j.aquaculture.2018.09.032

Soowannayan, C., Flegel, T.W., Sithigorngul, P., Slater, J., Hyatt, A., Cramerri, S., Wise, T.,
Crane, M.S., Cowley, J.A., McCulloch, R.J., Walker, P.J., 2003. Detection and
differentiation of yellow head complex viruses using monoclonal antibodies. Dis. Aquat.
Organ. 57(3), 193-200. https://doi.org/10.3354/dao057193

Suzuki, Y., Baidaliuk, A., Miesen, P., Frangeul, L., Crist, A.B., Merkling, S.H., Fontaine, A.,
Lequime, S., Moltini-Conclois, I., Blanc, H., van Rij, R.P., Lambrechts, L., Saleh, M.-C.,
2020. Non-retroviral Endogenous Viral Element Limits Cognate Virus Replication in
Aedes aegypti Ovaries. Curr. Biol. 30(18), 3495-3506.e3496. https://doi.org/10.101
6/j.cub.2020.06.057

Taengchaiyaphum, S., Wongkhaluang, P., Sittikankaew, K., Karoonuthaisiri, N., Flegel, T.W.,
Sritunyalucksana, K., 2022. Shrimp genome sequence contains independent clusters of

ancient and current Endogenous Viral Elements (EVE) of the parvovirus IHHNV. BMC
Genomics 23(1), 565. https://doi.org/10.1186/s12864-022-08802-3

Tang, K.F., Durand, S.V., White, B.L., Redman, R.M., Pantoja, C.R., Lightner, D.V., 2000.
Postlarvae and juveniles of a selected line of *Penaeus stylirostris* are resistant to infectious
hypodermal and hematopoietic necrosis virus infection. Aquaculture 190(3-4), 203-210.

Tang, K.F., Lightner, D.V., 2006. Infectious hypodermal and hematopoietic necrosis virus
(IHHNV)-related sequences in the genome of the black tiger prawn *Penaeus monodon* from
Africa and Australia. Virus Res. 118(1-2), 185-191.

Tang, K.F., Navarro, S.A., Lightner, D.V., 2007. PCR assay for discriminating between infectious
hypodermal and hematopoietic necrosis virus (IHHNV) and virus-related sequences in the
genome of *Penaeus monodon*. Dis. Aquat. Organ. 74(2), 165-170.

Uengwetwanit, T., Pootakham, W., Nookaew, I., Sonthirod, C., Angthong, P., Sittikankaew, K.,
Rungrassamee, W., Arayamethakorn, S., Wongsurawat, T., Jenjaroenpun, P., Sangsrakru,
D., Leelatanawit, R., Khudet, J., Koehorst, J.J., Schaap, P.J., Martins dos Santos, V., Tangy,
F., Karoonuthaisiri, N., 2021. A chromosome-level assembly of the black tiger shrimp
(*Penaeus monodon*) genome facilitates the identification of growth-associated genes. Mol.
Ecol. Resour. 21(5), 1620-1640. https://doi.org/https://doi.org/10.1111/1755-0998.13357

West E, Conacher C, Dexter J, Lee P, Heidenreich M, B., P., 2019. Aquaculture in Moreton Bay,
in: Tibbetts, I.R., Rothlisberg, P.C., Neil, D.T., Homburg, T.A., Brewer, D.T., Arthington,
A.H. (Eds.), Moreton Bay Quandamooka & Catchment: *Past, present, and future*. The
Moreton Bay Foundation, Brisbane, Australia.

Withyachumnarnkul, B., Chayaburakul, K., Lao-Aroon, S., Plodpai, P., Sritunyalucksana, K.,
Nash, G., 2006. Low impact of infectious hypodermal and hematopoietic necrosis virus
(IHHNV) on growth and reproductive performance of *Penaeus monodon*. Dis. Aquat.
Organ. 69(2-3), 129-136. https://doi.org/10.3354/dao069129

WOAH, 2008. Infection with infectious hypodermal and haematopoietic necrosis virus, in:
Aquatic Animal Health Code. World Organisation for Animal Health, the University of
California, USA, p. 307.

WOAH, 2019a. Infection with infectious hypodermal and haematopoietic necrosis virus, in:
Aquatic Animal Health Code. World Organisation for Animal Health, Paris, France, p. 324.

1225 WOAH, 2019b. Infection with infectious hypodermal and haematopoietic necrosis virus, in:

1226    Manual of Diagnostic Tests for Aquatic Animals. World Organisation for Animal Health,

1227    Paris, France.

1228 Wyban, J.A., 2007. Domestication of pacific white shrimp revolutionizes aquaculture, Global

1229    Aquaculture Advocate. Portsmouth, NH, USA, pp. 42-44.

1230 Wyban, J.A., Swingle, J.S., Sweeney, J.N., Pruder, G.D., 1992. Development and commerical

1231    performance of high health shrimp using specific pathogen free (SPF) broodstock *Penaeus*

1232    *vannamei*, in: Wyban, J.A. (Ed.) Proceedings of the special session on shrimp farming,

1233    World Aquaculture '92. World Aquaculture Society, Baton Rouge, LA, USA, pp. 254-260.

1234 Yang, Y.T., Lee, D.Y., Wang, Y., Hu, J.M., Li, W.H., Leu, J.H., Chang, G.D., Ke, H.M., Kang,

1235    S.T., Lin, S.S., Kou, G.H., Lo, C.F., 2014. The genome and occlusion bodies of marine

1236    *Penaeus monodon* nudivirus (PmNV, also known as MBV and PemoNPV) suggest that it

1237    should be assigned to a new nudivirus genus that is distinct from the terrestrial nudiviruses.

1238    BMC Genomics 15, 628. https://doi.org/10.1186/1471-2164-15-628.