Unveiling the ecology, taxonomy and metabolic capabilities of MBA03, a potential key player in anaerobic digestion

Roser Puchol-Royo[1], Javier Pascual[1], Asier Ortega-Legarreta[1,2], Pascal Otto[3], Jeroen Tideman[4], Sjoerd-Jan de Vries[4], Christian Abendroth[5], Kristie Tanner[1], Manuel Porcar[1,6] & Adriel Latorre-Perez[1]

[1]Darwin Bioprospecting Excellence, S.L. Parc Cientific Universitat de Valencia, Paterna, Valencia, Spain.

[2]Translational Bioinformatics Unit, Navarrabiomed, Complejo Hospitalario de Navarra (CHN), Universidad Pública de Navarra (UPNA), IdiSNA, Pamplona, Spain

[3]Technische Universität Dresden, Institute of Waste Management and Circular Economy, Pirna, Germany.

[4]Bioclear Earth B.V., Groningen, The Netherlands

[5]Brandenburgische Technische Universität Cottbus-Senftenberg, Chair of Circular Economy, Cottbus, Germany

[6]Institute for Integrative Systems Biology I2SysBio (University of Valencia - CSIC), Paterna, Spain

Corresponding author: A. Latorre-Pérez (alatorre@darwinbioprospecting.com)

Abstract

Biogas, a mix of CO_2, CH_4 and small proportions of other gases, is a biofuel obtained by anaerobic digestion (AD). Biogas production is often considered a black box process, as the role and dynamics of some of the microorganisms involved remain undisclosed. Previous metataxonomic studies in the frame of the MICRO4BIOGAS project (www.micro4biogas.eu) revealed that MBA03, an uncharacterised and uncultured bacterial taxon, was very prevalent and abundant in industrial full-scale AD plants. Surprisingly, no culturable specimen or genome of this taxon has ever been reported, so its role in AD has remained unclear. In the present work, thirty samples derived from anaerobic digesters were sequenced, allowing the reconstruction of 108 metagenome-assembled genomes (MAGs) potentially belonging to MBA03. According to phylogenetic analyses and genomic similarity indices, MBA03 constitutes a new bacterial order, proposed as *Darwinibacteriales* ord. nov., which includes *Darwinibacter acetoxidans* gen. nov., sp. nov. of the family *Darwinibacteriaceae* fam. nov., along with *Wallacebacter cryptica* gen. nov., sp. nov. of the *Wallacebacteriaceae* fam. nov. Ecotaxonomic studies determined that AD processes are the main ecological niche of *Darwinibacteriales*. Moreover, metabolic predictions identified *Darwinibacteraceae* members as putative syntrophic acetate oxidising bacteria (SAOB), as they encode for the reversed Wood-Ljungdahl pathway coupled to the glycine cleavage system. This suggests that *Darwinibacteraceae* members work in collaboration with hydrogenotrophic archaea to produce methane in industrial biogas plants. Overall, our findings present *Darwinibacteriales* as a potential key player in anaerobic digestion and pave the way towards the complete characterisation of this newly described bacterial taxa.

Keywords: Metagenomics, anaerobic digestion, biogas, SAOB, MBA03, *Darwinibacteriales*

1. Introduction

The evidence supporting the need for a transition from fossil to renewable energy is overwhelming and unanimous (Yaqoob et al., 2021). In recent years, biogas has arisen as one of the leading alternatives to non-renewable fossil fuels such as natural gas. Biogas is produced by anaerobic digestion (AD) of organic material, through a complex microbiological process in which several groups of microorganisms with diverse metabolic abilities work in close collaboration to obtain methane as the main product (Schnürer, 2016). In addition to the production of renewable methane and fertiliser, biogas production contributes to the utilisation of today's global waste streams of various origins as a source of organic matter. In particular, waste from agriculture, the food industry, wastewater and municipal solid waste can be treated. (Tsang et al., 2019).

Biogas primarily consists of methane (CH_4), carbon dioxide (CO_2) and other gases like H_2S, N_2, NH_3, and H_2 in lower concentrations (Yaqoob et al., 2021). Despite the complexity of the AD process, the microbial metabolic reactions that occur in this process and that lead to the synthesis of methane can be summarized in four key steps: hydrolysis, acidogenesis, acetogenesis and methanogenesis (Schnürer, 2016). Briefly, during hydrolysis, complex macromolecules are broken down into various oligo-, di- and monomers, the final concentrations of which will depend on the biomass source (Nwokolo et al., 2020). These molecules are uptaken and used by acidogenic bacteria to produce H_2, CO_2, alcohols and volatile fatty acids (VFAs) during acidogenesis, while in acetogenesis, acetogenic bacteria synthesize acetic acid through the Wood-Ljungdahl (W-L) pathway or autotrophically from H_2 and CO_2. Subsequently, methanogens metabolize acetate, hydrogen and/or methylated compounds to produce methane, creating an equilibrium and a co-dependence between bacteria and archaea (Nwokolo et al., 2020; Vanwonterghem et al., 2016). Two main pathways can occur in the production of methane from acetate: the first one, directly by acetoclastic methanogens, which split acetate to methane and CO_2, with the methyl group yielding methane and the carboxyl group yielding CO_2. The other pathway results from syntrophic acetate-oxidizing bacteria (SAOB) oxidizing acetate to CO_2 and H_2, which is subsequently used by hydrogenotrophic archaea to obtain methane. Thus, this pathway must be coupled with hydrogenotrophic methanogenesis (SAO-HM). Although less abundant in anaerobic digester plants, there is a third pathway, which is the methylotrophic pathway (Abendroth et al., 2015). This pathway is regarded as an ancestral pathway of methanogenesis (Wang et al., 2001), where the respective archaea are converting methylic C1 groups (Whitman et al., 2006).

The microbial community underlying AD shows very complex entangled ecologies, with the taxonomic composition additionally depending on many factors such as substrate, temperature or pH (De Vrieze et al., 2015; Lebuhn et al., 2015).

Despite the high economic potential, knowledge for the production of biogas with AD has remained stagnant, as the high variability of both the different operational parameters and the native microbial communities involved represent a challenge for its optimization and estandardization. Improving the feasibility of the process and achieving a consistent and controlled way to obtain biogas

would constitute a huge change in the energy industry. But reaching such an
ambitious goal requires gaining insights in the microbial machinery behind AD
and methanogenesis, and this goes through the study of a variety of anaerobic
digestion systems with different chemical, taxonomical and operational
parameters.

In this context, the EU-funded project MICRO4BIOGAS (https://micro4biogas.eu)
was born in 2021. The ultimate objective of the project is to improve the yield,
speed, quality and robustness of biogas production by optimising the AD
microbiome. In the frame of the project, 80 different samples were collected
from various AD plants in the Netherlands, Germany and Austria.

After metataxonomic analyses, a miscellaneous taxonomic group (i.e., MBA03)
was found to be very prevalent and abundant. Moreover, MBA03 was among the
taxa that correlated most strongly with operational (i.e., temperature) and
chemical parameters (i.e., acetic acid) (Otto et al., 2023), suggesting that it may
be a microorganism of great relevance for the AD process. MBA03 (AB114313.1)
was described for the first time by Tang et al. as an uncultured clone obtained
from a thermophilic anaerobic municipal solid-waste digester (Tang et al., 2004).
The proposal of MBA03 exclusively relied on the 16S rRNA gene sequence. Since
then, a series of metataxonomic studies have reported the presence of MBA03 in
anaerobic digester systems (Dyksma et al., 2020; FitzGerald, 2018; Laguillaumie
et al., 2022), representing up to 70% of the relative abundance in some samples
(FitzGerald, 2018). This taxon has been associated with a higher production of
methane and has been suggested as an indicator of the stability of the AD
process (Fang et al., 2022; Laguillaumie et al., 2022).

Surprisingly, MBA03 has not been properly taxonomically delimited. In fact, no
culturable specimen or genome of this taxon has ever been reported. However,
according to SILVA nr database, MBA03 represents a heterogenous group of
more than 150 16S rRNA sequences. Therefore, a deeper characterization of this
taxonomic group was urgently needed for unravelling its role in AD processes.

In this work, we used metagenomic data from MICRO4BIOGAS and other AD
studies to propose MBA03 as a member *Darwinibacteriales* ord. nov., a not-yet-
cultured bacterial taxa belonging to the class *Clostridia. Darwinibacteriales*
includes two different genera: *Darwinibacter acetoxidans* gen. nov., sp. nov., of
the *Darwinibacteriaceae* family, and *Wallacebacter cryptica* gen. nov., sp. nov.,
of the *Wallacebacteriacea* fam. nov. Based on our results, both *Darwinibacter*
and *Wallacebacter* predominantly inhabit AD environments. Furthermore, we
have inferred that *Darwinibacter* may play a key role in anaerobic digestion
processes related to biogas production, as it includes the reversed Wood-
Ljungdahl pathway coupled to the glycine cleavage system, which is required for
the synthesis of H_2 and CO_2 from acetate and the subsequent production of
methane by hydrogenotrophic methanogenesis.

2. Materials and methods

2.1. Sample description

As part of the MICRO4BIOGAS project, a total of 80 samples were collected from anaerobic digestion systems at 45 large-scale reactors with different operational conditions and feedstocks. These samples, together with the corresponding metadata, were used to investigate the chemical and taxonomic profiles (Otto et al., 2023). The first 40 samples were collected by members of the Technische Universität Dresden (TUD, Dresden, Germany) from 25 different biogas plants in Germany and Austria, while the second set of 40 samples was collected by the team members from Bioclear Earth B.V. (Groningen, Netherlands) from 20 different biogas plants in the Netherlands. A subset of 30 of these samples was selected to be analysed through metagenomic sequencing. This subset included samples enriched in MBA03, hydrolytic, acidogenic and acetogenic bacteria and/or methanogenic archaea. A full description of the samples can be found in Supplementary Table 1.

2.2. DNA extraction and sequencing

The sludge samples were washed before DNA extraction to reduce the number of inhibiting substances. Specifically, 5-10 mL of each sample was mixed with 40 mL of sterile phosphate-buffered saline (PBS). Subsequently, the samples were centrifuged at 20,000 g for 10 min and the supernatants were discarded. This process was repeated twice, and the final pellets were used for DNA extraction with the DNeasy® PowerSoil® Pro kit (QIAGEN, Germany) based on the manufacturer's instructions, with an extra incubation step at 65 °C for 10 min after adding the C1 solution to increase the efficiency of cell lysis. Qubit x1 dsDNA HS Assay kit (Qubit 4.0 Fluorometer, Thermo Fisher) was used for DNA quantification. The metagenomic DNA was randomly fragmented by sonication and DNA fragments were end polished, A-tailed, and ligated with the full-length adapters for Illumina sequencing: 5'-AGA TCG GAA GAG CGT CGT GTA GGG AAA GAG TGT AGA TCT CGG TGG TCG CCG TAT CAT T -3' (5' Adapter), and 5'- GAT CGG AAG AGC ACA CGT CTG AAC TCC AGT CAC GGA TGA CTA TCT CGT ATG CCG TCT TCT GCT TG -3' (3' adapter). PCR amplification with P5 and indexed P7 oligos was carried out, and the PCR products were purified with the AMPure XP system. Then, the libraries were checked for size distribution by an Agilent 2100 Bioanalyzer (Agilent Technologies, CA, USA), and quantified by real-time PCR. Libraries were sequenced at Novogene (UK) using the NovaSeq 6000 Illumina platform (150 bp x 2). The average sequencing depth was 45M reads/sample (min.: 35M reads; max.: 55M reads).

2.3. Metagenomic analysis

Adapters were trimmed from the raw reads using Cutadapt (v. 3.4; Martin, 2011), and a quality filtering step was performed with BBDuk (BBTools v.38.84; https://sourceforge.net/projects/bbtools/). Reads shorter than 75 bp or with a

mean quality lower than Q20 (in the PHRED scale) were discarded. Bases with a
quality below Q20 were also trimmed from both read ends. Human reads were
then removed with bowtie2 (v. 2.4.4; Langmead & Salzberg, 2012) by aligning
the reads against the human genome (GRCh38). The quality of the clean
sequences was checked with FastQC (v0.11.5;
http://www.bioinformatics.babraham.ac.uk/projects/fastqc). MEGAHIT was used
with default parameters for assembling the filtered metagenomic reads (v1.2.9;
(D. Li et al., 2015). Assembly statistics were calculated with QUAST (v. 5.0.2;
(Gurevich et al., 2013). Two different binning strategies were applied: MetaBAT
(Kang et al., 2019) and MaxBin (Wu et al., 2016). The MAGs obtained were
refined with Das Tool (Sieber et al., 2018) and classified according to their
quality: high quality (HQ, integrity ≥ 90% and contamination ≤ 5%), good quality
(GQ, integrity ≥ 80% and contamination ≤ 10%) and medium-low quality (LQ,
integrity ≤ 80% and/or contamination ≥ 10%), following the criteria of Bowers
et al., 2017 and Feng et al., 2021. After that, Centrifuge (Kim et al., 2016) was
used to study the taxonomic profile of the samples, while the resulting MAGs
were taxonomically annotated with MiGA (Rodriguez-R et al., 2018).

200 2.4. Recovery of MBA03 genomes via phylogenetic analyses

First of all, 16S rRNA sequences were extracted from the high-quality MAGs using
Prokka (v 1.14.6) (Seemann, 2014). These sequences were aligned together with
the 157 available MBA03 16S rRNA sequences from SILVA nr database (SSU
138.1 ref NR) with the SINA Aligner (v 1.2.11) (Pruesse et al., 2012), and
afterwards a phylogenetic tree was computed with FastTree (v 2.1.1) (Price
et al., 2009) and visualized with the Interactive Tree Of Life (iTOL) (Letunic &
Bork, 2021). Only the MAGs whose sequences formed a monophyletic group with
MBA03 were selected for further analysis. Due to the limitations associated to
the binning process, not all MAGs contained 16S rRNA sequences, while others
contained more than one sequence, so not all the MAGs corresponding to MBA03
could be identified by 16S rRNA.

213 2.5. MBA03 identification from phylogenomic data

After identifying the first MAGs hypothetically belonging to MBA03, a two-step
approach was followed to recover the remaining MAGs belonging to the MBA03
taxonomic group which did not contain rRNA operons.

As a first step, all the MAGs whose 16S rRNA gene formed a monophyletic group
with MBA03 according to the phylogenetic tree were annotated using the whole
genome classifier tool Genome Taxonomy Database toolkit (GTDB-tk; v 2.1.1)
(Chaumeil et al., 2022). Although GTDB does not contain any genome identified
as "MBA03", the resulting annotation was used as a first hypothesis to elucidate
which of the genomes actually corresponded to MBA03.

As a second step, all the HQ and GQ MAGs recovered in the project (n = 1053)
were also annotated GTDB-tk. In addition, a phylogenomic tree was constructed
using the UBCG pipeline (v 3.0) (Na et al., 2018). Once all the MAGs
corresponding to MBA03 were identified, genomes belonging to the MBA03 order

in the GTDB were downloaded and included in the dataset. Afterwards, genomes
from the Biogasmicrobiome repository (https://biogasmicrobiome.env.dtu.dk/),
and from (Dyksma et al., 2020) were annotated and incorporated into the
dataset.

2.6. Taxonomic delimitation of the MBA03 taxonomic clade

Prokka (v 1.14.6) (Seemann, 2014) was used again to obtain all the 16S rRNA
sequences from the MAGs and genomes identified as potential MBA03
representatives according to the previous phylogenomic analyses. To better
study the phylogenetic patterns, 16S rRNA sequences corresponding to the most
closely related taxonomic classes were set as outgroups. Sequences from class
Clostridia and *Limnochordia* were downloaded from the Living Tree Project (LTP)
and dereplicated using cd-hit (4.8.1) (W. Li & Godzik, 2006) at a 0.97 similarity
threshold. Sequences were aligned with SINA Aligner (v 1.2.11) (Pruesse et al.,
2012) and incorporated into the previous dataset. Finally, a phylogenetic tree
was computed with FastTree (Price et al., 2009), and visualized with iTOL
(Letunic & Bork, 2021). The reference sequence of the bacterial clone identified
for the first time as MBA03 was also downloaded from the NCBI (accession
number: AB114313.1) and included in the tree.

Following the same procedure as for the phylogenetic tree, genomes from
Clostridia and *Limnochordia* were downloaded from RefSeq NCBI Assembly
Database, filtering by "Assembly from Type Material" and "Complete Genomes"
(Supplementary Table 2). These genomes were analysed altogether with the
genomes identified as MBA03 and a phylogenomic tree was created with UBCG
(Na et al., 2018) and visualized with iTOL (Letunic & Bork, 2021).

2.7. Analysis of the ecological niche of MBA03

In order to explore the ecological niches of MBA03, an analysis with IMNGS
(Lagkouvardos et al., 2016) was carried out using a 97% identity cut-off. The
same sequences used for the phylogenetic analysis were dereplicated using cd-
hit (v 4.8.1) (W. Li & Godzik, 2006) at a 0.9 identity threshold. IMNGS results
were filtered with the objective of gathering all the samples by their sample type
to see which are the potentially relevant ecological niches of MBA03. As tags
describing ecological niches were very diverse and, in many cases, ambiguous, a
series of in-house Python scripts were needed to carry out a reclassification of
niche categories. Prior to this filtering, samples containing the 16S rRNA
sequence of interest in an abundance <0.1% were removed from the analyses.
To reclassify the remaining samples, two subsequent strategies were followed:
(1) description tags were filtered by the words "anaerobic", "digester", "biogas",
and "reactor", assigning all of them to the same tag: "anaerobic digestion"; (2)
the information of the remaining samples were scraped from NCBI. The output
was manually curated to determine which samples belonged to "anaerobic
digestion", and other general tags were also reassigned to more specific classes
(i.e., "metagenome" to "soil metagenome") depending on the project
description.

In order to determine the abundance of MBA03 among the samples, coverM
(https://github.com/wwood/CoverM) was used, comparing the raw reads of the
30 sequenced reactors against the genomes assigned to MBA03.

2.8. Functional annotation

MAGs belonging to MBA03 were filtered according to two criteria: completeness
> 90% and contamination < 5%. These MAGs were annotated with Bakta
(Schwengers et al., 2021) and they were analysed with Kofamscan (Aramaki et
al., 2020) and KEGG-Decoder (Graham et al., 2018).

As mentioned above, acetate as an intermediate is of particular importance
during anaerobic reactions, especially for SAOB bacteria and for
hydrogenotrophic methanogenesis. In order to elucidate if MBA03 was a SAOB,
the protein sequences of all the enzymes involved in the reversed Wood-
Ljungdahl (WL) pathway and the reversed WL pathway coupled to the glycine
cleavage system (GCS) (Supplementary Table 3) were downloaded from UniProt
(Coudert et al., 2023) (Swissprot), narrowing to all bacteria (taxonomy_id:2), and
a database was created with Diamond (Buchfink et al., 2015). A Diamond BLASTp
of the annotated protein sequences was performed against this database to
detect presence and absence of each gene in all MAGs with 50% coverage and
identity thresholds. For the whole metabolic analysis, any enzyme found in more
than 70% of the MAGs of each family was considered to be present in the
genome.

3. Results and discussion

3.1. Genomic reconstruction

MBA03 has proven abundant in a variety of anaerobic digesters (Dyksma et al.,
2020; Liu et al., 2023). Although there are more than 150 16S rRNA sequences
classified as MBA03 in the SILVA nr database (v. 138), no representative genome
of any bacteria belonging to this taxon has ever been described. Given its
importance in AD environments, this work focused on the recovery and
characterisation of the genome of this taxonomic group using 30 metagenomic
datasets derived from the MICRO4BIOGAS project and other genomic data
available in public databases.

After metagenomic sequencing, assembly and binning of sludge samples, the
metagenome-assembled genomes (MAGs) obtained were classified according to
their quality, and 1053 MAGs (600 of high quality (HQ) and 453 of good quality
(GQ)) were chosen for further analysis. After the annotation of the MAGs with
MIGA and the metagenomes with Centrifuge, in neither case, MBA03 was
reported despite being a highly abundant taxon according to the metataxonomic
results (Otto et al., 2023). This was expected as genomic databases did not
include MBA03 genomes, probably reflecting a lack of consensus on the
nomenclature of this taxonomic group.

3.2. Taxonomic analysis of MAGs

In order to identify the genomes belonging to MBA03, a phylogenetic tree was constructed using the 16S rRNA sequences extracted from the HQ MAGs and 157 sequences classified as MBA03 according to the SILVA database (SSU 138.1 ref NR) (Supplementary Figure 1). The sequences from SILVA formed a monophyletic clade, and only 17 sequences from the metagenomic dataset fell within this subtree. The MAGs corresponding to these 17 sequences were annotated with GTDB-tk (v 2.1.1) and 13 out of the 17 MAGs shared the same annotation: they belonged to phylum Bacillota, class *Limnochordia*, order DTU010. From here, two different families were identified: DTU010 and DTU012 (Supplementary Table 4).

After this preliminary analysis, the HQ and GQ MAGs (n=1053) were annotated with GTDB-tk (Supplementary Table 5). A total of 45 MAGs were classified as members of the order DTU010, and they were subsequently treated as potential representatives of MBA03. The phylogenomic tree of HQ and GQ MAGs (Figure 1) also revealed a large proportion of MAGs belonging to *Bacillota* (n = 503), while 104 MAGs were classified as *Bacteroidota,* 77 as *Actinobacteriota,* 63 as *Verrucomicrobiota* and 32 as *Chloroflexota,* while a total of 80 MAGs corresponded to Archaea. These results were expected considering previous studies reporting the average AD microbiome (Sundberg et al., 2013; Abendroth et al., 2015; Treu et al., 2016; Kirkegaard et al., 2017; Otto et al., 2023). Moreover, it was confirmed that MBA03 formed a monophyletic clade with all 45 sequences annotated as DTU010. Two different subclades could be clearly distinguished, corresponding to families DTU010 and DTU012 according to the GTDB taxonomy.

Figure 1. (A) Dendrogram of all the HQ and GQ MAGs from MICRO4BIOGAS (n=1053), computed with UBCG and visualized with iTOL. The legend shows a classification by different colours corresponding to different phyla, according to the annotation obtained with GTDB-tk (v 2.1.1), and a highlight on MBA03 (within *Bacillota*). (B) Tree with 108 leaves, computed with FastTree and visualized with iTOL, corresponding to the MAGs belonging to order DTU010 according to GTDB-tk (v 2.1.1). Two clades corresponding to two families (DTU010 in orange and DTU010 in yellow) can be seen, for which the names

Darwinibacteriaceae and *Wallaceae* are proposed, respectively. In both figures, bootstrap
values greater than 0.75 are represented by a small triangle in each branch.

For the enlargement of the dataset, 1804 MAGs from the Biogasmicrobiome
repository (https://biogasmicrobiome.env.dtu.dk/) and 16 from Dyksma et al.
were downloaded and annotated with GTDB-tk (Dyksma et al., 2016). After
filtering, a total of 64 potential MBA03 MAGs were compiled. Together with the
previous 45 MAGs (HQ and GQ) retrieved from MICRO4BIOGAS data, the final
dataset was composed of 108 genomes. These MAGs were used for creating a
phylogenomic tree (Figure 2), leading to an initial taxonomic delimitation of
MBA03: (i) there were two families, named DTU010 (which accounted for 82
MAGs) and DTU012 (with 26 MAGs) according to GTDB database; (ii) DTU010
clade included 20 different taxa annotated in GTDB. The most common was
DTU010 sp. 002391385, followed by sp. 900018335, and both formed
monophyletic clades; and (iii) there were 10 different taxa belonging to DTU012,
with the most common species being DTU012 sp.900019385.

Thereby, it was proposed the new order to be named *Darwinibacteriales* ord.
Nov., and the two families *Darwinibacteriaceae* fam. nov., and
Wallacebacteriacae fam. nov., after Charles Darwin and Alfred Wallace, who are
considered the fathers of the Theory of Evolution by natural selection.

3.3. Final taxonomic delimitation of MBA03

According to these results, the order *Darwinibacteriales* ord. nov. (DTU010
according to GTDB) was hypothesized to be equivalent to MBA03 (SILVA's name),
but further phylogenetic evidence was needed to get a proper delimitation of the
clade. For that reason, phylogenetic analyses based on the 16S rRNA gene were
carried out. A total of 52 16S rRNA sequences were identified by Prokka
(Seemann, 2014) in the 108 MAGs (i.e., not all MAGs contained 16S rRNA
sequences), and they were computed together with the 157 16S rRNA sequences
downloaded from the SILVA (SSU 138.1 ref NR) database in a phylogenetic tree
(n=209). Moreover, 772 sequences from *Clostridia* were added as outgroups.

The phylogenetic tree (Figure 2) confirmed the abovementioned hypothesis: the
sequences annotated as MBA03 in SILVA formed a monophyletic clade with a
clear subdivision of two clades, which included two distinct families
(*Darwinibacteraceae*/DTU010 and *Wallacebacteriaceae*/DTU012). Clade I was
formed by 52 sequences and included the MBA03 reference sequence, whereas
Clade II accounted for 44 sequences.

Figure 2. Phylogenetic tree based on 16S rRNA sequence from our own database and from SILVA ref NR SSU 138.1 database. The sequences in bold letter belong to the MAGs from our database, while the other sequences are the ones downloaded from SILVA. Coloured clades correspond to the outgroup sequences, and they have been collapsed for a better visualization of the MBA03 clade.

It is worth highlighting that, although both *Hydrogenispora* and *Darwinibacteriales* (MBA03) are annotated as class *Limnochordia* in the SILVA database, *Hydrogenispora* is annotated as *Clostridia* in the LPSN (List of Prokaryotic names with Standing in Nomenclature, Parte et al., 2020), which provides the internationally accepted and regulated taxonomic classification. This annotation, supported by other authors (FitzGerald, 2018; Westerholm et al., 2016), was confirmed once the tree was computed, as both *Hydrogenispora* and *Darwinibacteriales* appeared to be two different orders from class *Clostridia*. In other words; *Darwinibacterales* is proposed as a new order that belongs to class *Clostridia*.

Finally, a phylogenomic tree (Figure 3) was computed with the 108 genomes classified as MBA03 together with 206 genomes from *Clostridia* and one genome from *Limnochordia* as outgroups (Supplementary table 2). This confirmed the hypothesis that MBA03 belongs to a new order within class *Clostridia,* and it comprises two families.

Figure 3. Phylogenomic tree showing only the *Darwinibacteriales* (MBA03 in SILVA) clade, which includes two potential new families: *Darwinibacteriaceae* (DTU010) shaded in pink and *Wallacebacteriaceae* (DTU012) shaded in blue. Output groups from *Clostridia* and one genome from *Limnochordia* are collapsed in the tree for a better visualization. MAGs in green (n=9) belong to *Darwinibacteriales* (MBA03), as there is evidence at the genetic and genomic level. MAGs in red (n=9) may be MBA03, but their 16S rRNA phylogeny is not concordant. This could be explained by errors during the binning process. The remaining MAGs (n=90) are incomplete and/or 16S rRNA sequences could not recovered from the MAGs. Although they form a cluster with the rest of the MBA03 MAGs, they should remain as hypothetical members of this order.

According to these results, there is enough evidence to confirm that MBA03 belongs to a new taxonomical order, *Darwinibacteriales* ord. nov., which includes two potential new families, *Darwinibacteriaceae* fam. nov., (DTU010 according to GTDB) and *Wallacebacteriaceae* fam. nov.,(former DTU012). It is important to note that samples from MICRO4BIOGAS provided the higher number of MAGs belonging to MBA03, meaning that the samples and datasets from this project

marked a significant breakthrough towards the genetic isolation of MBA03 genomes. The formal description of these taxa can be found in Supplementary Material.

Once the two families from the order *Darwinibacteriales* (*Darwinibacteriaceae* and *Wallaceae*) were properly delimited, the read coverage of both families in the samples was calculated. 10 MAGs from DTU010 and 4 MAGs from DTU012 were used as reference for read mapping. The results showed that DTU010, with a mean abundance of 4.1% and a maximum abundance of near 18%, was considerably more abundant than DTU012, with a mean abundance of 0.6% (Supplementary Table 6).

3.5. Ecological distribution *Darwinibacteriales*

The ecological distribution of *Darwinibacteriales* was evaluated using IMNGS. This tool quantifies the prevalence and abundance of any 16S rRNA sequence of interest introduced as input, giving information about the number of coincidences with the samples included in its database. IMNGS also informs about the type of sample (e.g., soil, water, anaerobic digester, environmental) showing a hit with the input sequence. A total of 8856 SRA accessions containing 16S rRNA sequences from *Darwinibacteriales* were detected according to sequence similarity. Due to the high amount of data obtained, only the samples with an abundance of *Darwinibacteriales* higher than 0.1% were considered at first (Figure 4A). An exhaustive analysis of the sample metadata revealed that more than 99% of them belonged to anaerobic digestion studies, meaning that *Darwinibacteriales* is a very specialized bacterial group that grows and develops in this specific environment. This result reinforces the hypothesis that MBA03 *Darwinibacteriales* may play a central role in the AD process.

Figure 4. (A) Ecological niches of *Darwinibacteriaceae* based on coincidences of 16S rRNA sequences with samples of the IMNGS database with an abundance above 0.1%, (B) Ecological niches of *Darwinibacteriaceae* according to the samples with an abundance below 0.1% and excluding all the samples related to anaerobic digestion for a better insight into the wide range of environments from where it may be introduced to anaerobic reactors.

To study the ecological niches besides anaerobic digestion, a similar analysis
was performed considering only the samples with an abundance of
Darwinibacteriales-related sequences below 0.1%. The resulting 8650 hits were
then filtered in two steps: (i) first, those related to anaerobic digestion were
discarded, and (ii) results with less than 5 hits in total were removed from the
dataset, leaving 1636 samples that were categorised into different groups
(Figure 4B).

Soil was the second most common environment in which MBA03 could be found,
followed by rhizosphere and sediment samples. However, the abundance of
Darwinibacteriales in these habitats was substantially lower than in the case of
anaerobic digesters. Therefore, it can be hypothesized that these bacteria are
found in soil, plants and grounds, in a quiescent or inactive form. When
agricultural residues, soils, muds or sludges are taken as substrate for anaerobic
digestion, bacteria from the *Darwinibacteriales* order thrive in the anaerobic
digester environment. Similarly, the presence of MBA03 in gut and milk
microbiomes can be explained by the dietary habits of the animals. As
Darwinibacteriales exists in the soil, herbivorous and omnivorous animals may
ingest this group of bacteria during their feeding. Another hypothesis would be
that *Darwinibacteriales* is part of the native gut microbiota of the animals, but at
a very low abundance, and would thus be a reservoir which would guarantee the
prevalence in AD plants fed with those substrates. Therefore, when manure is
used as a substrate, *Darwinibacteriales* would be introduced into the bioreactors.

3.6. Unveiling the functional potential of *Darwinibacteriales*

In anaerobic digesters, the metabolic diversity of *Darwinibacteriales* remains
challenging due to the absence of known cultivable representatives. A total of 45
and 16 MAGs from *Darwinibacteriaceae* (former DTU010) and 16 from *Wallaceae*
(former DTU012), respectively, were analysed to gain insights into the potential
metabolic pathways of this bacterial order (Supplementary Table 7).

Darwinibacteriales are chemoorganoheterotrophs that derive energy from
anaerobic respiration through oxidative phosphorylation. They utilize inorganic
molecules of ferric iron (Fe^{3+}) as potential electron acceptors (K02012 and
K02011). These bacteria also possess sulphide:quinone oxidoreductase
(EC:1.8.5.4), an enzyme involved in the oxidation of sulphides, including
hydrogen sulphide, enabling them to be chemolitotrophic. Their ability to grow
on H_2S is interesting, as H_2S is usually not oxidized under anaerobic conditions.
However, there are some exceptions. For example, it has recently been
suggested to add reducing agents to control the H_2S level. In this regard, Jung et
al. (2020) highlighted the possibility to add magnetite. Jung et al. suggested that
the phenomenon of direct interspecies electron transfer (DIET) can not only
increase methanation activity, but can also reduce H_2S, which results in an
intracellular accumulation of S^0. In the present study, no hints for DIET have been
found so far. However, even in the case that DIET is not involved, it is
conceivable that electrons are transferred via alternative routes via the

mediated electron transfer (MIET). Due to the importance of sulfur-metabolism in
regard to industrial biogas formation, it would be of high interest, to study the
involvement of *Darwinibacteriales* in the sulfur-metabolism in further detail.
Apart from using anoxic electron donors, there is also the possibility to treat H_2S
due to micro-aeration in anaerobic digesters (van der Zee et al., 2007). Because
of this, plant operators inject sometimes small amounts of oxygen into anaerobic
digesters. In turn, this enables microbial oxidation of H_2S to sulfate, which
improves the overall biogas quality. The link to sulfur metabolism is particularly
interesting because some members of *Darwinibacteriales* appear to be
syntrophic microorganisms (as discussed later in 3.7). In fact, in recent years and
alongside Jung et al., several other articles have been examining the influence
between syntrophy and sulfur metabolism. To give here few examples: Igarashi
and Kuwabara (2016) applied a syntrophic community to dacite pumice to
induce oxygen supported H_2S oxidation. Shakeri Yekta et al. (2017) highlighted
the complex interplay between iron, sulfur metabolism, volatile fatty acid
turnover and syntrophic relation.

Darwinibacteriaceae and *Wallacebacteriaceae* have complete sets of genes for
glycolysis (Embden-Meyerhof pathway; M00001), gluconeogenesis (M00003),
and pyruvate oxidation (M00307) pathways. However, they lack certain essential
genes required for the tricarboxylic acid (TCA) cycle, specifically isocitrate
dehydrogenase. Both bacterial families can convert glycogen to glucose-6P.

Most MAGs from both families contain genes encoding beta-N-
acetylhexosaminidase for bacterial cell wall peptidoglycan degradation and
recycling, pullulanase for starch degradation, and beta-glucosidase for cellobiose
metabolism. *Darwinibacteriaceae* members also possess additional genes
encoding chitinase for chitin degradation, D-galacturonate isomerase for pectin
utilization, and alpha-amylase for starch hydrolysis. *Wallacebacteriaceae*
members carry the gene encoding oligogalacturonide lyase, allowing them to
degrade pectin. Both *Wallacebacteriaceae* and *Darwinibacteriaceae* exhibit the
ability to degrade catechol and methylcatechol through catechol 2,3-
dioxygenase enzymes (EC 1.13.11.2). However, the presence of these
compounds can be toxic to microorganisms in anaerobic digesters, inhibiting the
digestion process. These findings suggest that each bacterial family can exploit
different organic substrates as carbon and energy sources, maybe facilitating
their coexistence. This also indicates that the importance of MB03 does not only
restrict to one single phase of anaerobic digestion. Apart from acetogenesis and
syntrophic acetate oxidation, MB03 could also be involved in the hydrolysis
phase as well.

Members of *Darwinibacteriales* assimilate nitrogen by converting NH_3 to organic
nitrogen through the enzymatic activities of glutamine synthetase (EC 6.3.1.2)
and glutamate (NADPH) synthase (EC 1.4.1.13). However, genes associated with
other nitrogen metabolism processes, such as nitrate reduction, denitrification,
nitrogen fixation, nitrification, and anaerobic ammoniumoxidation, were not
observed. Most representatives of *Darwinibacteriaceae* have auxotrophic

requirements for various amino acids, while *Wallacebacteriaceae* representatives
are auxotrophic for a smaller set of amino acids (Supplementary Table 7).

Both bacterial families share several ABC transporters, including ABCB-BACATP
binding cassettes and NitT/TauT family transporters. They also possess multiple
sugar transport systems. *Darwinibacteriaceae* exhibits ABC transporters for iron
(III), phosphonate, alpha-glucoside, aldouronate, sn-glycerol 3-phosphate,
cobalamin, and haemolysin transport systems. Both *Darwinibacteriaceae* and
Wallacebacteriaceae show transport systems for maltose/maltodextrin,
arabinogalactan/maltooligosaccharide, raffinose/stachyose/melibiose,
glucose/mannose, trehalose/maltose, cellobiose, N,N'-diacetylchitobiose, L-
arabinose/D-xylose, rhamnose, ribose, fructooligosaccharides, sorbitol/mannitol,
arabinooligosaccharide, polygalacturonan/rhamnogalacturonan, pectin-derived
oligosaccharide, cobalt, spermidine/putrescine, nickel, and biotin transport
systems.

Members of *Darwinibacteriales* possess resistance genes against vancomycin
(vanW) and tetracycline (tetM and tetO). They also encode multidrug efflux
pumps (mdlA/smdA; mdlB/smdB; efrA/efrE) to withstand various antimicrobial
compounds.

Both families are mobile, as indicated by the presence of flagella synthesis
genes. However, *Darwinibacteriaceae* lacks genes for the MS/C ring type III
secretion system. Members of both taxa possess the enzymatic machinery for
chemotactic movement and detection and transport of D-ribose and D-galactose
via ABC transporters.

3.7 Members of *Darwinibacteriaceae* as potential SAOB

Members of the MBA03 lineage were previously hypothesised to be SAOB in AD
based on correlation studies (Perman et al., 2022; Zheng et al., 2019; Zeng
et al., 2021) together with other unclassified *Clostridia* (Dyksma et al., 2020);
however, the metabolism of this bacterial lineage, classified in this study as
Darwinibacteriaceae, remains poorly understood and characterised due to the
lack of cultured representatives. SAO bacteria possess either the known reversed
Wood-Ljungdahl (WL) pathway or an alternative route coupled to a glycine
cleavage system (GCS) (Zeng et al., 2021). Recently, the co-occurrence of
members of *MBA03* with other potential SAO bacteria, such as DTU014 and
Syntrophaceticus, and with hydrogenotrophic methanogens such as
Methanothermobacter, *Methanoculleus* and *Methanosarcina,* has been observed
(Otto et al., 2023). This suggests that several taxonomic groups identified as
potential SAOBs, including *Darwinibacteriaceae*, grow simultaneously with
methanogens to provide a continuous supply of H_2 and CO_2 for hydrogenotrophic
methanogenesis. Usually, hydrogenotrophic methanogens are associated with
high performance plants, whereas the acetoclastic methanogens *Methanothrix*
(formally *Methanosaeta*) are found specifically in low performance systems such
as sewage sludge digesters (Sunderberg et al. 2013; Abendroth et al., 2015).
Therefore, due to the co-occurrence between members of *Darwinibacteriaceae*
and hydrogenotrophic methanogens, we hypothesise that the presence of

Darwinibacteriaceae could be associated with a higher biogas production in
industrial AD plants, although further analyses are required.

To confirm whether members of the *Darwinibacteriaceae* and
Wallacebacteriaceae are SAOBs, their MAGs were mined to identify genes of the
reversed Wood-Ljungdahl pathway (Figure 5). In all members of the
Darwinibacteriaceae family, gene *Acka* was consistently detected, whereas *Pta*,
Acs, the *CODH* complex, and *MetF* were conspicuously absent. Most of the
Darwinibacteriaceae MAGs contained *FolD*, *Fhs*, and *Fdh* enzymes, which play
vital roles in the acquisition of CO_2 and H_2 from formate. The absence of gene *Pta*
and *Acs* could be justified by the presence of glycine reductase (*Grd*) in the
glycine cleavage system, which facilitates the conversion of glycine for entry into
the GCS (as reported by C. Li et al., 2022). Thus, the entire reverse WL-GCS
pathway was encoded by members of *Darwinibacteriaceae*, with only two genes,
namely *Pta* and *Acs* missing, which are in fact not necessary for the completion
of the pathway, as in the GCS, CH3-CO-Pi is directly converted to glycine through
the function of glycine reductase (Grd) (Li et al., 2022). The remaining enzymes
were identified in more than 70% of the *Darwinibacteriaceae* MAGs, with the
exception of *Dld*, which was present in only 60% of them.

In the case of the family *Wallacebacteriaceae*, only genes *AckA* and *Fhs* were
identified, being the other genes of the pathway were absent. The absence of
critical enzymes of the reversed Wood-Ljungdahl pathway, including *Sda*, *Grd*,
FolD, and *Fdh* suggests that members suggests that *Wallacebacteriaceae* are not
syntrophic acetate oxidising bacteria.

Based on the evidence found, it could be hypothesized that the family
Darwinibacteriaceae is a potential SAOB, as it encodes for the necessary genes
to perform the reversed Wood-Ljungdahl pathway coupled to the glycine
cleavage system. Within the intricate anaerobic digestion metabolic web, this
pathway would allow these bacteria to perform the oxidation of acetate to H_2 and
CO_2, reaction that would constitute a key step in anaerobic digestion, as
hydrogenotrophic archaea would take this H_2 as a substrate to ultimately
produce methane. However, the family *Wallacebacteriaceae* lacks several key
enzymatic of this pathway. Considering the relatively low abundance of this
taxon in the analyzed bioreactors (as detailed in Supplementary Table 6), our
hypothesis suggests that *Wallacebacteriaceae* may not play a significant role in
the functioning of anaerobic digesters, unlike *Darwinibacteriaceae*. This could be
attributed to the absence of a symbiotic relationship between
Wallacebacteriaceae and hydrogenotrophic archaea.

Figure 5. Mining of reverse WL pathway (black lines) coupled with glycine cleavage
system (purple lines) (WL-GCS) genes in MAGs of the families *Darwinibacteriaceae*
(n=45) and *Wallacebacteriaceae* (n=16). Red to green colors indicate the percentage of
MAGs in which the genes were identified. A gene was considered to be present in each
family if it was found in more than 70% of the MAGs. Bacteria from the family
Darwinibacteriaceae encode the genetic repertoire necessary to metabolise acetate to
produce CO2 and H2, which are then used by hydrogenotrophic archaea to produce
methane, whereas *Wallacebacteriaceae* showed a truncated pathway. Abbreviations:
AckA; acetate kinase, *Pta*; Phosphate acetyltransferase, *Acs*; Acetyl-coA synthetase, *Por*;
pyruvate-ferredoxin reductase, *Sda*; L-serine dehydratase, *GlyA*; glycine
hydroxymethyltransferase, Grd*;* glycine reductase, *GcvP*; glycine dehydrogenase,
GcvH/T; glycine cleavage system H/T protein, *Dld*; dihydrolipoyl dehydrogenase, *PflD*;
pyruvate-formate lyase, *AcsE;* iron sulfur protein methyltransferase, *metF*;
methylenetetrahydrofolate reductase, *FolD*; methylenetetrahydrofolate
dehydrogenase/cyclohydrolase, *Fhs*; formate-tetrahydrofolate ligase, *Fdh*; formate
dehydrogenase.

4. Conclusions

In this work, a cryptic bacterial group previously referred as MBA03 in SILVA
database, was described for the first time by recovering and analysing
representative genomes belonging to this taxon. Phylogenetic analyses
confirmed that MBA03 is a representative of a new taxonomic order with the
proposed name *Darwinibacteriales* ord. nov. Two different families can be
distinguished within this order: *Darwinibacteriaceae* fam. nov. (also known as
DTU010 according to the GTDB taxonomy) and *Wallaceae* fam. nov (also known
as DTU012).

Order *Darwinibacteriales,* particularly family *Darwinibacteriaceae,* is among the
most abundant genera in anaerobic digesters, regardless of operational and
chemical parameters such as feedstock and type of reactor. *Darwinibacteriales* is

a very specific bacterial group, being anaerobic digestion its main ecological niche. Bacteria from this order are also present in soils, sediments, plants and animal gut microbiomes at very low abundances. These materials are often used as substrates to feed anaerobic digesters, acting as a reservoir and an inoculum that allows *Darwinibacteriales* to thrive when the AD process begins.

The presence of the reversed Wood-Ljungdahl pathway coupled to the glycine cleavage system in the family *Darwinibacteriaceae* and the co-ocurrence of *Darwinibacteriales* with several hydrogenotrophic archaea (Otto et al., 2023) reveals that they are potential syntrophic acetate oxidising bacteria, and thus that they may act as competitors for acetate with acetoclastic archaea, favouring hydrogenotrophic methanogenesis. Furthermore, these bacteria may be important during hydrolysis, like other members of the *Clostridia* class, since *Darwinibacteriales* harbour a wide repertoire of hydrolytic enzymes. Nevertheless, it is important to highlight that this description has been entirely based on genome characterization, and so its culture and further laboratory analyses are needed to confirm what has been hypothesized.

The high abundance of *Darwinibacteriales* in AD environments strongly suggests an exaptation mechanism to this industrial process. In other words, *Darwinibacteriales* are naturally occurring hydrolytic and potentially SAOB bacteria that are pre-adapted to AD and are thus massively enriched in biogas producing and other AD facilities, where they may play an industrially relevant role that our results have only begun to reveal.

References

References

Abendroth, C., Vilanova, C., Günther, T., Luschnig, O., & Porcar, M. (2015).
Eubacteria and archaea communities in seven mesophile anaerobic
digester plants in Germany. Biotechnology for Biofuels, 8(1), 1-10.

Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., &
Ogata, H. (2020). KofamKOALA: KEGG Ortholog assignment based on
profile HMM and adaptive score threshold. Bioinformatics, 36(7), 2251-
2252. https://doi.org/10.1093/bioinformatics/btz859

Bowers, R. M., Kyrpides, N. C., Stepanauskas, R., Harmon-Smith, M., Doud, D.,
Reddy, T. B. K., Schulz, F., Jarett, J., Rivers, A. R., Eloe-Fadrosh, E. A.,
Tringe, S. G., Ivanova, N. N., Copeland, A., Clum, A., Becraft, E. D.,
Malmstrom, R. R., Birren, B., Podar, M., Bork, P., … Woyke, T. (2017).
Minimum information about a single amplified genome (MISAG) and a
metagenome-assembled genome (MIMAG) of bacteria and archaea.
Nature Biotechnology, 35(8), Article 8. https://doi.org/10.1038/nbt.3893

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein
alignment using DIAMOND. Nature Methods, 12(1), Article 1.
https://doi.org/10.1038/nmeth.3176

Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2022). GTDB-Tk v2:
Memory friendly classification with the genome taxonomy database.
Bioinformatics, 38(23), 5315-5316.
https://doi.org/10.1093/bioinformatics/btac672

Coudert, E., Gehant, S., de Castro, E., Pozzato, M., Baratin, D., Neto, T., Sigrist,
C. J. A., Redaschi, N., Bridge, A., & The UniProt Consortium. (2023).
Annotation of biologically relevant ligands in UniProtKB using ChEBI.
Bioinformatics, 39(1), btac793.
https://doi.org/10.1093/bioinformatics/btac793

De Vrieze, J., Saunders, A. M., He, Y., Fang, J., Nielsen, P. H., Verstraete, W., &
Boon, N. (2015). Ammonia and temperature determine potential clustering
in the anaerobic digestion microbiome. Water Research, 75,   312-323.
https://doi.org/10.1016/j.watres.2015.02.025

Dyksma, S., Jansen, L., & Gallert, C. (2020). Syntrophic acetate oxidation
replaces acetoclastic methanogenesis during thermophilic digestion of
biowaste. Microbiome, 8(1), 105. https://doi.org/10.1186/s40168-020-
00862-5

Fang, C., Zhou, L., Liu, Y., Xiong, J., Su, Y., Lan, Z., Han, L., & Huang, G. (2022).
Effect of micro-aerobic conditions based on semipermeable membrane-
covered on greenhouse gas emissions and bacterial community during
dairy manure storage at industrial scale. Environmental Pollution, 299,
118879. https://doi.org/10.1016/j.envpol.2022.118879

Feng, Y., Wang, Y., Zhu, B., Gao, G. F., Guo, Y., & Hu, Y. (2021). Metagenome-
assembled genomes and gene catalog from the chicken gut microbiome

aid in deciphering antibiotic resistomes. Communications Biology, 4(1),
Article 1. https://doi.org/10.1038/s42003-021-02827-2

FitzGerald, J. A. (2018). The microbial ecology of anaerobic digestion:
Characterising novel biogas configurations through molecular and
statistical methods. https://hdl.handle.net/10468/7573

Graham, E. D., Heidelberg, J. F., & Tully, B. J. (2018). Potential for primary
productivity in a globally-distributed bacterial phototroph. The ISME
Journal, 12(7), Article 7. https://doi.org/10.1038/s41396-018-0091-3

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality
assessment tool for genome assemblies. Bioinformatics, 29(8), 1072-
1075. https://doi.org/10.1093/bioinformatics/btt086

Igarashi, K., & Kuwabara, T. (2016). Hydrogen-Sulfide-Free Methane
Production by Fermenter–Methanogen Syntrophy Using Dacite Pumice
under Aerobic Gas Phase. Energy & Fuels, 30(6), 4945-4950.

Jung, H., Baek, G., & Lee, C. (2020). Magnetite-assisted in situ microbial
oxidation of H2S to S0 during anaerobic digestion: A new potential for
sulfide control. Chemical Engineering Journal, 397, 124982.

Kang, D. D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., & Wang, Z. (2019).
MetaBAT 2: An adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. PeerJ, 7, e7359.
https://doi.org/10.7717/peerj.7359

Kim, D., Song, L., Breitwieser, F. P., & Salzberg, S. L. (2016). Centrifuge: Rapid
and sensitive classification of metagenomic sequences. Genome
Research, 26(12), 1721-1729. https://doi.org/10.1101/gr.210641.116

Kirkegaard, R. H., McIlroy, S. J., Kristensen, J. M., Nierychlo, M., Karst, S. M.,
Dueholm, M. S., ... & Nielsen, P. H. (2017). The impact of immigration on
microbial community composition in full-scale anaerobic digesters.
Scientific reports, 7(1), 9343.

Lagkouvardos, I., Joseph, D., Kapfhammer, M., Giritli, S., Horn, M., Haller, D., &
Clavel, T. (2016). IMNGS: A comprehensive open resource of processed
16S rRNA microbial profiles for ecology and diversity studies. Scientific
Reports, 6(1), Article 1. https://doi.org/10.1038/srep33721

Laguillaumie, L., Rafrafi, Y., Moya-Leclair, E., Delagnes, D., Dubos, S.,
Spérandio, M., Paul, E., & Dumas, C. (2022). Stability of ex situ biological
methanation of H2/CO2 with a mixed microbial culture in a pilot scale
bubble column reactor. Bioresource Technology, 354, 127180.
https://doi.org/10.1016/j.biortech.2022.127180

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with
Bowtie   2.   Nature   methods,   9(4),   357-359.
https://doi.org/10.1038/nmeth.1923

Lebuhn, M., Weiß, S., Munk, B., & Guebitz, G. M. (2015). Microbiology and Molecular Biology Tools for Biogas Process Analysis, Diagnosis and Control. En G. M. Guebitz, A. Bauer, G. Bochmann, A. Gronauer, & S. Weiss (Eds.), Biogas Science and Technology (pp. 1-40). Springer International Publishing. https://doi.org/10.1007/978-3-319-21993-6_1

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for phylogenetic tree display and annotation. Nucleic Acids Research, 49(W1), W293-W296. https://doi.org/10.1093/nar/gkab301

Li, C., Hao, L., Lü, F., Duan, H., Zhang, H., & He, P. (2022). Syntrophic Acetate-Oxidizing Microbial Consortia Enriched from Full-Scale Mesophilic Food Waste Anaerobic Digesters Showing High Biodiversity and Functional Redundancy. mSystems, 7(5), e00339-22. https://doi.org/10.1128/msystems.00339-22

Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: An ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics, 31(10), 1674-1676. https://doi.org/10.1093/bioinformatics/btv033

Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics, 22(13), 1658-1659. https://doi.org/10.1093/bioinformatics/btl158

Liu, X., Zhu, X., Yellezuome, D., Liu, R., Liu, X., Sun, C., Abd-Alla, M. H., & Rasmey, A.-H. M. (2023). Effects of adding Thermoanaerobacterium thermosaccharolyticum in the hydrogen production stage of a two-stage anaerobic digestion system on hydrogen-methane production and microbial communities. Fuel, 342, 127831. https://doi.org/10.1016/j.fuel.2023.127831

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.Journal, 17(1), Article 1. https://doi.org/10.14806/ej.17.1.200

Na, S.-I., Kim, Y. O., Yoon, S.-H., Ha, S., Baek, I., & Chun, J. (2018). UBCG: Up-to-date bacterial core gene set and pipeline for phylogenomic tree reconstruction. Journal of Microbiology, 56(4), 280-285. https://doi.org/10.1007/s12275-018-8014-6

Nwokolo, N., Mukumba, P., Obileke, K., & Enebe, M. (2020). Waste to Energy: A Focus on the Impact of Substrate Type in Biogas Production. Processes, 8(10), Article 10. https://doi.org/10.3390/pr8101224

Perman, E., Schnürer, A., Björn, A., & Moestedt, J. (2022). Serial anaerobic digestion improves protein degradation and biogas production from mixed food waste. Biomass and Bioenergy, 161, 106478. https://doi.org/10.1016/j.biombioe.2022.106478

Price, M. N., Dehal, P. S., & Arkin, A. P. (2009). FastTree: Computing Large Minimum Evolution Trees with Profiles instead of a Distance Matrix.

Molecular   Biology   and   Evolution,   26(7),   1641-1650.
https://doi.org/10.1093/molbev/msp077

Pruesse, E., Peplies, J., & Glöckner, F. O. (2012). SINA: Accurate high-
throughput multiple sequence alignment of ribosomal RNA genes.
Bioinformatics,   28(14),   1823-1829.
https://doi.org/10.1093/bioinformatics/bts252

Rivière, D., Desvignes, V., Pelletier, E., Chaussonnerie, S., Guermazi, S.,
Weissenbach, J., Li, T., Camacho, P., & Sghir, A. (2009). Towards the
definition of a core of microorganisms involved in anaerobic digestion of
sludge.   The   ISME   Journal,   3(6),   Article   6.
https://doi.org/10.1038/ismej.2009.2

Rodriguez-R, L. M., Gunturu, S., Harvey, W. T., Rosselló-Mora, R., Tiedje, J. M.,
Cole, J. R., & Konstantinidis, K. T. (2018). The Microbial Genomes Atlas
(MiGA) webserver: Taxonomic and gene diversity analysis of Archaea and
Bacteria at the whole genome level. Nucleic Acids Research, 46(W1),
W282-W288. https://doi.org/10.1093/nar/gky467

Roopnarain, A., Rama, H., Ndaba, B., Bello-Akinosho, M., Bamuza-Pemu, E., &
Adeleke, R. (2021). Unravelling the anaerobic digestion 'black box':
Biotechnological approaches for process optimization. Renewable and
Sustainable   Energy   Reviews,   152,   111717.
https://doi.org/10.1016/j.rser.2021.111717

Schnürer, A. (2016). Biogas Production: Microbiology and Technology. En R.
Hatti-Kaul, G. Mamo, & B. Mattiasson (Eds.), Anaerobes in Biotechnology
(pp.   195-234).   Springer   International   Publishing.
https://doi.org/10.1007/10_2016_5

Schwengers, O., Jelonek, L., Dieckmann, M. A., Beyvers, S., Blom, J., &
Goesmann, A. (2021). Bakta: Rapid and standardized annotation of
bacterial genomes via alignment-free sequence identification. Microbial
Genomics, 7(11), 000685. https://doi.org/10.1099/mgen.0.000685

Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation.
Bioinformatics,   30(14),   2068-2069.
https://doi.org/10.1093/bioinformatics/btu153

Sieber, C. M. K., Probst, A. J., Sharrar, A., Thomas, B. C., Hess, M., Tringe, S.   G.,
& Banfield, J. F. (2018). Recovery of genomes from metagenomes via   a
dereplication, aggregation and scoring strategy. Nature Microbiology,
3(7), Article 7. https://doi.org/10.1038/s41564-018-0171-1

Sundberg, C., Al-Soud, W. A., Larsson, M., Alm, E., Yekta, S. S., Svensson, B. H.,
... & Karlsson, A. (2013). 454 pyrosequencing analyses of bacterial and
archaeal richness in 21 full-scale biogas digesters. FEMS microbiology
ecology, 85(3), 612-626.

Treu, L., Kougias, P. G., Campanaro, S., Bassani, I., & Angelidaki, I. (2016).
Deeper insight into the structure of the anaerobic digestion microbial

community; the biogas microbiome database is expanded with 157 new
genomes.   Bioresource   Technology,   216,   260-266.
https://doi.org/10.1016/j.biortech.2016.05.081

Tsang, Y. F., Kumar, V., Samadar, P., Yang, Y., Lee, J., Ok, Y. S., Song, H., Kim,
K.-H., Kwon, E. E., & Jeon, Y. J. (2019). Production of bioplastic through
food waste valorization. Environment International, 127, 625-644.
https://doi.org/10.1016/j.envint.2019.03.076

Van der Zee, F. P., Villaverde, S., Garcia, P. A., & Polanco, F. F. (2007). Sulfide
removal by moderate oxygenation of anaerobic sludge environments.
Bioresource technology, 98(3), 518-524.

Wang, Y., Wegener, G., Williams, T. A., Xie, R., Hou, J., Tian, C., ... & Xiao, X.
(2021). A methylotrophic origin of methanogenesis and early divergence
of anaerobic multicarbon alkane metabolism. Science advances, 7(27),
eabj1453.

Westerholm, M., Moestedt, J., & Schnürer, A. (2016). Biogas production through
syntrophic acetate oxidation and deliberate operating strategies for
improved   digester   performance.   Applied   Energy,   179,   124-135.
https://doi.org/10.1016/j.apenergy.2016.06.061

Whitman, W. B., Bowen, T. L., & Boone, D. R. (2006). The methanogenic
bacteria. Prokaryotes, 3(Chapter 9), 165-207.

Wu, Y.-W., Simmons, B. A., & Singer, S. W. (2016). MaxBin 2.0: An automated
binning algorithm to recover genomes from multiple metagenomic
datasets.   Bioinformatics,   32(4),   605-607.
https://doi.org/10.1093/bioinformatics/btv638

Yaqoob, H., Teoh, Y. H., Ud Din, Z., Sabah, N. U., Jamil, M. A., Mujtaba, M. A., &
Abid, A. (2021). The potential of sustainable biogas production from
biomass waste for power generation in Pakistan. Journal of Cleaner
Production, 307, 127250. https://doi.org/10.1016/j.jclepro.2021.127250

Yekta, S. S., Ziels, R. M., Björn, A., Skyllberg, U., Ejlertsson, J., Karlsson, A., ... &
Svensson, B. H. (2017). Importance of sulfide interaction with iron as
regulator of the microbial community in biogas reactors and its effect on
methanogenesis, volatile fatty acids turnover, and syntrophic long-chain
fatty acids degradation. Journal of bioscience and bioengineering, 123(5),
597-605.

Zeng, Y., Zheng, D., Gou, M., Xia, Z.-Y., Chen, Y.-T., Nobu, M. K., & Tang, Y.-Q.
(2021). Chasing the metabolism of novel syntrophic acetate-oxidizing
bacteria   in   thermophilic   methanogenic   chemostats   (p.
2021.07.06.451242). bioRxiv. https://doi.org/10.1101/2021.07.06.451242

Zheng, D., Wang, H.-Z., Gou, M., Nobu, M. K., Narihiro, T., Hu, B., Nie, Y., &
Tang, Y.-Q. (2019). Identification of novel potential acetate-oxidizing
bacteria in thermophilic methanogenic chemostats by DNA stable isotope

probing. Applied Microbiology and Biotechnology, 103(20), 8631-8645.
https://doi.org/10.1007/s00253-019-10078-9