Unveiling the ecology, taxonomy and metabolic capabilities of
MBA03, a potential key player in anaerobic digestion

Roser Puchol-Royo[1], Javier Pascual[1], Asier Ortega-Legarreta[1,2], Pascal Otto[3],
Jeroen Tideman[4], Sjoerd-Jan de Vries[4], Christian Abendroth[5], Kristie Tanner[1],
Manuel Porcar[1,6] & Adriel Latorre-Perez[1]

[1]Darwin Bioprospecting Excellence, S.L. Parc Cientific Universitat de Valencia,
Paterna, Valencia, Spain.

[2]Translational Bioinformatics Unit, Navarrabiomed, Complejo Hospitalario de
Navarra (CHN), Universidad Pública de Navarra (UPNA), IdiSNA, Pamplona, Spain

[3]Technische Universität Dresden, Institute of Waste Management and Circular
Economy, Pirna, Germany.

[4]Bioclear Earth B.V., Groningen, The Netherlands

[5]Brandenburgische Technische Universität Cottbus-Senftenberg, Chair of Circular
Economy, Cottbus, Germany

[6]Institute for Integrative Systems Biology I2SysBio (University of Valencia - CSIC),
Paterna, Spain

Corresponding author: A. Latorre-Pérez (alatorre@darwinbioprospecting.com)

Abstract

Biogas, a mix of CO_2, CH_4 and small proportions of other gases, is a biofuel obtained by anaerobic digestion (AD). Biogas production is often considered a black box process, as the role and dynamics of some of the microorganisms involved remain undisclosed. Previous metataxonomic studies in the frame of the MICRO4BIOGAS project (www.micro4biogas.eu) revealed that MBA03, an uncharacterised and uncultured bacterial taxon, was very prevalent and abundant in industrial full-scale AD plants. Surprisingly, no culturable specimen or genome of this taxon has ever been reported, so its role in AD has remained unclear. In the present work, thirty samples derived from anaerobic digesters were sequenced, allowing the reconstruction of 108 metagenome-assembled genomes (MAGs) potentially belonging to MBA03. According to phylogenetic analyses and genomic similarity indices, MBA03 constitutes a new bacterial order, proposed as *Darwinibacteriales* ord. nov., which includes *Darwinibacter acetoxidans* gen. nov., sp. nov. of the family *Darwinibacteriaceae* fam. nov., along with *Wallacebacter cryptica* gen. nov., sp. nov. of the *Wallacebacteriaceae* fam. nov. Ecotaxonomic studies determined that AD processes are the main ecological niche of *Darwinibacteriales*. Moreover, metabolic predictions identified *Darwinibacteraceae* members as putative syntrophic acetate oxidising bacteria (SAOB), as they encode for the reversed Wood-Ljungdahl (W-L) pathway coupled to the glycine cleavage system. This suggests that *Darwinibacteraceae* members work in collaboration with hydrogenotrophic archaea to produce methane in industrial biogas plants. Overall, our findings present *Darwinibacteriales* as a potential key player in anaerobic digestion and pave the way towards the complete characterisation of this newly described bacterial taxa.

Keywords: Metagenomics, anaerobic digestion, biogas, SAOB, MBA03, *Darwinibacteriales*

1 Introduction

The evidence supporting the need for a transition from fossil to renewable energy is overwhelming and unanimous (Yaqoob et al., 2021). In recent years, biogas has arisen as one of the leading alternatives to non-renewable fossil fuels such as natural gas. Biogas is produced by anaerobic digestion (AD) of organic material, through a complex microbiological process in which several groups of microorganisms with diverse metabolic abilities work in close collaboration to obtain methane as the main product (Schnürer, 2016). In addition, biogas production contributes to the utilisation of today's global waste streams of various origins as a source of organic matter (Tsang et al., 2019).

Biogas primarily consists of methane (CH_4), carbon dioxide (CO_2) and other gases like sulfuric acid (H_2S), nitrogen (N_2), ammonia (NH_3), and hydrogen (H_2) in lower concentrations (Yaqoob et al., 2021). Despite the complexity of the AD process, the microbial metabolic reactions that occur in this process can be summarized in four key steps: hydrolysis, acidogenesis, acetogenesis and methanogenesis (Schnürer, 2016). Briefly, during hydrolysis, complex macromolecules are broken down into various oligo-, di- and monomers, the final concentrations of which will depend on the biomass source (Nwokolo et al., 2020). These molecules are uptaken and used by acidogenic bacteria to produce H_2, CO_2, alcohols and volatile fatty acids (VFAs) during acidogenesis, while in acetogenesis, acetogenic bacteria synthesize acetic acid through the Wood-Ljungdahl (W-L) pathway or autotrophically from H_2 and CO_2. Subsequently, methanogens metabolize acetate, hydrogen and/or methylated compounds to produce methane, creating an equilibrium and a co-dependence between bacteria and archaea (Vanwonterghem et al., 2016; Nwokolo et al., 2020). Two pathways can occur in the production of methane from acetate: the first one, directly by acetoclastic methanogens, which split acetate to methane and CO_2, with the methyl group yielding methane and the carboxyl group yielding CO_2. The other pathway results from syntrophic acetate-oxidizing bacteria (SAOB) oxidizing acetate to CO_2 and H_2, which is subsequently used by hydrogenotrophic archaea to obtain methane. Thus, this pathway must be coupled with hydrogenotrophic methanogenesis (SAO-HM).

Despite the high economic potential, knowledge for the production of biogas with AD has remained stagnant, as the high variability of both the different operational parameters and the native microbial communities involved represent a challenge for its optimization and standardization. Improving the feasibility of the process and achieving a consistent and controlled way to obtain biogas would constitute a huge change in the energy industry (Nielsen & Angelidaki, 2008). But reaching such an ambitious goal requires gaining insights in the microbial machinery behind AD and methanogenesis, and this goes through the study of a variety of AD systems with different chemical, taxonomical and operational parameters.

In this context, the EU-funded project MICRO4BIOGAS (https://micro4biogas.eu) was born in 2021. The ultimate objective of the project is to improve the yield, speed, quality and robustness of biogas production by optimising the AD microbiome. In the frame of the project, 80 different samples were collected from various full-scale AD plants in the Netherlands, Germany and Austria.

After metataxonomic analyses of the 80 samples during MICRO4BIOGAS project, a miscellaneous taxonomic group (i.e., MBA03) was found to be very prevalent and abundant, with some samples reaching abundances up to 38% of total bacteria. Moreover, MBA03 was among the taxa that correlated most strongly with operational (i.e., temperature) and chemical parameters (i.e., acetic acid) (Otto et al., 2023), suggesting that it may be a microorganism of great relevance for the AD process. MBA03 (AB114313.1) was described for the first time by Tang et al. as an uncultured clone obtained from a thermophilic anaerobic municipal solid-waste digester (Tang et al., 2004). The proposal of MBA03 exclusively relied on the 16S rRNA gene sequence. Since then, a series of metataxonomic studies have reported the presence of MBA03 in anaerobic digester systems (Dyksma et al., 2020; FitzGerald, 2018; Laguillaumie et al., 2022), representing up to 70% of the relative abundance in some samples (FitzGerald, 2018). This taxon has been associated with a higher production of methane and has been suggested as an indicator of the stability of the AD process (Fang et al., 2022; Laguillaumie et al., 2022).

Surprisingly, MBA03 has not been properly taxonomically delimited and no culturable specimen or genome of this taxon has ever been reported. However, according to SILVA database (SSU 138.1 Ref NR), MBA03 represents a heterogenous group of more than 150 16S rRNA sequences. Therefore, a deeper characterization of this taxonomic group was urgently needed for unravelling its role in AD processes.

In this work, we used metagenomic data from 30 MICRO4BIOGAS samples and other AD studies to propose MBA03 as a member *Darwinibacteriales* ord. nov., a not-yet-cultured bacterial taxa belonging to the class *Clostridia*. *Darwinibacteriales* includes two different genera: *Darwinibacter acetoxidans* gen. nov., sp. nov., of the *Darwinibacteriaceae* family, and *Wallacebacter cryptica* gen. nov., sp. nov., of the *Wallacebacteriacea* fam. nov. Based on our results, both *Darwinibacter* and *Wallacebacter* predominantly inhabit AD environments. Furthermore, we have inferred that *Darwinibacter* may play a key role in AD processes related to biogas production, as it includes the reversed Wood-Ljungdahl pathway coupled to the glycine cleavage system, which is required for the synthesis of H_2 and CO_2 from acetate and the subsequent production of methane by hydrogenotrophic methanogenesis.

2. Materials and methods

2.1. Sample description, DNA extraction and metagenomic sequencing

As part of the MICRO4BIOGAS project, a total of 80 samples were collected from anaerobic digestion systems at 45 large-scale reactors across Germany, Austria and the Netherlands (Otto et al., 2023). A subset of 30 samples was analysed through metagenomic sequencing. This subset included samples enriched in MBA03, hydrolytic, acidogenic and acetogenic bacteria and/or methanogenic archaea. A full description of the samples can be found in Supplementary Table 1.

The sludge samples were washed with PBS, and the DNeasy® PowerSoil® Pro kit
(QIAGEN, Germany) was used for DNA extraction. The metagenomic DNA was
fragmented, end polished, A-tailed, and ligated with Illumina adapters. Libraries
were sequenced at Novogene (UK) using the NovaSeq 6000 Illumina platform
(150 bp x 2). The average sequencing depth was 45M reads/sample (min.: 35M
reads; max.: 55M reads). Additional information about DNA extraction and
sequencing can be found at Supplementary Methods.

147 2.3. Metagenomic analysis

Reads were filtered and quality checked as described at Supplementary
Methods. MEGAHIT (v1.2.9) was used with default parameters for assembling the
filtered reads (D. Li et al., 2015). Two different binning strategies were applied:
MetaBAT (Kang et al., 2019) and MaxBin (Wu et al., 2016). The MAGs obtained
were refined with Das Tool (Sieber et al., 2018) and classified according to their
quality with CheckM (Parks et al., 2014): high quality (HQ, completeness ≥ 90%
and contamination ≤ 5%), good quality (GQ, completeness ≥ 80% and
contamination ≤ 10%) and medium-low quality (LQ, completeness ≤ 80% and/or
contamination ≥ 10%), following the criteria of Bowers et al., 2017 and Feng
et al., 2021. After that, Centrifuge (Kim et al., 2016) was used to study the
taxonomic profile of the samples, while the resulting MAGs were taxonomically
annotated with Microbial Genome Atlas (MiGA) (Rodriguez-R et al., 2018).

161 2.4. Recovery of MBA03 genomes via phylogenetic analyses

First of all, 16S rRNA sequences were extracted from the high-quality MAGs using
Prokka (v1.14.6) (Seemann, 2014). These sequences were aligned together with
the 157 available MBA03 16S rRNA sequences from SILVA database (SSU 138.1
Ref NR) with the SINA Aligner (v1.2.11) (Pruesse et al., 2012), and afterwards a
phylogenetic tree was computed with FastTree (v2.1.1) (Price et al., 2009) and
visualized with the Interactive Tree Of Life (iTOL) (v5, Letunic & Bork, 2021). Only
the MAGs whose sequences formed a monophyletic group with MBA03 were
selected for further analysis. Due to the limitations associated to the binning
process, not all MAGs contained 16S rRNA sequences, while others contained
more than one sequence, so not all the MAGs corresponding to MBA03 could be
identified by 16S rRNA.

174 2.5. MBA03 identification from phylogenomic data

After identifying the first MAGs hypothetically belonging to MBA03, a two-step
approach was followed to recover the remaining MAGs belonging to the MBA03
taxonomic group which did not contain rRNA operons.

As a first step, all the MAGs whose 16S rRNA gene formed a monophyletic group
with MBA03 according to the phylogenetic tree were annotated using the whole
genome classifier tool Genome Taxonomy Database toolkit (GTDB-tk; v2.1.1)
(Chaumeil et al., 2022). Although GTDB does not contain any genome identified

as "MBA03", the resulting annotation was used as a first hypothesis to elucidate
which of the genomes actually corresponded to MBA03.

As a second step, all the HQ and GQ MAGs recovered in the project (n = 1053)
were also annotated GTDB-tk. In addition, a phylogenomic tree was constructed
using the UBCG pipeline (v3.0) (Na et al., 2018). Once all the MAGs
corresponding to MBA03 were identified, genomes belonging to the MBA03 order
in the GTDB were downloaded and included in the dataset. Afterwards, genomes
from the Biogasmicrobiome repository (https://biogasmicrobiome.env.dtu.dk/),
and from (Dyksma et al., 2020) were annotated and incorporated into the
dataset.

193　2.6. Taxonomic delimitation of the MBA03 taxonomic clade

Prokka (v1.14.6) (Seemann, 2014) was used again to obtain all the 16S rRNA
sequences from the MAGs and genomes identified as potential MBA03
representatives according to the previous phylogenomic analyses. To better
study the phylogenetic patterns, 16S rRNA sequences corresponding to the most
closely related taxonomic classes were set as outgroups. Sequences from class
Clostridia and *Limnochordia* were downloaded from the Living Tree Project (LTP)
and dereplicated using cd-hit (4.8.1) (W. Li & Godzik, 2006) at a 0.97 similarity
threshold. Sequences were aligned with SINA Aligner (v1.2.11) (Pruesse et al.,
2012) and incorporated into the previous dataset. Finally, a phylogenetic tree
was computed with FastTree (Price et al., 2009), and visualized with iTOL (v5;
Letunic & Bork, 2021). The reference sequence of the bacterial clone identified
for the first time as MBA03 was also downloaded from the NCBI (AB114313.1)
and included in the tree.

Following the same procedure as for the phylogenetic tree, genomes from
Clostridia and *Limnochordia* were downloaded from RefSeq NCBI Assembly
Database, filtering by "Assembly from Type Material" and "Complete Genomes"
(Supplementary Table 2). These genomes were analysed altogether with the
genomes identified as MBA03 and a phylogenomic tree was created with UBCG
(Na et al., 2018) and visualized with iTOL (v5; Letunic & Bork, 2021).

214　 2.7. Analysis of the ecological niche of MBA03

In order to explore the ecological niches of MBA03, an analysis with IMNGS (v1)
(Lagkouvardos et al., 2016) was carried out using a 97% identity cut-off. The
same sequences used for the phylogenetic analysis were dereplicated using cd-
hit (v4.8.1) (W. Li & Godzik, 2006) at a 0.9 identity threshold. IMNGS results were
filtered with a series of in-house Python scripts and manual curation
(https://github.com/roserpr/IMNGSanalysis.git). More information about the
filtering process can be found at Supplementary Methods.

In order to determine the abundance of MBA03 among the samples, coverM
(v0.5.0, https://github.com/wwood/CoverM) was used, comparing the raw reads
of the 30 sequenced reactors against the genomes assigned to MBA03.

2.8. Functional annotation

MAGs belonging to MBA03 with the higher qualities were selected (completeness > 90% and contamination limited to < 3%, both calculated with CheckM (Parks et al., 2014)). These MAGs were annotated with Bakta (Schwengers et al., 2021) and they were analysed with Kofamscan (Aramaki et al., 2020) and KEGG-Decoder (Graham et al., 2018).

As mentioned above, acetate as an intermediate is of particular importance during anaerobic reactions, especially for SAOB bacteria and hydrogenotrophic methanogenesis. In order to elucidate if MBA03 was a SAOB, the protein sequences of all the enzymes involved in the reversed WL pathway and the reversed WL pathway coupled to the glycine cleavage system (GCS) (Supplementary Table 3) were downloaded from UniProt (Coudert et al., 2023) (Swissprot), narrowing to all bacteria (taxonomy_id:2), and a database was created with Diamond (Buchfink et al., 2015). A Diamond BLASTp of the annotated protein sequences was performed against this database to detect presence and absence of each gene in all MAGs with 50% coverage and identity thresholds. For the whole metabolic analysis, any enzyme found in more than 70% of the MAGs of each family was considered to be present in the genome.

3. Results and discussion

3.1. Genomic reconstruction

MBA03 has proven abundant in a variety of anaerobic digesters (Dyksma et al., 2020; Liu et al., 2023). Given its importance in AD environments, this work focused on the recovery and characterisation of the genome of this taxonomic group using 30 metagenomic datasets derived from the MICRO4BIOGAS project and other genomic data available in public databases.

After metagenomic sequencing, assembly and binning of sludge samples, the MAGs obtained were classified according to their quality, and 1053 MAGs (HQ and GQ) were chosen for further analysis. After the annotation of the MAGs with MIGA and the metagenomes with Centrifuge, in neither case MBA03 was reported despite being a highly abundant taxon according to the metataxonomic results (Otto et al., 2023). This was expected as genomic databases did not include MBA03 genomes, probably reflecting a lack of consensus on the nomenclature of this taxonomic group.

3.2. Taxonomic analysis of MAGs

In order to identify the genomes belonging to MBA03, a phylogenetic tree was constructed using the 16S rRNA sequences extracted from the HQ MAGs and 157 sequences classified as MBA03 according to the SILVA database (SSU 138.1 Ref NR) (Supplementary Figure 1). The sequences from SILVA formed a monophyletic clade, and only 16 sequences from the metagenomic dataset fell within this subtree. The MAGs corresponding to these 16 sequences were annotated with GTDB-tk (v2.1.1) and 12 out of the 16 MAGs shared the same annotation: they

belonged to phylum Bacillota, class *Limnochordia*, order DTU010. From here, two
different families were identified: DTU010 and DTU012 (Supplementary Table 4).

After this preliminary analysis, the HQ and GQ MAGs (n=1053) were annotated
with GTDB-tk (Supplementary Table 5). A total of 45 MAGs were classified as
members of the order DTU010, and they were subsequently treated as potential
representatives of MBA03. The phylogenomic tree of HQ and GQ MAGs (Figure 1)
also revealed a large proportion of MAGs belonging to *Bacillota* (n = 503), while
104 MAGs were classified as *Bacteroidota,* 77 as *Actinobacteriota,* 63 as
Verrucomicrobiota and 32 as *Chloroflexota,* while 80 MAGs corresponded to
Archaea. These results were expected considering previous studies reporting the
average AD microbiome (Sundberg et al., 2013; Abendroth et al., 2015; Treu
et al., 2016; Kirkegaard et al., 2017; Otto et al., 2023). Moreover, it was
confirmed that MBA03 formed a monophyletic clade with all 45 sequences
annotated as DTU010. Two different subclades could be clearly distinguished,
corresponding to families DTU010 and DTU012 according to the GTDB taxonomy.

Figure 1. (A) Dendrogram of all the HQ and GQ MAGs from MICRO4BIOGAS (n=1053),
computed with UBCG and visualized with iTOL (v5). The legend shows a classification by
different colours corresponding to different phyla, according to the annotation obtained
with GTDB-tk (v2.1.1), and a highlight on MBA03 (within *Bacillota*). (B) Tree with 108
leaves, computed with FastTree and visualized with iTOL (v5), corresponding to the MAGs
belonging to order DTU010 according to GTDB-tk. Two clades corresponding to two
families (DTU010 in orange and DTU010 in yellow) can be seen, for which the names
Darwinibacteriaceae and *Wallacebacteriaceae* are proposed, respectively. In both
figures, bootstrap values greater than 0.75 are represented by a small triangle in each
branch.
294

For the enlargement of the dataset, 2237 MAGs from the Biogasmicrobiome
repository (https://biogasmicrobiome.env.dtu.dk/), 16 from Dyksma et al. and 19
from GTDB (release 207) were downloaded and annotated with GTDB-tk
(Dyksma et al., 2016). After filtering, a total of 63 potential MBA03 MAGs were
compiled. Together with the previous 45 MAGs (HQ and GQ) retrieved from
MICRO4BIOGAS data, the final dataset was composed of 108 genomes. These
MAGs were used for creating a phylogenomic tree (Figure 2), leading to an initial

taxonomic delimitation of MBA03: (i) there were two families, named DTU010 (which accounted for 82 MAGs) and DTU012 (with 26 MAGs) according to GTDB database; (ii) DTU010 clade included 20 different taxa annotated in GTDB-tk. The most common was DTU010 sp. 002391385, followed by sp. 900018335, and both formed monophyletic clades; and (iii) there were 10 different taxa belonging to DTU012, with the most common species being DTU012 sp.900019385.

Thereby, it was proposed the new order to be named *Darwinibacteriales* ord. nov., and the two families *Darwinibacteriaceae* fam. nov., and *Wallacebacteriaceae* fam. nov., after Charles Darwin and Alfred Wallace, who are considered the fathers of the Theory of Evolution by natural selection.

3.3. Final taxonomic delimitation of MBA03

According to these results, the order *Darwinibacteriales* ord. nov. (DTU010 according to GTDB) was hypothesized to be equivalent to MBA03 (SILVA's name), but further phylogenetic evidence was needed to get a proper delimitation of the clade. For that reason, phylogenetic analyses based on the 16S rRNA gene were carried out. A total of 52 16S rRNA sequences were identified by Prokka (Seemann, 2014) in the 108 MAGs (i.e., not all MAGs contained 16S rRNA sequences), and they were computed together with the 157 16S rRNA sequences downloaded from the SILVA (SSU 138.1 ref NR) database in a phylogenetic tree (n=209). Moreover, 772 sequences from *Clostridia* were added as outgroups.

The phylogenetic tree (Figure 2) confirmed the abovementioned hypothesis: the sequences annotated as MBA03 in SILVA formed a monophyletic clade with a clear subdivision of two clades, which included two distinct families (*Darwinibacteraceae*/DTU010 and *Wallacebacteriaceae*/DTU012). Clade I was formed by 52 sequences and included the MBA03 reference sequence, whereas Clade II accounted for 44 sequences.

Figure 2. Phylogenetic tree based on 16S rRNA sequence from our own database and from SILVA SSU 138.1 Ref NR database. The sequences in bold letter belong to the MAGs from our database, while the other sequences are the ones downloaded from SILVA. Output groups were collapsed for a better visualization of the MBA03 clade.

It is worth highlighting that, although both *Hydrogenispora* and *Darwinibacteriales* (MBA03) are annotated as class *Limnochordia* in the SILVA database, *Hydrogenispora* is annotated as *Clostridia* in the LPSN (List of Prokaryotic names with Standing in Nomenclature, Parte et al., 2020), which provides the internationally accepted and regulated taxonomic classification. This annotation, supported by other authors (FitzGerald, 2018; Westerholm et al., 2016), was confirmed once the tree was computed, as both *Hydrogenispora* and *Darwinibacteriales* appeared to be two different orders from class *Clostridia*. In other words; *Darwinibacterales* is proposed as a new order that belongs to class *Clostridia*.

Finally, a phylogenomic tree (Figure 3) was computed with the 108 genomes classified as MBA03 together with 206 genomes from *Clostridia* and one genome from *Limnochordia* as outgroups (Supplementary table 2). This confirmed the hypothesis that MBA03 belongs to a new order within class *Clostridia*, and it comprises two families.

Figure 3. Phylogenomic tree showing only the *Darwinibacteriales* (MBA03 in SILVA) clade, which includes two potential new families: *Darwinibacteriaceae* (DTU010) shaded in pink and *Wallacebacteriaceae* (DTU012) shaded in green. MAGs in green (n=9) belong to *Darwinibacteriales* (MBA03), as there is evidence at the genetic and genomic level. MAGs in red (n=9) may be MBA03, but their 16S rRNA phylogeny is not concordant. This could be explained by errors during the binning process. The remaining MAGs (n=90) are incomplete and/or 16S rRNA sequences could not recovered from the MAGs. Although they form a cluster with the rest of the MBA03 MAGs, they should remain as hypothetical members of this order. Output groups were collapsed for a better visualization of the MBA03 clade.

According to these results, there is sufficient evidence to confirm that MBA03 belongs to a new taxonomical order, *Darwinibacteriales* ord. nov., which includes two potential new families, *Darwinibacteriaceae* fam. nov., (DTU010 according to GTDB) and *Wallacebacteriaceae* fam. nov.,(former DTU012). It is important to note that samples from MICRO4BIOGAS provided the largest number of MAGs belonging to MBA03, meaning that the samples and datasets from this project

marked a significant breakthrough towards the genetic isolation of MBA03
genomes. The formal description of these taxa can be found in Supplementary
Material.

Once the two families from the order *Darwinibacteriales* (*Darwinibacteriaceae*
and *Wallacebacteriaceae*) were properly delimited, the read coverage of both
families in the samples was calculated. 10 MAGs from DTU010 and 4 MAGs from
DTU012 were used as reference for read mapping. The results showed that
DTU010, with a mean abundance of 4.1% and a maximum abundance of near
18%, was considerably more abundant than DTU012, with a mean abundance of
0.6% (Supplementary Table 6).

3.5. Ecological distribution *Darwinibacteriales*

The ecological distribution of *Darwinibacteriales* was evaluated using IMNGS.
This tool quantifies the prevalence and abundance of any 16S rRNA sequence of
interest introduced as input, giving information about the number of
coincidences with the samples included in its database. IMNGS also informs
about the type of sample (e.g., soil, water, anaerobic digester, environmental)
showing a hit with the input sequence. A total of 8856 SRA accessions containing
16S rRNA sequences from *Darwinibacteriales* were detected according to
sequence similarity. Due to the high amount of data obtained, only the samples
with an abundance of *Darwinibacteriales* higher than 0.1% were considered at
first (Figure 4A). An exhaustive analysis of the sample metadata revealed that
more than 99% of them belonged to anaerobic digestion studies, meaning that
Darwinibacteriales is a very specialized bacterial group that grows and develops
in this specific environment. This result reinforces the hypothesis that MBA03
Darwinibacteriales may play a central role in the AD process.

Figure 4. (A) Ecological niches of *Darwinibacteriales* based on coincidences of 16S rRNA
sequences with samples of the IMNGS database with an abundance above 0.1%, (B)
Ecological niches of *Darwinibacteriales* according to the samples with an abundance
below 0.1% and excluding all the samples related to anaerobic digestion for a better
insight into the wide range of environments from where it may be introduced to
anaerobic reactors.

To study the ecological niches besides anaerobic digestion, a similar analysis
was performed considering only the samples with an abundance of
Darwinibacteriales-related sequences below 0.1%. The resulting 8650 hits were
then filtered in two steps: (i) first, those related to anaerobic digestion were
discarded, and (ii) results with less than 5 hits in total were removed from the
dataset, leaving 1636 samples that were categorised into different groups
(Figure 4B).

Soil was the second most common environment in which MBA03 could be found,
followed by rhizosphere and sediment samples. However, the abundance of
Darwinibacteriales in these habitats was substantially lower than in the case of
anaerobic digesters. Therefore, it can be hypothesized that these bacteria are
found in soil, plants and grounds, in a quiescent or inactive form. When
agricultural residues, soils or sludges are taken as substrate for anaerobic
digestion, bacteria from the *Darwinibacteriales* order thrive in the AD
environment. Similarly, the presence of MBA03 in gut and milk microbiomes can
be explained by the dietary habits of the animals. As *Darwinibacteriales* exists in
the soil, herbivorous and omnivorous animals may ingest this group of bacteria
during their feeding. Another hypothesis would be that *Darwinibacteriales* is part
of the native gut microbiota of the animals, but at a very low abundance, and
would thus be a reservoir which would guarantee the prevalence in AD plants fed
with those substrates. Therefore, when manure is used as a substrate,
Darwinibacteriales would be introduced into the bioreactors.

3.6. Unveiling the functional potential of *Darwinibacteriales*

In anaerobic digesters, the metabolic diversity of *Darwinibacteriales* remains
challenging due to the absence of known cultivable representatives. A total of 45
and 16 MAGs from *Darwinibacteriaceae* (former DTU010) and 16 from
Wallacebacteriaceae (former DTU012), respectively, were analysed to gain
insights into the potential metabolic pathways of this bacterial order
(Supplementary Table 7).

Darwinibacteriales are chemoorganoheterotrophs that derive energy from
anaerobic respiration through oxidative phosphorylation. They utilize inorganic
molecules of ferric iron (Fe^{3+}) as potential electron acceptors (K02012 and
K02011). These bacteria also possess sulphide:quinone oxidoreductase
(EC:1.8.5.4), an enzyme involved in the oxidation of sulphides, including
hydrogen sulphide, enabling them to be chemolitotrophic. Their ability to grow
on H_2S is interesting, as H_2S is usually not oxidized under anaerobic conditions.
However, there are some exceptions.

Darwinibacteriaceae and *Wallacebacteriaceae* have complete sets of genes for
glycolysis (Embden-Meyerhof pathway; M00001), gluconeogenesis (M00003),
and pyruvate oxidation (M00307) pathways. However, they lack certain essential
genes required for the tricarboxylic acid (TCA) cycle, specifically isocitrate
dehydrogenase. Both bacterial families can convert glycogen to glucose-6P.

MAGs from both families contain genes encoding beta-N-acetylhexosaminidase for bacterial cell wall peptidoglycan degradation and recycling, pullulanase for starch degradation, and beta-glucosidase for cellobiose metabolism. *Darwinibacteriaceae* members also possess additional genes encoding chitinase for chitin degradation, D-galacturonate isomerase for pectin utilization, and alpha-amylase for starch hydrolysis. *Wallacebacteriaceae* members carry the gene encoding oligogalacturonide lyase, allowing them to degrade pectin. Both *Wallacebacteriaceae* and *Darwinibacteriaceae* exhibit the ability to degrade catechol and methylcatechol through catechol 2,3-dioxygenase enzymes (EC 1.13.11.2). However, the presence of these compounds can be toxic to microorganisms in anaerobic digesters, inhibiting the digestion process. These findings suggest that each bacterial family can exploit different organic substrates as carbon and energy sources, maybe facilitating their coexistence. This also indicates that the importance of MB03 does not only restrict to one single phase of anaerobic digestion. Apart from acetogenesis and syntrophic acetate oxidation, MB03 could also be involved in the hydrolysis phase as well.

Members of *Darwinibacteriales* assimilate nitrogen by converting NH_3 to organic nitrogen through the enzymatic activities of glutamine synthetase (EC 6.3.1.2) and glutamate (NADPH) synthase (EC 1.4.1.13). However, genes associated with other nitrogen metabolism processes, such as nitrate reduction, denitrification, nitrogen fixation, nitrification, and anaerobic ammoniumoxidation, were not observed. Most representatives of *Darwinibacteriaceae* have auxotrophic requirements for various amino acids, while *Wallacebacteriaceae* representatives are auxotrophic for a smaller set of amino acids (Supplementary Table 7).

Both bacterial families share several ABC transporters, including ABCB-BACATP binding cassettes and NitT/TauT family transporters. They also possess multiple sugar transport systems. *Darwinibacteriaceae* exhibits ABC transporters for iron (III), phosphonate, alpha-glucoside, aldouronate, sn-glycerol 3-phosphate, cobalamin, and haemolysin transport systems. Both *Darwinibacteriaceae* and *Wallacebacteriaceae* show transport systems for maltose/maltodextrin, arabinogalactan/maltooligosaccharide, raffinose/stachyose/melibiose, glucose/mannose, trehalose/maltose, cellobiose and other tansport systems (See Supplementary Table 7).

Members of *Darwinibacteriales* possess resistance genes against vancomycin (vanW) and tetracycline (tetM and tetO). They also encode multidrug efflux pumps (mdlA/smdA; mdlB/smdB; efrA/efrE) to withstand various antimicrobial compounds.

Both families are motile, as indicated by the presence of flagella synthesis genes. However, *Darwinibacteriaceae* lacks genes for the MS/C ring type III secretion system. Members of both taxa possess the enzymatic machinery for chemotactic movement and detection and transport of D-ribose and D-galactose via ABC transporters.

3.7 Members of *Darwinibacteriaceae* as potential SAOB

Members of the MBA03 lineage were previously hypothesised to be SAOB in AD
based on correlation studies (Zheng et al., 2019; Zeng et al., 2021; Perman
et al., 2022); however, the metabolism of this bacterial lineage, classified in this
study as *Darwinibacteriaceae*, remains poorly understood and characterised due
to the lack of cultured representatives. Recently, the co-occurrence of members
of *MBA03* with other potential SAO bacteria, such as DTU014 and
Syntrophaceticus, and with hydrogenotrophic methanogens such as
Methanothermobacter, *Methanoculleus* and *Methanosarcina,* has been observed
(Otto et al., 2023). This suggests that several taxonomic groups identified as
potential SAOBs, including *Darwinibacteriaceae*, grow simultaneously with
methanogens to provide a continuous supply of H_2 and CO_2 for hydrogenotrophic
methanogenesis. Usually, hydrogenotrophic methanogens are associated with
high performance plants, whereas the acetoclastic methanogens *Methanothrix*
(formally *Methanosaeta*) are found specifically in low performance systems such
as sewage sludge digesters (Sunderberg et al. 2013; Abendroth et al., 2015).
Therefore, due to the co-occurrence between members of *Darwinibacteriaceae*
and hydrogenotrophic methanogens, we hypothesise that the presence of
Darwinibacteriaceae could be associated with a higher biogas production in
industrial AD plants, although further analyses are required.

It has been seen that SAO bacteria possess either the known reversed WL
pathway or an alternative route coupled to the GCS (Zeng et al., 2021). To
confirm whether members of the *Darwinibacteriaceae* and *Wallacebacteriaceae*
are SAOBs, their MAGs were mined to identify genes of the reversed WL pathway
and the GCS (Figure 5). In all members of the *Darwinibacteriaceae* family, gene
Acka was consistently detected, whereas *Pta*, *Acs*, the *CODH* complex, and *MetF*
were conspicuously absent. Most of the *Darwinibacteriaceae* MAGs contained
FolD, *Fhs*, and *Fdh* enzymes, which play vital roles in the acquisition of CO_2 and
H_2 from formate. The absence of gene *Pta* and *Acs* could be justified by the
presence of glycine reductase (*Grd*) in the glycine cleavage system, which
facilitates the conversion of glycine for entry into the GCS (Li et al., 2022). Thus,
the entire reverse WL-GCS pathway was encoded by members of
Darwinibacteriaceae, with only two genes, namely *Pta* and *Acs* missing, which
are in fact not necessary for the completion of the pathway, as in the GCS, CH3-
CO-Pi is directly converted to glycine through the function of glycine reductase
(Grd) (Li et al., 2022). The remaining enzymes were identified in more than 70%
of the *Darwinibacteriaceae* MAGs, with the exception of *Dld*, which was present
in only 60% of them.

In the case of the family *Wallacebacteriaceae*, only genes *AckA* and *Fhs* were
identified, while the other genes of the pathway were absent. The absence of
critical enzymes of the reversed Wood-Ljungdahl pathway, including *Sda*, *Grd*,
FolD, and *Fdh* suggests that members of *Wallacebacteriaceae* are not SAOB.

Based on the evidence found, it could be hypothesized that members of the
family *Darwinibacteriaceae* are potential SAOBs, as they encode for the
necessary genes to perform the reversed WL pathway coupled to the GCS.
Within the intricate anaerobic digestion metabolic web, this pathway would allow
these bacteria to perform the oxidation of acetate to H_2 and CO_2, reaction that
would constitute a key step in AD, as hydrogenotrophic archaea would take this

H_2 as a substrate to produce methane. However, the family *Wallacebacteriaceae*
lacks several key enzymatic of this pathway. Considering the relatively low
abundance of this taxon in the analyzed bioreactors (as detailed in
Supplementary Table 6), our hypothesis suggests that *Wallacebacteriaceae* may
not play a significant role in the functioning of anaerobic digesters, unlike
Darwinibacteriaceae. This could be attributed to the absence of a symbiotic
relationship between *Wallacebacteriaceae* and hydrogenotrophic archaea.

Figure 5. Mining of reverse WL pathway (black lines) coupled with glycine cleavage
system (purple lines) (WL-GCS) genes in MAGs of the families *Darwinibacteriaceae*
(n=45) and *Wallacebacteriaceae* (n=16). Red to green colors indicate the percentage of
MAGs in which the genes were identified. A gene was considered to be present in each
family if it was found in more than 70% of the MAGs. Bacteria from the family
Darwinibacteriaceae encode the genetic repertoire necessary to metabolise acetate to
produce CO2 and H2, which are then used by hydrogenotrophic archaea to produce
methane, whereas *Wallacebacteriaceae* showed a truncated pathway. Abbreviations:
AckA; acetate kinase, *Pta*; Phosphate acetyltransferase, *Acs*; Acetyl-coa synthetase, *Por*;
pyruvate-ferredoxin reductase, *Sda*; L-serine dehydratase, *GlyA*; glycine
hydroxymethyltransferase, Grd*;* glycine reductase, *GcvP*; glycine dehydrogenase,
GcvH/T; glycine cleavage system H/T protein, *Dld*; dihydrolipoyl dehydrogenase, *PflD*;
pyruvate-formate lyase, *AcsE;* iron sulfur protein methyltransferase, *metF*;
methylenetetrahydrofolate reductase, *FolD*; methylenetetrahydrofolate
dehydrogenase/cyclohydrolase, *Fhs*; formate-tetrahydrofolate ligase, *Fdh*; formate
dehydrogenase.

4. Conclusions

In this work, a cryptic bacterial group, previously referred as MBA03 in SILVA
database, was described for the first time by recovering and analysing
representative genomes belonging to this taxon. Phylogenetic analyses
confirmed that MBA03 is a representative of a new taxonomic order with the

proposed name *Darwinibacteriales* ord. nov. Two different families can be
distinguished within this order: *Darwinibacteriaceae* fam. nov. (also known as
DTU010 according to the GTDB taxonomy) and *Wallacebacteriaceae* fam. nov.
(also known as DTU012).

Order *Darwinibacteriales,* particularly family *Darwinibacteriaceae,* is among the
most abundant genera in anaerobic digesters, regardless of operational and
chemical parameters such as feedstock and type of reactor. *Darwinibacteriales* is
a very specific bacterial group, being anaerobic digestion its main ecological
niche. Bacteria from this order are also present in soils, sediments, plants and
animal gut microbiomes at very low abundances. These materials are often used
as substrates to feed anaerobic digesters, acting as a reservoir and an inoculum
that allows *Darwinibacteriales* to thrive when the AD process begins.

The presence of the reversed WL pathway coupled to the GCS in the family
Darwinibacteriaceae and the co-ocurrence of *Darwinibacteriales* with several
hydrogenotrophic archaea (Otto et al., 2023) reveals that they are potential
SAOB, and thus that they may act as competitors for acetate with acetoclastic
archaea, favouring hydrogenotrophic methanogenesis. Furthermore, these
bacteria seem to be important during hydrolysis, since *Darwinibacteriales*
harbour a wide repertoire of hydrolytic enzymes (beta-N-acetylhexosaminidase,
pullulanase, beta-glucosidase, chitinase, alpha-amylase). Nevertheless, it is
important to highlight that this description has been entirely based on genome
characterization, so its culture and further laboratory analyses are needed to
confirm the hypotheses made.

The high abundance of *Darwinibacteriales* in AD environments strongly suggests
an exaptation mechanism to this industrial process. In other words,
Darwinibacteriales are naturally occurring hydrolytic and potentially SAO
bacteria that are pre-adapted to AD and are thus massively enriched in biogas
producing and other AD facilities, where they may play an industrially relevant
role that our results have only begun to reveal.

References

Abendroth, C., Vilanova, C., Günther, T., Luschnig, O., & Porcar, M. (2015). Eubacteria and archaea communities in seven mesophile anaerobic digester plants in Germany. Biotechnology for Biofuels, 8(1), 1-10.

Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., & Ogata, H. (2020). KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics, 36(7), 2251-2252. https://doi.org/10.1093/bioinformatics/btz859

Bowers, R. M., Kyrpides, N. C., Stepanauskas, R., Harmon-Smith, M., Doud, D., Reddy, T. B. K., Schulz, F., Jarett, J., Rivers, A. R., Eloe-Fadrosh, E. A., Tringe, S. G., Ivanova, N. N., Copeland, A., Clum, A., Becraft, E. D., Malmstrom, R. R., Birren, B., Podar, M., Bork, P., … Woyke, T. (2017). Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. Nature Biotechnology, 35(8), Article 8. https://doi.org/10.1038/nbt.3893

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. Nature Methods, 12(1), Article 1. https://doi.org/10.1038/nmeth.3176

Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2022). GTDB-Tk v2: Memory friendly classification with the genome taxonomy database. Bioinformatics, 38(23), 5315-5316. https://doi.org/10.1093/bioinformatics/btac672

Coudert, E., Gehant, S., de Castro, E., Pozzato, M., Baratin, D., Neto, T., Sigrist, C. J. A., Redaschi, N., Bridge, A., & The UniProt Consortium. (2023). Annotation of biologically relevant ligands in UniProtKB using ChEBI. Bioinformatics, 39(1), btac793. https://doi.org/10.1093/bioinformatics/btac793

De Vrieze, J., Saunders, A. M., He, Y., Fang, J., Nielsen, P. H., Verstraete, W., & Boon, N. (2015). Ammonia and temperature determine potential clustering in the anaerobic digestion microbiome. Water Research, 75, 312-323. https://doi.org/10.1016/j.watres.2015.02.025

Dyksma, S., Jansen, L., & Gallert, C. (2020). Syntrophic acetate oxidation replaces acetoclastic methanogenesis during thermophilic digestion of biowaste. Microbiome, 8(1), 105. https://doi.org/10.1186/s40168-020-00862-5

Fang, C., Zhou, L., Liu, Y., Xiong, J., Su, Y., Lan, Z., Han, L., & Huang, G. (2022). Effect of micro-aerobic conditions based on semipermeable membrane-covered on greenhouse gas emissions and bacterial community during dairy manure storage at industrial scale. Environmental Pollution, 299, 118879. https://doi.org/10.1016/j.envpol.2022.118879

Feng, Y., Wang, Y., Zhu, B., Gao, G. F., Guo, Y., & Hu, Y. (2021). Metagenome-
assembled genomes and gene catalog from the chicken gut microbiome
aid in deciphering antibiotic resistomes. Communications Biology, 4(1),
Article 1. https://doi.org/10.1038/s42003-021-02827-2

FitzGerald, J. A. (2018). The microbial ecology of anaerobic digestion:
Characterising novel biogas configurations through molecular and
statistical methods. https://hdl.handle.net/10468/7573

Graham, E. D., Heidelberg, J. F., & Tully, B. J. (2018). Potential for primary
productivity in a globally-distributed bacterial phototroph. The ISME
Journal, 12(7), Article 7. https://doi.org/10.1038/s41396-018-0091-3

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality
assessment tool for genome assemblies. Bioinformatics, 29(8), 1072-
1075. https://doi.org/10.1093/bioinformatics/btt086

Igarashi, K., & Kuwabara, T. (2016). Hydrogen-Sulfide-Free Methane
Production by Fermenter–Methanogen Syntrophy Using Dacite Pumice
under Aerobic Gas Phase. Energy & Fuels, 30(6), 4945-4950.

Jung, H., Baek, G., & Lee, C. (2020). Magnetite-assisted in situ microbial
oxidation of H2S to S0 during anaerobic digestion: A new potential for
sulfide control. Chemical Engineering Journal, 397, 124982.

Kang, D. D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., & Wang, Z. (2019).
MetaBAT 2: An adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. PeerJ, 7, e7359.
https://doi.org/10.7717/peerj.7359

Kim, D., Song, L., Breitwieser, F. P., & Salzberg, S. L. (2016). Centrifuge: Rapid
and sensitive classification of metagenomic sequences. Genome
Research, 26(12), 1721-1729. https://doi.org/10.1101/gr.210641.116

Kirkegaard, R. H., McIlroy, S. J., Kristensen, J. M., Nierychlo, M., Karst, S. M.,
Dueholm, M. S., ... & Nielsen, P. H. (2017). The impact of immigration on
microbial community composition in full-scale anaerobic digesters.
Scientific reports, 7(1), 9343.

Lagkouvardos, I., Joseph, D., Kapfhammer, M., Giritli, S., Horn, M., Haller, D., &
Clavel, T. (2016). IMNGS: A comprehensive open resource of processed
16S rRNA microbial profiles for ecology and diversity studies. Scientific
Reports, 6(1), Article 1. https://doi.org/10.1038/srep33721

Laguillaumie, L., Rafrafi, Y., Moya-Leclair, E., Delagnes, D., Dubos, S.,
Spérandio, M., Paul, E., & Dumas, C. (2022). Stability of ex situ biological
methanation of H2/CO2 with a mixed microbial culture in a pilot scale
bubble column reactor. Bioresource Technology, 354, 127180.
https://doi.org/10.1016/j.biortech.2022.127180

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with
Bowtie 2. Nature methods, 9(4), 357-359.
https://doi.org/10.1038/nmeth.1923

Lebuhn, M., Weiß, S., Munk, B., & Guebitz, G. M. (2015). Microbiology and
Molecular Biology Tools for Biogas Process Analysis, Diagnosis and
Control. En G. M. Guebitz, A. Bauer, G. Bochmann, A. Gronauer, & S.
Weiss (Eds.), Biogas Science and Technology (pp. 1-40). Springer
International Publishing. https://doi.org/10.1007/978-3-319-21993-6_1

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for
phylogenetic tree display and annotation. Nucleic Acids Research,
49(W1), W293-W296. https://doi.org/10.1093/nar/gkab301

Li, C., Hao, L., Lü, F., Duan, H., Zhang, H., & He, P. (2022). Syntrophic Acetate-
Oxidizing Microbial Consortia Enriched from Full-Scale Mesophilic Food
Waste Anaerobic Digesters Showing High Biodiversity and Functional
Redundancy. mSystems, 7(5), e00339-22.
https://doi.org/10.1128/msystems.00339-22

Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: An
ultra-fast single-node solution for large and complex metagenomics
assembly via succinct de Bruijn graph. Bioinformatics, 31(10), 1674-1676.
https://doi.org/10.1093/bioinformatics/btv033

Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing
large sets of protein or nucleotide sequences. Bioinformatics, 22(13),
1658-1659. https://doi.org/10.1093/bioinformatics/btl158

Liu, X., Zhu, X., Yellezuome, D., Liu, R., Liu, X., Sun, C., Abd-Alla, M. H., &
Rasmey, A.-H. M. (2023). Effects of adding Thermoanaerobacterium
thermosaccharolyticum in the hydrogen production stage of a two-stage
anaerobic digestion system on hydrogen-methane production and
microbial communities. Fuel, 342, 127831.
https://doi.org/10.1016/j.fuel.2023.127831

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.Journal, 17(1), Article 1.
https://doi.org/10.14806/ej.17.1.200

Na, S.-I., Kim, Y. O., Yoon, S.-H., Ha, S., Baek, I., & Chun, J. (2018). UBCG: Up- to-
date bacterial core gene set and pipeline for phylogenomic tree
reconstruction. Journal of Microbiology, 56(4), 280-285.
https://doi.org/10.1007/s12275-018-8014-6

Nwokolo, N., Mukumba, P., Obileke, K., & Enebe, M. (2020). Waste to Energy: A
Focus on the Impact of Substrate Type in Biogas Production. Processes,
8(10), Article 10. https://doi.org/10.3390/pr8101224

Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2014. Assessing
the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. Genome Research, 25: 1043-1055.

Perman, E., Schnürer, A., Björn, A., & Moestedt, J. (2022). Serial anaerobic digestion improves protein degradation and biogas production from mixed food waste. Biomass and Bioenergy, 161, 106478. https://doi.org/10.1016/j.biombioe.2022.106478

Price, M. N., Dehal, P. S., & Arkin, A. P. (2009). FastTree: Computing Large Minimum Evolution Trees with Profiles instead of a Distance Matrix. Molecular Biology and Evolution, 26(7), 1641-1650. https://doi.org/10.1093/molbev/msp077

Pruesse, E., Peplies, J., & Glöckner, F. O. (2012). SINA: Accurate high-throughput multiple sequence alignment of ribosomal RNA genes. Bioinformatics, 28(14), 1823-1829. https://doi.org/10.1093/bioinformatics/bts252

Rivière, D., Desvignes, V., Pelletier, E., Chaussonnerie, S., Guermazi, S., Weissenbach, J., Li, T., Camacho, P., & Sghir, A. (2009). Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. The ISME Journal, 3(6), Article 6. https://doi.org/10.1038/ismej.2009.2

Rodriguez-R, L. M., Gunturu, S., Harvey, W. T., Rosselló-Mora, R., Tiedje, J. M., Cole, J. R., & Konstantinidis, K. T. (2018). The Microbial Genomes Atlas (MiGA) webserver: Taxonomic and gene diversity analysis of Archaea and Bacteria at the whole genome level. Nucleic Acids Research, 46(W1), W282-W288. https://doi.org/10.1093/nar/gky467

Roopnarain, A., Rama, H., Ndaba, B., Bello-Akinosho, M., Bamuza-Pemu, E., & Adeleke, R. (2021). Unravelling the anaerobic digestion 'black box': Biotechnological approaches for process optimization. Renewable and Sustainable Energy Reviews, 152, 111717. https://doi.org/10.1016/j.rser.2021.111717

Schnürer, A. (2016). Biogas Production: Microbiology and Technology. En R. Hatti-Kaul, G. Mamo, & B. Mattiasson (Eds.), Anaerobes in Biotechnology (pp. 195-234). Springer International Publishing. https://doi.org/10.1007/10_2016_5

Schwengers, O., Jelonek, L., Dieckmann, M. A., Beyvers, S., Blom, J., & Goesmann, A. (2021). Bakta: Rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. Microbial Genomics, 7(11), 000685. https://doi.org/10.1099/mgen.0.000685

Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. Bioinformatics, 30(14), 2068-2069. https://doi.org/10.1093/bioinformatics/btu153

Sieber, C. M. K., Probst, A. J., Sharrar, A., Thomas, B. C., Hess, M., Tringe, S. G., & Banfield, J. F. (2018). Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nature Microbiology, 3(7), Article 7. https://doi.org/10.1038/s41564-018-0171-1

Sundberg, C., Al-Soud, W. A., Larsson, M., Alm, E., Yekta, S. S., Svensson, B. H.,
... & Karlsson, A. (2013). 454 pyrosequencing analyses of bacterial and
archaeal richness in 21 full-scale biogas digesters. FEMS microbiology
ecology, 85(3), 612-626.

Treu, L., Kougias, P. G., Campanaro, S., Bassani, I., & Angelidaki, I. (2016).
Deeper insight into the structure of the anaerobic digestion microbial
community; the biogas microbiome database is expanded with 157 new
genomes. Bioresource Technology, 216, 260-266.
https://doi.org/10.1016/j.biortech.2016.05.081

Tsang, Y. F., Kumar, V., Samadar, P., Yang, Y., Lee, J., Ok, Y. S., Song, H., Kim,
K.-H., Kwon, E. E., & Jeon, Y. J. (2019). Production of bioplastic through
food waste valorization. Environment International, 127, 625-644.
https://doi.org/10.1016/j.envint.2019.03.076

Van der Zee, F. P., Villaverde, S., Garcia, P. A., & Polanco, F. F. (2007). Sulfide
removal by moderate oxygenation of anaerobic sludge environments.
Bioresource technology, 98(3), 518-524.

Wang, Y., Wegener, G., Williams, T. A., Xie, R., Hou, J., Tian, C., ... & Xiao, X.
(2021). A methylotrophic origin of methanogenesis and early divergence
of anaerobic multicarbon alkane metabolism. Science advances, 7(27),
eabj1453.

Westerholm, M., Moestedt, J., & Schnürer, A. (2016). Biogas production through
syntrophic acetate oxidation and deliberate operating strategies for
improved digester performance. Applied Energy, 179, 124-135.
https://doi.org/10.1016/j.apenergy.2016.06.061

Whitman, W. B., Bowen, T. L., & Boone, D. R. (2006). The methanogenic
bacteria. Prokaryotes, 3(Chapter 9), 165-207.

Wu, Y.-W., Simmons, B. A., & Singer, S. W. (2016). MaxBin 2.0: An automated
binning algorithm to recover genomes from multiple metagenomic
datasets. Bioinformatics, 32(4), 605-607.
https://doi.org/10.1093/bioinformatics/btv638

Yaqoob, H., Teoh, Y. H., Ud Din, Z., Sabah, N. U., Jamil, M. A., Mujtaba, M. A., &
Abid, A. (2021). The potential of sustainable biogas production from
biomass waste for power generation in Pakistan. Journal of Cleaner
Production, 307, 127250. https://doi.org/10.1016/j.jclepro.2021.127250

Yekta, S. S., Ziels, R. M., Björn, A., Skyllberg, U., Ejlertsson, J., Karlsson, A., &
Svensson, B. H. (2017). Importance of sulfide interaction with iron as
regulator of the microbial community in biogas reactors and its effect on
methanogenesis, volatile fatty acids turnover, and syntrophic long-chain
fatty acids degradation. Journal of bioscience and bioengineering, 123(5),
597-605.

Zeng, Y., Zheng, D., Gou, M., Xia, Z.-Y., Chen, Y.-T., Nobu, M. K., & Tang, Y.-Q.
(2021). Chasing the metabolism of novel syntrophic acetate-oxidizing

bacteria in thermophilic methanogenic chemostats (p.
2021.07.06.451242). bioRxiv. https://doi.org/10.1101/2021.07.06.451242

Zheng, D., Wang, H.-Z., Gou, M., Nobu, M. K., Narihiro, T., Hu, B., Nie, Y., &
Tang, Y.-Q. (2019). Identification of novel potential acetate-oxidizing
bacteria in thermophilic methanogenic chemostats by DNA stable isotope
probing. Applied Microbiology and Biotechnology, 103(20), 8631-8645.
https://doi.org/10.1007/s00253-019-10078-9

Supplementary Methods

Sample description

As part of the MICRO4BIOGAS project, a total of 80 samples were collected from
anaerobic digestion systems at 45 large-scale reactors with different operational
conditions and feedstocks. These samples, together with the corresponding
metadata, were used to investigate the chemical and taxonomic profiles (Otto et
al., 2023). The first 40 samples were collected by members of the Technische
Universität Dresden (TUD, Dresden, Germany) from 25 different biogas plants in
Germany and Austria, while the second set of 40 samples was collected by the
team members from Bioclear Earth B.V. (Groningen, Netherlands) from 20
different biogas plants in the Netherlands. A subset of 30 of these samples was
selected to be analysed through metagenomic sequencing. This subset included
samples enriched in MBA03, hydrolytic, acidogenic and acetogenic bacteria
and/or methanogenic archaea. A full description of the samples can be found in
Supplementary Table 1.

DNA extraction and sequencing

The sludge samples were washed before DNA extraction to reduce the number of
inhibiting substances. Specifically, 5-10 mL of each sample was mixed with 40
mL of sterile phosphate-buffered saline (PBS). Subsequently, the samples were
centrifuged at 20,000 g for 10 min and the supernatants were discarded. This
process was repeated twice, and the final pellets were used for DNA extraction
with the DNeasy® PowerSoil® Pro kit (QIAGEN, Germany) based on the
manufacturer's instructions, with an extra incubation step at 65 °C for 10 min
after adding the C1 solution to increase the efficiency of cell lysis. Qubit x1
dsDNA HS Assay kit (Qubit 4.0 Fluorometer, Thermo Fisher) was used for DNA
quantification. The metagenomic DNA was randomly fragmented by sonication
and DNA fragments were end polished, A-tailed, and ligated with the full-length
adapters for Illumina sequencing: 5'-AGA TCG GAA GAG CGT CGT GTA GGG AAA
GAG TGT AGA TCT CGG TGG TCG CCG TAT CAT T -3' (5' Adapter), and 5'- GAT
CGG AAG AGC ACA CGT CTG AAC TCC AGT CAC GGA TGA CTA TCT CGT ATG CCG
TCT TCT GCT TG -3' (3' adapter). PCR amplification with P5 and indexed P7 oligos
was carried out, and the PCR products were purified with the AMPure XP system.
Then, the libraries were checked for size distribution by an Agilent 2100
Bioanalyzer (Agilent Technologies, CA, USA), and quantified by real-time PCR.
Libraries were sequenced at Novogene (UK) using the NovaSeq 6000 Illumina
platform (150 bp x 2). The average sequencing depth was 45M reads/sample
(min.: 35M reads; max.: 55M reads).

IMNGS filtering

IMNGS results were filtered with the objective of gathering all the samples by
their sample type to see which were the potentially relevant ecological niches of
MBA03. As tags describing ecological niches were very diverse and, in many
cases, ambiguous, a series of in-house Python scripts were needed to carry out a

reclassification of niche categories
(https://github.com/roserpr/IMNGSanalysis.git).

To briefly explain the filtering process, samples containing the 16S rRNA
sequence of interest in an abundance <0.1% were removed from the analyses.
To reclassify the remaining samples, two subsequent strategies were followed:
(1) description tags were filtered by the words "anaerobic", "digester", "biogas",
and "reactor", assigning all of them to the same tag: "anaerobic digestion"; (2)
the information of the remaining samples were scraped from NCBI. The output
was manually curated to determine which samples belonged to "anaerobic
digestion", and other general tags were also reassigned to more specific classes
(i.e., "metagenome" to "soil metagenome") depending on the project
description.