Cohesin reconstitution and homologous recombination repair of DNA double strand breaks in late mitosis

Jessel Ayra-Plasencia[1,2,#,†,*], Sara Medina-Suárez[1,2,†], Esperanza Hernández-Carralero[1], Jonay García-Luis[2], Lorraine S Symington[3], and Félix Machín[1,2,4,*]

[1] Hospital Universitario Nuestra Señora de Candelaria, Instituto de Investigación Sanitaria de Canarias (IISC), 38010 Santa Cruz De Tenerife, Spain.
[2] Instituto of Tecnologías Biomédicas, Universidad de La Laguna, 38200 San Cristóbal de La Laguna, Spain.
[3] Department of Microbiology and Immunology, Columbia University Medical Center, New York, United States.
[4] Facultad de Ciencias de la Salud, Universidad Fernando Pessoa Canarias, Las Palmas de Gran Canaria, Spain.

[#] Present address: Unidad de Investigación, Hospital Universitario de Canarias, 38320 San Cristóbal de La Laguna, Spain

[†] These authors contributed equally.

[*] Co-corresponding authors: jayra@fciisc.es; fmachin@fciisc.es

Abstract

The cohesin complex maintains sister chromatid cohesion from S phase to anaphase onset. Cohesin also plays roles in chromosome structure and DNA repair. At anaphase onset, the cohesin subunit Scc1 is cleaved to allow segregation in an orderly manner, although some residual cohesin subunits remain to maintain chromosome structure. Efficient DNA double strand break (DSB) repair by homologous recombination (HR) with the sister chromatid also depends on cohesin. Here, we have examined the role of residual cohesin in DSB repair in telophase (late mitosis). We have found that Scc1 returns in telophase after DSBs, and that it partially reconstitutes a chromatin-bound cohesin complex with Smc1 and an acetylated pool of Smc3 after a single HO-induced DSB at the *MAT* locus. However, this new cohesin is neither required for the HR-driven *MAT* switching nor binds to the *MAT* locus after the DSB.

Keywords

DNA double strand break; anaphase; telophase; cohesin; homologous recombination; Scc1/Mcd1; acetylated Smc3

Introduction

Cell survival and genome integrity depend on the faithful segregation of chromosomes in anaphase. From the S phase until the anaphase onset, the highly conserved cohesin complex plays an essential role in the structural maintenance of sister chromatids. Cohesin is a multiprotein complex composed in yeast of Smc1, Smc3, Scc1 (also known as Mcd1), Scc3 and Pds5 subunits (Hirano, 2000; Michaelis et al., 1997; Sjögren and Nasmyth, 2001). At the core of the complex, Smc1, Smc3 and Scc1 form a heterotrimeric ring that embraces and holds the replicated sister chromatids together and well-aligned until reaching G_2/M (Díaz-Martínez et al., 2008; Koshland and Guacci, 2000; Laloraya et al., 2000; Michaelis et al., 1997; Strunnikov et al., 1993). Cohesin subunits are loaded onto chromosomes in G1 by the Scc2-Scc3 complex (Ciosk et al., 2000). When cells enter S phase, the acetyl-transferase Eco1 acetylates Smc3, which inhibits the ATPase activity of Smc1-Smc3 heads and prevents the opening of the Smc3-Scc1 interface (Çamdere et al., 2015; Chan et al., 2012; Huber et al., 2016; Murayama and Uhlmann, 2015; Ström et al., 2007; Unal et al., 2007). This embraces the nascent sister chromatids and establishes cohesion (Uhlmann and Nasmyth, 1998). Once sister chromatids are ready for segregation, the APC^{Cdc20} (Anaphase Promoting Complex associated with its regulatory cofactor Cdc20) initiates anaphase by degrading securin/Pds1, so that the separase/Esp1 becomes active. Separase then cleaves the Scc1 subunit by proteolysis, releasing sister chromatids from cohesion (Cohen-Fix et al., 1996; Michaelis et al., 1997; Nasmyth and Haering, 2009; Uhlmann et al., 1999; Yamamoto et al., 1996). The fragmented Scc1 is then rapidly degraded (Rao et al., 2001), while a pool of Smc1-Smc3 dimers appears to remain through anaphase (Renshaw et al., 2010; Tanaka et al., 1999). This pool however becomes loose as Smc3 is deacetylated by Hos1 in anaphase (Chan et al., 2012; Huber et al., 2016; Murayama and Uhlmann, 2015). Interestingly, some cohesin-dependent cohesion remains at chromosome arms during anaphase, suggesting that residual cohesin is present despite separase activation (Djeghloul et al., 2020; Garcia-Luis et al., 2022; Renshaw et al., 2010).

In addition to its role in sister chromatid cohesion, cohesin has also been involved in chromosome structure and DNA repair. On the one hand, cohesin can hold two DNA segments within the same chromatin forming an extruded loop (Lazar-Stefanita et al., 2017; Rao et al., 2017; Schalbetter et al., 2017). On the other hand, DNA double strand breaks (DSBs) produce the so-called damage-induced cohesion (DI-cohesion) (Kim et al., 2010). Following DNA damage, cohesin is recruited and accumulated both along 50 – 100 kb surrounding the DSB site and genome-wide (Ström et al., 2004; Ünal et al., 2004). The cohesin recruitment creates a firm anchoring of two well-aligned sister chromatids, which in turn facilitates DSB repair by homologous recombination (HR) (Hou et al., 2022; Phipps and Dubrana, 2022). In this regard, DNA damage kinases such as Mec1 and Chk1 phosphorylate Scc1 to initiate the cohesion establishment in post-replicative cells (Heidinger-Pauli et al., 2008).

HR is a DSB repair mechanism that uses a homologous template for restoring the original broken DNA. HR is highly reliable when the correct template is used, which in mitotic cells is the sister chromatid. Hence, HR is the preferred repair mechanism from S to G2/M, when the sister chromatid is available in close proximity (Langerak and Russell, 2011; Mathiasen and Lisby, 2014; Symington et al., 2014). Before DNA replication, in G1, the alternative error-prone non-homologous end joining (NHEJ) is used instead. Cells coordinate

the choice of either NHEJ or HR based on the activity of cyclin dependent kinase (CDK).
Low CDK keeps cells in G1 and favours NHEJ, whereas high CDK is present in S and G2/M
and activates HR. Noticeably, there is a paradox in this link between high CDK and HR. In
the cell cycle window that spans from anaphase to the telophase-to-G$_1$ transition (we will
refer to this window as late mitosis), high CDK is set to favour HR despite a sister chromatid
is not in proximity (Machín and Ayra-Plasencia, 2020). In a previous study, we showed that
sister loci can indeed move closer and coalesce, and that HR still appears important for DSB
survival in late mitosis (Ayra-Plasencia and Machín, 2019). Considering all these
observations, we wondered whether the residual cohesin is affected by DSBs in late mitosis
(telophase) and whether this cohesin pool is important for HR repair.

Results and Discussion
103

Scc1 becomes stable after DSBs in late mitosis.

At the anaphase onset, Esp1 cleaves Scc1, opening the cohesin ring and releasing
sister chromatids from cohesion. Thereafter, Scc1 needs to be translated *de novo* since
cleaved fragments are unstable and degraded by the proteasome (Rao et al., 2001).
However, it is known that activation of Esp1 can be blocked by DDC kinases (Yam et al.,
2020). In addition, DDC kinases also phosphorylate Scc1 to facilitate DI-cohesion at a post-
replicative stage (Heidinger-Pauli et al., 2008). Thus, it is feasible that DSB generation in late
mitosis can stabilize *de novo* Scc1 as well as rendering it cohesive. Noteworthy, Smc1 and
Smc3 subunits remain attached to chromatin after segregation (Garcia-Luis et al., 2022;
Renshaw et al., 2010; Tanaka et al., 1999).

To assess how cohesin subunits behave after DSBs in late mitosis, *cdc15-2* strains
that bear Scc1 tagged with myc epitopes and Smc1/3 tagged with HA epitopes were first
blocked in telophase at 34 ºC for 3 hours (Tel) (Fig. 1a). The *cdc15-2* allele encodes a
conditionally thermosensitive Cdc15 kinase, which drives telophase-to-G1 transition (Machín
and Ayra-Plasencia, 2020). These strains also harbour an inducible HO endonuclease,
which cuts the HO cutting site (*HOcs*) at the *MAT* locus in chromosome III. The HO gene is
under the control of a promoter system that responds to β-estradiol (*lexO4p:HO*:plus LexA-
ER-B112-T) (Gnügge and Symington, 2020; Ottoz et al., 2014). In wild type strains, the HO
DSB is repaired by HR with the intramolecular ectopic loci *HMR* and *HML*. The locus
election depends on the *MAT* allele (the *MAT*a HO DSB repairs with *HML*) and results in a
gene conversion (e.g., from *MAT*a to *MAT*α). In order to undoubtedly attribute the effects on
cohesin to the DSB and not to downstream events of the repair process, we used a set of
strains unable to repair the HO-mediated DSB (Δ*hml* Δ*hmr* double mutant). After the Tel
arrest, the cultures were divided into three. The first subculture served as a mock control, the
second was used to generate the single HO DSB (+HO), and the third was used to generate
multiple random DSBs by means of the radiomimetic drug phleomycin (+phle). Noteworthy,
these latter DSBs are repairable in principle. Incubation of the subcultures was prolonged at
34 ºC for 2 extra hours.

The presence or absence of Scc1-3myc, Smc1-6HA, and Smc3-6HA was checked
by Western blot. As controls, we included asynchronous samples from the same strains,

before the telophase arrest, as well as a parallel arrest at G2/M by the drug nocodazole (Nz).
As expected, Smc1, Smc3 and Scc1 bands were detected in both cycling (asyn) cells and
cells blocked in G2/M (Fig. 1b-d). The Scc1 signal disappeared almost entirely when cells
were arrested in telophase and remained low in the mock control two hours later. Strikingly,
full-length Scc1 was restored after single and multiple DSBs generation (Fig. 1b,e).
Compared to the mock condition, phleomycin treatment increased Scc1 levels fivefold,
reaching approximately half of the levels observed in G2/M. In the case of HO DSBs, Scc1
levels surpassed those of the mock by more than fifteenfold and were even higher than in
G2/M. In contrast, Smc1 and Smc3 levels remained more stable in both types of DSBs (Fig.
1b-d).

The cohesin complex is reconstituted and binds to chromatin after the HO DSBs in
telophase.

The above findings suggest that the cohesin complex could be newly formed and
loaded onto the breaks, promoting DI-cohesion, which could in turn ease the
recombinational events. To assess whether cohesin subunits are reassembled after DSB
induction in telophase, we performed a co-immunoprecipitation (co-IP) experiment using
Smc1-6HA as the bait protein in a strain that also expresses Scc1-3myc (Fig. 2a and S1). In
our experimental setup, in addition to the anti-HA and anti-myc antibodies, we included an
antibody for the acetylated form of Smc3 at lysine residues K112 and K113 (acSmc3), which
have been linked to the chromatin-bound pool of the Smc1-Smc3 dimer (Beckouët et al.,
2010; Minamino et al., 2015; Rolef Ben-Shahar et al., 2008; Unal et al., 2008). In
asynchronous cultures, both Scc1-3myc and acSmc3 co-immunoprecipitated with Smc1-
6HA, indicating that we were able to pull down the chromatin-bound pool (acetylated Smc3)
of the canonical Smc1-Smc3-Scc1 complex. A parallel co-IP using untagged Smc1 did not
recover the other subunits, confirming the specificity of the interaction. As expected, Scc1
was not detected in the pull-down under the mock condition (no DSBs); however, it was
recovered after generation of the HO DSB. Surprisingly, this was not the case when random
DSBs were induced by phleomycin. Incidentally, acSmc3 appeared as two bands in the
asynchronous culture but as a single upper band in all telophase conditions, suggesting a
difference in posttranslational modifications at this cell cycle stage.

The reconstitution of the complex and the detection of the acetylated form of Smc3,
prompted us to confirm whether the newly formed Scc1 was bound to chromatin, since the
coexistence of two pools could not be ruled out based on the co-IP experiments alone. For
instance, a soluble Smc1-Smc3-Scc1 complex could coexist with a chromatin-bound Smc1-
acSmc3 dimer, which could represent up to 30% of the total Smc1-Smc3 in anaphase
(Beckouët et al., 2010). To this end, we performed biochemical fractionation of the
chromatin-bound proteins, observing an increase in all cohesin subunits in the chromatin
fraction after the HO DSB (Fig. 2b and S2). The increase was approximately twofold for
Smc1-acSmc3 and up to sevenfold for Scc1. These results confirmed that the new Scc1
binds to a reconstituted chromatin-bound cohesin complex following the HO DSB. In
contrast, this chromatin recruitment was not observed after phleomycin treatment. The
differences in co-IP and fractionation due to the source and nature of the DSBs are
intriguing. In this experimental setup, the HO DSB is irreparable, whereas DSBs caused by

phleomycin can eventually be repaired by anaphase regression and sister loci coalescence
(Ayra-Plasencia and Machín, 2019). In addition, the HO DSB leaves clean ends that can be
easily resected, while the DSBs after phleomycin are chemically modified and require
extended processing.

Next, we used chromatin immunoprecipitation (ChIP) to investigate whether the HO
DSB recruits more cohesin to the vicinity of the *HOcs*. This recruitment has been reported
during DSBs that exert a G2/M arrest by the DDC (Ström et al., 2004), Because the level of
Scc1 changes so dramatically after the HO DSB, we selected one of the subunits that does
not for the ChIP. An Smc3-HA strain unable to repair the HO DSB (Δ*hml* Δ*hmr* double
mutant) was arrested in both G2/M and telophase. We confirmed an increase in the Smc3-
bound pool around the *HOcs* in G2/M after the DSB (Fig. S3), however such an increase
was not observed in telophase (Fig. 2c). We thus conclude that the core of the cohesin
SMCs (Smc1-Smc3) are not further recruited to the HO DSB in late mitosis. This lack of local
recruitment could be due to several non-exclusive scenarios, including the inability to
increase the acSmc3 pools by Eco1, whose levels drop after G2/M (Lyons and Morgan,
2011), as well as condensin binding to chromosome arms during anaphase (Leonard et al.,
2015), which might act as an eviction barrier for cohesin loading. However, reassembly of
the full cohesin complex at residual Smc1-acSmc3 dimers may shift the organization of the
surrounding chromosome regions and impinges on the HO DSB repair in telophase.

197
198
199 **The HO-mediated DSB is efficiently repaired by HR in late mitosis.**
200
In a previous report, we showed from both genetic and cytological points of view that
HR was still active in late mitosis (Ayra-Plasencia and Machín, 2019). Having observed that
cohesin is reconstituted and chromatin-bound after the HO DSB in telophase, we next
wondered about the functional role of the complex in this context.
We first tested whether the HO-induced DSB is efficiently resected into 3' ssDNA
tails in late mitosis. Resection is necessary to form the protruding 3' Rad51 nucleoprotein
filaments that invade the donor homologous sequence to restore the break by HR (Peng et
al., 2021; Symington, 2016). To confirm and quantitate the formation of ssDNA flanking the
HOcs, we used a qPCR approach whereby primers can only amplify a target DNA cut with
the restriction enzyme *StyI* if the target has been rendered single-stranded by resection (Fig.
S4) (Gnügge et al., 2018; Zierhut and Diffley, 2008). We used the Δ*hml* Δ*hmr* double mutant
to hamper the normal HR flow from resection to invasion and thus boosts ssDNA detection.
Two different locations downstream of the *HOcs* were monitored to determine both the initial
resection (0.7 Kb from *HOcs*) and the kinetics further away (5.7 Kb). In addition, cells
arrested in either G2/M or telophase were directly compared to each other. Cells were first
arrested for 3 h, the HO was induced afterwards, and samples were taken every hour during
a 4 h time course while sustaining the HO induction (Fig. 3a). As expected, cells in G2/M
efficiently resected the *HOcs* ends. The time to reach half of the maximum resection ($t_{1/2}$)
was ~1 h at 0.7 Kb and ~2.5 h at 5.7 Kb from the DSB, respectively. Resection in telophase
was only slightly delayed at 0.7 Kb ($t_{1/2}$ ~1.5 h), whereas the delay was more pronounced at
5.7 Kb ($t_{1/2}$ ~ 3.5 h). The delay may be due to either the fact that CDK activity in late mitosis,
though still high, is slightly inferior than that at G2/M or the different chromosome
compaction degrees between the arrests. Together with the samples for the qPCR, samples

for Western blot of Rad53 were taken. Rad53 is an effector kinase of the DDC that gets activated by hyperphosphorylation after being recruited to the 3' ssDNA overhands (Branzei and Foiani, 2006). In agreement with the resection profile, Rad53 hyperphosphorylation occurred in late mitosis, although with ~1 h delay (Fig. 3b). This delay is also consistent with previous studies on Rad53 activation upon an unrepairable HO DSB (Pellicioli et al., 2001).

Once resection begins, cells are normally committed to recombine because long ssDNA fragments do not work as efficient substrates for NHEJ. In our haploid strain, HR at the *MAT* locus after the HO-driven DSB implies the gene conversion from the *MAT*a to the *MAT*α allele, whereas repair by NHEJ results in reconstitution of the *MAT*a allele (Fig. 3c) (Yamaguchi and Haber, 2021). This gene conversion can be detected by Southern blot because of a restriction fragment length polymorphism for *StyI*. Cells that are in G1 can barely repair the HO DSB, whereas those in G2/M do so through HR (Ira et al., 2004). To check whether HR was efficient at the HO DSB in late mitosis, we performed an experimental setup similar to the one shown above, but with two important differences. On the one hand, the strain retains the *HML* locus, enabling HR if engaged, and on the other hand, the HO endonuclease was transiently induced by 1 h β-estradiol pulse, allowing time for the complete repair of the induced DSB. Samples were taken at the time of the arrest, after the pulse, and all through the repair window. We confirmed by Western blot that HO (HO tagged with the Flag epitope) is produced after the pulse, and is rapidly degraded afterwards (Fig. 3d, HO-Flag blot). Rad53 hyperphosphorylation followed the ~1 h delay shown above, and became prominent by the time the DSB inducer had been removed (Fig. 3d, Rad53 blot), suggesting extensive resection even in this repairable setup. The Southern blot showed that (i) most *MAT*a locus in the cell population has been efficiently cut after the HO pulse (Fig. 3e and f; decrease of the *MAT*a band intensity and rise of that of the cut *HOcs*); and (ii) most DSBs are repaired afterwards exclusively through HR (Fig. 3e and f; drop of the HO cut band intensity and rise of that of the *MAT*α). No signs of DSB repair by NHEJ were observed (the remaining *MAT*a band stays constant throughout). This was further confirmed by measuring the repair kinetics of a derivative strain deficient in NHEJ (*yku70*Δ; Fig. S2a-c), and is consistent with our previous studies with a HR-deficient *rad52*Δ strain (Medina-Suárez et al., 2024). Interestingly, HR kinetics were not altered in cells unable to activate the DDC (*rad9*Δ; Fig. S2d-f) or to tether the DSB ends (*mre11*Δ; Fig. S2g-i). These results are also consistent with previous studies in G2/M cells, in which HR-driven repair of the HO-DSB appeared more robust than other types of DSBs (Lazzaro et al., 2008; Moreau et al., 2001).

Overall, we concluded that HR is the chosen repair pathway in late mitosis. This is in agreement with predictions based on CDK activity and also fits well with the genetic and cytological data we have obtained before (Ayra-Plasencia and Machín, 2019; Machín and Ayra-Plasencia, 2020; Medina-Suárez et al., 2024). By contrast, this emphasizes the paradoxes of having HR in the context of mostly segregated sister chromatids and without the cohesin complex holding them together as expected. We have now explored the second paradox further as we have just shown that cohesin is reconstituted after DSBs in telophase, and DI-cohesion appears important for efficient DSB repair by HR (Hou et al., 2022; Phipps and Dubrana, 2022).

The MAT switching in late mitosis does not depend on cohesin.

In cells arrested in *cdc15-2*, the Smc1-Smc3 pool mostly resides at centromeres where it brings together adjacent regions in both chromosome arms (Garcia-Luis et al., 2022). Of note, this configuration could favour the physical interaction in *cis* of the *MAT* and the *HML* loci (Piazza et al., 2021), which may in turn promote a rapid and efficient HR-driven *MAT* switching. Thus, we tested whether *MAT* switching was dependent on this residual centromeric cohesin. To this aim, we tagged the Smc3 subunit with the auxin-mediated degron system (aid*). To check that the system was working correctly, a serial dilution spot assay was carried out (Fig. 4a). The strain expressing Smc3-aid* was plated on YPDA and YPDA plus 8 mM of the auxin indole-acetic acid (IAA). Alongside, we also plated a derivative strain where we had removed the preceptive F-box protein OsTIR1, which forms the functional ubiquitin ligase responsible for targeting the aid* for degradation upon IAA addition (Morawska and Ulrich, 2013). Since cohesin is an essential complex for vegetative growth, the lack of growth in the *SMC3-AID* OsTIR1* strain strongly points out that Smc3-aid* is degraded efficiently.

Next, we used this strain to degrade Smc3 after cells have been arrested in late mitosis and before the HO DSB. After IAA addition, there was a clear decline in Smc3-aid* levels, including the chromatin-bound acSmc3-aid* pool (Fig. S3a). After Smc3 depletion, a 1 h pulse of HO expression was applied, which allowed us to follow both DSB generation and its subsequent repair without residual cohesin. More than 90% of the Smc3-aid* had disappeared by the time the HO promoter was shut down (Fig. S3a). The addition of IAA drastically delayed gene conversion from *MAT*a to *MAT*α (Fig. S3b,c), although the cut efficiency was lower than in the wild type in the absence of auxin (~ 50 % *MAT*a remained uncut in the *SMC3:aid* OsTIR1* strain). Also, the detection of DNA damage through Rad53 phosphorylation seems fainter and slightly delayed (Fig. S3a). To unequivocally determine whether *MAT* switching kinetics were affected by Smc3 depletion, and not by an off-target effect of IAA, we compared strains that could either degrade or retain Smc3 in the presence of IAA. Four isogenic strains were analysed, comprising all possible combinations of Smc3-aid* and OsTir1, as confirmed by Western blot (Fig. 4b; Tel samples). The addition of IAA induced Smc3-aid* degradation only in the strain co-expressing OsTIR1 (Fig. 4b; +IAA samples). In the wild type strain (no tagged Smc3 and no OsTIR1), the cutting efficiency of *HOcs* and the yield of gene conversion to *MAT*α were both reduced in the presence of IAA, confirming that IAA itself negatively impacts *MAT* switching (Fig. 4c, compared two leftmost lanes with Fig. 3e). However, when the yield of the *MAT* conversion after the recovery time was normalized to the cut *HOcs* when HO was repressed, no significant differences were observed among the four strains (Fig. 4c,d). We therefore conclude that the core of the cohesin complex is not required for the HR-driven repair of the HO-DSB in telophase. Of note, it was previously reported that cohesin was not required for the *MAT* switching in G2/M-arrested cells, despite it being essential for efficient post-replicative DSB repair (Ünal et al., 2004).

Conclusions and perspectives

In this work we have addressed whether HR was molecularly efficient in yeast cells arrested in late mitosis. We have used the well-established *MAT* switching system, and found that HR is indeed working in late mitosis, complementing previous results that pointed

in this direction both genetically and cytologically (Ayra-Plasencia and Machín, 2019). In
addition, we have addressed the paradox of having HR in a cell mitotic stage with no sister
chromatid cohesion. Our results suggest that residual cohesin, or at least residual
chromatin-bound Smc1-Smc3 dimers, are complemented by incorporating *de novo* Scc1
(Fig. 4e). This fact suggests that chromosome structure may undergo significant changes
after late DSBs. The nature of these changes remains unexplored and we cannot currently
rule out that some kind of DI-cohesion could be established, perhaps globally. Alternatively,
telophase chromatin may acquire new intramolecular loops as it occurs in G2/M-arrested
cells (Piazza et al., 2021). Either of these may favour repair by HR with the intact sister or
intramolecularly, although it seems still dispensable for *MAT* switching at this late cell cycle
stage. Further research will be needed to confirm these hypotheses, as well as the
whereabouts of cohesin and the consequences for late segregation after DSB repair.

Material and Methods

Strains and experimental conditions.

The table 1 contains the *S. cerevisiae* strains used in this work. Genetic backgrounds
are either W303 or YPH499. Strain construction was performed by lithium acetate-based
transformation on pre-made frozen competent cells (Knop et al., 1999).

Table 1. Strains used in this work.

Strain	Relevant Genotype[1]	Origin	Figure
FM2344	(YPH499) *MATa ura3-52 lys2-801 ade2-101 trp1-Δ63 his3-Δ200 leu2-Δ1 bar1-Δ; cdc15-2:9myc::Hph; HMLα Δhmr::HIS3MX*	Machín lab	(parental)
LSY4319-1C	(W303) *MATα leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15 RAD5 Δhml Δhmr; leu2-3::lexO4p:HO:FLAG:CYC1t::ACT1p:LexA-ER-B112-T:CYC1p::LEU2MX* (pRG646)	Symington lab	(parental)
FM2450	LSY4319-1C; *cdc15-2:9myc::Hph*	This work	3a,b
FM2520	FM2450; *SCC1:3myc::HIS3MX*	This work	1b,e
FM2531	FM2344; *leu2-3::lexO4p:HO:FLAG:CYC1t::ACT1p:LexA-ER-B112-T:CYC1p::LEU2MX*	This work	3d-f; 4b-d
FM2635	FM2450; *SMC1:6HA:HIS3MX*	This work	1c,e
FM2662	FM2531; *Δrad9::KanMX4*	This work	S5
FM2663	FM2531; *Δyku70::KanMX4*	This work	S5
FM2668	FM2531; *Δmre11::KanMX4*	This work	S5

FM2672	FM2531; *SMC3:AID*:9myc::KanMX*	This work	4a-d
FM2674	FM2450; *SMC3:3HA::HIS3MX*	This work	1d; 2c; S2; S3
FM2680	FM2672; *ura3-52::ADH1p:OsTIR1:9myc::URA3*	This work	4a-d; S6
FM3235	FM2520; *SMC1:6HA::NatNT2*	This work	2a,b; S1
FM3294	FM2531; *ura3-52::ADH1p:OsTIR1:9myc::URA3*	This work	4b-d

[1] Semicolons separate genetic modifications obtained through sequential transformation
steps. Intermediate strains are omitted.

Strains were cultured overnight in the YPD medium (10 g·L^{-1} yeast extract, 20 g·L^{-1}
peptone and 20 g·L^{-1} glucose) with moderate shaking (180 rpm). The day after, 10-100 µL of
grown cells were diluted into a flask containing an appropriate YPD volume and grown
overnight at 25 ºC again. Finally, the exponentially growing culture was adjusted to OD$_{600}$ =
0.5 to start the experiment.

In general, cells were first arrested in late mitosis by incubating the culture at 34 ºC
for 3 hours (all strains carry the *cdc15-2* thermosensitive allele). Then, the culture was split
in two, one subculture remained untreated (mock), and DSBs were generated in the second
one. When a G2/M arrest was required, 15 µg·mL^{-1} nocodazole (Nz; Sigma-Aldrich, M1404)
was added to the asynchronous culture, which was then incubated at 25°C for 3 h (with a 7.5
µg·mL^{-1} Nz shot at 2 h). To conditionally degrade the Smc3-aid* variant, 8 mM of the auxin
3-indol-acetic acid (IAA; Sigma-Aldrich, I2886) was added to the arrested cells 1 h prior to
generating DSBs.

DSB generation was accomplished by incubating the culture for 1 to 3 hours with
either 10 µg·mL^{-1} phleomycin (random DSBs) or 2 µM of β-estradiol (HO-driven DSBs). For
the *HOcs* DSB, the corresponding strain harbours an integrative system based on a β-
estradiol inducible promoter (Ottoz et al., 2014). These strains were constructed by
transforming with the pRG464 plasmid.

Western blot for protein levels and phosphorylation

Western blotting was performed as reported before (Ayra-Plasencia and Machín,
2019). The trichloroacetic acid TCA method was used for protein extraction. Total protein
was quantified in a Qubit 4 Fluorometer (Thermo Fisher Scientific, Q33227). Proteins were
resolved in 7.5% SDS-PAGE gels and transferred to PVFD membranes (Pall Corporation,
PVM020C099). The membrane was stained with Ponceau S solution (PanReac AppliChem,
A2935) as a loading reference. The following primary antibodies were used for
immunoblotting: mouse monoclonal α-HA (1:1,000; Sigma-Aldrich, H9658), mouse
monoclonal α-myc (1: 5,000; Sigma-Aldrich, M4439), mouse monoclonal α-Pgk1 (1:5,000;
Thermo Fisher Scientific, 22C5D8), mouse monoclonal α-miniaid (1:500; MBL, M214-3),
mouse monoclonal α-Flag® (1:5000; Sigma-Aldrich, F3165), mouse monoclonal α-Rad53
(1:1000; Abcam, ab166859), and mouse monoclonal α-acSmc3 (1:5000; a gift from
Katsuhiko Shirahige). The secondary antibody was a horseradish peroxidase polyclonal goat
anti-mouse (from 1:5,000 to 1:10,000, depending on the primary antibody; Promega,

W4021). Proteins were detected with the ECL chemiluminescence reagent (GE Healthcare, RPN2232), and visualized in a Vilber-Lourmat Fusion Solo S chamber.

Co-immunoprecipitation (co-IP) for the reassembly of the cohesin complex.

For coI-P assays, approximately 100 OD_{600} units from the corresponding conditions were collected. Cell disruption was performed by resuspending the frozen pellets in 200 µL of Buffer A (50 mM HEPES-Na, 150 mM KCl, 1.5 mM $MgCl_2$, 0.5 mM DTT, 0.5% Triton X-100, 1× EDTA-free protease inhibitor cocktail) together with glass beads. Samples were vortexed vigorously for 5 min, and additional lysis cycles were performed as needed, allowing 1 min incubation on ice between rounds. Lysates were transferred to a clean tube. 400 µL of Buffer A was used to wash the beads and pooled with the lysates. Samples were then centrifuged (12,000 rpm, 5 min, 4°C), and the clarified lysates were transferred to new tubes. An aliquot of 70 µL was saved as whole cell extract (WCE) for western blot analysis. The remaining lysate was incubated with pre-equilibrated Pierce™ Anti-HA Magnetic Beads (Thermo Fisher Scientific; 88836) for 2 h at 4°C on a rotating wheel. After incubation, beads were placed on a magnetic rack, and washed five times with 200 µL of IPP150 buffer (10 mM Tris-Cl pH 7.5, 150 mM NaCl, 0.5% Triton X-100).

For elution, beads were resuspended in 25 µL of 1× SR buffer (8% SDS, 0.5 M Tris-Cl pH 6.8) and incubated at 37 °C for 4 min. The supernatant was collected using a magnetic rack and transferred to a new tube containing 8.3 µL of 4× SS buffer (20% sucrose, 0.05% bromophenol blue, 0.1% sodium azide) supplemented with β-mercaptoethanol. Samples were boiled for 2 min at 95°C, centrifuged (max speed, 30 sec) and analyzed by SDS-PAGE and western blotting.

Chromatin fractionation for the chromatin-bound cohesin.

Biochemical fractionation of cells was performed as described before with little modifications (Cuevas-Bermúdez et al., 2020). Briefly, cells were collected and resuspended in 400 µl of buffer 1 (20 mM HEPES pH 8, 60 mM KCl, 15 mM NaCl, 10 mM $MgCl_2$, 1 mM $CaCl_2$, 0.8% Triton X-100, 1.25 M sucrose, 0.5 mM spermine). Breakage of cell walls was done by adding ~ 200 mg of glass beads and vortexing for 4 min. After centrifuging at 500 g for 5 min., supernatant was transfer to a new tube and 40 µl were kept as Input. Samples were centrifuge at 18000 g for 20 min, the supernatant was kept as S1, whereas the pellet was resuspended in 200 µl of Buffer 1 (50 µl were saved as P1). Samples were centrifuged again 18000 g for 20 min, the supernatant was kept as S2, and the pellet was resuspended in 200 µl of buffer 2 (20 mM HEPES pH 7.6, 45 mM NaCl, 7.5 mM $MgCl_2$, 20 mM EDTA pH 8, 10% glycerol, 1% NP-40, 2 M urea, 0.5 M sucrose, protease inhibitors) and 50 µl were saved as P2. Finally, samples were centrifuged one last time at the same conditions, keeping the supernatant as S3, and the pellet was resuspended in 50 µl of Laemli buffer, constituting the chromatin-enriched fraction (P3).

Chromatin immunoprecipitation (ChIP-qPCR) for the cohesin loading onto the HO DSB.

For ChIP–qPCR, a total of 100 OD_{600} units of the treated cultures were collected and crosslinked with 1.4% formaldehyde for 15 min. Crosslinking was quenched with glycine (final concentration 125 mM) for 7 min. Cells were harvested by centrifugation (4000 rpm, 1 min), washed with PBS, transferred to screw-cap tubes, and frozen at –80°C.

Frozen pellets were resuspended in 300 µL of IP buffer (150 mM NaCl, 50 mM Tris-
HCl pH 7.5, 5 mM EDTA, 1% Triton X-100, 0.05% NP-40) supplemented with 1 mM PMSF
and protease inhibitor cocktail (EDTA-free, Roche). Disruption was performed by bead
beating with 500 µL of glass beads using a Bertin Precellys homogenizer for nine 20 s cycles
at hard setting. Lysates were transferred to new tubes and supplemented with an additional
100 µL of IP buffer containing PMSF and protease inhibitors. Samples were clarified by
centrifugation (15,000 rpm, 10 min, 4 °C). Pellets were resuspended in 1 mL IP buffer with
PMSF and protease inhibitors and sonicated in a Diagenode Bioruptor Plus (30 cycles: 30 s
ON 30 s OFF, high power, 4°C). After sonication, samples were centrifuged (15,000 rpm,
10 min), and the supernatant was collected. A 200 µL aliquot was reserved as input. Input
DNA was precipitated with 0.3 M sodium acetate and 2.5 volumes of cold ethanol,
centrifuged (15,000 rpm, 30 min), washed with 70% ethanol, and air-dried.

For immunoprecipitation, 400 µL of the sonicated chromatin was incubated with 40 µg
of anti-HA antibody (Roche, 11666606001) in the Bioruptor at low power for 30 min (30 s
ON/30 s OFF cycles). Antibody-chromatin complexes were pelleted (13,000 rpm, 5 min), and
the supernatant was incubated with 60 µL of Dynabeads Protein G (Invitrogen), pre-
equilibrated in IP buffer, then incubated for 2 h at 4°C on a rotating wheel, and finally washed
five times with IP buffer using a magnetic rack.

Both input and IP samples were resuspended in de-crosslinking buffer (1× TE, 1%
SDS, 10 µg/mL RNase A, 1 mg/mL proteinase K) and incubated overnight at 65°C and
purified using High Pure PCR product purification kit (Roche) for the qPCR.

434
435   **qPCR for DSB resection.**
436
In the resection experiments, qPCR was performed on genomic DNA (gDNA)
extracted by the glass beads/phenol method (Hoffman and Winston, 1987). Experimental
details on the resection assay can be found in (Gnügge et al., 2018). Briefly, each gDNA
sample was divided into two and one aliquot was digested with StyI-HF (NEB, R3500S).
Then, qPCR reactions were mounted using the PowerUp™ SYBR™ Green Master Mix
(Thermo Scientific, A25741) and run in a BioRad CFX384 Real-Time PCR instrument (10 µl
final volume in 384-well block plates).
Resection was calculated as a normalized fraction ($f_{resected}$) to that of the HOcs
effectively cut by HO (f). These two parameters were calculated by their corresponding
formulas (Gnügge et al., 2018).
447
Southern blot for the MAT switching assay.

In this case, gDNA was extracted by digesting with 50U lyticase (Sigma-Aldrich,
L4025) a cell pellet previously resuspended in 200 µl of digestion buffer (1 % SDS, 100 mM
NaCl, 50 mM Tris-HCl, and 10 mM EDTA). Then, the gDNA was phase separated by
phenol:chloroform (PanReac AppliChem, A0944), precipitated with ethanol, resuspended in
TE 1X with 10 µg·mL^{-1} RNase A, precipitated again, and resuspended in TE 1X. Then, the
gDNA was digested with StyI, the restriction fragments separated on a 1.2% low EEOO LS
Agarose gel, and finally Southern blotted. Southern blot was carried out by a saline
downwards transference onto a positively charged nylon membrane (Hybond-N+,
Amersham-GE; RPN303B) (García-Luis and Machín, 2014). DNA probes against *ACT1* and
MAT loci were made using Fluorescein-12-dUTP Solution (ThermoFisher; R0101) and the

Expand™ High Fidelity PCR System (Roche; 11732641001). Hybridization with fluorescein-
labeled probes was performed overnight at 68°C. The following day, the membrane was
incubated with an anti-fluorescein antibody coupled to alkaline phosphatase (Roche;
11426338910), and the chemiluminescent signal detected using CDP-star (Amersham;
RPN3682). Blots were visualized in a Vilber-Lourmat Fusion Solo S chamber.

For band quantification, each individual band was normalized to the *ACT1* signal in
the lane. Then, a second normalization to the *MAT*a band at the arrest was performed. To
determine the yield of gene conversion to *MAT*α specifically, the *MAT*α band after recovery
from the DSB (2-3 h after β-estradiol removal) was normalized to the amount of the cut
HOcs band immediately before recovery (1h after β-estradiol addition).

Data representation and statistics.

Three types of graphs were used to represent the data: bar charts, marker line
graphs and box plots. In box plots, the center line represents the medians, box limits
represent the 25th and 75th percentile, the whiskers extend to the 5th and 95th percentiles,
and the dots represent outliers. Error bars in all bar and line charts represent the standard
error of the mean (s.e.m.) of three independent biological replicates. Individual values are
also represented as dots in the bar charts. Graphpad Prism 9 was used for generating the
charts and for statistical analysis. Differences between experimental data points were
generally estimated using the Mann-Whitney U test for box plots and one-way ANOVA with
Tukey post hoc for bar charts.

Acknowledgements

We are grateful to Katsuhiko Shirahige for the anti-acSmc3 antibody. We would like
to thank other members from Machín's and Symington's labs for enriching and fruitful
discussion. This work was supported by the Spanish Ministry of Science, Innovation and
Universities (MICIU/AEI/10.13039/501100011033; research grants BFU2017-83954-R and
PID2021-123716OB-I00 to F.M.; co-funded by the EU-ERDF "A way of making Europe") and
the National Institute of Health (research grant NIH R35 GM126997 to L.S.). The Agencia
Canaria de Investigación, Innovación y Sociedad de la Información (ACIISI) supported S.
M.-S. through a predoctoral fellowship (TESIS2020010028; co-funded by the ESF+), and the
Servicio Público de Empleo Estatal (SEPE) supported E.H-C. through an INVESTIGO
program contract (co-funded by the ESF+).

Author contributions

J Ayra-Plasencia: methodology, investigation, formal analysis, visualization, curation,
writing–original draft, writing–review and editing.

S Medina-Suárez: methodology, investigation, formal analysis, visualization, writing–review
and editing.

E Hernández-Carralero: methodology, investigation, formal analysis, visualization writing–
review and editing.

J García-Luis: methodology, investigation, formal analysis, visualization writing–review and
editing.

L Symington: supervision, project administration, funding acquisition, writing–review and
editing.

F Machín: conceptualization, visualization, resources, supervision, project administration,
funding acquisition, curation, writing–original draft, writing–review and editing.

References

Ayra-Plasencia J, Machín F. 2019. DNA double-strand breaks in telophase lead to
coalescence between segregated sister chromatid loci. *Nat Commun* **10**:2862.
doi:10.1038/s41467-019-10742-8

Beckouët F, Hu B, Roig MB, Sutani T, Komata M, Uluocak P, Katis VL, Shirahige K,
Nasmyth K. 2010. An Smc3 Acetylation Cycle Is Essential for Establishment of Sister
Chromatid Cohesion. *Mol Cell* **39**:689–699. doi:10.1016/j.molcel.2010.08.008

Branzei D, Foiani M. 2006. The Rad53 signal transduction pathway: Replication fork
stabilization, DNA repair, and adaptation. *Exp Cell Res* **312**:2654–9.
doi:10.1016/j.yexcr.2006.06.012

Çamdere G, Guacci V, Stricklin J, Koshland D. 2015. The ATPases of cohesin interface with
regulators to modulate cohesin-mediated DNA tethering. *Elife* **4**.
doi:10.7554/eLife.11315

Chan K-L, Roig MB, Hu B, Beckouët F, Metson J, Nasmyth K. 2012. Cohesin's DNA Exit
Gate Is Distinct from Its Entrance Gate and Is Regulated by Acetylation. *Cell* **150**:961–
974. doi:10.1016/j.cell.2012.07.028

Ciosk R, Shirayama M, Shevchenko Anna, Tanaka T, Toth A, Shevchenko Andrej, Nasmyth
K. 2000. Cohesin's binding to chromosomes depends on a separate complex consisting
of Scc2 and Scc4 proteins. *Mol Cell* **5**:243–254. doi:10.1016/S1097-2765(00)80420-7

Cohen-Fix O, Peters JM, Kirschner MW, Koshland D. 1996. Anaphase initiation in
Saccharomyces cerevisiae is controlled by the APC-dependent degradation of the
anaphase inhibitor Pds1p. *Genes Dev* **10**:3081–93. doi:10.1101/GAD.10.24.3081

Cuevas-Bermúdez A, Garrido-Godino AI, Gutiérrez-Santiago F, Navarro F. 2020. A Yeast
Chromatin-enriched Fractions Purification Approach, yChEFs, from Saccharomyces
cerevisiae. *Bio-protocol* **10**:e3471. doi:10.21769/BioProtoc.3471

Díaz-Martínez L a, Giménez-Abián JF, Clarke DJ. 2008. Chromosome cohesion - rings,
knots, orcs and fellowship. *J Cell Sci* **121**:2107–14. doi:10.1242/jcs.029132

Djeghloul D, Patel B, Kramer H, Dimond A, Whilding C, Brown K, Kohler A-C, Feytout A,
Veland N, Elliott J, Bharat TAM, Tarafder AK, Löwe J, Ng BL, Guo Y, Guy J, Huseyin
MK, Klose RJ, Merkenschlager M, Fisher AG. 2020. Identifying proteins bound to native
mitotic ESC chromosomes reveals chromatin repressors are important for compaction.
Nat Commun **11**:4118. doi:10.1038/s41467-020-17823-z

Garcia-Luis J, Bordelet H, Thierry A, Koszul R, Aragon L. 2022. Depletion or cleavage of

cohesin during anaphase differentially affects chromatin structure and segregation. *Elife*
11:1–22. doi:10.7554/eLife.80147

García-Luis J, Machín F. 2014. Mus81-Mms4 and Yen1 resolve a novel anaphase bridge
formed by noncanonical Holliday junctions. *Nat Commun* **5**:5652.
doi:10.1038/ncomms6652

Gnügge R, Oh J, Symington LS. 2018. Processing of DNA Double-Strand Breaks in Yeast.
Methods Enzymol **600**:1–24. doi:10.1016/bs.mie.2017.11.007

Gnügge R, Symington LS. 2020. Efficient DNA double-strand break formation at single or
multiple defined sites in the Saccharomyces cerevisiae genome. *Nucleic Acids Res*
48:e115–e115. doi:10.1093/nar/gkaa833

Heidinger-Pauli JM, Ünal E, Guacci V, Koshland D. 2008. The Kleisin Subunit of Cohesin
Dictates Damage-Induced Cohesion. *Mol Cell* **31**:47–56.
doi:10.1016/j.molcel.2008.06.005

Hirano T. 2000. Chromosome cohesion, condensation, and separation. *Annu Rev Biochem*
69:115–44. doi:10.1146/annurev.biochem.69.1.115

Hoffman CS, Winston F. 1987. A ten-minute DNA preparation from yeast efficiently releases
autonomous plasmids for transformaion of Escherichia coli. *Gene* **57**:267–272.
doi:10.1016/0378-1119(87)90131-4

Hou W, Li Y, Zhang J, Xia Y, Wang X, Chen H, Lou H. 2022. Cohesin in DNA damage
response and double-strand break repair. *Crit Rev Biochem Mol Biol* **57**:333–350.
doi:10.1080/10409238.2022.2027336

Huber RG, Kulemzina I, Ang K, Chavda AP, Suranthran S, Teh J-T, Kenanov D, Liu G,
Rancati G, Szmyd R, Kaldis P, Bond PJ, Ivanov D. 2016. Impairing Cohesin Smc1/3
Head Engagement Compensates for the Lack of Eco1 Function. *Structure* **24**:1991–
1999. doi:10.1016/j.str.2016.09.001

Ira G, Pellicioli A, Balijja A, Wang X, Fiorani S, Carotenuto W, Liberi G, Bressan D, Wan L,
Hollingsworth NM, Haber JE, Foiani M. 2004. DNA end resection, homologous
recombination and DNA damage checkpoint activation require CDK1. *Nature*
431:1011–7. doi:10.1038/nature02964

Kim B-J, Li Yehua, Zhang J, Xi Y, Li Yumei, Yang T, Jung SY, Pan X, Chen R, Li W, Wang
Y, Qin J. 2010. Genome-wide Reinforcement of Cohesin Binding at Pre-existing
Cohesin Sites in Response to Ionizing Radiation in Human Cells. *J Biol Chem*
285:22784–22792. doi:10.1074/jbc.M110.134577

Knop M, Siegers K, Pereira G, Zachariae W, Winsor B, Nasmyth K, Schiebel E. 1999.
Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved
practical routines. *Yeast* **15**:963–72. doi:10.1002/(SICI)1097-
0061(199907)15:10B<963::AID-YEA399>3.0.CO;2-W

Koshland DE, Guacci V. 2000. Sister chromatid cohesion: the beginning of a long and
beautiful relationship. *Curr Opin Cell Biol* **12**:297–301. doi:10.1016/s0955-
0674(00)00092-2

Laloraya S, Guacci V, Koshland D. 2000. Chromosomal addresses of the cohesin
component Mcd1p. *J Cell Biol* **151**:1047–56. doi:10.1083/JCB.151.5.1047

Langerak P, Russell P. 2011. Regulatory networks integrating cell cycle control with DNA
damage checkpoints and double-strand break repair. *Philos Trans R Soc Lond B Biol*
Sci **366**:3562–71. doi:10.1098/rstb.2011.0070

Lazar-Stefanita L, Scolari VF, Mercy G, Muller H, Guérin TM, Thierry A, Mozziconacci J,
Koszul R. 2017. Cohesins and condensins orchestrate the 4D dynamics of yeast
chromosomes during the cell cycle. *EMBO J* **36**:2684–2697.
doi:10.15252/embj.201797342

Lazzaro F, Sapountzi V, Granata M, Pellicioli A, Vaze M, Haber JE, Plevani P, Lydall D,
Muzi-Falconi M. 2008. Histone methyltransferase Dot1 and Rad9 inhibit single-stranded
DNA accumulation at DSBs and uncapped telomeres. *EMBO J* **27**:1502–1512.
doi:10.1038/emboj.2008.81

Leonard J, Sen N, Torres R, Sutani T, Jarmuz A, Shirahige K, Aragón L. 2015. Condensin
Relocalization from Centromeres to Chromosome Arms Promotes Top2 Recruitment
during Anaphase. *Cell Rep* **13**:2336–2344. doi:10.1016/j.celrep.2015.11.041

Lyons NA, Morgan DO. 2011. Cdk1-Dependent Destruction of Eco1 Prevents Cohesion
Establishment after S Phase. *Mol Cell* **42**:378–389. doi:10.1016/j.molcel.2011.03.023

Machín F, Ayra-Plasencia J. 2020. Are Anaphase Events Really Irreversible? The Endmost
Stages of Cell Division and the Paradox of the DNA Double-Strand Break Repair.
BioEssays **42**:2000021. doi:10.1002/bies.202000021

Mathiasen DP, Lisby M. 2014. Cell cycle regulation of homologous recombination in
Saccharomyces cerevisiae. *FEMS Microbiol Rev* **38**:172–184. doi:10.1111/1574-
6976.12066

Medina-Suárez S, Ayra-Plasencia J, Pérez-Martínez L, Butter F, Machín F. 2024. Msc1 is a
nuclear envelope protein that reinforces DNA repair in late mitosis. *iScience* **27**:110250.
doi:10.1016/j.isci.2024.110250

Michaelis C, Ciosk R, Nasmyth K. 1997. Cohesins: Chromosomal proteins that prevent
premature separation of sister chromatids. *Cell* **91**:35–45. doi:10.1016/S0092-
8674(01)80007-6

Minamino M, Ishibashi M, Nakato R, Akiyama K, Tanaka H, Kato Y, Negishi L, Hirota T,
Sutani T, Bando M, Shirahige K. 2015. Esco1 Acetylates Cohesin via a Mechanism
Different from That of Esco2. *Curr Biol* **25**:1694–706. doi:10.1016/j.cub.2015.05.017

Morawska M, Ulrich HD. 2013. An expanded tool kit for the auxin-inducible degron system in
budding yeast. *Yeast* **30**:341–51. doi:10.1002/yea.2967

Moreau S, Morgan EA, Symington LS. 2001. Overlapping functions of the Saccharomyces
cerevisiae Mre11, Exo1 and Rad27 nucleases in DNA metabolism. *Genetics* **159**:1423–
33. doi:10.1093/genetics/159.4.1423

Murayama Y, Uhlmann F. 2015. DNA Entry into and Exit out of the Cohesin Ring by an
Interlocking Gate Mechanism. *Cell* **163**:1628–1640. doi:10.1016/j.cell.2015.11.030

Nasmyth K, Haering CH. 2009. Cohesin: Its Roles and Mechanisms. *Annu Rev Genet*
43:525–558. doi:10.1146/annurev-genet-102108-134233

Ottoz DSM, Rudolf F, Stelling J. 2014. Inducible, tightly regulated and growth condition-

independent transcription factor in Saccharomyces cerevisiae. *Nucleic Acids Res* **42**.
doi:10.1093/nar/gku616

Pellicioli A, Lee SE, Lucca C, Foiani M, Haber JE. 2001. Regulation of Saccharomyces
Rad53 checkpoint kinase during adaptation from DNA damage-induced G2/M arrest.
Mol Cell **7**:293–300. doi:10.1016/s1097-2765(01)00177-0

Peng H, Zhang S, Chen X. 2021. Monitoring 5'-End Resection at Site-Specific Double-
Strand Breaks by Southern Blot Analysis. *Methods Mol Biol* **2196**:245–255.
doi:10.1007/978-1-0716-0868-5_20

Phipps J, Dubrana K. 2022. DNA Repair in Space and Time: Safeguarding the Genome with
the Cohesin Complex. *Genes (Basel)* **13**. doi:10.3390/genes13020198

Piazza A, Bordelet H, Dumont A, Thierry A, Savocco J, Girard F, Koszul R. 2021. Cohesin
regulates homology search during recombinational DNA repair. *Nat Cell Biol* **23**:1176–
1186. doi:10.1038/s41556-021-00783-x

Rao H, Uhlmann F, Nasmyth K, Varshavsky A. 2001. Degradation of a cohesin subunit by
the N-end rule pathway is essential for chromosome stability. *Nature* **410**:955–9.
doi:10.1038/35073627

Rao SSP, Huang S-C, Glenn St Hilaire B, Engreitz JM, Perez EM, Kieffer-Kwon K-R,
Sanborn AL, Johnstone SE, Bascom GD, Bochkov ID, Huang X, Shamim MS, Shin J,
Turner D, Ye Z, Omer AD, Robinson JT, Schlick T, Bernstein BE, Casellas R, Lander
ES, Aiden EL. 2017. Cohesin Loss Eliminates All Loop Domains. *Cell* **171**:305-320.e24.
doi:10.1016/j.cell.2017.09.026

Renshaw MJ, Ward JJ, Kanemaki M, Natsume K, Nédélec FJ, Tanaka TU. 2010.
Condensins promote chromosome recoiling during early anaphase to complete sister
chromatid separation. *Dev Cell* **19**:232–44. doi:10.1016/j.devcel.2010.07.013

Rolef Ben-Shahar T, Heeger S, Lehane C, East P, Flynn H, Skehel M, Uhlmann F. 2008.
Eco1-dependent cohesin acetylation during establishment of sister chromatid cohesion.
Science **321**:563–6. doi:10.1126/science.1157774

Schalbetter SA, Goloborodko A, Fudenberg G, Belton J-M, Miles C, Yu M, Dekker J, Mirny
L, Baxter J. 2017. SMC complexes differentially compact mitotic chromosomes
according to genomic context. *Nat Cell Biol* **19**:1071–1080. doi:10.1038/ncb3594

Sjögren C, Nasmyth K. 2001. Sister chromatid cohesion is required for postreplicative
double-strand break repair in Saccharomyces cerevisiae. *Curr Biol* **11**:991–5.
doi:10.1016/s0960-9822(01)00271-8

Ström L, Karlsson C, Lindroos HB, Wedahl S, Katou Y, Shirahige K, Sjögren C. 2007.
Postreplicative formation of cohesion is required for repair and induced by a single DNA
break. *Science* **317**:242–5. doi:10.1126/science.1140649

Ström L, Lindroos HB, Shirahige K, Sjögren C. 2004. Postreplicative Recruitment of Cohesin
to Double-Strand Breaks Is Required for DNA Repair. *Mol Cell* **16**:1003–1015.
doi:10.1016/j.molcel.2004.11.026

Strunnikov A V., Larionov VL, Koshland D. 1993. SMC1: an essential yeast gene encoding a
putative head-rod-tail protein is required for nuclear division and defines a new
ubiquitous protein family. *J Cell Biol* **123**:1635–48. doi:10.1083/jcb.123.6.1635

Symington LS. 2016. Mechanism and regulation of DNA end resection in eukaryotes. *Crit*
Rev Biochem Mol Biol **51**:195–212. doi:10.3109/10409238.2016.1172552

Symington LS, Rothstein R, Lisby M. 2014. Mechanisms and Regulation of Mitotic
Recombination in Saccharomyces cerevisiae. *Genetics* **198**:795–835.
doi:10.1534/genetics.114.166140

Tanaka T, Cosma MP, Wirth K, Nasmyth K. 1999. Identification of cohesin association sites
at centromeres and along chromosome arms. *Cell* **98**:847–858. doi:10.1016/S0092-
8674(00)81518-4

Uhlmann F, Lottspeich F, Nasmyth K. 1999. Sister-chromatid separation at anaphase onset
is promoted by cleavage of the cohesin subunit Scc1. *Nature* **400**:37–42.
doi:10.1038/21831

Uhlmann F, Nasmyth K. 1998. Cohesion between sister chromatids must be established
during DNA replication. *Curr Biol* **8**:1095–101. doi:10.1016/s0960-9822(98)70463-4

Ünal E, Arbel-Eden A, Sattler U, Shroff R, Lichten M, Haber JE, Koshland D. 2004. DNA
Damage Response Pathway Uses Histone Modification to Assemble a Double-Strand
Break-Specific Cohesin Domain. *Mol Cell* **16**:991–1002.
doi:10.1016/j.molcel.2004.11.027

Unal E, Heidinger-Pauli JM, Kim W, Guacci V, Onn I, Gygi SP, Koshland DE. 2008. A
molecular determinant for the establishment of sister chromatid cohesion. *Science*
321:566–9. doi:10.1126/science.1157880

Unal E, Heidinger-Pauli JM, Koshland D. 2007. DNA double-strand breaks trigger genome-
wide sister-chromatid cohesion through Eco1 (Ctf7). *Science* **317**:245–8.
doi:10.1126/science.1140637

Yam CQX, Chia DB, Shi I, Lim HH, Surana U. 2020. Dun1, a Chk2-related kinase, is the
central regulator of securin-separase dynamics during DNA damage signaling. *Nucleic*
Acids Res **48**:6092–6107. doi:10.1093/nar/gkaa355

Yamaguchi M, Haber JE. 2021. Monitoring Gene Conversion in Budding Yeast by Southern
Blot Analysis. *Methods Mol Biol* **2153**:221–238. doi:10.1007/978-1-0716-0644-5_16

Yamamoto A, Guacci V, Koshland D. 1996. Pds1p is required for faithful execution of
anaphase in the yeast, Saccharomyces cerevisiae. *J Cell Biol* **133**:85–97.
doi:10.1083/jcb.133.1.85

Zierhut C, Diffley JFX. 2008. Break dosage, cell cycle stage and DNA replication influence
DNA double strand break response. *EMBO J* **27**:1875–85. doi:10.1038/emboj.2008.111

FIGURE LEGENDS

Figure 1. Scc1 returns after DSBs in late mitosis. (**a**) Schematic of the experimental
procedure. Cells in logarithmic growth phase were first arrested in telophase (Tel) by
incubating them at 34⁰C for 3 h (strains bear the *cdc15-2* allele). Then, the culture was
divided into three subcultures. One served as a mock control, and the other two were treated
to generate DSBs, one with β-estradiol (to express the HO endonuclease) and the other with

phleomycin. The strain is unable to repair the HO-DSB break by HR-driven gene conversion (Δ*hmr* Δ*hml*). (**b**) Western blot against Scc1-3myc. This experiment compares Scc1-3myc levels in an asynchronous, G2- and telophase-blocked cultures. The leftmost lane (No tag) is a control strain for Scc1 without the 3myc epitope tag. Pgk1 protein served as a housekeeping. Ponceau S staining is also shown as a loading control. Asyn.: Asynchronous. Tel: Telophase. +Phle: 10 mg·mL^{-1} phleomycin. +HO: 2 µM β-estradiol. *: Unspecific band detected by the α-myc antibody just over the Scc1-3myc signal. (**c**) Like in (**b**) but against Smc1-6HA. (**d**) Like in (**c**) but against Smc3-6HA. (**e**) Quantification of Scc1-3myc and Smc1-6HA levels in all conditions (mean ± sem, n=3). Statistical comparisons are shown for selected pairs (**, p<0.01; one-way ANOVA, Tukey's post hoc).

Figure 2. Scc1 forms a reconstituted chromatin-bound cohesin complex after the HO DSB in late mitosis. Cells were treated as in Figure 1a. (**a**) Co-IP of the core cohesin complex. Smc1-6HA was used as the bait protein; the leftmost lane includes a control with an untagged Smc1. (**b**) Chromatin fractionation of the strain expressing Smc1-6HA and Scc1-3myc. Relative protein levels (to the mock sample) are indicated under the blots. Pgk1 and histone H3 (HH3) were included as reporters of the cytosolic and chromatin-bound fractions, respectively. The asterisk (*) indicates an unspecific band in the whole cell extract (WCE), that is absent from the chromatin-bound fraction. (**c**) Cohesin does not bind the HO DSB in telophase. Chromatin immunoprecipitation (ChIP) of Smc3-6HA with (+HO) and without (-HO) HO induction (mean ± sem, n=3).

Figure 3. Homologous recombination repairs the HO-DSB in telophase. (**a,b**) Resection in late mitosis is almost as efficient as in G2/M. Cells were arrested in G2/M or telophase (Tel), and then the HO DSB was generated by adding β-estradiol. Samples were taken at the indicated times to monitor the kinetics of resection. This strain is unable to repair the HO-DSB break by HR-driven gene conversion (Δ*hmr* Δ*hml*). (**a**) Charts depicting the resection kinetics for two different amplicons located at 726 bp and 5.7 kb downstream the *HOcs* break (mean ± sem, n=3; $f_{resected}$ is the proportion of resected DNA). (**b**) Representative western blots against Rad53 to follow the sensing of DNA damage detection in G2/M versus late mitosis. (**c-f**) Yeast cells use HR to repair the HO DSB in late mitosis. Cells were first blocked in telophase (Tel). Then, the HO DSB was generated by adding β-estradiol. After 1 hour, the β-estradiol was washed away, and samples were taken to monitor the repair for 3 hours. The strain can repair the HO-DSB break by HR-driven gene conversion (Δ*hmr* HML). (**c**) Schematic of the fragments obtained after a StyI digestion for both *MAT*a and *MAT*α sequences. The probe to detect the fragments by Southern blot is in blue. When the *MAT*a locus is intact, the digestion gives rise to a 0.9 kb fragment. The HO cutting site (*HOcs*) is located within the StyI-digested *MAT*a locus. Thus, the HO-driven DSB shortens the fragment to 0.7 kb. HR leads to a gene conversion to *MAT*α, which results in the loss of a *StyI* restriction site and a new fragment of 1.8 kb. (**d**) Representative Western blot analyses for HO induction and subsequent degradation (tagged with Flag epitope) and the DSB sensing through Rad53 hyperphosphorylation. A Pgk1 western blot served as a housekeeping, and Ponceau S staining of the membrane as a loading control for all lanes. The leftmost lane in the Ponceau S corresponds to the protein weight marker. (**e**) Representative Southern blot for the *MAT* switching assay in late mitosis. Alongside the *MAT* probe, a second probe against the *ACT1* gene (1.1 kb fragment) was included for normalization. The *MAT* probe also recognizes an allele-independent *MAT* distal fragment (2.2 kb). (**f**) Quantification of relative band intensities in the *MAT* switching Southern blots

(mean ± sem, n=3). Individual values were normalized to the *ACT1* signals. Then, every lane
was normalized to *MAT*a at the arrest. Tel: Telophase. +βE: β-Estradiol addition.

Figure 4. Cohesin removal does not alter HR at the *MAT* locus in late mitosis. (**a**) Serial
dilution spot assay for the *smc3:aid** strains. The fitness effect of Smc3-aid* degradation was
tested in the *smc3:aid* OsTIR1* strain in the presence of 8 mM IAA. The same strain without
the ubiquitin-ligase *OsTIR1* was also used as the control. (**b-d**) Cells of four strains with all
possible combination of the *smc3:aid** and *OsTIR1* pairs were treated as in Figure 1a but
including a 1 h IAA step between the *cdc15-2* arrest and the induction of HO. Then, HO was
induced for 1h with β-estradiol (1h+βE sample), after which it was washed off and the cells
were allowed to recover from the DSB for 2h (3h-βE sample). IAA and the Tel arrest were
maintained throughout the experiment. (**b**) Representative Western blot of the four strains at
the time of the arrest (Tel) and 1h after IAA addition (+IAA). Note the confirmation of the
different genotypic combinations, as well as the Smc3-aid* decline after IAA addition. (**c**)
Representative Southern blot for the *MAT* switching assay of the four strains in the presence
of 8 mM IAA. Note that HO cutting is less efficient than without IAA (see Figures 3e and
S6b,c) but gene conversion is still possible after degrading Smc3-aid* in late mitosis. (**d**)
Quantification of relative *MAT* switching (mean ± sem, n=3). Gene conversion to *MAT*α in
the 3h-βE sample was normalized to the cut *HOcs* in the 1h+βE sample. Then, gene
conversion yield was normalized to the wild type (no tagged Smc3, no OsTir1) strain. Tel:
Telophase. +βE: β-Estradiol addition. -βE: β-Estradiol removal.+IAA: Indole-acetic acid. (**e**)
Model of how DSBs reconstitute cohesin in late mitosis. The cohesin complex (Smc1-Smc3-
Scc1) entraps sister chromatids before anaphase. (1) At anaphase onset, Esp1 (separase)
is activated to cleave Scc1 and release sister chromatids for segregation. (2) If DSBs occur
before cytokinesis, Scc1 returns and reconstitutes the cohesin complex. At least a fraction of
the complex binds to chromatin. The reappearance of Scc1 likely occurs through the
inhibition of Esp1 and/or post-translational protection of Scc1 by the DNA damage
checkpoint.

a

2h after Tel arrest

	Asyn	Asyn	mock	+Phle	+HO
Smc1-6HA					
Scc1-3myc					
acSmc3					

SMC1　　　　*SMC1:6HA*

b

Chromatin bound

	Mock	+Phle	+HO
Scc1-3myc			
Rel. Scc1:	1.00	0.74	7.43
Smc1-6HA			
Rel. Smc1:	1.00	1.07	1.95
Pgk1			
HH3			

WCE

Scc1-3myc				*
Smc1-6HA			
Pgk1			

c

Cohesin reconstitution and homologous recombination repair of DNA double strand breaks in late mitosis.

Jessel Ayra-Plasencia[1,2,#,†,*], Sara Medina-Suárez[1,2,†], Esperanza Hernández-Carralero[1], Jonay García-Luis[2], Lorraine S Symington[3], and Félix Machín[1,2,4,*]

[1] Hospital Universitario Nuestra Señora de Candelaria, Instituto de Investigación Sanitaria de Canarias (IISC), 38010 Santa Cruz De Tenerife, Spain.
[2] Instituto of Tecnologías Biomédicas, Universidad de La Laguna, 38200 San Cristóbal de La Laguna, Spain.
[3] Department of Microbiology and Immunology, Columbia University Medical Center, New York, United States.
[4] Facultad de Ciencias de la Salud, Universidad Fernando Pessoa Canarias, Las Palmas de Gran Canaria, Spain.

[#] Present address: Unidad de Investigación, Hospital Universitario de Canarias, 38320 San Cristóbal de La Laguna, Spain

[†] These authors contributed equally.

[*] Co-corresponding authors: jayra@fciisc.es; fmachin@fciisc.es

Supplemental Figures

Figure S1. Co-IP of the core cohesin complex. This is a biological replicate of the experiment shown in Figure 2a.

Figure S2. Assessment of chromatin-bound cohesin after DSBs in late mitosis. Related to Figure 2b. (**a**) Optimization of the chromatin fractionation using the chromatin-bound pool of Smc3 (acetylated Smc3; acSmc3) and the histone H3 (HH3). The best results were obtained after three successive rounds of fractionation (pellet #3, P3; see Material and Methods). S, supernatant; P, pellet. (**b**) The strain expressing Smc3-6HA was subjected to the same experiment and chromatin fractionation performed in Figure 2b for Smc1-6HA and Scc1-3myc. Note that there are more chromatin-bound acSmc3 after DSBs, especially after the HO DSB.

Figure S3. Chromatin immunoprecipitation (ChIP) of Smc3-6HA in G2/M. Related to Figure 2c. The ChiP was performed upon cells arrested in G2/M with nocodazole and further subdivided into two subcultures, one in which the HO DSB was generated (+HO) and one without the DSB (-HO). The strain is unable to repair the break by gene conversion (Δ*hmr* Δ*hml*) and the HO induction lasted 2 h. Note the increase in Smc3 binding in the vicinity of *HOcs* after the DSB. This experiment was performed in parallel to the second repetition for Tel arrested cells shown in Figure 2c.

Figure S4. The principle of the qPCR resection assay. Related to Figure 3a,b. (**a**) Schematic representation of the HO resection assay. Primers (blue arrows) are designed to amplify a sequence that contains a *StyI* target site adjacent to the HO cutting site (*HOcs*). When this restriction enzyme is used on the extracted genomic DNA, amplification is inhibited. If resection extends beyond the *StyI* site, StyI does not cut and the primers can amplify. (**b**) Summary table of the amplification yield obtained after the StyI digestion during *HOcs* resection.

Figure S5. Features of HR to repair the *HOcs* DSB in late mitosis. Related to Figure 3c-f. Cells were first blocked in the *cdc15-2* arrest at 34 °C for 3 hours (Tel). Then, the HO DSB was generated by adding β-estradiol. After 1 hour, the β-estradiol was washed away, and samples were taken to monitor the repair for 3 hours. (**a-c**) Kinetics of the *HOcs* DSB repair in *yku70Δ*. Yku70 is part of a conserved complex that recognizes a DSB and drives its repair towards NHEJ. (**d-f**) Kinetics of the *HOcs* DSB repair in *rad9Δ*. Rad9 is a mediator in the DDC that promotes activation of the effector kinase Rad53. (**g-i**) Kinetics of the *HOcs* DSB repair in *mre11Δ*. Mre1 is part of the MRX complex, which tethers the DSB ends and facilitates end resection. (**a,d,g**) Representative Western blot analyses for HO induction and subsequent degradation (tagged with Flag epitope) and the DSB sensing through Rad53 hyperphosphorylation (note that Rad53 does not get hyperphosphorylated in *rad9Δ*). A PGK1 western blot served as a housekeeping and Ponceau S staining of the membrane as a loading control for all lanes. The leftmost lane in the Ponceau S corresponds to the protein weight marker. (**b,e,h**) Representative Southern blots for the MAT switching assays in late mitosis. Alongside the MAT probe, a second probe against the *ACT1* gene (1.1 kb fragment) was included for normalization. The MAT probe also recognizes an allele-independent MAT distal fragment (2.2 kb). (**c,f,i**) Quantification of relative band intensities in the MAT switching Southern blots (mean ± sem, n=3). Individual values were normalized to the *ACT1* signals. Then, every lane was normalized to *MAT*a at the arrest. Note that repair kinetics is similar to that of the wild type in Figure 3e,f. Tel: Telophase. +βE: β-estradiol addition.

Figure S6. MAT switching yield of Smc3-aid* OsTIR1 upon IAA addition. Cells were treated as in Figure 3d-f but including a 1 h IAA step between the *cdc15-2* arrest and the induction of HO. IAA was then maintained throughout the experiment. (**a**) Representative Western blot as in Figure 3d but also including Smc3-aid* decline after IAA addition. Note that the chromatin-bound acSmc3 is degraded as well. (**b**) Representative Southern blot for the MAT switching assay as in Figure 3e. (**c**) Quantification of relative band intensities for MAT switching Southern blots (mean ± sem, n=3). Values were normalized as in Figure 3f. The MAT switching in the wild type strain without IAA is represented alongside for comparison. Note that there is a decrease in the cutting efficiency of HO (*MAT*a levels remained higher) as well as in the gene conversion into *MAT*α. However, these phenotypic changes are due to IAA rather than cohesin depletion (see Figure 4c). Tel: Telophase. +βE: β-Estradiol addition. +IAA: Indole-acetic acid.