Shorter reported sleep duration in men is associated with worse virtual spatial navigation performance

Emre Yavuz[1], Alpar S Lazar[2], Christoffer J. Gahnstrom[3], Sarah Goodroe[3], Antoine Coutrot[4], Michael Hornberger[2], Hugo J. Spiers[1]

1. Institute of Behavioural Neuroscience, Department of Experimental Psychology, Division of Psychology and Language Sciences, University College London, London, UK

2. Norwich Medical School, University of East Anglia, Norwich, UK

3. Department of Psychology, University of Pennsylvania, Pennsylvania, USA

4. LIRIS - CNRS - University of Lyon, France

* corresponding authors: emre.yavuz.22@ucl.ac.uk, h.spiers@ucl.ac.uk

Abstract

Sleep has been shown to impact navigation ability. However, it remains unclear how different sleep-related variables may be independently associated with spatial navigation performance, and as to whether gender may play a role in these associations. We used a mobile video game app, Sea Hero Quest (SHQ), to measure wayfinding ability in US-based participants. Wayfinding performance on SHQ has been shown to correlate with real-world wayfinding. Participants were asked to report their sleep duration, quality, daytime sleepiness and nap frequency and duration (n = 737, 409 men, 328 women, mean age = 27.1 years, range = 18-59 years). A multivariate linear regression was used to identify which sleep variables were independently associated with wayfinding performance. For men, longer sleep durations were associated with better wayfinding performance. For women, no such association was found. While other sleep variables showed trends of association with wayfinding performance, none of these were significantly associated with wayfinding performance in our regression model. These findings from younger U.S.-based participants suggest that a longer sleep duration may be an important contributor to successful navigation ability in men.

Introduction

Being able to maintain a sense of direction and location in order to find our way in different environments is a fundamental cognitive function that relies on multiple cognitive domains (Newcombe et al., 2022; Spiers et al., 2023). Understanding these individual differences in navigation ability is crucial given that deficits in navigation may constitute the earliest signs of Alzheimer's Disease (Coughlan et al., 2018), and are apparent in other conditions such as traumatic brain injury (Seton et al., 2023). There are also negative effects of disorientation, such as getting lost in everyday environments, leading to distress for patients and family members and in extreme cases death from exposure (Coughlan et al., 2018). Understanding these individual differences in navigation will also advance the field of spatial cognition at large (Newcombe et al., 2022; Spiers et al., 2023). Being able to create a valid test of navigation that accounts for the wide variation in navigation performance is challenging, given the large sample sizes needed and the high levels of environmental manipulation and experimental control required in standard research settings (Newcombe et al., 2022). With the recent evolution of widespread touch-screen technology on both tablet and mobile devices and virtual reality (VR), our team developed a series of navigation tests in the form of a mobile video game app Sea Hero Quest (SHQ)(Spiers et al., 2023). SHQ has since been used to test the navigation ability of 4 million people globally, has good test-retest reliability and is predictive of real-world navigation performance (Coughlan et al., 2020; Coutrot et al., 2018; Coutrot et al., 2019).

Various studies have shown that subjective and objective measures of sleep quality are associated with poorer virtual spatial navigation performance (Coutrot et al., 2022a; Maybrier et al., 2022; Valera et al., 2016; Varga et al., 2016), including sleep duration (Coutrot et al., 2022a; Maybrier et al., 2022), fragmented sleep (Maybrier et al., 2022; Varga et al., 2016) and insomnia-like symptoms (Valera et al., 2016). Experimental sleep deprivation studies have also shown that sleep deprivation results in poorer spatial navigation performance (Feld et al., 2022; Ferrara et al., 2006; 2008; Nguyen et al., 2013), although other studies report no significant changes in spatial navigation performance following sleep

deprivation (Orban et al., 2006; Simon et al., 2022). In a recent study we reported a U-shaped association between self-reported sleep duration and virtual spatial navigation performance on SHQ, where mid-range sleep duration was associated with optimal performance (Coutrot et al., 2022a). This latter study corroborates broader findings of a U-shaped association between self-reported sleep duration and other cognitive domains aside spatial navigation (Fjell et al., 2023; Li et al., 2022; Lo et al., 2015), even when controlling for other sleep-related variables such as daytime sleepiness, sleep apnea and insomnia symptoms (Kyle et al., 2017; Tai et al., 2022). However, it remains unclear as to how different sleep-related variables may independently be associated with spatial navigation performance.

Gender may also play a role in the association between sleep and spatial navigation performance. Many studies have reported gender differences in sleep-related variables, where sleep duration is greater in women when based on both self-report and objective measures (Coutrot et al., 2022a; Su et al., 2022), and have found that the association between self-reported sleep duration and cognitive function differs by gender, where shorter and longer sleep durations are associated with worse cognitive performance only in men (Jackowska & Cadar, 2021). Women also report poorer subjective sleep quality than men (Burgard and Ailshire, 2013; Fatima et al., 2016; Kyle et al., 2017; Tang et al., 2017). Many studies also demonstrate gender differences in both virtual and real-world navigation ability in favour of males (Nazareth et al., 2019; Spiers et al., 2023; Coutrot et al, 2018). However, it remains unclear as to how gender influences the association between different sleep-related variables and navigation performance.

With both poorer sleep quality and poorer spatial navigation performance being associated with a higher risk of developing various neurodegenerative disorders (Coughlan et al., 2018; Joza et al., 2023; Lucey et al., 2021), a deeper understanding of the association between sleep and spatial navigation performance at the population level will provide greater insight into the factors that may contribute to cognitive decline.

Bringing together these findings, we hypothesised that sleep duration would show a U-shaped (quadratic) association with navigation ability that would differ by gender, where men who report sleeping for mid-range durations would have significantly better navigation ability than women. We also hypothesised that sleep duration would show a significant linear association with spatial navigation ability that would also differ by gender, where the linear relationship between sleep duration and navigation ability would be stronger in males. In addition to our hypotheses, we explored whether other self-reported sleep-related variables aside sleep duration would be independently associated with navigation ability and whether these associations would differ by gender.

Methods

Participants

903 participants living in the US aged 18 and above (365 men, 459 women, mean age = 27.1 years, SD = 8.0 years, range = 18-66 years, mean number of years spent in formal education = 16.1 years, mean BMI = 26.4, 285 currently living cities, 556 currently living outside cities) were recruited using the Prolific database (www.prolific.co, 2023) and reimbursed for their time. Ethical approval was obtained from the University College London Review Board conforming to Helsinki's Declaration. All participants provided informed written consent. We removed 17 participants who selected 'other' for gender, as we were primarily interested in the differences in the associations between the sleep variables and wayfinding distance in men and women. We then excluded outliers in 2 stages. We first removed participants whose time spent in bed was less than 4 hours and greater than 12 hours, as we believed these participants' responses were not genuine ot considered them to be either extremely short or extremely long sleepers respectively, based on a criteria used by a previous study (Mohlenhoff et al., 2018). This resulted in the removal of 62 participants, and a remaining sample of 824 participants. We then further identified outliers in this remaining sample of 824 participants using Mahalanobis' Distance, a method shown to have high

sensitivity and specificity and a minimal change in bias in simulated and real datasets when removing outliers based on questionnaire data compared to other methods (Curran, 2016; Zijlstra et al., 2011)(Supplementary Materials). 97 participants were removed as outliers. This resulted in a final sample of 724 participants (328 men, 409 women, mean age = 26.4 years, SD = 7.2 years, range = 18-59 years, mean number of years spent in formal education = 16.1, mean Body Mass Index (BMI) = 26.0, 248 currently living cities, 476 currently living outside cities). Demographic information for the final sample is summarised in Table 1. Data analysis was completed using R studio (version 1.4.1564) and Python (version 3.9.12). T-tests were used to determine whether sleep-related variables differed in men and women for the continuous sleep variables. A chi-squared analysis was used to determine whether the frequency of alarm use differed in men and women given that this was a categorical variable.

Statistical Power Analysis

A power analysis was conducted using G* power (Erdfelder et al., 1996). Given that a previous study looking at the association between sleep quality and spatial navigation ability found a medium effect size for this association (Varga et al., 2016), it was determined that 737 participants were sufficient to acheive a medium effect size (Cohen's f2 = 0.15) at an alpha threshold of 0.05 with 95% power (Selya et al., 2012).

Experimental Procedure

Self-report questionnaires

To characterize self-reported sleep patterns, we used a series of questions about habitual bedtime, wake-up time, difficulty waking up, overall sleep quality, number of awakenings during the night, time spent awake during the night, sleep duration, sleepiness on waking (sleep inertia), nap duration, frequency of

naps, and frequency of alarm use (Supplementary Materials). As sleep quality was originally coded in the opposite direction to the other variables (1 = maximum, 10 = minimum), we reverse coded this variable to be consistent with the other variables with the coding strategy used forthe other variables we collected (1 = minimum, 10 = maximum). We also calculated three additional sleep-based measures and the sleepiness resolution index (SRI). The SRI was calculated as the ratio of the reported sleepiness upon bedtime over that upon wake-up time (reference). Thus, an SRI value of greater than 1 would indicate that one was more sleepy when going to bed than when waking up. Time in bed was calculated as the time difference between reported bedtime and wakeup time.

To characterize the individual video game use, participants were asked to indicate the number of hours per week spent playing video games on all kind of devices and the number of hours of phone use per week.

Participants were also asked to report their age, gender, highest education level, height (in feet and inches), smoking frequency (currently, previous or never smoked), weekly units of alcohol intake, daily cups of caffeine, frequency of daily strenuous physical activity. BMI was calculated as weight (kg) divided by height squared (m^2).

Please see the Supplementary Materials for the full set of questionnaires.

Sea Hero Quest Task

Sea Hero Quest (SHQ) is a VR-based video game for mobile and tablet devices which requires participants to navigate through a three-dimensional world on a boat to search for sea creatures in order to photograph them, with the environment consisting of rivers and lakes (Coutrot et al., 2018). Although navigational abilities on SHQ have been assessed using Wayfinding, Path Integration and Radial Arm

Maze measures, we focused on wayfinding in this study (Figure 1). Please refer to Yavuz et al., 2023 for specific details.

Figure 1. Outline of the wayfinding task.

(A) At the start of each game level participants were presented with a map indicating the goals which they had to navigate to and the order of visitation, where '1' indicates goal number 1 etc. (the map from level 32 is shown as an example). Level 1 (not shown) provided one goal and a simple river with no choices to reach the goal as training. (B) Participants selected to close the map by

pressing 'close', at which point the participant had to start navigating to the goals using their memory of the map. First-person view of the environment is shown, where the participant tapped left or right of the boat to steer it. (C) Examples of individual player trajectories (level 32) from the start location to the final goal. Trajectories are ordered by performance, with the top left providing the best performance (shortest trajectory length), through to bottom right who has the worst performance (longest trajectory length). Adapted from Coutrot et al. (2018).

Data Analysis

To investigate the relationship between sleep and spatial navigation, as measured by the self-reported questions and wayfinding distance in SHQ, we first ran partial spearman's correlations, controlling for age, gender, highest level of education achieved, daily hours of sunlight, daily cups of caffeine, daily hours of strenuous physical activity, weekly units of alcohol and Body Mass Index (BMI). We included these covariates to control for factors associated with sleep or wayfinding distance aside the sleep-related variables (Bisson and Lachman, 2023; Coutrot et al., 2022b; Shattuck and Matsangas et al., 2017; Nunez et al., 2020; Yavuz et al., 2023; Zheng et al., 2020). We also included weekly hours of phone use as a covariate given that SHQ is a mobile phone-based video game. To measure the associations between the frequency of alarm use and wayfinding distance, we ran a chi-squared analysis given that the frequency of alarm use was a categorical variable. For these chi-squared analyses, a higher Cramer's V effect size indicated a stronger association between binary categorical variables (Kerney, 2017).

We then entered the five self-reported sleep variables with the strongest effect sizes (i.e., the strongest association with wayfinding distance as indicated by the spearman's correlation coefficient) into a multivariate linear regression model. This was to determine the independent associations between each of the sleep variables and navigation performance when controlling for other variables. Age, gender, highest level of education, weekly hours of video gaming, weekly hours of phone use, daily hours of sunlight,

daily cups of caffeine, daily hours of strenuous physical activity and Body Mass Index (BMI) were also included as covariates in this multivariate model for the reasons mentioned above. A sleep variable x gender interaction term for each sleep variable was included in the model as we were interested in the role played by gender in the association between the sleep-related variables and navigation. Given that sleep duration has been shown to have both a U-shaped association with spatial navigation ability on SHQ (Coutrot et al., 2022b) and a linear association with spatial navigation ability on other tasks (Maybrier et al., 2022), we specified two models. In the first model, sleep duration was included as a linear term (Table 3). In the second model, sleep duration was included as a quadratic term (Table 4).

As a check for multicollinearity, we calculated the variance inflation factor (*VIF*) for each predictor variable in each model (Supplementary Tables S1 and S2). A *VIF* value of <5 indicates that multicollinearity is not a concern (Fox and Monet, 1992; Kim, 2019). Post-hoc t-tests bonferroni-corrected for multiple comparisons to control for type 1 errors were carried out where main effects and interactions were significant. When conducting post-hoc t-tests for sleep duration, we binned sleep duration into <=7 hours and >7 hours of sleep to match the approach taken by other studies (Jackowska & Cadar, 2020; Fatima et al., 2016; Hirshkowitz et al., 2015; Li et al., 2022). Given that only 5 participants slept for more than 9 hours on a typical night, we grouped those who slept for 9 hours or 10 hours on a typical night into one group (>= 9 hours on a typical night) when visualising the data.

	Female	Male
n	410	314
Weighted wayfinding distance	12.21 (4.03)	11.06 (4.88)
Age	25.27 (6.83)	27.98 (7.38)

Highest level of education achieved (n, %)		
Some formal education	5 (1.2)	2 (0.6)
High school	109 (26.6)	111 (35.4)
2-year college/uni	57 (13.9)	43 (13.7)
4-year college/uni	178 (43.4)	117 (37.3)
Masters	55 (13.4)	30 (9.6)
PhD/MD	6 (1.5)	11 (3.5)
Hours of phone use per week	35.60 (23.08)	28.59 (21.54)
Weekly hours of video gaming on all devices	4.67 (5.23)	9.67 (6.19)
BMI	26.44 (7.24)	25.34 (5.67)
Daily hours of sunlight	1.24 (1.11)	1.38 (1.24)
Daily cups of coffee	1.12 (1.09)	1.44 (1.48)
Weekly units of alcohol	3.09 (5.14)	3.04 (5.16)
Frequency of daily strenuous physical activity (%)		
Never	9.8	10.5
Rarely	37.8	38.9
Sometimes	37.8	37.9
Often	12.7	9.9
Always	2.0	2.9

Smoking frequency (%)

Never smoked	85.1	72.0
Former smoker	11.5	16.2
Current smoker	3.4	11.8

Table 1. Summary of basic descriptive statistics for the variables included in the model (weighted wayfinding distance, age, highest level of education achieved, weekly hours of video game use on all devices, weekly hours of phone use, BMI, daily hours of sunlight, daily cups of coffee, weekly units of alcohol and frequency of daily strenuous physical activity) in men and women separately. Mean and standard deviation (SD) values are shown. Men were significantly older than women, spent significantly greater weekly hours video gaming and had significantly fewer weekly hours of phone use.

Results

Self-reported sleep-related characteristics

Please see Table 2 for self-reported sleep-related characteristics in men and women.

Variable	Female	Male	*t*	χ2	*p*
Sleep duration (hours) (mean ± SD)	7.23 ± 0.97	7.05 ± 1.03	2.33		0.020
Nap duration (mins) (mean ± SD)	47.32 ± 41.13	35.82 ± 37	1.41		0.158
Number of times one wakes up during the	1.34 ± 1.15	1.06 ± 1.09	-1.08		0.282

night (mean ± SD)				
Sleep quality (mean ± SD)	7.42 ± 1.86	7.39 ± 2.03	2.3	0.022
Sleepiness on going to bed (mean ± SD)	5.79 ± 2.45	5.81 ± 2.36	0.25	0.803
Sleepiness resolution index (mean ± SD)	1.2 ± 0.88	1.27 ± 0.93	3.95	**<0.001**
Sleep inertia (mean ± SD)	6.1 ± 2.36	5.7 ± 2.27	3.43	**0.001**
Time in bed (hours) (mean ± SD)	8.28 ± 1.47	7.77 ± 1.4	4.69	**<0.001**
Time spent awake during the night (mins)	21.2 ± 30.78	18.13 ± 27.47	-0.12	0.907
Frequency of alarm use			*30.15	*<0.00
Frequency of alarm use - never (%)	5.9	16.6		
Frequency of alarm use - rarely (%)	10.0	13.1		
Frequency of alarm use - often (%)	21.5	21.3		
Frequency of alarm use - sometimes (%)	28.5	27.1		
Frequency of alarm use - always (%)	34.1	22		

Table 2. Gender differences in sleep-related variables. Significant P-values are highlighted in bold when applying a bonferroni-corrected alpha threshold of 0.005. *For the categorical variable of frequency of alarm use, a single effect size and p-value are printed given that each of the categories were compared with one another.

Multivariate analysis - in males, shorter reported sleep was associated with worse navigation

Based on an initial bivariate analysis, difficulty waking up, sleep inertia, sleepiness resolution index, time spent awake during the night and sleep quality were chosen as sleep variables of interest to include in our

multivariate linear regression model, given that these variables had the strongest associations with weighted wayfinding distance (see Supplementary Table 3 for the bivariate analysis results). Interactions between gender and each of the sleep variables were included in the model. As for the partial correlations, age, gender, highest level of education achieved, hours of weekly video gaming, hours of weekly phone use, daily hours of sunlight, BMI, weekly units of alcohol, daily cups of caffeine, smoking frequency and frequency of daily strenuous physical activity were included as covariates in the model (Table 3). Before running the regression, as a check for multicollinearity, we calculated the variance inflation factor (VIF) for each predictor variable in the model. This revealed that all the predictor variables had a VIF value that was less than 5, indicating that multicollinearity was not a concern (Supplementary Table S1). Please see Figure 2 for correlations between each of the sleep variables included in the model.

	Weighted wayfinding distance	Difficulty waking up	Sleep duration (hours)	Sleepiness on waking	Sleep quality	Time spent awake during the night (mins)	Sleepiness resolution index
Weighted wayfinding distance	1	-0.21	-0.07	-0.18	0.13	0.05	0.13
Difficulty waking up	-0.21	1	0.01	0.67	-0.34	0.09	-0.52
Sleep duration (hours)	-0.07	0.01	1	-0.08	0.23	-0.17	0.05
Sleepiness on waking	-0.18	0.67	-0.08	1	-0.32	0.1	-0.71
Sleep quality	0.13	-0.34	0.23	-0.32	1	-0.31	0.27
Time spent awake during the night (mins)	0.05	0.09	-0.17	0.1	-0.31	1	-0.13
Sleepiness resolution index	0.13	-0.52	0.05	-0.71	0.27	-0.13	1

Figure 2. Associations between each of the sleep variables included in the final model and wayfinding distance. Pearson's correlation coefficients for the associations between each of the sleep variables (sleep quality, sleep inertia, sleepiness resolution index, sleep duration (hours), time spent awake during the night and difficulty waking up) included in the model and and wayfinding distance (across game levels across participants). Values highlighted in red represent significant associations ($p < 0.05$).

Model

Weighted wayfinding distance ~ Sleep duration + Sleep duration*Gender + Difficulty waking up +

Difficulty waking up*Gender + Sleepiness resolution index + Sleepiness resolution index*Gender +

Sleep quality + Sleep quality*Gender + Time spent awake during the night + Time spent awake during

the night*Gender + Sleep inertia + Sleep inertia*Gender + Gender + Age + Weekly hours of video

gaming + Weekly hours of phone use + BMI + Daily hours of sunlight + Highest education level

achieved + Daily cups of caffeine + Weekly units of alcohol + Frequency of daily strenuous physical

activity + Smoking frequency

Table 3. Model specification with sleep duration as a linear term.

The outputs from the multivariate linear regression model are as follows:

Sleep-related variables

There was a significant interaction between sleep duration and gender, with longer sleep being associated

with shorter wayfinding distance and therefore better navigation performance in men (β = -0.11, $f2$ =

0.01, p = 0.032, CI = [-0.22, -0.01])(Table 4 and Figure 3). Bonferroni-corrected post-hoc t-tests showed

that men who slept for more than 7 hours were significantly better at navigating than women (t = 3.56, p

< 0.001). However, in those who slept for 7 hours or less, this gender difference in navigation ability was

no longer significant (t = 1.31, p = 0.191)(Table 5). There were no significant associations between sleep

quality, difficulty waking up, sleep inertia, sleepiness resolution index or time spent awake during the

night and wayfinding distance (p > 0.05)(Table 4 and Figure 3).

Variable	β	95% CI	t	p	sig	f2
(Intercept)	0.07	[-0.05, 0.20]	1.16			
Age	0.12	[0.06, 0.18]	3.94	**<0.001**	***	0.04
Male gender	-0.11	[-0.23, <0.01]	-1.95	0.052	.	0.02
Difficulty waking up	-0.06	[-0.15, 0.03]	-1.35	0.177		0.03
Sleep duration (hours)	0.01	[-0.07, 0.08]	0.14	0.890		<0.01
Sleep inertia	-0.03	[-0.14, 0.08]	-0.52	0.600		<0.01
Sleep quality	0.01	[-0.06, 0.09]	0.36	0.719		0.01
Time spent awake during the night (mins)	0.03	[-0.04, 0.10]	0.83	0.408		0.01
Sleepiness resolution index	0.00	[-0.10, 0.10]	0.05	0.959		<0.01
Male gender*Difficulty waking up	-0.03	[-0.17, 0.10]	-0.48	0.629		<0.01
Male gender*Sleep duration (hours)	-0.11	[-0.22, -0.01]	-2.15	**0.032**	*	0.01
Male gender*Sleep inertia	-0.06	[-0.22, 0.11]	-0.66	0.509		<0.01
Male gender*Sleep quality	0.08	[-0.03, 0.20]	1.41	0.158		<0.01
Male gender*Time spent awake during the night (mins)	0.04	[-0.07, 0.15]	0.76	0.446		<0.01
Male gender*Sleepiness resolution index	-0.05	[-0.19, 0.09]	-0.66	0.511		<0.01
Weekly hours of video gaming on all devices	-0.11	[-0.17, -0.06]	-4.06	**<0.001**	***	0.02

Weekly hours of phone use	0.09	[0.04, 0.14]	3.55	**<0.001**	***	0.02
Daily hours of sunlight	0.02	[-0.03, 0.07]	0.70	0.485		<0.01
Highest level of education achieved	0.03	[-0.09, 0.14]	0.48	0.633		<0.01
BMI	-0.06	[-0.11, -0.01]	-2.26	**0.024**	*	0.01
Weekly units of alcohol	0.02	[-0.04, 0.07]	0.59	0.553		<0.01
Daily cups of caffeine	0.02	[-0.03, 0.08]	0.84	0.404		<0.01
Smoking frequency	-0.10	[-0.23, 0.04]	-1.42	0.155		<0.01
Frequency of daily strenuous physical activity	-0.10	[-0.20, <0.01]	-1.91	0.057	.	<0.01

Table 4. Results of the multiple linear regression model associating sleep-related variables with wayfinding distance when controlling for associated covariates. P-values for the significant associations are highlighted in bold. See Yavuz et al (2023) for consideration of the impact of video games and phone use variables.

Sleep Duration	*t*	*p*
<=7 hours	1.31	0.191
>7 hours	3.56	**<0.001**

Table 5. Bonferroni-corrected post-hoc t-tests comparing wayfinding distance in those who reported sleeping <=7 hours with those who slept >7 hours on a typical night. P-values for the significant associations are highlighted in bold when applying a bonferroni-corrected alpha threshold of 0.025.

A)

B)

C)

D)

E)   F)

Figure 3 Associations between each of the sleep-related variables and wayfinding distance in men and women. Higher wayfinding distance indicates worse navigation ability. VR-m = Virtual reality meters. (A) The asterix indicates the significant interaction between gender and sleep duration at 5 hours of sleep duration (* = p < 0.05). (B) A value of 1 indicates that the level of sleepiness when going to bed = the level of sleepiness when waking up, a value exceeding 1 indicates that the level of sleepiness on going to bed > the level of sleepiness when waking up and a value of less than 1 indicates that the level of sleepiness on going to bed < level of sleepiness when waking waking up. (C) 1 = minimum difficulty waking up, 10 = maximum difficulty waking up. (D) 1 = minimum sleep inertia, 10 = maximum sleep inertia. (E) 1 = minimum sleep quality, 10 = maximum sleep quality. (A-F) Data points represent the mean wayfinding distance across game levels across participants. Bars represent one standard error above and below the mean wayfinding distance across game levels across participants.

Complementary analysis

Conducting the model with sleep duration as a quadratic term

When conducting the model with sleep duration specified as a quadratic term (Table 6), there was no significant interaction between sleep duration and gender ($\beta = 0.04$, $f2 = {<}0.01$, $p = 0.251$, $CI = [-0.03, 0.12]$)(Table 7).

Model

Weighted wayfinding distance ~ Sleep duration^2 + Sleep duration^2*Gender + Difficulty waking up + Difficulty waking up*Gender + Sleepiness resolution index + Sleepiness resolution index*Gender + Sleep quality + Sleep quality*Gender + Time spent awake during the night + Time spent awake during the night*Gender + Sleep inertia + Sleep inertia*Gender + Gender + Age + Weekly hours of video gaming + Weekly hours of phone use + BMI + Daily hours of sunlight + Highest education level achieved + Daily cups of caffeine + Weekly units of alcohol + Frequency of daily strenuous physical activity + Smoking frequency

Table 6. Model specification with sleep duration as a quadratic term.

Variable	*β*	*95% CI*	*t*	*p*	sig	*f2*
(Intercept)	0.04	[-0.09, 0.17]	0.57	0.572		0.04
Age	0.12	[0.06, 0.18]	4.03	**<0.001**	***	0.02
Male gender	-0.15	[-0.28, -0.01]	-2.12	**0.034**	*	0.03

Difficulty waking up	-0.06	[-0.15, 0.03]	-1.35	0.179		0.01
Sleep duration (hours)^2	0.03	[-0.02, 0.08]	1.21	0.227		<0.01
Sleep inertia	-0.03	[-0.13, 0.08]	-0.47	0.641		0.01
Sleep quality	0.02	[-0.06, 0.10]	0.47	0.637		0.01
Time spent awake during the night (mins)	0.02	[-0.04, 0.09]	0.71	0.476		<0.01
Sleepiness resolution index	0.01	[-0.09, 0.11]	0.16	0.871		0.03
Male gender*Difficulty waking up	-0.04	[-0.18, 0.09]	-0.62	0.536		<0.01
Male gender*Sleep duration (hours)^2	0.04	[-0.03, 0.12]	1.15	0.251		<0.01
Male gender*Sleep inertia	-0.03	[-0.19, 0.13]	-0.34	0.734		<0.01
Male gender*Sleep quality	0.08	[-0.03, 0.20]	1.44	0.152		<0.01
Male gender*Time spent awake during the night (mins)	0.06	[-0.05, 0.17]	1.11	0.267		<0.01
Male gender*Sleepiness resolution index	-0.04	[-0.19, 0.10]	-0.60	0.546		<0.01
Weekly hours of video gaming on all devices	-0.12	[-0.18, -0.07]	-4.36	**<0.001**	***	0.02
Weekly hours of phone use	0.09	[0.04, 0.14]	3.43	**0.001**	***	<0.01
Daily hours of sunlight	0.02	[-0.03, 0.07]	0.67	0.502		<0.01
Highest level of education achieved	0.03	[-0.08, 0.15]	0.59	0.556		0.01
BMI	-0.06	[-0.11, -0.01]	-2.23	**0.026**	*	<0.01
Weekly units of alcohol	0.02	[-0.04, 0.07]	0.61	0.540		<0.01
Daily cups of caffeine	0.03	[-0.02, 0.08]	1.07	0.284		<0.01

Smoking frequency	-0.09	[-0.23, 0.04]	-1.38	0.168		<0.01
Frequency of daily strenuous physical activity	-0.10	[-0.20, <0.01]	-1.89	0.059	.	<0.01

Table 7. Results of the multiple regression model associating sleep-related variables with wayfinding distance, specifying sleep duration as a quadratic term, when controlling for associated covariates. P-values for the significant associations are highlighted in bold.

Conducting the linear model with sleep duration as a linear term in a smaller subsample excluding those spending <5 and > 10 hours in bed (n = 658)

We also ran the same model on a smaller subsample of 658 participants after excluding those who slept for >10 hours or <5 hours (rather than >12 or <4 hours in the current sample), given that our recent study associating sleep duration with navigation ability also used this exclusion criteria (Coutrot et al., 2022b). The interaction between sleep duration and gender in this model was not significant (β = 0.08, $f2$ = <0.01, p = 0.167, CI = [-0.03, 0.19])(Supplementary Table S6). However, sleep duration (hours) showed a significant negative association with wayfinding distance (β = -0.10, $f2$ = <0.01, p = 0.018, CI = [-0.18, -0.02])(Supplementary Table S6), where greater hours of sleep were associated with a shorter wayfinding distance and thus better navigation performance. Additionally, the time spent awake during the night (minutes) showed a significant positive association with wayfinding distance (β = 0.09, $f2$ = 0.01, p = 0.045, CI = [<0.01, 0.17])(Supplementary Table S6), where a greater time spent awake during the night was associated with a longer wayfinding distance and thus worse navigation ability.

Results Summary

Taken together, there are 3 key findings: 1) greater sleep duration was associated with significantly better navigation performance only in men when specifying sleep duration as a linear term, 2) sleep duration did not show a significant interaction with gender when specifying sleep duration as a quadratic term, 3) neither difficulty waking up, sleep inertia, sleepiness resolution index, sleep quality or time spent awake during the night were significant independent predictors of navigation performance. We also found that when restricting the sleep range to match a previous analysis that only included sleep duration, greater difficulty waking up and a shorter sleep duration .

Discussion

Here, we investigated the association between multiple sleep-related variables and human spatial navigation performance. Men reporting 7 hours or less sleep had worse wayfinding performance. This supports our hypothesis that sleep duration would show a significant linear association with navigation ability that would differ by gender. Accordingly, this finding aligns with previous findings of a gender difference in the effect of sleep duration on cognitive function in a large cohort (Jackowska & Cadar, 2020), but not with studies where no association has been found in both navigation and non-navigation-related domains (Ngyuen et al., 2013; Vallat et al., 2022; van Oostrom et al., 2018). Several factors may have accounted for these discrepancies, such as circadian rhythms and the time one spends awake during the day, which have been found to influence gender differences in cognitive function and which were not examined here (Santhi et al., 2016). Additionally, a longer sleep duration has been associated with greater levels of physical activity the following day (Bisson & Lachman, 2023), which in turn may have been associated with better navigation ability. However, we controlled for the frequency of daily strenuous physical activity in our analysis, suggesting that this explanation is unlikely.

When conducting a model with sleep duration specified as a quadratic term, we did not find a significant interaction between sleep duration and gender, contradicting our hypothesis that men would have better

navigation performance than women specifically at the mid-range sleep duration. This finding may contradict previous findings showing that those with mid-range sleep durations have better cognitive performance than those with shorter and longer sleep durations, in both navigation-related (Coutrot et al., 2022b) and non-navigation-related cognitive tasks (Fjell et al., 2023; Jackowska & Cadar, 2021; Lucey et al., 2021; Ma et al., 2020; Maybrier et al., 2022; Mohlenhoff et al., 2018; Van Dongen et al., 2003; Van Oostrom et al., 2018). Several factors may have accounted for these discrepancies. Our sample was solely US-based, whilst Coutrot et al., 2022b used participants from all around the world. Age may also be another key factor. Our sample was a relatively young sample (mean age = 26.4 years), whilst Coutrot et al., 2022b used participants with a mean age of 38.7 years. Other studies have also shown that participants in the middle range of the cohort in terms of age had better cognitive performance at shorter and longer sleep durations than both younger and older participants (Mohlenhoff et al., 2018), as well as showing that in participants of a similar age to ours, sleep can facilitate performance on memory-based tasks but not spatial navigation performance (Simon et al., 2022), and that sleep fragmentation is associated with poorer wayfinding performance only in older participants but not in younger participants (Maybrier et al., 2022). Thus, the relatively young age of our sample may have accounted for discrepancies between our findings and previous findings. Nonetheless, younger participants have previously been found to have significantly improved wayfinding performance following a night's sleep whilst older participants did not (Varga et al., 2016). Thus, further investigation in samples of a wider age range is warranted.

We found no significant associations between our exploratory sleep-related measures of sleep inertia, difficulty waking up, sleep quality, sleepiness resolution index or time spent awake during the night and wayfinding distance. These findings may seem to contradict previous findings showing that increased sleep fragmentation is associated with poorer wayfinding performance (Varga et al., 2016), and that poorer sleep quality is associated with a reduced ability to use shortcuts during wayfinding and thus less efficient wayfinding performance (Valera et al., 2016). However, our smaller subsample analysis (conducted to match Coutrot et al., 2022a) did find that less time spent awake during the night was

significantly associated with better navigation performance, providing some support towards the possibility that these variables influence navigation performance. Additionally, poorer sleep may not always lead to changes in cognitive function, with a study demonstrating that greater sleep fragmentation was significantly associated with reduced glucose metabolism but not cognitive function (Andre et al., 2019). In our study, men with greater sleep inertia, greater difficulty waking up and a lower sleepiness resolution index trended towards having better navigation performance, albeit non-significant associations. This may suggest that poorer sleep may in fact be associated with better navigation depending on the facets of sleep examined. In line with this, converting to Lewy-Body Dementia and Parkinson's Disease Dementia is associated with improved sleep quality, as indicated by a lower severity of insomnia, and having frequent insomnia symptoms has been found to be associated with better cognitive performance (Joza et al., 2023; Kyle et al., 2017).

Our finding of a significant associations between the time spent awake during the night and sleep duration with navigation ability in our smaller subsample analysis might suggest that these aspects of sleep play a more important in role in shaping one's navigation ability than others, and provides a platform for further studies determining whether certain aspects of sleep may offer more protection against cognitive decline. The non-significant trends found in our study, as well as differences in the experimental task, study design (cross-sectional vs longitudinal) and samples used in our study compared with former studies warrants further investigation to firmly establish the directionality of any associations between sleep and navigation ability.

Limitations and Future Directions

While our study allowed assessment of over 700 participants on an ecologically valid (Coutrot et al., 2019) and reliable cognitive test of navigation (Couglan et al. 2020), there are a number limitations to consider that future studies could address. Our study relied on self-report measures of sleep duration,

resulting in participants being excluded for inaccurately reporting sleep-related measures, as well

potential differences in findings due to a discrepancy between self-reported and objective sleep-related

measures (Fatima et al., 2016). Smaller sample sizes for those who slept for shorter and longer sleep

durations, where only 5 participants reported sleeping more than 9 hours on a typical night, may have led

to an underrepresentation of the association between sleep and navigation ability. Given the cross-

sectional nature of our study, future studies should consider an interventional and longitudinal design (Li

et al., 2022), especially considering that gender can predict night-to-night variation in sleep duration and

differences in sleep duration on weekdays compared with weekends (Willoughby et al., 2023).

Incorporating objective sleep-based measures and neural recordings in such studies will help identify

individuals who may be more likely to benefit from sleep-based therapeutic interventions and effectively

target those at greater risk of cognitive decline. Additionally, conducting these studies across countries

and cultures will help determine the extent to which associations between sleep and navigation

performance may differ across different populations, as has been shown for both self-reported sleep

duration (Willoughby et al., 2023) and self-reported navigation ability (Walkowiak et al., 2023). Such

future studies would also benefit from using a broader range of virtual navigation and spatial memory

tests, that extend to real-world environments (Coutrot et al., 2019; He et al., 2021; Hill et al., 2023;

Howard et al., 2014; Javadi et al., 2019; Patai et al., 2019). This would be useful to help explore our

incidental observation that a higher BMI was associated with worse navigation performance, a finding

that mirrors evidence that high BMI is associated with worse episodic memory (Cheke et al. 2016).

Conclusion

Overall, our findings suggest that the effects of sleep duration on navigation ability may differ by gender.

Accordingly, this research will serve as a platform for future research looking into how actively changing

one's sleep habits may improve their navigation ability and provide a greater understanding as to why

individual differences in navigation ability may exist.

Acknowledgements

We would like to thank all the participants who volunteered to take part in this research. This research is part of the Sea Hero Quest initiative funded and supported by Deutsche Telekom. Alzheimer's Research UK (ARUK-DT2016-1) funded the analysis; Glitchers designed and produced the game; and Saatchi and Saatchi London managed its creation. EY was funded by The Leverhulme Trust. We thank Masoud Tasmanian for helpful comments on this manuscript.

Data availability

A dataset containing the preprocessed trajectory lengths and demographic information is available at https://osf.io/7nqw6/?view_only=6af022f2a7064d4d8a7e586913a1f157. Due to its considerable size (around 1 terabyte), a dataset with the full trajectories is available on a dedicated server: https://shqdata.z6.web.core.windows.net/. We also set up a portal where researchers can invite a targeted group of participants to play SHQ and generate data about their spatial navigation capabilities. Those invited to play the game will be sent a unique participant key, generated by the SHQ system according to the criteria and requirements of a specific project decided by the experimenter. https://seaheroquest.alzheimersresearchuk.org/. Access to the portal will be granted for non-commercial purposes. Future publications based on this dataset should add 'Sea Hero Quest Project' as a co-author.

Code availability

The code used to produce this data is accessible at: https://osf.io/d5q4r/

References

André, C., Tomadesso, C., Flores, R., Branger, P., Rehel, S., Mézenge, F., Landeau, B., Sayette, V., Eustache, F., Chételat, G., & Rauchs, G. (2019). Brain and cognitive correlates of sleep fragmentation in elderly subjects with and without cognitive deficits. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, *11*(1), 142–150. https://doi.org/10.1016/j.dadm.2018.12.009

Bisson, A. N. S., & Lachman, M. E. (2023). The relationship of daily physical activity and sleep in adults: variations by age, sex, and race. *Journal of Behavioral Medicine*. https://doi.org/10.1007/s10865-022-00387-2

Burgard, S. A., & Ailshire, J. A. (2013). Gender and Time for Sleep among U.S. Adults. *American Sociological Review*, *78*(1), 51–69. https://doi.org/10.1177/0003122412472048

Cheke, L. G., Simons, J. S., & Clayton, N. S. (2016). Higher body mass index is associated with episodic memory deficits in young adults. *Quarterly Journal of Experimental Psychology*, *69*(11), 2305-2316.

Coughlan, G., Coutrot, A., Khondoker, M., Minihane, A.-M., Spiers, H., & Hornberger, M. (2019). Toward personalized cognitive diagnostics of at-genetic-risk Alzheimer's disease. *Proceedings of the National Academy of Sciences*, *116*(19), 9285–9292. https://doi.org/10.1073/pnas.1901600116

Coughlan, G., Laczó, J., Hort, J., Minihane, A.-M., & Hornberger, M. (2018). Spatial navigation deficits — overlooked cognitive marker for preclinical Alzheimer disease? *Nature Reviews Neurology*, *14*(8), 496–506. https://doi.org/10.1038/s41582-018-0031-x

Coughlan, G., Puthusseryppady, V., Lowry, E., Gillings, R., Spiers, H., Minihane, A.-M., & Hornberger, M. (2020). Test-retest reliability of spatial navigation in adults at-risk of Alzheimer's disease. *PLOS ONE*, *15*(9), e0239077. https://doi.org/10.1371/journal.pone.0239077

Coutrot, A., Lazar, A. S., Richards, M., Manley, E., Wiener, J. M., Dalton, R. C., Hornberger, M., & Spiers, H. J. (2022a). Reported sleep duration reveals segmentation of the adult life-course into three phases. *Nature Communications*, *13*(1), 7697. https://doi.org/10.1038/s41467-022-34624-8

Coutrot, A., Manley, E., Goodroe, S., Gahnstrom, C., Filomena, G., Yesiltepe, D., Dalton, R. C., Wiener, J. M., Hölscher, C., Hornberger, M., & Spiers, H. J. (2022b). Entropy of city street networks linked to future spatial navigation ability. *Nature*, *604*(7904), 104–110. https://doi.org/10.1038/s41586-022-04486-7

Coutrot, A., Schmidt, S., Coutrot, L., Pittman, J., Hong, L., Wiener, J. M., Hölscher, C., Dalton, R. C., Hornberger, M., & Spiers, H. J. (2019). Virtual navigation tested on a mobile app is predictive of real-world wayfinding navigation performance. *PLOS ONE*, *14*(3), e0213272. https://doi.org/10.1371/journal.pone.0213272

Coutrot, A., Silva, R., Manley, E., de Cothi, W., Sami, S., Bohbot, V. D., Wiener, J. M., Hölscher, C., Dalton, R. C., Hornberger, M., & Spiers, H. J. (2018). Global Determinants of Navigation Ability. *Current Biology*, *28*(17), 2861-2866.e4. https://doi.org/10.1016/j.cub.2018.06.009

Creery, J. D., Brang, D. J., Arndt, J. D., Bassard, A., Towle, V. L., Tao, J. X., Wu, S., Rose, S., Warnke, P. C., Issa, N. P., & Paller, K. A. (2022). Electrophysiological markers of memory consolidation in the human brain when memories are reactivated during sleep. *Proceedings of the National Academy of Sciences*, *119*(44). https://doi.org/10.1073/pnas.2123430119

Ekstrom, A. D., Spiers, H. J., Bohbot, V. D., & Rosenbaum, R. S. (2018). *Human spatial navigation.* Princeton University Press.

Erdfelder, E., Faul, F., & Buchner, A. (1996). GPOWER: A general power analysis program. *Behavior Research Methods, Instruments, & Computers*, *28*(1), 1–11. https://doi.org/10.3758/bf03203630

Fatima, Y., Doi, S. A. R., Najman, J. M., & Mamun, A. A. (2016). Exploring Gender Difference in Sleep Quality of Young Adults: Findings from a Large Population Study. *Clinical Medicine & Research*, *14*(3-4), 138–144. https://doi.org/10.3121/cmr.2016.1338

Feld, G., Bernard, M., Rawson, A. R., & Spiers, H. J. (2022). Sleep targets highly connected global and local nodes to aid consolidation of learned graph networks. *Scientific Reports*, *12*(1). https://doi.org/10.1038/s41598-022-17747-2

Ferrara, M., Iaria, G., De Gennaro, L., Guariglia, C., Curcio, G., Tempesta, D., & Bertini, M. (2006). The role of sleep in the consolidation of route learning in humans: A behavioural study. *Brain Research Bulletin*, *71*(1-3), 4–9. https://doi.org/10.1016/j.brainresbull.2006.07.015

Ferrara, M., Iaria, G., Tempesta, D., Curcio, G., Moroni, F., Marzano, C., De Gennaro, L., & Pacitti, C. (2008). Sleep to find your way: The role of sleep in the consolidation of memory for navigation in humans. *Hippocampus*, *18*(8), 844–851. https://doi.org/10.1002/hipo.20444

Fjell, A. M., Øystein Sørensen, Wang, Y., Amlien, I. K., William, Bartrés-Faz, D., Bertram, L., Carl-Johan Boraxbekk, Brandmaier, A. M., Demuth, I., Drevon, C. A., Ebmeier, K. P., Paolo Ghisletta, Kievit, R. A., Kühn, S., Kathrine Skak Madsen, Mowinckel, A. M., Nyberg, L., Sexton, C. E., & Solé-Padullés, C. (2022). Sleep duration and brain structure – phenotypic associations and genotypic covariance. *BioRxiv (Cold Spring Harbor Laboratory)*. https://doi.org/10.1101/2022.02.15.480501

Fjell, A. M., Sørensen, Ø., Amlien, I. K., Bartrés-Faz, D., Brandmaier, A. M., Buchmann, N., Demuth, I., Drevon, C. A., Düzel, S., Ebmeier, K. P., Ghisletta, P., Idland, A.-V., Kietzmann, T. C., Kievit, R. A., Kühn, S., Lindenberger, U., Magnussen, F., Macià, D., Mowinckel, A. M., & Nyberg, L. (2020). Poor Self-Reported Sleep is Related to Regional Cortical Thinning in Aging but not Memory Decline—Results From the Lifebrain Consortium. *Cerebral Cortex*, *31*(4), 1953–1969. https://doi.org/10.1093/cercor/bhaa332

Fjell, A.M., Øystein Sørensen, Wang, Y., Amlien, I.K., William F.C. Baaré, Bartrés-Faz, D., Carl-Johan Boraxbekk, Brandmaier, A.M., Demuth, I., Drevon, C.A., Ebmeier, K.P., Paolo

Ghisletta, Kievit, R., Kühn, S., Kathrine Skak Madsen, Nyberg, L., Solé-Padullés, C., Didac Vidal-Piñeiro, Wagner, G. and Leiv Otto Watne (2023). Is short sleep bad for the brain? Brain structure and cognitive function in short sleepers [Preprint]. p.JN-22. doi:https://doi.org/10.1523/jneurosci.2330-22.2023.

Grandner, M. A., Schopfer, E. A., Sands-Lincoln, M., Jackson, N., & Malhotra, A. (2015). Relationship between sleep duration and body mass index depends on age. *Obesity*, *23*(12), 2491–2498. https://doi.org/10.1002/oby.21247

He, Q., Beveridge, E. H., Starnes, J., Goodroe, S. C., & Brown, T. I. (2021). Environmental overlap and individual encoding strategy modulate memory interference in spatial navigation. *Cognition*, *207*, 104508. https://doi.org/10.1016/j.cognition.2020.104508

Hill, P. F., McAvan, A. S., Garren, J. D., Grilli, M. D., Barnes, C. A., & Ekstrom, A. D. (2023). *Age differences in spatial memory are mitigated during naturalistic navigation.* https://doi.org/10.1101/2023.01.23.525279

Hirshkowitz, M., Whiton, K., Albert, S. M., Alessi, C., Bruni, O., DonCarlos, L., Hazen, N., Herman, J., Katz, E. S., Kheirandish-Gozal, L., Neubauer, D. N., O'Donnell, A. E., Ohayon, M., Peever, J., Rawding, R., Sachdeva, R. C., Setters, B., Vitiello, M. V., Ware, J. C., & Adams Hillard, P. J. (2015). National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*, *1*(1), 40–43. https://doi.org/10.1016/j.sleh.2014.12.010

Howard, Lorelei R., Javadi, A., Yu, Y., Mill, Ravi D., Morrison, Laura C., Knight, R., Loftus, Michelle M., Staskute, L., & Spiers, Hugo J. (2014). The Hippocampus and Entorhinal Cortex Encode the Path and Euclidean Distances to Goals during Navigation. *Current Biology*, *24*(12), 1331–1340. https://doi.org/10.1016/j.cub.2014.05.001

Jackowska, M., & Cadar, D. (2020). The mediating role of low-grade inflammation on the prospective association between sleep and cognitive function in older men and women: 8-year

follow-up from the English Longitudinal Study of Ageing. *Archives of Gerontology and Geriatrics*, *87*, 103967. https://doi.org/10.1016/j.archger.2019.103967

Javadi, A.-H., Patai, E. Z., Marin-Garcia, E., Margolis, A., Tan, H.-R. M., Kumaran, D., Nardini, M., Penny, W., Duzel, E., Dayan, P., & Spiers, H. J. (2019). Prefrontal Dynamics Associated with Efficient Detours and Shortcuts: A Combined Functional Magnetic Resonance Imaging and Magnetoencenphalography Study. *Journal of Cognitive Neuroscience*, *31*(8), 1227–1247. https://doi.org/10.1162/jocn_a_01414

Joza, S., Hu, M. T., Jung, K.-Y., Kunz, D., Stefani, A., Petr Dušek, Terzaghi, M., Arnaldi, D., Aleksandar Videnovic, Schiess, M. C., Hermann, W., Jee Young Lee, Ferini-Strambi, L., Lewis, S. J. G., Laurène Leclair-Visonneau, Oertel, W. H., Antelmi, E., Friederike Sixel-Döring, De, C., & Liguori, C. (2023). Progression of clinical markers in prodromal Parkinson's disease and dementia with Lewy bodies: a multicentre study. *Brain*. https://doi.org/10.1093/brain/awad072

Knutson, K. L., & von Schantz, M. (2018). Associations between chronotype, morbidity and mortality in the UK Biobank cohort. *Chronobiology International*, *35*(8), 1–9. https://doi.org/10.1080/07420528.2018.1454458

Kyle, S. D., Sexton, C. E., Feige, B., Luik, A. I., Lane, J., Saxena, R., Anderson, S. G., Bechtold, D. A., Dixon, W., Little, M. A., Ray, D., Riemann, D., Espie, C. A., Rutter, M. K., & Spiegelhalder, K. (2017). Sleep and cognitive performance: cross-sectional associations in the UK Biobank. *Sleep Medicine*, *38*, 85–91. https://doi.org/10.1016/j.sleep.2017.07.001

Li, Y., Sahakian, B. J., Kang, J., Langley, C., Zhang, W., Xie, C., Xiang, S., Yu, J., Cheng, W., & Feng, J. (2022). The brain structure and genetic mechanisms underlying the nonlinear association between sleep duration, cognition and mental health. *Nature Aging*, *2*(5), 425–437. https://doi.org/10.1038/s43587-022-00210-2

Lo, J., Cheng, G., Loh, K., Leong, R., & Chee, M. (2015). Associations between self-reported sleep duration and cognitive performance in older adults: A systematic review and meta-analysis. *Sleep Medicine*, *16*, S31–S32. https://doi.org/10.1016/j.sleep.2015.02.078

Lucey, B. P., Wisch, J., Boerwinkle, A. H., Landsness, E. C., Toedebusch, C. D., McLeland, J. S., Butt, O. H., Hassenstab, J., Morris, J. C., Ances, B. M., & Holtzman, D. M. (2021). Sleep and longitudinal cognitive performance in preclinical and early symptomatic Alzheimer's disease. *Brain*, *144*(9), 2852–2862. https://doi.org/10.1093/brain/awab272

Ma, Y., Liang, L., Zheng, F., Shi, L., Zhong, B., & Xie, W. (2020). Association Between Sleep Duration and Cognitive Decline. *JAMA Network Open*, *3*(9), e2013573.

https://doi.org/10.1001/jamanetworkopen.2020.13573

Maybrier, H. R., Ben J.A. Palanca, & Head, D. (2022). Associations of environmental and lifestyle factors with spatial navigation in younger and older adults. *Journal of the International Neuropsychological Society*, 1–11. https://doi.org/10.1017/s1355617722000303

Mohlenhoff, B. S., Insel, P. S., Mackin, R. S., Neylan, T. C., Flenniken, D., Nosheny, R., Richards, A., Maruff, P., & Weiner, M. W. (2018). Total Sleep Time Interacts With Age to Predict Cognitive Performance Among Adults. *Journal of Clinical Sleep Medicine*, *14*(09), 1587–1594. https://doi.org/10.5664/jcsm.7342

Mortaza Afshani, Mahmoodi-Aznaveh, A., Noori, K., Masoumeh Rostampour, Mojtaba Zarei, Spiegelhalder, K., Habibolah Khazaie, & Tahmasian, M. (2023). Discriminating Paradoxical and Psychophysiological Insomnia Based on Structural and Functional Brain Images: A Preliminary Machine Learning Study. *Brain Sci.* , *13*(4), 672–672. https://doi.org/10.3390/brainsci13040672

Nazareth, A., Huang, X., Voyer, D., & Newcombe, N. (2019). A meta-analysis of sex differences in human navigation skills. *Psychonomic Bulletin & Review*, *26*(5), 1503–1528. https://doi.org/10.3758/s13423-019-01633-6

Newcombe, N. S., Hegarty, M., & Uttal, D. (2022). Building a Cognitive Science of Human Variation: Individual Differences in Spatial Navigation. *Topics in Cognitive Science*, *15*(1), 6–14. https://doi.org/10.1111/tops.12626

Nguyen, N. D., Tucker, M. A., Stickgold, R., & Wamsley, E. J. (2013). Overnight Sleep Enhances Hippocampus-Dependent Aspects of Spatial Memory. *Sleep*, *36*(7), 1051–1057. https://doi.org/10.5665/sleep.2808

Nuñez, A. N., Rhee, J. U., Haynes, P., Chakravorty, S., Patterson, F., Killgore, W. D. S., Gallagher, R. A., Hale, L., Branas, C., Carrazco, N., Alfonso-Miller, P., Gehrels, J.-A., & Grandner, M. A. (2020). Smoke at night and sleep worse? The associations between cigarette smoking with insomnia severity and sleep duration. *Sleep Health*, *7*(2). https://doi.org/10.1016/j.sleh.2020.10.006

Orban, P., Rauchs, G., Balteau, E., Degueldre, C., Luxen, A., Maquet, P., & Peigneux, P. (2006). Sleep after spatial learning promotes covert reorganization of brain activity. *Proceedings of the National Academy of Sciences*, *103*(18), 7124–7129. https://doi.org/10.1073/pnas.0510198103

Patai, E. Z., & Spiers, H. J. (2021). The Versatile Wayfinder: Prefrontal Contributions to Spatial Navigation. *Trends in Cognitive Sciences*, *25*(6), 520–533. https://doi.org/10.1016/j.tics.2021.02.010

Peigneux, P., Laureys, S., Fuchs, S., Collette, F., Perrin, F., Reggers, J., Phillips, C., Degueldre, C., Del Fiore, G., Aerts, J., Luxen, A., & Maquet, P. (2004). Are Spatial Memories Strengthened in the Human Hippocampus during Slow Wave Sleep? *Neuron*, *44*(3), 535–545. https://doi.org/10.1016/j.neuron.2004.10.007

Puthusseryppady, V., Morrissey, S., Spiers, H., Patel, M., & Hornberger, M. (2022). Predicting real world spatial disorientation in Alzheimer's disease patients using virtual reality navigation tests. *Scientific Reports*, *12*(1). https://doi.org/10.1038/s41598-022-17634-w

Santhi, N., Lazar, A.S., McCabe, P.J., Lo, J.C., Groeger, J.A. and Dijk, D.-J. (2016). Sex differences in the circadian regulation of sleep and waking cognition in humans. *Proceedings of the National Academy of Sciences*, 113(19), pp.E2730–E2739. doi:https://doi.org/10.1073/pnas.1521637113.

Seton, C., Coutrot, A., Hornberger, M., Spiers, H. J., Knight, R., & Whyatt, C. (2023). Wayfinding and path integration deficits detected using a virtual reality mobile app in patients with traumatic brain injury. *PLOS ONE* , *18*(3), e0282255–e0282255. https://doi.org/10.1371/journal.pone.0282255

Shattuck, N. L., & Matsangas, P. (2017). Sunlight Exposure, Work Hours, Caffeine Consumption, and Sleep Duration in the Naval Environment. *Aerospace Medicine and Human Performance*, *88*(6), 579–585. https://doi.org/10.3357/amhp.4721.2017

Simon, K. C., Clemenson, G. D., Zhang, J., Sattari, N., Shuster, A. E., Clayton, B., Alzueta, E., Dulai, T., de Zambotti, M., Stark, C., Baker, F. C., & Mednick, S. C. (2022). Sleep facilitates spatial memory but not navigation using the Minecraft Memory and Navigation task. *Proceedings of the National Academy of Sciences*, *119*(43). https://doi.org/10.1073/pnas.2202394119

Spiers, H. J., Coutrot, A., & Hornberger, M. (2023). Explaining World-Wide Variation in Navigation Ability from Millions of People: Citizen Science Project Sea Hero Quest. *Topics in Cognitive Science*. https://doi.org/10.1111/tops.12590

Su, S., Li, X., Xu, Y., McCall, W. V., & Wang, X. (2022). Epidemiology of accelerometer-based sleep parameters in US school-aged children and adults: NHANES 2011–2014. *Scientific Reports*, *12*(1). https://doi.org/10.1038/s41598-022-11848-8

Tai, X. Y., Chen, C., Manohar, S., & Husain, M. (2022). Impact of sleep duration on executive function and brain structure. *Communications Biology*, *5*(1). https://doi.org/10.1038/s42003-022-03123-3

Tang, J., Liao, Y., Kelly, B. C., Xie, L., Xiang, Y.-T., Qi, C., Pan, C., Hao, W., Liu, T., Zhang, F., & Chen, X. (2017). Gender and Regional Differences in Sleep Quality and Insomnia: A General Population-based Study in Hunan Province of China. *Scientific Reports*, *7*. https://doi.org/10.1038/srep43690

Valera, S., Guadagni, V., Slone, E., Burles, F., Ferrara, M., Campbell, T., & Iaria, G. (2016). Poor sleep quality affects spatial orientation in virtual environments. *Sleep Science*, *9*(3), 225–231. https://doi.org/10.1016/j.slsci.2016.10.005

Vallat, R., Berry, S. E., Tsereteli, N., Capdevila, J., Khatib, H. A., Valdes, A. M., Delahanty, L. M., Drew, D. A., Chan, A. T., Wolf, J., Franks, P. W., Spector, T. D., & Walker, M. P. (2022). How people wake up is associated with previous night's sleep together with physical activity and food intake. *Nature Communications*, *13*(1), 7116. https://doi.org/10.1038/s41467-022-34503-2

Van Dongen, H. P. A., Maislin, G., Mullington, J. M., & Dinges, D. F. (2003). The Cumulative Cost of Additional Wakefulness: Dose-Response Effects on Neurobehavioral Functions and Sleep Physiology From Chronic Sleep Restriction and Total Sleep Deprivation. *Sleep*, *26*(2), 117–126. https://doi.org/10.1093/sleep/26.2.117

van Oostrom, S. H. , Nooyens, A. C., J., van Boxtel, M. P., J., & Monique Verschuren, W., M.,. (2018). Long sleep duration is associated with lower cognitive function among middle-age adults – the Doetinchem Cohort Study. *Sleep Medicine*, *41*, 78–85. https://doi.org/10.1016/j.sleep.2017.07.029

Varga, A. W., Ducca, E. L., Kishi, A., Fischer, E., Parekh, A., Koushyk, V., Yau, P. L., Gumb, T., Leibert, D. P., Wohlleber, M. E., Burschtin, O. E., Convit, A., Rapoport, D. M., Osorio, R. S., & Ayappa, I. (2016). Effects of aging on slow-wave sleep dynamics and human spatial navigational memory consolidation. *Neurobiology of Aging*, *42*, 142–149. https://doi.org/10.1016/j.neurobiolaging.2016.03.008

Walkowiak, S., A. Coutrot, Hegarty, M., Velasco, P. F., Wiener, J. M., Dalton, R. C., C. Hölscher, Hornberger, M., Spiers, H. J., & Manley, E. (2023). Cultural determinants of the gap between self-estimated navigation ability and wayfinding performance: evidence from 46 countries. *Scientific Reports*, *13*(1). https://doi.org/10.1038/s41598-023-30937-w

Webster, L., Costafreda, S. G., Powell, K., & Livingston, G. (2022). How do care home staff use non-pharmacological strategies to manage sleep disturbances in residents with dementia: The

SIESTA qualitative study. *PLOS ONE*, *17*(8), e0272814.

https://doi.org/10.1371/journal.pone.0272814

Weisberg, S. M., & Newcombe, N. S. (2016). How do (some) people make a cognitive map?

Routes, places, and working memory. *Journal of Experimental Psychology: Learning, Memory,*

and Cognition, *42*(5), 768–785. https://doi.org/10.1037/xlm0000200

Weisberg, S. M., Schinazi, V. R., Newcombe, N. S., Shipley, T. F., & Epstein, R. A. (2014).

Variations in cognitive maps: Understanding individual differences in navigation. *Journal of*

Experimental Psychology: Learning, Memory, and Cognition, *40*(3), 669–682.

https://doi.org/10.1037/a0035261

West, G. L., Patai, Z. E., Coutrot, A., Hornberger, M., Bohbot, V. D., & Spiers, H. J. (2023).

Landmark-dependent Navigation Strategy Declines across the Human Life-Span: Evidence from

Over 37,000 Participants. *Journal of Cognitive Neuroscience*, *35*(3), 452–467.

https://doi.org/10.1162/jocn_a_01956

Whitmore, N., & Paller, K. A. (2022). Sleep Disruption Selectively Weakens Reactivated

Memories. *Learn. Mem.* https://doi.org/10.1101/2022.04.10.487316

Willoughby, A. R., Iman Alikhani, Karsikas, M., Xin Yu Chua, & Michael W.L. Chee. (2023).

Cross-Country Differences in Nocturnal Sleep Duration and Variability Across the Week.

[Preprint]. https://doi.org/10.31234/osf.io/8ahsu

Wolbers, T., & Hegarty, M. (2010). What determines our navigational abilities? *Trends in*

Cognitive Sciences, *14*(3), 138–146. https://doi.org/10.1016/j.tics.2010.01.001

Zheng, D., Yuan, X., Ma, C., Liu, Y., VanEvery, H., Sun, Y., Wu, S., & Gao, X. (2020). Alcohol

consumption and sleep quality: a community-based study. *Public Health Nutrition*, *24*(15), 1–23.

https://doi.org/10.1017/S1368980020004553