**Spatial molecular profiling of mixed invasive ductal-lobular breast cancers reveals**

**heterogeneity in intrinsic molecular subtypes, oncogenic signatures, and mutations**

Osama Shiraz Shah[1,3], Azadeh Nasrazadani[1,4], Julia Foldi[7], Jennifer M. Atkinson[1,2], Celina G Kleer[8],

Priscilla F. McAuliffe[1,6,] Tyler J Johnston[10], Wayne Stallaert[10],  Edaise M da Silva[9], Pier Selenica[9],

Higinio Dopeso[9], Fresia Pareja[9], Diana Mandelker[9], Britta Weigelt[9], Jorge S. Reis-Filho[9$], Rohit

Bhargava[4], Peter C. Lucas[5], Adrian V. Lee*[1,2], Steffi Oesterreich*[1,2]

[1]Womens Cancer Research Center at UPMC Hillman Cancer Center and Magee Women's

Research Institute, Pittsburgh, PA.

[2]Department of Pharmacology & Chemical Biology, University of Pittsburgh, Pittsburgh, PA.

[3]Integrative Systems Biology Program, University of Pittsburgh, Pittsburgh PA.

[4]Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, PA

[5]Department of Laboratory Medicine and Pathology, Mayo Clinic College of Medicine and

Science

[6]Division of Surgical Oncology, Department of Surgery, University of Pittsburgh School of

Medicine, Pittsburgh, PA

[7]Division of Hematology-Oncology, Department of Medicine, University of Pittsburgh,

Pittsburgh, PA.

[8]Department of Pathology and Rogel Cancer Center, University of Michigan Ann Arbor, MI

48109, USA

[9]Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center

(MSKCC), New York, NY, USA

[10]Department of Computational and Systems Biology, University of Pittsburgh School of

Medicine, Pittsburgh, PA

[$]Current affiliation: AstraZeneca

**Keywords:** Mixed invasive ductal and lobular carcinoma, mDLC, digital spatial profiling, DSP,

breast cancer, tumor heterogeneity, multi-omics, precision medicine

* Steffi Oesterreich and Adrian Lee share senior authorship

**Corresponding Authors:** Steffi Oesterreich, University of Pittsburgh, 5051 Centre Ave,

Pittsburgh, PA 15213. Phone: 412-641-8555, Fax: 412-641-7724, E-mail:

oesterreichs@upmc.edu.  Adrian Lee, University of Pittsburgh, 5051 Centre Ave, Pittsburgh, PA

15213. E-mail: leeav@upmc.edu.

**Abstract**

Mixed invasive ductal and lobular carcinoma (MDLC) is a rare histologic subtype of breast cancer displaying both E-cadherin positive ductal and E-cadherin negative lobular morphologies within the same tumor, posing challenges with regard to anticipated clinical management. It remains unclear whether these distinct morphologies also have distinct biology and risk of recurrence. Our spatially-resolved transcriptomic, genomic, and single-cell profiling revealed clinically significant differences between ductal and lobular tumor regions including distinct intrinsic subtype heterogeneity (e.g., MDLC with TNBC/basal ductal and ER+/luminal lobular regions), distinct enrichment of senescence/dormancy and oncogenic (ER and MYC) signatures, genetic and epigenetic *CDH1* inactivation in lobular, but not ductal regions, and single-cell ductal and lobular sub-populations with unique oncogenic signatures further highlighting intra-regional heterogeneity. Altogether, we demonstrated that the intra-tumoral morphological/histological heterogeneity within MDLC is underpinned by intrinsic subtype and oncogenic heterogeneity which may result in prognostic uncertainty and therapeutic dilemma.

## Significance

MDLC displays both ductal and lobular tumor regions. Our multi-omic profiling approach revealed that these morphologically distinct tumor regions harbor distinct intrinsic subtypes and oncogenic features that may cause prognostic uncertainty and therapeutic dilemma. Thus histopathological/molecular profiling of individual tumor regions may guide clinical decision making and benefit patients with MDLC, particularly in the advanced setting where there is increased reliance on next generation sequencing.

**Main Text**

**Introduction**

Breast cancer is a prevalent disease worldwide[1] and is the most frequently diagnosed cancer in

women in the US, with approximately 300,000 cases per year[2]. The majority (approximately

75%) of cases are classified histologically as invasive breast carcinoma of no special type (IBC-

NST), previously described as invasive ductal cancer (IDC). Invasive lobular carcinoma (ILC) is the

most common special type accounting for approximately 10-15% of all invasive breast cancer

diagnoses[3,4]. Apart from the well-defined histologic and tissue histological differences, there

are numerous notable molecular distinctions between ILC and IBC-NST. ILC exhibits enrichment

of *CDH1*, *FOXA1*, *TBX3*, *ERBB2* and *PTEN* alterations, while IBC-NST shows a higher frequency of

*GATA3*, *MAP3K1*, *TP53* and *MYC* alterations[5]. As a result of *CDH1* alterations, ILC lacks E-

cadherin expression, an important cell adhesion molecule[6], and display cytoplasmic p120

staining while IBC-NST retains membranous expression for both molecules[7]. Clinically, ILC is

more common in elderly women, presents as larger tumors and is frequently multicentric and

multifocal compared to IBC-NST[8,9]. ILC is usually detected later and leads to higher rates of

mastectomy [10–14]. ILC is also associated with more frequent metastases to urogenital and

gastrointestinal tracts as well as to serosal surfaces than IBC-NST[15,16]. Despite presenting with

markers of favorable prognosis, such as low grade, low proliferation index, estrogen receptor

positive and human epidermal growth factor receptor (HER2) low/negative, patients with ILC

have a higher rate of late recurrences and worse long-term prognosis than those with IBC-

NST[14,17,18]. Given its distinct prognosis and outcome compared to IBC-NST, ILC is increasingly

being recognized as a distinct disease, yet it is still treated similarly to IBC-NST in the clinic due

to a lack of specific treatment guidelines[14,19,20]. Ongoing clinical trials focused on patients with

ILC may lead to the development of more tailored and effective treatment options in the

future[20].

Apart from ILC, there are numerous other special types of breast cancers such as mixed invasive

breast carcinoma. This is an elusive group of breast cancers which the World Health

Organization (WHO) defines as tumors presenting with areas of 10-90% of special histology

(such as ILC) admixed with at least 10% of IBC-NST[21]. Importantly, this current WHO definition

of mixed invasive breast cancers MDLC differs considerably from its previous definition i.e.,

tumors presenting with areas of 10-49% of the special subtype and at least 50% IBC-NST[22],

indicating the fluid interpretation of this disease. Mixed tumors comprising of both ILC and IBC-

NST are the most common (~5-10% of all breast cancer cases[23–28]) and have different

terminologies in the literature including mixed invasive ductal and lobular breast carcinoma

(MDLC)[29] (used throughout this study), mixed invasive ductolobular breast cancer (MIDLC)[30] ),

invasive ductolobular carcinomas (IDLC)[31], and more. Despite increasing appreciation of MDLC

disease[32,33], it remains understudied, leading to diagnostic and prognostic uncertainty and

clinical management challenges for the patients with this elusive disease. Although there are

limited reports focused on MDLC, a few seminal studies have provided insights into its complex

pathobiology. Immunohistochemical (IHC) studies have confirmed while that ductal regions of

MDLC retain membranous E-cadherin, while lobular regions either lose or display aberrant E-

cadherin IHC staining[34–36]. Notably, the current MDLC diagnosis guidelines from WHO only

consider the presence of mixed histological patterns and lack recommendations for support

from E-cadherin IHC. However, a recent study suggests that the use of E-cadherin IHC can

improve diagnostic accuracy of lobular tumor components in ILC and MDLC tumors[37]. Clinical

studies have often yielded contradictory findings[24–26,33,34,36,38,39], but overall, they suggest that

MDLC is clinically more similar to pure ILC than IBC-NST[29,40,41 29,40]. These discrepancies may be

due to the non-standardized and fluid definition of MDLC disease, limited consideration of its

underlying tumor regions and their distinct features and proportions, and the vast intratumoral

heterogeneity of this disease[29,33,42,43]. Findings from previous reports point to several subtypes

of MDLC tumors based on their histological and E-cadherin staining patterns: 1)

intermixed/intermingled type where the E-cadherin positive ductal and E-cadherin

negative/aberrant lobular tumor cells appear closely intermixed/intermingled together[31,44], 2)

collision type where the E-cadherin positive ductal and E-cadherin negative/aberrant lobular

tumor regions appear in spatially distinct regions which may collide into each other forming

intermixed regions[31,45], and 3) lobular-like IBC-NST type where there are histologically distinct

ductal and lobular tumor regions but the overall tumor is predominately E-cadherin positive[46].

Moreover, molecular studies with profiling of individual tumor regions have been limited. The

largest molecular study on MDLC tumors was from the TCGA group[47], where 88 MDLC were

classified into lobular-like and ductal-like groups based on transcriptomic and mutational

similarity to ILC and IBC-NST tumors. The lobular-like group comprised of ~25% of MDLC cases

and showed enrichment of mutations in and lower expression of *CDH1* gene. However, these

findings may not have any biological relevance as these samples lacked comprehensive

annotation and may have been incorrectly defined as MDLC as well as the study lacked

individual tumor region analysis. Additional molecular studies have largely focused on analysis

of somatic genetic alterations[44,45,48]. While these have provided valuable insights into the

tumorigenesis pathways and distinct mutational landscape of ductal and lobular tumor

regions[36,49], comprehensive studies delineating transcriptomic differences between these

distinct regions have not been performed.

To fill this knowledge gap, we employed spatially resolved transcriptomic, genomic, and single

cell profiling approaches to comprehensively investigate MDLC and highlight its intratumoral

heterogeneity and complexity. We delineate the biological and intrinsic subtype differences

between its underlying ductal and lobular tumor regions. Notably, our findings highlight that in

some MDLC cases distinct tumor regions with distinct biology may pose distinct progression

risk. Hence histopathological and molecular profiling MDLC and its individual tumor regions

may inform treatment decisions.

## Results

**Patient Specimens and Their Clinicopathologic Characteristics**

A panel of three molecular pathologists identified invasive breast cancer cases diagnosed as "mixed invasive ductal and lobular carcinoma" and shortlisted three cases for this pilot study. All three cases were reported as estrogen receptor positive (ER+) and HER2 negative/low and their ages ranged from 46-63 years (Fig. 1A, Supplementary Table S1A). Assessment of H&E and E-cadherin/p120 dual staining confirmed that each case exhibited histomorphic features in line with the latest MDLC definition i.e., admix of E-cadherin negative lobular tumor regions with at least 10-90% of E-cadherin positive ductal tumor regions[21] (Supplementary Table S1B). Importantly, overall, these cases exhibited collision of spatially distinct ductal and lobular tumor regions[31,45] along with some regions of intermixing and hence could be broadly defined as MDLC of collision type. These cases were particularly amenable for downstream multi-omic profiling compared to those that exhibited more intermingling or intermixing with close proximity of histologic tumor regions. Clinical diagnostic and treatment timelines of these cases, source of their tumor tissue used in this study, and their histomorphic features are illustrated in Fig. 1A. Both MDLC-2 and MDLC-3 eventually developed recurrent disease resulting in patient death in the case of MDLC-2. As of the last follow-up in 2022, in both MDLC-3 and MDLC-1 cases, the patients were alive with tumors, and in the case of MDLC-1 the patient was free of recurrent disease.

**Whole-Transcriptome Spatial Profiling of Ductal and Lobular Regions**

To better understand the biological heterogeneity and transcriptomic features of ductal and

lobular regions, we performed whole-transcriptome digital spatial profiling (DSP) of these three

MDLC cases. In total, we analyzed 26 regions of interest (ROIs) with at least 6 ROIs per case.

Selected ROIs from a representative case (MDLC-3) are shown in Fig. 1B. We segmented ductal

and lobular areas using E-cadherin and Pan-CK staining (Fig. 1C). This enabled profiling of pure

ductal and lobular areas from heterogeneous regions that were predominantly of one histology

or a mixture of both (Fig. 1D). E-cadherin expression in segmented ductal and lobular regions

showed expected results (Fig. 1E). All profiled regions had high RNA sequencing saturation (>

94%) and low noise (Q3 count > Negative probe count). However, three regions had low read

count (< 0.3 million) and were excluded from further analysis (Supplementary Fig. S1A-C). Our

final DSP dataset is comprised of 13 ductal and 13 lobular regions.

**Transcriptomic Clusters, Intrinsic Molecular Subtypes and Differential Transcriptomic**

**Features**

Exploration using top variable genes revealed that regions from each tumor clustered based

upon their histology (Fig. 2A; Supplementary Fig. S1D). MDLC-3 ductal regions showed the most

distinct mRNA expression compared to all other regions. PAM50 analysis showed both inter and

intra-tumor heterogeneity of prognostically relevant intrinsic molecular subtypes (IMS). MDLC-

2 ductal regions were a mix of Luminal A and HER2-Enriched subtypes while lobular regions

were HER2-Enriched (but stained negative for HER2 IHC – Supplementary Table S1A). MDLC-1

ductal and lobular regions were both of luminal IMS, with the former being Luminal A and the

latter being Luminal B. Finally, in line with mRNA expression patterns, MDLC-3 ductal and

lobular regions showed stark IMS differences (former Luminal A and later Basal-Like). MDLC-3

ductal region was the only TNBC/Basal-Like tumor (and had low *ESR1*, *PGR* and *ERBB2* mRNA

expression – Supplementary Fig. S1E). We further confirmed the TNBC status of MDLC-3 ductal

regions using ER and HER2 IHC (Supplementary Table S1A). Overall, our transcriptomic

clustering and IMS analysis suggest that ductal and lobular tumor regions within MDLC may

have distinct transcriptomes and intrinsic subtypes.

To investigate global transcriptomic differences between ductal and lobular regions across all

cases, we performed differential gene expression analysis in a pan-patient manner. To avoid

biases caused by fundamental differences between ER+ and ER- disease, we excluded MDLC-3

ER-/TNBC ductal regions from this analysis and later downstream analyses (Supplementary Fig.

S2C, Supplementary Table S5). In total, we identified 168 significantly up-regulated and 198

significantly down-regulated differentially expressed genes (DEGs) in the lobular compared to

ductal regions across all cases (Fig. 2B, Supplementary Table S5). These pan-patient DEGs

showed strong overlap with patient specific DEGs (Supplementary Fig. S2A-C) indicating that

our pan-patient analysis captured most if not all strong global differences between ductal and

lobular tumor regions in each case.

To assess transcriptomic similarity of MDLC ductal and lobular regions to pure counterparts,

i.e., IBC-NST (IDC) and ILC, we compared MDLC ductal vs. lobular region DEGs with those from

the comparison of TCGA ER+ IBC-NST (N=327) vs ILC (N=113), respectively (Supplementary Fig

S2D). We found *KLK11, KLK10, NRAP, SHROOM1, BTG2,* and *PDK4* were upregulated, while

*CDH1, DCD, SERPINA1*, and *CPB1* were downregulated in both comparisons (i.e., MDLC ductal vs

lobular tumor regions and IBC-NST vs ILC tumors). This overlap confirmed the well-known

down-regulation of *CDH1* as a marker of the lobular phenotype as well as potential novel

markers.

**Differential Enrichment of ER and MYC Signatures & Recurring Biological Themes**

To understand the distinct biology of ductal and lobular regions, we scored each DSP region for

hallmark signatures[50]. Twenty signatures were differentially enriched between ductal and

lobular regions with p-value ≤ 0.05 (Fig. 3A, Supplementary Table S6). Of note, many of these

pathways can also be found differentially activated when comparing ER+ IBC-NST and ER+ ILC in

TCGA – for example, Kras and TNFA signaling as well as Hypoxia are among significantly

enriched pathways in ER+ ILC TCGA samples, whereas Myc targets are enriched in ER+ IBC-NST

TCGA samples (Fig. 3B). In line with differential enrichment of Estrogen Response and Myc

Target signatures, our transcription factor (TF) signature analysis revealed enrichment of

*ESR1/FOXA* TF signatures in lobular and *MYC* TF signatures in ductal regions (Supplementary Fig.

S3).

To determine recurring biological themes, we expanded our differential enrichment analysis to

include all GO[51], REACTOME[52] and KEGG[53] genesets. Signatures differentially enriched between

ductal and lobular regions with p-value ≤ 0.05 were further assessed for robustness against

background permutations. Robust signatures were grouped into signature clusters based on

similarity of overlapping genes in ductal and lobular regions. All signature clusters were further

grouped into broad biological themes based on the underlying biological terms (Fig. 3C, left

panel). The mean score of all signature clusters in ductal and lobular regions is shown in Fig. 3C,

right panel. Lobular regions were enriched for TP53 & senescence, collagen & extracellular

matrix (ECM), MAPK/ERK targets (in line with KRAS signatures shown in Fig. 3A), while ductal

regions were enriched in double stranded break & homologous repair (DSB/HR), mTOR

signaling (in line with Mtorc1/Mtor signatures shown in Fig. 3A) among other signatures

(Supplementary Table S7).

Notably, several genes including *MYC, G3BP1* and *SNRPD1* overlapped between Myc Targets V1

and G2m Checkpoint signatures (both enriched in ductal regions). Both of these signatures had

no overlapping genes with DNA damage and repair (D1: DBS/HR) signatures (Supplementary

Fig. S4A) indicating that Myc related signatures in ductal regions are associated with

proliferation. Importantly, various senescence and cell death related genes (*CDKN1A/2B/2C,*

*TP53, BCL6* among others) overlapped between the L1: Cell cycle and L2: *TP53* & Senescence

signatures indicating negative regulation of cell cycle via *TP53* signaling and cell cycle inhibitors

(Supplementary Fig. S4B). Key genes associated with these notable and other biological

signatures are highlighted in Supplementary Fig. S4C. In summary, we identified various

differentially enrichment of various oncogenic and biological signatures between ductal vs.

lobular regions including cell cycle (Myc and G2m genes in ductal, and TP53 and cell cycle arrest

genes in lobular), immune, collagen & ECM and metabolism among others.

To further investigate some of the notable distinctions in gene expression signatures between

ductal and lobular tumor regions, we performed multiplexed immunofluorescence to obtain

single-cell proteomic measurements of key cell cycle and senescence effectors in ductal and

lobular tumor cells, identified by staining for pan-cytokeratin 26 and E-cadherin (Fig. 3D,

<mark>Supplementary Table S12). Individual cells within ductal regions exhibited significantly higher</mark>

<mark>expression of proliferative markers including cMYC, cyclin E2, and cyclin A1, while lobular</mark>

<mark>regions had significantly higher expression of the senescence biomarker p16 (Fig. 3E). These</mark>

<mark>findings confirmed the activation of Myc signaling and proliferative pathways in ductal, and</mark>

<mark>activation of cell cycle arrest/senescence in lobular tumor regions.</mark>

**Genomic Landscape and Clinically Relevant Genomic Alterations**

To explore the genomic landscape of ductal and lobular regions, we employed the FDA-

approved targeted MSK-IMPACT panel. We micro-dissected the ductal and lobular tumor

regions from FFPE blocks of each case and sequenced them at a median depth of 416x (range

287–563) (Supplementary Fig. S5A). In all cases, at least one pathogenic mutation was detected

in known breast cancer driver genes, including *CDH1, PIK3CA, TP53, GATA3, KMT2C, MAPK31,*

*ARID1A,* and *CBFB,* in both ductal and lobular regions, as depicted in Fig. 4A. Furthermore, all

pathogenic mutations had a high cancer cell fraction (Supplementary Fig. S5B).

Loss of E-cadherin is an established driver of lobular tumorigenesis[54]. Typically, it results from

biallelic *CDH1* inactivation, through a combination of truncating mutation and loss of

heterozygosity (LOH)[28]. All lobular regions exhibited loss of E-cadherin, with both MDLC-1 and

MDLC-3 harboring *CDH1* truncating mutations and LOH (Fig. 4B; Fig. 1E; Supplementary Fig.

S1E). The lobular region of MDLC-2 did not have *CDH1* genomic alterations but showed

epigenetic inactivation via methylation of its promoter. A putative passenger mutation in *CDH1*

was also observed in the MDLC-3 ductal region, which had no effect on mRNA or E-cadherin

levels (Fig. 1E; Supplementary Fig. S1E). The loss of E-cadherin is consistent with the

downstream down-regulation of Adherens Junction signatures in all lobular regions (Fig. 3A).

Hotspot *PIK3CA* mutations, which are early oncogenic driver events frequently observed in ER+

BC[55], were present in all cases except MDLC-3. The MDLC-1 ductal region had mutations in

*GATA3* (critical transcriptional binding partner of ERalpha[56]) and *CBFB* (forms complex with

*RUNX1* to suppress NOTCH signaling[57]). The MDLC-3 ductal region had loss of *TP53*, an event

associated with more aggressive types of breast cancer like basal/TNBC[58] and consistent with its

basal-like intrinsic subtype. MDLC-2 ductal region had truncating mutation in *KMT2C*, an

important regulator of ER alpha activity[59]. MDLC-3 lobular region had mutations in *ATRX* and

*ARID1A,* two chromatin remodeling and DNA repair genes[60]. In addition, while MDLC-1 and

MDLC-3 ductal and lobular regions did not share any somatic clonal mutations, MDLC-2 ductal

and lobular regions shared several somatic clonal mutations in *PIK3CA* and *MAPK31* genes

indicating shared tumor origin.

To investigate the copy number (CN) landscape of ductal and lobular regions, we performed CN

analysis using FACETS[61] (Fig. 4B). All regions exhibited breast cancer characteristic chromosome

1q gains and 16q losses[58]. Overall, ductal tumor regions showed more frequent aneuploidy

events than lobular tumor regions. MDLC-3 ductal region showed the most aberrant CN profile

consistent with its basal/TNBC etiology[58]. Moreover, chromosome 22 aneuploidy was only

observed in ductal tumor regions. Notably, both histologic regions of MDLC-3 and ductal region

of MDLC-2 had *NOTCH2* gains or amplifications (Supplementary Fig. S5C).

**Mutational Processes Driving Tumor Evolution**

To determine the mutational processes underlying the evolution of ductal and lobular tumor

regions within MDLC, we performed mutational signature analysis of all cases (Fig. 4C). Both

histologic regions in MDLC-2 were associated with an "aging" signature. In MDLC-3, ductal

region was associated with homologous repair deficiency (HRD) while the lobular regions with

APOBEC signature. Tumors with an APOBEC signatures have a higher frequency of mutations in

chromatin remodeling and DNA repair genes[62], as seen in MDLC-3 lobular region. We did not

identify any signatures for MDLC-1 ductal and lobular regions. The putative evolution of each

MDLC case and its histologic region based upon the mutations, *CDH1* alterations and

mutational signatures is shown in Fig. 4D. Notably, the MDLC-2 ductal and lobular regions not

only shared clonal pathogenic (*PIK3CA*) and passenger (*MST1R* and *MAP3K1*) mutations but also

the "aging" mutational signature. Loss of function *CDH1* alterations were hallmark of all lobular

regions across cases.

**Single-Cell RNA-Sequencing and Identification of Ductal and Lobular Populations**

To explore the heterogeneity of MDLC at a single-cell level, we performed scRNAseq

sequencing of a fresh tumor specimen from MDLC-1. The resulting scRNAseq comprised of

4,671 cells and had sequencing saturation, reads in cell and median genes per cell as shown in

Supplementary Fig. S6A. Cell-type annotation based on canonical markers (see details in

Supplementary Methods) revealed various tumor and non-tumor cell types (Supplementary Fig.

S7B). scRNAseq inferred copy number revealed that only the epithelial tumor cells had

aneuploid copy number status further confirming their nature (Supplementary Fig. S7C). For the

downstream analyses, we only focused on the tumor cell cluster. Clustering of this revealed two

transcriptionally distinct tumor cell populations with similar cell cycle profiles; however, we

were unable to annotate them as either ductal or lobular tumors using *CDH1* mRNA expression

(Fig. 5A). Notably, while the MDLC-1 lobular regions showed loss of E-cadherin protein (Fig. 1E),

the *CDH1* p.Q23* truncating mutation in these regions (Fig. 4A) didn't alter *CDH1* mRNA

expression (Supplementary Fig. S1E). Previous studies suggest that some protein truncating

mutations can escape non-sense mediated decay and thereby maintain gene expression levels

but lack a functional protein[63–65]. Indeed, similar effects of Q23 mutations were seen in the

TCGA cohort, where *CDH1* mRNA levels in Q23* mutant tumors were similar to those in *CDH1*

WT tumors, while the E-cadherin protein levels in Q23* mutant tumors were similar to the

tumors with all other *CDH1* truncating mutations (Supplementary Fig. S6).[63–65]

To identify ductal and lobular single populations, we derived ductal and lobular transcriptome

signatures (Supplementary Table S5) from differential gene expression analysis of the MDLC-1

DSP ductal vs lobular tumor regions and used these signatures to score each single cell in

MDLC-1 scRNAseq dataset. Using this deconvolution approach, we successfully annotated

ductal and lobular tumor populations in the scRNAseq dataset (Fig. 5B). scRNAseq inferred copy

number profiles of ductal and lobular cells revealed various breast cancer specific

aberrant/aneuploid events in chromosomes 1, 8, 11, 16, 18 among others (Supplementary Fig.

S7E). Overall, the copy number profiles of ductal tumor cells were more aberrant than those of

lobular tumor cells. Ductal and lobular tumor regions DEGs in scRNAseq and DSP datasets

showed significant overlap (Supplementary Fig. S7D,F). Differential enrichment of several

hallmark signatures and recurring biological themes in ductal and lobular regions from DSP data

was recapitulated in our scRNAseq data (Supplementary Fig. S8A,C). Notably, we saw

enrichment of Estrogen and senescence signatures in lobular and Myc signature in ductal tumor

cells in both DSP and scRNAseq datasets.

**Single-Cell Heterogeneity and Mapping of Single-Cells to Spatial Regions**

To investigate single-cell heterogeneity of ductal and lobular populations, we increased

resolution of clustering to identify sub-clusters. Each sub-cluster had unique marker genes (Fig.

5C) and several of these also showed differential enrichment of hallmark signatures and

recurring biological theme clusters (Supplementary Fig. S8B,D), indicating intra-tumor

heterogeneity within ductal and lobular tumor populations. To investigate whether single-cell

transcriptomic heterogeneity was associated with spatial heterogeneity, we mapped scRNAseq

sub-clusters to DSP regions. To assess similarity between each region and sub-cluster, we

scored all regions for sub-cluster marker genesets using GSVA (Fig. 5D). Each region was

mapped to a sub-cluster for which it had most similarity (i.e., highest GSVA score) (Fig. 5E). The

top five marker genes of each scRNAseq sub-cluster (Supplementary Fig. S7G) are shown beside

each sub-cluster and spatial region link. These findings perhaps suggest that the single-cells in

each sub-cluster came from a spatially distinct niche in the tumor.

## Discussion

Despite an increasing incidence of MDLC[32,33], this special subtype of breast cancer remains poorly understood, with limited knowledge about its overall biology and that of its underlying ductal and lobular tumor regions, as well as their clinical implications.

Spatially resolved and single-cell profiling techniques have revolutionized our understanding of physiological and pathological processes and tissue architecture[66]. Efforts such as Human Cell Atlas[67] have successfully uncovered the major cell types and their spatial organization in both normal breast tissue[68,69] and breast cancer[70]. Spatial and single-cell profiling of primary breast cancer subtypes and metastatic disease have been instrumental in identifying novel molecular features and their prognostic value[71–75]. However, to date genome-wide spatial and single-cell investigation of MDLC has not been reported. To address this gap, we employed spatially resolved transcriptomics, genomics, and single-cell profiling to investigate the biology of individual tumor regions within MDLC of collision type. Our study confirmed various previously reported observations of genetic inactivation of *CDH1* and concomitant loss of E-cadherin in lobular regions[31,34–36] and the presence of distinct driver alterations in individual tumor regions[31,36]. We also uncovered novel findings that highlight the vast heterogeneity within MDLC, several of which may be indicative of distinct prognosis, hence highlighting the significance of reporting this heterogeneity in clinical diagnosis. Ductal and lobular tumor regions had distinct biological signatures, in particular cell cycle arrest/senescence, ER and *MYC* signaling, and intrinsic molecular subtypes (IMS). The IMS distinctions were most prominent in the case of MDLC-3, which was clinically reported as ER+ tumor but molecular profiling revealed

that the underlying ductal region was ER-/TNBC with basal-like IMS while the lobular region was

ER+ with luminal-A IMS. Notably, both ER- status and basal-like IMS are associated with worse

prognosis and outcome than ER+ and luminal IMS[76,77]. In this case, the patient had brain and

lung metastases, both of which were also ER-, suggesting progression of the primary ER- ductal

tumor. Another notable finding was enrichment of cell cycle arrest/senescence and ER signaling

signatures in lobular while *MYC* signatures in ductal tumor regions. Multiplex

immunofluorescence staining of various effector proteins confirmed that Myc and Cell

proliferation signaling were active in ductal tumor cells, while cell cycle arrest/senescence

signaling was active in lobular regions. Further analysis at single-cell RNA level revealed that ER

signatures were enriched in only one lobular subcluster (L2), highlighting molecular

heterogeneity even within individual histologically distinct tumor regions. Majority of ILC are

strongly ER+ and respond well to endocrine therapies, while they are mostly less proliferative,

responding poorly to chemotherapy[14,20,28,78,79]. Whether this is also true for lobular regions

within MDLC is an important question and must be further investigated. Perhaps, future trials

can study whether tumors with predominantly ductal vs. lobular MDLC respond differently to

endocrine and chemotherapies.

Development of robust mutational signature analysis tools such as Signature Multivariate

Analysis (SigMa) have allowed accurate identification of mutational signatures associated with

such HRD and APOBEC defects from targeted sequencing panels such as MSK-impact[80–84].

Utilization of SigMa algorithm in our study revealed that individual tumors regions with distinct

alterations may share common (Aging in MDLC-2) or distinct (APOBEC or HRD in MDLC-3)

mutational signatures. Collectively, our results indicate ductal and lobular regions in MDLC-2

have shared origin, while MDLC-3 is a clear collision of two independent tumors. The etiology of

MDLC-1 is questionable – while there are no shared mutations, we note that this interpretation

is based upon a small panel of genes measured by the MSK IMPACT panel with strict filtering.

We cannot exclude the possibility of shared mutations in other cancer-related genes not

covered by MSK-impact panel. Larger efforts using whole genome profiling of more MDLC

tumors are needed to comprehensively investigate the mutational signatures and origins of

ductal and lobular tumor regions in mixed tumors.

The presence of distinct clinically actionable alterations in individual tumor regions within

MDLC is of clinical relevance and has been previously reported MDLC[45]. Tumor regions with

distinct alterations can complicate clinical management. Alterations in components of ER

signaling (such as *GATA3, KMT2C*) as well as Notch signaling (*CBFB, NOTCH2*) can lead to

endocrine resistance[85–87], while alterations in DNA repair response genes (*ARID1A, ATRX*) may

confer increased sensitivity to chemotherapy and other DNA repair pathway targeting drugs

including PARP inhibitors[88,89]. Future studies will need to address the important question

whether treatment decisions for patients with MDLC benefits from considering the distinct

alterations in individual tumor regions.

For a long time, it was unclear whether ductal and lobular regions within MDLC share any

molecular similarity with their pure counterparts (i.e., ILC and IBC-NST tumors). To fill in the

gap, in our study, we compared the molecular features of individual tumor regions within MDLC

with their pure counterparts (i.e., ILC and IBC-NST tumors) in the TCGA cohort[28] and discovered

a set of shared markers of lobular phenotype. There was some but limited overlap of the differentially expressed genes between ductal and lobular tumor regions in MDLC cases with differential gene expression in ER+ ILC vs ER+ IBC-NST in the TCGA cohort.  This could be due to technical reasons such as bulk sequencing of frozen tumor samples (in TCGA) vs. spatial sequencing of epithelial tumor regions from FFPE tissues (in this study). Furthermore, this also could also be reflective of true biology i.e., unique gene expression patterns in MDLC tumor regions compared to pure ILC and IBC-NST tumors. As expected, one of overlapping genes that was downregulated in both MDLC lobular tumor regions and ILC tumors was *CDH1*, an important driver of lobular tumorigenesis[54]. [2854]This primarily occurs via genetic inactivation of *CDH1* in ILC tumors[28], similar to what was observed in the case of lobular regions in MDLC-1 and MDLC-3. However, in the lobular region of MDLC-2, *CDH1* promoter methylation was observed. *CDH1* promoter methylation is also associated with epithelial-to-mesenchymal transition (EMT), however, EMT is not a typical feature of lobular tumors and is primarily associated more complex subtypes of breast cancers such as metaplastic and claudin-low subtypes[90–93]. EMT markers[90] (Supplementary Fig. S1F)  were not consistently upregulated in MDLC-2 lobular region hence ruling out EMT-associated *CDH1* promoter methylation. This is an intriguing observation, suggesting that the pathogenesis of lobular regions within MDLC may occur via both genetic and epigenetic inactivation of *CDH1,* unlike ILC where genetic inactivation of *CDH1* is predominant[6,31,94–96]. A recent study revealed the synthetic lethality in E-cadherin deficient breast cancers with inhibition of *ROS1*, a tyrosine kinase[97], and trials are ongoing testing efficacy of *ROS1* inhibitors in ER+ ILC disease. Future studies could assess *ROS1*

inhibitors or other drugs in treating patients with predominantly lobular MDLC disease.

Additional studies are also warranted to investigate whether E-cadherin deficient tumors with

distinct etiology, such as *CDH1* epigenetic inactivation, have distinct biology potentially

associated with differential treatment response.

Our study focused primarily on profiling MDLC of collision type as these were the most

amenable for our molecular profiling approaches i.e., we could clearly annotate ductal and

lobular tumor regions based on histological and E-cadherin staining patterns and cleanly

capture RNA/DNA from these individual regions. Notably, we did attempt to profile ductal and

lobular tumor cells separately from intermixed regions identified in MDLC-2, however, the

resulting data was of poor quality due to low tumor cellularity. Although our study presents a

unique and novel analysis of MDLC of collision type, it has some limitations, including a small

sample size and lack of analysis of other subtypes (such as lobular-like IBC-NST[46] and

intermixed/intermingled MDLC[31,44]) and tumor microenvironment of individual tumor regions

within MDLC. Future studies, using rapidly improving spatial sequencing technologies with

increased single cell capabilities will be necessary to validate and expand our observations in

larger and more diverse cohorts of MDLC. Generating a comprehensive MDLC cohort for future

studies may not be an easy task, given the challenges associated with the diagnosis of this

mixed entity. However, the use of E-cadherin IHC in histological assessments and evaluation by

a panel of pathologists can ensure the generation of a high-confidence and well-annotated

MDLC cohort for future studies. Moreover, the application of digital pathology and artificial

intelligence will make this more feasible[98]. These tools will not only improve diagnostic

accuracy[99] but will also allow scalable annotation of ductal and lobular tumor regions across

heterogenous tumor samples[100].

In summary, our study provides new insights into the molecular heterogeneity of MDLC and its

underlying ductal and lobular tumor regions. This is the first report on genome-wide spatially

resolved and single-cell molecular profiling of MDLC. We discovered distinct biological

signatures and intrinsic molecular subtypes between these regions that may have important

implications for patient prognosis and clinical management. Our findings also highlight the need

for comprehensive histopathological and molecular profiling of individual tumor regions within

MDLC to better understand its progression and response to cytotoxic, endocrine, and other

targeted therapies.

## Methods

**Sample Collection**

Informed patient consent was obtained for tissue collection, and institutional review board approval was obtained from the University of Pittsburgh prior to initiation of the study. Patients diagnosed with "mixed invasive ductal and lobular" were identified from the UPMC Breast Cancer Registry and Formalin-Fixed Paraffin-Embedded (FFPE) blocks requested from Pitt Biospecimen Core. Four-um-thick FFPE sections were used for H&E, ER-alpha and E-cadherin IHC staining. A panel of pathologists reviewed the H&E and E-cadherin staining to confirm MDLC diagnosis in line with recent WHO guidelines[21] (10-90% of ILC and at least 10% IBC-NST) and shortlisted MDLC cases amenable for multi-omic profiling i.e., those with good separation of ductal and lobular tumor regions to avoid spill over and contamination of individual ductal and lobular tumor regions by neighboring lobular and ductal tumor cells, respectively. Hence, we selected three collision type MDLC cases that showed both clear separation of ductal and lobular tumor regions along with some intermixed/intermingled regions. These included cases MDLC-1, MDLC-2 & MDLC-3 (clinical features and pathology review notes are shown in Supplementary Table S1A & S1B). All cases underwent NanoString digital spatial, MSK-IMPACT mutation and *CDH1* promoter methylation profiling. Fresh tumor was only available for MDLC-1 and was utilized for single-cell RNAseq. See details in Supplementary Methods.

**Multiomic Profiling**

Digital spatial profiling (DSP) was performed using NanoString GeoMX whole transcriptome atlas platform. Mutation profiling was performed using MSK-IMPACT targeted NGS panel. *CDH1*

471 promoter methylation was assessed using ddPCR system. Finally, single cell RNA sequencing

472 was performed using 10X genomics platform based on 3'end chemistry v3. Please see

473 supplementary methods for details.

**Multiplexed immunofluorescence**

Tissue sections were prepared and iteratively stained as described previously[101]. Briefly, tissue

sections were baked overnight at 60⯑C, deparaffinized with xylene, rehydrated by decreasing

ethanol washes, followed by proprietary antigen retrieval optimized for multiple antigens (Leica

Biosystems). Tissues were incubated in 0.3% Triton X-100 in PBS for 10 min followed by

blocking in 10% (w/v) donkey serum and 3% (w/v) bovine serum albumin (BSA) in PBS for 1 h at

room temperature. Primary antibodies were diluted in blocking solution and incubated for 1 h

at room temperature (anti-E-cadherin, Cell Signaling Technology, 96743; anti-pan-cytokeratin,

Millipore Sigma, C5992; anti-c-Myc Alexa Fluor 647, Abcam, ab190560; anti-p16 Alexa Fluor

647, Abcam, ab192054; anti-cyclin E2 Alexa Fluor 488, Abcam, ab207336; anti-cyclin A1, R&D

Systems, MAB7046; anti-p21, R&D Systems, AF1047; anti-phospho-H2AX Alexa Fluor 555,

Abcam, ab206900). In-house primary antibody fluorescent conjugates were generated using

the Alexa Fluor 647 labelling kit (Invitrogen, A20186) or the Alexa Fluor 555 labelling kit

(Invitrogen, A20187). Nuclear labelling was performed using Hoechst (2 ug/ml, Invitrogen,

H3570) for 10 min at room temperature. Imaging was performed in 50% glycerol (v/v) in PBS.

After imaging, dye inactivation was performed in an alkali solution containing H2O2 for 15 min

with agitation followed by a PBS wash. Samples were reimaged following dye inactivation to

measure residual fluorescence. Additional rounds of antibody staining and imaging were

performed as described above, starting with primary antibody incubation. Image acquisition, flat-field correction, autofluorescence removal, and registration were performed using the Cell DIVE Imager (Leica Biosystems). Single-cell segmentation was performed using Cellpose[102] and intensity measurements were extracted using scikit-image[103]. Measurements greater than 1.5x the interquartile range were identified as outliers and removed from analysis.

**Figures & Illustrations**

DSP spatial images were exported from GeoMx® Data Analysis Suite (DSPDA) and used directly in the manuscript without graphical modifications. Most bioinformatic analysis visualizations were generated using R. These visualizations were aesthetically improved in power point to produce the final figures. Illustrations were prepared using BioRender.com.

**Data Availability**

Data and code generated in this study is shared in the Supplementary Data files, Mendeley (https://data.mendeley.com/datasets/btv7g7n9ys/1) and GitHub (https://github.com/osamashiraz/MDLC_Spatial_Analysis_2023).

## Authors' Disclosures

PCL has an equity interest in Amgen (outside the submitted work). BW reports research funding by Repare Therapeutics, outside the submitted work. JSR-F reports having received personal/consultancy fees from Goldman Sachs, Bain Capital, Repare Therapeutics, Saga Diagnostics and Paige.AI, membership of the scientific advisory boards of VolitionRx, REPARE Therapeutics and Paige.AI, membership of the Board of Directors of Grupo Oncoclinicas, and ad hoc membership of the scientific advisory boards of AstraZeneca, Merck, Daiichi Sankyo, Roche Tissue Diagnostics and Personalis, outside the scope of this study. No disclosures were reported by other authors.

## Authors' Contributions

**O.S. Shah:** Conceptualization, data curation, software, formal analysis, investigation, visualization, methodology, project administration, writing–original draft, writing–review and editing. **A. Nasrazadani:** resources, funding acquisition, writing-review and editing. **J.M. Atkinson:** resources, supervision, project administration, writing–review and editing. **J. Foldi:** resources, investigation, data curation, writing–review and editing. **C.G. Kleer:** investigation, data curation, writing–review and editing. **P.F. McAuliffe:** investigation, resources, data curation, writing–review and editing. **T.J. Johnston**: investigation, resources, data curation, software, formal analysis, writing–review and editing. **W. Stallaert**: investigation, resources, data curation, software, formal analysis, writing–review and editing. **E.M. da Silva:** investigation, resources, data curation, software, formal analysis, writing–review and editing. **P. Selenica:** investigation, resources, data curation, software, formal analysis, writing–review and editing. **H. Dopeso:** investigation, resources, data curation, writing–review and editing. **F. Pareja:** investigation, resources, data curation, writing–review and editing. **D. Mandelker:** investigation, resources, data curation, writing–review and editing. **B. Weigelt:** conceptualization, investigation, methodology, resources, writing–review and editing. **J.S. Reis-Filho:** conceptualization, investigation, methodology, resources, writing–review and editing. **R. Bhargava:** conceptualization, investigation, resources, data curation, methodology, data curation, writing–review and editing. **P.C. Lucas:** conceptualization, investigation, resources, data curation, methodology, data curation, writing–review and editing. **A.V. Lee:** conceptualization, resources, methodology, supervision, funding acquisition, project

536 administration, writing–review and editing. **S. Oesterreich:** conceptualization, resources,

537 methodology, supervision, funding acquisition, project administration, writing–review and

538 editing.

## Acknowledgments

This work was primarily supported by the Dynami Foundation (Flora Migyanka) and BCRF grants to **S. Oesterreich** and in part by various other grants including the ASCO Gianni Bonadonna Breast Cancer Research Fellowship grant to **A. Nasrazadani** and Conquer Cancer ASCO Young Investigator Award to **J. Foldi**. We would also like to thank NanoString DSP Technology Access Program for providing early access to their whole transcriptome spatial technology, UPMC Hillman Cancer Center and Tissue and Research Pathology/Pitt Biospecimen Core shared resources for providing access to valuable clinical specimens and technical support (award P30CA047904), University of Pittsburgh Center for Research Computing, RRID:SCR_022735 (HTC cluster, NIH Award S10OD028483) , Pitt Genomics Research Core and UPMC Genome Center for technical support in preparing scRNAseq libraries and sequencing, and Ms. Morgan DeBerry, HTL(ASCP)CM, QIHCCM for excellent technical assistance in tissue sectioning, staining and other logistics. Research reported in this publication was partly funded by an NIH/NCI Cancer Center Support Grant (P30CA008748, MSKCC). BW, FP and JSR-F were funded in part by an NIH/NCI P50 CA247749 01 and Breast Cancer Research Foundation grants.

**Figure 1. Digital Spatial Profiling of MDLC Ductal and Lobular Tumor Regions.** (A) Top panel shows clinical timelines of the studied cases including the time of core biopsies, surgical interventions, treatments, metastasis and last known vital status. The source of the tumor tissue used in this study is highlighted with * in red color. The bottom panels show high (40X) magnification H&E and Ecad/p120 IHC images highlighting histomorphic and E-cadherin features of ductal and lobular regions across each case. (B) E-cadherin and Pan-CK immunofluorescence staining of a representative case (MDLC-3). Selected regions of interests (ROIs) are shown with dotted blue (for ductal) and red (for lobular). (C) Segmentation of ductal and lobular cells using E-cadherin and Pan-CK staining masks. (D) Segmented areas containing only ductal or lobular tumor cells defined using ductal and lobular masks illustrated in panel C. (E) E-cadherin staining intensity in representative ductal and lobular regions profiled across cases.

**Figure 2. Molecular Subtypes and Differentially Expressed Genes in Ductal and Lobular Tumor Regions.** (A) Profiled ductal and lobular tumor regions separated by consensus clusters. All regions are annotated by cases, histology, and consensus clusters. Gene expression heatmap shows top variable genes and followed by PAM50 scores for each subtype across ductal and lobular regions. (B) Differentially expressed genes (DEGs) between ER+ ductal and lobular regions. The top panel shows DEG heatmap, while bottom panel shows volcano plot along with number of significantly upregulated (N=168) and downregulated (N=198) DEGs in ductal vs lobular regions.

**Figure 3. Distinct Hallmark and Biological Signatures in Ductal and Lobular Tumor Regions.** (A)

Differentially enriched cancer hallmark signatures. The heatmap shows GSVA scores of each

differentially enriched pathway in ductal vs lobular tumor regions. Pathways with bold red and

blue names are the ones that also were significant in the scRNAseq dataset (Supplementary Fig.

S8A). Enrichment score (ES) for a pathway shows mean differences of GSVA score for that

pathway in lobular – ductal regions and the -log10 of FDR-adjusted P-value (P-val) shows

statistical significance of enrichment based on T test. (P-val score definitions: 4 = P ≤ 0.0001, 3 =

P ≤ 0.001, 2 = P ≤ 0.01 and 1 = P ≤ 0.05). (B) The heatmap shows GSVA scores of differentially

enriched signatures between TCGA ER+ IBC-NST vs. ILC that overlap with MDLC ductal vs lobular

tumor region hallmark signatures in panel A. (C) Recurrent biological themes in ductal and

lobular tumor regions. The left panel shows the Jaccard similarity between all signatures.

Signatures are grouped based on their Jaccard similarity into signature clusters. Each cluster is

associated with a broad biological theme (shown in middle). The right panel shows the mean

GSVA score of each signature cluster in ductal and lobular regions. (D) E-cadherin (ECad) and

pan-cytokeratin (pan-CK26) staining in representative image of adjacent ductal and lobular

tumor regions. (E) Representative immunofluorescence images (right panel) and single-cell

quantification (left panel) of cell cycle and senescence biomarkers from all ER+ ductal and

lobular regions. Statistical significance is determined using a student's t-test (****: P ≤ 0.0001).

ECM: Extracellular matrix, DSB: Double stranded break, HR: Homologous recombination.

**Figure 4. Overview of Genomic Alterations in Ductal and Lobular Tumor Regions.** (A) Somatic

mutations identified using MSK-IMPACT panel in ductal and lobular components across

samples. Gene names are followed by amino acid change due to mutation. Genes in black font

had somatic mutations in breast cancer driver genes while those in gray color had

passenger/non-driver mutations. (B) Somatic copy number changes in ductal and lobular

components across samples. For each case, genome-wide ductal and lobular copy number

profiles are shown. Black line shows total copy number, red line shows minor allele copy

number and off-white regions show aneuploidy (change in minor allele copy number including

LOH). (C) Mutational signatures in MDLC-2 and MDCL-3 ductal and lobular tumor regions. Pie

charts show percentage similarity of mutational patterns of each tumor component with

established mutational patterns (associated with different means/definition). (D) Predicted

disease evolution. Tumors with overlapping mutations were linked to common root node

indicating common progenitor or precursor neoplasm (N). Those without common root node

likely indicate evolution from independent precursors. Key mutations and events (*CDH1*

promoter methylation and amplifications) are shown on the branches.

**Figure 5: Single Cell Heterogeneity of Ductal and Lobular Tumor Populations.** (A) Tumor

populations in MDLC-1 scRNAseq dataset (left) and assessment of CDH1 expression in MDLC-1

scRNAseq dataset (middle and right plot). Statistical significance was evaluated using the T test.

(B) Deconvolution of scRNAseq using DSP derived ductal and lobular signatures. GSVA scores

for ductal and lobular signatures are shown on the right panel. (C) Sub-clustering and

identification of ductal and lobular sub-populations (left) with unique marker genes (right). (D)

Scoring spatial regions using GSVA of sub-cluster marker genes. (E) Mapping of single cell

populations to spatial regions based on GSVA scores in panel D.

**Supplementary Figure S1: Digital Spatial Profiling RNAseq Dataset Quality Control.** (A) Scatter plot showing sequencing saturation (y-axis) and area of interest (AOI) surface area (mm2) across cases. Dashed red lines show the quality control (QC) cutoffs (sequencing saturation >50% and area > 6000mm$^2$) for selecting high quality samples. (B) Overview of signal (Q3-normalized counts) to noise (negative probe counts). Dashed red line shows cutoff i.e., signal > noise. (C) Evaluation of Q3 normalized counts (for 1000 randomly sampled genes) across profiled regions to identify samples with low quality data showing count artifacts. (D) tSNE plot showing clustering of profiled samples that passed QC cutoffs. (E) *CDH1, ESR1, ERBB2* and *PGR* gene expression in ductal vs lobular regions across the three profiled cases. Statistical significance was evaluated using the T test. (F) Expression of EMT marker genes in ductal and lobular tumor regions across each case. EMT marker genes were adapted from McCart Reed et al[34].

**Supplementary Figure S2: Patient Wise and Pan-Patient MDLC Ductal vs Lobular Differential Analysis.** Volcano plots (top) showing per patient or patient-wise differentially expressed genes (DEGs) between ductal vs lobular tumor regions and overlap of these patient-wise DEGs with pan-patient (all patients combined) DEGs for (A) MDLC-1, (B) MDLC-2 and (C) MDLC-3. A RF > 1 and p < 0.01 shows significantly strong overlap. (D) Overlap of MDLC pan-patient ductal vs. lobular DEGs with TCGA ER+ IBC-NST vs ILC DEGs. Statistical significance of overlap between genesets was computed using overlap_stats tool (http://nemates.org/MA/progs/overlap_stats.html). RF: Representation Factor. For details on

RF calculation see overlap_stats webpage. Briefly, RF is the number of overlapping genes

divided by the expected number of overlapping genes drawn from two independent groups. A

RF > 1 indicates more overlap than expected for independent genesets, a RF < 1 indicates less

overlap than expected, and a RF = 1 indicates that the two genesets by the number of genes

expected for independent groups of genes.

**Supplementary Figure S3: Transcription Factor Enrichment Analysis using Pan-Patient Ductal**

**vs Lobular DEGs.** Ductal vs Lobular DEGs were separately input into eXpression2Kinase (X2K)

web tool (https://maayanlab.cloud/X2K/) to identify enriched transcription factors and kinases

in ductal (left) and lobular (right) tumor regions. The top layer with blue nodes shows kinases,

middle layer shows intermediate proteins and bottom layer with red nodes shows transcription

factors. An image inset is shown highlighting MYC and ER signaling related transcription factors

enriched in ductal and lobular tumor regions, respectively.

**Supplementary Figure S4: Key Genes Associated with Notable Differentially Enriched**

**Oncogenic and Biological Signatures.** (A) Venn diagram showing overlap between D1: Double

stranded break/homologous recombination (DBS/HR), Hallmark G2m Checkpoint and Hallmark

Myc Targets V1. (B). Venn diagram showing overlap between Hallmark Myc Targets V1, L1: Cell

Cycle and L2: TP53 & Senescence. Details of all genes in each signature shown in (A) and (B) are

listed in Supplementary Tables S6 and S7. The genes used for comparing biological theme

signature clusters (D1: DBS/HR, L1: Cell Cycle and L2: TP53 & Senescence) are defined as union

of all individual signatures present in each of these signature clusters. (C) Expression of select

genes from Hallmark Myc Targets V1, L1: Cell Cycle, L2: TP53/Senescence, L5: Collagen & ECM,

L7: Collagen metabolism, L9: MAPK/ERK Targets, and D5: mTOR in ductal vs. lobular tumor

regions. Statistical significance was evaluated using the T test.

**Supplementary Figure S5: MSK-IMPACT Extended Results.** (A) Tumor purity, sequencing

coverage and ploidy of ductal and lobular tumor regions across profiled cases. (B) Somatic

mutations and their cancer cell fraction (CCF). High CCF indicate clonal mutations while low CCF

indicates sub-clonal mutations. (C) Copy number of key breast cancer genes in ductal vs lobular

regions across each case.

**Supplementary Figure S6: Effect of Q23 mutation on CDH1 RNA and E-cadherin protein**

**expression in ER+ Breast Tumors.** Bar plots showing CDH1 RNA (top) and E-cadherin protein

(bottom) expression in ER+ patient tumor samples (N=434) with CDH1 Q23* mutations (N=5),

all other CDH1 truncating mutations (N=71) and CDH1 WT (N=358) in ER+ breast tumors in

TCGA cohort. To avoid confounding effects of CDH1 copy number, we excluded all tumor

samples with CDH1 gains, amplification, and deep deletions. Statistical significance is

determined using student's t-test (****: P ≤ 0.0001, **: P ≤ 0.01).

**Supplementary Figure S7: MDLC-1 Ductal and Lobular Tumor scRNAseq Heterogeneity.** (A)

MDLC-1 single-cell RNAseq (scRNAseq) dataset quality control metrics from cell ranger

software. (B) tSNE plot showing identified cell types in the scRNAseq dataset including T-cells,

Endothelial cells (Endo), Macrophages (Macro), Fibroblasts (Fibro), Smooth muscle cells,

luminal tumor epithelial cells (Tumor cells), luminal epithelial cells (Luminal cells) and basal

epithelial cells (Basal cells). (C) CopyKat algorithm predictions for copy number status of

scRNAseq dataset to confirm aneuploidy status of tumor cells and diploid status of non-tumor

cells. (D) MDLC-1 scRNAseq ductal vs lobular DEG analysis. (E) Heatmap of CopyKat inferred

copy number values for ductal and lobular tumor cell populations. (F) Overlap of MDLC-1 ductal

vs lobular DEGs between scRNAseq and digital spatial profiling (DSP) datasets. A RF > 1 and p <

0.01 shows significantly strong overlap. Statistical significance of DEG overlap between DEGs

from two technologies was computed using overlap_stats tool as described previously in

Supplementary Fig. S2. (G) Top five marker genes of each ductal and lobular sub-cluster

identified in Fig. 5C.

**Supplementary Figure S8: Distinct Biological Signatures in MDLC-1 Ductal and Lobular Tumor**

**scRNAseq.** (A) GSVA based hallmark signature enrichment analysis in scRNAseq dataset.

Enrichment score (ES) for a pathway shows mean differences of GSVA score for that pathway in

lobular – ductal regions and the -log10 of FDR-adjusted P-value (P-val) shows statistical

significance of enrichment based on T test. (P-val score definitions: 4 = $P \leq 0.0001$, 3 = $P \leq 0.001$,

2 = $P \leq 0.01$ and 1 = $P \leq 0.05$). (B) Barplots showing mean GSVA scores of hallmark signatures in

ductal and lobular subclusters. (C) Recurring biology themes in ductal vs lobular cells in

scRNAseq dataset. (D) Mean GSVA score for select differentially enriched biology themes across

ductal and lobular sub-clusters.

References

1. Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. **68**, 394–424 (2018).

2. Siegel, R. L., Miller, K. D., Wagle, N. S. & Jemal, A. Cancer statistics, 2023. *CA Cancer J Clin* **73**, 17–48 (2023).

3. Sledge, G. W., Chagpar, A. & Perou, C. Collective wisdom: lobular carcinoma of the breast. *Am Soc Clin Oncol Educ B* **35**, 18–21 (2016).

4. Reed, M. E. M. C., Kutasovic, J. R., Lakhani, S. R. & Simpson, P. T. Invasive lobular carcinoma of the breast: morphology, biomarkers and 'omics. **17**, 12 (2015).

5. Ciriello, G. *et al.* Comprehensive molecular portraits of invasive lobular breast cancer. *Cell* **163**, 506–519 (2015).

6. Reed, M. E. M. C., Kutasovic, J. R., Lakhani, S. R. & Simpson, P. T. Invasive lobular carcinoma of the breast: morphology, biomarkers and 'omics. **17**, 12 (2015).

7. Dabbs, D. J., Bhargava, R. & Chivukula, M. Lobular versus ductal breast neoplasms: the diagnostic utility of p120 catenin. *Am J Surg Pathol* **31**, 427–37 (2007).

8. Arpino, G. *et al.* Infiltrating lobular carcinoma of the breast: tumor characteristics and clinical outcome. **6**, (2004).

9. Oesterreich, S. *et al.* Clinicopathological Features and Outcomes Comparing Patients With Invasive Ductal and Lobular Breast Cancer. *J Natl Cancer Inst* **114**, 1511–1522 (2022).

10. Z, C. *et al.* Invasive lobular carcinoma of the breast: A special histological type compared with invasive ductal carcinoma. **12**, e0182397 (2017).

11. Yeatman, T. J. *et al.* Tumor biology of infiltrating lobular carcinoma. Implications for management. *Ann Surg* **222**, 549–559 (1995).

12. Helvie, M. A., Paramagul, C., Anderson, K. L. & Adler, D. D. Invasive lobular carcinoma. Imaging features and clinical detection. *Invest Radiol* **28**, 202–207 (1993).

13. Arpino, G. *et al.* Infiltrating lobular carcinoma of the breast: tumor characteristics and clinical outcome. **6**, (2004).

14. Oesterreich, S. *et al.* Clinicopathological Features and Outcomes Comparing Patients With Invasive Ductal and Lobular Breast Cancer. *J Natl Cancer Inst* **114**, 1511–1522 (2022).

15. Mathew, A. *et al.* Distinct Pattern of Metastases in Patients with Invasive Lobular Carcinoma of the Breast. *Geburtshilfe Frauenheilkd* (2017) doi:10.1055/s-0043-109374.

16. Lamovec, J. & Bračko, M. Metastatic pattern of infiltrating lobular carcinoma of the breast: an autopsy study. **48**, 28–33 (1991).

17. Djerroudi, L., Cabel, L., Bidard, F. C. & Vincent-Salomon, A. Invasive Lobular Carcinoma of the Breast: Toward Tailoring Therapy? **114**, 1434–1436 (2022).

18. K, V. B. *et al.* Current and future diagnostic and treatment strategies for patients with invasive lobular breast cancer. *Annals of Oncology* **33**, 769–785 (2022).

19. Luveta, J., Parks, R. M., Heery, D. M., Cheung, K.-L. & Johnston, S. J. Invasive Lobular
Breast Cancer as a Distinct Disease: Implications for Therapeutic Strategy. *Oncol Ther* **8**,
1–11 (2020).

20. K, V. B. *et al.* Current and future diagnostic and treatment strategies for patients with
invasive lobular breast cancer. *Annals of Oncology* **33**, 769–785 (2022).

21. Hoon Tan, P. *et al.* The 2019 WHO classification of tumours of the breast. *Histopathology*
(2020) doi:10.1111/his.14091.

22. Sinn, H. P. & Kreipe, H. *A Brief Overview of the WHO Classification of Breast Tumors,*
*4th Edition, Focusing on Issues and Updates from the 3rd Edition. Breast Care* vol. 8 149
(Karger Publishers, 2013).

23. Metzger-Filho, O. *et al. The Diagnosis of Mixed Invasive Ductal and Lobular Carcinoma*
*(IDC-L) in Clinical Practice Is Often Associated with Uncertainty Related to Its Prognosis*
*and Response to Systemic Therapies. With the Increasing Recognition of Invasive Lobular*
*Carc*. vol. 24 (Oxford University Press (OUP), 2019).

24. Zengel, B. *et al.* Comparison of the clinicopathological features of invasive ductal,
invasive lobular, and mixed (invasive ductal + invasive lobular) carcinoma of the breast.
*Breast Cancer* (2015) doi:10.1007/s12282-013-0489-8.

25. Xiao, Y. *et al.* Mixed invasive ductal and lobular carcinoma has distinct clinical features
and predicts worse prognosis when stratified by estrogen receptor status. *Sci Rep* (2017)
doi:10.1038/s41598-017-10789-x.

26. Duraker, N., Hot, S., Akan, A. & Ozay Nayir, P. A Comparison of the Clinicopathological
Features, Metastasis Sites and Survival Outcomes of Invasive Lobular, Invasive Ductal
and Mixed Invasive Ductal and Lobular Breast Carcinoma. *Eur J Breast Health* **16**, 22–31
(2020).

27. Nasrazadani, A. *et al.* Mixed invasive ductal lobular carcinoma is clinically and
pathologically more similar to invasive lobular than ductal carcinoma. *British Journal of*
*Cancer 2023* 1–10 (2023) doi:10.1038/s41416-022-02131-8.

28. Ciriello, G. *et al.* Comprehensive molecular portraits of invasive lobular breast cancer.
*Cell* **163**, 506–519 (2015).

29. Nasrazadani, A. *et al.* Mixed invasive ductal lobular carcinoma is clinically and
pathologically more similar to invasive lobular than ductal carcinoma. *British Journal of*
*Cancer 2023* 1–10 (2023) doi:10.1038/s41416-022-02131-8.

30. Lohani, K. R. *et al.* Lobular-Like Features and Outcomes of Mixed Invasive Ductolobular
Breast Cancer (MIDLC): Insights from 54,403 Stage I-III MIDLC Patients. *Ann Surg*
*Oncol* **31**, 936–946 (2024).

31. Pérez-Mies, B. *et al.* The Clonal Relationship Between the Ductal and Lobular
Components of Mixed Ductal-Lobular Carcinomas Suggested a Ductal Origin in Most
Tumors. *American Journal of Surgical Pathology* **46**, 1545–1553 (2022).

32. Li, C. I., Anderson, B. O., Daling, J. R. & Moe, R. E. Trends in Incidence Rates of
Invasive Lobular and Ductal Breast Carcinoma. *J Am Med Assoc* (2003)
doi:10.1001/jama.289.11.1421.

33. Suryadevara, A., Paruchuri, L. P., Banisaeed, N., Dunnington, G. & Rao, K. A. The
clinical behavior of mixed ductal/lobular carcinoma of the breast: A clinicopathologic
analysis. *World J Surg Oncol* (2010) doi:10.1186/1477-7819-8-51.

34. Rakha, E. A. *et al.* The biological and clinical characteristics of breast carcinoma with
mixed ductal and lobular morphology. *Breast Cancer Res Treat* (2009)
doi:10.1007/s10549-008-0007-4.

35. Acs, G., Lawton, T. J., Rebbeck, T. R., LiVolsi, V. A. & Zhang, P. J. Differential
Expression of E-Cadherin in Lobular and Ductal Neoplasms of the Breast and Its Biologic
and Diagnostic Implications. *Am J Clin Pathol* (2001) doi:10.1309/FDHX-L92R-BATQ-
2GE0.

36. McCart Reed, A. E. *et al.* Mixed ductal-lobular carcinomas: evidence for progression from
ductal to lobular morphology. *Journal of Pathology* (2018) doi:10.1002/path.5040.

37. Christgen, M. *et al.* Inter-observer agreement for the histological diagnosis of invasive
lobular breast carcinoma. *J Pathol Clin Res* **8**, 191–205 (2022).

38. Metzger□Filho, O. *et al.* Mixed Invasive Ductal and Lobular Carcinoma of the Breast:
Prognosis and the Importance of Histologic Grade. *Oncologist* **24**, (2019).

39. Zhao, H. The prognosis of invasive ductal carcinoma, lobular carcinoma and mixed ductal
and lobular carcinoma according to molecular subtypes of the breast. *Breast Cancer* **28**,
187–195 (2021).

40. Arps, D. P., Healy, P., Zhao, L., Kleer, C. G. & Pang, J. C. Invasive ductal carcinoma with
lobular features: a comparison study to invasive ductal and invasive lobular carcinomas of
the breast. *Breast Cancer Res Treat* **138**, 719 (2013).

41. Zels, G. *et al.* Metastases of primary mixed no-special type and lobular breast cancer
display an exclusive lobular histology. *Breast* **75**, 103732 (2024).

42. M, C. *et al.* Lobular breast cancer: Histomorphology and different concepts of a special
spectrum of tumors. **13**, 3695 (2021).

43. Christgen, M., Kandt, L. D., Antonopoulos, W. & al., et. Inter-observer agreement for the
histological diagnosis of invasive lobular breast carcinoma. *J Pathol Clin Res* **8**, 191–205
(2022).

44. Tang, X. *et al.* Mixed ductal–lobular carcinoma: an analysis of CDH1 DNA copy number
variation and mutation. *Breast Cancer* **28**, 1318–1327 (2021).

45. McCart Reed, A. E. *et al.* Mixed ductal-lobular carcinomas: evidence for progression from
ductal to lobular morphology. *Journal of Pathology* (2018) doi:10.1002/path.5040.

46. Yu, J. *et al.* Clinicopathologic and genomic features of lobular like invasive mammary
carcinoma: is it a distinct entity? *npj Breast Cancer 2023 9:1* **9**, 1–13 (2023).

47. Ciriello, G. *et al.* Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer.
*Cell* **163**, 506–519 (2015).

48. Suryadevara, A., Paruchuri, L. P., Banisaeed, N., Dunnington, G. & Rao, K. A. The
clinical behavior of mixed ductal/lobular carcinoma of the breast: A clinicopathologic
analysis. *World J Surg Oncol* (2010) doi:10.1186/1477-7819-8-51.

49.    Pérez-Mies, B. *et al.* The Clonal Relationship Between the Ductal and Lobular
817           Components of Mixed Ductal-Lobular Carcinomas Suggested a Ductal Origin in Most
818           Tumors. *American Journal of Surgical Pathology* **46**, 1545–1553 (2022).

819    50.    Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection.
820           *Cell Syst* **1**, 417–425 (2015).

821    51.    Carbon, S. *et al.* The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res*
822           **49**, D325–D334 (2021).

823    52.    Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res* **50**,
824           D687–D692 (2022).

825    53.    Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG
826           for taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res* **51**, D587–
827           D592 (2023).

828    54.    Cleton-Jansen, A. M. E-cadherin and loss of heterozygosity at chromosome 16 in breast
829           carcinogenesis: different genetic pathways in ductal and lobular breast cancer? *Breast*
830           *Cancer Res* **4**, 5–8 (2002).

831    55.    André, F. *et al.* Alpelisib for PIK3CA -Mutated, Hormone Receptor–Positive Advanced
832           Breast Cancer . *New England Journal of Medicine* **380**, 1929–1940 (2019).

833    56.    Theodorou, V., Stark, R., Menon, S. & Carroll, J. S. GATA3 acts upstream of FOXA1 in
834           mediating ESR1 binding by shaping enhancer accessibility. *Genome Res* **23**, 12 (2013).

835    57.    Malik, N. *et al.* The transcription factor CBFB suppresses breast cancer through
836           orchestrating translation and transcription. *Nature Communications 2019 10:1* **10**, 1–15
837           (2019).

838    58.    Comprehensive molecular portraits of human breast tumours. *Nature* **490**, (2012).

839    59.    Gala, K. *et al.* KMT2C mediates the estrogen dependence of breast cancer through
840           regulation of ERα enhancer function. *Oncogene 2018 37:34* **37**, 4692–4710 (2018).

841    60.    Glaser, A. P. *et al.* APOBEC-mediated mutagenesis in urothelial carcinoma is associated
842           with improved survival, mutations in DNA damage response genes, and immune response.
843           *Oncotarget* **9**, 4537–4548 (2017).

844    61.    Shen, R. & Seshan, V. E. FACETS: allele-specific copy number and clonal heterogeneity
845           analysis tool for high-throughput DNA sequencing. *Nucleic Acids Res* **44**, (2016).

846    62.    Glaser, A. P. *et al.* APOBEC-mediated mutagenesis in urothelial carcinoma is associated
847           with improved survival, mutations in DNA damage response genes, and immune response.
848           *Oncotarget* **9**, 4537–4548 (2017).

849    63.    Klonowski, J., Liang, Q., Coban-Akdemir, Z., Lo, C. & Kostka, D. aenmd: annotating
850           escape from nonsense-mediated decay for transcripts with protein-truncating variants.
851           *Bioinformatics* **39**, (2023).

852    64.    Khajavi, M., Inoue, K. & Lupski, J. R. Nonsense-mediated mRNA decay modulates
853           clinical outcome of genetic disease. *European Journal of Human Genetics 2006 14:10* **14**,
854           1074–1081 (2006).

855    65.    Litchfield, K. *et al.* Escape from nonsense-mediated decay associates with anti-tumor
856           immunogenicity. *Nature Communications 2020 11:1* **11**, 1–11 (2020).

857    66.    Moffitt, J. R., Lundberg, E. & Heyn, H. The emerging landscape of spatial profiling
858           technologies. *Nature Reviews Genetics 2022 23:12* **23**, 741–759 (2022).

859    67.    van Amerongen, R. Behind the Scenes of the Human Breast Cell Atlas Project. *J
860           Mammary Gland Biol Neoplasia* **26**, 67 (2021).

861    68.    Nguyen, Q. H. *et al.* Profiling human breast epithelial cells using single cell RNA
862           sequencing identifies cell diversity. *Nat Commun* (2018) doi:10.1038/s41467-018-04334-
863           1.

864    69.    Gray, G. K. *et al.* A human breast atlas integrating single-cell proteomics and
865           transcriptomics. *Dev Cell* **57**, 1400-1420.e7 (2022).

866    70.    Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nature
867           Genetics 2021 53:9* **53**, 1334–1347 (2021).

868    71.    Sun, H. *et al.* The relevance between hypoxia-dependent spatial transcriptomics and the
869           prognosis and efficacy of immunotherapy in claudin-low breast cancer. *Front Immunol*
870           **13**, 7828 (2023).

871    72.    Bassiouni, R. *et al.* Spatial Transcriptomic Analysis of a Diverse Patient Cohort Reveals a
872           Conserved Architecture in Triple-Negative Breast Cancer. *Cancer Res* **83**, 34–48 (2023).

873    73.    Johnson, B. E. *et al.* An omic and multidimensional spatial atlas from serial biopsies of an
874           evolving metastatic breast cancer. *Cell Rep Med* **3**, 100525 (2022).

875    74.    Kulasinghe, A. *et al.* Spatial Profiling Identifies Prognostic Features of Response to
876           Adjuvant Therapy in Triple Negative Breast Cancer (TNBC). *Front Oncol* **11**, 1 (2021).

877    75.    Andersson, A. *et al.* Spatial deconvolution of HER2-positive breast cancer delineates
878           tumor-associated cell type interactions. *Nature Communications 2021 12:1* **12**, 1–14
879           (2021).

880    76.    Badve, S. *et al.* Basal-like and triple-negative breast cancers: a critical review with an
881           emphasis on the implications for pathologists and oncologists. *Modern Pathology* **24**,
882           157–167 (2011).

883    77.    Perou, C. M. Molecular Stratification of Triple☐Negative Breast Cancers. *Oncologist* **16**,
884           61–70 (2011).

885    78.    Michaut, M. *et al.* Integration of genomic, transcriptomic and proteomic data identifies
886           two biologically distinct subtypes of invasive lobular breast cancer. *Sci Rep* (2016)
887           doi:10.1038/srep18517.

888    79.    Djerroudi, L., Cabel, L., Bidard, F. C. & Vincent-Salomon, A. Invasive Lobular
889           Carcinoma of the Breast: Toward Tailoring Therapy? **114**, 1434–1436 (2022).

890    80.    Gulhan, D. C., Lee, J. J. K., Melloni, G. E. M., Cortés-Ciriano, I. & Park, P. J. Detecting
891           the mutational signature of homologous recombination deficiency in clinical samples. *Nat
892           Genet* **51**, 912–919 (2019).

893    81.    Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**,
894           415–421 (2013).

82. Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature*
**578**, 94–101 (2020).

83. Vahdatinia, M. *et al.* KIT genetic alterations in breast cancer. *J Clin Pathol* (2022)
doi:10.1136/JCP-2022-208611.

84. Geyer, F. C. *et al.* Genetic analysis of a morphologically heterogeneous ovarian
endometrioid carcinoma. *Histopathology* **71**, 480 (2017).

85. Azim, H. A., Nguyen, B., Brohée, S., Zoppoli, G. & Sotiriou, C. Genomic aberrations in
young and elderly breast cancer patients. *BMC Med* **13**, 1–7 (2015).

86. Afzaljavan, F., Sadr, A. S., Savas, S. & Pasdar, A. GATA3 somatic mutations are
associated with clinicopathological features and expression profile in TCGA breast cancer
patients. *Scientific Reports 2021 11:1* **11**, 1–13 (2021).

87. BeLow, M. & Osipo, C. Notch Signaling in Breast Cancer: A Role in Drug Resistance.
*Cells 2020, Vol. 9, Page 2204* **9**, 2204 (2020).

88. Hopkins, J. L., Lan, L. & Zou, L. DNA repair defects in cancer and therapeutic
opportunities. *Genes Dev* **36**, 278 (2022).

89. Voutsadakis, I. A. & Stravodimou, A. Homologous Recombination Defects and Mutations
in DNA Damage Response (DDR) Genes Besides BRCA1 and BRCA2 as Breast Cancer
Biomarkers for PARP Inhibitors and Other DDR Targeting Therapies. *Anticancer Res* **43**,
967–981 (2023).

90. McCart Reed, A. E. *et al.* An epithelial to mesenchymal transition programme does not
usually drive the phenotype of invasive lobular carcinomas. *Journal of Pathology* **238**,
489–494 (2016).

91. Herschkowitz, J. I. *et al.* Identification of conserved gene expression features between
murine mammary carcinoma models and human breast tumors. *Genome Biol* **8**, 1–17
(2007).

92. Pan, C. *et al.* Research progress of Claudin-low breast cancer. *Front Oncol* **13**, 1226118
(2023).

93. Dias, K. *et al.* Claudin-Low Breast Cancer; Clinical & Pathological Characteristics. *PLoS*
*One* **12**, (2017).

94. McCart Reed, A. E. *et al.* Mixed ductal-lobular carcinomas: evidence for progression from
ductal to lobular morphology. *Journal of Pathology* (2018) doi:10.1002/path.5040.

95. Kutasovic, J. R., McCart Reed, A. E., Sokolova, A., Lakhani, S. R. & Simpson, P. T.
Morphologic and genomic heterogeneity in the evolution and progression of breast cancer.
*Cancers* vol. 12 Preprint at https://doi.org/10.3390/cancers12040848 (2020).

96. Ciriello, G. *et al.* Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer.
*Cell* (2015) doi:10.1016/j.cell.2015.09.033.

97. Bajrami, I. *et al.* E-Cadherin/ROS1 Inhibitor Synthetic Lethality in Breast Cancer. *Cancer*
*Discov* **8**, 498–515 (2018).

98. Jiang, Y., Yang, M., Wang, S., Li, X. & Sun, Y. Emerging role of deep learning-based
artificial intelligence in tumor pathology. *Cancer Commun* **40**, 154–166 (2020).

935 99. Couture, H. D. *et al.* Image analysis with deep learning to predict breast cancer grade, ER
936   status, histologic subtype, and intrinsic subtype. *npj Breast Cancer 2018 4:1* **4**, 1–8
937   (2018).

938 100. Chan, R. C. K. *et al.* Artificial intelligence in breast cancer histopathology.
939   *Histopathology* **82**, 198–210 (2023).

940 101. Gerdes, M. J. *et al.* Highly multiplexed single-cell analysis of formalin-fixed, paraffin-
941   embedded cancer tissue. *Proc Natl Acad Sci U S A* **110**, 11982–11987 (2013).

942 102. Pachitariu, M. & Stringer, C. Cellpose 2.0: how to train your own model. *Nat Methods* **19**,
943   1634–1641 (2022).

944 103. Van Der Walt, S. *et al.* scikit-image: image processing in Python. *PeerJ* **2**, (2014).

**Figure 1. Digital Spatial Profiling of mDLC Ductal and Lobular Tumor Regions.**

**Figure 2. Molecular Subtypes and Differentially Expressed Genes in Ductal and Lobular Tumor Regions.**

**Figure 3. Distinct Biological Signatures in Ductal and Lobular Tumor Regions**

**Figure 4. Overview of Genomic Alterations in Ductal and Lobular Tumor Regions.**

**Figure 5: Single Cell Heterogeneity of Ductal and Lobular Tumor Populations.**

**Supplementary Figure S1: Digital Spatial Profiling RNAseq Dataset Quality Control.**

**Supplementary Figure S2: Patient Wise and Pan-Patient mDLC Ductal vs Lobular Differential Analysis.**

**Ductal Regulatory Network**

**Lobular Regulatory Network**

Enriched Kinases

Intermediate Protein

Enriched Transcription Factors

MYC MAX

ESR1 FOXA2 FOXA1

**Key Ductal Transcription Factors**

**Key Lobular Transcription Factors**

**Supplementary Figure S3: Transcription Factor Enrichment Analysis using Pan-Patient Ductal vs Lobular DEGs.**

**Supplementary Figure S4: Key Genes Associated with Notable Differentially Enriched Oncogenic and Biological Signatures.**

A

| Sample | | Purity | Depth | Ploidy |
|---|---|---|---|---|
| **MDLC-1** | Ductal | 0.36 | 428x | 2.02 |
| | Lobular | 0.45 | 523x | 1.97 |
| **MDLC-2** | Ductal | 0.51 | 416x | 2.04 |
| | Lobular | 0.31 | 339x | 2.02 |
| **MDLC-3** | Ductal | 0.62 | 378x | 3.35 |
| | Lobular | 0.4 | 490x | 2.01 |

C

B

**Supplementary Figure S5: MSK-IMPACT Extended Results.**

**Supplementary Figure S6: Effect of Q23 mutation on *CDH1* RNA and E-cadherin protein expression in ER+ Breast Tumors.**

**Supplementary Figure S7: MDLC-1 Ductal and Lobular Tumor scRNAseq Heterogeneity.**

**Supplementary Figure S8: Distinct Biological Signatures in MDLC-1 Ductal and Lobular Tumor scRNAseq.**