Spatial molecular profiling of mixed invasive ductal-lobular breast cancers reveals

heterogeneity in intrinsic molecular subtypes, oncogenic signatures and actionable driver

mutations

Osama Shiraz Shah[1,3], Azadeh Nasrazadani[1,4], Julia Foldi[7], Jennifer M. Atkinson[1,2], Celina G Kleer[8],

Priscilla F. McAuliffe[1,6,] Edaise M da Silva[9], Pier Selenica[9], Higinio Dopeso[9], Fresia Pareja[9], Diana

Mandelker[9], Britta Weigelt[9], Jorge S. Reis-Filho[9], Rohit Bhargava[4], Peter C. Lucas[1,4,5], Adrian V.

Lee*[1,2], Steffi Oesterreich*[1,2]

[1]Womens Cancer Research Center at UPMC Hillman Cancer Center and Magee Women's

Research Institute, Pittsburgh, PA.

[2]Department of Pharmacology & Chemical Biology, University of Pittsburgh, Pittsburgh, PA.

[3]Integrative Systems Biology Program, University of Pittsburgh, Pittsburgh PA.

[4]Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, PA

[5]NSABP Foundation, Inc., Pittsburgh, PA

[6]Division of Surgical Oncology, Department of Surgery, University of Pittsburgh School of

Medicine, Pittsburgh, PA

[7]Division of Hematology-Oncology, Department of Medicine, University of Pittsburgh,

Pittsburgh, PA.

[8]Department of Pathology and Rogel Cancer Center, University of Michigan Ann Arbor, MI

48109, USA

[9]Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center

(MSKCC), New York, NY, USA

Keywords: Mixed invasive ductal and lobular carcinoma, mDLC, digital spatial profiling, DSP,

breast cancer, tumor heterogeneity, multi-omics, precision medicine

* Steffi Oesterreich and Adrian Lee share senior authorship

Corresponding Authors: Steffi Oesterreich, University of Pittsburgh, 5051 Centre Ave,

Pittsburgh, PA 15213. Phone: 412-641-8555, Fax: 412-641-7724, E-mail:

oesterreichs@upmc.edu. Adrian Lee, University of Pittsburgh, 5051 Centre Ave, Pittsburgh, PA

15213. E-mail: leeav@upmc.edu.

Abstract [N=135]

Mixed invasive ductal and lobular carcinoma (mDLC) is a rare histologic subtype of breast cancer displaying both E-cadherin positive ductal and E-cadherin negative lobular morphologies within the same tumor, posing challenges with regard to anticipated clinical management. It remains unclear whether these distinct morphologies also have distinct biology and risk of recurrence. Our spatially-resolved transcriptomic, genomic, and single-cell profiling revealed clinically significant differences between ductal and lobular tumor regions including distinct intrinsic subtype heterogeneity (e.g., mDLC with TNBC/basal ductal and ER+/luminal lobular regions), distinct enrichment of senescence/dormancy and oncogenic (ER and MYC) signatures, genetic and epigenetic *CDH1* inactivation in lobular, but not ductal regions, and 4) single-cell ductal and lobular sub-populations with unique oncogenic signatures further highlighting intra-regional heterogeneity. Altogether, we demonstrated that the intra-tumoral morphological/histological heterogeneity within mDLC is underpinned by intrinsic subtype and oncogenic heterogeneity which may result in prognostic uncertainty and therapeutic dilemma.

Significance [N=58]

mDLC displays both ductal and lobular tumor regions. Our multi-omic profiling approach revealed that these morphologically distinct tumor regions harbor distinct intrinsic subtypes and oncogenic features that may cause prognostic uncertainty and therapeutic dilemma. Thus histopathological/molecular profiling of individual tumor regions may guide clinical decision making and benefit patients with mDLC, particularly in the advanced setting where there is increased reliance on next generation sequencing.

Main Text

Introduction

Breast cancer is a prevalent disease worldwide[1] and is the most frequently diagnosed cancer in women in the US, with approximately 300,000 cases per year[2]. The majority (approximately 75%) of cases are classified histologically as invasive ductal carcinoma of no special type (IDC-NST). Invasive lobular carcinoma (ILC) is the most common special type accounting for approximately 10-15% of all invasive breast cancer diagnoses[3,4]. Apart from the well-defined histologic and tissue histological differences, there are numerous notable molecular distinctions between ILC and IDC-NST. ILC exhibits enrichment of *CDH1*, *FOXA1*, *TBX3*, *ERBB2* and *PTEN* alterations, while IDC-NST shows a higher frequency of *GATA3*, *MAP3K1*, *TP53* and *MYC* alterations[5]. As a result of *CDH1* alterations, ILC lacks E-cadherin expression, an important cell adhesion molecule[6], and display cytoplasmic p120 staining while IDC-NST retains membranous expression for both molecules[7]. Clinically, ILC is more common in elderly women, presents as larger tumors and is frequently multicentric and multifocal compared to IDC-NST[8,9]. ILC is usually detected later and leads to higher rates of mastectomy [10–14]. ILC is also associated with more frequent metastases to urogenital and gastrointestinal tracts as well as to serosal surfaces than IDC-NST[15,16]. Despite presenting with markers of favorable prognosis, such as low grade, low proliferation index, estrogen receptor positive and human epidermal growth factor receptor (HER2) low/negative, patients with ILC have a higher rate of late recurrences and worse long-term prognosis than those with IDC-NST[14,17,18]. Given its distinct prognosis and outcome compared to IDC-NST, ILC is increasingly being recognized as a distinct disease, yet it is still treated similarly to IDC-NST in the

clinic due to a lack of specific treatment guidelines[14,19,20]. Ongoing clinical trials focused on

patients with ILC may lead to the development of more tailored and effective treatment options

in the future[20].

Apart from ILC, there are numerous other special types of breast cancers such as mixed invasive

breast carcinoma. This is an elusive group of breast cancers which the World Health Organization

(WHO) defines as tumors presenting with areas of 10-90% of special histology (such as ILC)

admixed with at least 10% of IDC-NST[21]. Mixed tumors comprising of both ILC and IDC-NST are

the most common (~5-10% of all breast cancer cases[22–27]) and are termed mixed invasive ductal

and lobular breast carcinoma (mDLC)[28]. Despite increasing incidence of mDLC[29,30], it remains

understudied, leading to diagnostic and prognostic uncertainty and clinical management

challenges for the patients with this elusive disease. Although there are limited reports focused

on mDLC, a few seminal studies have provided insights into its complex pathobiology.

Immunohistochemical (IHC) studies have confirmed while that ductal regions of mDLC retain

membranous E-cadherin, while lobular regions either lose or display aberrant E-cadherin IHC

staining[31–33]. Clinical studies have often yielded contradictory findings[23–25,30,31,33–35], but overall,

they suggest that mDLC is clinically more similar to pure ILC than IDC-NST[28,36]. These

discrepancies may be due to the non-standardized and fluid definition of mDLC, limited

consideration of its underlying tumor regions and their distinct features and proportions, and the

vast intratumoral heterogeneity of this disease[28,30,37,38]. Misdiagnosis is also possible, particularly

if only one of the two histologic regions is identified in the diagnostic biopsies[28,30,37,38]. Notably,

accurate diagnosis of mDLC remains a challenge[28,30,37,38] and traditional diagnostic methods, such

as mammography, may prove inadequate, as they are often less sensitive in detecting lobular

tumors owing to their diffuse growth pattern[13,14,39–41]. However, the use of magnetic resonance

imaging (MRI) and E-cadherin IHC assessment in biopsy specimens can improve diagnostic

accuracy[20,42,43]. Moreover, molecular studies with profiling of individual tumor regions have been

limited and largely focused on analysis of somatic genetic alterations. While these have provided

valuable insights into the tumorigenesis pathways and distinct mutational landscape of ductal

and lobular tumor regions[33,44], comprehensive studies delineating transcriptomic differences

between these distinct regions have not been performed.

To fill this knowledge gap, we employed spatially resolved transcriptomic, genomic, and single

cell profiling approaches to comprehensively investigate mDLC and highlight its intratumoral

heterogeneity and complexity. We delineate the biological and intrinsic subtype differences

between its underlying ductal and lobular tumor regions. Notably, our findings highlight that in

some mDLC cases distinct tumor regions with distinct biology may pose distinct progression risk.

Hence histopathological and molecular profiling mDLC and its individual tumor regions may

inform treatment decisions.

Results

Patient Specimens and Their Clinicopathologic Characteristics

A panel of three molecular pathologists identified invasive breast cancer cases diagnosed as "mixed invasive ductal and lobular carcinoma" and shortlisted three cases for this pilot study. All three cases were reported as estrogen receptor positive (ER+) and HER2 negative/low and their ages ranged from 46-63 years (Fig. 1A, Supplementary Table S1A). Assessment of H&E and E-cadherin/p120 dual staining confirmed that each case exhibited histomorphic features in line with the latest mDLC definition i.e., admix of E-cadherin negative lobular tumor regions with at least 10-90% of E-cadherin positive ductal tumor regions (Supplementary Table S1B). Importantly, overall, these cases exhibited collision of spatially distinct ductal and lobular tumor regions[68,78] along with some regions of intermixing and hence could be broadly defined as mDLC of "collision" type. These cases were particularly amenable for downstream multi-omic profiling compared to those that exhibited more intermingling or intermixing with close proximity of histologic tumor regions. Clinical diagnostic and treatment timelines of these cases, source of their tumor tissue used in this study, and their histomorphic features are illustrated in Fig. 1A. Both MDLC-2 and MDLC-3 eventually developed recurrent disease resulting in patient death in the case of MDLC-2. As of the last follow-up in 2022, both MDLC-3 and MDLC-1 are alive, and MDLC-1 remains free of recurrent disease.

Whole-Transcriptome Spatial Profiling of Ductal and Lobular Regions

To better understand the biological heterogeneity and transcriptomic features of ductal and lobular regions, we performed whole-transcriptome digital spatial profiling (DSP) of these three

mDLC cases. In total, we analyzed 26 regions of interest (ROIs) with at least 6 ROIs per case.

Selected ROIs from a representative case (MDLC-3) are shown in Fig. 1B. We segmented ductal

and lobular areas using E-cadherin and Pan-CK staining (Fig. 1C). This enabled profiling of pure

ductal and lobular areas from heterogeneous regions that were predominantly of one histology

or a mixture of both (Fig. 1D). E-cadherin expression in segmented ductal and lobular regions

showed expected results (Fig. 1E). All profiled regions had high RNA sequencing saturation (>

94%) and low noise (Q3 count > Negative probe count). However, three regions had low read

count (< 0.3 million) and were excluded from further analysis (Supplementary Fig. S1A-C). Our

final DSP dataset is comprised of 13 ductal and 13 lobular regions.

Transcriptomic Clusters and Intrinsic Molecular Subtypes

Exploration using top variable genes revealed that regions from each tumor clustered based upon

their histology (Fig. 2A; Supplementary Fig. S1D). MDLC-3 ductal regions showed the most distinct

mRNA expression compared to all other regions. PAM50 analysis showed both inter and intra-

tumor heterogeneity of prognostically relevant intrinsic molecular subtypes (IMS). MDLC-2 ductal

regions were a mix of Luminal A and HER2-Enriched subtypes while lobular regions were HER2-

Enriched (but stained negative for HER2 IHC – Supplementary Table S1A). MDLC-1 ductal and

lobular regions were both of luminal IMS, with the former being Luminal A and the latter being

Luminal B. Finally, in line with mRNA expression patterns, MDLC-3 ductal and lobular regions

showed stark IMS differences (former Luminal A and later Basal-Like). MDLC-3 ductal region was

the only TNBC/Basal-Like tumor (and had low *ESR1*, *PGR* and *ERBB2* mRNA expression –

Supplementary Fig. S1E). We further confirmed the TNBC status of MDLC-3 ductal regions using

ER and HER2 IHC (Supplementary Table S1A). Overall, our transcriptomic clustering and IMS

analysis suggest that ductal and lobular tumor regions within mDLC may have distinct

transcriptomes and intrinsic subtypes.

Differential Transcriptomic Features and Comparison with Pure ILC and IDC-NST

To investigate global transcriptomic differences between ductal and lobular regions across all

cases, we performed differential gene expression analysis in a pan-patient manner. To avoid

biases caused by fundamental differences between ER+ and ER- disease, we excluded MDLC-3

ER-/TNBC ductal regions from this analysis and later downstream analyses (Supplementary Fig.

S2C). In total, we identified 168 significantly up-regulated and 198 significantly down-regulated

differentially expressed genes (DEGs) in the lobular compared to ductal regions across all cases

(Fig. 2B). These pan-patient DEGs showed strong overlap with patient specific DEGs

(Supplementary Fig. S2A,B,C) indicating that our pan-patient analysis captured most if not all

strong global differences between ductal and lobular tumor regions.

To assess transcriptomic similarity of mDLC ductal and lobular regions to pure counterparts, i.e.,

IDC-NST and ILC, we compared mDLC ductal vs. lobular region DEGs with those from the

comparison of TCGA ER+ IDC-NST (N=327) vs ILC (N=113), respectively (Fig. 2C). We found *KLK11,*

KLK10, NRAP, SHROOM1, BTG2, and *PDK4* were upregulated, while *CDH1*, *DCD, SERPINA1*, and

CPB1 were downregulated in both comparisons (i.e., mDLC lobular vs ductal tumor regions and

ILC vs IDC-NST tumors). This overlap confirmed the well-known (i.e., down-regulation of *CDH1*)

markers of the lobular phenotype as well as potential novel markers.

Differential Enrichment of ER and MYC Signatures & Recurring Biological Themes

To understand the distinct biology of ductal and lobular regions, we scored each DSP region for

hallmark signatures[45]. Twenty signatures were differentially enriched between ductal and lobular

regions with p-value ≤ 0.05 (Fig. 2D, Supplementary Table S6). Findings for several signatures,

including Kras Signaling Up/Dn, Estrogen Response Early/Late, Tnfa Signaling, and Hypoxia in the

lobular regions, and Myc Targets V1, Protein Secretion and Adherens Junction in ductal regions,

were largely recapitulated in MDLC-1 scRNAseq data (Supplementary Fig. S6D). In line with

differential enrichment of Estrogen Response and Myc Target signatures, our transcription factor

(TF) signature analysis revealed enrichment of *ESR1* TF signatures in lobular and *MYC* TF

signatures in ductal regions (Supplementary Fig. S3). In summary, our findings show enrichment

of distinct oncogenic signatures in ductal vs. lobular regions.

To determine recurring biological themes, we expanded our differential enrichment analysis to

include all GO[46], REACTOME[47] and KEGG[48] genesets. Signatures differentially enriched between

ductal and lobular regions with p-value ≤ 0.05 were further assessed for robustness against

background permutations. Robust signatures were grouped into signature clusters based on

similarity of overlapping genes. This was done separately for signatures enriched in ductal and

lobular regions. All signature clusters were further grouped into broad biological themes based

on the underlying biological terms (Fig. 2E, left panel). The mean score of all signature clusters in

ductal and lobular regions is shown in Fig. 2E, right panel. Lobular regions were enriched for TP53

& senescence, collagen & extracellular matrix, MAPK/ERK targets (in line with KRAS signatures

shown in Fig. 2D) among other signatures. In contrast, ductal regions were enriched in double

stranded break & homologous repair, mTOR signaling (in line with Mtorc1/Mtor signatures

shown in Fig. 2D) among other signatures (Supplementary Table S7). Additionally, the expression

of senescence or cell cycle arrest genes (*CDKN1A/2B*) aligned with L2, collagen genes

(*COL4A1/4/5*) aligned with L5, and ERK/MAPK target genes (*DUSP4/6*) aligned with L9 signature

clusters were upregulated in lobular regions (Supplementary Fig. S4A,B,D). Enrichment of both

ERK/MAPK (L9) and KRAS signatures (from hallmark analysis) indicate higher activity of

KRAS/ERK/MAPK signaling in lobular regions, whereas enrichment of mTOR (D5) and

Mtorc1/Mtor signaling (from hallmark analysis) indicate higher activity of mTOR signaling in

ductal regions. In summary, we found differential enrichment of various cell cycle, immune,

collagen & ECM, metabolism, and oncogenic signatures between ductal vs. lobular regions.

Genomic Landscape and Clinically Relevant Genomic Alterations

To explore the genomic landscape of ductal and lobular regions, we employed the FDA-approved

targeted MSK-IMPACT panel. We micro-dissected the ductal and lobular tumor regions from FFPE

blocks of each case and sequenced them at a median depth of 416x (range 287–563)

(Supplementary Fig. S5A). In all cases, at least one pathogenic mutation was detected in known

breast cancer driver genes, including *CDH1, PIK3CA, TP53, GATA3, KMT2C, MAPK31, ARID1A,* and

CBFB, in both ductal and lobular regions, as depicted in Fig. 3A. Furthermore, all pathogenic

mutations had a high cancer cell fraction (Supplementary Fig. S5B).

Loss of E-cadherin is an established driver of lobular tumorigenesis[49]. Typically, it results from

biallelic *CDH1* inactivation, through a combination of truncating mutation and loss of

heterozygosity (LOH)[27]. All lobular regions exhibited loss of E-cadherin, with both MDLC-1 and

MDLC-3 harboring *CDH1* truncating mutations and LOH (Fig. 3B; Fig. 1E; Supplementary Fig. S1E).

The lobular region of MDLC-2 did not have *CDH1* genomic alterations but showed epigenetic

inactivation via methylation of its promoter. A putative passenger mutation in *CDH1* was also

observed in the MDLC-3 ductal region, which had no effect on mRNA or E-cadherin levels (Fig.

1E; Supplementary Fig. S1E). The loss of E-cadherin is consistent with the downstream down-

regulation of Adherens Junction signatures in all lobular regions (Fig. 2D). Hotspot *PIK3CA*

mutations, which are early oncogenic driver events frequently observed in ER+ BC[50], were

present in all cases except MDLC-3. The MDLC-1 ductal region had mutations in *GATA3* (critical

transcriptional binding partner of ERalpha[51]) and *CBFB* (forms complex with *RUNX1* to suppress

NOTCH signaling[52]). The MDLC-3 ductal region had loss of *TP53*, an event associated with more

aggressive types of breast cancer like basal/TNBC[53] and consistent with its basal-like intrinsic

subtype. MDLC-2 ductal region had truncating mutation in *KMT2C*, an important regulator of ER

alpha activity[54]. MDLC-3 lobular region had mutations in *ATRX* and *ARID1A,* two chromatin

remodeling and DNA repair genes[55]. In addition, while MDLC-1 and MDLC-3 ductal and lobular

regions did not share any somatic clonal mutations, MDLC-2 ductal and lobular regions shared

several somatic clonal mutations in *PIK3CA* and *MAPK31* genes indicating shared tumor origin.

To investigate the copy number (CN) landscape of ductal and lobular regions, we performed CN

analysis using FACETS[56] (Fig. 3B). All regions exhibited breast cancer characteristic chromosome

1q gains and 16q losses[53]. Overall, ductal tumor regions showed more frequent aneuploidy

events than lobular tumor regions. MDLC-3 ductal region showed the most aberrant CN profile

consistent with its basal/TNBC etiology[53]. Moreover, chromosome 22 aneuploidy was only

observed in ductal tumor regions. Notably, both histologic regions of MDLC-3 and ductal region

of MDLC-2 had *NOTCH2* gains or amplifications (Supplementary Fig. S5C).

Mutational Processes Driving Tumor Evolution

To determine the mutational processes underlying the evolution of ductal and lobular tumor

regions within mDLC, we performed mutational signature analysis of all cases (Fig. 3C). Both

histologic regions in MDLC-2 were associated with an "aging" signature. In MDLC-3, ductal region

was associated with homologous repair deficiency (HDR) while the lobular regions with APOBEC

signature. Tumors with an APOBEC signatures have a higher frequency of mutations in chromatin

remodeling and DNA repair genes[57], as seen in MDLC-3 lobular region. We did not identify any

signatures for MDLC-1 ductal and lobular regions. The putative evolution of each mDLC case and

its histologic region based upon the mutations, *CDH1* alterations and mutational signatures is

shown in Fig. 3D. Notably, the MDLC-2 ductal and lobular regions not only shared clonal

pathogenic (*PIK3CA*) and passenger (*MST1R* and *MAP3K1*) mutations but also the "aging"

mutational signature. Loss of function *CDH1* alterations were hallmark of all lobular regions

across cases.

Single-Cell RNA-Sequencing and Identification of Ductal and Lobular Populations

To explore the heterogeneity of mDLC at a single-cell level, we performed scRNAseq sequencing

of a fresh tumor specimen from MDLC-1. The resulting scRNAseq comprised of 4,671 cells and

had sequencing saturation, reads in cell and median genes per cell as shown in Supplementary

Fig. S6A. Clustering revealed two transcriptionally distinct tumor populations; however, we were

unable to annotate these populations as ductal and lobular tumors using *CDH1* mRNA (Fig. 4A).

Notably, while the lobular regions showed loss of E-cadherin protein (Fig. 1G), the *CDH1* p.Q23*

mutation in MDLC-1 (Fig. 3A) didn't alter *CDH1* mRNA levels (Supplementary Fig. S1E). Therefore,

we derived ductal and lobular transcriptome signatures (Supplementary Table S5) from MDLC-1

DSP dataset and used these signatures to score each single cell in MDLC-1 scRNAseq dataset.

Using this deconvolution approach, we successfully annotated ductal and lobular tumor

populations in the scRNAseq dataset (Fig. 4B). There was significant overlap of ductal vs. lobular

DEGs between scRNAseq and DSP datasets (Supplementary Fig. S6B,C). Differential enrichment

of several hallmark signatures and recurring biological themes in ductal and lobular regions from

DSP data was recapitulated in our scRNAseq data (Supplementary Fig. S7A,C). Notably, we saw

enrichment of Estrogen and senescence signatures in lobular and Myc signature in ductal tumor

cells in both DSP and scRNAseq datasets.

Single-Cell Heterogeneity and Mapping of Single-Cells to Spatial Regions

To investigate single-cell heterogeneity of ductal and lobular populations, we increased

resolution of clustering to identify sub-clusters. Each sub-cluster had unique marker genes (Fig.

4C) and several of these also showed differential enrichment of hallmark signatures and recurring

biological theme clusters (Supplementary Fig. S7B,D), indicating intra-tumor heterogeneity

within ductal and lobular tumor populations. To investigate whether single-cell transcriptomic

heterogeneity was associated with spatial heterogeneity, we mapped scRNAseq sub-clusters to

DSP regions. To assess similarity between each region and sub-cluster, we scored all regions for

sub-cluster marker genesets using GSVA (Fig. 4D). Each region was mapped to a sub-cluster for

which it had most similarity (i.e., highest GSVA score) (Fig. 4E). The top five marker genes of each

scRNAseq sub-cluster (Supplementary Fig. S6D) are shown beside each sub-cluster and spatial

region link. These findings perhaps suggest that the single-cells in each sub-cluster came from a

spatially distinct niche in the tumor.

Discussion

Despite an increasing incidence of mDLC[29,30], this special subtype of breast cancer remains poorly understood, with limited knowledge about its overall biology and that of its underlying ductal and lobular tumor regions, as well as their clinical implications.

Spatially resolved and single-cell profiling techniques have revolutionized our understanding of physiological and pathological processes and tissue architecture[58]. Efforts such as Human Cell Atlas (HCA)[59] and those from other groups have successfully uncovered the major cell types and their spatial organization in both normal breast tissue[60,61] and breast cancer[62]. Spatial and single-cell profiling of primary breast cancer subtypes and metastatic disease have been instrumental in identifying novel molecular features and their prognostic value[63–67]. However, to date genome-wide spatial and single-cell investigation of mDLC has not been reported. To address this gap, we employed a state-of-the-art methodology, comprising of spatially resolved transcriptomics, genomics, and single-cell profiling to investigate the biology of individual tumor regions within mDLC of "collision" type. Our study confirmed various previously reported observations of genetic inactivation of *CDH1* and concomitant loss of E-cadherin in lobular regions[31–33,68] and the presence of distinct driver alterations in individual tumor regions[33,68]. Moreover, we also uncovered novel findings that highlight the vast heterogeneity within mDLC, several of which were indicative of distinct prognosis. Ductal and lobular tumor regions had distinct biological signatures, in particular senescence, ER and *MYC* signaling, and intrinsic molecular subtypes (IMS). These distinctions were most prominent in sample MDLC-3, which was clinically reported as ER+ tumor but molecular profiling revealed that the underlying ductal region was ER-/TNBC

with basal-like IMS while the lobular region was ER+ with luminal-A IMS. Notably, both ER- status

and basal-like IMS are associated with worse prognosis and outcome than ER+ and luminal

IMS[69,70]. This patient had brain and lung metastasis, both of which were also ER-, possibly

indicating progression of the primary ER- ductal tumor. This observation is clinically significant,

indicating that individual tumor regions within mDLC may have distinct clinical and biological

features and that one may pose higher recurrence risk than the other.

Another notable finding was enrichment of senescence and ER signaling signatures in lobular

while *MYC* signatures in ductal tumor regions. Further analysis at single-cell level revealed that

these signatures were enriched in only one lobular subcluster (L2), highlighting molecular

heterogeneity even within individual histologically distinct tumor regions. Most patients with ILC

are strongly ER+ and respond well to endocrine therapies, while they are less proliferative and

respond poorly to chemotherapy[14,20,27,71,72]. Whether this is also true for lobular regions within

mDLC is an important question and must be further investigated. Perhaps, future trials can study

whether patients with predominantly ductal vs. lobular mDLC respond differently to standard of

care endocrine and chemotherapies. However, this will be difficult to do as frequency of mDLC

cases is low and a specific trial dedicated to patients with this disease may not be feasible.

The presence of distinct clinically actionable alterations in individual tumor regions within mDLC

is a clinically important observation. We show that these alterations may emerge via common

(Aging in MDLC-2) or distinct (APOBEC or HRD in MDLC-3) mutational processes. Tumor regions

with distinct alterations can possibly complicate patient prognosis and clinical management.

Alterations in components of ER signaling (such as *GATA3, KMT2C*) as well as Notch signaling

(*CBFB*, *NOTCH2*) can lead to endocrine resistance[73–75], while alterations in DNA repair response

genes (*ARID1A*, *ATRX*) may confer increased sensitivity to chemotherapy and other DNA repair

pathway targeting drugs including PARP inhibitors[76,77]. Treatment regimens decisions for patients

with mDLC can greatly benefit from considering the distinct alterations in individual tumor

regions to avoid drug resistance by rational selection of therapies.

For a long time, it was unclear whether ductal and lobular regions within mDLC share any

molecular similarity with their pure counterparts (i.e., ILC and IDC-NST tumors). In our study, we

compared the molecular features of individual tumor regions within mDLC with their pure

counterparts in the TCGA cohort[27] and discovered a set of shared markers of lobular phenotype.

One of these marker genes was *CDH1*, whose inactivation drives lobular tumorigenesis[49]. This

primarily occurs via genetic alterations[27] which was the case in lobular regions of MDLC-1 and

MDLC-3. However, in lobular regions of MDLC-2, *CDH1* promoter methylation was observed,

indicating an alternative but less studied route of lobular tumor formation via epigenetic

inactivation. This is an intriguing observation, suggesting that the pathogenesis of lobular regions

within mDLC maybe more complex and diverse than previously thought[6,68,78–80]. Our expression

analysis of various previously reported[81] epithelial to mesenchymal (EMT) markers

(Supplementary Fig. S1F) excluded the role of EMT in driving this event.

A recent study revealed the synthetic lethality in E-cadherin deficient breast cancers with

inhibition of *ROS1*, a tyrosine kinase[82]. The ongoing ROSALINE trial is investigating efficacy of

ROS1 inhibitors in early setting of ER+ ILC disease. If successful, future studies can assess the

value of *ROS1* inhibitors in treating patients with predominantly lobular mDLC disease. Additional

studies are also warranted to investigate whether E-cadherin deficient tumors with distinct

etiology, such as *CDH1* epigenetic or genetic inactivation, have distinct biology potentially

associated with differential treatment response.

Although our study presents a unique and novel analysis of mDLC of "collision" type, it has some

limitations, including small sample size and a lack of analysis of other subtypes (such as lobular-

like IDC-NST[83] and intermixed/intermingled mDLC[68]) and tumor microenvironment of individual

tumor regions within mDLC. Future studies will be necessary to validate our observations in larger

and more diverse cohorts of mDLC. Generating a comprehensive mDLC cohort for future studies

may not be an easy task, given the challenges associated with the diagnosis of this mixed entity.

However, the use of E-cadherin IHC in histological assessments and evaluation by a panel of

pathologists can ensure the generation of a high-confidence and well-annotated mDLC cohort

for future studies. Moreover, the application of digital pathology and artificial intelligence will

make this more feasible[84]. These tools will not only improve diagnostic accuracy[85] but will also

allow scalable annotation of ductal and lobular tumor regions across heterogenous tumor

samples[86].

In summary, our study provides new insights into the molecular heterogeneity of mDLC and its

underlying ductal and lobular tumor regions. This is the first report on genome-wide spatially

resolved and single-cell molecular profiling of mDLC. We discovered distinct biological signatures

and intrinsic molecular subtypes between these regions that may have important implications

for patient prognosis and clinical management. Our findings also highlight the need for

comprehensive histopathological and molecular profiling of individual tumor regions within

mDLC to better understand its progression and response to standard of care hormone and

cytotoxic therapies.

Methods

Sample Collection

Institutional review board approval was obtained from the University of Pittsburgh prior to initiation of the study. Patients diagnosed with "mixed invasive ductal and lobular" were identified from the UPMC Breast Cancer Registry and Formalin-Fixed Paraffin-Embedded (FFPE) blocks requested from Pitt Biospecimen Core. Four-um-thick FFPE sections were used for H&E, ER-alpha and E-cadherin IHC staining. A panel of pathologists reviewed the H&E and E-cadherin staining to confirm mDLC diagnosis in line with recent WHO guidelines (10-90% of ILC and at least 10% IDC-NST). For our pilot study, we only selected mDLC cases amenable for multi-omic profiling i.e., those with good separation of ductal and lobular tumor regions to avoid spill over and contamination of individual ductal and lobular tumor regions by neighboring lobular and ductal tumor cells, respectively. Hence, we selected three "collision" type mDLC cases that showed both clear separation of ductal and lobular tumor regions along with some intermixed regions. These included cases MDLC-1, MDLC-2 & MDLC-3 (clinical features and pathology review notes are shown in Supplementary Table S1A & S1B). All cases underwent NanoString digital spatial, MSK-IMPACT mutation and *CDH1* promoter methylation profiling. Fresh tumor was only available for MDLC-1 and was utilized for single-cell RNAseq. See details in Supplementary Methods.

Digital Spatial Profiling (DSP)

DSP was performed by NanoString GeoMX as described previously[87]. Briefly, five-μm-thick FFPE tumor sections were mounted on single positively charged slide. Slides underwent deparaffinization, rehydration, heat-induced epitope retrieval, and enzymatic digestion.

Oligonucleotide probe mix from whole-transcriptome atlas (WTA) panel were hybridized and

incubated overnight. The slide was blocked and incubated with fluorescent antibodies against

morphology markers and DNA dye (Syto83 dye; Invitrogen Cat # S11364) for 1 hour. Morphology

markers included E-cadherin (BD Pharmingen Cat# 560062), Pan-cytokeratin (Novus Cat# NBP2-

33200AF488) and Vimentin (Santa Cruz Cat# sc-373717 AF594). Finally, the slide was loaded into

the GeoMx platform and scanned for an immunofluorescent signal. Together with the aid of H&E

and morphology marker stains, our panel of pathologist selected 26 rectangle ROIs (12 lobular

ROIs, 11 ductal ROIs and 3 mixed ROIs) with an area of 518,100 μm^2. To profile ductal and lobular

cells separately the following segmentation approach was used; 1) ductal cells a E-cadherin +ive

& Pan- cytokeratin +ive, and 2) lobular cells a E-cadherin -ive & Pan- cytokeratin +ive. Above

segmentation resulted in total of 29 areas (14 ductal and 15 lobular). UV light was directed at

each segmented region to release the oligonucleotides that were collected and then sequenced

using Illumina NextSeq 2000. The resulting FASTQ files were processed using GeoMX NGS

pipeline 2.3.4 which provided the count data for all target probe across profiled regions. All 29

profiled regions had high sequencing saturation (> 94%) and low signal-to-noise ratio (Q3

count/NegProbe count ≥ 1) (Supplementary Fig. S1A-C; Supplementary Table S2,S3). Further

downstream analysis is described in the Supplementary Methods.

MSK-IMPACT Targeted Next Generation Sequencing

Eight-μm-thick sections from representative FFPE tumor and matched normal tissue sections

from MDLC-3, MDLC-2 & MDLC-1 were stained with nuclear fast red and microdissected using a

sterile needle under a stereomicroscope (Olympus SZ61) to enrich tumor content, as previously

described[88,89]. Histologically distinct tumor components were microdissected separately.

Genomic DNA of each component and of matched normal tissue was extracted using the DNeasy

Blood and Tissue Kit (Qiagen) according to manufacturers' instructions. Tumor and normal DNA

from each case were subjected to massively parallel sequencing using the Food and Drug

Administration (FDA) approved Memorial Sloan Kettering Integrated Mutation Profiling of

Actionable Cancer Targets (MSK-IMPACT) multigene panel targeting all coding regions of 515

cancer-related genes as previously described[90–92]. Somatic alterations were identified as

previously described[93,94] (Supplementary Table S8). The median depth of coverage of tumor and

normal samples was 416x (range 287–563) and 210x (range 175-217), respectively

(Supplementary Fig. S5A). In brief, reads were aligned to the reference human genome GRCh37

using the BWA v0.7.15[95]. Local realignment, duplicate removal and base quality recalibration

were performed using the GATK. V3.1.1[96]. Further downstream analysis is described in the

Supplementary Methods.

Figures & Illustrations

424 DSP spatial images were exported from Most bioinformatic analysis visualizations were

425 generated using R. These visualizations were aesthetically improved in power point to produce

426 the final figures. Illustrations were prepared using BioRender.com.

Data Availability

Data generated in this study is shared and can be included in the Supplementary Data and GitHub.

R code for data analysis can be accessed at

https://github.com/osamashiraz/mDLC_Spatial_Analysis.

Authors' Disclosures

PCL has an equity interest in Amgen (outside the submitted work). No disclosures were reported

by other authors.

Authors' Contributions

O.S. Shah: Conceptualization, data curation, software, formal analysis, investigation, visualization, methodology, project administration, writing–original draft, writing–review and editing. **A. Nasrazadani:** resources, funding acquisition, writing-review and editing. **J.M. Atkinson:** resources, supervision, project administration, writing–review and editing. **J. Foldi:** resources, investigation, data curation, writing–review and editing. **C.G. Kleer:** investigation, data curation, writing–review and editing. **P.F. McAuliffe:** investigation, resources, data curation, writing–review and editing. **E.M. da Silva:** investigation, resources, data curation, software, formal analysis, writing–review and editing. **P. Selenica:** investigation, resources, data curation, software, formal analysis, writing–review and editing. **H. Dopeso:** investigation, resources, data curation, writing–review and editing. **F. Pareja:** investigation, resources, data curation, writing–review and editing. **D. Mandelker:** investigation, resources, data curation, writing–review and editing. **B. Weigelt:** conceptualization, investigation, methodology, resources, writing–review and editing. **J.S. Reis-Filho:** conceptualization, investigation, methodology, resources, writing–review and editing. **R. Bhargava:** conceptualization, investigation, resources, data curation, methodology, data curation, writing–review and editing. **P.C. Lucas:** conceptualization, investigation, resources, data curation, methodology, data curation, writing–review and editing. **A.V. Lee:** conceptualization, resources, methodology, supervision, funding acquisition, project administration, writing–review and editing. **S. Oesterreich:** conceptualization, resources, methodology, supervision, funding acquisition, project administration, writing–review and editing.

Acknowledgments

We would also like to thank NanoString DSP Technology Access Program for providing early

access to their whole transcriptome spatial technology, Pitt Biospecimen Core at University of

Pittsburgh for providing access to valuable clinical specimens and technical support, Pitt

Genomics Research Core and UPMC Genome Center for technical support in preparing scRNAseq

libraries and sequencing, and Ms. Morgan DeBerry, HTL(ASCP)CM, QIHCCM for excellent

technical assistance in tissue sectioning, staining and other logistics. This work was primarily

supported by Dynami and BCRF grants to **S. Oesterreich** and in part by various other grants

including the ASCO Gianni Bonadonna Breast Cancer Research Fellowship grant to **A.**

Nasrazadani and Conquer Cancer ASCO Young Investigator Award to **J. Foldi**.

 Figure Legends

Figure 1. Digital Spatial Profiling of mDLC Ductal and Lobular Tumor Regions. (A) Top panel

shows clinical timelines of the studied cases including the time of core biopsies, surgical

interventions, treatments, metastasis and last known vital status. The source of the tumor tissue

used in this study is highlighted with * in red color. The bottom panels show high (40X)

magnification H&E and Ecad/p120 IHC images highlighting histomorphic and E-cadherin features

of ductal and lobular regions across each case. (B) E-cadherin and Pan-CK immunofluorescence

staining of a representative case (MDLC-3). Selected regions of interests (ROIs) are shown with

dotted blue (for ductal) and red (for lobular). (C) Segmentation of ductal and lobular cells using

E-cadherin and Pan-CK staining masks. (D) Segmented areas containing only ductal or lobular

tumor cells defined using ductal and lobular masks illustrated in panel C. (E) E-cadherin staining

intensity in representative ductal and lobular regions profiled across cases.

Figure 2. Molecular Subtypes and Distinct Biological Signatures of Ductal and Lobular Tumor

Regions. (A) Profiled ductal and lobular tumor regions separated by consensus clusters. All

regions are annotated by patients, histology, and consensus clusters. Gene expression heatmap

shows top variable genes and followed by PAM50 scores for each subtype across ductal and

lobular regions. (B) Differentially expressed genes (DEGs) between ER+ ductal and lobular

regions. The top panel shows DEG heatmap, while bottom panel shows volcano plot along with

number of significantly upregulated (N=168) and downregulated (N=198) DEGs in lobular vs

ductal regions. (C) Overlap of ductal and lobular region DEGs with TCGA ILC vs IDC-NST DEGs. (D)

Differentially enriched cancer hallmark pathways. The heatmap shows GSVA scores of each

differentially enriched pathway in ductal vs lobular tumor regions. Pathways with bold red and

blue names are the ones that also were significant in the scRNAseq dataset (Supplementary Fig.

S7A). Enrichment score (ES) for a pathway shows mean differences of GSVA score for that

pathway in lobular – ductal regions and -log10 of FDR-adjusted P-value (P-val) shows statistical

significance of enrichment based on T test. (E) Recurrent biological themes in ductal and lobular

tumor regions. The left panel shows the Jaccard similarity between all signatures. Signatures are

grouped based on their Jaccard similarity into signature clusters. Each cluster is associated with

a broad biological theme (shown in middle). The right panel shows the mean GSVA score of each

signature cluster in ductal and lobular regions.

Figure 3. Overview of Genomic Alterations in Ductal and Lobular Tumor Regions. (A) Somatic

mutations identified using MSK-IMPACT panel in ductal and lobular components across samples.

Gene names are followed by amino acid change due to mutation. Genes in black font had somatic

mutations in breast cancer driver genes while those in gray color had passenger/non-driver

mutations. (B) Somatic copy number changes in ductal and lobular components across samples.

For each case, genome-wide ductal and lobular copy number profiles are shown. Black line shows

total copy number, red line shows minor allele copy number and off-white regions show

aneuploidy (change in minor allele copy number including LOH). (C) Mutational signatures in

MDLC-2 and MDCL-3 ductal and lobular tumor regions. Pie charts show percentage similarity of

mutational patterns of each tumor component with established mutational patterns (associated

with different means/definition). (D) Predicted disease evolution. Tumors with overlapping

mutations were linked to common root node indicating common progenitor or precursor

neoplasm (N). Those without common root node likely indicate evolution from independent

precursors. Key mutations and events (CDH1 promoter methylation and amplifications) are

shown on the branches.

Figure 4: Single Cell Heterogeneity of Ductal and Lobular Tumor Populations. (A) Tumor

populations in MDLC-1 scRNAseq dataset (left) and assessment of CDH1 expression in MDLC-1

scRNAseq dataset (middle and right plot). Statistical significance was evaluated using the T test.

(B) Deconvolution of scRNAseq using DSP derived ductal and lobular signatures. GSVA scores for

ductal and lobular signatures are shown on the right panel. (C) Sub-clustering and identification

of ductal and lobular sub-populations (left) with unique marker genes (right). (D) Scoring spatial

regions using GSVA of sub-cluster marker genes. (E) Mapping of single cell populations to spatial

regions based on GSVA scores in D.

 Supplementary Figure Legends

Supplementary Figure S1: Digital Spatial Profiling RNAseq Dataset Quality Control. (A) Scatter

plot showing sequencing saturation (y-axis) and area of interest (AOI) surface area (mm2) across

cases. Dashed red lines show the quality control (QC) cutoffs (sequencing saturation >50% and

area > 6000mm2) for selecting high quality samples. (B) Overview of signal (Q3-normalized

counts) to noise (negative probe counts). Dashed red line shows cutoff i.e., signal > noise. (C)

Evaluation of Q3 normalized counts (of 1000 randomly sampled genes) across profiled regions to

identify samples with low quality data showing count artifacts. (D) tSNE plot showing clustering

of profiled samples that passed QC cutoffs. (E) CDH1, ESR1, ERBB2 and PGR gene expression in

ductal vs lobular regions across the three profiled cases. Statistical significance was evaluated

using the T test. (F) Expression of EMT marker genes in ductal and lobular tumor regions across

each case. EMT marker genes were adapted from McCart Reed et al [33].

Supplementary Figure S2: Patient Wise Differential Analysis and Comparison with Pan-Patient

Differential Analysis. Differentially expressed genes (DEGs) identified in ductal vs lobular in each

patient; MDLC-1 (A), MDLC-2 (B) and MDLC-3 (C) and overlap of patient-wise DEGs with pan-

patient (all patients combined) DEGs. Statistical significance of overlap between patient DEGs

and pan-patient DEGs was test using overlap_stats tool

(http://nemates.org/MA/progs/overlap_stats.html). A representation score > 1 and p < 0.05

shows significantly strong overlap.

Supplementary Figure S3: Transcription Factor Enrichment Analysis using Pan-Patient Lobular

vs Ductal DEGs. Lobular vs Ductal DEGs were separately input into https://maayanlab.cloud/X2K/

web portal to identify enriched transcription factors and kinases in lobular (left) and ductal (right)

tumor regions. The top layer with blue nodes shows kinases, middle layer shows intermediate

proteins and bottom layer with red nodes shows transcription factors. An image inset is shown

highlighting MYC and ER signaling related transcription factors enriched in ductal and lobular

tumor regions, respectively.

Supplementary Figure S4: Genes Associated with Key Differentially Enriched Biological

Signatures. Select differentially expressed genes in various differentially enriched biological

signature clusters between ductal vs. lobular tumor regions including (A) L2: TP53/Senescence,

(B) L5: Collagen & ECM, (C) L7: Collagen metabolism, (D) L9: MAPK/ERK Targets, (E) D5: mTOR,

(F) D4: Oxidative Stress, (G) D6: Acetyltransferase, (H) D7: Protein Maturation and (I) D8:

Phosphatidylinositol. Statistical significance was evaluated using the T test.

Supplementary Figure S5: MSK-IMPACT Extended Results. (A) Tumor purity, sequencing

coverage and ploidy of ductal and lobualr tumor regions across profiled cases. (B) Somatic

mutations and their cancer cell fraction (CCF). High CCF indicate clonal mutations while low CCF

indicates sub-clonal mutations. (C) Copy number of key breast cancer genes in ductal vs lobular

regions across each case.

Supplementary Figure S6: Single-cell RNAseq DEG and Pathway Analysis of MDLC-1. (A) MDLC-

1 single-cell RNAseq (scRNAseq) dataset quality control meterics from cell ranger software. (B)

MDLC-1 scRNAseq lobular vs ductal DEG analysis. (C) Overlap of MDLC-1 lobular vs ductal DEGs

between scRNseq and digital spatial profiling (DSP) datasets. Statistical significance of DEG

overlap   between   two   technologies   was   test   using   overlap_stats   tool

(http://nemates.org/MA/progs/overlap_stats.html). A representation score > 1 and p < 0.05

shows significantly strong overlap. (D) Top five marker genes of each lobular and ductal sub-

cluster identified in Fig. 4C.

Supplementary Figure S7: MDLC-1 Single-cell RNAseq Pathway Analysis. (A) GSVA based

hallmark signature enrichment analysis in scRNAseq dataset. Enrichment score (ES) for a pathway

shows mean differences of GSVA score for that pathway in lobular – ductal regions and -log10 of

FDR-adjusted P-value (P-val) shows statistical significance of enrichment based on T test. (B)

Hallmark signature enrichment in ductal and lobular subclusters. (C) Recurrening biology themes

in ductal vs lobular cells in scRNAseq dataset. (D) Mean GSVA score for select differentially

enriched biology themes across sub-clusters.

References

1.   Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and
mortality worldwide for 36 cancers in 185 countries. **68**, 394–424 (2018).

2.   Siegel, R. L., Miller, K. D., Wagle, N. S. & Jemal, A. Cancer statistics, 2023. *CA Cancer J Clin*
73, 17–48 (2023).

3.   Sledge, G. W., Chagpar, A. & Perou, C. Collective wisdom: lobular carcinoma of the breast.
Am Soc Clin Oncol Educ B **35**, 18–21 (2016).

4.   Reed, M. E. M. C., Kutasovic, J. R., Lakhani, S. R. & Simpson, P. T. Invasive lobular carcinoma
of the breast: morphology, biomarkers and 'omics. **17**, 12 (2015).

5.   Ciriello, G. *et al.* Comprehensive molecular portraits of invasive lobular breast cancer. *Cell*
163, 506–519 (2015).

6.   Reed, M. E. M. C., Kutasovic, J. R., Lakhani, S. R. & Simpson, P. T. Invasive lobular carcinoma
of the breast: morphology, biomarkers and 'omics. **17**, 12 (2015).

7.   Dabbs, D. J., Bhargava, R. & Chivukula, M. Lobular versus ductal breast neoplasms: the
diagnostic utility of p120 catenin. *Am J Surg Pathol* **31**, 427–37 (2007).

8.   Arpino, G. *et al.* Infiltrating lobular carcinoma of the breast: tumor characteristics and
clinical outcome. **6**, (2004).

9.   Oesterreich, S. *et al.* Clinicopathological Features and Outcomes Comparing Patients With
Invasive Ductal and Lobular Breast Cancer. *J Natl Cancer Inst* **114**, 1511–1522 (2022).

10.   Z, C. *et al.* Invasive lobular carcinoma of the breast: A special histological type compared
with invasive ductal carcinoma. **12**, e0182397 (2017).

11.   Yeatman, T. J. *et al.* Tumor biology of infiltrating lobular carcinoma. Implications for
management. *Ann Surg* **222**, 549–559 (1995).

12.   Helvie, M. A., Paramagul, C., Anderson, K. L. & Adler, D. D. Invasive lobular carcinoma.
Imaging features and clinical detection. *Invest Radiol* **28**, 202–207 (1993).

13.   Arpino, G. *et al.* Infiltrating lobular carcinoma of the breast: tumor characteristics and
clinical outcome. **6**, (2004).

14.   Oesterreich, S. *et al.* Clinicopathological Features and Outcomes Comparing Patients With
Invasive Ductal and Lobular Breast Cancer. *J Natl Cancer Inst* **114**, 1511–1522 (2022).

15.   Mathew, A. *et al.* Distinct Pattern of Metastases in Patients with Invasive Lobular
Carcinoma of the Breast. *Geburtshilfe Frauenheilkd* (2017) doi:10.1055/s-0043-109374.

16.   Lamovec, J. & Bračkko, M. Metastatic pattern of infiltrating lobular carcinoma of the
breast: an autopsy study. **48**, 28–33 (1991).

17.   Djerroudi, L., Cabel, L., Bidard, F. C. & Vincent-Salomon, A. Invasive Lobular Carcinoma of
the Breast: Toward Tailoring Therapy? **114**, 1434–1436 (2022).

18. K, V. B. *et al.* Current and future diagnostic and treatment strategies for patients with invasive lobular breast cancer. *Annals of Oncology* **33**, 769–785 (2022).

19. Luveta, J., Parks, R. M., Heery, D. M., Cheung, K.-L. & Johnston, S. J. Invasive Lobular Breast Cancer as a Distinct Disease: Implications for Therapeutic Strategy. *Oncol Ther* **8**, 1–11 (2020).

20. K, V. B. *et al.* Current and future diagnostic and treatment strategies for patients with invasive lobular breast cancer. *Annals of Oncology* **33**, 769–785 (2022).

21. The 2019 World Health Organization classification of tumours of the breast - Tan - 2020 - Histopathology - Wiley Online Library.

22. Metzger-Filho, O. *et al. The diagnosis of mixed invasive ductal and lobular carcinoma (IDC-L) in clinical practice is often associated with uncertainty related to its prognosis and response to systemic therapies. With the increasing recognition of invasive lobular carc*. vol. 24 (Oxford University Press (OUP), 2019).

23. Zengel, B. *et al.* Comparison of the clinicopathological features of invasive ductal, invasive lobular, and mixed (invasive ductal + invasive lobular) carcinoma of the breast. *Breast Cancer* (2015) doi:10.1007/s12282-013-0489-8.

24. Xiao, Y. *et al.* Mixed invasive ductal and lobular carcinoma has distinct clinical features and predicts worse prognosis when stratified by estrogen receptor status. *Sci Rep* (2017) doi:10.1038/s41598-017-10789-x.

25. Duraker, N., Hot, S., Akan, A. & Ozay Nayir, P. A Comparison of the Clinicopathological Features, Metastasis Sites and Survival Outcomes of Invasive Lobular, Invasive Ductal and Mixed Invasive Ductal and Lobular Breast Carcinoma. *Eur J Breast Health* **16**, 22–31 (2020).

26. Nasrazadani, A. *et al.* Mixed invasive ductal lobular carcinoma is clinically and pathologically more similar to invasive lobular than ductal carcinoma. *British Journal of Cancer 2023* 1–10 (2023) doi:10.1038/s41416-022-02131-8.

27. Ciriello, G. *et al.* Comprehensive molecular portraits of invasive lobular breast cancer. *Cell* **163**, 506–519 (2015).

28. Nasrazadani, A. *et al.* Mixed invasive ductal lobular carcinoma is clinically and pathologically more similar to invasive lobular than ductal carcinoma. *British Journal of Cancer 2023* 1–10 (2023) doi:10.1038/s41416-022-02131-8.

29. Li, C. I., Anderson, B. O., Daling, J. R. & Moe, R. E. Trends in Incidence Rates of Invasive Lobular and Ductal Breast Carcinoma. *J Am Med Assoc* (2003) doi:10.1001/jama.289.11.1421.

30. Suryadevara, A., Paruchuri, L. P., Banisaeed, N., Dunnington, G. & Rao, K. A. The clinical behavior of mixed ductal/lobular carcinoma of the breast: A clinicopathologic analysis. *World J Surg Oncol* (2010) doi:10.1186/1477-7819-8-51.

31. Rakha, E. A. *et al.* The biological and clinical characteristics of breast carcinoma with mixed ductal and lobular morphology. *Breast Cancer Res Treat* (2009) doi:10.1007/s10549-008-0007-4.

32. Acs, G., Lawton, T. J., Rebbeck, T. R., LiVolsi, V. A. & Zhang, P. J. Differential Expression of
E-Cadherin in Lobular and Ductal Neoplasms of the Breast and Its Biologic and Diagnostic
Implications. *Am J Clin Pathol* (2001) doi:10.1309/FDHX-L92R-BATQ-2GE0.

33. McCart Reed, A. E. *et al.* Mixed ductal-lobular carcinomas: evidence for progression from
ductal to lobular morphology. *Journal of Pathology* (2018) doi:10.1002/path.5040.

34. Metzger-Filho, O. *et al.* Mixed Invasive Ductal and Lobular Carcinoma of the Breast:
Prognosis and the Importance of Histologic Grade. *Oncologist* **24**, (2019).

35. Zhao, H. The prognosis of invasive ductal carcinoma, lobular carcinoma and mixed ductal
and lobular carcinoma according to molecular subtypes of the breast. *Breast Cancer* **28**,
187–195 (2021).

36. Arps, D. P., Healy, P., Zhao, L., Kleer, C. G. & Pang, J. C. Invasive ductal carcinoma with
lobular features: a comparison study to invasive ductal and invasive lobular carcinomas of
the breast. *Breast Cancer Res Treat* **138**, 719 (2013).

37. M, C. *et al.* Lobular breast cancer: Histomorphology and different concepts of a special
spectrum of tumors. **13**, 3695 (2021).

38. Christgen, M., Kandt, L. D., Antonopoulos, W. & al., et. Inter-observer agreement for the
histological diagnosis of invasive lobular breast carcinoma. *J Pathol Clin Res* **8**, 191–205
(2022).

39. Z, C. *et al.* Invasive lobular carcinoma of the breast: A special histological type compared
with invasive ductal carcinoma. **12**, e0182397 (2017).

40. Yeatman, T. J. *et al.* Tumor biology of infiltrating lobular carcinoma. Implications for
management. *Ann Surg* **222**, 549–559 (1995).

41. Helvie, M. A., Paramagul, C., Anderson, K. L. & Adler, D. D. Invasive lobular carcinoma.
Imaging features and clinical detection. *Invest Radiol* **28**, 202–207 (1993).

42. Mann, R. M. The effectiveness of MR imaging in the assessment of invasive lobular
carcinoma of the breast. *Magn Reson Imaging Clin N Am* **18**, 259–276 (2010).

43. Christgen, M., Kandt, L. D., Antonopoulos, W. & al., et. Inter-observer agreement for the
histological diagnosis of invasive lobular breast carcinoma. *J Pathol Clin Res* **8**, 191–205
(2022).

44. Pérez-Mies, B. *et al.* The Clonal Relationship Between the Ductal and Lobular Components
of Mixed Ductal-Lobular Carcinomas Suggested a Ductal Origin in Most Tumors. *American
Journal of Surgical Pathology* **46**, 1545–1553 (2022).

45. Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell
Syst* **1**, 417–425 (2015).

46. Carbon, S. *et al.* The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res* **49**,
D325–D334 (2021).

47. Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res* **50**,
D687–D692 (2022).

48.   Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG for
taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res* **51**, D587–D592
(2023).

49.   Cleton-Jansen, A. M. E-cadherin and loss of heterozygosity at chromosome 16 in breast
carcinogenesis: different genetic pathways in ductal and lobular breast cancer? *Breast*
Cancer Res **4**, 5–8 (2002).

50.   André, F. *et al.* Alpelisib for PIK3CA -Mutated, Hormone Receptor–Positive Advanced
Breast Cancer . *New England Journal of Medicine* **380**, 1929–1940 (2019).

51.   Theodorou, V., Stark, R., Menon, S. & Carroll, J. S. GATA3 acts upstream of FOXA1 in
mediating ESR1 binding by shaping enhancer accessibility. *Genome Res* **23**, 12 (2013).

52.   Malik, N. *et al.* The transcription factor CBFB suppresses breast cancer through
orchestrating translation and transcription. *Nature Communications 2019 10:1* **10**, 1–15
(2019).

53.   Comprehensive molecular portraits of human breast tumours. *Nature* **490**, (2012).

54.   Gala, K. *et al.* KMT2C mediates the estrogen dependence of breast cancer through
regulation of ERα enhancer function. *Oncogene 2018 37:34* **37**, 4692–4710 (2018).

55.   Glaser, A. P. *et al.* APOBEC-mediated mutagenesis in urothelial carcinoma is associated
with improved survival, mutations in DNA damage response genes, and immune response.
Oncotarget **9**, 4537–4548 (2017).

56.   Shen, R. & Seshan, V. E. FACETS: allele-specific copy number and clonal heterogeneity
analysis tool for high-throughput DNA sequencing. *Nucleic Acids Res* **44**, (2016).

57.   Glaser, A. P. *et al.* APOBEC-mediated mutagenesis in urothelial carcinoma is associated
with improved survival, mutations in DNA damage response genes, and immune response.
Oncotarget **9**, 4537–4548 (2017).

58.   Moffitt, J. R., Lundberg, E. & Heyn, H. The emerging landscape of spatial profiling
technologies. *Nature Reviews Genetics 2022 23:12* **23**, 741–759 (2022).

59.   van Amerongen, R. Behind the Scenes of the Human Breast Cell Atlas Project. *J Mammary*
Gland Biol Neoplasia **26**, 67 (2021).

60.   Nguyen, Q. H. *et al.* Profiling human breast epithelial cells using single cell RNA sequencing
identifies cell diversity. *Nat Commun* (2018) doi:10.1038/s41467-018-04334-1.

61.   Gray, G. K. *et al.* A human breast atlas integrating single-cell proteomics and
transcriptomics. *Dev Cell* **57**, 1400-1420.e7 (2022).

62.   Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nature*
Genetics 2021 53:9 **53**, 1334–1347 (2021).

63.   Sun, H. *et al.* The relevance between hypoxia-dependent spatial transcriptomics and the
prognosis and efficacy of immunotherapy in claudin-low breast cancer. *Front Immunol* **13**,
7828 (2023).

64.   Bassiouni, R. *et al.* Spatial Transcriptomic Analysis of a Diverse Patient Cohort Reveals a
721   Conserved Architecture in Triple-Negative Breast Cancer. *Cancer Res* **83**, 34–48 (2023).

722   65.   Johnson, B. E. *et al.* An omic and multidimensional spatial atlas from serial biopsies of an
723   evolving metastatic breast cancer. *Cell Rep Med* **3**, 100525 (2022).

724   66.   Kulasinghe, A. *et al.* Spatial Profiling Identifies Prognostic Features of Response to Adjuvant
725   Therapy in Triple Negative Breast Cancer (TNBC). *Front Oncol* **11**, 1 (2021).

726   67.   Andersson, A. *et al.* Spatial deconvolution of HER2-positive breast cancer delineates
727   tumor-associated cell type interactions. *Nature Communications 2021 12:1* **12**, 1–14
728   (2021).

729   68.   Pérez-Mies, B. *et al.* The Clonal Relationship Between the Ductal and Lobular Components
730   of Mixed Ductal-Lobular Carcinomas Suggested a Ductal Origin in Most Tumors. *American
731   Journal of Surgical Pathology* **46**, 1545–1553 (2022).

732   69.   Badve, S. *et al.* Basal-like and triple-negative breast cancers: a critical review with an
733   emphasis on the implications for pathologists and oncologists. *Modern Pathology* **24**, 157–
734   167 (2011).

735   70.   Perou, C. M. Molecular Stratification of Triple-Negative Breast Cancers. *Oncologist* **16**, 61–
736   70 (2011).

737   71.   Michaut, M. *et al.* Integration of genomic, transcriptomic and proteomic data identifies
738   two biologically distinct subtypes of invasive lobular breast cancer. *Sci Rep* (2016)
739   doi:10.1038/srep18517.

740   72.   Djerroudi, L., Cabel, L., Bidard, F. C. & Vincent-Salomon, A. Invasive Lobular Carcinoma of
741   the Breast: Toward Tailoring Therapy? **114**, 1434–1436 (2022).

742   73.   Azim, H. A., Nguyen, B., Brohée, S., Zoppoli, G. & Sotiriou, C. Genomic aberrations in young
743   and elderly breast cancer patients. *BMC Med* **13**, 1–7 (2015).

744   74.   Afzaljavan, F., Sadr, A. S., Savas, S. & Pasdar, A. GATA3 somatic mutations are associated
745   with clinicopathological features and expression profile in TCGA breast cancer patients.
746   *Scientific Reports 2021 11:1* **11**, 1–13 (2021).

747   75.   BeLow, M. & Osipo, C. Notch Signaling in Breast Cancer: A Role in Drug Resistance. *Cells
748   2020, Vol. 9, Page 2204* **9**, 2204 (2020).

749   76.   Hopkins, J. L., Lan, L. & Zou, L. DNA repair defects in cancer and therapeutic opportunities.
750   *Genes Dev* **36**, 278 (2022).

751   77.   Voutsadakis, I. A. & Stravodimou, A. Homologous Recombination Defects and Mutations
752   in DNA Damage Response (DDR) Genes Besides BRCA1 and BRCA2 as Breast Cancer
753   Biomarkers for PARP Inhibitors and Other DDR Targeting Therapies. *Anticancer Res* **43**,
754   967–981 (2023).

755   78.   McCart Reed, A. E. *et al.* Mixed ductal-lobular carcinomas: evidence for progression from
756   ductal to lobular morphology. *Journal of Pathology* (2018) doi:10.1002/path.5040.

79. Kutasovic, J. R., McCart Reed, A. E., Sokolova, A., Lakhani, S. R. & Simpson, P. T. Morphologic and genomic heterogeneity in the evolution and progression of breast cancer. *Cancers* vol. 12 Preprint at https://doi.org/10.3390/cancers12040848 (2020).

80. Ciriello, G. *et al.* Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell* (2015) doi:10.1016/j.cell.2015.09.033.

81. McCart Reed, A. E. *et al.* An epithelial to mesenchymal transition programme does not usually drive the phenotype of invasive lobular carcinomas. *Journal of Pathology* (2016) doi:10.1002/path.4668.

82. Bajrami, I. *et al.* E-Cadherin/ROS1 Inhibitor Synthetic Lethality in Breast Cancer. *Cancer Discov* **8**, 498–515 (2018).

83. da Silva, E. M. *et al.* Abstract P2-23-15: Histologic, immunohistochemical and genomic comparison between classic Invasive lobular carcinomas and lobular-like invasive ductal carcinomas. *Cancer Res* **83**, P2-23–15 (2023).

84. Jiang, Y., Yang, M., Wang, S., Li, X. & Sun, Y. Emerging role of deep learning-based artificial intelligence in tumor pathology. *Cancer Commun* **40**, 154–166 (2020).

85. Couture, H. D. *et al.* Image analysis with deep learning to predict breast cancer grade, ER status, histologic subtype, and intrinsic subtype. *npj Breast Cancer 2018 4:1* **4**, 1–8 (2018).

86. Chan, R. C. K. *et al.* Artificial intelligence in breast cancer histopathology. *Histopathology* **82**, 198–210 (2023).

87. Merritt, C. R. *et al.* Multiplex digital spatial profiling of proteins and RNA in fixed tissue. *Nature Biotechnology 2020 38:5* **38**, 586–599 (2020).

88. da Silva, E. M. *et al.* Mesonephric and mesonephric-like carcinomas of the female genital tract: molecular characterization including cases with mixed histology and matched metastases. *Mod Pathol* **34**, 1570–1587 (2021).

89. da Silva, E. M. *et al.* TERT promoter hotspot mutations and gene amplification in metaplastic breast cancer. *NPJ Breast Cancer* **7**, (2021).

90. Selenica, P. *et al.* APOBEC mutagenesis, kataegis, chromothripsis in EGFR-mutant osimertinib-resistant lung adenocarcinomas. *Ann Oncol* **33**, 1284–1295 (2022).

91. Da Cruz Paula, A. *et al.* Genomic profiling of primary and recurrent adult granulosa cell tumors of the ovary. *Mod Pathol* **33**, 1606–1617 (2020).

92. da Silva, E. M. *et al.* Mesonephric and mesonephric-like carcinomas of the female genital tract: molecular characterization including cases with mixed histology and matched metastases. *Mod Pathol* **34**, 1570–1587 (2021).

93. Zehir, A. *et al.* Mutational landscape of metastatic cancer revealed from prospective clinical sequencing of 10,000 patients. *Nat Med* **23**, 703–713 (2017).

94. Cheng, D. T. *et al.* Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A Hybridization Capture-Based Next-Generation

Sequencing Clinical Assay for Solid Tumor Molecular Oncology. *The Journal of Molecular*
Diagnostics **17**, 251–264 (2015).

95. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* (2009) doi:10.1093/bioinformatics/btp324.

96. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res* **20**, 1297–1303 (2010).

Figure 1. Digital Spatial Profiling of mDLC Ductal and Lobular Tumor Regions.

Figure 2. Molecular Subtypes and Distinct Biological Signatures of Ductal and Lobular Tumor Regions.

Figure 3. Overview of Genomic Alterations in Ductal and Lobular Tumor Regions.

Figure 4: Single Cell Heterogeneity of Ductal and Lobular Tumor Populations.

Supplementary Figure S1: Digital Spatial Profiling RNAseq Dataset Quality Control.

Supplementary Figure S2: Patient Wise Differential Analysis and Comparison with Pan-Patient Differential Analysis.

Supplementary Figure S3: Transcription Factor Enrichment Analysis using Pan-Patient Lobular vs Ductal DEGs.

Supplementary Figure S4: Genes Associated with Key Differentially Enriched Biological Signatures.

A

	Sample	Purity	Depth	Ploidy
MDLC-1	Ductal	0.36	428x	2.02
	Lobular	0.45	523x	1.97
MDLC-2	Ductal	0.51	416x	2.04
	Lobular	0.31	339x	2.02
MDLC-3	Ductal	0.62	378x	3.35
	Lobular	0.4	490x	2.01

Supplementary Figure S5: MSK Impact Extended Results.

A

Metric	Value
Estimated Number of Cells	4,671
Fraction of Reads in Cells	90.6%
Mean Reads per Cell	91,174
Median Genes per Cell	1,481
Total Genes Detected	24,726
Median UMI Counts per Cell	4,480
Number of reads	425 million
Valid Barcodes	96%
Sequencing Saturation	78.4%

B

C

DEGs downregulated in Lobular

445 | 77 | 368

DSP   scRNAseq

Significance of Overlap
Representation factor: 6.1
p < 1.946e-38

DEGs Upregulated in Lobular

514 | 86 | 189

DSP   scRNAseq

Significance of Overlap
Representation factor: 9.7
p < 3.107e-61

D

Supplementary Figure S6: MDLC-1 Single-cell RNAseq DEG Analysis and Subcluster Marker Genes.

Supplementary Figure S7: MDLC-1 Single-cell RNAseq Pathway Analysis.