Laws of genome nucleotide composition

Zhang Zhang[1,2,3,*]

[1] National Genomics Data Center & CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China

[2] China National Center for Bioinformation, Beijing 100101, China

[3] University of Chinese Academy of Sciences, Beijing 100049, China

*Corresponding author (zhangzhang@big.ac.cn).

Running head: Laws of genome nucleotide composition

Abstract

Genome nucleotide composition is of fundamental significance in molecular evolution, genome organization and synthetic biology. Albeit it has been studied for decades, very few theoretical efforts have been devoted to exploring whether there is any law underlying variable genome nucleotide composition across different species. From the mathematical viewpoint, here we propose three laws of genome nucleotide composition and validate their effectiveness on a large collection of empirical genome sequences across three domains of life. The laws together provide a unifying framework that is able to unravel the mystery of genome-wide nucleotide composition variation across diverse species, paving the way toward a general theory with quantitative insights for deciphering basic principles of life and further advancing theoretical biology.

One-Sentence Summary: Laws of genome nucleotide composition

Introduction

Genome nucleotide composition, one of the most important characteristics at the genome-wide level, is usually expressed in terms of the proportions of four bases as well as their combinations in DNA molecule. It has been studied for decades (*1-3*) that genomes of different species are highly variable in their nucleotide composition (*4, 5*), as demonstrated that guanine-plus-cytosine content varies widely with a broader range from ~20% to ~80% (*6-9*). A body of empirical evidence has further accumulated that heterogeneity of genome-wide nucleotide composition in different species associates closely with several factors, such as, genome size (*10*), phylogeny (*11, 12*), growth temperature (*13*), environment (*14*), origin of replication (*15*), transposable element (*16*), bacterial land colonization (*17*), codon/amino acid usage (*18-20*), cell fate control (*21*), and natural selection (*22-25*), etc. Contrastingly, very few theoretical efforts (*26*) have been devoted to exploring whether there is any law underlying such variable genome nucleotide composition across different species. Theoretically, such law(s) would be desirable to be used as a fundamental framework for better understanding genome composition dynamics, molecular evolution, genome organization and synthetic biology. Built upon previous findings, here we propose three laws of genome nucleotide composition in a mathematical manner and demonstrate their effectiveness to formulate diverse genome nucleotide compositions in a large collection of 17,873 complete genome sequences across three domains of life.

First law: the law of base pairing

The first law is the Chargaff's rules (*27-29*) that adenine (A) pairs with thymine (T) and guanine (G) pairs with cytosine (C), leading to $P(A) = P(T)$ and $P(G) = P(C)$ (Eqs. 1 and 2), where P is the proportion (probability) of any base as well as their combination. Such

base pairing symmetry in any double-strand genome and each single strand corresponds to

the Chargaff's first parity rule and second parity rule, respectively. With biological,

chemical and physical significances, the rules played a crucial role in the discovery of the

double helix structure of DNA in 1953 (*30*) and laid profound foundations in advancing

molecular biology and genomics (*31*). Despite the debate on the Chargaff's second parity

rule (*32, 33*), the first law holds valid as testified by nearly perfect linear regression in a

wide diversity of genomes across the three domains of life (Fig. 1A and 1B).

$$P(A) = P(T) \tag{1}$$

$$P(G) = P(C) \tag{2}$$

Second law: the law of base equality

The second law states the base equality between purines (A and G) and pyrimidines (T and

C). As a consequence of the base-pairing nature of DNA double helix, a 1:1 stoichiometric

ratio of purines (R) and pyrimidines (Y) can be deduced. In other words, the proportion of

R approximates the proportion of Y, namely, $P(R) = P(Y)$. Additionally considering

guanine-plus-cytosine (S or GC) and adenine-plus-thymine (W) contents, $P(A) + P(G) +$

$P(T) + P(C) = 100\%$ can be further formulated as $P(R) + P(Y) = 2P(R) = 2P(Y) = P(S) +$

$P(W) = 100\%$ (Eqs. 3 and 4).

$$P(R) = P(Y) = 50\% \tag{3}$$

$$P(S) + P(W) = 100\% \tag{4}$$

Conforming with the second law as well as previous findings (*4, 7, 8*), $P(R)$ is observed to

approximate $P(Y)$ and fluctuate slightly around 50% in diverse genomes belonging to

Archaea, Bacteria and Eukarya (Fig. 1C and 1D). Meanwhile, considering that

independence of two variables means that the occurrence of one variable does not affect

the probability of the other, S, varying from ~20% to ~80%, is believed to be statistically

independent from R, as indicated by linear regression slope very close to the optimum value

of zero and intercept near 0.5 (Fig. 1D).

Third law: the law of base composition

The third law expresses the principle of base composition. Suppose that the universal set

of the four bases is Ω={A, T, G, C} and X and Y are two subsets of U, there are three

common operations in set theory: (1) union of X and Y, viz., $X \cup Y$, is the set of all elements

that are a member of X or Y or both; (2) intersection of X and Y, viz., $X \cap Y$, is the set of

all elements that are members of both X and Y; (3) complement of X relative to Ω, viz.,

X^c, is the set of all members that are not members of X. Thus, GC and purine contents can

be denoted as $S = G \cup C$ and $R = A \cup G$, respectively. Likewise, $W = A \cup T = S^c$ and $Y =$

$T \cup C = R^c$. Because S and R form an independent pair as mentioned above, thus $P(S \cap R)$

can be quantitatively expressed as $P(S)$ multiplied by $P(R)$, namely, $P(S \cap R) = P(S) \times P(R)$,

which is also applicable to the other three pairs: S^c and R, S and R^c, and S^c and R^c (for

details see our previous study (*34*)). As a result, the proportion of each base can be

quantitatively formulated as

$$P(A) = P(S^c \cap R) = P(S^c) \times P(R), \tag{5}$$

$$P(T) = P(S^c \cap R^c) = P(S^c) \times P(R^c), \tag{6}$$

$$P(G) = P(S \cap R) = P(S) \times P(R), \tag{7}$$

$$P(C) = P(S \cap R^c) = P(S) \times P(R^c), \tag{8}$$

where $P(S^c) = P(W) = 1 - P(S)$ and $P(R^c) = P(Y) = 1 - P(R)$ according to Eqs. 3-4.

Strikingly, if $P(R) = 0.5$, then $P(A) = P(T) = P(S^c)/2$ and $P(G) = P(C) = P(S)/2$, which is a

special case equivalent to the first law or Chargaff's rules (*27, 28*). Based on our empirical

datasets, the third law is effective to quantify the base proportions very close to the

observed ones, as signified by squared correlation coefficients very close to the optimum

value of 1.0, linear regression slopes near the optimum value of 1.0 and intercepts

approaching the optimum value of zero (Fig. 1E to 1H).

Concluding thoughts

The first law of base pairing uncovers the complementary nature of DNA that is essential

for its structure, stability and function, the second law of base equality reveals the equality

relationship between purines and pyrimidines as well as the independence relationship

between GC content and purine content, and the third law deduces the mathematical

principle of each base composition that is of great utility for designing genomes for any

specified composition. Together, the three laws provide a unifying framework for studying

genome nucleotide composition (Fig. 2), which is not only able to unravel the mystery of

various genome-wide nucleotide compositions across diverse organisms as validated in

large-scale empirical genome data across the three domains of life, but also to detect

organisms with unusual genome-wide nucleotide composition that potentially experience

complex evolutionary processes adapted to extreme habitats (e.g., strong disparity of 14.97%

G and 29.83% C in *Candidatus* Chazhemtobacterium aquaticus Ch65 (*35*)).

Nowadays (and in the foreseeable future), we are drowning in the deluge of multi-omics

data growing at exponential rates, thirsting for fundamental theories to explain or predict a

range of biological phenomena and to derive general laws from a number of empirical

observations and experiments. From this perspective, the three laws, as the union of

theoretical and empirical work, potentially pave the way toward a more general theory with

quantitative insights for studying genome organization and evolution, driving synthetic

genome engineering and further advancing theoretical biology as a long-term objective on

a par with theoretical physics.

Authors' contributions

126   ZZ formulated this theoretical idea, wrote the manuscript and supervised the research. The

127   author read and approved the final manuscript.

Acknowledgments

129   I sincerely thank Haipeng Li, Yalong Guo, Yong Zhang, Jun Yu, Songnian Hu, Jingfa Xiao

130   and Lina Ma for their valuable discussions on this work as well as Shuai Jiang and Zhao

131   Li for their assistance in data collection. This work was supported by the National Natural

132   Science Foundation of China (32030021) and International Partnership Program of the

133   Chinese Academy of Sciences (153F11KYSB20160008).

References

1.   N. Sueoka, On the genetic basis of variation and heterogeneity of DNA base
composition. *Proc Natl Acad Sci U S A* **48**, 582-592 (1962).
2.   S. Karlin, A. M. Campbell, J. Mrazek, Comparative DNA analysis across diverse
genomes. *Annu Rev Genet* **32**, 185-225 (1998).
3.   A. Nekrutenko, W. H. Li, Assessment of compositional heterogeneity within and
between eukaryotic genomes. *Genome Res* **10**, 1986-1995 (2000).
4.   E. Haywood-Farmer, S. P. Otto, The evolution of genomic base composition in
bacteria. *Evolution* **57**, 1783-1792 (2003).
5.   S. Karlin, J. Mrazek, Compositional differences within and between eukaryotic
genomes. *Proc Natl Acad Sci U S A* **94**, 10227-10232 (1997).
6.   A. Muto, S. Osawa, The guanine and cytosine content of genomic DNA and
bacterial evolution. *Proc Natl Acad Sci U S A* **84**, 166-169 (1987).
7.   J. Hu, X. Zhao, Z. Zhang, J. Yu, Compositional dynamics of guanine and cytosine
content in prokaryotic genomes. *Res Microbiol* **158**, 363-370 (2007).
8.   X. Zhao, Z. Zhang, J. Yan, J. Yu, GC content variability of eubacteria is governed
by the pol III alpha subunit. *Biochem Biophys Res Commun* **356**, 20-25 (2007).
9.   H. Wu, Z. Zhang, S. Hu, J. Yu, On the molecular mechanism of GC content
variation among eubacterial genomes. *Biol Direct* **7**, 2 (2012).
10.   H. Nishida, Evolution of genome base composition and genome size in bacteria.
Front Microbiol **3**, 420 (2012).

11.   A. O. Mooers, E. C. Holmes, The evolution of base composition and phylogenetic
inference. *Trends Ecol Evol* **15**, 365-369 (2000).
12.   E. R. Reichenberger, G. Rosen, U. Hershberg, R. Hershberg, Prokaryotic nucleotide
composition is shaped by both phylogeny and the environment. *Genome Biol Evol*
7, 1380-1389 (2015).
13.   H. Musto *et al.*, Correlations between genomic GC levels and optimal growth
temperatures in prokaryotes. *FEBS Lett* **573**, 73-77 (2004).
14.   K. U. Foerstner, C. von Mering, S. D. Hooper, P. Bork, Environments shape the
nucleotide composition of genomes. *EMBO Rep* **6**, 1208-1213 (2005).
15.   J. R. Lobry, Asymmetric substitution patterns in the two DNA strands of bacteria.
Mol Biol Evol **13**, 660-665 (1996).
16.   S. Boissinot, On the Base Composition of Transposable Elements. *International*
journal of molecular sciences **23**, (2022).
17.   H. Wu, Y. Fang, J. Yu, Z. Zhang, The quest for a unified view of bacterial land
colonization. *The ISME journal* **8**, 1358-1369 (2014).
18.   R. Hershberg, D. A. Petrov, Selection on codon bias. *Annu Rev Genet* **42**, 287-299
(2008).
19.   G. A. Singer, D. A. Hickey, Nucleotide bias causes a genomewide bias in the amino
acid composition of proteins. *Mol Biol Evol* **17**, 1581-1588 (2000).
20.   A. Garcia, S. Gonzalez, F. Antequera, Nucleosomal organization and DNA base
composition patterns. *Nucleus* **8**, 469-474 (2017).
21.   R. Pantier *et al.*, SALL4 controls cell fate in response to DNA base composition.
Mol Cell **81**, 845-858 e848 (2021).
22.   F. Hildebrand, A. Meyer, A. Eyre-Walker, Evidence of selection upon genomic
GC-content in bacteria. *PLoS Genet* **6**, e1001107 (2010).
23.   H. Long *et al.*, Evolutionary determinants of genome-wide nucleotide composition.
Nat Ecol Evol **2**, 237-240 (2018).
24.   M. Bulmer, The selection-mutation-drift theory of synonymous codon usage.
Genetics **129**, 897-907 (1991).
25.   S. L. Chen, W. Lee, A. K. Hottes, L. Shapiro, H. H. McAdams, Codon usage
between genomes is constrained by genome-wide mutational processes. *Proc Natl*
Acad Sci U S A **101**, 3480-3485 (2004).
26.   R. A. Elton, Theoretical models for heterogeneity of base composition in DNA. *J*
Theor Biol **45**, 533-553 (1974).
27.   E. Chargaff, Chemical specificity of nucleic acids and mechanism of their
enzymatic degradation. *Experientia* **6**, 201-209 (1950).
28.   E. Chargaff, Some recent studies on the composition and structure of nucleic acids.
J Cell Physiol Suppl **38**, 41-59 (1951).
29.   E. Chargaff, R. Lipshitz, C. Green, Composition of the desoxypentose nucleic acids
of four genera of sea-urchin. *J Biol Chem* **195**, 155-160 (1952).
30.   J. D. Watson, F. H. Crick, Molecular structure of nucleic acids; a structure for
deoxyribose nucleic acid. *Nature* **171**, 737-738 (1953).
31.   D. R. Forsdyke, J. R. Mortimer, Chargaff's legacy. *Gene* **261**, 127-137 (2000).
32.   D. Mitchell, R. Bridge, A test of Chargaff's second rule. *Biochem Biophys Res*
Commun **340**, 90-94 (2006).
33.   Y. Guo, An overview on the DNA nucleotide compositions across kingdoms. *Gene*
Reports **8**, 45-48 (2017).

34. Z. Zhang, J. Yu, Modeling compositional dynamics based on GC and purine
contents of protein-coding sequences. *Biol Direct* **5**, 63 (2010).
35. V. V. Kadnikov, A. V. Mardanov, A. V. Beletsky, O. V. Karnachuk, N. V. Ravin,
Complete Genome of a Member of a New Bacterial Lineage in the Microgenomates
Group Reveals an Unusual Nucleotide Composition Disparity Between Two
Strands of DNA and Limited Metabolic Potential. *Microorganisms* **8**, 320 (2020).
208
209

 Figure Legends

Fig. 1. Genome-wide proportions of nucleotide composition across three domains of

life.

(A) Proportion of adenine (A) vs. proportion of thymine (T); **(B)** Proportion of guanine (G)

vs proportion of cytosine (C); **(C)** Proportion of guanine-plus-cytosine content (S) vs.

difference between proportion of purine content (R) and proportion of pyrimidine content

(Y); **(D)** Proportion of S vs. proportion of R; **(E-H)** Observed vs. estimated proportions for

A, T, G, and C, respectively. Observed proportions of the four bases were derived from a

large collection of 17,873 complete genome sequences in NCBI RefSeq, including 313 in

Archaea, 17,289 in Bacteria, and 271 in Eukarya. Estimated proportions of the four bases

were quantified according to Eqs. 5-8. In all panels, each point represents the composition

for a specific genome. All detailed information were summarized in Supplementary Table

S1.

Fig. 2. Schematic representation of laws of genome nucleotide composition.

The laws are illustrated by equations and operations of intersection ($\cap$), union ($\cup$) and

complement (^{c}) on four bases—adenine (A), thymine (T), guanine (G) and cytosine (C),

where P is the proportion (probability) of any base as well as their combination. The first

law of base pairing states the complementary nature of DNA that leads to

$P(A)=P(T)=P(W)/2$ and $P(G)=P(C)=P(S)/2$, where $W=A\cup T$, $S=G\cup C$, $R=A\cup G$, and

$Y=T\cup C$. The second law of base equality reveals the quantitative relationships of

$P(R)=P(Y)=50\%$ and $P(S)+P(W)=100\%$, where $R\cup Y$ or $S\cup W$ is the universal set $\Omega=\{A,$

T, G, C$\}$, $R\cup W=\{A, T, G\}=C^c$ and $Y\cup S=\{T, G, C\}=A^c$. The third law deduces the

mathematical principle of each base composition given the independence relationship

between S and R, viz., $P(G)=P(S \cap R)=P(S)P(R)$, $P(A)=P(W \cap R)=P(W)P(R)$, $P(C)=P(S$

$\cap Y)=P(S)P(Y)$, and $P(T)=P(W \cap Y)=P(W)P(Y)$, where $W=S^c$ and $Y=R^c$.

Figures

Fig. 1

Fig. 2

Law of Base Pairing   Law of Base Equality   Law of Base Composition

Supplementary information

244 **Table S1** Statistics of genome-wide nucleotide composition as well as other related meta

245 information of 17,873 complete genome sequences across three domains of life.

Laws of Genome Nucleotide Composition

Law of Base Pairing

$P(A)=P(T)=P(W)/2$
$P(G)=P(C)=P(S)/2$

∪	A	C
G	R	S
T	W	Y

Law of Base Equality

$P(R)=P(Y)=0.5$
$P(S)+P(W)=1$

∪	S	R
W	Ω	C^c
Y	A^c	Ω

Law of Base Composition

$P(G)=P(S)P(R)$ $P(C)=P(S)P(Y)$
$P(A)=P(W)P(R)$ $P(T)=P(W)P(Y)$

∩	R	Y
S	G	C
W	A	T