Genomic Insights into the Population History and Adaptive Traits of Latin American Criollo Cattle

James A. Ward[1], Said I. Ng'ang'a[2,3], Imtiaz A. S. Randhawa[4], Gillian P. McHugo[1], John F. O'Grady[1], Julio M. Flórez[5,6], John A. Browne[1], Ana M. Pérez O'Brien[5], Antonio J. Landaeta-Hernández[7], José F. Garcia[6], Tad S. Sonstegard[5], Laurent A. F. Frantz[2,3], Michael Salter-Townshend[8], David E. MacHugh[1,9]

[1] Animal Genomics Laboratory, UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[2] Palaeogenomics Group, Department of Veterinary Sciences, Ludwig Maximilian University, Munich D-80539, Germany.

[3] School of Biological and Chemical Sciences, Queen Mary University of London, London E1 4NS, United Kingdom.

[4] Faculty of Science, The University of Queensland, Gatton, QLD 4343, Australia

[5] Acceligen, 3388 Mike Collins Drive, Eagan, MN, 55121, USA.

[6] Department of Preventive Veterinary Medicine and Animal Reproduction, School of Agricultural and Veterinarian Sciences, São Paulo State University (Unesp), Jaboticabal, Brazil.

[7] Unidad de Investigaciones Zootécnicas, Facultad de Ciencias Veterinarias, Universidad del Zulia, Maracaibo, Venezuela.

[8] UCD School of Mathematics and Statistics, University College Dublin, Dublin, D04 V1W8, Ireland.

[9] UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Dublin, D04 V1W8, Ireland.

Author for correspondence

David E. MacHugh

e-mail: david.machugh@ucd.ie

ABSTRACT

Criollo cattle, descendants of animals brought by Iberian colonists to the Americas, have been subject to centuries of natural and human-mediated selection in novel tropical agroecological zones. Consequently, these breeds have evolved distinct characteristics such as resistance to disease and exceptional heat tolerance. In addition to European taurine (*Bos taurus*) ancestry, it has been proposed that gene flow from African taurine and Asian zebu (*Bos indicus*) cattle has shaped the ancestry of Criollo cattle. In this study, we analysed Criollo breeds from Colombia and Venezuela using whole genome sequencing data to examine population structure and admixture at very high resolution. We did not detect evidence of African taurine admixture in Criollo cattle; however, there was evidence of longstanding zebu ancestry in one breed (Hartón del Valle). In addition, we detected selection signatures associated with a myriad of adaptive traits, revealing genes linked to thermotolerance, reproduction, fertility, immunity, and distinct coat and skin coloration traits. This study underscores the remarkable adaptability of Criollo cattle and highlights the genetic richness and potential of these breeds in the face of climate change, habitat flux, and disease challenges. Further research is warranted to leverage these findings for more effective and sustainable cattle breeding programmes.

Keywords: cattle, Latin America, microevolution, population genomics, selection, thermotolerance

INTRODUCTION

The term "Criollo" is derived from the Portuguese word "crioulo" and was originally used in the colonial period to distinguish people born overseas from those born in the home country. Over time, this term was extended to domestic livestock (cattle, sheep, horses, and goats) with genetic roots in the Iberian Peninsula. The first introductions of cattle (taurine – *Bos taurus*) to the Americas were relatively recent, dating back to the end of the 15[th] century with animals transported during Columbus' second voyage and brought initially to the island of Hispaniola [1]. Early estimates suggest that fewer than 1,000 cattle were introduced to the Americas during these initial expeditions [2]. However, over the ensuing five centuries, increasing numbers of cattle were exported to North and South America particularly as the process of European colonization intensified during the 18[th] and 19[th] centuries [3]. A comparable narrative has been reported for sheep [4], and for horses, as both historical and extant horses in North America exhibit pronounced genetic affinities with Iberian horse populations [5]. More recently, zebu cattle (*Bos indicus*) were directly introduced from South Asia—primarily the region encompassed by present-day India and Pakistan [6].

The extent of contributions from cattle outside Europe and the Indian subcontinent to the South and Central American cattle gene pools remains uncertain. Modern Iberian cattle populations, whose ancestors likely contributed the bulk of South American Criollo cattle ancestry have an African taurine cattle genomic component [7-11]; however, this could post-date the initial introductions to the Americas. In parallel to this, it has also been posited that African cattle may have been directly introduced into South America alongside the initial imports [12]. This hypothesis is supported by the presence of the T1-c mtDNA haplotype, which is prevalent in African cattle, rarely found in Iberian animals, yet appears with an approximate frequency of 8% in Criollo cattle. [13]. Relatively few studies have explored potential autosomal contributions from African cattle. Preliminary findings, however, indicate that the degree of African ancestry in Criollo cattle is contingent upon the specific breed. Notably, the Guadeloupe breed, indigenous to the Caribbean,

exhibits a substantial African genetic influence, with almost 50% of its genetic ancestry estimated

to trace back to African origins [14, 15]. The potential direct or indirect influence of zebu cattle is

also an active area of research. While there is no question on whether zebu cattle have been directly

imported from South Asia, it is unclear whether this accounts for all of the zebu ancestry evident in

South and Central American cattle [14]. In Africa a gradient of zebu ancestry is present, escalating

across the continent from West to East Africa [16-20], Therefore, it is plausible that some of the

zebu ancestry for certain Criollo breeds could be attributed to historical African cattle importation

[15].

The cattle that were brought to the Americas from Iberia in the 15th and 16th centuries were

adapted to Mediterranean agroecologies and, over the succeeding centuries, these populations have

evolved adaptations to tropical and arid environments, emerging as distinctive breeds with unique

heat tolerance and disease resistance traits [11, 21-29]. Despite this, Criollo cattle are not highly

valued in modern production systems and are slowly being replaced with more productive

commercial breeds, which is eroding indigenous cattle genetic resources in South and Central

America [30-32]. Therefore, using genome-scale data, the primary aims of this study were to

evaluate *B. taurus* and *B. indicus* ancestry and detect historical gene flow from African cattle into

the Blanco Orejinegro, Hartón del Valle, and Limonero Criollo breeds. In addition, we use high-

resolution analyses of genetic structure, and selection signatures to provide new information about

the microevolutionary histories of these populations.

MATERIALS AND METHODS

Cattle samples, library preparation, high-throughput sequencing, and variant calling

In total, 34 Criollo cattle were used to generate high-quality de novo whole-genome sequence

(WGS) data sets with the following methods. Coat hair follicles obtained from 24 Colombian and

Venezuelan cattle and semen straws from 10 Colombian AI bulls were used to obtain genomic

DNA using standard commercial procedures for livestock DNA extraction and purification

(Weatherbys Scientific, Co. Kildare, Ireland). For the coat hair follicle-derived samples, the quality

and concentration of DNA was low; therefore, library preparation was performed in-house using the

Nextera DNA Flex Library Prep kit (now available as the Illumina® DNA Prep kit; Illumina,

Cambridge, UK). The WGS data generation was performed by sequencing to a mean depth of 20×

using two commercial service providers (Novogene, Cambridge, UK and Macrogen, Seoul, South

Korea). The geographical origins of the three primary Criollo breeds used in this study are shown in

figure *1*.

Publicly available WGS data was also assembled for 37 European taurine (EUT) (NCBI

Bioproject IDs: PRJNA210521; PRJNA318089; PRJNA343262; and PRJNA514237), 13 American

indicine (ASI) (NCBI Bioproject ID: PRJNA343262), nine African taurine (AFT), and 13 American

Criollo (AMC) cattle (NCBI Bioproject IDs: PRJNA471656 and PRJNA422135). Sequence reads

were mapped to the bovine reference genome (ARS-UCD1.2) [33] using the default parameters for

bwa-mem2 v.2.2.1 [34]. Samtools v.1.13 [35] was used to sort bam files and create index files.

Potential PCR duplicates were identified using the "MarkDuplicates" function of Picard v.2.27.1

(http://broadinstitute.github.io/picard). The "BaseRecalibrator" and "ApplyBQSR" functions of the

genome analysis toolkit (GATK) v.4.2.6.1 [36] were used to perform base quality score

recalibration.

Figure 1. Approximate geographical origins of the animals from three Criollo breeds that were sampled in Colombia and Venezuela and used to generate new whole-genome sequence data. Map modified from original by Mapswire (https://mapswire.com) (Creative Commons Attribution 4.0 International License CC-BY 4.0). Individual cattle art images modified from Felius [37] with permission of the author.

Candidate SNPs were called from the bam files using the "HaplotypeCaller" function in GATK with the "--emit-ref-confidence GVCF" option. Individual GVCF files were merged using the "GenomicsDBImport" function. SNPs from the genomics database were called and selected using "GenotypeGVCFs" and the "SelectVariants" function. The "VariantFiltration" function of GATK was used to avoid possible false-positive calls according to GATK best practices: SNP clusters with "--cluster-size 3" and "--cluster-window-size 10", quality by depth "QD <2", read position rank sum test "ReadPosRankSum < -8", phred-scaled variant quality score "QUAL < 30",

Fisher strand "FS > 60", mapping quality "MQ < 40", mapping quality rank sum test

"MQRankSum < -12.5" and strand odds ratio "SOR > 3". Triallelic SNPs and those with a minor

allele frequency < 0.01 and/or missing genotype rates over > 0.1 were filtered out. The SNPs

remaining in each WGS data set were then annotated according to their positions using SnpEff [38].

The combined panel of 106 cattle samples with WGS data are detailed in electronic supplementary

material, table S1.

BovineHD 777k array data

For broader comparative population genomics analyses, publicly available BovineHD

777k SNP array data for 45 Iberian cattle representing 14 breeds [39], and 12 Criollo cattle

representing one breed, the Senepol were obtained from the WIDDE database [40]. This SNP panel

was originally mapped to the UMD3.1 assembly [41]; therefore, we updated SNP positions to ARS-

UCD1.2 following a protocol detailed by Riggio et al. [42]. Firstly, BovineHD data were converted

from the TOP coding format to the FOR coding format using the SNPchiMp v.3 suite of tools [43].

Secondly, SNP positions from UMD3.1 were updated to the ARS-UCD1.2 genome using updated

BovineHD array coordinates available from the NAGRP Data Repository [44] and SNPchiMp v.3

data for the BovineHD array with positions updated using the liftOver implementation of the

rtracklayer R package [45]. Thirdly, reference and alternative alleles for the data were updated

using BovineHD array SNPs that were extracted from WGS data aligned to the ARS-UCD1.2

genome and files of updated alleles also available from the NAGRP Data Repository. Processing of

these data was conducted using PLINK v1.90b6.25 [46] and R version 4.1.3 [47]. The combined

panel of 57 cattle samples with BovineHD 777k SNP data are detailed in electronic supplementary

material, table S2.

Population differentiation and genetic structure analyses

Principal component analysis (PCA) for all animals (WGS and imputed BovineHD genotype

data) were performed using PLINK v1.90b6.25 and the results were plotted using ggplot2 v3.4.0

[48] in the R v4.1.3 environment. The genetic structure of each population was visualized using

fastSTRUCTURE v1.0 and ggplot2 v3.4.0 with $K = 2–5$ unsupervised modelled ancestries [49]. We

used VCFtools v.0.1.16 [50] to calculate nucleotide diversity (π) in windows of 10 Mb.

Detection of historical admixture

Using ADMIXTOOLS [51] with default parameters, we calculated f_3 statistics using the

whole genome data set to test for evidence that the Blanco Orejinegro, Hartón del Valle, and

Limonero breeds are derived from the admixture of two sets of populations (European taurine

populations and American zebu cattle, and European taurine populations and African taurine

cattle). A significant negative f_3 value is considered evidence of historical admixture in the target

population (Criollo). For visualization of the f_3 statistics, we plotted the f_3 values against the

population trios with error bars indicating the standard errors using ggplot2 v3.4.0 [48].

Test statistics for detection of selection signatures

For the selection signatures analysis, only the WGS data set was used, which included

approximately 9.7 million SNPs genotyped in 106 cattle. We implemented three selection tests to

investigate if genomic regions were under selection. SNP-by-SNP F_{ST} values [52] were calculated

for each Criollo breed versus a panel of Holstein cattle using VCFtools v.0.1.16 [50]. XP-EHH

statistics were calculated using the program selscan v.1.2.0 [53] using the default parameters, where

each Criollo breed was also compared to the Holstein panel. The unstandardized XP-EHH values

were then normalized using the --norm function in selscan. We calculated the ΔSAF (directional

change in selected allele frequency) using the --freq function of VCFtools to generate allele

frequencies, then using a custom R script we calculated the ΔSAF for each Criollo breed versus the

Holstein panel. Similarly to the XPEHH values, ΔSAF values were standardized to $Z \sim N(0,1)$ as

described by Randhawa et al. [54].

Implementing the composite selection signals (CSS) methodology

To obtain composite selection signals (CSS) statistics [54] for each Criollo breed, the F_{ST}, XP-EHH, and ΔSAF statistics were combined by SNP ID (chromosome:position), which produced final sets of SNPs for the CSS analysis of 9,344,833, 9,465,007 and 9,660,647 million SNPs, for the Blanco Orejinegro, Hartón del Valle, and Limonero, respectively. Following this, the MINOTAUR package in R [55] was used to calculate CSS statistics as described by Randhawa et al. [54]. The resulting statistics were then averaged over 20 kb windows to reduce spurious signals. To further reduce the likelihood of spurious signals, a signal was only considered significant if at least one SNP in the 0.1% of CSS scores was flanked by at least five other SNPs in the top 1% of CSS values [54, 56-58].

Identification of genomic regions under selection

From the top 0.1% of SNPs identified, genomic intervals in the format chromosome:start-end were extracted and the BovineMine resource (v1.6) [59] was then used to find genes ± 500 kb from the start and end of these regions. The functional relevance of these CSS-derived gene sets was then evaluated using the scientific literature and the g:Profiler software tool [60] was used to identify overrepresented gene ontology (GO) terms for each of the CSS-derived gene sets by breed. Following this, BovineMine v1.6 was used to systematically identify previously described quantitative trait loci (QTLs) for production, health, and welfare traits that are located within the CSS cluster regions for each breed.

RESULTS

Population differentiation, genetic structure, and admixture in Criollo cattle

Merging of the WGS and remapped BovineHD 777K BeadChip data sets resulted in a shared set of 367,339 SNPs for 163 individual animals. A PCA of these SNPs demonstrated that PC1 and PC2 accounted for 22.5% and 9.5% of the variation observed, with PC1 representing the split between the *B. taurus* (taurine) and *B. indicus* (zebu) lineages and PC2 the split between the

European and African taurine (figure *2a*). The genetic structure of the cattle populations was

analysed using fastSTRUCTURE, wherein the inferred number of clusters (K) was set to range

between two and five (K=2–5) (figure *2b*). At $K = 2$, the initial partition delineated the taurine and

zebu lineages, which mirrored PC1 in the PCA. At $K = 3$, a second partition differentiated the

European and African taurine lineages, which reflects PC2 in the PCA. The subsequent split at $K =$

4 discerned an additional cluster within the European taurine populations, and $K = 5$ primarily

segregated the Iberian taurine and Criollo populations from the other groups.

Figure 2. Genetic diversity and admixture in African, Asian, Criollo, and European cattle. (*a*) Results

of the principal component analysis (PCA) for 163 animals (106 from whole-genome sequence

data and 57 from high-density SNP array data) with a shared set of 367,339 SNPs. The PCA plot

shows the coordinates for each animal based on the first two principal components. Principal

component 1 (PC1) differentiates the *Bos taurus* and *Bos indicus* lineages, whereas PC2 separates

the African and European taurine groups. A histogram plot shows the variance contributions for

the first 20 PCs, with PC1 and PC2 accounting for 22.5% and 9.5% of the variance, respectively. (*b*)

Unsupervised genetic structure plot for European and Iberian taurine populations, Criollo cattle

populations, African taurine, and Asian zebu populations. Results for ancestry clusters ranging

from *K*=2-5 are shown.

Further examination of the PCA and structure plots in figure *2a* and *2b* showed that the

Criollo cattle samples tend to cluster with the Iberian taurine cattle samples. However, some breeds,

notably the Senepol and the Hartón del Valle, display a consistent and relatively uniform ancestry

component from the *B. indicus* zebu lineage. The other Criollo breeds (Blanco Orejinegro, Costeño

con Cuernos, Limonero, and Sanmartinero) are more heterogeneous with some individual animals

showing substantial zebu admixture and other animals with an exclusively taurine genetic

composition. Calculation of f_3 statistics, which used the WGS data, indicate that there is no strong

evidence of admixture in the Blanco Orejinegro (figure *3a*) and Limonero (figure *3c*) breeds when

using any of the European taurine reference populations and the American indicine as the second

reference. However, there is some evidence of admixture in the Hartón del Valle breed (figure *3b*)

as the f_3 statistics for all population trios that used any European taurine reference population with

the American indicine as the second reference population produced negative f_3 values. This result is

reflected in the positions of the Hartón del Valle animals on the PCA plot (figure *2a*), and their

genetic composition shown in the structure plot (figure *2b*).

Selection signals across Criollo cattle breed genomes

The CSS test was used to identify selection signals across the genomes of three cattle breeds

that had WGS data and sufficient sample sizes: the Blanco Orejinegro, Limonero, and Hartón del

Valle breeds. For each breed, the test highlighted several selection signals spanning multiple

chromosomes. For the Blanco Orejinegro breed, the CSS test detected 31 clustered regions

spanning 17 chromosomes, with 442 genes located within these regions and extended ±500 kb

(figure *4a*). Similarly, for the Hartón del Valle breed, the CSS test revealed 31 distinct clusters

distributed over 17 chromosomes, encompassing 410 genes (figure *4b*). Lastly, for the Limonero

breed, the CSS test detected 38 cluster regions, also extending across 17 chromosomes, and that

contained 451 genes in total (figure *4c*). When these results were consolidated, 76 genes within

selection signature regions are shared across the three breeds, and BTA18 was the chromosome

with the largest number of these genes (28 genes; 36.8% of the total). The CSS cluster regions and

the genes within these regions are detailed by breed in electronic supplementary material, table S3).

Figure 3. Three-population tests using Blanco Orejinegrio (*a*), Hartón del Valle (*b*) and Limonero (*c*) as target populations. Each coloured data point shows the f_3 statistics value and the horizontal line represents plus or minus the standard error. Three-letter breed/population codes are detailed in Table S1. All tests were conducted on whole-genome sequencing data. Individual cattle art images modified from Felius [37] with permission of the author.

Gene ontology (GO) term enrichment analysis

The functional significance of the genes within the CSS cluster regions for each breed was evaluated using GO term overrepresentation analysis with the g:Profiler software tool [60]. After adjusting for multiple tests and using a *P*-value adjusted for the false discovery rate (FDR-$P_{\text{adj.}}$ < 0.05), overrepresented GO terms were catalogued for each breed. The results of the GO term overrepresentation analysis are shown in electronic supplementary material, tables S4–S6 and figure *5*. For both the Blanco Orejinegro (figure *5a*) and Limonero (figure *5b*) breeds, the GO terms related primarily to immunobiology, encompassing functions such as immunoglobulin receptor binding, phagocytosis, complement activation, and antigen binding. Conversely, for the Hartón del

Valle breed (figure *5c*), the predominant GO terms were associated with skin epidermis and

keratinization processes.

Figure 4. Chromosome-wide circular plots of the smoothed composite selection signals (CSS)

scores for Blanco Orejinegro (*a*), Hartón del Valle (*b*) and Limonero (*c*). The dashed red lines

indicate the genome-wide 0.1% thresholds of the empirical CSS scores. Green coloured dots are

those deemed significant by being surrounded at least five SNPs among the top 1% of CSS scores.

Individual cattle art images modified from Felius [37] with permission of the author.

Figure 5. Gene ontology (GO) overrepresentation analysis results for the Blanco Orejinegrio (*a*), Hartón del Valle (*b*) and Limonero (*c*). The y-axis represents the negative $\log_{10}$ FDR-$P_{\text{adj.}}$ from the overrepresentation analysis. Individual cattle art images modified from Felius [37] with permission of the author.

Identification of quantitative trait loci (QTLs) within CSS cluster regions

Quantitative trait loci (QTLs) within the CSS cluster regions were systematically identified using BovineMine v1.6 [59]. The number of QTLs identified varied across the breeds, with 519 QTLs found in Blanco Orejinegro, 332 in Hartón del Valle, and 370 in Limonero. Interestingly, only 11 QTLs were common to all three breeds, of which a QTL associated with tick resistance [61] was particularly noteworthy. In the case of Blanco Orejinegro, most of the identified QTLs (291 out of 519) in the CSS cluster regions were associated with calving ease. Additionally, 46 QTLs were linked to milk protein percentage, 40 to milk yield, and 24 to coat texture. For the Limonero breed, there were 292 QTLs located in the CSS cluster regions related to calving ease; however, no other traits had more than five SNP markers among the QTLs. In the case of the Hartón del Valle breed, the most notable traits represented among the CSS cluster region QTLs were milk related. Specifically, 86 QTLs were associated with milk protein percentage, 45 with milk yield, and 35 with milk fat percentage. Additionally, 27 QTLs in Hartón del Valle were associated with coat texture. The results of the QTL search using BovineMine v1.6 are detailed in electronic supplementary material, table S7.

DISCUSSION

Population structure and admixture

The differentiation of cattle samples across PC1 of the PCA plot (figure *2a*) and the primary split in the structure plot (figure *2b*) for $K = 2$ reflects the evolutionary divergence between the *B. taurus* and *B. indicus* lineages, which occurred 150–500 kya [62-64]. The second division evident for PC2 in the PCA plot (figure *2a*) and for $K = 3$ in the structure plot differentiates the European and African taurine groups, a separation that has been well documented in several previous studies using genome-wide SNPs and other autosomal genetic markers [10, 17-19, 65, 66]. Extending these observations further, complex patterns of admixture due to human-mediated and unmanaged crossbreeding between taurine and zebu cattle have also been described in cattle populations across several geographical regions, particular in sub-Saharan Africa [10, 18, 19, 65].

Previous studies have also highlighted the presence of African taurine autosomal admixture in the Criollo cattle breeds of the Caribbean [11, 12, 14, 15]. Results from the present study, which are captured in the PCA plot (figure *2a*) and structure analysis (figure *2b*) suggest this is not the case for the Criollo breeds and individual animals analysed here (Blanco Orejinegro, Costeño con Cuernos, Hartón del Valle, Limonero, Sanmartinero, and Senepol). Therefore, the presence of a significant proportion of African taurine autosomal ancestry is highly breed-specific and may be confined to the Caribbean region [14, 15]. With regard to *B. indicus* ancestry, all of the South American Criollo cattle populations examined here do exhibit varying degrees of zebu admixture (figure *2b*). Interestingly, however, the Hartón del Valle and the Blanco Orejinegro, which both originate in Colombia, do not cluster together, which is a consequence of the relatively modest zebu ancestry in the Blanco Orejinegro versus a significantly larger zebu component in the Hartón del Valle. The Limonero breed samples in this study largely cluster together except for four animals that exhibit a significant proportion of indicine (zebu) ancestry. These results, taken together with the f_3 statistics for the Blanco Orejinegro, Hartón del Valle, and Limonero (figure *3*), and the histories of these breeds (the Blanco Orejinegro breed has been in a conservation programme since

1940 [67]), suggest that zebu gene flow has had a long-term impact on breeds such as the Hartón

del Valle and Senepol, which exhibit relatively uniform levels of zebu admixture (figure *2b*). This

has implications for understanding the selection signatures that are described in the following

sections (detailed in figure *4* and electronic supplementary material, table S3). In some cases, the

selection signals detected may be caused by adaptive introgression of *B. indicus* haplotypes. Details

for several of the genes identified through the CSS analysis are provided in supplementary material,

table S8.

Genomic signatures of human-driven selection in Criollo cattle

Blanco Orejinegro, translating to "white with black ears" designates a breed of cattle

predominantly found in the Andean region of Colombia. They stand out among cattle in the

Americas due to their distinctive coat and skin coloration; their coats are typically white, often

speckled with black spots, and their skin is highly pigmented, as exemplified by their black ears.

This coat-skin pattern also occurs in British White and Chillingham/White Park cattle [68], and the

White Fulani cattle of West Africa [69]. For White Park cattle, it has been demonstrated that their

coat colour variation is caused by a duplication in the *KIT* gene and an aberrantly inserted *KIT* gene

on BTA29 [70]. Other genes that harbour cattle coat colour polymorphisms include *MC1R*,

TWIST2, *MITF*, *PAX3*, *SLC45A2*, *COPA*, *TYR*, *TYRP1*, *KITLG*, and *ASIP*; however, in addition to

this group of genes, there are more than 100 other genes that have been shown to determine coat

colour traits in mammals [71-73]. For the Blanco Orejinegro breed, the genes *PAX3*, *SLC45A2*, and

KITLG are in genome regions with evidence for directional selection based on the CSS results.

Mutations in *PAX3* have been shown to result in a spectrum of phenotypic outcomes in horses from

white spotting to a uniform white coat trait [74]. Similarly, mutations in the bovine *SLC45A2* gene

have been shown to cause oculocutaneous albinism [75], while the *KITLG* gene has been associated

with the roan coat type in cattle [76].

Interestingly, for the Hartón del Valle and Limonero breeds, *MC1R* was part of a BTA18 CSS

cluster region. The protein receptor encoded by *MC1R* regulates tyrosinase levels in melanocytes,

an enzyme critical to melanin synthesis. High concentrations of tyrosinase facilitate the production

of eumelanin, contributing to darker brown or black shades, whereas lower concentrations drive the

synthesis of phaeomelanin, yielding lighter red or yellow tones [77]. It is therefore noteworthy that

Limonero and the Hartón del Valle cattle generally possess a "bayo" or bay coat, which is a yellow

to light brown/red colour [78-80]. Though not in a CSS cluster region in the Blanco Orejinegro, a

notable peak close to *MC1R* is present on BTA18. This gene has been observed to be under positive

selection in *B. indicus* cattle resulting in a light grey to white coat in the Brahman and Nelore

breeds [81, 82]. The adaptive significance of this observation is that lighter coat colours, determined

by functional polymorphisms at M*C1R*, reflect a significant portion of incident solar radiation [83],

which makes these cattle more suited to tropical environments [84].

Selection for heat tolerance in Criollo cattle

Criollo cattle breeds such as the Blanco Orejinegro, Hartón del Valle, and Limonero have

evolved over five centuries to thrive in the challenging tropical agroecologies of Colombia and

Venezuela. In the context of this environmental adaptation, a key finding for all three breeds is the

presence of a CSS peak on BTA20 that encompasses the *PRLR* gene, most notably in Blanco

Orejinegro and Hartón del Valle breeds, where it was statistically significant. Mutations in this gene

have been shown to cause the "slick" phenotype in in Criollo breeds such as the Blanco Orejinegro,

Harton del Valle, Limonero, Romosinuano, Costeño con Cuernos, Mexican Criollo Lechero

Tropical, and also Criollo-influenced composite breeds such as the Carora and Senepol [27, 28, 85].

Of particular interest for understanding how convergent evolution can act in livestock populations is

the existence of several different *PRLR* mutations that can produce the "slick" phenotype in Criollo

cattle [85]. Mutations in the bovine *PRLR* gene can have major effects on the length and the

structure of hair coats providing improved thermotolerance and concomitant increase on fertility

and milk yields in cattle populations that inhabit dry and tropical conditions [22, 78, 86-88]. In

addition, it has been shown that these mutations can act pleiotropically and cause other

physiological changes [89].

Variation at other genes may also confer enhanced thermotolerance in Criollo cattle; for

example, *MVD*, previously implicated as being under selection in North African cattle [90], was

found in a BTA18 CSS cluster region for all three breeds. In addition, *CCM2* which regulates heart

and blood vessel formation and integrity, is located in a Limonero BTA4 CSS cluster region and

was also highlighted by Ben☐Jemaa and colleagues in North African cattle [90]. This may

represent an adaptation that increases heat transfer from the interior to the skin, which is also

supported by the enhanced vascularisation of the dermis that has been documented in Limonero

cattle [78]. Another gene, *SESN2*, located in a BTA2 CSS cluster detected in the Hartón del Valle

and the Blanco Orejinegro breeds may also be associated with adaptation to tropical environments.

This gene was observed to be downregulated in a functional genomics study of heat-stress

responses in dermal fibroblast cells from zebu and zebu × taurine crossbred cattle [91].

Heat-induced stress is a form of proteotoxicity that can cause protein denaturation; if these

denatured proteins aggregate, they can become cytotoxic [92]. Many heat shock proteins (HSPs)

serve as molecular chaperones, aiding in the proper folding of proteins, and thereby preventing such

potentially harmful protein aggregation [93]. In this regard, *FBXO4* located in a BTA11 Limonero

CSS cluster region encodes a protein that regulates body temperature through interactions with

HSPB6 (previously HSP20) [84, 94]. We also observed that *DNAJA2* was in a Blanco Orejinegro

BTA18 CSS cluster region. This gene encodes a member of the DNAJ/HSP40 family of proteins

and shows increased expression in peripheral blood from Holstein calves exposed to heat stress

[95]. Similarly, *HIGD1A* (BTA22 CSS cluster) and *CBFA2T3* (a BTA18 CSS cluster distinct from

that containing *DNAJA2*), encode proteins associated with cellular responses to hypoxia [96-98],

which could be indicative of adaptation to oxidative stress caused by high temperatures [99].

Genomic signatures of selection for fertility and reproductive traits in Criollo cattle

Compared to taurine breeds from temperate zones, Criollo cattle are renowned for high

fertility and excellent reproductive performance under tropical conditions [21, 29, 100, 101] and

using the CSS method we detected several genes associated with these traits. For example, in the

Blanco Orejinegro breed a BTA7 CSS cluster region contains the *CATSPER3* gene exclusively

expressed in the testis, which encodes a sperm-specific ion channel protein directly linked to sperm

motility and male fertility in many species, including cattle [102]. Another CSS cluster region in

the Blanco Orejinegro on BTA20 contained *SPEF2*, which is involved in the formation and

functionality of sperm flagella [103]. Additionally, the *CREM* gene, implicated in spermatogenesis,

was located within a Blanco Orejinegro CSS cluster region on BTA13 [104]. Other genes related to

reproductive physiology were located within CSS cluster regions for the Hartón del Valle breed

including *HKDC1* on BTA28 that exhibits high expression in the testes [105], and *MISP3* on BTA7,

which is implicated in spermatogenesis [106]. For the Limonero breed, CSS cluster regions

included one on BTA10 that contains *FSIP1*; this gene is crucial for normal spermiogenesis and

flagella development [107]. In addition, the *PTGES* gene within a Limonero CSS cluster region on

BTA11 encodes an enzyme involved in the synthesis of prostaglandin E2 (*PGE2*), which is a

signalling molecule with a crucial role in reproductive processes [108]. In parallel to this, the QTL

analysis (electronic supplementary material, table S7) for the Blanco Orejinegro breed identified

QTLs predominantly associated with reproduction, notably calving ease, and lactation. These

findings corroborate previous results obtained using SNP array data for Blanco Orejinegro cattle

[25, 109]. The results of the QTL analysis in the CSS cluster regions for the Limonero breed were

similar to those for the Blanco Orejinegro breed.

Genomic adaptations to infectious diseases in Criollo cattle

Comparable to reproductive and fertility traits, desirable disease resistance and tolerance traits

have been highlighted as important features of Criollo cattle populations. However, empirical

evidence for the genetic basis of these traits is often lacking; instead, much of the evidence base is

anecdotal and derived from practical knowledge and experiences of veterinarians, farmers and

breeders [79]. In the present study, however, we have begun to address this knowledge gap through

identification of several immune genes in CSS cluster regions for the Blanco Orejinegro, Hartón del

Valle, and Limonero breeds. For example, the *CBFA2T3* gene located in the BTA18 CSS cluster

region detected for all three breeds and discussed in the context of hypoxia, has a role in the bovine

immune response to mammary gland inflammation [110]. Similarly, *CXCL14*, located in a BTA7

CSS cluster region for the Blanco Orejinegro and Limonero breeds has also been implicated in the

immune response to bovine mastitis caused by *Staphylococcus aureus* [111]. In addition, for the

Limonero, a BTA11 CSS cluster region contains the *LRRC8A* gene, which is essential for

development and function of T lymphocytes [112].

In Colombia and the surrounding tropical and subtropical regions, there is a significant

prevalence of vector-borne haemoparasitic infections [113] that induce anaemia, fever, wasting, and

reproductive dysfunction [114]. These diseases are primarily caused by eukaryotic parasites within

the *Babesia* [115] and *Trypanosoma* [116] genera, and bacterial pathogens from the *Anaplasma*

genus [117]. It is noteworthy, therefore, that the *EPB41* gene located within a Blanco Orejinegro

and Hartón del Valle BTA2 CSS cluster region is associated with resistance to anaemia and

trypanotolerance in African cattle [118]. Also, for the Blanco Orejinegro and Limonero, *RPS27* and

RPS20 are located in CSS cluster regions on BTA3 and BTA14, respectively and mutations in these

genes can cause Diamond-Blackfan anaemia [119, 120]. Furthermore, in the Blanco Orejinegro

breed, the *ZFPM1* gene within a BTA18 CSS cluster region, has been shown to play a role in

cardiac and hematopoietic development. In a murine *Trypanosoma congolense* infection model,

mice exhibiting decreased *ZFPM1* expression recovered more effectively from anaemia [121].

Gene GO term analysis in the Criollo CSS cluster regions revealed a marked

overrepresentation of immunobiological processes for the Blanco Orejinegro and Limonero breeds,

many of which overlapped between the two breeds (figure *5a* and *5c* and electronic supplementary

material, tables S4 and S5). The statistically significant overrepresented biological process GO

terms in the Blanco Orejinegro and Limonero included *phagocytosis* (GO:0006909), *phagocytosis,*

engulfment (GO:0006911), *phagocytosis, recognition* (GO:0006910), *complement activation,*

classical pathway (GO:0006958), and the *B cell receptor signalling pathway* (GO:0050853) For

molecular function, statistically significant overrepresented GO terms in the two breeds included

immunoglobulin receptor binding (GO:0034987), *antigen binding* (GO:0003823), and *RAGE*

receptor binding (GO:0050786). Again, immunobiology was prominent in the statistically

significant overrepresented cellular component GO terms for the two breeds, which included

immunoglobulin complex (GO:0019814) and *immunoglobulin complex, circulating* (GO:0042571).

Criollo cattle populations have adapted over several centuries to novel agroecological environments

with significant parasite and pathogen challenges. With regard to specific infectious disease

selection pressures and these overrepresented immunological GO terms, it is noteworthy that

Blanco Orejinegro cattle have a documented resistance to brucellosis [122, 123], a disease

characterized by abortion and retention of the placenta, which is principally caused by *Brucella*

abortus an intracellular bacterium that subverts phagocytic pathways and processes to enter and

manipulate host cells [124, 125].

CONCLUSIONS

464  This study provides valuable insights into the genomic basis of microevolutionary change in

Criollo cattle as they have adapted to the tropical environments of South America. Evidence of

strong selective pressure was apparent, particularly for the distinct coat and skin coloration traits

observed in these breeds, which are advantageous in cattle populations exposed to significant levels

of incident solar radiation. Notably, we also discovered genomic selection signatures that may be

associated with thermotolerance, again underscoring adaptation to hot climates. In addition, some of

the selection signatures we identified align with documented fertility traits in Criollo cattle.

Moreover, functional overrepresentation analysis revealed many genes related to immune function,

which could reflect resilience to multiple infectious disease challenges. Taken together, our results,

show the remarkable adaptability of Criollo cattle, which has been driven by natural and human-

mediated selection and that underscores the genetic richness and value of these breeds for future

breeding programmes.

SUPPLEMENTARY MATERIAL

Table S1: Cattle samples with whole-genome sequence (WGS) data.

Table S2: Cattle samples with BovineHD 777K SNP array data.

Table S3: Composite selection signatures (CSS) cluster regions and genes for all breeds

Table S4: Composite selection signatures (CSS) cluster gene ontology (GO) term overrepresentation analysis for the Blanco Orejinegro breed.

Table S5: Composite selection signatures (CSS) cluster gene ontology (GO) term overrepresentation analysis for the Hartón del Valle breed.

Table S6: Composite selection signatures (CSS) cluster gene ontology (GO) term overrepresentation analysis for the Limonero breed.

Table S7: Composite selection signatures (CSS) cluster QTLs identified for all breeds.

Table S8: Composite selection signatures (CSS) genes and functional inference.

Ethics: The hair and semen samples used in this research were collected as part of a national cattle
breeding programme, a process that conforms to routine animal husbandry procedures. No
experimental procedure was performed on live animals.

Data accessibility: The new whole-genome sequence data generated from 34 Criollo cattle for this
study is available at the European Nucleotide Archive (ENA) with the project accession ID
PRJEB65887.

Authors' contributions: J.A.W.: conceptualization, data curation, formal analysis, methodology,
software, visualization, and writing—original draft, and writing—review and editing; S.I.N.: data
curation, formal analysis, methodology, and writing—review and editing; I.A.S.R., G.P.H., and
J.F.O'G.: formal analysis, methodology, software, and writing—review and editing; J.M.F.,
A.M.P.O'B., and A.J.L-H.: conceptualization, resources, and writing—review and editing; J.A.B.:
methodology, and writing—review and editing; T.S.S.: conceptualization, funding acquisition,
project administration, resources, and writing—review and editing; L.A.F.F.: conceptualization,
data curation, methodology, resources, and writing—review and editing; M.S-T.; conceptualization,
methodology, supervision, and writing—review and editing; D.E.M.: conceptualization, funding
acquisition, project administration, resources, supervision, writing—original draft and writing—
review and editing.

Conflict of interest declaration: The authors declare no competing interests.

Funding: J.A.W. was supported by Science Foundation Ireland (SFI) and Acceligen/Recombinetics
Inc. through the SFI Centre for Research Training in Genomics Data Science under grant no.
18/CRT/6214. G.P.M. and D.E.M. were supported by an SFI Investigator Award (SFI/15/IA/3154).
T.S.S. and D.E.M were supported by a United States Department of Agriculture (USDA) and
Department of Agriculture, Food, and the Marine (DAFM) US-Ireland Research and Development
Partnership grant (TARGET-TB, 17/RD/USROI/52).

Acknowledgements: We thank Thomas Hall for advice on bioinformatics and computational
genomics methodologies.

REFERENCES

[1] Rouse, J.E. 1977 *The Criollo: Spanish Cattle in the Americas*, University of Oklahoma Press.

[2] Rodero, A., Delgado, J.V. & Rodero. 1992 Primitive Andalusian livestock and their implications in the discovery of America. *Arch. Zootec.* **41**, 383-400.

[3] Willham, R.L. 1982 Genetic improvement of beef cattle in the United States: cattle, people and their interaction. *J. Anim. Sci.* **54**, 659-666. (doi:10.2527/jas1982.543659x).

[4] Tobar, K.M.C., Álvarez, D.C. & Franco, L.A. 2020 Genome-wide association studies in sheep from Latin America. Review. *Revista Mexicana de Ciencias Pecuarias* **11**. (doi:10.22319/rmcp.v11i3.5372sander).

[5] Taylor, W.T.T., Librado, P., Hunska Tašunke Icu, M., Shield Chief Gover, C., Arterberry, J., Luta Wi☐, A., Nujipi, A., Omniya, T., Gonzalez, M., Means, B., et al. 2023 Early dispersal of domestic horses into the Great Plains and northern Rockies. *Science* **379**, 1316-1323. (doi:10.1126/science.adc9691).

[6] Sanders, J.O. 1980 History and development of zebu cattle in the United States. *J. Anim. Sci.* **50**, 1188-1200. (doi:10.2527/jas1980.5061188x).

[7] Cymbron, T., Loftus, R.T., Malheiro, M.I. & Bradley, D.G. 1999 Mitochondrial sequence variation suggests an African influence in Portuguese cattle. *Proc. Biol. Sci.* **266**, 597-603. (doi:10.1098/rspb.1999.0678).

[8] Miretti, M.M., Dunner, S., Naves, M., Contel, E.P. & Ferro, J.A. 2004 Predominant African-derived mtDNA in Caribbean and Brazilian Creole cattle is also found in Spanish cattle (*Bos taurus*). *J. Hered.* **95**, 450-453. (doi:10.1093/jhered/esh070).

[9] da Fonseca, R.R., Ureña, I., Afonso, S., Pires, A.E., Jørsboe, E., Chikhi, L. & Ginja, C. 2019 Consequences of breed formation on patterns of genomic diversity and differentiation: the case of highly diverse peripheral Iberian cattle. *BMC Genomics* **20**, 334. (doi:10.1186/s12864-019-5685-2).

[10] Decker, J.E., McKay, S.D., Rolf, M.M., Kim, J., Molina Alcala, A., Sonstegard, T.S., Hanotte, O., Gotherstrom, A., Seabury, C.M., Praharani, L., et al. 2014 Worldwide patterns of ancestry, divergence, and admixture in domesticated cattle. *PLoS Genet.* **10**, e1004254. (doi:10.1371/journal.pgen.1004254).

[11] Pitt, D., Bruford, M.W., Barbato, M., Orozco-terWengel, P., Martínez, R. & Sevane, N. 2019 Demography and rapid local adaptation shape Creole cattle genome diversity in the tropics. *Evol. Appl.* **12**, 105-122. (doi:10.1111/eva.12641).

[12] Magee, D.A., Meghen, C., Harrison, S., Troy, C.S., Cymbron, T., Gaillard, C., Morrow, A., Maillard, J.C. & Bradley, D.G. 2002 A partial African ancestry for the Creole cattle populations of the Caribbean. *J. Hered.* **93**, 429-432. (doi:10.1093/jhered/93.6.429).

[13] Ginja, C., Gama, L.T., Cortés, O., Burriel, I.M., Vega-Pla, J.L., Penedo, C., Sponenberg, P., Cañón, J., Sanz, A., do Egito, A.A., et al. 2019 The genetic ancestry of American Creole cattle inferred from uniparental and autosomal genetic markers. *Sci. Rep.* **9**, 11486. (doi:10.1038/s41598-019-47636-0).

[14] Gautier, M. & Naves, M. 2011 Footprints of selection in the ancestral admixture of a New World Creole cattle breed. *Mol. Ecol.* **20**, 3128-3143. (doi:10.1111/j.1365-294X.2011.05163.x).

[15] Ben-Jemaa, S., Adam, G., Boussaha, M., Bardou, P., Klopp, C., Mandonnet, N. & Naves, M. 2023 Whole genome sequencing reveals signals of adaptive admixture in Creole cattle. *Sci. Rep.* **13**, 12155. (doi:10.1038/s41598-023-38774-7).

[16] Bradley, D.G., MacHugh, D.E., Cunningham, P. & Loftus, R.T. 1996 Mitochondrial diversity
and the origins of African and European cattle. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 5131-5135.
(doi:10.1073/pnas.93.10.5131).

[17] MacHugh, D.E., Shriver, M.D., Loftus, R.T., Cunningham, P. & Bradley, D.G. 1997
Microsatellite DNA variation and the evolution, domestication and phylogeography of taurine and
zebu cattle (*Bos taurus* and *Bos indicus*). *Genetics* **146**, 1071-1086.
(doi:10.1093/genetics/146.3.1071).

[18] Hanotte, O., Bradley, D.G., Ochieng, J.W., Verjee, Y., Hill, E.W. & Rege, J.E.O. 2002 African
pastoralism: Genetic imprints of origins and migrations. *Science* **296**, 336-339.
(doi:10.1126/science.1069878).

[19] Kim, K., Kwon, T., Dessie, T., Yoo, D., Mwai, O.A., Jang, J., Sung, S., Lee, S., Salim, B.,
Jung, J., et al. 2020 The mosaic genome of indigenous African cattle as a unique genetic resource
for African pastoralism. *Nat. Genet.* **52**, 1099-1110. (doi:10.1038/s41588-020-0694-2).

[20] Ward, J.A., McHugo, G.P., Dover, M.J., Hall, T.J., Sonstegard, T.S., Bradley, D.G., Frantz,
L.A.F., Salter-Townshend, M. & MacHugh, D.E. 2022 Genome-wide local ancestry and evidence
for mitonuclear coadaptation in African hybrid cattle populations. *iScience* **25**.
(doi:10.1016/j.isci.2022.104672).

[21] de Alba, J. 1987 Criollo Cattle of Latin America. In *Animal Genetic Resources: Strategies for*
Improved Use and Conservation (ed. J. Hodges), pp. 17-39, Food and Agriculture Organization of
the United Nations.

[22] Landaeta-Hernández, A.J., Zambrano-Nava, S., Verde, O., Pinto-Santini, L., Montero-
Urdaneta, M., Hernández-Fonseca, J.P., Fuenmayor-Morales, C., Sonstegard, T.S., Huson, H.J. &
Olson, T.A. 2021 Heat stress response in slick vs normal-haired Criollo Limonero heifers in a
tropical environment. *Trop. Anim. Health Prod.* **53**, 445. (doi:10.1007/s11250-021-02856-3).

[23] De León, C., Martínez, R., Rocha, J.F. & Darghan, A.E. 2021 Selection of genomic regions
and genes associated with adaptation and fertility traits in two Colombian Creole cattle breeds.
Genet. Mol. Res. **20**. (doi:10.4238/gmr18882).

[24] Martínez, A.M., Gama, L.T., Cañón, J., Ginja, C., Delgado, J.V., Dunner, S., Landi, V.,
Martín-Burriel, I., Penedo, M.C., Rodellar, C., et al. 2012 Genetic footprints of Iberian cattle in
America 500 years after the arrival of Columbus. *PLoS ONE* **7**, e49066.
(doi:10.1371/journal.pone.0049066).

[25] Caivio-Nasner, S., López-Herrera, A., González-Herrera, L.G. & Rincón, J.C. 2021 Frequency
of genotypic markers for genetic disorders, colour, polledness, and major genes in Blanco
Orejinegro cattle. *Trop. Anim. Health Prod.* **53**, 546. (doi:10.1007/s11250-021-02990-y).

[26] Rocha, J.F., Martínez, R., López-Villalobos, N. & Morris, S.T. 2019 Tick burden in *Bos taurus*
cattle and its relationship with heat stress in three agroecological zones in the tropics of Colombia.
Parasit. Vectors **12**, 73. (doi:10.1186/s13071-019-3319-9).

[27] Littlejohn, M.D., Henty, K.M., Tiplady, K., Johnson, T., Harland, C., Lopdell, T., Sherlock,
R.G., Li, W., Lukefahr, S.D., Shanks, B.C., et al. 2014 Functionally reciprocal mutations of the
prolactin signalling pathway define hairy and slick cattle. *Nat. Commun.* **5**, 5861.
(doi:10.1038/ncomms6861).

[28] Porto-Neto, L.R., Bickhart, D.M., Landaeta-Hernandez, A.J., Utsunomiya, Y.T., Pagan, M.,
Jimenez, E., Hansen, P.J., Dikmen, S., Schroeder, S.G., Kim, E.S., et al. 2018 Convergent evolution
of slick coat in cattle through truncation mutations in the prolactin receptor. *Front. Genet.* **9**, 57.
(doi:10.3389/fgene.2018.00057).

[29] Anderson, D.M., Estell, R.E., Gonzalez, A.L., Cibils, A.F. & Torell, L.A. 2015 Criollo cattle:
heritage genetics for arid landscapes. *Rangelands* **37**, 62-67. (doi:10.1016/j.rala.2015.01.006).

[30] Encina Ruiz, R., Saucedo-Uriarte, J.A., Portocarrero-Villegas, S.M., Quispe-Ccasa, H.A. &
Cayo-Colca, I.S. 2021 Zoometric characterization of Creole cows from the southern Amazon region
of Peru. *Diversity* **13**, 510. (doi:10.3390/d13110510).

[31] Ginja, C., Gama, L.T., Cortes, O., Delgado, J.V., Dunner, S., García, D., Landi, V., Martín-
Burriel, I., Martínez-Martínez, A., Penedo, M.C.T., et al. 2013 Analysis of conservation priorities of
Iberoamerican cattle based on autosomal microsatellite markers. *Genet. Sel. Evol.* **45**, 35.
(doi:10.1186/1297-9686-45-35).

[32] Food and Agriculture Organization. 2015 *The Second Report on the State of the World's*
Animal Genetic Resources for Food and Agriculture. Rome, Italy, edited by B.D. Scherf & D.
Pilling. FAO Commission on Genetic Resources for Food and Agriculture Assessments.

[33] Rosen, B.D., Bickhart, D.M., Schnabel, R.D., Koren, S., Elsik, C.G., Tseng, E., Rowan, T.N.,
Low, W.Y., Zimin, A., Couldrey, C., et al. 2020 De novo assembly of the cattle reference genome
with single-molecule sequencing. *Gigascience* **9**. (doi:10.1093/gigascience/giaa021).

[34] Vasimuddin, M., Misra, S., Li, H. & Aluru, S. 2019 Efficient architecture-aware acceleration
of BWA-MEM for multicore ssemystems. *2019 IEEE International Parallel and Distributed*
Processing Symposium (IPDPS), 314-324. (doi:10.1109/IPDPS.2019.00041).

[35] Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., et al. 2021 Twelve years of SAMtools and BCFtools.
Gigascience **10**. (doi:10.1093/gigascience/giab008).

[36] Van der Auwera, G.A. & O'Connor, B.D. 2020 *Genomics in the Cloud: using Docker, GATK,*
and WDL in Terra. First edition. ed. Sebastopol, CA, O'Reilly Media.

[37] Felius, M. 1995 *Cattle Breeds: An Encyclopedia*. Doetinchem, Netherlands, Misset.

[38] Cingolani, P., Platts, A., Wang le, L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X. &
Ruden, D.M. 2012 A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-
3. *Fly (Austin)* **6**, 80-92. (doi:10.4161/fly.19695).

[39] Upadhyay, M.R., Chen, W., Lenstra, J.A., Goderie, C.R.J., MacHugh, D.E., Park, S.D.E.,
Magee, D.A., Matassino, D., Ciani, F., Megens, H.J., et al. 2017 Genetic origin, admixture and
population history of aurochs (*Bos primigenius*) and primitive European cattle. *Heredity* **118**, 169-
176. (doi:10.1038/hdy.2016.79).

[40] Sempéré, G., Moazami-Goudarzi, K., Eggen, A., Laloë, D., Gautier, M. & Flori, L. 2015
WIDDE: a Web-Interfaced next generation database for genetic diversity exploration, with a first
application in cattle. *BMC Genomics* **16**, 940. (doi:verit).

[41] Zimin, A.V., Delcher, A.L., Florea, L., Kelley, D.R., Schatz, M.C., Puiu, D., Hanrahan, F.,
Pertea, G., Van Tassell, C.P., Sonstegard, T.S., et al. 2009 A whole-genome assembly of the
domestic cow, *Bos taurus*. *Genome Biol.* **10**, R42. (doi:10.1186/gb-2009-10-4-r42).

[42] Riggio, V., Tijjani, A., Callaby, R., Talenti, A., Wragg, D., Obishakin, E.T., Ezeasor, C.,
Jongejan, F., Ogo, N.I., Aboagye-Antwi, F., et al. 2022 Assessment of genotyping array
performance for genome-wide association studies and imputation in African cattle. *Genet. Sel. Evol.*
54, 58. (doi:10.1186/s12711-022-00751-5).

[43] Nicolazzi, E.L., Caprera, A., Nazzicari, N., Cozzi, P., Strozzi, F., Lawley, C., Pirani, A., Soans,
C., Brew, F., Jorjani, H., et al. 2015 SNPchiMp v.3: integrating and standardizing single nucleotide
polymorphism data for livestock species. *BMC Genomics* **16**, 283. (doi:10.1186/s12864-015-1497-
1).

[44] Schnabel, R.D. 2018 ARS-UCD1.2 Cow Genome Assembly: mapping of all existing variants.
www.animalgenome.org/repository/cattle/UMC_bovine_coordinates.

[45] Lawrence, M., Gentleman, R. & Carey, V. 2009 rtracklayer: an R package for interfacing with
genome browsers. *Bioinformatics* **25**, 1841-1842. (doi:10.1093/bioinformatics/btp328).

[46] Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M. & Lee, J.J. 2015 Second-
generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7.
(doi:10.1186/s13742-015-0047-8).

[47] R Core Team. 2022 R: A language and environment for statistical computing. (Vienna,
Austria, R Foundation for Statistical Computing. http://www.R-project.org.

[48] Wickham, H. 2016 *ggplot2: Elegant Graphics for Data Analysis*, Springer Publishing
Company, New York, USA; wri p.

[49] Raj, A., Stephens, M. & Pritchard, J.K. 2014 fastSTRUCTURE: variational inference of
population structure in large SNP data sets. *Genetics* **197**, 573-589.
(doi:10.1534/genetics.114.164350).

[50] Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker,
R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. 2011 The variant call format and VCFtools.
Bioinformatics **27**, 2156-2158. (doi:10.1093/bioinformatics/btr330).

[51] Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T.,
Webster, T. & Reich, D. 2012 Ancient admixture in human history. *Genetics* **192**, 1065-1093.
(doi:10.1534/genetics.112.145037).

[52] Wright, S. 1950 Genetical structure of populations. *Nature* **166**, 247-249.
(doi:10.1038/166247a0).

[53] Szpiech, Z.A. & Hernandez, R.D. 2014 selscan: An efficient multithreaded program to perform
EHH-based scans for positive selection. *Mol. Biol. Evol.* **31**, 2824-2827.
(doi:10.1093/molbev/msu211).

[54] Randhawa, I.A.S., Khatkar, M.S., Thomson, P.C. & Raadsma, H.W. 2014 Composite selection
signals can localize the trait specific genomic regions in multi-breed populations of cattle and
sheep. *BMC Genet.* **15**, 34. (doi:10.1186/1471-2156-15-34).

[55] Verity, R., Collins, C., Card, D.C., Schaal, S.M., Wang, L. & Lotterhos, K.E. 2017 minotaur:
A platform for the analysis and visualization of multivariate results from genome scans with R
Shiny. *Mol. Ecol. Resour.* **17**, 33-43. (doi:sosa).

[56] Han, H., McGivney, B.A., Farries, G., Katz, L.M., MacHugh, D.E., Randhawa, I.A.S. & Hill,
E.W. 2020 Selection in Australian Thoroughbred horses acts on a locus associated with early two-
year old speed. *PLoS ONE* **15**, e0227212. (doi:10.1371/journal.pone.0227212).

[57] Browett, S., McHugo, G., Richardson, I.W., Magee, D.A., Park, S.D.E., Fahey, A.G., Kearney,
J.F., Correia, C.N., Randhawa, I.A.S. & MacHugh, D.E. 2018 Genomic characterisation of the
indigenous Irish Kerry cattle breed. *Front. Genet.* **9**. (doi:10.3389/fgene.2018.00051).

[58] Han, H., Randhawa, I.A.S., MacHugh, D.E., McGivney, B.A., Katz, L.M., Dugarjaviin, M. &
Hill, E.W. 2023 Selection signatures for local and regional adaptation in Chinese Mongolian horse
breeds reveal candidate genes for hoof health. *BMC Genomics* **24**, 35. (doi:10.1186/s12864-023-
09116-8).

[59] Elsik, C.G., Unni, D.R., Diesh, C.M., Tayal, A., Emery, M.L., Nguyen, H.N. & Hagen, D.E.
2016 Bovine Genome Database: new tools for gleaning function from the *Bos taurus* genome.
Nucleic Acids Res. **44**, D834-D839. (doi:10.1093/nar/gkv1077).

[60] Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H. & Vilo, J. 2019
g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019
update). *Nucleic Acids Res.* **47**, W191-W198. (doi:10.1093/nar/gkz369).

[61] Sollero, B.P., Junqueira, V.S., Gomes, C.C.G., Caetano, A.R. & Cardoso, F.F. 2017 Tag SNP
selection for prediction of tick resistance in Brazilian Braford and Hereford cattle breeds using
Bayesian methods. *Genet. Sel. Evol.* **49**, 1-15. (doi:10.1186/s12711-017-0325-2).

[62] Chen, N., Cai, Y., Chen, Q., Li, R., Wang, K., Huang, Y., Hu, S., Huang, S., Zhang, H., Zheng,
Z., et al. 2018 Whole-genome resequencing reveals world-wide ancestry and adaptive introgression
events of domesticated cattle in East Asia. *Nat. Commun.* **9**, 2337. (doi:10.1038/s41467-018-04737-
0).

[63] Wang, K., Lenstra, J.A., Liu, L., Hu, Q., Ma, T., Qiu, Q. & Liu, J. 2018 Incomplete lineage
sorting rather than hybridization explains the inconsistent phylogeny of the wisent. *Commun. Biol.*
1, 169. (doi:10.1038/s42003-018-0176-6).

[64] Wu, D.D., Ding, X.D., Wang, S., Wojcik, J.M., Zhang, Y., Tokarska, M., Li, Y., Wang, M.S.,
Faruque, O., Nielsen, R., et al. 2018 Pervasive introgression facilitated domestication and
adaptation in the *Bos* species complex. *Nat. Ecol. Evol.* **2**, 1139-1145. (doi:10.1038/s41559-018-
0562-y).

[65] Gebrehiwot, N.Z., Strucken, E.M., Aliloo, H., Marshall, K. & Gibson, J.P. 2020 The patterns
of admixture, divergence, and ancestry of African cattle populations determined from genome-wide
SNP data. *BMC Genomics* **21**, 869. (doi:10.1186/s12864-020-07270-x).

[66] Freeman, A.R., Meghen, C.M., MacHugh, D.E., Loftus, R.T., Achukwi, M.D., Bado, A.,
Sauveroche, B. & Bradley, D.G. 2004 Admixture and diversity in West African cattle populations.
Mol. Ecol. **13**, 3477-3487. (doi:10.1111/j.1365-294X.2004.02311.x).

[67] Toro, R.J.E. 2020 Evaluación de herramientas para desarrollar un programa de mejoramiento
genético en la raza Blanco Orejinegro (BON) [PhD thesis]. Medellín, Colombia, University of
Antioquia.

[68] Pearson, L. 1968 A note on the history of black-eared white cattle. *Agric. Hist. Rev.* **16**, 159-
160.

[69] Oke, O.E., Oso, O.M., Logunleko, M.O., Uyanga, V.A., Akinyemi, F., Okeniyi, F.A., Akosile,
O.A., Baloyi, J.J. & Onagbesan, O.M. 2022 Adaptation of the White Fulani cattle to the tropical
environment. *J. Therm. Biol.* **110**, 103372. (doi:10.1016/j.jtherbio.2022.103372).

[70] Brenig, B., Beck, J., Floren, C., Bornemann-Kolatzki, K., Wiedemann, I., Hennecke, S.,
Swalve, H. & Schütz, E. 2013 Molecular genetics of coat colour variations in White Galloway and
White Park cattle. *Anim. Genet.* **44**, 450-453. (doi:10.1111/age.12029).

[71] Senczuk, G., Guerra, L., Mastrangelo, S., Campobasso, C., Zoubeyda, K., Imane, M., Marletta,
D., Kusza, S., Karsli, T., Gaouar, S.B.S., et al. 2020 Fifteen shades of grey: combined analysis of
genome-wide SNP data in Steppe and Mediterranean grey cattle sheds new light on the molecular
basis of coat color. *Genes (Basel)* **11**, 932. (doi:10.3390/genes11080932).

[72] Senczuk, G., Landi, V., Mastrangelo, S., Persichilli, C., Pilla, F. & Ciani, E. 2022 Seven
shades of grey: A follow-up study on the molecular basis of coat colour in indicine grey cattle using
genome-wide SNP data. *Genes (Basel)* **13**, 1601. (doi:10.3390/genes13091601).

[73] Han, S.-H., Cho, I.-C., Kim, J.-H., Ko, M.-S., Kim, Y.-H., Kim, E.-Y., Park, S.-P. & Lee, S.-S.
2011 Coat color patterns and genotypes of *Extension* and *Agouti* in Hanwoo and Jeju black cattle. *J.*
Life Sci. **21**. (doi:10.5352/JLS.2011.21.4.494).

[74] Druml, T., Grilz-Seger, G., Neuditschko, M., Horna, M., Ricard, A., Pausch, H. & Brem, G.
2018 Novel insights into Sabino1 and splashed white coat color patterns in horses. *Anim. Genet.* **49**,
249-253. (doi:10.1111/age.12657).

[75] Rothammer, S., Kunz, E., Seichter, D., Krebs, S., Wassertheurer, M., Fries, R., Brem, G. &
Medugorac, I. 2017 Detection of two non-synonymous SNPs in *SLC45A2* on BTA20 as candidate

causal mutations for oculocutaneous albinism in Braunvieh cattle. *Genet. Sel. Evol.* **49**, 73. (doi:10.1186/s12711-017-0349-7).

[76] Voß, K., Blaj, I., Tetens, J.L., Thaller, G. & Becker, D. 2022 Roan coat color in livestock. *Anim. Genet.* **53**, 549-556. (doi:10.1111/age.13240).

[77] Goud, T.S., Upadhyay, R.C., Onteru, S.K., Pichili, V.B.R. & Chadipiralla, K. 2020 Identification and sequence characterization of melanocortin 1 receptor gene (*MC1R*) in *Bos indicus* versus (*Bos taurus* X *Bos indicus*). *Anim. Biotechnol.* **31**, 283-294. (doi:10.1080/10495398.2019.1585866).

[78] Landaeta-Hernández, A., Zambrano-Nava, S., Hernández-Fonseca, J.P., Godoy, R., Calles, M., Iragorri, J.L., Añez, L., Polanco, M., Montero-Urdaneta, M. & Olson, T. 2011 Variability of hair coat and skin traits as related to adaptation in Criollo Limonero cattle. *Trop. Anim. Health Prod.* **43**, 657-663. (doi:10.1007/s11250-010-9749-1).

[79] Landaeta-Hernández, A., Zambrano, S., Marin, N. & Moreno, Y. 2011 Criollos Latinoamericanos. In *Workshop Jornadas de Actualización en Ganadería Tropical* (pp. 52-62, Asociación de Ganaderos Alberto Adriani (ASODEGAA).

[80] Parra-Cortés, R.I. & Magaña-Magaña, M.Á. 2021 Características técnico-económicas de sistemas de producción bovina de las razas criollas colombianas Romosinuano y Hartón del Valle. *Rev. MVZ Cordoba* **26**, e2079. (doi:10.21897/rmvz.2079).

[81] Xu, L., Bickhart, D.M., Cole, J.B., Schroeder, S.G., Song, J., Van Tassell, C.P., Sonstegard, T.S. & Liu, G.E. 2015 Genomic signatures reveal new evidences for selection of important traits in domestic cattle. *Mol. Biol. Evol.* **32**, 711-725. (doi:10.1093/molbev/msu333).

[82] Mei, C., Wang, H., Liao, Q., Wang, L., Cheng, G., Wang, H., Zhao, C., Zhao, S., Song, J., Guang, X., et al. 2018 Genetic architecture and selection of Chinese cattle revealed by whole genome resequencing. *Mol. Biol. Evol.* **35**, 688-699. (doi:10.1093/molbev/msx322).

[83] Hansen, P.J. 2004 Physiological and cellular adaptations of zebu cattle to thermal stress. *Anim. Reprod. Sci.* **82-83**, 349-360. (doi:10.1016/j.anireprosci.2004.04.011).

[84] Hansen, P.J. 2020 Prospects for gene introgression or gene editing as a strategy for reduction of the impact of heat stress on production and reproduction in cattle. *Theriogenology* **154**, 190-202. (doi:10.1016/j.theriogenology.2020.05.010).

[85] Flórez Murillo, J.M., Landaeta-Hernández, A.J., Kim, E.-S., Bostrom, J.R., Larson, S.A., Pérez O'Brien, A.M., Montero-Urdaneta, M.A., Garcia, J.F. & Sonstegard, T.S. 2021 Three novel nonsense mutations of prolactin receptor found in heat-tolerant *Bos taurus* breeds of the Caribbean Basin. *Anim. Genet.* **52**, 132-134. (doi:10.1111/age.13027).

[86] Olson, T.A., Lucena, C., Chase, C.C., Jr. & Hammond, A.C. 2003 Evidence of a major gene influencing hair length and heat tolerance in *Bos taurus* cattle. *J. Anim. Sci.* **81**, 80-90. (doi:10.2527/2003.81180x).

[87] Dikmen, S., Alava, E., Pontes, E., Fear, J.M., Dikmen, B.Y., Olson, T.A. & Hansen, P.J. 2008 Differences in thermoregulatory ability between slick-haired and wild-type lactating Holstein cows in response to acute heat stress. *J. Dairy Sci.* **91**, 3395-3402. (doi:10.3168/jds.2008-1072).

[88] Dikmen, S., Khan, F.A., Huson, H.J., Sonstegard, T.S., Moss, J.I., Dahl, G.E. & Hansen, P.J. 2014 The *SLICK* hair locus derived from Senepol cattle confers thermotolerance to intensively managed lactating Holstein cows. *J. Dairy Sci.* **97**, 5508-5520. (doi:10.3168/jds.2014-8087).

[89] Sosa, F., Santos, J.E.P., Rae, D.O., Larson, C.C., Macchietto, M., Abrahante, J.E., Amaral, T.F., Denicol, A.C., Sonstegard, T.S. & Hansen, P.J. 2022 Effects of the SLICK1 mutation in *PRLR* on regulation of core body temperature and global gene expression in liver in cattle. *Animal* **16**, 100523. (doi:sing).

[90] Ben-Jemaa, S., Mastrangelo, S., Lee, S.-H., Lee, J.H. & Boussaha, M. 2020 Genome-wide scan
for selection signatures reveals novel insights into the adaptive capacity in local North African
cattle. *Sci. Rep.* **10**, 19466. (doi:10.1038/s41598-020-76576-3).

[91] Singh, A.K., Upadhyay, R.C., Chandra, G., Kumar, S., Malakar, D., Singh, S.V. & Singh,
M.K. 2020 Genome-wide expression analysis of the heat stress response in dermal fibroblasts of
Tharparkar (zebu) and Karan-Fries (zebu × taurine) cattle. *Cell Stress Chaperones* **25**, 327-344.
(doi:druml).

[92] Feder, M.E. & Hofmann, G.E. 1999 Heat-shock proteins, molecular chaperones, and the stress
response: evolutionary and ecological physiology. *Annu. Rev. Physiol.* **61**, 243-282.
(doi:10.1146/annurev.physiol.61.1.243).

[93] Hu, C., Yang, J., Qi, Z., Wu, H., Wang, B., Zou, F., Mei, H., Liu, J., Wang, W. & Liu, Q. 2022
Heat shock proteins: Biological functions, pathological roles, and therapeutic opportunities.
MedComm (2020) **3**, e161. (doi:10.1002/mco2.161).

[94] Peng, J., Li, Y., Wang, X., Deng, S., Holland, J., Yates, E., Chen, J., Gu, H., Essandoh, K., Mu,
X., et al. 2018 An Hsp20-FBXO4 axis regulates adipocyte function through modulating PPARγ
ubiquitination. *Cell Rep.* **23**, 3607-3620. (doi:10.1016/j.celrep.2018.05.065).

[95] Srikanth, K., Kwon, A., Lee, E. & Chung, H. 2017 Characterization of genes and pathways that
respond to heat stress in Holstein calves through transcriptome analysis. *Cell Stress Chaperones* **22**,
29-42. (doi:10.1007/s12192-016-0739-8).

[96] Hayashi, T., Asano, Y., Shintani, Y., Aoyama, H., Kioka, H., Tsukamoto, O., Hikita, M.,
Shinzawa-Itoh, K., Takafuji, K., Higo, S., et al. 2015 Higd1a is a positive regulator of cytochrome c
oxidase. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 1553-1558. (doi:10.1073/pnas.1419767112).

[97] Zhang, J., Long, K., Wang, J., Zhang, J., Jin, L., Tang, Q., Li, X., Ma, J., Li, M. & Jiang, A.
2022 Yak miR-2285o-3p attenuates hypoxia-induced apoptosis by targeting caspase-3. *Anim.*
Genet. **53**, 49-57. (doi:10.1111/age.13153).

[98] Terefe, E., Belay, G., Han, J., Hanotte, O. & Tijjani, A. 2022 Genomic adaptation of Ethiopian
indigenous cattle to high altitude. *Front. Genet.* **13**, 960234. (doi:10.3389/fgene.2022.960234).

[99] Zhang, X., Wang, D. & Liu, J. 2023 Hypoxia-inducible factor-1α is involved in the response to
heat stress in lactating dairy cows. *J. Therm. Biol.* **112**, 103460.
(doi:10.1016/j.jtherbio.2023.103460).

[100] Armstrong, E., Rodriguez Almeida, F.A., McIntosh, M.M., Poli, M., Cibils, A.F., Martínez-
Quintana, J.A., Félix-Portillo, M. & Estell, R.E. 2022 Genetic and productive background of Criollo
cattle in Argentina, Mexico, Uruguay and the United States. *J. Arid Environ.* **200**, 104722.
(doi:10.1016/j.jaridenv.2022.104722).

[101] Rojas-Espinoza, R., Macedo, R., Suaña, A., Delgado, A., Manrique, Y.P., Rodríguez, H.,
Quispe, Y.M., Perez-Guerra, U.H., Pérez-Durand, M.G. & García-Herreros, M. 2023 Phenotypic
characterization of Creole cattle in the Andean highlands using bio-morphometric measures and
zoometric indices. *Animals (Basel)* **13**. (doi:10.3390/ani13111843).

[102] Johnson, G.P., English, A.-M., Cronin, S., Hoey, D.A., Meade, K.G. & Fair, S. 2017
Genomic identification, expression profiling, and functional characterization of CatSper channels in
the bovine. *Biol. Reprod.* **97**, 302-312. (doi:10.1093/biolre/iox082).

[103] Guo, F., Yang, B., Ju, Z.H., Wang, X.G., Qi, C., Zhang, Y., Wang, C.F., Liu, H.D., Feng,
M.Y., Chen, Y., et al. 2014 Alternative splicing, promoter methylation, and functional SNPs of
sperm flagella 2 gene in testis and mature spermatozoa of Holstein bulls. *J. Reprod. Fertil.* **147**,
241-252. (doi:10.1530/REP-13-0343).

[104] Nantel, F., Monaco, L., Foulkes, N.S., Masquilier, D., LeMeur, M., Henriksén, K., Dierich,
A., Parvinen, M. & Sassone-Corsi, P. 1996 Spermiogenesis deficiency and germ-cell apoptosis in
CREM-mutant mice. *Nature* **380**, 159-162. (doi:10.1038/380159a0).

[105] Zapater, J.L., Lednovich, K.R., Khan, M.W., Pusec, C.M. & Layden, B.T. 2022 Hexokinase
domain-containing protein-1 in metabolic diseases and beyond. *Trends Endocrinol. Metab.* **33**, 72-
84. (doi:10.1016/j.tem.2021.10.006).

[106] Ren, F., Xi, H., Qiao, P., Li, Y., Xian, M., Zhu, D. & Hu, J. 2022 Single-cell transcriptomics
reveals male germ cells and Sertoli cells developmental patterns in dairy goats. *Front. Cell Dev.*
Biol. **10**, 944325. (doi:10.3389/fcell.2022.944325).

[107] Gamallat, Y., Fang, X., Mai, H., Liu, X., Li, H., Zhou, P., Han, D., Zheng, S., Liao, C., Yang,
M., et al. 2021 Bi-allelic mutation in *Fsip1* impairs acrosome vesicle formation and attenuates
flagellogenesis in mice. *Redox Biol.* **43**, 101969. (doi:10.1016/j.redox.2021.101969).

[108] Berisha, B., Rodler, D., Schams, D., Sinowatz, F. & Pfaffl, M.W. 2019 Prostaglandins in
superovulation induced bovine follicles during the preovulatory period and early corpus luteum.
Front. Endocrinol. (Lausanne) **10**, 467. (doi:10.3389/fendo.2019.00467).

[109] Caivio-Nasner, S., López-Herrera, A., González-Herrera, L.G. & Rincón, J.C. 2021 Diversity
analysis, runs of homozygosity and genomic inbreeding reveal recent selection in Blanco
Orejinegro cattle. *J. Anim. Breed. Genet.* **138**, 613-627. (doi:10.1111/jbg.12549).

[110] Alshawi, A., Essa, A., Al-Bayatti, S. & Hanotte, O. 2019 Genome analysis reveals genetic
admixture and signature of selection for productivity and environmental traits in Iraqi cattle. *Front.*
Genet. **10**, 609. (doi:10.3389/fgene.2019.00609).

[111] Fang, L., Hou, Y., An, J., Li, B., Song, M., Wang, X., Sørensen, P., Dong, Y., Liu, C., Wang,
Y., et al. 2016 Genome-wide transcriptional and post-transcriptional regulation of innate immune
and defense responses of bovine mammary gland to *Staphylococcus aureus*. *Front. Cell. Infect.*
Microbiol. **6**, 193. (doi:10.3389/fcimb.2016.00193).

[112] Kumar, L., Chou, J., Yee, C.S., Borzutzky, A., Vollmann, E.H., von Andrian, U.H., Park,
S.Y., Hollander, G., Manis, J.P., Poliani, P.L., et al. 2014 Leucine-rich repeat containing 8A
(LRRC8A) is essential for T lymphocyte development and function. *J. Exp. Med.* **211**, 929-942.
(doi:10.1084/jem.20131379).

[113] Rodriguez-Morales, A.J., Bonilla-Aldana, D.K., Idarraga-Bedoya, S.E., Garcia-Bustos, J.J.,
Cardona-Ospina, J.A. & Faccini-Martínez, Á.A. 2018 Epidemiology of zoonotic tick-borne diseases
in Latin America: Are we just seeing the tip of the iceberg? *F1000Res.* **7**, 1988.
(doi:10.12688/f1000research.17649.2).

[114] Ristic, M. & McIntyre, I. 1981 *Diseases of Cattle in the Tropics: Economic and Zoonotic*
Relevance. The Hague, Netherlands, Martinus Nijhoff Publishers.

[115] Jaimes-Dueñez, J., Triana-Chávez, O., Holguín-Rocha, A., Tobon-Castaño, A. & Mejía-
Jaramillo, A.M. 2018 Molecular surveillance and phylogenetic traits of *Babesia bigemina* and
Babesia bovis in cattle (*Bos taurus*) and water buffaloes (*Bubalus bubalis*) from Colombia. *Parasit.*
Vectors **11**, 510. (doi:10.1186/s13071-018-3091-2).

[116] Fetene, E., Leta, S., Regassa, F. & Büscher, P. 2021 Global distribution, host range and
prevalence of *Trypanosoma vivax*: a systematic review and meta-analysis. *Parasit. Vectors* **14**, 80.
(doi:10.1186/s13071-021-04584-x).

[117] Jaimes-Dueñez, J., Triana-Chávez, O. & Mejía-Jaramillo, A.M. 2018 Genetic, host and
environmental factors associated with a high prevalence of *Anaplasma marginale*. *Ticks Tick Borne*
Dis. **9**, 1286-1295. (doi:10.1016/j.ttbdis.2018.05.009).

[118] Naval-Sánchez, M., Porto-Neto, L.R., Cardoso, D.F., Hayes, B.J., Daetwyler, H.D., Kijas, J.
& Reverter, A. 2020 Selection signatures in tropical cattle are enriched for promoter and coding
regions and reveal missense mutations in the damage response gene *HELB*. *Genet. Sel. Evol.* **52**, 27.
(doi:10.1186/s12711-020-00546-6).

[119] Bhar, S., Zhou, F., Reineke, L.C., Morris, D.K., Khincha, P.P., Giri, N., Mirabello, L.,
Bergstrom, K., Lemon, L.D., Williams, C.L., et al. 2020 Expansion of germline *RPS20* mutation
phenotype to include Diamond-Blackfan anemia. *Hum. Mutat.* **41**, 1918-1930.
(doi:10.1002/humu.24092).

[120] Wang, R., Yoshida, K., Toki, T., Sawada, T., Uechi, T., Okuno, Y., Sato-Otsubo, A., Kudo,
K., Kamimaki, I., Kanezaki, R., et al. 2015 Loss of function mutations in *RPL27* and *RPS27*
identified by whole-exome sequencing in Diamond-Blackfan anaemia. *Br. J. Haematol.* **168**, 854-
864. (doi:10.1111/bjh.13229).

[121] Noyes, H.A., Alimohammadian, M.H., Agaba, M., Brass, A., Fuchs, H., Gailus-Durner, V.,
Hulme, H., Iraqi, F., Kemp, S., Rathkolb, B., et al. 2009 Mechanisms controlling anaemia in
Trypanosoma congolense infected mice. *PLoS ONE* **4**, e5170. (doi:10.1371/journal.pone.0005170).

[122] Martínez, R., Toro, R., Montoya, F., Burbano, M. & Ariza, F. 2005 Evaluación genética para
resistencia a brucelosis en ganado criollo colombiano BON. *Archivos de Zootecnia* **54**.

[123] Martínez, R., Dunner, S., Toro, R., Tobón, J., Gallego, J. & Cañón, J. 2010 Effect of
polymorphisms in the *Slc11a1* coding region on resistance to brucellosis by macrophages in vitro
and after challenge in two *Bos* breeds (Blanco Orejinegro and Zebu). *Genet. Mol. Biol.* **33**, 463-470.
(doi:10.1590/S1415-47572010000300014).

[124] Celli, J. 2019 The intracellular life cycle of *Brucella* spp. *Microbiol. Spectr.* **7**.
(doi:10.1128/microbiolspec.BAI-0006-2019).

[125] Huy, T.X.N., Nguyen, T.T., Kim, H., Reyes, A.W.B. & Kim, S. 2022 *Brucella* phagocytosis
mediated by pathogen-host interactions and their intracellular survival. *Microorganisms* **10**.
(doi:10.3390/microorganisms10102003).