Genomic Insights into the Population History and Adaptive Traits of Latin American Criollo Cattle

James A. Ward[1], Said I. Ng'ang'a[2,3], Imtiaz A. S. Randhawa[4], Gillian P. McHugo[1], John F. O'Grady[1], Julio M. Flórez[5,6], John A. Browne[1], Ana M. Pérez O'Brien[5], Antonio J. Landaeta-Hernández[7], José F. Garcia[6], Tad S. Sonstegard[5], Laurent A. F. Frantz[2,3], Michael Salter-Townshend[8], David E. MacHugh[1,9]

[1] Animal Genomics Laboratory, UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[2] Palaeogenomics Group, Department of Veterinary Sciences, Ludwig Maximilian University, Munich D-80539, Germany.

[3] School of Biological and Chemical Sciences, Queen Mary University of London, London E1 4NS, United Kingdom.

[4] Faculty of Science, The University of Queensland, Gatton, QLD 4343, Australia

[5] Acceligen, 3388 Mike Collins Drive, Eagan, MN, 55121, USA.

[6] Department of Preventive Veterinary Medicine and Animal Reproduction, School of Agricultural and Veterinarian Sciences, São Paulo State University (Unesp), Jaboticabal, Brazil.

[7] Unidad de Investigaciones Zootécnicas, Facultad de Ciencias Veterinarias, Universidad del Zulia, Maracaibo, Venezuela.

[8] UCD School of Mathematics and Statistics, University College Dublin, Dublin, D04 V1W8, Ireland.

[9] UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Dublin, D04 V1W8, Ireland.

Author for correspondence

David E. MacHugh

e-mail: david.machugh@ucd.ie

ABSTRACT

Criollo cattle, descendants of animals brought by Iberian colonists to the Americas, have been subject to centuries of natural and human-mediated selection in novel tropical agroecological zones. Consequently, these breeds have evolved distinct characteristics such as resistance to disease and exceptional heat tolerance. In addition to European taurine (*Bos taurus*) ancestry, it has been proposed that gene flow from African taurine and Asian indicine (*Bos indicus*) cattle has shaped the ancestry of Criollo cattle. In this study, we analysed Criollo breeds from Colombia and Venezuela using whole-genome sequencing (WGS) and single-nucleotide polymorphism (SNP) array data to examine population structure and admixture at high resolution. Analysis of genetic structure and ancestry components provided evidence for African taurine and Asian indicine admixture in Criollo cattle. In addition, using WGS data, we detected selection signatures associated with a myriad of adaptive traits, revealing genes linked to thermotolerance, reproduction, fertility, immunity, and distinct coat and skin coloration traits. This study underscores the remarkable adaptability of Criollo cattle and highlights the genetic richness and potential of these breeds in the face of climate change, habitat flux, and disease challenges. Further research is warranted to leverage these findings for more effective and sustainable cattle breeding programmes.

Keywords: cattle, Latin America, microevolution, population genomics, selection, thermotolerance

INTRODUCTION

The term "Criollo", with origins in the Portuguese "Crioulo", historically distinguished people born in the New World from those native to Iberia and was subsequently extended to livestock (cattle, sheep, horses, and goats). European livestock production in the Americas can be traced back to Columbus' second voyage in 1493, where cattle, among other animals, sourced from La Gomera in the Canary Islands were brought to the island of Hispaniola [1-3]. Adoption and growth of cattle agriculture on Hispaniola was rapid; however, the spread of animals to the South American continent was more gradual. This was documented by early colonial accounts, illustrating a transformative ecological and economic process that started in the Caribbean, then Mexico, and eventually reaching the Ilanos of Colombia and Venezuela, where by 1600, cattle ranching had become a significant industry [4, 5]. Over the ensuing five centuries, increasing numbers of cattle were exported to North and South America particularly as the process of European colonization intensified during the 18th and 19th centuries [6]. Evidence for a similar Iberian origin narrative has been described for sheep [7], pigs [8], and horses. For example, recent work has shown that historical and extant North American horse populations exhibit pronounced genetic affinities with Iberian horse populations [9]. The ancestry of modern South American cattle has also been shaped by indicine (zebu) cattle (*Bos indicus*) that were directly introduced from South Asia—primarily present-day India and Pakistan [10].

The extent of contributions from cattle outside Europe and the Indian subcontinent to the South and Central American cattle gene pools remains uncertain. Modern Iberian cattle populations, whose ancestors likely contributed the bulk of Latin American Criollo cattle ancestry have an African taurine cattle genomic component [11-13]. In parallel to this, it has also been posited that African cattle may have been directly introduced into South America alongside imports from Europe [14]. This hypothesis is supported by the presence of the T1-c mtDNA haplotype at a frequency of 8% in Criollo cattle; T1-c is prevalent in African cattle and found at low frequencies in Iberian populations [15]. It is also supported by recent findings of colonial era mitogenomes which reveal the presence of African

haplogroups in 17ᵗʰ century cattle from Mexico [16]. Recent work suggests that the degree of African

ancestry in Criollo cattle depends on the breed. Notably, the Guadeloupe breed, indigenous to the

Caribbean, traces almost 35% of its genetic ancestry back to African taurine cattle [17].

The cattle that were brought to the Americas from Iberia in the 15ᵗʰ and 16ᵗʰ centuries were

adapted to Mediterranean agroecologies and, over the following centuries, these populations have

evolved adaptations to tropical and arid environments, emerging as distinctive breeds with unique

heat tolerance and disease resistance traits [12, 18-21]. However, Criollo cattle are undervalued in

modern production systems and face gradual replacement by more productive commercial breeds,

which is eroding indigenous cattle genetic resources in Latin America [22]. Using genome-scale data,

this study aims to evaluate *B. taurus* and *B. indicus* ancestry and detect historical gene flow from

African cattle into the Blanco Orejinegro, Hartón del Valle, and Limonero Criollo breeds.

Additionally, we use high-resolution analyses of genetic structure, and selection signatures to provide

new information about the microevolutionary histories of these populations.

MATERIALS AND METHODS

Cattle samples, library preparation, high-throughput sequencing, and variant calling

In total, 34 Criollo cattle were used to generate high-quality de novo whole-genome sequence

(WGS) data sets. Coat hair follicles obtained from 24 Colombian and Venezuelan cattle and semen

straws from 10 Colombian AI bulls were used to obtain genomic DNA using standard commercial

procedures for livestock DNA extraction and purification (Weatherbys Scientific, Co. Kildare,

Ireland). The WGS data generation was performed by sequencing to a mean depth of 20× using two

commercial service providers (Novogene, Cambridge, UK and Macrogen, Seoul, South Korea).

Figure 1 displays the geographical origins of the three main Criollo breeds in this study.

Publicly available WGS data was also assembled for 37 European taurine, 13 American

indicine, and 13 American Criollo cattle. Sequence reads were mapped to the bovine reference

genome (ARS-UCD1.2) [23] using the default parameters for bwa-mem2 v.2.2.1 [24]. Samtools

v.1.13 [25] was used to sort bam files and create index files. Potential PCR duplicates were identified

using the "MarkDuplicates" function of Picard v.2.27.1 (http://broadinstitute.github.io/picard). The

"BaseRecalibrator" and "ApplyBQSR" functions of the Genome Analysis Toolkit (GATK) v.4.2.6.1

[26] were used to perform base quality score recalibration.

Figure 1. Approximate geographical origins of the animals from three Criollo breeds that were used

to generate new whole-genome sequence data. Map modified from original by Mapswire

(https://mapswire.com) (Creative Commons Attribution 4.0 International License CC-BY 4.0).

Individual cattle art images modified from Felius [27] with permission of the author.

Candidate SNPs were called from the bam files using the "HaplotypeCaller" function in GATK

with the "--emit-ref-confidence GVCF" option. Individual GVCF files were merged using the

"GenomicsDBImport" function. SNPs from the genomics database were called and selected using

"GenotypeGVCFs" and the "SelectVariants" function. The "VariantFiltration" function of GATK

was used to avoid possible false-positive calls according to GATK best practices: SNP clusters with

"--cluster-size 3" and "--cluster-window-size 10", quality by depth "QD <2", read position rank sum

test "ReadPosRankSum < -8", phred-scaled variant quality score "QUAL < 30", Fisher strand "FS >

60", mapping quality "MQ < 40", mapping quality rank sum test "MQRankSum < -12.5" and strand

odds ratio "SOR > 3". Triallelic SNPs and those with a minor allele frequency < 0.01 and/or missing

genotype rates over > 0.1 were filtered out. The SNPs remaining in each WGS data set were then

annotated according to their positions using SnpEff [28]. The combined panel of 97 cattle samples

with WGS data are detailed in electronic supplementary material, table S1.

BovineHD 777k array data

For broader comparative population genomics analyses, publicly available BovineHD

777k SNP array data were used. These were 91 European taurine cattle representing two breeds, 82

Asian and American indicine cattle representing four breeds [29, 30], 45 Iberian cattle representing

13 breeds [31], and 31 African taurine cattle representing two breeds [29, 32]. In addition, 12 Criollo

cattle representing the Senepol breed were obtained from the WIDDE database [33]. SNP positions

were updated from the UMD3.1 assembly [34] to ARS-UCD1.2 following Riggio et al. [35].

Subsequent processing of these data was conducted using PLINK v1.90b6.25 [36] and R version 4.1.3

[37]. The combined panel of 261 cattle samples with BovineHD 777k SNP data are detailed in

electronic supplementary material, table S2.

Population differentiation, genetic structure, and demographic analyses

Principal component analysis (PCA) for all animals (WGS and BovineHD genotype data) were

performed using PLINK v1.90b6.25 and the results were plotted using ggplot2 v3.4.0 [38] in the R

v4.1.3 environment. The genetic structure of each population was visualized using ADMIXTURE

v1.3.0 [39] with $K = 3$–6 unsupervised modelled ancestries and ggplot2 v3.4.0. Demographic

modelling of current and historical effective population size (N_e) trends was conducted using the

SNeP software tool [40] with WGS data for the Blanco Orejinegro, Hartón del Valle, Limonero,

N'Dama, and Holstein populations.

Detection of historical admixture

Using ADMIXTOOLS [41] with default parameters, we calculated f_3 statistics using the

combined WGS and BovineHD genotype data set to test for evidence that three Criollo breeds

(Blanco Orejinegro, Hartón del Valle, and Limonero) and three Iberian breeds (Limia, Pajuna, and

Sayaguesa) are derived from the admixture of two sets of populations (European taurine populations

and indicine cattle, and European taurine populations and African taurine cattle). A significant

negative f_3 value is considered evidence of historical admixture in the target population (Criollo or

Iberian). For visualization of the f_3 statistics, we plotted the f_3 values against the population trios with

error bars indicating the standard errors using ggplot2 v3.4.0 [38].

Test statistics for detection of selection signatures

For the selection signatures analysis, only the WGS data set was used, which included

approximately 9.7 million SNPs genotyped in 97 cattle. SNP-by-SNP F_{ST} values were calculated for

each Criollo breed versus a panel of Holstein cattle using VCFtools v.0.1.16 [42]. XP-EHH statistics

were calculated using the program selscan v.1.2.0 [43] with default parameters, where each Criollo

breed was also compared to the Holstein panel. The unstandardized XP-EHH values were then

normalized using the --norm function in selscan. We calculated the ΔSAF (directional change in

selected allele frequency) using the --freq function of VCFtools to generate allele frequencies, then

using a custom R script we calculated the ΔSAF for each Criollo breed versus the Holstein panel.

Similarly to the XPEHH values, ΔSAF values were standardized to $Z \sim N(0,1)$ as described by

Randhawa et al. [44].

Implementing the composite selection signals (CSS) methodology

To obtain composite selection signals (CSS) statistics [44] for each Criollo breed, the F_{ST}, XP-EHH, and ΔSAF statistics were combined by SNP ID (chromosome:position), which produced final sets of SNPs for the CSS analysis of 9,344,833, 9,465,007 and 9,660,647 million SNPs, for the Blanco Orejinegro, Hartón del Valle, and Limonero breeds, respectively. Following this, the MINOTAUR R package [45] was used to calculate CSS statistics, which were then averaged over 20 kb windows to reduce spurious signals. In addition, a signal was only considered significant if at least one SNP in the 0.1% of CSS scores was flanked by at least five other SNPs in the top 1% of CSS values [44, 46].

Identification of genomic regions under selection

From the top 0.1% of SNPs identified, genomic intervals in the format chromosome:start-end were extracted and the BovineMine resource (v1.6) [47] was then used to find genes ± 500 kb from the start and end of these regions. The functional relevance of these CSS-derived gene sets was then evaluated using the scientific literature and the g:Profiler software tool [48], which was used to identify overrepresented gene ontology (GO) terms for each of the CSS-derived gene sets by breed. Following this, BovineMine v1.6 was used to systematically identify previously described quantitative trait loci (QTLs) for production, health, and welfare traits that are located within the CSS cluster regions for each breed.

RESULTS

Population differentiation, genetic structure, and admixture in Criollo cattle

The merged WGS and remapped BovineHD 777K BeadChip datasets yielded 302,012 shared SNPs for 308 individual animals. A PCA of these SNPs demonstrated that PC1 and PC2 accounted for 44.7% and 12.9% of the variation observed across the first 20 PCs, with PC1 representing the split between the *B. taurus* and *B. indicus* lineages and PC2 the split between the European and African taurine lineages (figure 2*a*). Genetic structure was analysed using ADMIXTURE, with cluster numbers (K) ranging from 3 to 6 (figure 2*b*). At $K = 3$, we recovered the three clusters observed in the PCA, which delineated European *B. taurus*, African *B. taurus*, and *B. indicus* ancestry components, respectively. The subsequent split at $K = 4$ captured an additional cluster within the European taurine populations that differentiated Jersey cattle. At $K = 5$ a Holstein cattle ancestry component was evident and at $K = 6$ a split was observed that divided *B. indicus* ancestry into two clusters.

Figure 2. Genetic diversity and admixture in Criollo and comparator cattle populations. (*a*) Results

of the principal component analysis (PCA) for 308 animals (47 from whole-genome sequence data

and 261 from high-density SNP array data) with a shared set of 302,012 SNPs. The PCA plot shows

the coordinates for each animal based on the first two principal components with the major breed

groupings highlighted. Principal component 1 (PC1) differentiates the *Bos taurus* and *Bos indicus*

lineages, whereas PC2 separates the African and European taurine groups. A histogram plot also

shows the variance contributions for the first 20 PCs, with PC1 and PC2 accounting for 44.7% and

12.9% of this variance, respectively. (*b*) Unsupervised genetic structure plot for European and

Iberian taurine, Criollo cattle, African taurine, and American and Asian indicine populations. Results

for ancestry clusters ranging from *K* = 3–6 are shown.

Further examination of the PCA and structure plots in figure 2*a* and 2*b* showed that the Criollo

cattle samples cluster with the Iberian taurine cattle samples, all of which show relatively uniform

levels of African taurine admixture at $K = 3$ and $K = 4$. In addition, the Senepol and Hartón del Valle

Criollo breeds display a consistent and relatively uniform ancestry component from the *B. indicus*

lineage, while the Blanco Orejinegro, Costeño con Cuernos, Limonero, and Sanmartinero breeds

exhibit more heterogeneous patterns of indicine admixture. Conversely, f_3 statistics [41] indicate that

there is no evidence of admixture from either African taurine or indicine cattle in the Blanco

Orejinegro and Limonero breeds when using any of the European taurine reference populations and

African taurine or indicine breeds as the second reference (electronic supplementary material, figure

S1*a* and *c*). However, there is some evidence of indicine admixture in the Hartón del Valle as the f_3

statistics for all population trios that used any European taurine reference population with an indicine

breed as the second reference population produced negative f_3 values for this breed (electronic

supplementary material, figure S1*b*). This is also evident from the PCA (figure 2*a*), and the structure

plot (figure 2*b*).

Selection signals across Criollo cattle breed genomes

The CSS test identified selection signals for the three Criollo cattle breeds that had WGS data

and sufficient sample sizes (Blanco Orejinegro, Hartón del Valle, and Limonero). For each breed, the

test highlighted several selection signals spanning multiple chromosomes. For the Blanco Orejinegro

breed, the CSS test detected 31 cluster regions spanning 17 chromosomes, with 442 genes located

within these regions and extended ±500 kb (figure 3*a*). Similarly, for the Hartón del Valle breed, the

CSS test revealed 31 distinct clusters distributed over 17 chromosomes, encompassing 410 genes

(figure 3*b*). Lastly, for the Limonero breed, the CSS test detected 38 cluster regions, also extending

across 17 chromosomes, and that contained 451 genes in total (figure 3*c*). When these results were

consolidated, 76 genes within selection signature regions are shared across the three breeds, and

BTA18 was the chromosome with the largest number of these genes (28 genes; 36.8% of the total).

The CSS cluster regions and the genes within these regions are detailed by breed in electronic

supplementary material, tables S3 and S4).

Gene ontology (GO) term enrichment analysis

The functional significance of the genes within the CSS cluster regions for each breed was

evaluated using GO term overrepresentation analysis with the g:Profiler software tool. After adjusting

for multiple tests and using a P-value adjusted for the false discovery rate (FDR-$P_{adj.}$ < 0.05),

overrepresented GO terms were catalogued for each breed. The results of the GO term

overrepresentation analysis are shown in the electronic supplementary material, tables S5–S7 and

figure 4. For both the Blanco Orejinegro (figure 4*a*) and Limonero (figure 4*b*) breeds, the GO terms

related primarily to immunobiology, encompassing functions such as immunoglobulin receptor

binding, phagocytosis, complement activation, and antigen binding. Conversely, for the Hartón del

Valle breed (figure 4*c*), the predominant GO terms were associated with skin epidermis and

keratinization processes.

Figure 3. Chromosome-wide Manhattan plots of the smoothed composite selection signals (CSS) scores for Blanco Orejinegrio (*a*), Hartón del Valle (*b*) and Limonero (*c*). The dashed red lines indicate the genome-wide 0.1% thresholds of the empirical CSS scores. Green data points represent SNPs classified as significant and flanked by at least five additional SNPs among the top 1% of CSS scores. Notable candidate genes are indicated and discussed further in the text. Individual cattle art images modified from Felius [27] with permission of the author.

Figure 4. Gene ontology (GO) overrepresentation analysis results for the Blanco Orejinegrio (*a*), Hartón del Valle (*b*) and Limonero (*c*). The y-axis represents the negative $\log_{10}$ FDR-$P_{adj.}$ from the overrepresentation analysis. Individual cattle art images modified from Felius [27] with permission of the author.

Identification of quantitative trait loci (QTLs) within CSS cluster regions

QTLs within the CSS cluster regions were systematically identified using BovineMine v1.6.

The number of QTLs identified varied across the breeds: 519 in Blanco Orejinegro, 332 in Hartón

del Valle, and 370 in Limonero. Interestingly, only 11 QTLs were common to all three breeds and a

QTL associated with tick resistance [49] was particularly noteworthy. In the case of Blanco

Orejinegro, most of the identified QTLs (291 out of 519) in the CSS cluster regions were associated

with calving ease. Additionally, 46 QTLs were linked to milk protein percentage, 40 to milk yield,

and 24 to coat texture. For the Limonero breed, there were 292 QTLs located in the CSS cluster

regions related to calving ease; however, no other traits had more than five SNP markers among the

QTLs. In the case of the Hartón del Valle breed, the most notable traits represented among the CSS

cluster region QTLs were milk related. Specifically, 86 QTLs were associated with milk protein

percentage, 45 with milk yield, and 35 with milk fat percentage. Additionally, 27 QTLs in Hartón del

Valle were associated with coat texture. The results of the QTL search using BovineMine v1.6 are

detailed in electronic supplementary material, table S8.

DISCUSSION

Population structure and admixture

The differentiation of cattle samples across PC1 of the PCA plot (figure 2*a*) reflects the evolutionary divergence between the *B. taurus* and *B. indicus* lineages, which occurred 150–500 kya [50, 51] The second division evident for PC2 in the PCA plot (figure 2*a*) differentiates the European and African taurine groups, a separation that has been well documented in several previous studies using genome-wide SNPs and other autosomal genetic markers [52-55]. These PCs are mirrored by the clusters observed at $K = 3$ in the structure analysis (figure 2*b*), which differentiate the European taurine, African taurine, and indicine groups and reveal patterns of shared ancestry among the major groups. Extending these observations further, complex patterns of admixture due to human-mediated and unmanaged crossbreeding between taurine and indicine cattle have also been described in cattle populations across several geographical regions, particular in sub-Saharan Africa [53-55].

Previous studies have noted African taurine autosomal admixture in Criollo cattle breeds [12, 17]. Results from the present study, which are captured in the PCA plot (figure 2*a*) and structure analysis (figure 2*b*) suggest this is also the case for the Criollo breed population samples analysed here (Blanco Orejinegro, Costeño con Cuernos, Hartón del Valle, Limonero, Sanmartinero, and Senepol), albeit at lower levels compared to that observed for Guadeloupe cattle [17]. Conversely, the positive f_3 statistics observed for the three Criollo breeds examined using this method (Blanco Orejinegro, Hartón del Valle, and Limonero) do not support this (electronic supplementary material, figure S1*a–c*). It should be noted, however, that a positive f_3 value does not necessarily imply a lack of admixture, since excessive genetic drift in the target population can obscure an admixture signal [41]. In this regard, demographic modelling of historical effective population sizes (N_e values) and comparison with the Holstein, a commercial dairy population with a markedly low N_e [56], indicates that founder effects and subsequent genetic drift may have impacted the Blanco Orejinegro, Hartón del Valle, and Limonero breeds (electronic supplementary material, figure S2).

For *B. indicus* ancestry, all the Latin American Criollo cattle populations examined here exhibit

varying degrees of indicine admixture (figure 2*b*). Interestingly, however, the Blanco Orejinegro and

Hartón del Valle both originate in Colombia but do not cluster together, which is presumably a

consequence of the relatively modest indicine ancestry in the Blanco Orejinegro. The Limonero breed

samples in this study largely cluster together except for four animals that exhibit a significant

proportion of indicine ancestry. These results, taken together with the f_3 statistics for the Blanco

Orejinegro, Hartón del Valle, and Limonero (electronic supplementary material, figure S1*a–c*), and

the histories of these breeds (the Blanco Orejinegro breed has been in a conservation programme

since 1940 [57]), suggest that indicine gene flow has only had a long-term impact on breeds such as

the Hartón del Valle which exhibits relatively uniform levels of indicine admixture compared to the

Limonero, where it is more heterogeneous (figure 2*b*). This has implications for understanding the

selection signatures that are described in the following sections: in some cases, the selection signals

detected may be caused by adaptive introgression of *B. indicus* haplotypes.

Despite the inclusion of a diverse panel of European taurine, African taurine, Asian indicine,

and admixed cattle in the design and validation of the BovineHD SNP array [58], we acknowledge

that ascertainment bias could have influenced results from some of the analyses of population

differentiation and genetic ancestry. However, we think that this is unlikely because ascertainment

bias typically affects analyses such as selection signal detection that use individual SNP locus

frequency-based statistics (e.g., F_{ST}) substantially more than genome-wide multi-locus dimension

reduction tools like ADMIXTURE and PCA [59-62]. Importantly, in this regard, we used only WGS

data for the CSS analyses (figure 3 and electronic supplementary material, table S3).

Genomic signatures of human-driven selection in Criollo cattle

Blanco Orejinegro are distinctive among Criollo cattle for their white coats, often speckled with

black spots, and their skin is highly pigmented, as exemplified by their black ears. This coat-skin

pattern also occurs in British White and White Park cattle [63], and for White Park, it has been

demonstrated that this is caused by a duplication in the *KIT* gene and an aberrantly inserted *KIT* gene

on BTA29 [64]. Additional genes that harbour cattle coat colour polymorphisms include *MC1R*,

TWIST2, *MITF*, *PAX3*, *SLC45A2*, *COPA*, *TYR*, *TYRP1*, *KITLG*, and *ASIP*. For the Blanco Orejinegro

breed, the genes *PAX3*, *SLC45A2*, and *KITLG* are in genome regions with evidence for directional

selection based on the CSS results. Mutations in *PAX3* have been shown to result in a spectrum of

phenotypic outcomes in horses from white spotting to a uniform white coat trait [65]. Similarly,

mutations in the bovine *SLC45A2* gene have been shown to cause oculocutaneous albinism [66],

while the *KITLG* gene has been associated with the roan coat type in cattle [67].

Interestingly, for the Hartón del Valle and Limonero breeds, *MC1R* was part of a BTA18 CSS

cluster region. The protein receptor encoded by *MC1R* regulates tyrosinase levels in melanocytes, an

enzyme critical to melanin synthesis. High concentrations of tyrosinase facilitate the production of

eumelanin, contributing to darker brown or black shades, whereas lower concentrations drive the

synthesis of phaeomelanin, yielding lighter red or yellow tones [68]. It is therefore noteworthy that

Limonero and Hartón del Valle cattle generally possess a "bayo" or bay coat, which is a yellow to

light brown/red colour [69, 70]. Though not in a CSS cluster region in the Blanco Orejinegro, a

notable peak close to *MC1R* is present on BTA18. This gene has been observed to be under positive

selection in *B. indicus* cattle resulting in a light grey to white coat in the Brahman and Nelore breeds

[71]. The adaptive significance of functional polymorphisms at *MC1R* is that lighter coat colours

reflect a significant portion of incident solar radiation, which makes these cattle more suited to

tropical environments [72].

337   Selection for heat tolerance in Criollo cattle

Criollo cattle breeds such as the Blanco Orejinegro, Hartón del Valle, and Limonero have

evolved over five centuries to thrive in challenging tropical agroecologies. In the context of this

environmental adaptation, a key finding for all three breeds is the presence of a CSS peak on BTA20

that encompasses the *PRLR* gene, most notably in the Blanco Orejinegro and Hartón del Valle breeds,

where it was statistically significant. Mutations in this gene have been shown to cause the "slick"

phenotype in in Criollo breeds such as the Blanco Orejinegro, Hartón del Valle, Limonero,

Romosinuano, Costeño con Cuernos, Mexican Criollo Lechero Tropical, and also Criollo-influenced

composite breeds such as the Carora and Senepol [73]. Of particular interest for understanding how

convergent evolution can act in livestock populations is the existence of several different *PRLR*

mutations that can produce the "slick" phenotype in Criollo cattle [73]. Mutations in the bovine *PRLR*

gene can have major effects on the length and the structure of hair coats providing improved

thermotolerance and concomitant increase on fertility and milk yields in cattle populations that inhabit

dry and tropical conditions [18, 69, 74]. In addition, it has been shown that these mutations can act

pleiotropically and cause other physiological changes [75].

Variation at other genes may also confer enhanced thermotolerance in Criollo cattle; for

example, *MVD*, previously implicated as being under selection in North African cattle [76], was found

in a BTA18 CSS cluster region for all three breeds. In addition, *CCM2* which regulates heart and

blood vessel formation and integrity, is located in a Limonero BTA4 CSS cluster region and was also

highlighted by Ben-Jemaa and colleagues in North African cattle [76]. This may represent an

adaptation that increases heat transfer from the interior to the skin, which is also supported by the

enhanced vascularisation of the dermis that has been documented in Limonero cattle [69]. Another

gene, *SESN2*, located in a BTA2 CSS cluster detected in the Hartón del Valle and the Blanco

Orejinegro breeds may also be associated with adaptation to tropical environments. This gene was

observed to be downregulated in a functional genomics study of heat-stress responses in dermal

fibroblast cells from indicine and indicine × taurine crossbred cattle [77].

The BTA11 Limonero CSS cluster region contains *FBXO4* that encodes a protein, which

regulates body temperature through interactions with the heath shock protein HSPB6 (previously

HSP20) [72, 78]. We also observed that *DNAJA2* was in a Blanco Orejinegro BTA18 CSS cluster

region. This gene encodes a member of the DNAJ/HSP40 family of proteins and shows increased

expression in peripheral blood from Holstein calves exposed to heat stress [79]. Similarly, *HIGD1A*

(BTA22 CSS cluster) and *CBFA2T3* (a BTA18 CSS cluster distinct from that containing *DNAJA2*),

encode proteins associated with cellular responses to hypoxia [80, 81], which could be indicative of

adaptation to oxidative stress caused by high temperatures [82].

371 Genomic signatures of selection for fertility and reproductive traits in Criollo cattle

Compared to taurine breeds from temperate zones, Criollo cattle are renowned for high fertility

and excellent reproductive performance [21, 83] and using the CSS method we detected several genes

associated with these traits. For example, in the Blanco Orejinegro breed a BTA7 CSS cluster region

contains the *CATSPER3* gene, which encodes a sperm-specific ion channel protein directly linked to

sperm motility and male fertility in many species, including cattle [84]. Another CSS cluster region

in the Blanco Orejinegro on BTA20 contained *SPEF2*, which is involved in the formation and

functionality of sperm flagella [85]. Additionally, the *CREM* gene, implicated in spermatogenesis,

was located within a Blanco Orejinegro CSS cluster region on BTA13 [86]. Other genes related to

reproductive physiology were located within CSS cluster regions for the Hartón del Valle breed

including *HKDC1* on BTA28 that exhibits high expression in the testes [87], and *MISP3* on BTA7,

which is implicated in spermatogenesis [88]. For the Limonero breed, CSS cluster regions included

one on BTA10 that contains *FSIP1* a gene crucial for normal spermiogenesis and flagella

development [89]. In addition, the *PTGES* gene within a Limonero CSS cluster region on BTA11

encodes an enzyme involved in the synthesis of prostaglandin E2 (*PGE2*), which is a signalling

molecule with a crucial role in reproductive processes [90]. In parallel to this, the QTL analysis

(electronic supplementary material, table S8) for the Blanco Orejinegro breed identified QTLs

predominantly associated with reproduction, notably calving ease, and lactation. These findings

corroborate previous results obtained using SNP array data for Blanco Orejinegro cattle [20, 91]. The

results of the QTL analysis in the CSS cluster regions for the Limonero breed were similar to those

for the Blanco Orejinegro breed.

392 Genomic adaptations to infectious diseases in Criollo cattle

Comparable to reproductive and fertility traits, desirable disease resistance and tolerance traits

have been highlighted as important features of Criollo cattle populations. However, empirical

evidence for the genetic basis of these traits is often lacking; instead, much of the evidence base is

anecdotal and derived from practical knowledge and experiences of veterinarians, farmers and

breeders [70]. In the present study, however, we have begun to address this knowledge gap through

identification of several immune genes in CSS cluster regions for the Blanco Orejinegro, Hartón del

Valle, and Limonero breeds. For example, the *CBFA2T3* gene located in the BTA18 CSS cluster

region detected for all three breeds and discussed in the context of hypoxia, has a role in the bovine

immune response to mammary gland inflammation [92]. Similarly, *CXCL14*, located in a BTA7 CSS

cluster region for the Blanco Orejinegro and Limonero breeds has also been implicated in the immune

response to bovine mastitis caused by *Staphylococcus aureus* [93]. In addition, for the Limonero, a

BTA11 CSS cluster region contains the *LRRC8A* gene, which is essential for development and

function of T lymphocytes [94].

In tropical and subtropical regions, there is a significant prevalence of vector-borne

haemoparasitic infections [95] that induce anaemia, fever, wasting, and reproductive dysfunction

[96]. These diseases are primarily caused by eukaryotic parasites within the *Babesia* [97] and

Trypanosoma [98] genera, and bacterial pathogens from the *Anaplasma* genus [99]. It is noteworthy,

therefore, that the *EPB41* gene located within a Blanco Orejinegro and Hartón del Valle BTA2 CSS

cluster region is associated with resistance to anaemia and trypanotolerance in African cattle [100].

Also, for the Blanco Orejinegro and Limonero, *RPS27* and *RPS20* are located in CSS cluster regions

on BTA3 and BTA14, respectively and mutations in these genes can cause Diamond-Blackfan

anaemia [101, 102]. Furthermore, in the Blanco Orejinegro breed, the *ZFPM1* gene within a BTA18

CSS cluster region, has been shown to play a role in cardiac and hematopoietic development. In a

murine *Trypanosoma congolense* infection model, mice exhibiting decreased *ZFPM1* expression

recovered more effectively from anaemia [103].

Gene GO term analysis in the Criollo CSS cluster regions revealed a marked overrepresentation

of immunobiological processes for the Blanco Orejinegro and Limonero breeds, many of which

overlapped between the two breeds (figure 4*a* and 4*c* and electronic supplementary material, tables

S5 and S7). The statistically significant overrepresented biological process GO terms in the Blanco

Orejinegro and Limonero included several terms related to intracellular pathogens, including

phagocytosis (GO:0006909), *phagocytosis, engulfment* (GO:0006911), and *phagocytosis,*

recognition (GO:0006910). It is therefore noteworthy that Blanco Orejinegro cattle have a

documented resistance to brucellosis [104], a disease characterized by abortion and retention of the

placenta, which is principally caused by *Brucella abortus* an intracellular bacterium that subverts

phagocytic pathways and processes to enter and manipulate host cells [105].

CONCLUSIONS

This study provides valuable insights into the genomic basis of microevolutionary change in

Criollo cattle as they have adapted to the tropical environments of Latin America. Evidence of strong

selective pressure was apparent, particularly for the distinct coat and skin coloration traits observed

in these breeds, which are advantageous in cattle populations exposed to significant levels of incident

solar radiation. Notably, we also discovered genomic selection signatures that may be associated with

thermotolerance, again underscoring adaptation to hot climates. In addition, some of the selection

signatures we identified align with documented fertility traits in Criollo cattle. Moreover, functional

overrepresentation analysis revealed many genes related to immune function, which could reflect

resilience to multiple infectious disease challenges. Taken together, our results, show the remarkable

adaptability of Criollo cattle, which has been driven by natural and human-mediated selection and

that underscores the genetic richness and value of these breeds for future breeding programmes.

Acknowledgements: We thank Thomas Hall for advice on bioinformatics and computational
genomics methodologies.

Funding: J.A.W. was supported by Science Foundation Ireland (SFI) and Acceligen/Recombinetics
Inc. through the SFI Centre for Research Training in Genomics Data Science under grant no.
18/CRT/6214. G.P.M. and D.E.M. were supported by an SFI Investigator Award (SFI/15/IA/3154).
T.S.S. and D.E.M were supported by a United States Department of Agriculture (USDA) and
Department of Agriculture, Food, and the Marine (DAFM) US-Ireland Research and Development
Partnership grant (TARGET-TB, 17/RD/USROI/52).

REFERENCES

[1] Rouse, J.E. 1977 *The Criollo: Spanish Cattle in the Americas*, University of Oklahoma Press.

[2] Armenteros Martínez, I. 2018 The Canary Islands as an area of interconnectivity between the Mediterranean and the Atlantic (Fourteenth-Sixteenth Centuries). In *Entre mers—Outre-mer: Spaces, Modes and Agents of Indo-Mediterranean Connectivity* (eds. N. Jaspert & S. Kolditz), pp. 201-217. Heidelberg, Germany, Heidelberg University Publishing.

[3] Rodero, A., Delgado, J.V. & Rodero. 1992 Primitive Andalusian livestock and their implications in the discovery of America. *Arch. Zootec.* **41**, 383-400.

[4] Dunmire, W.W. 2004 *Gardens of New Spain: How Mediterranean Plants and Foods Changed America*. 1st ed. Austin, University of Texas Press; xviii, 375 p. p.

[5] Crosby, A.W. 2003 *The Columbian Exchange: Biological and Cultural Consequences of 1492*. 30th anniversary ed. Westport, Conn., Praeger; 283 p. p.

[6] Willham, R.L. 1982 Genetic improvement of beef cattle in the United States: cattle, people and their interaction. *J. Anim. Sci.* **54**, 659-666. (doi:10.2527/jas1982.543659x).

[7] Tobar, K.M.C., Álvarez, D.C. & Franco, L.A. 2020 Genome-wide association studies in sheep from Latin America. Review. *Rev. Mex. Cienc. Pecu.* **11**. (doi:10.22319/rmcp.v11i3.5372sander).

[8] Revidatti, M.A., Gama, L.T., Martin Burriel, I., Cortes Gardyn, O., Cappello Villada, J.S., Carolino, M.I., Canon, F.J., Ginja, C., Sponenberg, P., Vicente, A.P., et al. 2021 On the origins of American Criollo pigs: A common genetic background with a lasting Iberian signature. *PLoS ONE* **16**, e0251879. (doi:10.1371/journal.pone.0251879).

[9] Taylor, W.T.T., Librado, P., American Horse, C.J., Shield Chief Gover, C., Arterberry, J., Afraid of Bear-Cook, A.L., Left Heron, H., Yellow Hair, R.M., Gonzalez, M., Means, B., et al. 2023 Early dispersal of domestic horses into the Great Plains and northern Rockies. *Science* **379**, 1316-1323. (doi:10.1126/science.adc9691).

[10] Sanders, J.O. 1980 History and development of zebu cattle in the United States. *J. Anim. Sci.* **50**, 1188-1200. (doi:10.2527/jas1980.5061188x).

[11] da Fonseca, R.R., Ureña, I., Afonso, S., Pires, A.E., Jørsboe, E., Chikhi, L. & Ginja, C. 2019 Consequences of breed formation on patterns of genomic diversity and differentiation: the case of highly diverse peripheral Iberian cattle. *BMC Genomics* **20**, 334. (doi:10.1186/s12864-019-5685-2).

[12] Pitt, D., Bruford, M.W., Barbato, M., Orozco-terWengel, P., Martínez, R. & Sevane, N. 2019 Demography and rapid local adaptation shape Creole cattle genome diversity in the tropics. *Evol. Appl.* **12**, 105-122. (doi:10.1111/eva.12641).

[13] Upadhyay, M., Bortoluzzi, C., Barbato, M., Ajmone-Marsan, P., Colli, L., Ginja, C., Sonstegard, T.S., Bosse, M., Lenstra, J.A., Groenen, M.A.M., et al. 2019 Deciphering the patterns of genetic admixture and diversity in southern European cattle using genome-wide SNPs. *Evol. Appl.* **12**, 951-963. (doi:10.1111/eva.12770).

[14] Magee, D.A., Meghen, C., Harrison, S., Troy, C.S., Cymbron, T., Gaillard, C., Morrow, A., Maillard, J.C. & Bradley, D.G. 2002 A partial African ancestry for the Creole cattle populations of the Caribbean. *J. Hered.* **93**, 429-432. (doi:10.1093/jhered/93.6.429).

[15] Ginja, C., Gama, L.T., Cortés, O., Burriel, I.M., Vega-Pla, J.L., Penedo, C., Sponenberg, P., Cañón, J., Sanz, A., do Egito, A.A., et al. 2019 The genetic ancestry of American Creole cattle inferred from uniparental and autosomal genetic markers. *Sci. Rep.* **9**, 11486. (doi:10.1038/s41598-019-47636-0).

[16] Delsol, N., Stucky, B.J., Oswald, J.A., Cobb, C.R., Emery, K.F. & Guralnick, R. 2023 Ancient
DNA confirms diverse origins of early post-Columbian cattle in the Americas. *Sci. Rep.* **13**, 12444.
(doi:10.1038/s41598-023-39518-3).

[17] Ben-Jemaa, S., Adam, G., Boussaha, M., Bardou, P., Klopp, C., Mandonnet, N. & Naves, M.
2023 Whole genome sequencing reveals signals of adaptive admixture in Creole cattle. *Sci. Rep.* **13**,
12155. (doi:10.1038/s41598-023-38774-7).

[18] Landaeta-Hernández, A.J., Zambrano-Nava, S., Verde, O., Pinto-Santini, L., Montero-Urdaneta,
M., Hernández-Fonseca, J.P., Fuenmayor-Morales, C., Sonstegard, T.S., Huson, H.J. & Olson, T.A.
2021 Heat stress response in slick vs normal-haired Criollo Limonero heifers in a tropical
environment. *Trop. Anim. Health Prod.* **53**, 445. (doi:10.1007/s11250-021-02856-3).

[19] Martínez, A.M., Gama, L.T., Cañón, J., Ginja, C., Delgado, J.V., Dunner, S., Landi, V., Martín-
Burriel, I., Penedo, M.C., Rodellar, C., et al. 2012 Genetic footprints of Iberian cattle in America 500
years after the arrival of Columbus. *PLoS ONE* **7**, e49066. (doi:10.1371/journal.pone.0049066).

[20] Caivio-Nasner, S., López-Herrera, A., González-Herrera, L.G. & Rincón, J.C. 2021 Frequency
of genotypic markers for genetic disorders, colour, polledness, and major genes in Blanco Orejinegro
cattle. *Trop. Anim. Health Prod.* **53**, 546. (doi:10.1007/s11250-021-02990-y).

[21] Anderson, D.M., Estell, R.E., Gonzalez, A.L., Cibils, A.F. & Torell, L.A. 2015 Criollo cattle:
heritage genetics for arid landscapes. *Rangelands* **37**, 62-67. (doi:10.1016/j.rala.2015.01.006).

[22] Food and Agriculture Organization. 2015 *The Second Report on the State of the World's Animal
Genetic Resources for Food and Agriculture*. Rome, Italy, edited by B.D. Scherf & D. Pilling. FAO
Commission on Genetic Resources for Food and Agriculture Assessments.

[23] Rosen, B.D., Bickhart, D.M., Schnabel, R.D., Koren, S., Elsik, C.G., Tseng, E., Rowan, T.N.,
Low, W.Y., Zimin, A., Couldrey, C., et al. 2020 De novo assembly of the cattle reference genome
with single-molecule sequencing. *Gigascience* **9**. (doi:10.1093/gigascience/giaa021).

[24] Vasimuddin, M., Misra, S., Li, H. & Aluru, S. 2019 Efficient architecture-aware acceleration of
BWA-MEM for multicore ssemystems. *2019 IEEE International Parallel and Distributed
Processing Symposium (IPDPS)*, 314-324. (doi:10.1109/IPDPS.2019.00041).

[25] Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., et al. 2021 Twelve years of SAMtools and BCFtools.
Gigascience **10**. (doi:10.1093/gigascience/giab008).

[26] Van der Auwera, G.A. & O'Connor, B.D. 2020 *Genomics in the Cloud: using Docker, GATK,
and WDL in Terra*. 1st ed. Sebastopol, CA, O'Reilly Media.

[27] Felius, M. 1995 *Cattle Breeds: An Encyclopedia*. Doetinchem, Netherlands, Misset.

[28] Cingolani, P., Platts, A., Wang le, L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X. &
Ruden, D.M. 2012 A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-
3. *Fly (Austin)* **6**, 80-92. (doi:10.4161/fly.19695).

[29] Bahbahani, H., Tijjani, A., Mukasa, C., Wragg, D., Almathen, F., Nash, O., Akpa, G.N., Mbole-
Kariuki, M., Malla, S., Woolhouse, M., et al. 2017 Signatures of selection for environmental
adaptation and zebu x taurine hybrid fitness in East African shorthorn zebu. *Front. Genet.* **8**, 68.
(doi:10.3389/fgene.2017.00068).

[30] Verdugo, M.P., Mullin, V.E., Scheu, A., Mattiangeli, V., Daly, K.G., Maisano Delser, P., Hare,
A.J., Burger, J., Collins, M.J., Kehati, R., et al. 2019 Ancient cattle genomics, origins, and rapid
turnover in the Fertile Crescent. *Science* **365**, 173-176. (doi:10.1126/science.aav1002).

[31] Upadhyay, M.R., Chen, W., Lenstra, J.A., Goderie, C.R.J., MacHugh, D.E., Park, S.D.E., Magee,
D.A., Matassino, D., Ciani, F., Megens, H.J., et al. 2017 Genetic origin, admixture and population

history of aurochs (*Bos primigenius*) and primitive European cattle. *Heredity* **118**, 169-176.
(doi:10.1038/hdy.2016.79).

[32] Ward, J.A., McHugo, G.P., Dover, M.J., Hall, T.J., Sonstegard, T.S., Bradley, D.G., Frantz,
L.A.F., Salter-Townshend, M. & MacHugh, D.E. 2022 Genome-wide local ancestry and evidence for
mitonuclear   coadaptation   in   African   hybrid   cattle   populations.   *iScience*   **25**.
(doi:10.1016/j.isci.2022.104672).

[33] Sempéré, G., Moazami-Goudarzi, K., Eggen, A., Laloë, D., Gautier, M. & Flori, L. 2015
WIDDE: a Web-Interfaced next generation database for genetic diversity exploration, with a first
application in cattle. *BMC Genomics* **16**, 940. (doi:verit).

[34] Zimin, A.V., Delcher, A.L., Florea, L., Kelley, D.R., Schatz, M.C., Puiu, D., Hanrahan, F.,
Pertea, G., Van Tassell, C.P., Sonstegard, T.S., et al. 2009 A whole-genome assembly of the domestic
cow, *Bos taurus*. *Genome Biol.* **10**, R42. (doi:10.1186/gb-2009-10-4-r42).

[35] Riggio, V., Tijjani, A., Callaby, R., Talenti, A., Wragg, D., Obishakin, E.T., Ezeasor, C.,
Jongejan, F., Ogo, N.I., Aboagye-Antwi, F., et al. 2022 Assessment of genotyping array performance
for genome-wide association studies and imputation in African cattle. *Genet. Sel. Evol.* **54**, 58.
(doi:10.1186/s12711-022-00751-5).

[36] Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M. & Lee, J.J. 2015 Second-
generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7.
(doi:10.1186/s13742-015-0047-8).

[37] R Core Team. 2022 R: A language and environment for statistical computing. (Vienna, Austria,
R Foundation for Statistical Computing. http://www.R-project.org.

[38] Wickham, H. 2016 *ggplot2: Elegant Graphics for Data Analysis*, Springer Publishing Company,
New York, USA; wri p.

[39] Alexander, D.H., Novembre, J. & Lange, K. 2009 Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res.* **19**, 1655-1664. (doi:10.1101/gr.094052.109).

[40] Barbato, M., Orozco-terWengel, P., Tapio, M. & Bruford, M.W. 2015 SNeP: a tool to estimate
trends in recent effective population size trajectories using genome-wide SNP data. *Front. Genet.* **6**,
109. (doi:10.3389/fgene.2015.00109).

[41] Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T.,
Webster, T. & Reich, D. 2012 Ancient admixture in human history. *Genetics* **192**, 1065-1093.
(doi:10.1534/genetics.112.145037).

[42] Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
Lunter, G., Marth, G.T., Sherry, S.T., et al. 2011 The variant call format and VCFtools.
Bioinformatics **27**, 2156-2158. (doi:10.1093/bioinformatics/btr330).

[43] Szpiech, Z.A. & Hernandez, R.D. 2014 selscan: An efficient multithreaded program to perform
EHH-based   scans   for   positive   selection.   *Mol.   Biol.   Evol.*   **31**,   2824-2827.
(doi:10.1093/molbev/msu211).

[44] Randhawa, I.A.S., Khatkar, M.S., Thomson, P.C. & Raadsma, H.W. 2014 Composite selection
signals can localize the trait specific genomic regions in multi-breed populations of cattle and sheep.
BMC Genet. **15**, 34. (doi:10.1186/1471-2156-15-34).

[45] Verity, R., Collins, C., Card, D.C., Schaal, S.M., Wang, L. & Lotterhos, K.E. 2017 minotaur: A
platform for the analysis and visualization of multivariate results from genome scans with R Shiny.
Mol. Ecol. Resour. **17**, 33-43. (doi:10.1111/1755-0998.12579).

[46] Han, H., Randhawa, I.A.S., MacHugh, D.E., McGivney, B.A., Katz, L.M., Dugarjaviin, M. &
Hill, E.W. 2023 Selection signatures for local and regional adaptation in Chinese Mongolian horse

breeds reveal candidate genes for hoof health. *BMC Genomics* **24**, 35. (doi:10.1186/s12864-023-09116-8).

[47] Elsik, C.G., Unni, D.R., Diesh, C.M., Tayal, A., Emery, M.L., Nguyen, H.N. & Hagen, D.E. 2016 Bovine Genome Database: new tools for gleaning function from the *Bos taurus* genome. *Nucleic Acids Res.* **44**, D834-D839. (doi:10.1093/nar/gkv1077).

[48] Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H. & Vilo, J. 2019 g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191-W198. (doi:10.1093/nar/gkz369).

[49] Sollero, B.P., Junqueira, V.S., Gomes, C.C.G., Caetano, A.R. & Cardoso, F.F. 2017 Tag SNP selection for prediction of tick resistance in Brazilian Braford and Hereford cattle breeds using Bayesian methods. *Genet. Sel. Evol.* **49**, 1-15. (doi:10.1186/s12711-017-0325-2).

[50] Chen, N., Cai, Y., Chen, Q., Li, R., Wang, K., Huang, Y., Hu, S., Huang, S., Zhang, H., Zheng, Z., et al. 2018 Whole-genome resequencing reveals world-wide ancestry and adaptive introgression events of domesticated cattle in East Asia. *Nat. Commun.* **9**, 2337. (doi:10.1038/s41467-018-04737-0).

[51] Wu, D.D., Ding, X.D., Wang, S., Wojcik, J.M., Zhang, Y., Tokarska, M., Li, Y., Wang, M.S., Faruque, O., Nielsen, R., et al. 2018 Pervasive introgression facilitated domestication and adaptation in the *Bos* species complex. *Nat. Ecol. Evol.* **2**, 1139-1145. (doi:10.1038/s41559-018-0562-y).

[52] MacHugh, D.E., Shriver, M.D., Loftus, R.T., Cunningham, P. & Bradley, D.G. 1997 Microsatellite DNA variation and the evolution, domestication and phylogeography of taurine and zebu cattle (*Bos taurus* and *Bos indicus*). *Genetics* **146**, 1071-1086. (doi:10.1093/genetics/146.3.1071).

[53] Kim, K., Kwon, T., Dessie, T., Yoo, D., Mwai, O.A., Jang, J., Sung, S., Lee, S., Salim, B., Jung, J., et al. 2020 The mosaic genome of indigenous African cattle as a unique genetic resource for African pastoralism. *Nat. Genet.* **52**, 1099-1110. (doi:10.1038/s41588-020-0694-2).

[54] Hanotte, O., Bradley, D.G., Ochieng, J.W., Verjee, Y., Hill, E.W. & Rege, J.E.O. 2002 African pastoralism: Genetic imprints of origins and migrations. *Science* **296**, 336-339. (doi:10.1126/science.1069878).

[55] Decker, J.E., McKay, S.D., Rolf, M.M., Kim, J., Molina Alcala, A., Sonstegard, T.S., Hanotte, O., Gotherstrom, A., Seabury, C.M., Praharani, L., et al. 2014 Worldwide patterns of ancestry, divergence, and admixture in domesticated cattle. *PLoS Genet.* **10**, e1004254. (doi:10.1371/journal.pgen.1004254).

[56] Makanjuola, B.O., Miglior, F., Abdalla, E.A., Maltecca, C., Schenkel, F.S. & Baes, C.F. 2020 Effect of genomic selection on rate of inbreeding and coancestry and effective population size of Holstein and Jersey cattle populations. *J. Dairy Sci.* **103**, 5183-5199. (doi:10.3168/jds.2019-18013).

[57] Toro, R.J.E. 2020 Evaluación de herramientas para desarrollar un programa de mejoramiento genético en la raza Blanco Orejinegro (BON) [PhD thesis]. Medellín, Colombia, University of Antioquia.

[58] Illumina. 2015 Data sheet: BovineHD Genotyping BeadChip. (San Diego, CA. www.illumina.com/Documents/products/datasheets/datasheet_bovineHD.pdf.

[59] Albrechtsen, A., Nielsen, F.C. & Nielsen, R. 2010 Ascertainment biases in SNP chips affect measures of population divergence. *Mol. Biol. Evol.* **27**, 2534-2547. (doi:10.1093/molbev/msq148).

[60] Malomane, D.K., Reimer, C., Weigend, S., Weigend, A., Sharifi, A.R. & Simianer, H. 2018 Efficiency of different strategies to mitigate ascertainment bias when using SNP panels in diversity studies. *BMC Genomics* **19**, 22. (doi:10.1186/s12864-017-4416-9).

[61] McTavish, E.J. & Hillis, D.M. 2015 How do SNP ascertainment schemes and population
demographics affect inferences about population history? *BMC Genomics* **16**, 266.
(doi:10.1186/s12864-015-1469-5).

[62] McVean, G. 2009 A genealogical interpretation of principal components analysis. *PLoS Genet.*
5, e1000686. (doi:10.1371/journal.pgen.1000686).

[63] Pearson, L. 1968 A note on the history of black-eared white cattle. *Agric. Hist. Rev.* **16**, 159-160.

[64] Brenig, B., Beck, J., Floren, C., Bornemann-Kolatzki, K., Wiedemann, I., Hennecke, S., Swalve,
H. & Schütz, E. 2013 Molecular genetics of coat colour variations in White Galloway and White Park
cattle. *Anim. Genet.* **44**, 450-453. (doi:10.1111/age.12029).

[65] Druml, T., Grilz-Seger, G., Neuditschko, M., Horna, M., Ricard, A., Pausch, H. & Brem, G.
2018 Novel insights into Sabino1 and splashed white coat color patterns in horses. *Anim. Genet.* **49**,
249-253. (doi:10.1111/age.12657).

[66] Rothammer, S., Kunz, E., Seichter, D., Krebs, S., Wassertheurer, M., Fries, R., Brem, G. &
Medugorac, I. 2017 Detection of two non-synonymous SNPs in *SLC45A2* on BTA20 as candidate
causal mutations for oculocutaneous albinism in Braunvieh cattle. *Genetics Selection Evolution* **49**,
73. (doi:10.1186/s12711-017-0349-7).

[67] Voß, K., Blaj, I., Tetens, J.L., Thaller, G. & Becker, D. 2022 Roan coat color in livestock. *Anim.*
Genet. **53**, 549-556. (doi:10.1111/age.13240).

[68] Goud, T.S., Upadhyay, R.C., Onteru, S.K., Pichili, V.B.R. & Chadipiralla, K. 2020 Identification
and sequence characterization of melanocortin 1 receptor gene (*MC1R*) in *Bos indicus* versus (*Bos*
taurus X *Bos indicus*). *Anim. Biotechnol.* **31**, 283-294. (doi:10.1080/10495398.2019.1585866).

[69] Landaeta-Hernández, A., Zambrano-Nava, S., Hernández-Fonseca, J.P., Godoy, R., Calles, M.,
Iragorri, J.L., Añez, L., Polanco, M., Montero-Urdaneta, M. & Olson, T. 2011 Variability of hair coat
and skin traits as related to adaptation in Criollo Limonero cattle. *Trop. Anim. Health Prod.* **43**, 657-
663. (doi:10.1007/s11250-010-9749-1).

[70] Landaeta-Hernández, A., Zambrano, S., Marin, N. & Moreno, Y. 2011 Criollos
Latinoamericanos. In *Workshop Jornadas de Actualización en Ganadería Tropical* (pp. 52-62,
Asociación de Ganaderos Alberto Adriani (ASODEGAA).

[71] Mei, C., Wang, H., Liao, Q., Wang, L., Cheng, G., Wang, H., Zhao, C., Zhao, S., Song, J., Guang,
X., et al. 2018 Genetic architecture and selection of Chinese cattle revealed by whole genome
resequencing. *Mol. Biol. Evol.* **35**, 688-699. (doi:10.1093/molbev/msx322).

[72] Hansen, P.J. 2020 Prospects for gene introgression or gene editing as a strategy for reduction of
the impact of heat stress on production and reproduction in cattle. *Theriogenology* **154**, 190-202.
(doi:10.1016/j.theriogenology.2020.05.010).

[73] Flórez Murillo, J.M., Landaeta-Hernández, A.J., Kim, E.-S., Bostrom, J.R., Larson, S.A., Pérez
O'Brien, A.M., Montero-Urdaneta, M.A., Garcia, J.F. & Sonstegard, T.S. 2021 Three novel nonsense
mutations of prolactin receptor found in heat-tolerant *Bos taurus* breeds of the Caribbean Basin. *Anim.*
Genet. **52**, 132-134. (doi:10.1111/age.13027).

[74] Dikmen, S., Khan, F.A., Huson, H.J., Sonstegard, T.S., Moss, J.I., Dahl, G.E. & Hansen, P.J.
2014 The *SLICK* hair locus derived from Senepol cattle confers thermotolerance to intensively
managed lactating Holstein cows. *J. Dairy Sci.* **97**, 5508-5520. (doi:10.3168/jds.2014-8087).

[75] Sosa, F., Santos, J.E.P., Rae, D.O., Larson, C.C., Macchietto, M., Abrahante, J.E., Amaral, T.F.,
Denicol, A.C., Sonstegard, T.S. & Hansen, P.J. 2022 Effects of the SLICK1 mutation in *PRLR* on
regulation of core body temperature and global gene expression in liver in cattle. *Animal* **16**, 100523.
(doi:10.3168/jds.2022-22272).

[76] Ben-Jemaa, S., Mastrangelo, S., Lee, S.-H., Lee, J.H. & Boussaha, M. 2020 Genome-wide scan
for selection signatures reveals novel insights into the adaptive capacity in local North African cattle.
Sci. Rep. **10**, 19466. (doi:10.1038/s41598-020-76576-3).

[77] Singh, A.K., Upadhyay, R.C., Chandra, G., Kumar, S., Malakar, D., Singh, S.V. & Singh, M.K.
2020 Genome-wide expression analysis of the heat stress response in dermal fibroblasts of Tharparkar
(zebu) and Karan-Fries (zebu × taurine) cattle. *Cell Stress Chaperones* **25**, 327-344. (doi:druml).

[78] Peng, J., Li, Y., Wang, X., Deng, S., Holland, J., Yates, E., Chen, J., Gu, H., Essandoh, K., Mu,
X., et al. 2018 An Hsp20-FBXO4 axis regulates adipocyte function through modulating PPARγ
ubiquitination. *Cell Rep.* **23**, 3607-3620. (doi:10.1016/j.celrep.2018.05.065).

[79] Srikanth, K., Kwon, A., Lee, E. & Chung, H. 2017 Characterization of genes and pathways that
respond to heat stress in Holstein calves through transcriptome analysis. *Cell Stress Chaperones* **22**,
29-42. (doi:10.1007/s12192-016-0739-8).

[80] Hayashi, T., Asano, Y., Shintani, Y., Aoyama, H., Kioka, H., Tsukamoto, O., Hikita, M.,
Shinzawa-Itoh, K., Takafuji, K., Higo, S., et al. 2015 Higd1a is a positive regulator of cytochrome c
oxidase. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 1553-1558. (doi:10.1073/pnas.1419767112).

[81] Zhang, J., Long, K., Wang, J., Zhang, J., Jin, L., Tang, Q., Li, X., Ma, J., Li, M. & Jiang, A.
2022 Yak miR-2285o-3p attenuates hypoxia-induced apoptosis by targeting caspase-3. *Anim. Genet.*
53, 49-57. (doi:10.1111/age.13153).

[82] Zhang, X., Wang, D. & Liu, J. 2023 Hypoxia-inducible factor-1α is involved in the response to
heat stress in lactating dairy cows. *J. Therm. Biol.* **112**, 103460. (doi:10.1016/j.jtherbio.2023.103460).

[83] Armstrong, E., Rodriguez Almeida, F.A., McIntosh, M.M., Poli, M., Cibils, A.F., Martínez-
Quintana, J.A., Félix-Portillo, M. & Estell, R.E. 2022 Genetic and productive background of Criollo
cattle in Argentina, Mexico, Uruguay and the United States. *J. Arid Environ.* **200**, 104722.
(doi:10.1016/j.jaridenv.2022.104722).

[84] Johnson, G.P., English, A.-M., Cronin, S., Hoey, D.A., Meade, K.G. & Fair, S. 2017 Genomic
identification, expression profiling, and functional characterization of CatSper channels in the bovine.
Biol. Reprod. **97**, 302-312. (doi:10.1093/biolre/iox082).

[85] Guo, F., Yang, B., Ju, Z.H., Wang, X.G., Qi, C., Zhang, Y., Wang, C.F., Liu, H.D., Feng, M.Y.,
Chen, Y., et al. 2014 Alternative splicing, promoter methylation, and functional SNPs of sperm
flagella 2 gene in testis and mature spermatozoa of Holstein bulls. *J. Reprod. Fertil.* **147**, 241-252.
(doi:10.1530/REP-13-0343).

[86] Nantel, F., Monaco, L., Foulkes, N.S., Masquilier, D., LeMeur, M., Henriksén, K., Dierich, A.,
Parvinen, M. & Sassone-Corsi, P. 1996 Spermiogenesis deficiency and germ-cell apoptosis in
CREM-mutant mice. *Nature* **380**, 159-162. (doi:10.1038/380159a0).

[87] Zapater, J.L., Lednovich, K.R., Khan, M.W., Pusec, C.M. & Layden, B.T. 2022 Hexokinase
domain-containing protein-1 in metabolic diseases and beyond. *Trends Endocrinol. Metab.* **33**, 72-
84. (doi:10.1016/j.tem.2021.10.006).

[88] Ren, F., Xi, H., Qiao, P., Li, Y., Xian, M., Zhu, D. & Hu, J. 2022 Single-cell transcriptomics
reveals male germ cells and Sertoli cells developmental patterns in dairy goats. *Front. Cell Dev. Biol.*
10, 944325. (doi:10.3389/fcell.2022.944325).

[89] Gamallat, Y., Fang, X., Mai, H., Liu, X., Li, H., Zhou, P., Han, D., Zheng, S., Liao, C., Yang,
M., et al. 2021 Bi-allelic mutation in *Fsip1* impairs acrosome vesicle formation and attenuates
flagellogenesis in mice. *Redox Biol.* **43**, 101969. (doi:10.1016/j.redox.2021.101969).

[90] Berisha, B., Rodler, D., Schams, D., Sinowatz, F. & Pfaffl, M.W. 2019 Prostaglandins in
superovulation induced bovine follicles during the preovulatory period and early corpus luteum.
Front. Endocrinol. (Lausanne) **10**, 467. (doi:10.3389/fendo.2019.00467).

[91] Caivio-Nasner, S., López-Herrera, A., González-Herrera, L.G. & Rincón, J.C. 2021 Diversity analysis, runs of homozygosity and genomic inbreeding reveal recent selection in Blanco Orejinegro cattle. *J. Anim. Breed. Genet.* **138**, 613-627. (doi:10.1111/jbg.12549).

[92] Alshawi, A., Essa, A., Al-Bayatti, S. & Hanotte, O. 2019 Genome analysis reveals genetic admixture and signature of selection for productivity and environmental traits in Iraqi cattle. *Front. Genet.* **10**, 609. (doi:10.3389/fgene.2019.00609).

[93] Fang, L., Hou, Y., An, J., Li, B., Song, M., Wang, X., Sørensen, P., Dong, Y., Liu, C., Wang, Y., et al. 2016 Genome-wide transcriptional and post-transcriptional regulation of innate immune and defense responses of bovine mammary gland to *Staphylococcus aureus*. *Front. Cell. Infect. Microbiol.* **6**, 193. (doi:10.3389/fcimb.2016.00193).

[94] Kumar, L., Chou, J., Yee, C.S., Borzutzky, A., Vollmann, E.H., von Andrian, U.H., Park, S.Y., Hollander, G., Manis, J.P., Poliani, P.L., et al. 2014 Leucine-rich repeat containing 8A (LRRC8A) is essential for T lymphocyte development and function. *J. Exp. Med.* **211**, 929-942. (doi:10.1084/jem.20131379).

[95] Rodriguez-Morales, A.J., Bonilla-Aldana, D.K., Idarraga-Bedoya, S.E., Garcia-Bustos, J.J., Cardona-Ospina, J.A. & Faccini-Martínez, Á.A. 2018 Epidemiology of zoonotic tick-borne diseases in Latin America: Are we just seeing the tip of the iceberg? *F1000Res.* **7**, 1988. (doi:10.12688/f1000research.17649.2).

[96] Ristic, M. & McIntyre, I. 1981 *Diseases of Cattle in the Tropics: Economic and Zoonotic Relevance*. The Hague, Netherlands, Martinus Nijhoff Publishers.

[97] Jaimes-Dueñez, J., Triana-Chávez, O., Holguín-Rocha, A., Tobon-Castaño, A. & Mejía-Jaramillo, A.M. 2018 Molecular surveillance and phylogenetic traits of *Babesia bigemina* and *Babesia bovis* in cattle (*Bos taurus*) and water buffaloes (*Bubalus bubalis*) from Colombia. *Parasit. Vectors* **11**, 510. (doi:10.1186/s13071-018-3091-2).

[98] Fetene, E., Leta, S., Regassa, F. & Büscher, P. 2021 Global distribution, host range and prevalence of *Trypanosoma vivax*: a systematic review and meta-analysis. *Parasit. Vectors* **14**, 80. (doi:10.1186/s13071-021-04584-x).

[99] Jaimes-Dueñez, J., Triana-Chávez, O. & Mejía-Jaramillo, A.M. 2018 Genetic, host and environmental factors associated with a high prevalence of *Anaplasma marginale*. *Ticks Tick Borne Dis.* **9**, 1286-1295. (doi:10.1016/j.ttbdis.2018.05.009).

[100] Naval-Sánchez, M., Porto-Neto, L.R., Cardoso, D.F., Hayes, B.J., Daetwyler, H.D., Kijas, J. & Reverter, A. 2020 Selection signatures in tropical cattle are enriched for promoter and coding regions and reveal missense mutations in the damage response gene *HELB*. *Genet. Sel. Evol.* **52**, 27. (doi:10.1186/s12711-020-00546-6).

[101] Bhar, S., Zhou, F., Reineke, L.C., Morris, D.K., Khincha, P.P., Giri, N., Mirabello, L., Bergstrom, K., Lemon, L.D., Williams, C.L., et al. 2020 Expansion of germline *RPS20* mutation phenotype to include Diamond-Blackfan anemia. *Hum. Mutat.* **41**, 1918-1930. (doi:10.1002/humu.24092).

[102] Wang, R., Yoshida, K., Toki, T., Sawada, T., Uechi, T., Okuno, Y., Sato-Otsubo, A., Kudo, K., Kamimaki, I., Kanezaki, R., et al. 2015 Loss of function mutations in *RPL27* and *RPS27* identified by whole-exome sequencing in Diamond-Blackfan anaemia. *Br. J. Haematol.* **168**, 854-864. (doi:10.1111/bjh.13229).

[103] Noyes, H.A., Alimohammadian, M.H., Agaba, M., Brass, A., Fuchs, H., Gailus-Durner, V., Hulme, H., Iraqi, F., Kemp, S., Rathkolb, B., et al. 2009 Mechanisms controlling anaemia in *Trypanosoma congolense* infected mice. *PLoS ONE* **4**, e5170. (doi:10.1371/journal.pone.0005170).

[104] Martínez, R., Dunner, S., Toro, R., Tobón, J., Gallego, J. & Cañón, J. 2010 Effect of polymorphisms in the *Slc11a1* coding region on resistance to brucellosis by macrophages in vitro and

766 after challenge in two *Bos* breeds (Blanco Orejinegro and Zebu). *Genet. Mol. Biol.* **33**, 463-470.
767 (doi:10.1590/S1415-47572010000300014).

768 [105] Celli, J. 2019 The intracellular life cycle of *Brucella* spp. *Microbiol. Spectr.* **7**.
769 (doi:10.1128/microbiolspec.BAI-0006-2019).