SMARTdb: an integrated database for exploring single-cell multi-omics data of reproductive medicine

Authors: Zekai Liu[1,†], Zhen Yuan[1,†], Yunlei Guo[1,†], Yusheng Guan[1,3], Ruilin Wang[1], Zhanglian Wang[1], Yunan Chen[1], Tianlu Wang[1], Meining Jiang[1], Shuhui Bian[1,2,3]*

[1]State Key Laboratory of Reproductive Medicine and Offspring Health, Nanjing Medical University, Nanjing 211166, P.R. China

[2]Collaborative Innovation Center for Cancer Personalized Medicine, Nanjing Medical University, Nanjing 211166, P.R. China

[3]School of Basic Medical Sciences, Nanjing Medical University, Nanjing 211166, P.R. China

*Correspondence: bianshuhui@njmu.edu.cn (S. Bian)

†Co-first authors.

Abstract

Single-cell multi-omics sequencing has greatly accelerated reproduction research in recent years, and the data are continually growing. However, utilizing those data resources is challenging for wet-lab researchers, and therefore a comprehensive platform for exploring single-cell multi-omics data related to reproduction is urgently needed. Here we introduce SMARTdb (single-cell multi-omics atlas of reproduction), which is an integrative and user-friendly platform for exploring molecular dynamics of reproductive development, aging, and disease, covering multi-omics, multi-species, and multi-stages. We have curated and analyzed single-cell transcriptome and epigenome data of nearly 1.8 million cells from 5 species across whole lifespan. A series of powerful functionalities are provided, such as "Query gene expression", "DIY expression plot", "DNA methylation plot", and "Epigenome browser". With SMARTdb, we found that the male germ-cell-specific expression pattern of *RPL39L* and *RPL10L* is conserved between human and other model animals. Female germ cells also highly express RPL39L in fetal stage, but gradually decrease RPL39L expression in adult stage. Moreover, DNA hypomethylation and open chromatin may regulate the specific expression pattern of RPL39L collectively in both male and female germ cells. In summary, SMARTdb is a powerful platform for convenient data mining and gaining novel insights into reproductive development, aging and disease.

SMARTdb is publicly available at https://smart-db.cn.

Key words: single-cell multi-omics, reproduction, embryo, germ cell, gonad

Introduction

Single-cell multi-omics technique has greatly promoted our molecular understanding

of early embryos[1–14], fetal germ cells and gonadal somatic cells[15–22], developing testis

and ovary[23–30], adult testis and ovary[26,29,31–38], aging testis and ovary[39–41],

endometrium[42,43], maternal-fetal interface[44–46], as well as reproductive diseases[47–52].

Multiple molecular layers, including gene expression patterns, DNA methylation

dynamics, and open chromatin features, have been profiled with high-throughput

sequencing technology, and the data have been growing continually. It is noted that

previous single-cell multi-omics studies typically provide comprehensive atlas rather

than digging into individual genes or genomic sites, but they offer valuable resources

for more wet-lab researchers to perform further investigations into specific questions,

genes, and genomic sites of interest. For example, it is important for researchers using

animal models to know if the expression pattern of a specific gene is conserved in

human, and whether knocking out of the gene could lead to important phenotypic

changes.

However, analyzing the massive and complex single-cell multi-omics data requires

experienced skills and a large number of computing resources, which is challenging

and time-consuming for wet-lab researchers especially. For instance, as each of those

studies typically focuses on a narrow time span, researchers who wish to query the

expression levels of a particular gene in child, young adult, aging adult and infertile

man need to download and process large amount of data from several separate studies,

and they might find this process difficult and laborious due to a lack of analytical skills

and computing resources. This inconvenience hinders the optimal utilization of the

published data, thus limiting potential novel discoveries. Hence, integrated and user-

friendly platforms are needed to bridge the gap between rich data resource and wet-lab

researchers.

Endeavors such as Human Cell Atlas (HCA, https://www.humancellatlas.org/) and

Human Protein Atlas (HPA, https://www.proteinatlas.org/) have served as excellent

platforms for exploring massive single-cell data conveniently. However, these

databases do not contain much reproduction-related data. Currently, there are a few

small-scale reproduction-related websites available, which typically only include data

of their own study, are limited in data amount and time span, only provide simple

function such as querying gene expression of a specific gene, and lack long-term

maintenance. As a result, there is still an urgent need for an integrated and user-friendly

platform which focuses on promoting efficient utilization of public single-cell multi-

omics data related to reproduction across full life-cycle.

Here we introduce SMARTdb (single-cell multi-omics atlas of reproduction), which is

a manually curated, integrative and interactive platform aiming to bridge the gap

between mass data resources and wet-lab researchers, and promote exploring molecular

dynamics of reproductive development, aging, and disease.

Data collection and processing

Data collection

We curated single-cell transcriptome and epigenome (DNA methylation and chromatin

accessibility) data related to reproduction from published studies (Table S1). The gene

expression matrix, DNA methylation, and chromatin accessibility data were

downloaded from NCBI GEO (https://www.ncbi.nlm.nih.gov/geo), EMBL-EBI

(https://www.ebi.ac.uk/), and Zenodo (https://zenodo.org/). In addition to data of

developmental stages and normal physiological states, data of aging and human

infertility (such as non-obstructive azoospermia, NOA) are also incorporated. In

summary, the single-cell data of nearly 1.8 million individual cells across five species

(human, monkey, mouse, pig, and buffalo) were curated, involving early embryos, fetal

germ cells and gonadal somatic cells, developing testis and ovary, adult testis and ovary,

aging testis and ovary, endometrium, maternal-fetal interface, and human infertility.

These data cover the entire life cycle, and span >100 specific time points of 5 species

(Table S1).

Processing of single-cell transcriptome data and cell clustering

To reproduce the cell clustering results of original studies as much as possible, for the

datasets with metadata provided by the original study, only the cells with cell type

identities were used in downstream analyses. For the datasets without provided

metadata, the cells were filtered according to the quality control standards mentioned

in the original studies. For those studies not providing concrete cutoff values, only cells

with >500 expressed genes and <20% mitochondrial gene ratios were retained. Seurat[53]

(version 4.3.0) was used to perform the following processing steps. Briefly, the UMI

count of each gene for each cell was first divided by the total counts for that cell and

multiplied by 10,000 (or author-designated parameters), and was then natural-log

transformed. The top highly variable genes among cells were selected with

"FindVariableFeatures" function with default or author-designated parameters. The

normalized data were then centered and scaled, resulting in that the mean expression of

each gene across all cells was zero and the standard deviation was one. Principal

component analysis (PCA) was performed on the scaled data with the previously

determined highly variable genes used as input. Then the cells were clustered with

Louvain algorithm. The uniform manifold approximation and projection (UMAP) was

used to visualize the cell clustering in low-dimensional space. For the datasets with

batch effects, the anchors of different samples were identified and then different

samples were integrated to correct batch effects. For the datasets without obvious batch

effects, no additional correction was performed.

Cell type annotation and differentially expressed gene analysis of single-cell

transcriptome data

A series of analyses were performed to ensure the accuracy of cell identities for each

dataset. For the datasets with metadata provided by the original study, the cells were

annotated correspondingly, and classical marker genes for each cell type were checked

to confirm accuracy. For the datasets without metadata, the top differentially expressed

genes (DEGs) of each cell cluster were identified using the Wilcoxon rank sum test

with the "FindAllMarkers" function in Seurat[53] (version 4.3.0) with default parameters

(logfc.threshold=0.25, min.pct=0.1, return.thresh=0.01), and their consistency with

classical marker genes for each cell type was carefully checked. After these steps, the

cell type identities can be annotated accurately.

Visualization of gene expression levels

First, Seurat objects were converted into "anndata" objects with the R package sceasy (version 0.0.7). Then, the dot plots, heatmaps and violin plots were generated with Scanpy[54] (version 1.9.3). For dot plots and heatmaps, both the relative expression levels with and without standard scaling are provided. The standard scale normalization strategy in Scanpy was performed with parameters "standard_scale = 'var'", which standardized the expression level of a given gene between 0 and 1 by subtracting the minimum and dividing each by its maximum.

Identification of transcription factor motifs

The *de novo* motif discovery was performed with HOMER software (version 4.11)[55] with default parameters.

Processing of single-cell epigenome data

The processed single-cell DNA methylation and chromatin accessibility data provided by the original study were downloaded from GEO database, and were further processed to standard BigWig format with bedtools (version 2.30.0), UCSC bedGraphToBigWig (version 2.9) and bigWigMerge (version 2) for downstream usage. For the data generated by scBS-seq[56] technique, only CpG sites were retained. For the data generated by scCOOL-seq[8] technique, WCG (W=A, T) sites were used for DNA methylation analysis and GCH (H=A, C, T) sites were used for chromatin accessibility analysis.

Considering the sparse coverage of single-cell epigenome data, we merged single-cell epigenome data according to their cell type identities and stages (e.g. 2-cell embryo, 8-week female mitotic germ cells, etc.), and sequencing techniques (e.g. scBS-seq and scCOOL-seq) using custom scripts. Briefly, for the genomic sites only covered by one cell, they were directly retained in the merged files. For the genomic sites covered by multiple cells in the group, the mean values of covered cells were calculated to represent the final states of the sites. For single-cell DNA methylation data, only the genomic sites with DNA methylation levels <0.1 or >0.9 in each individual cell were used in downstream merging. The merged single-cell data of cell groups are more convenient and efficient for downstream exploration.

The DNA methylation "Tanghulu" plot was implemented by a custom Python script to visualize DNA methylation state of each CpG site in a narrow genomic region of interest. For single-cell DNA methylation data, only the genomic sites with values <0.1 or >0.9 in each individual cell will be plotted. The WashU epigenome browser[57] was incorporated into SMARTdb for convenient visualization of single-cell DNA methylation levels and chromatin accessibility states.

Implementation of the database website

SMARTdb was developed utilizing Python 3 in conjunction with the Django framework (https://djangoproject.com). The data housed within SMARTdb is systematically organized and accessible through PostgreSQL (https://www.postgresql.org). The user-friendly front-end interface was created using NG-ZORRO and Angular. The platform is hosted using the Nginx web server (http://nginx.org) and is freely accessible online at https://smart-db.cn without the need for registration or login. For optimal performance, SMARTdb is best used on major web browsers such as Google Chrome, Microsoft Edge, Safari, and Mozilla Firefox, with JavaScript enabled.

Database content and usage

Overall design

SMARTdb is an integrative and interactive platform for exploring single-cell multi-omics data related to reproduction. It was designed to have three main features (Figures 1 and S1). First, single-cell multi-omics data of nearly 1.8 million individual cells were manually curated and analyzed (Table S1). Users can access gene expression levels, DNA methylation levels and chromatin accessibility levels at single-cell resolution conveniently. Second, the main stages of full life-cycle were covered, including zygote, early embryo, fetus, infant, child, puberty, adult, and aging. More than 100 specific time points are included in total. Besides, data of human infertility are also incorporated. Users can not only explore molecular dynamics during full life-cycle, but also compare normal physiological state with disease. Third, considering cross-species conservation analyses are very important, data from five species were included, including human, non-human primate (monkey), mouse, pig, and buffalo. For example, the researchers

using mouse as animal model can find out if the expression pattern of the gene that they
decide to investigate is conserved in human, which will help select more meaningful
research targets.
As a user-friendly platform, SMARTdb offers multiple featured function modules to
users for exploring the rich single-cell multi-omics data resources, such as "Query gene
expression", "DIY expression plot", "DNA methylation plot" and "Epigenome
browser". The featured function modules are described below.
206

"Overview" and "Query gene expression" modules

In the "Overview" module, SMARTdb facilitates users to get a quick overview of cell
clustering, sample origins, cell counts, and DEGs of each cell type for each scRNA-seq
dataset (Figure 2A). In the "Query gene expression" module, users can conveniently
plot the expression levels of a selected gene across the cell types in the chosen dataset
with UMAP and violin plots (Figure 2B). Additionally, a concise introduction for each
human gene is also provided in this module.
214

"DIY expression plot" module

This module provides a very practical function for users to customize expression plots
according to their personalized requirements (Figure 2C). Users can select multiple
genes and multiple cell types of interest to plot simultaneously, and choose appropriate
plot forms for visualization. Three main visualization plots are offered in this module,
including dot plots, heatmaps, and violin plots. Moreover, the orders of genes and cell
types displayed in the plots can be designated by users freely. Users can also choose
whether to standardly scale the expression levels within each gene when using dot plots
and heatmaps. The generated figures are of publication quality.
224

"DNA methylation plot" module

This module provides "Tanghulu" plots to visualize DNA methylation status of each
cytosine site in CpG or WCG context (Figure 3A). Users can conveniently generate a
customized "Tanghulu" plot by selecting datasets and cell types of interest, and
designating specific genomic region range. In "Tanghulu" plot, each row represents an
individual cell, and each circle represents a cytosine site in CpG or WCG context. Black
circles stand for methylated sites, while the white circles refer to unmethylated sites.

"Epigenome browser" module

This module incorporates the WashU epigenome browser tool[57] and offers a state-of-

the-art solution for browsing epigenomic datasets (Figure 3B). Users can select single-

cell DNA methylation and chromatin accessibility datasets of various cell types based

on their interest, such as oocytes, sperms, early embryos, fetal germ cells, and gonadal

somatic cells. The DNA methylation and chromatin accessibility levels of selected

samples can be directly and conveniently visualized in the browser. Especially, since

single-cell data are typically sparse, the single-cell data of each cell type or sample type

are merged for optimal visualization and enabling more efficient exploration. The

genomic region locator allows users to navigate to a specific gene or designated

genomic region, zoom in and out, and move to new regions by dragging the mouse left

and right. Additionally, the browser interface is allowed to be customized to meet

personalized requirements, such as display mode (bar plot or heatmap), RGB colors,

axis scales, and text labels.

Examples for using SMARTdb

Here we illustrate how to make good use of SMARTdb to facilitate exploration of

single-cell multi-omics data and promote new findings. A recent study has revealed that

RPL39L is a male germ-cell-specific ribosome protein and plays critical roles in

spermatogenesis and male fertility in mouse[58]. Deletion of *Rpl39l* in mice can lead to

defective sperm formation and reduced fertility[58]. With high-purity enrichment and

proteomic analyses for mouse spermatogonia cells, pachytene spermatocytes, round

spermatids, elongated spermatids, and somatic cells (Sertoli cells and interstitial cells),

the study showed that RPL39L was mainly expressed in germ cells, whereas the

corresponding core ribosome protein RPL39 displayed decreased expression levels in

germ cells during spermatogenesis and was mainly expressed in Sertoli cells and

interstitial cells. The study also mentioned that the expression of RPL10L was testis-

specific in bulk level, but its expression levels in specific subtypes of male germ cells

were not revealed. Inspired by the study, we further asked:

(1) Whether the germ-cell-specific expression pattern is conserved in testis of human,

non-human primates as well as other animal models?

(2) When does the germ-cell-specific expression pattern of RPL39L form? In fetal stage, peri-puberty or adulthood?

(3) Does epigenetic regulation function in regulating the male germ-cell-specific expression pattern? We demonstrate how to explore the three questions with SMARTdb below.

Cross-species conservation

To answer the first question, we utilized the "DIY expression plot" module to plot the expression levels of the ribosomal large subunit protein genes *RPL39L* and *RPL10L*, and the corresponding core ribosome protein genes *RPL39* and *RPL10* in human[36], as well as their homologous genes in mouse[37], monkey[32], and pig[28] (Figures 2C and 4A). For human adult testis[36], taking the advantage of accurate cell type identification with single-cell RNA-seq data, we found that the expression levels of *RPL39L* and *RPL10L* peak at spermatocyte (especially pachytene and/or diplotene spermatocyte), and then reduce in later stages of spermatogenesis at mRNA level (Figure 2, B and C). In contrast, *RPL39* and *RPL10* decreased their expression levels during spermatogenesis, and mainly expressed in somatic cells (Figure 2C). Moreover, the germ-cell-specific expression pattern of *RPL39L* and *RPL10L*, and decreasing of *RPL39* and *RPL10* during spermatogenesis are conserved in human and other species (Figure 4A).

Developing dynamics

To clarify the second question, we plotted the expression levels of *RPL39L* in fetal stage[18] and peri-pubertal stage[24] with the "DIY expression plot" module (Figure 4B).We found that the germ cells in fetal stages have started to highly express *RPL39L*, as early as in migrating primordial germ cells (PGCs). Meanwhile, although the somatic niche cells in fetal stage relatively lowly expressed *RPL39L*, their expressed percentages were still higher than those of postnatal stages (peri-puberty and adulthood), indicating that the germ-cell specific pattern of *RPL39L* had started to establish but had not been completely established in fetal stage at mRNA level. In human male peri-puberty, the germ-cell-specific pattern of *RPL39L* has been established, with high expression in germ cells and low expression in somatic cells (both in the expressed percentages and mean expression levels). Additionally, for the corresponding core ribosome protein gene *RPL39*, some subtypes of fetal germ cells still have relatively

high expression levels of *RPL39*, such as male migrating PGCs, and both male and female mitotic germ cells. Meanwhile, in human male peri-pubertal stage, the expression levels and expressed percentage of *RPL39* decreased gradually from spermatogonia to spermatid, leaving only somatic cells with relatively high expression levels.

Moreover, we noticed that *RPL39L* transcripts are highly expressed in both male and female germ cells in human fetal stage, but diverge between male and female in adulthood. The fully-grown, MI, and MII oocytes sharply decrease the expression levels of *RPL39L*, leaving only growing oocytes in human adult ovary[34] maintaining relatively high expression levels of *RPL39L* (Figure 5A). As a result, *RPL39L* is expressed in male germ cells and female growing oocytes in human adulthood. The divergence in the *RPL39L* expression among oocytes of various stages in human adult ovary was detected in mRNA level, indicating potential transcriptional regulation of the gene.

Regulation mechanism

Then we further explored the transcriptional regulatory mechanism using the "Epigenome browser" module. With epigenetic data of human fetal stage[15], we found that germ cells in fetal stage have hypomethylated and more accessible regions around the promoter of *RPL39L* compared with somatic cells (Figures 3B and S2), which is consistent with the relatively high expression of *RPL39L* of fetal germ cells (Figure 4B). In human adult female[34], growing oocytes displayed similar DNA methylation and chromatin accessibility pattern with fetal germ cells (Figure 5B). In contrast, fully-grown, MI, and MII oocytes up-regulate the DNA methylation levels around *RPL39L* promoter, and down-regulate the chromatin accessibility levels around *RPL39L* promoter (Figure 5B), which may explain the decreased expression level of RPL39L in fully-grown, MI, and MII oocytes. Hence, epigenetic regulation potentially plays important roles in shaping the expression pattern of *RPL39L* in both male and female human germ cells during both fetal and adult stages.

In summary, we validated that *RPL39L* and *RPL10L* are germ-cell-specific at mRNA level, which is consistent with previous studies[58]. Moreover, we obtained multiple

novel findings with SMARTdb. We found that the male germ-cell-specific expression pattern is conserved between human and other model animals, illustrating its importance during evolution. The germ-cell-specific pattern of *RPL39L* at mRNA level has started to establish but has not been completely established in fetal gonad. In human adulthood, *RPL39L* is relatively highly expressed in male germ cells and female growing oocytes, but lowly expressed in fully-grown, MI, and MII oocytes. Furthermore, DNA methylation and chromatin accessibility may cooperate in regulating the divergence of *RPL39L* expression in germ cells of human adult at mRNA level. These results demonstrate the power of SMARTdb, providing valuable clues for further investigations.

Discussion

The mystery of reproductive development, aging and disease has long been explored. Single-cell multi-omics sequencing technique has revolutionized researches in reproduction in recent years, and has led to accelerated accumulation of massive datasets. The public data provide valuable resources for researchers to perform further data mining. However, difficulties in processing big data for broad wet-lab researchers prevent them from being effectively utilized. To deal with the situation of growing data and promote a more efficient usage, we present SMARTdb – a single-cell multi-omics platform for reproduction research. To the best of our knowledge, SMARTdb is the first integrated platform with multiple powerful functionalities to explore single-cell multi-omics data in the field of reproductive medicine. Moreover, it will be continuously updated by adding high-quality single-cell multi-omics data from reproductive research.

Our database will promote reproductive medicine research in several aspects. For example, researchers using animal models are able to speculate whether knocking out of a specific gene could potentially cause important phenotypic changes. Users can know whether the expression pattern of a specific gene is conserved in human, which helps select more meaningful research targets. For researcher using human samples, cross-species conservation analyses provide important clues for selecting suitable animal models for further validation. Cross-stage dynamic analysis helps select the

right time window for further investigations. Moreover, epigenetics data resources will

provide novel insights for potential regulatory mechanisms.

We showed examples for using SMARTdb and demonstrated the power of it by

exploring germ-cell-specific *RPL39L* and *RPL10L*. We not only validated the germ-

cell-specific expression of ribosome gene *RPL39L*, but also gained a deeper

understanding of its cross-species conservation, cross-stage expression dynamics, and

potential epigenetic regulation. Moreover, taking the advantages of single-cell

transcriptome sequencing, the gene expression patterns in multiple cell types can be

observed more accurately and in detail. For example, we found that *RPL39L* was not

only specifically expressed in male germ cells in human adult testis, but also highly

expressed in the growing oocytes (one subtype of oocytes) in human adult ovary.

In conclusion, we firmly believe that with the assistance of SMARTdb, researchers in

the field of reproduction can easily access and explore massive single-cell multi-omics

data, and hopefully, inspire novel ideas and new findings.

References

1. Guo, H. *et al.* The DNA methylation landscape of human early embryos. *Nature* **511**, 606–610 (2014).

2. Zhu, P. *et al.* Single-cell DNA methylome sequencing of human preimplantation embryos. *Nat. Genet.* **50**, 12–19 (2018).

3. Blakeley, P. *et al.* Defining the three cell lineages of the human blastocyst by single-cell RNA-seq. *Development* **142**, 3613–3613 (2015).

4. Yan, L. *et al.* Single-cell RNA-seq profiling of human preimplantation embryos and embryonic stem cells. *Nat. Struct. Mol. Biol.* **20**, 1131–1139 (2013).

5. Zhou, F. *et al.* Reconstituting the transcriptome and DNA methylome landscapes of human implantation. *Nature* **572**, 660–664 (2019).

6. Petropoulos, S. *et al.* Single-cell RNA-seq reveals lineage and X chromosome dynamics in human preimplantation embryos. *Cell* **165**, 1012–1026 (2016).

7. Guo, H. *et al.* Single-cell methylome landscapes of mouse embryonic stem cells and early embryos analyzed using reduced representation bisulfite sequencing. *Genome Res.* **23**, 2126–2135 (2013).

8. Guo, F. *et al.* Single-cell multi-omics sequencing of mouse early embryos and embryonic stem cells. *Cell Res.* **27**, 967–988 (2017).

9. Jiang, S. *et al.* Single-cell chromatin accessibility and transcriptome atlas of mouse embryos. *Cell Rep.* **42**, 112210 (2023).

10. Wang, Y. *et al.* Single-cell multi-omics sequencing reveals the functional regulatory landscape of early embryos. *Nat. Commun.* **12**, 1247 (2021).

11. Scialdone, A. *et al.* Resolving early mesoderm diversification through single-cell expression profiling. *Nature* **535**, 289–293 (2016).

12. Argelaguet, R. *et al.* Multi-omics profiling of mouse gastrulation at single-cell resolution. *Nature* **576**, 487–491 (2019).

13. Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature* **566**, 490–495 (2019).

14. Li, L. *et al.* Single-cell multi-omics sequencing of human early embryos. *Nat. Cell Biol.* **20**, 847–858 (2018).

15. Li, L. *et al.* Dissecting the epigenomic dynamics of human fetal germ cell development at single-cell resolution. *Cell Res.* **31**, 463–477 (2021).

16. Chitiashvili, T. *et al.* Female human primordial germ cells display X-chromosome dosage compensation despite the absence of X-inactivation. *Nat. Cell Biol.* **22**, 1436–1446 (2020).

17. Garcia-Alonso, L. *et al.* Single-cell roadmap of human gonadal development. *Nature* **607**,
540–547 (2022).

18. Li, L. *et al.* Single-cell RNA-seq analysis maps development of human germline cells and
gonadal niche interactions. *Cell Stem Cell* **20**, 858-873.e4 (2017).

19. Guo, F. *et al.* The transcriptome and DNA methylome landscapes of human primordial
germ cells. *Cell* **161**, 1437–1452 (2015).

20. Stévant, I. *et al.* Dissecting cell lineage specification and sex fate determination in gonadal
somatic cells using single-cell transcriptomics. *Cell Rep.* **26**, 3272-3283.e3 (2019).

21. Chen, M. *et al.* Integration of single-cell transcriptome and chromatin accessibility of early
gonads development among goats, pigs, macaques, and humans. *Cell Rep.* **41**, 111587
(2022).

22. Stévant, I. *et al.* Deciphering cell lineage specification during male sex determination with
single-cell RNA sequencing. *Cell Rep.* **22**, 1589–1599 (2018).

23. Guo, J. *et al.* Single-cell analysis of the developing human testis reveals somatic niche cell
specification and fetal germline stem cell establishment. *Cell Stem Cell* **28**, 764-778.e4
(2021).

24. Guo, J. *et al.* The dynamic transcriptional cell atlas of testis development during human
puberty. *Cell Stem Cell* **26**, 262-276.e4 (2020).

25. Zhao, J. *et al.* Cell-fate transition and determination analysis of mouse male germ cells
throughout development. *Nat. Commun.* **12**, 6839 (2021).

26. Ernst, C., Eling, N., Martinez-Jimenez, C. P., Marioni, J. C. & Odom, D. T. Staged
developmental mapping and X chromosome transcriptional dynamics during mouse
spermatogenesis. *Nat. Commun.* **10**, 1251 (2019).

27. Huang, L. *et al.* Single-cell RNA sequencing uncovers dynamic roadmap and cell-cell
communication during buffalo spermatogenesis. *iScience* **26**, 105733 (2022).

28. Zhang, L. *et al.* Single-cell RNA-seq analysis of testicular somatic cell development in pigs.
J. Genet. Genomics **49**, 1016–1028 (2022).

29. Sohni, A. *et al.* The neonatal and adult human testis defined at the single-cell level. *Cell*
Rep. **26**, 1501-1517.e4 (2019).

30. Gu, C., Liu, S., Wu, Q., Zhang, L. & Guo, F. Integrative single-cell analysis of
transcriptome, DNA methylome and chromatin accessibility in mouse oocytes. *Cell Res.*
29, 110–123 (2019).

31. Chen, Y. *et al.* Single-cell RNA-seq uncovers dynamic processes and critical regulators in
mouse spermatogenesis. *Cell Res.* **28**, 879–896 (2018).

32. Shami, A. N. *et al.* Single-cell RNA sequencing of human, macaque, and mouse testes
uncovers conserved and divergent features of mammalian spermatogenesis. *Dev. Cell* **54**,
529-547.e12 (2020).

33. Fan, X. *et al.* Single-cell reconstruction of follicular remodeling in the human adult ovary.
Nat. Commun. **10**, 3164 (2019).

34. Yan, R. *et al.* Decoding dynamic epigenetic landscapes in human oocytes using single-cell
multi-omics sequencing. *Cell Stem Cell* **28**, 1641-1656.e7 (2021).

35. Zhang, Y. *et al.* Transcriptome landscape of human folliculogenesis reveals oocyte and
granulosa cell interactions. *Mol. Cell* **72**, 1021-1034.e4 (2018).

36. Guo, J. *et al.* The adult human testis transcriptional cell atlas. *Cell Res.* **28**, 1141–1157
(2018).

37. Green, C. D. *et al.* A comprehensive roadmap of murine spermatogenesis defined by single-
cell RNA-seq. *Dev. Cell* **46**, 651-667.e10 (2018).

38. Hermann, B. P. *et al.* The mammalian spermatogenesis single-cell transcriptome, from
spermatogonial stem cells to spermatids. *Cell Rep.* **25**, 1650-1667.e8 (2018).

39. Nie, X. *et al.* Single-cell analysis of human testis aging and correlation with elevated body
mass index. *Dev. Cell* **57**, 1160-1176.e5 (2022).

40. Huang, D. *et al.* A single-nucleus transcriptomic atlas of primate testicular aging reveals
exhaustion of the spermatogonial stem cell reservoir and loss of Sertoli cell homeostasis.
Protein Cell pwac057 (2022).

41. Wang, S. *et al.* Single-cell transcriptomic atlas of primate ovarian aging. *Cell* **180**, 585-
600.e19 (2020).

42. Garcia-Alonso, L. *et al.* Mapping the temporal and spatial dynamics of the human
endometrium in vivo and in vitro. *Nat. Genet.* **53**, 1698–1711 (2021).

43. Wang, W. *et al.* Single-cell transcriptomic atlas of the human endometrium during the
menstrual cycle. *Nat. Med.* **26**, 1644–1653 (2020).

44. Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal–fetal interface in
humans. *Nature* **563**, 347–353 (2018).

45. Arutyunyan, A. *et al.* Spatial multi-omics map of trophoblast development in early
pregnancy. *Nature* **616**, 143–151 (2023).

46. Greenbaum, S. *et al.* A spatially resolved timeline of the human maternal–fetal interface.
Nature **619**, 595–605 (2023).

47. Zhao, L. *et al.* Single-cell analysis of developing and azoospermia human testicles reveals
central role of Sertoli cells. *Nat. Commun.* **11**, 5683 (2020).

48. Di Persio, S. *et al.* Single-cell RNA-seq unravels alterations of the human spermatogonial
stem cell compartment in patients with impaired spermatogenesis. *Cell Rep. Med.* **2**,
100395 (2021).

49. Alfano, M. *et al.* Aging, inflammation and DNA damage in the somatic testicular niche
with idiopathic germ cell aplasia. *Nat. Commun.* **12**, 5205 (2021).

50. Mahyari, E. *et al.* Comparative single-cell analysis of biopsies clarifies pathogenic
mechanisms in Klinefelter syndrome. *Am. J. Hum. Genet.* **108**, 1924–1945 (2021).

51. Chen, Y. *et al.* Deciphering the molecular characteristics of human idiopathic
nonobstructive azoospermia from the perspective of germ cells. *Adv. Sci.* 2206852 (2023).

52. Ferrero, H. *et al.* Single-cell RNA sequencing of oocytes from ovarian endometriosis
patients reveals a differential transcriptomic profile associated with lower quality. *Hum.*
Reprod. **34**, 1302–1312 (2019).

53. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
(2021).

54. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biol.* **19**, 15 (2018).

55. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-
regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589
(2010).

56. Smallwood, S. A. *et al.* Single-cell genome-wide bisulfite sequencing for assessing
epigenetic heterogeneity. *Nat. Methods* **11**, 817–820 (2014).

57. Li, D. *et al.* Washu epigenome browser update 2022. *Nucleic Acids Res.* **50**, W774–W781
(2022).

58. Li, H. *et al.* A male germ-cell-specific ribosome controls male fertility. *Nature* **612**, 725–
731 (2022).
506
507

Acknowledgments

We thank professors Fuchou Tang, Jingchu Luo, Zhibin Hu, Jiahao Sha, and Ran Huo for discussion and suggestions. This work was supported by Young Elite Scientists Sponsorship Program by China Association for Science and Technology (CAST) (2020QNRC001) and research start-up funding from Nanjing Medical University (KY116RC20200007).

Author contributions

Shuhui Bian: Conceptualization, supervision, Writing - original draft, Writing - review & editing, Visualization, Data curation, Methodology, Formal analysis, Funding acquisition, Investigation, Project administration. **Zekai Liu:** Data curation, Formal analysis, Methodology, Software, Visualization, Writing - original draft. **Zhen Yuan:** Data curation, Formal analysis, Methodology, Software, Visualization, Writing - original draft. **Yunlei Guo:** Formal analysis, Methodology, Software, Visualization, Validation. **Yusheng Guan:** Data curation, Methodology, Validation.

Competing interests

The authors declare that they have no competing interests.

Data availability

SMARTdb is publicly available at https://smart-db.cn.

Figure 1

Figure 1. A brief schema describing data curation, web construction and featured

functionalities. The single-cell data of nearly 1.8 million individual cells across five species

(human, monkey, mouse, pig, and buffalo) were curated, covering the entire life cycle and

spanning >100 specific time points of 5 species. Multiple featured function modules are

provided, such as "Query gene expression", "DIY expression plot", "DNA methylation plot"

and "Epigenome browser".

Figure 2

Figure 2. Featured functionalities for exploring single-cell transcriptome datasets.

(A) The "Overview" module provides UMAP plots showing cell type and sample

identity, bar plots showing cell number and proportion of each cell type, and tables

showing differentially expressed genes for each single-cell dataset. The single-cell

RNA-seq data of human adult testis (GSE112013) was shown as an example.

(B) The "Query gene expression" module enables visualization of the expression levels

of each selected gene with UMAP plot (left) and violin plot (right). The expression level

of a germ-cell specific ribosomal gene *RPL39L* in human adult testis was shown.

(C) The "DIY expression plot" module allows users to create customized plots

displaying the expression levels of selected genes in selected cell types. Three types of

plots (dot plot, violin plot and heatmap) are provided. The expression levels of

representative marker genes and four ribosomal genes in major cell types of human

adult testis were shown.

The description of cell type abbreviation are listed below. SSC, spermatogonia stem

cell. SPG, spermatogonia. L, leptotene spermatocyte. Z, zygotene spermatocyte. P,

pachytene spermatocyte. D, diplotene spermatocyte. SPC7, a mixture of diakinesis,

metaphase, anaphase, telophase, and secondary spermatocytes.

Figure 3

A DNA methylation plot

DNA demethylation region in morula
(chr2:155396001-155396500)

de novo DNA methylation region in morula
(chr16:72930501-72931000)

B Epigenome browser

● DNA methylation

PGC and soma (before 11 weeks)

● Chromatin accessibility

PGC and soma (before 11 weeks)

Figure 3. Featured functionalities for exploring single-cell epigenome datasets.

(A) The representative DNA demethylation and *de novo* methylation regions between

human 8-cell embryo and morula were shown. Each row in the "Tanghulu" plot

corresponds to an individual cell, and each circle represents a CpG site. Methylated

CpG sites are represented by black dots, while unmethylated CpG sites are indicated by

white dots. The grey dots at the bottom indicate covered CpG sites in selected cells.

The CpG sites which were not covered by any cell were omitted. The single-cell DNA

methylation data of human early embryo (GSE100272) was used.

(B) The screenshots of epigenome browser displaying the DNA methylation levels and

chromatin accessibility levels around the promoter region of *RPL39L* in primordial

germ cells and somatic cells. The grey rectangles highlight regions showing significant

differences between somatic niche cells and germline cells. PGC, primordial germ cell.

Figure 4

A

Cross-species conservation of model animals

Mouse testis (GSE112393)

Rpl39l Rpl10l Rpl39 Rpl10

Adult germ cell
- Spermatogonia
- Spermatocyte
- Round spermatid
- Elongat'spermatid

Soma
- Sertoli cell
- Leydig cell
- Myoid cell
- Endothelial cell
- Macrophage

Monkey testis (GSE142585)

RPL39L RPL10

Adult germ cell
- Spermatogonia
- Spermatocyte
- Round spermatid
- Elongat'spermatid

Soma
- Immature Leydig
- Myoid
- Endothelial
- Macrophage
- T cell

Pig testis (GSE174782)

RPL10L RPL10

Postnatal germ cell
- Spermatogonia
- Spermatocyte
- Spermatid

Soma
- Sertoli cell
- Leydig cell
- Myoid
- Endothelial
- Macrophage
- T cell

Relative expression level (standard scale)

Low — High

Expressed percentage

20% 40% 60% 80% 100%

B

Cross-stage analysis

Human fetal stage (UMAP)

- ♂ Migrating FGC
- Mitotic FGC ♀ & ♂
- ♀ RA-responsive FGC
- Endothelial
- ♂ Mitotic-arrest FGC
- Other somatic
- Sertoli
- ♀ Meiotic FGC
- ♀ Oogenesis FGC

UMAP_2 / UMAP_1

Human peri-pubertal stage (UMAP)

- Macrophage
- Sertoli
- Smooth muscle
- Spermatogonia
- Spermatid
- Spermatocyte
- PMC & Leydig
- Endothelial

UMAP_2 / UMAP_1

Human fetal gonad (Dotplot)

RPL39L RPL39

♀FGC
- Mitotic FGC
- RA-response FGC
- Meiotic FGC
- Oogenesis FGC

♂FGC
- Migrating FGC
- Mitotic FGC
- Mitotic-arrest FGC

Soma
- Early granulosa
- Mural granulosa
- Late granulosa
- Sertoli cell
- Leydig precursor
- Differentiated leydig
- Endothelial

Human peri-pubertal testis (Dotplot)

RPL39L RPL39

♂ Germ cells
- Spermatogonia
- Spermatocyte
- Spermatid

Soma
- Sertoli cell
- PMC & Leydig
- Smooth muscle
- Endothelial cell
- Macrophage

Expressed percentage

20% 40% 60% 80% 100%

Relative expression level (standard scale)

Low — High

Figure 4. Cross-species and cross-stage analyses of ribosomal genes using
SMARTdb.

(A) The dot plots show the expression levels of four ribosomal genes in mouse,
cynomolgus monkey, and pig. PMC, peritubular myoid cell. IP, interstitial progenitor
cell.

(B) The UMAP plots show the cell type identities of each cell cluster. The expression
levels of two ribosomal genes (*RPL39L* and *RPL39*) in human germline and somatic
niche cells of fetal stage and peri-pubertal stage were shown in dot plots. FGC, fetal
germ cell. PMC, peritubular myoid cell.

Figure 5

A

B

● DNA methylation

● Chromatin accessibility

Figure 5. The gene expression, DNA methylation and chromatin accessibility

patterns of *RPL39L* in human ovary.

(A) The leftmost UMAP plot shows the cell type identities of human adult ovary. The

expression levels of *RPL39L* in various cell types of human adult ovary are shown on

the middle UMAP plot and the rightmost dot plot.

(B) The screenshots of epigenome browser displaying the DNA methylation levels and

chromatin accessibility levels around the promoter region of *RPL39L* in human adult

ovary. The grey rectangles highlight regions showing changes in fully grown, MI, and

MII oocytes compared with growing oocytes. GO, growing oocyte. FGO, fully grown

oocyte. MI, metaphase I. MII, metaphase II.

Figure S1

Stage/state

Development

Young adult

Aging

Disease

Sample type

Embryo

Fetal gonad

Infant testis

Prepubertal testis

Prepubertal ovary

Pubertal testis

Adult testis

Adult epididymis

Adult ovary

Endometrium

Maternal-fetal interface

Aging testis

Aging ovary

Adult testis of diabetic

OA

NOA (idiopathic)

NOA (AZF deletion)

Cryptozoospermia

Kinefelter syndrome

Species

Human

Monkey

Mouse

Pig

Buffalo

Omics

Transcriptome

Epigenome

588 **Figure S1. The Sankey plot shows the composition of datasets included in**
589 **SMARTdb.** OA, obstructive azoospermia. NOA, Non-obstructive azoospermia.
590

Figure S2

● DNA methylation

Human fetal germ cells and soma (after 11 weeks)

● Chromatin accessibility

Human fetal germ cells and soma (after 11 weeks)

Figure S2. The DNA methylation levels and chromatin accessibility levels around

the *RPL39L* promoter in fetal germ cells after 11 weeks.

The screenshots of epigenome browser displaying the DNA methylation levels and

chromatin accessibility levels around the *RPL39L* promoter in fetal germ cells after 11

weeks post fertilization. The grey rectangles highlight regions showing significant

differences between germ cells and somatic niche cells. The mean chromatin

accessibility levels of the regions which are highlighted by grey rectangles are labelled.