Title page

Authors

Shohei Shibata[1], Takeshi Furuichi[1]

Title

Comparative analysis of intragroup intermale relationships: A study of wild bonobos

(*Pan paniscus*) in Wamba, Democratic Republic of Congo and chimpanzees (*Pan*

troglodytes) in Kalinzu Forest Reserve, Uganda

Affiliations and Address

[1] Wildlife Research Center, Kyoto University, Inuyama, Aichi 484-8506, Japan

Correspondence

Name: Shohei Shibata

ORCID: 0000-0002-3605-2502

Address: Kyoto University Wildlife Research Center, 41-2, Kanrin, Inuyama, Aichi

484-8506, Japan

Email: shibata.shohei.d12@kyoto-u.jp

16
17

Abstract

Although chimpanzees (*Pan troglodytes*) and bonobos (*Pan paniscus*) share a multi-

male/multi-female societal organization and form male-philopatric groups, disparities in

terms of male aggression and stability of temporary parties are thought to exist among

them. However, existing research in bonobos has mainly focused on the high social

status, prolonged receptivity, and characteristic sexual behaviors of females, leaving the

behaviors of males understudied. Moreover, prior comparative studies on *Pan* suffer

from methodological inconsistencies. This study addresses these gaps by employing a

uniform observation method to explore party attendance and aggressive interactions

among male bonobos in Wamba and male chimpanzees in Kalinzu. Unlike male

chimpanzees that exhibit dispersion in the absence of receptive females in the group,

male bonobos showed a lesser degree of such dispersion. Although the overall

frequency of aggressive interactions per observation unit did not significantly differ

between the two species, the nature of these interactions varied. Notably, severe

aggressive behaviors such as physical confrontations among adult males were absent in

bonobos, with most aggression occurring between the sons of the two highest-ranking

females. Additionally, in bonobos, females actively engaged in polyadic aggressive

behavior as aggressors, while all instances of coalitionary aggression in chimpanzees

originated from male aggressors. These findings underscore the substantial impact of

female behaviors on the observed distinctions in male aggressive interactions between

the two species.

Keywords

Pan paniscus, Pan troglodytes, aggression, fission–fusion dynamics, party size,

dominance rank, Wamba, Kalinzu Forest Reserve

Introduction

Factors concerning intermale mating competition primarily influence the reproductive strategies and social relationships of primate males. These factors include the number of receptive females, mating seasonality, operational sex ratio, dominance rank among males in a group, and relationships with other groups or out-group males. Among male chimpanzees (*Pan troglodytes*), the dominance hierarchy is relatively linear, and aggressive behaviors from dominant individuals to subordinate individuals are commonly observed in many dyads (Nishida and Hosaka 1996; Newton-Fisher 2004). The aggression among males seems to be related to male reproductive strategies because the presence of receptive females increases male aggression (Muller and Wrangham 2004). The aggressive behaviors are sometimes very violent and may result in the killing of adult males of the same group (Fawcett and Muhumuza 2000; Watts 2004; Kaburu et al. 2013; Wilson et al. 2014). On the other hand, some male members of a chimpanzee group form coalitions, which may help them monopolize mating opportunities against other male members (Watts 1998; Duffy et al. 2007; Mitani 2009; Schülke et al. 2010).

Bonobos (*Pan paniscus*) form male-philopatric groups that are very similar to those of chimpanzees (Kano 1982; Gerloff et al. 1999; Hashimoto et al. 2008; Ishizuka

et al. 2019). Therefore, we can expect to see similar behaviors in males, including intense aggression against other males in the group, an increase in the aggressive interactions in the presence of receptive females, coalitions among males for the monopolization of receptive females. However, these behaviors are seldom observed or are observed with less intense aggression among males within a group (Kano 1992; Furuichi and Ihobe 1994; Furuichi 1997; Surbeck et al. 2012; Samuni et al. 2022), and it raises the question of why bonobos and chimpanzees show substantial differences in their aggression and behavior patterns. Moreover, a significant question arises regarding the prevalence of reproductive skew toward high-ranking males in bonobos compared to chimpanzees, even though the intermale relationships in bonobos appear to be less despotic (Surbeck et al. 2017b; Ishizuka et al. 2018; Surbeck et al. 2019; Yokoyama and Furuichi 2022; Ishizuka 2023).

While answering this question requires observations of males from both species, studies focusing on behavioral interactions among male bonobos have been relatively scarce (Kuroda 1980; Ihobe 1992; Furuichi and Ihobe 1994; Surbeck et al. 2017a; Cheng et al. 2022). This scarcity is likely attributed to the less conspicuous nature of male bonobos behaviors compared to those of chimpanzees. Additionally, the heightened conspicuousness of female behavioral traits in bonobos may contribute to the focus on

the latter (Parish 1994; White and Wood 2007; Furuichi 2011, 2023). Furthermore, most

previous studies comparing the behavior of wild chimpanzees and bonobos did not rely

on data obtained through uniform observation methods. Therefore, to clarify whether

there are significant differences in within-group male relationships between

chimpanzees and bonobos, we conducted studies using the same observation methods

under natural conditions of male bonobos in Wamba, Democratic Republic of the Congo

and male chimpanzees in Kalinzu Forest Reserve, Uganda.

In contrast to chimpanzees, bonobos are not male dominant but rather show

male-female co dominance (Furuichi 1997; Stevens et al. 2007). The patterns of

association and behavior of male bonobos have been considered to be strongly affected

by prolonged sexual receptivity of females and the constant presence of multiple

receptive females in parties (Furuichi and Hashimoto 2002; Hashimoto et al. 2022),

moderation of mating competition by the presence of multiple receptive females (Kano

1992; Furuichi and Hashimoto 2002; Furuichi 2023), long-lasting mother–son

relationships (Furuichi 1989; Kano 1992; Furuichi 2011), and strong support from

dominant mothers in mating competition and acquisition of alpha status (Surbeck et al.

2010; Furuichi 2011; Ishizuka et al. 2018; Furuichi 2019; Surbeck et al. 2019; Shibata

and Furuichi 2023). Based on these previous reports, we hypothesized that in bonobos,

unlike chimpanzees, the prolonged sexual receptivity of females and the support from

high-ranking mothers strongly influence agonistic relationships among males so that the

importance of intermale aggressive interactions is relatively lower, and the frequency

and intensity of male aggression are reduced. In this study, we tested four predictions

derived from our hypothesis.

1.   Male bonobos are more likely to find a higher number of receptive females in

parties to which they attend than do male chimpanzees.

2.   In bonobos, instances of aggressive interactions among males are less frequent or

involve less physical contact than in chimpanzees.

3.   Aggressive interactions between males over access to receptive females are less

frequent in bonobos than in chimpanzees.

4.   In bonobos, aggressive interactions between males more commonly involve their

mothers as coalition partners compared to the patterns observed in chimpanzees.

Methods

Study sites and subjects

Bonobos in Wamba

Field assistants and author S.S. observed wild bonobos of the E1 group in

Wamba in the northern sector of the Luo Scientific Reserve in DR Congo, where long-

term research has been conducted since 1973 (Kano 1992; Furuichi et al. 2012). The

bonobos were observed for 37 days from September 19 to December 29, 2019. All the

individuals in the E1 group were identified and habituated before the beginning of this

study. At the time of this research, the group comprised approximately 40 individuals,

including eight adult males aged over 14 years, three adolescent males aged 8–14 years,

13 adult females, a few immigrant adolescent females, and many juveniles and infants.

The age classes of subject males were categorized according to Hashimoto (1997). The

study subjects were eight adult and three adolescent males. The individual names,

dominance rank, estimated birth year, and their mothers are shown in Table 1. The

dominance rank of males was calculated using David's score (David 1987) based on

dyadic aggressive interactions (see below for the definition of aggressive interactions).

The individuals were divided into one of the following four categories, namely the

alpha male (1), high ranking (2–3), middle ranking (4–7), and low ranking (8–11).

Chimpanzees in Kalinzu

Field assistants and author S.S. observed the wild chimpanzees of the M group

in the Kalinzu Forest Reserve, Uganda, where long-term research has been conducted

since 1992 (Hashimoto 1995; Hashimoto and Furuichi 2006). The chimpanzees were

observed for 112 days from February 3 to April 18, from June 23 to September 1, 2018,

and from March 11 to May 24 2019. During this study, the group consisted of

approximately 100 individuals, including 15 adult males aged over 15 years (Goodall

1986), 29 adult females, a few adolescent males and females, and many juveniles and

infants. All adult males of the M group were identified and habituated by the beginning

of the study period. The study subjects of the current study comprised 10 adult males.

The individual names, dominance ranks, and estimated birth years of each adult male

are shown in Table 2. The dominance rank of males was calculated using David's score

(David 1987) based on dyadic aggressive interactions. The individuals were divided

into one of the following four categories, namely the alpha male (1), high ranking (2–4),

middle ranking (5–7), and low ranking (8–10) (Table 2).

Data collection

Bonobos in Wamba were followed from approximately 0600 h from their

previous night's sleep site. Chimpanzees in Kalinzu were followed daily from

approximately 0700 h. We conducted observations using the same methods for both

bonobos and chimpanzees. We used the focal sampling method (Altmann 1974), in

which a focal animal was followed for as long as possible. We followed the first male

we found in the morning unless we had followed that individual on the previous day.

When we lost sight of the focal individual, we continued to observe if the individual

was found again within 30 min. However, if we could not find the individual for 30

min, we stopped the focal observation for that day.

Party composition and observation unit

Every 60 min during the focal following, we recorded the party composition that the

focal animal belonged to using the 1-h party method (Hashimoto et al. 2001; Mulavwa

et al. 2008). We recorded individuals within the visible range at the beginning of each

hour and included new individuals that joined the party until the end of the hour. Each

hour-long observation was considered a unit for analysis and referred to as one

observational hour unit (OHU) when focal animals were observed for > 30 min within

that hour. To exclude the potential for overestimating data on behaviors that were easily

observed, we excluded all observation data for days with < 3 OHUs (11 OHUs for

bonobos and 22 OHUs for chimpanzees). In total, 217 OHUs were analyzed for all the

subject bonobos with a mean of 19.7 ± 5.0 SD OHUs (range: 11–30) for each individual

(Table 1). The mean duration of OHU was 53.5 min (range: 30–60). For all the

chimpanzee subjects, 677 OHUs were analyzed with a mean of 67.7 ± 11.1 SD OHUs

(range: 53–90) for each individual (Table 2). The mean duration of OHU was 52.1 min

(range: 30–60).

Aggressive interactions

We recorded all the observed intragroup aggressive interactions in each OHU

using *ad libitum* sampling, including cases where the focal animal was not involved. We

defined "aggressive interactions" as involving at least one aggressive behavior toward

one or more individuals: aggressive physical contact such as kicking, beating, and

biting; chasing; charging; directed aggressive displaying (including branch dragging and

provocative behaviors for bonobos); non-vocal threatening. Undirected aggressive

displays, namely jumping in chimpanzees and branch dragging in bonobos, were only

recorded as aggressive interactions when other individuals responded by screaming or

fleeing after that behavior. Interactions involving only pant-grunts in chimpanzees were

not recorded as aggressive interactions because the frequency of such interactions was

very low in chimpanzees in Kalinzu, and bonobos did not exhibit corresponding

behaviors. Although we could not record some aggressive interactions due to bad

visibility in the bush or on the high canopy, all recorded aggressive interactions that

involved subject males were included in the analyses. The visibility was not

substantially different between the two study sites; therefore, it did not bias comparisons

between bonobos and chimpanzees in this study.

Presence of females with maximum swelling and comparison of party attendance by adult males

The swelling status of the sexual skin of each female in each party was

recorded as one of the following two categories: non-swelling and maximum swelling

(firmness 3 defined by (Furuichi 1987), hereinafter MS).

We defined all OHUs on days when we observed at least one female with MS

as "OHU under the presence of females with MS." In bonobos, females resume

receptivity in the early stage of postpartum infertility; therefore, usually multiple

females show MS and sexual receptivity (Kano 1992; Furuichi and Hashimoto 2002;

Hashimoto et al. 2022). In the current study, bonobo females with MS were observed in

35 of 37 (97%) observation days, and 197 of 217 (90.8%) OHUs were defined as "OHU

under the presence of females with MS." Therefore, we included all 217 OHUs in

bonobos as a single dataset in the analysis of the party attendance by adult males. In

chimpanzees, females with MS were observed in 62 of 112 (55.4%) observation days;

376 out of 677 (55.5%) OHUs were defined as "OHU under the presence of females

with MS," and the remaining 301 OHUs were defined as "OHU under the absence of

females with MS." We compared the number of adult males observed in a party in three

different group conditions, namely bonobos, chimpanzees under the presence of females

with MS, and chimpanzees under the absence of females with MS (Fig. 1). To

determine whether the datasets of the three conditions followed the same distribution,

we first normalized each dataset and then conducted the Kolmogorov–Smirnov two-

sample tests for three combinations by using the "ks.test()" function. We used R

(version 4.2.2; R Foundation for Statistical Computing, Vienna, Austria, http://www.r-

project.org [accessed on 13 December 2022]) for all statistical analyses.

Intragroup aggressive interactions among males

We recorded 92 aggressive interactions among male bonobos. Among these, 76

cases were dyadic aggressive interactions between males, comprising 43 cases in 12

dyads between adult males and 33 cases in 10 dyads between adult and adolescent

males. The remaining 16 cases were aggression toward males from multiple individuals

(including at least one male as an aggressor), comprising 14 cases toward adult males

and two cases toward adolescent males. For chimpanzees, we recorded 201 aggressive

interactions among adult males, including 177 dyadic aggressive interactions in 65

dyads and 24 polyadic interactions. Polyadic interactions include both cases of

coalitions and cases when there were multiple victims. We divided aggressive

interactions into two conditions for chimpanzees based on the presence/absence of

females with MS in the group. The aggressive interactions were divided into four

categories according to their intensity, namely aggressive physical contact, chasing,

directed aggressive display, and others including non-vocal threatening. All the

aggressive interactions among all the subject bonobos (including adolescents) were

summarized in the aggressive interaction matrix.

We calculated the frequencies of aggressive interactions in two different ways. First, we

calculated the frequency of overall aggressive interactions per OHU by dividing the

number of overall aggressive interactions by the number of OHUs. Secondly, we

calculated the frequencies of aggressive interactions per male in an OHU by dividing

the total number of males involved in aggressive interactions (including both aggressors

and receivers) by the total number of males observed during each OHU. To examine the

difference in the frequency of aggressive interactions between each group/condition, we

conducted a chi-square test with a Bonferroni-corrected *post hoc* test using the

"pairwise.prop.test()" function.

Results

Availability of receptive females and grouping patterns of adult males

A large difference between bonobos and chimpanzees in the availability of

receptive females with MS for males was observed. In bonobos, 1.22 ± 1.15 (range 0–4)

females with MS were observed in each OHU. On the other hand, in chimpanzees, 0.47

± 0.68 (range 0–3) females with MS were observed in each OHU.

Figure 1 shows the number of adult males observed in each OHU. In

chimpanzees, the number of males in parties differed considerably between days when

females with MS were observed (Fig. 1b) and days when they were not observed (Fig.

1c). In the absence of females with MS, the modal value of the number of males was

one, indicating that the males were highly dispersed. For bonobos, we drew only one

pooled figure because females with MS were absent only in two days (Fig. 1a). The

tendency of a high degree of dispersion was not found in bonobos. The party size

distribution differed significantly in all three pairs (Kolmogorov–Smirnov two sample

tests, chimpanzees under the absence of females with MS vs. chimpanzees under the

presence of them, D = 0.125, p = 0.011; bonobos vs. chimpanzees under the absence of

females with MS, D = 0.163, p = 0.003; bonobos vs. chimpanzees under the presence of

females with MS, D = 0.145, p = 0.006). However, the shape of the distribution for male

bonobos and the one for male chipanzees under the presence of females with MS were

similar with modal number of males of three to four (37.5–40.0% of adult male

members) for bonobos and six to seven (40.0–46.7%) for chimpanzees, suggesting that

both male bonobos and chimpanzees show similar tendency of party attendance when

females with MS are present in the group.

Aggressive interactions among male bonobos and chimpanzees

Table 3 shows the number of dyadic aggressive interactions expressed and

received in each dyad among 11 adult and adolescent male bonobos. We included those

involving adolescent males for bonobos because those aggressive interactions played

substantial roles in the dominance struggles between mother-son groups. Dyadic

aggressive interactions were observed in 22 out of 55 dyads. Of the 76 dyadic

aggressive interactions, 37 cases were directed at two middle and low-ranking males,

Jiro and Jo, and 30 out of the 37 were from one of the three high-ranking males (Kitaro,

Nobita, and Kiyota). In the 16 cases of polyadic aggressive interactions involving more

than two individuals (including 14 cases among adult males and two cases involving

adolescent males), the aggressee was a single male in all these cases and 15 cases were

directed at Jiro or Jo. Kitaro, Nobita, and Kiyota are the sons of the alpha female Kiku,

and Jiro and Jo are the sons of the second-ranking female Jacky (Table 1). During this

observation period, the sons of Jacky were challenging the high status of the sons of

Kiku by provoking them, and aggressive interactions among them frequently involved

their mothers and brothers (Shibata, personal observation).

Table 4 shows the numbers of dyadic aggressive interactions expressed and

received in each dyad among 10 adult male chimpanzees. Dyadic aggressive

interactions were observed in 32 out of 45 dyads. All the individuals except the alpha

male were attacked multiple times by higher-ranking males. Of the all 100 aggressive

interactions among adult males, 47 were expressed by the alpha male, Goku.

Comparison of the type and frequency of adult male aggressive interactions between

bonobos and chimpanzees

Table 5 shows the number of OHUs observed, the mean number of adult males

in the party, the total number of adult males observed in each OHU, and the number of

aggressive interactions among adult male bonobos of the E1 group and adult male

chimpanzees of the M group (note that this comparison did not include aggressive

interactions involving adolescent male bonobos).

Among bonobos, > 50% of aggressive interactions were dyadic directed

displays. Chasing behaviors were observed with a similar frequency, but more than half

of them were polyadic aggressive interactions (shown in parentheses in Table 5).

Aggressive physical contact among adult males was not observed during this study.

Among chimpanzees, chasing was the most frequent aggressive interaction and

accounted for approximately 60% of the total aggressive interactions. Aggressive

physical contact accounted for 15% of all aggressive interactions, and > 70% of these

occurred under the presence of females with MS.

In bonobos, all 14 observed polyadic aggressive interactions among adult

males involved chasing behaviors, with multiple individuals pursuing one individual,

and at least one adult male participating as an aggressor. Three or four individuals

joined as aggressors in eight of the 14 cases, and the aggressee of these interactions was

a single male, Jiro in 13 of 14 cases. Kitaro or/and Nobita joined in 12 of these 13 cases

of aggression toward Jiro. Females joined as aggressors in six of the 14 cases. Although

Kiku joined as an aggressor with Nobita in only one case, probably because she had a

newborn baby during this study period, Nao, an old female ally of Kiku, joined in three

cases to attack Jiro.

Out of all 24 polyadic aggressive interactions among chimpanzees, 10 cases

were coalitionary aggression comprising nine chasing behaviors and one directed

aggressive display from two males. All of these coalitionary aggressions occurred under

the presence of females with MS, except for one case of chasing behavior. Female

chimpanzees did not join any of the coalitionary aggression as aggressors. Chimpanzees

also showed 14 aggressive interactions from one male toward multiple individuals.

These interactions include six cases of chasing directed at multiple individuals and eight

cases of aggressive displays after which other multiple males responded by screaming

or fleeing. Such aggressive interactions from one individual to multiple individuals

were not observed in bonobos.

The frequency of overall aggressive interactions per OHU in bonobos (0.263)

was lower than that of chimpanzees (0.297) but the difference was not statistically

significant (chi-square test: $\chi^2 = 0.778$, df = 1, $p = 0.378$). When data on chimpanzees

was categorized into two groups based on the presence/absence of females with MS, the

frequency of aggressive interactions followed the order: chimpanzees under the

presence of females MS (0.404) > bonobos (0.263) > chimpanzees under the absence of

females with MS (0.163). These differences were statistically significant (chi-square

test: $\chi^2 = 48.41$, df = 2, $p < 0.001$). The results of the Bonferroni-corrected *post hoc* tests

showed that the differences were significant in all three pairs: bonobos vs. chimpanzees

with females with MS ($p < 0.01$), bonobos vs. chimpanzees without females with MS (p

< 0.001), chimpanzees with females with MS vs. chimpanzees without females with

MS ($p < 0.001$) (Table 6).

However, when the difference in the mean number of males in the party was

considered, these differences between the species/conditions changed. The frequency of

aggressive interactions per male in bonobos (0.162) was higher than that of

chimpanzees (0.112), and the difference between the species was statistically significant

(chi-square test: $\chi^2 = 4.821$, df = 1, $p = 0.028$). It seemed to be because a larger number

of males attended OHU in chimpanzees, but aggressive interactions specifically

occurred among some higher-ranking males, lowering the per-male frequency of

aggressive interactions in chimpanzees. When data on chimpanzees were categorized

based on the presence/absence of females with MS, the relationships in the frequencies

of aggressive interactions per male was in the order: bonobos (0.162) > chimpanzees

under the presence of females with MS (0.136) > chimpanzees under the absence of

females with MS (0.072), and the differences were statistically significant (chi-square

test: $\chi^2 = 20.471$, df = 2, $p < 0.001$). The results of the Bonferroni-corrected *post hoc*

tests showed that the differences were significant in bonobos vs. chimpanzees under the

absence of females with MS ($p < 0. 001$) and chimpanzees under the presence of

females with MS vs. chimpanzees under the absence of them ($p < 0.001$), but not in

bonobos vs. chimpanzees under the presence of females with MS ($p = 0.410$) (Table 6).

Discussion

In this study, we have focused on the difference in the frequency and intensity

of male aggression between chimpanzees and bonobos. We hypothesized that the

patterns of association and behaviors of male bonobos are strongly affected by the

prolonged sexual receptivity of females and the constant presence of multiple receptive

females (females with MS) in the group (Kano 1992; Furuichi and Hashimoto 2002;

Hashimoto et al. 2022; Furuichi 2023), and the long-lasting mother–son relationships

and strong support from mothers for mating competition and acquisition of the alpha

status (Furuichi 1989; Kano 1992; Surbeck et al. 2010; Furuichi 2011; Ishizuka et al.

2018; Furuichi 2019; Surbeck et al. 2019; Shibata and Furuichi 2023). We examined the

four predictions derived from this hypothesis.

The results of the current study support the first prediction that male bonobos

find a higher number of receptive females in their parties than male chimpanzees do.

Chimpanzee and bonobos males found 0.47 ± 0.68 and 1.22 ± 1.15 receptive females in

each OHU, respectively, probably because a large proportion of females in the group are

receptive showing MS due to the higher proportion of the receptive period of females in

bonobos, especially caused by the resumption of MS in the early stage of postpartum

amenorrhea (Furuichi and Hashimoto 2002; Hashimoto et al. 2022).

While at least one female with MS was found in 94.6% of all observation days

in bonobos, they were found in only 55.4 % of observation days in chimpanzees. On the

days when females with MS were not observed in the group, the modal value of the

number of male chimpanzees in parties was one, indicating that the males were highly

dispersed. In chimpanzees, low-ranking males were substantially more likely to range

alone or in parties without other males than high- and middle-ranking males in the

absence of receptive females, which might help them to reduce the probability of being

attacked by other males (Shibata et al. 2022). However, on the days when females with

MS were observed, nearly half of the male chimpanzees were observed in the same

party, similar to male bonobos in pooled data. In such days when females with MS were

observed in the group, no significant relationship between solitary ranging and male

rank was observed in chimpanzees, suggesting active participation by even low-ranking

males in the party (Shibata et al. 2022). Despite a significant difference in the

distribution of the number of adult males in the party between bonobos and

chimpanzees, both species seem to show a similar tendency to attend mixed-sex parties

when females with MS were present. This result underscores the significant impact of

the presence (in chimpanzees) or absence (in bonobos) of days without females with

MS on the social dynamics among males.

The analysis of aggressive interactions among adult males did not fully support the second prediction, which posited that the frequency of aggressive interactions is lower in bonobos than in chimpanzees (Tables 5 and 6). In particular, while the frequency of aggressive interactions per OHU showed no significant difference between bonobos and chimpanzees, the frequency of aggressive interactions per OHU per male was significantly higher in bonobos than in chimpanzees. When comparing the frequency of aggressive interactions in bonobos with those in chimpanzees under the presence of females with MS, the frequency per OHU was significantly lower in bonobos than in chimpanzees but there was no significant difference in the frequency per OHU per male (Tables 5 and 6). In the days when females with MS were not observed in chimpanzees, both the frequency of aggressive interactions per OHU and per OHU per male were significantly lower than those in pooled data of bonobos and those in chimpanzees under the presence of females with MS (Tables 5 and 6), probably because aggressive interactions did not occur over access to MS females, and low-ranking males frequently range alone or in parties without other males to avoid aggression of higher-ranking males under such circumstances (Shibata et al. 2022).

Conversely, prediction 2 was supported when examining the type of aggression. In bonobo males, aggressive interactions were primarily characterized by chasing or

aggressive displays, and notably, unlike in chimpanzees, no instance of aggressive

physical contact was observed during this study. This may explain why lower-ranking

bonobo males do not need to leave the party to avoid attacks by dominant males.

Although a single male was targeted in all polyadic aggressive interactions in bonobos,

one male targeted multiple males in 14 of 24 cases of polyadic aggressive interactions in

chimpanzees. These interactions occurred while other males were resting or grooming

in proximity, and the alpha male was the aggressor in nine of these cases. This behavior

aligns with what has been previously termed as "separating intervention," in other

studies, wherein alpha males prevent other males from forming affiliative relations and

coalitions (de Waal 1982; Nishida and Hosaka 1996). This underscores the intricacies of

male competition among chimpanzees and the substantial impact of aggressive

behaviors from dominant individuals.

The results of this study support the third prediction, indicating that aggressive

interactions between males over access to receptive females are less frequent in bonobos

than in chimpanzees. Previous research on chimpanzees has documented an increase in

male aggression in the presence of receptive females (Muller and Wrangham 2004).

Similarly, in the current study, we observed a rise in the frequency of more severe

aggressive interactions, such as aggressive physical contact and chasing behavior,

among chimpanzees when females with MS were present. Notably, nine of 10 cases of

coalitionary aggressive interactions in chimpanzees occurred under the presence of

females with MS. These results suggest that aggressive interactions among male

chimpanzees intensify under the presence of receptive females, and males strategically

form coalitionary attacks for mating competition. In contrast, a previous study at

Wamba reported a relatively low frequency of aggressive behavior that interferes with

the mating of subordinates in bonobos (27 of 515 cases of copulations (Kano 1996); 2

of 36 cases of agonistic interactions (Furuichi 1997)). Similarly, in this study, out of 73

cases of aggressive interactions in clear contexts, only seven cases were observed over

access to females with MS (Shibata, unpublished data).

The findings of the current study also corroborate the fourth prediction,

indicating that aggressive interactions between males involve their mothers more

frequently in bonobos than in chimpanzees. In chimpanzees, aggressive interactions

occurred in 32 out of 45 dyads of adult males, with all males except the alpha male

being aggressed multiple times by higher-ranking males. However, females did not

participate in any of the aggressive interactions among males. On the contrary, in

bonobos, dyadic aggressive interactions were observed in only 22 out of 55 dyads of

adult and adolescent males. Notably, dyadic aggression was directed toward two

specific males in as many as 37 out of 76 cases. During this study period, the two sons

of the second-ranking female Jacky (Jo and Jiro) frequently challenged the sons of Kiku

(Kitaro, Nobita, and Kiyota) and faced counter-attacks. Jiro, in particular, was targeted

in 13 out of 14 polyadic aggressive interactions, and Kitaro or/and Nobita joined in 12

of these 13 cases of aggression toward Jiro, highlighting the intensity of competition for

the alpha male position. These observations suggest that a significant proportion of

aggressive interactions among bonobo males involved competition for the alpha male

position. In all 45 cases of aggressive interactions between the sons of Kiku and the

sons of Jacky, Kiku or her female ally, Nao, joined in four cases to attack Jiro. In the PE

group neighboring the E1 group, the mother of one of the males participated in

approximately 6% of aggressive interactions between males (Tokuyama, personal

communication). Another study at LuiKotale reported that mothers joined

approximately 22% of aggressive interactions over access to receptive females (Surbeck

et al. 2010). The current study, along with these preceding reports, indicates that bonobo

females participate in aggressive interactions more frequently than female chimpanzees

do, intervening in mating competitions or competitions for alpha-male status involving

their sons.

It should be noted that although a part of the second prediction was not

supported due to the equal or even higher frequency of aggressive interactions among

male bonobos compared to male chimpanzees, the frequency of aggressive interactions

among male bonobos substantially decreases when interactions among these specific

males are excluded.

In conclusion, the scrutiny of the four predictions aligns with our hypothesis

that the prolonged sexual receptivity of females exert a significant influence on male

agonistic relationships in bonobos. As documented in previous studies, the operational

sex ratio (proportion of adult males to sexually receptive females) in bonobos is notably

reduced due to the early resumption of sexual swelling and receptivity during the

infertile nursing period (Kano 1992; Furuichi and Hashimoto 2002; Hashimoto et al.

2022; Furuichi 2023). This study reveals that this reduction increases the presence of

females with MS in the group, along with the number of such females that male

bonobos encounter in the parties they attend. This heightened opportunity for mating

appears to contribute to less frequent aggressive interactions, particularly those related

to access to receptive females, and a decrease in the severity of aggressive behaviors

among males.

Additionally, our findings highlight that a significant portion of aggressive

interactions among male bonobos revolves around struggles for the alpha-male position

and that female bonobos sometimes attend to aggressive interactions to support their

sons. The alpha female, Kiku, was observed to participate in these interactions only

once during this study, likely influenced by her caring for a newborn infant. However,

Kiku's female ally, Nao, consistently joined in aggressive interactions to support Kiku's

sons. Examining the historical alpha status dynamics in the E1 group over nearly the

past 40 years (1983–2020) emphasizes the substantial impact of the mother's status

among females and their intervention in struggles for alpha status among male bonobos

(Furuichi 1997, 2011; Shibata and Furuichi 2023). This seems to be a reason why

relatively young males (late adolescent or young adult) with mothers in their prime age

tend to acquire the alpha male status in bonobos (Furuichi 1997, 2019; Shibata and

Furuichi 2023)

In contrast, in chimpanzees, age emerges as a pivotal factor in acquiring alpha

status among males. Hasegawa and Kutsukake (2015), based on data from three long-

term study sites, reported that male chimpanzees typically reach their highest potential

to become alpha at an age of approximately 25 years, except in cases where the alpha

status is achieved through coalition formation. Eastern chimpanzees in Mahale acquire

alpha status at the age of 19 to 37 years (Nishida 2008). In Gombe, middle-aged males

reach their highest rank, dominating most, if not all, younger and older males. While

cases have been reported of males with supportive elder brothers becoming alpha, and

mothers occasionally support their offspring in fights, the influence of such maternal

support on the rank of adult males remains unclear (Bygott 1979).

The variations in the impact of maternal support observed between the two

species may stem from differences in mother–son relationships. In chimpanzees, males

significantly decrease their association with mothers when mothers resume cyclic

receptivity, eventually severing their relationship to associate with male adults (Pusey

1983; Hayaki 1988). Conversely, male bonobos maintain close associations with their

mothers into adulthood, remaining in the same party, providing ample opportunities for

maternal support (Furuichi 1989; Kano 1992; Surbeck et al. 2010; Schubert et al. 2013;

Hashimoto and Furuichi 2015). However, it remains unclear whether the prolonged

mother–son associations enhance the influence of mothers on intermale competition or

if the evolution of female behaviors to support their sons leads to such enduring

mother–son associations.

As highlighted in the Introduction, a question in male reproductive competition

is the more pronounced reproductive skew toward high-ranking males in bonobos

compared to chimpanzees, despite seemingly less despotic intermale relationships

(Surbeck et al. 2017b; Ishizuka et al. 2018; Surbeck et al. 2019). Particularly in the E1

group, 10 out of 11 infants with confirmed paternity were sired by alpha males at that

time, with nine cases by Nobita and one by Kitaro, both sons of the alpha female Kiku

(Ishizuka et al. 2018; Ishizuka 2023). As explained above, one of the reasons for the

high reproductive skew in bonobos might be the direct support of high-ranking mothers

in the mating competition to support their high-ranking sons. Although aggressive

interactions among males do not occur frequently, the presence of high-ranking mothers

or their aggressive intervention may suppress the mating behaviors of their sons' rivals

(Surbeck et al. 2010; Furuichi 2023). Another reason might be the characteristic spatial

distribution of bonobos within the mixed-sex party. Unlike female chimpanzees, female

bonobos gather and stay at the center of the party, and males tend to remain peripheral

(Kuroda 1979; Furuichi 1989; Hashimoto and Furuichi 2015). However, due to the

strong mother-son association, sons of high-ranking females tend to stay in the

gathering of females with their mothers and have more opportunities or easier access to

receptive females even if they do not suppress mating behaviors of lower-ranking males

by aggressive interactions (Surbeck et al. 2012; Yokoyama and Furuichi 2022; Furuichi

2023). Although these possibilities are yet to be confirmed in future studies, such high

influences of support by high-ranking females seem to diminish the significance of

direct aggressive interactions among males and reduce the severity of aggressive

behaviors. Additionally, female behaviors for supporting their sons might have evolved

in bonobos for their own reproductive success, aiming to increase the number of their

grandchildren (Furuichi 1997; Surbeck et al. 2019; Furuichi 2023).

The present study faces a limitation stemming from the truncated observation

period for bonobos, a consequence of the COVID-19 pandemic. In contrast to the

comprehensive eight-month study duration for chimpanzees covering a majority of the

year, the investigation into bonobos spanned a condensed four-month interval,

specifically from September to December. This temporal discrepancy raises concerns,

particularly if male behavioral patterns are influenced by seasonality, particularly in

relation to fruit abundance. Studies on a neighboring group at Wamba, PE, indicated

that intergroup encounters were most frequent during the early peak of fruit abundance,

coinciding with a reduction in aggression between males within the group (Sakamaki et

al. 2018; Tokuyama et al. 2019). Conversely, a study at Kokolopori, a nearby site to

Wamba, reported an increase in aggression and cortisol levels in males with rising fruit

abundance (Cheng et al. 2021). Fortunately, the four-month period of our study aligned

with the transition from the highest to the lowest fruiting season at Wamba (Mulavwa et

al. 2008); therefore, the current results may not significantly biased by the seasonal

change in fruit abundance. However, through studies with with extended periods, we

must assess the influences of seasonality in the abundance and distribution of food

resources on the type and frequency of intermale aggressive interactions.

It is crucial also to note that the behavioral tendencies observed in this study

cannot be universally generalized across populations of bonobos and chimpanzees.

Intermale relationships are likely influenced by various factors beyond fruiting

seasonality, including group size (Mitani et al. 2000; Watts 2018), the personalities of

high-ranking males and females (Goodall 1986; Takahata 1990; Shibata and Furuichi

2023), and differences between chimpanzee subspecies (Boesch 1996). Notably,

intermale relationships in bonobos at Wamba vary across years depending on the

presence or absence of competing mother–son pairs and their respective personalities.

Hence, additional studies involving multiple groups of bonobos and chimpanzees across

various years are essential for a comprehensive understanding of behavioral tendencies

common to each species and the variations in intermale relationships within male-

philopatric societies of the *Pan* species.

Acknowledgments

We thank the National Forestry Authority of Uganda, the Uganda National

Council for Science and Technology, and the Ministry of Scientific Research of the

Democratic Republic of the Congo for their permission to conduct this research. We

thank the local assistants for their help with field observations. We thank Nahoko

Tokuyama, Tetsuya Sakamaki, Kazuya Toda, Chie Hashimoto, Hiroyuki Takemoto,

Daisuke Shimizu, Akito Toge, and Shintaro Ishizuka for their help during the field

research and their valuable discussions. This research was financially supported by the

Leading Program in Primatology and Wildlife Science of the Kyoto University, the

Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for JSPS Fellows, and

JSPS Grants-in-Aid for Scientific Research (to TF, 16H02753, 25304019, and

26257408). We also appreciate the anonymous reviewers for their helpful comments on

this paper.

Author contributions

SS designed this study, collected and analyzed behavioral data, and wrote and

revised the manuscript. TF supervised SS in designing this study. Both authors revised

the manuscript and gave the final approval for publication.

Ethical approval

This study was approved by the National Forestry Authority of Uganda and the

Uganda National Council for Science and Technology.

References

Altmann J (1974) Observational study of behavior: sampling methods. Behaviour 49:227-266.
https://doi.org/DOI:10.1163/156853974X00534
Boesch C (1996) Social grouping in Tai chimpanzees. In: McGrew WC, Marchant LF, Nishida T
(eds) Great Ape Societies. Cambridge Univ Press, Cambridge, pp 101-113
Bygott JD (1979) Agonistic behavior, dominance, and social structure in wild chimpanzees of
the Gombe National Park. In: Hamburg DA, McCown ER (eds) The great apes.
Benjamin/Cummings, California, pp 405-428
Cheng L, Lucchesi S, Mundry R, Samuni L, Deschner T, Surbeck M (2021) Variation in
aggression rates and urinary cortisol levels indicates intergroup competition in wild
bonobos. Horm Behav 128:104914. https://doi.org/10.1016/j.yhbeh.2020.104914
Cheng L, Samuni L, Lucchesi S, Deschner T, Surbeck M (2022) Love thy neighbour:
behavioural and endocrine correlates of male strategies during intergroup encounters in
bonobos. Anim Behav 187:319-330. https://doi.org/10.1016/j.anbehav.2022.02.014
David HA (1987) Ranking from unbalanced paired-comparison data. Biometrika 74:432-436.
https://doi.org/10.1093/biomet/74.2.432
de Waal FBM (1982) Chimpanzee politics: power and sex among apes. Jonathan Cape,London
Duffy KG, Wrangham RW, Silk JB (2007) Male chimpanzees exchange political support for
mating opportunities. Curr Biol 17:R586-R587.
https://doi.org/10.1016/j.cub.2007.06.001

Fawcett K, Muhumuza G (2000) Death of a wild chimpanzee community member: possible
608   outcome of intense sexual competition. American Journal of primatology 51:243-247.
609   https://doi.org/10.1002/1098-2345(200008)51:4<243::AID-AJP3>3.0.CO;2-P

610   Furuichi T (1987) Sexual swelling receptivity and grouping of wild pygmy chimpanzee females
611   at Wamba, Zaire. Primates 28:309-318. https://doi.org/doi.org/10.1007/BF02381014

612   Furuichi T (1989) Social interactions and the life history of female *Pan paniscus* in Wamba,
613   Zaire. Int J Primatol 10:173-197. https://doi.org/10.1007/BF02735199

614   Furuichi T (1997) Agonistic interactions and matrifocal dominance rank of wild bonobos (*Pan*
615   *paniscus*) at Wamba. Int J Primatol 18:855-875.
616   https://doi.org/10.1023/A:1026327627943

617   Furuichi T (2011) Female contributions to the peaceful nature of bonobo society. Evol Anthropol
618   20:131-142. https://doi.org/10.1002/Evan.20308

619   Furuichi T (2019) Bonobo and chimpanzee: the lessons of social coexistence. Primatology
620   Monographs. Springer

621   Furuichi T (2023) Prolonged sexual receptivity in females and its impact on the evolution of
622   bonobos. In: Furuichi T, Idani G, Kimura D, Ihobe H, Hashimoto C (eds) Bonobos and
623   people at Wamba: 50 years of research. Springer, Singapore, pp 545-564

624   Furuichi T, Hashimoto C (2002) Why female bonobos have a lower copulation rate during estrus
625   than chimpanzees. In: Boesch C, Hohmann G, Marchant LF (eds) Behavioural diversity
626   in chimpanzees and bonobos. Cambridge University Press, New York, pp 156-167

627   Furuichi T, Idani G, Ihobe H, Hashimoto C, Tashiro Y, Sakamaki T, Mulavwa MN, Yangozene K,
628   Kuroda S (2012) Long-term studies on wild bonobos at Wamba, Luo Scientific Reserve,
629   D. R. Congo: towards the understanding of female life history in a male-philopatric
630   species. In: Kappeler PM, Watts DP (eds) Long-term field studies of primates. Springer-
631   Verlag, Berlin, Heidelberg, pp 413-433

632   Furuichi T, Ihobe H (1994) Variation in male relationships in bonobos and chimpanzees.
633   Behaviour 130:211-228. https://doi.org/10.1163/156853994X00532

634   Gerloff U, Hartung B, Fruth B, Hohmann G, Tautz D (1999) Intracommunity relationships,
635   dispersal pattern and paternity success in a wild living community of bonobos (Pan
636   paniscus) determined from DNA analysis of faecal samples. Proc R Soc Lond B
637   266:1189-1195. https://doi.org/10.1098/rspb.1999.0762

638   Goodall J (1986) The chimpanzees of Gombe: patterns of behavior. Harvard University
639   Press,Cambridge

640   Hasegawa M, Kutsukake N (2015) Bayesian competitiveness estimation predicts dominance
641   turnover among wild male chimpanzees. Behav Ecol Sociobiol 69:89-99.
642   https://doi.org/10.1007/s00265-014-1821-9

Hashimoto C (1995) Population census of the chimpanzees in the Kalinzu Forest, Uganda: comparison between methods with nest counts. Primates 36:477-488. https://doi.org/10.1007/BF02382870

Hashimoto C (1997) Context and development of sexual behavior of wild bonobos (*Pan paniscus*) at Wamba, Zaire. Int J Primatol 18:1-21. https://doi.org/10.1023/A:1026384922066

Hashimoto C, Furuichi T (2006) Comparison of behavioral sequence of copulation between chimpanzees and bonobos. Primates 47:51-55. https://doi.org/https://doi.org/10.1007/s10329-005-0144-x

Hashimoto C, Furuichi T (2015) Sex differences in ranging and association patterns in chimpanzees in comparison with bonobos. In: Furuichi T, Yamagiwa J, Aureli F (eds) Dispersing primate females: life history and social strategies in male-philopatric species. Springer Japan, Tokyo, pp 105-126

Hashimoto C, Furuichi T, Tashiro Y (2001) What factors affect the size of chimpanzee parties in the Kalinzu Forest, Uganda? examination of fruit abundance and number of estrous females. Int J Primatol 22:947-959. https://doi.org/10.1023/A:1012061504420

Hashimoto C, Ryu H, Mouri K, Shimizu K, Sakamaki T, Furuichi T (2022) Physical, behavioral, and hormonal changes in the resumption of sexual receptivity during postpartum infertility in female bonobos at Wamba. Primates 63:109-121. https://doi.org/10.1007/s10329-021-00968-w

Hashimoto C, Tashiro Y, Hibino E, Mulavwa M, Yangozene K, Furuichi T, Idani G, Takenaka O (2008) Longitudinal structure of a unit-group of bonobos: Male philopatry and possible fusion of unit-groups. In: Furuichi T, Thompson J (eds) The bonobos: behavior, ecology and conservation. Springer, New York, pp 107-119

Hayaki H (1988) Association partners of young chimpanzees in the Mahale Mountains National Park Tanzania. Primates 29:147-161. https://doi.org/10.1007/BF02381119

Ihobe H (1992) Male-male relationships among wild bonobos (*Pan paniscus*) at Wamba, Republic of Zaire. Primates 33:163-179. https://doi.org/10.1007/BF02382747

Ishizuka S (2023) Within-and between-group kin structure of wild bonobos in comparison to other African great apes. In: Furuichi T, Idani G, Kimura D, Ihobe H, Hashimoto C (eds) Bonobos and people at Wamba: 50 years of research. Springer, Singapore, pp 93-111

Ishizuka S, Kawamoto Y, Sakamaki T, Tokuyama N, Toda K, Okamura H, Furuichi T (2018) Paternity and kin structure among neighbouring groups in wild bonobos at Wamba. Roy Soc Open Sci 5:171006. https://doi.org/10.6084/m9.figshare.c.3980784

Ishizuka S, Toda K, Furuichi T (2019) Genetic analysis of migration pattern of female bonobos
(Pan paniscus) among three neighboring groups. International Journal of Primatology:1-
14

Kaburu S, Inoue S, Newton-Fisher N (2013) Death of the alpha: within-community lethal
violence among chimpanzees of the Mahale Mountains National Park. American
Journal of Primatology 75:789-797. https://doi.org/10.1002/ajp.22135

Kano T (1982) The social group of pygmy chimpanzees (*Pan paniscus*) of Wamba. Primates
23:171-188. https://doi.org/10.1007/BF02381159

Kano T (1992) The Last Ape: Pygmy Chimpanzee Behavior and Ecology. Stanford University
Press,Stanford, California

Kano T (1996) Male rank order and copulation rate in a unit-group of bonobos at Wamba, Zaire.
In: McGrew WC, Marchant LF, Nishida T (eds) Great ape societies. Cambridge Univ
Press, Cambridge, pp 135-145

Kuroda S (1979) Grouping of the pygmy chimpanzees. Primates 20:161-183.
https://doi.org/10.1007/BF02373371

Kuroda S (1980) Social behavior of the pygmy chimpanzees. Primates 21:181-197.
https://doi.org/10.1007/BF02374032

Mitani J (2009) Cooperation and competition in chimpanzees: current understanding and future
challenges. Evol Anthropol 18:215-227. https://doi.org/10.1002/evan.20229

Mitani JC, Merriwether DA, Zhang CB (2000) Male affiliation, cooperation and kinship in wild
chimpanzees. Animal Behaviour 59:885-893

Mulavwa M, Furuichi T, Yangozene K, Yamba-Yamba M, Motema-Salo B, Idani G, Ihobe H,
Hashimoto C, Tashiro Y, Mwanza N (2008) Seasonal changes in fruit production and
party size of bonobos at Wamba. In: Furuichi T, Thompson J (eds) The bonobos:
behavior, ecology, and conservation. Springer, New York, pp 121-134

Muller MN, Wrangham RW (2004) Dominance, aggression and testosterone in wild
chimpanzees: a test of the ' challenge hypothesis '. Animal Behaviour 67:113-123.
https://doi.org/10.1016/j.anbehav.2003.03.013

Nishida T (2008) Chimpanzee no sekai (the society of chimpanzees) (in Japanese). Toho
Shuppan,Osaka, Japan

Nishida T, Hosaka K (1996) Coalition strategies among adult male chimpanzees of the Mahale
Mountains, Tanzania. In: McGrew W, Marchant L, Nishida T (eds) Great ape societies.
Cambridge University Press, Cambridge, pp 114-134

Parish AR (1994) Sex and food control in the uncommon chimpanzee: how bonobo females
overcome a phylogenetic legacy of male dominance. Ethol Sociobiol 15:157-179.
https://doi.org/10.1016/0162-3095(94)90038-8

Pusey AE (1983) Mother-offspring relationships in chimpanzees after weaning. Anim Behav 31:363-377. https://doi.org/10.1016/S0003-3472(83)80055-4

Sakamaki T, Ryu H, Toda K, Tokuyama N, Furuichi T (2018) Increased frequency of intergroup encounters in wild bonobos (*Pan paniscus*) around the yearly peak in fruit abundance at Wamba. Int J Primatol 39:685-704. https://doi.org/10.1007/s10764-018-0058-2

Samuni L, Langergraber KE, Surbeck MH (2022) Characterization of Pan social systems reveals in-group/out-group distinction and out-group tolerance in bonobos. Proceedings of the National Academy of Sciences 119:e2201122119

Schubert G, Vigilant L, Boesch C, Klenke R, Langergraber K, Mundry R, Surbeck M, Hohmann G (2013) Co–residence between males and their mothers and grandmothers is more frequent in bonobos than chimpanzees. PLoS One 8:e83870. https://doi.org/10.1371/journal.pone.0083870

Schülke O, Bhagavatula J, Vigilant L, Ostner J (2010) Social bonds enhance reproductive success in male macaques. Curr Biol 20:2207-2210. https://doi.org/10.1016/j.cub.2010.10.058

Shibata S, Furuichi T (2023) Intermale relationships in wild bonobos at Wamba. In: Furuichi T, Idani G, Kimura D, Ihobe H, Hashimoto C (eds) Bonobos and people at Wamba: 50 years of research. Springer, Singapore, pp 285-300

Shibata S, Furuichi T, Hashimoto C (2022) Adaptive Party Choice of Low-Ranking Males in Fission–Fusion Dynamics of Chimpanzees in Kalinzu Forest Reserve, Uganda. Animals 12:2240. https://doi.org/10.3390/ani12172240

Stevens JMG, Vervaecke H, de Vries H, van Elsacker L (2007) Sex differences in the steepness of dominance hierarchies in captive bonobo groups. Int J Primatol 28:1417-1430. https://doi.org/10.1007/s10764-007-9186-9

Surbeck M, Beschner T, Shubert G, Weltring A, Hohmann G (2012) Mate competition, testosterone and intersexual relationships in bonobos, *Pan paniscus*. Anim Behav 83:659-669. https://doi.org/10.1016/j.anbehav.2011.12.010

Surbeck M, Boesch C, Girard-Buttoz C, Crockford C, Hohmann G, Wittig RM (2017a) Comparison of male conflict behavior in chimpanzees (*Pan troglodytes*) and bonobos (*Pan paniscus*), with specific regard to coalition and post-conflict behavior. Am J Primatol 79:e22641. https://doi.org/10.1002/ajp.22641

Surbeck M, Boesch C, Thompson ME, Furuichi T, Fruth B, Hohmann G, Ishizuka S, Machanda Z, Muller MN, Pusey A, Sakamaki T, Tokuyama N, K. W, Wrangham R, Wroblewski E, Zuberbuhler K, Vigilant L, langergraber K (2019) Males with a mother living in their group have higher paternity success in bonobos but not chimpanzees. Curr Biol 29:R341-R357. https://doi.org/10.1016/j.cub.2019.03.040

Surbeck M, Langergraber KE, Fruth B, Vigilant L, Hohmann G (2017b) Male reproductive skew
is higher in bonobos than chimpanzees. Current Biology 27:R640-R641.
https://doi.org/https://doi.org/10.1016/j.cub.2017.05.039

Surbeck M, Mundry R, Hohmann G (2010) Mothers matter! Maternal support, dominance status
and mating success in male bonobos (*Pan paniscus*). Proc R Soc B 278:590-598.
https://doi.org/10.1098/rspb.2010.1572

Takahata Y (1990) Social relationships among adult males. In: Nishida T (ed) The chimpanzees
of the Mahale Mountains: sexual and life history strategies. University of Tokyo Press,
Tokyo, pp 149-170

Tokuyama N, Sakamaki T, Furuichi T (2019) Inter-group aggressive interaction patterns indicate
male mate defense and female cooperation across bonobo groups at Wamba,
Democratic Republic of the Congo. Am J Phys Anthropol 170:535-550.
https://doi.org/10.1002/ajpa.23929

Watts DP (1998) Coalitionary mate guarding by male chimpanzees at Ngogo, Kibale National
Park, Uganda. Behav Ecol Sociobiol 44:43-55. https://doi.org/10.1007/s002650050513

Watts DP (2004) Intracommunity coalitionary killing of an adult male chimpanzee at Ngogo,
Kibale National Park, Uganda. Int J Primatol 25:507-521.
https://doi.org/10.1023/B:IJOP.0000023573.56625.59

Watts DP (2018) Male dominance relationships in an extremely large chimpanzee community at
Ngogo, Kibale National Park, Uganda. Behaviour 155:969-1009

White FJ, Wood KD (2007) Female feeding priority in bonobos, *Pan paniscus*, and the question
of female dominance. Am J Primatol 69:837-850. https://doi.org/10.1002/ajp.20387

Wilson ML, Boesch C, Fruth B, Furuichi T, Gilby IC, Hashimoto C, Hobaiter CL, Hohmann G,
Itoh N, Koops K, Lloyd JN, Matsuzawa T, Mitani JC, Mjungu DC, Morgan D, Muller MN,
Mundry R, Nakamura M, Pruetz J, Pusey AE, Riedel J, Sanz C, Schel AM, Simmons N,
Waller M, Watts DP, White F, Wittig RM, Zuberbuhler K, Wrangham RW (2014) Lethal
aggression in *Pan* is better explained by adaptive strategies than human impacts.
Nature 513:414-417. https://doi.org/10.1038/nature13727

Yokoyama T, Furuichi T (2022) Why bonobos show a high reproductive skew towards high-
ranking males: analyses for association and mating patterns concerning female sexual
states. Primates 63:483-494. https://doi.org/10.1007/s10329-022-01004-1

Figure 1. The number of OHUs in which the party included the respective number of adult males.

a) Adult male bonobos (N = 217). b) Adult male chimpanzees under the presence of females with

MS (N = 376). c) Adult male chimpanzees under the absence of females with MS (N = 301).

Table 1. Subject males, dominance rank, birth year, and the number of observation hour units (OHU)

of bonobos of the E1 group.

Name (abbreviation)	Dominance rank	Birth year	Mother	OHU
Kitaro (KT)	Alpha	2004	Kiku	30
Nobita (NB)	High	1988	Kiku	18
Kiyota* (KY)	High	2009	Kiku	20
Loboko (LB)	Middle	1994**	-	11
Jiro (JR)	Middle	2002	Jackey	24
Dai (DI)	Middle	1975	-	23
Ten (TN)	Middle	1974	-	20
Tawashi (TW)	Low	1970	-	16
Gauche (GC)	Low	1988**	-	18
Hachiro* (HC)	Low	2009	Hoshi	16
Jo* (JO)	Low	2007	Jackey	21

* Adolescent male

** Year estimated

Table 2. Subject males, dominance rank, birth year, and the number of observation hour units (OHU)

of chimpanzees of the M group.

Name (abbreviation)	Dominance rank	Birth year	OHU
Goku (GK)	Alpha	1993*	90
Ponta (PO)	High	1995*	53
Ichiro (IC)	High	1980s*	76
Buru (BR)	High	1970s*	65
Prince (PR)	Middle	1997*	59
Taiki (TK)	Middle	1999	77
Deo (DO)	Middle	1970s*	62
Pietan (PT)	Low	2001	65
Black (BL)	Low	1998*	72
Jo (JO)	Low	2000*	58

* Year estimated

Table 3. Aggressive interaction matrix with numbers of aggression expressed and received among 11 adult and adolescent male bonobos of the E1 group.

		Receiver									
		KT	NB	KY*	LB	JR	DI	TW	TN	GC	HC*	JO*
Aggressor	KT	-	2	0	0	5	1	0	0	3	0	5
	NB	0	-	0	0	7	0	0	0	5	1	6
	KY*	0	0	-	1	1	0	0	0	0	0	6
	LB	0	0	0	-	0	3	0	0	0	0	0
	JR	0	4	0	0	-	1	0	0	4	0	0
	DI	0	0	0	0	0	-	0	4	1	0	1
	TW	0	0	0	0	0	0	-	0	0	0	0
	TN	0	0	0	0	0	0	0	-	0	1	0
	GC	0	0	0	0	1	1	1	0	-	0	5
	HC*	0	0	0	0	0	0	0	1	0	-	0
	JO*	0	0	1	0	0	0	0	0	2	2	-

* Adolescent male

Table 4. Aggressive interaction matrix with numbers of aggression expressed and received among 10

adult chimpanzees of the M group.

		GK	PO	BR	IC	PR	TK	DO	PT	BL	JO
						Receiver					
Aggressor	GK	-	3	4	9	3	6	5	6	2	9
	PO	0	-	0	0	5	2	0	3	2	2
	BR	0	0	-	0	3	0	0	1	1	0
	IC	0	0	0	-	5	2	1	1	0	2
	PR	0	0	1	2	-	1	2	0	0	0
	TK	0	0	0	0	1	-	3	0	1	1
	DO	0	0	0	2	1	0	-	0	0	0
	PT	0	0	0	1	0	0	1	-	0	0
	BL	0	1	0	1	2	0	0	1	-	1
	JO	0	0	0	0	0	0	0	0	0	-

Table 5. Type and frequency of aggressive interactions among adult males compared between bonobos of the E1 group and chimpanzees of the M group

| Groups | Total number of OHUs observed | Mean number of males in the party | Total number of males observed | Number of male-male aggressive interactions* | | | | | Total number of males involved in aggressive interactions | Frequency of aggressive interactions per OHU | Frequency of aggressive interaction per OHU per male |
				Physical contact	Chasing	Directed display	Others	Total			
Bonobos	217	3.60 ± 1.80	782	0 (0)	25 (14)	31 (0)	1 (0)	57 (14)	127	0.263	0.162
Chimpanzees	677	5.67 ± 3.09	3840	30 (0)	126 (15)	42 (9)	3 (0)	201 (24)	429	0.297	0.112
Chimpanzees (under the presence of females with MS)	376	6.35 ± 2.98	2389	23 (0)	99 (13)	29 (6)	1 (0)	152 (19)	325	0.404	0.136
Chimpanzees (under the absence of females with MS)	301	4.82 ± 3.03	1451	7 (0)	27 (2)	13 (3)	2 (0)	49 (5)	104	0.163	0.072

Numbers in parentheses represent the number of polyadic aggressive interactions.

* The 35 agonistic interactions involving adolescent males in bonobos are not included in this table.

Table 6. The results of the Bonferroni-corrected post hoc tests.

Group comparison	Adjusted p-value	
	Aggression per OHU	Aggression per OHU per male
Bonobos vs. Chimpanzees	NS	<0.05
Bonobos vs. Chimpanzees (under the presence of females with MS)	<0.01	NS
Bonobos vs. Chimpanzees (under the absence of females with MS)	<0.001	<0.001
Chimpanzees (under the presence of females with MS) vs. Chimpanzees (under the absence of females with MS)	<0.001	<0.001