Title page

Authors

Shohei Shibata[1], Takeshi Furuichi[1]

Title

Intragroup intermale relationships in wild bonobos (*Pan paniscus*) at Wamba,

Democratic Republic of Congo, compared to those of chimpanzees (*Pan troglodytes*) at

Kalinzu Forest Reserve, Uganda

Affiliations and Address

[1] Wildlife Research Center, Kyoto University, Inuyama, Aichi 484-8506, Japan

Correspondence

Name: Shohei Shibata

ORCID: 0000-0002-3605-2502

Address: Inuyama campus, Kyoto University, Kanrin 41, Inuyama, Aichi 484-8506,

Japan

Email: shibata.shohei.d12@kyoto-u.jp

Phone Number: +81 (0)568 63 0537

Abstract

Although both chimpanzees (*Pan troglodytes*) and bonobos (*P. paniscus*) live in

multi-male/multi-female societies and form male-philopatric groups, they differ in male

aggression and the stability of temporary parties. Because the high social status,

prolonged receptivity, and characteristic sexual behaviors of females gathered the

attention of researchers, the behaviors of male bonobos have only been the focus of a

small number of studies. In addition, most studies comparing the behaviors of

chimpanzees and bonobos have been based on data obtained using different methods. In

this study, using the same observation method, we investigated party attendance and

aggressive interactions among male bonobos in Wamba and male chimpanzees in

Kalinzu. Male bonobos did not show the tendencies of a high degree of dispersion that

male chimpanzees showed in the absence of receptive females. The aggressive behaviors

between males within a group were not significantly different in the frequency of overall

aggressive interactions per observation unit between the two species. However, the types

of aggressive behaviors observed differed between the species. In bonobos, severe

aggressive interactions such as aggressive physical contact among adult males were not

observed, and most aggression occurred between the sons of the two highest-ranking

females. In bonobos, females participated in polyadic aggressive behaviors as the

aggressor, whereas all cases of coalitionary aggression in chimpanzees were from male

aggressors. These results suggest the considerable influence of female behaviors on the

substantial differences in aggressive interactions between the two species. This study

highlights both similarities and differences in male aggression between bonobos and

chimpanzees using the same method.

Keywords

Pan paniscus, Pan troglodytes, aggression, fission-fusion dynamics, party size,

dominance rank, , Wamba, Kalinzu Forest Reserve

Introduction

Factors concerning intermale mating competition primarily influence the socio-sexual behaviors of primate males. These include the number of receptive females, mating seasonality, operational sex ratio, dominance rank among males in a group, and relationships with other groups or out-group males. In addition to these factors, various forms of coalition or mutual tolerance may improve the mating success of males (van Hooff and van Schaik 1994). A coalition among males of the same group may benefit them through intergroup competition over reproductive females and food resources (Wilson et al. 2001; Mitani et al. 2010; Xiang et al. 2014). Furthermore, the coalition among some male members of a group may help them monopolize mating opportunities against other male members (Watts 1998; Duffy et al. 2007; Mitani 2009; Schülke et al. 2010).

The development of male bonding and cooperation in chimpanzees is not conditional on close genetic relatedness (Langergraber et al. 2007; Mitani 2009). This means that cooperation or coalition among males benefits these males, even if the benefit to their kin is not considered. However, these studies do not deny the possibility that cooperation or coalitions among related males are more beneficial in male-philopatric societies, in which most male members are kin-related to some extent

(Strier et al. 2014). The male-philopatric group of chimpanzees (*P. troglodytes*) is an

illustrative example. Males from the same group are more closely related within groups

than between groups, and they cooperatively patrol the boundary of the group range and

attack and sometimes kill males from different groups (Mitani and Watts 2005;

Langergraber et al. 2017). Such coalitionary attacks may result in the expansion of their

territories and increased access to food resources and females (Goodall 1986;

Wrangham 1999; Mitani et al. 2010; Nishida 2012). Furthermore, males sometimes

form coalitions with other males within the group to achieve alpha status and share

mating opportunities (de Waal 1982; Nishida 1983; Nishida and Hosaka 1996; Mitani

2009). Chimpanzees perform various affiliative behaviors with other males to enhance

their alliance (Nishida 1970; Goodall 1986; de Waal 1989). Such behaviors have also

been reported in spider monkeys (*Ateles* spp.), which usually form male-philopatric

groups (Santorelli et al. 2013). Males of white-bellied spider monkeys (*A. belzebuth*

belzebuth) often travel along the boundary area in alliance with other males (Shimooka

2005). Males of Geoffroy's spider monkeys (*A. geoffroyi yucatanensis*) have been

observed raiding the territory of the other groups while progressing in single file on the

ground in unusual silence (Aureli et al. 2006). Both of these behaviors are highly

similar to the patrolling and raiding behaviors of chimpanzees. Males of Geoffroy's

spider monkeys also show coalitionary aggression against subadult males within their

group (Campbell 2006). This is similar to the alliance of some males against others

within a group of chimpanzees.

Bonobos (*P. paniscus*) also form male-philopatric groups that are very similar

to those of chimpanzees (Kano 1982; Gerloff et al. 1999; Hashimoto et al. 2008;

Ishizuka et al. 2019). Therefore, we can expect to see similar behaviors in males,

including coalitionary aggression against other males in the group, coalitions among

males during intergroup encounters, and male cooperation for boundary patrols or

raiding the territories of other groups. However, these behaviors are seldom observed or

are observed with less aggression (Idani 1990; Kano 1992; Sakamaki et al. 2018;

Tokuyama et al. 2019), and it raises the question why bonobos and chimpanzees show

substantial differences in their aggression and behavior patterns, while both species

have male-philopatric groups. Although observations of males of both species are

necessary to answer this question, behavioral interactions among male bonobos have

been the focus of relatively few studies (Kuroda 1980; Ihobe 1992; Surbeck et al. 2012a,

b, 2017; Tokuyama et al. 2019; Cheng et al. 2022). This is likely because they are less

conspicuous than those in chimpanzees and because the female behavioral traits in

bonobos are more conspicuous (Parish 1994; Furuichi 1997, 2011; White and Wood

2007). Surbeck et al. (2012a) reported a positive correlation between the dominance

rank of males and the aggression frequency in which higher-ranking males showed

higher aggression frequencies than lower-ranking males. In Wamba, Ihobe (1992)

reported the frequency of aggressive interactions and that dominant–subordinate

relationships between a male–male dyad were easily recognized from their aggressive

interactions with few exceptional cases. However, this study was carried out during a

period when observations were made using artificial provisioning, which was

terminated in 1996 when a civil war interrupted research in the area (Hashimoto et al.

2008; Furuichi et al. 2012). In addition, most previous studies comparing the behaviors

of wild chimpanzees and bonobos have been based on published information for at least

one of the two species. Therefore, the data for these comparisons were not obtained

using the same observation methods. To clarify whether there are substantial differences

in male behaviors between chimpanzees and bonobos, we conducted the research using

the same observation methods under the natural conditions of male bonobos at Wamba,

Democratic Republic of the Congo (DR Congo), and male chimpanzees at Kalinzu

Forest Reserve, Uganda.

The patterns of association and behaviors of male bonobos are strongly

affected by prolonged sexual receptivity of females and the constant presence of

multiple receptive females in parties (Furuichi and Hashimoto 2002), moderation of

mating competition by the presence of multiple receptive females (Kano 1992; Furuichi

and Hashimoto 2002; Hashimoto et al. 2022), long-lasting mother–son relationships

(Furuichi 1989; Kano 1992), and strong support from dominant mothers in mating

competition and acquisition of alpha status (Furuichi 2011, 2019; Surbeck et al. 2011,

2019; Ishizuka et al. 2018). From these previous reports, we hypothesized that in

bonobos, unlike chimpanzees, the prolonged sexual receptivity of females and the

dominance status of the mother strongly influence mating competition among males so

that the importance of intermale aggressive interactions is relatively lower, and the

frequency and intensity of male aggression are reduced. In this study, we examined the

following four predictions expected from the hypothesis.

1. Male bonobos attend mixed-sex parties where they find receptive females more

    frequently than male chimpanzees.

2. Among bonobos, aggressive interactions between males are less frequent or involve

    less physical contact than in chimpanzees.

3. Aggressive interactions between males over access to receptive females are less

    frequent in bonobos than in chimpanzees.

4. In bonobos, aggressive interactions between males involve their mothers more

frequently than in chimpanzees.

Methods

Study sites and subjects

Bonobos in Wamba, Luo Scientific Reserve

Field assistants and author S.S. observed the wild bonobos of the E1 group at

Wamba in the northern sector of the Luo Scientific Reserve in DR Congo, where

long-term research has been conducted since 1973 (Kano 1992; Furuichi et al. 2012).

The bonobos were followed from September 19 to December 29, 2019. At the time of

this research, the group comprised approximately 40 individuals, including eight adult

males, three adolescent males, 13 adult females, a few immigrant adolescent females,

and many juveniles and infants. The age classes of subjects were categorized according

to Hashimoto (1997) and Toda and Furuichi (2020). All the individuals in the E1 group

were identified and habituated before the beginning of this study. The study subjects

comprised eight adult males aged over 14 years and three adolescent males aged 8–14

years which were categorized according to the methods of Hashimoto (1997). The

individual names, dominance rank, estimated birth year, and their mothers are shown in

Table 1. The dominance rank of males was calculated using David's score (David 1987)

based on dyadic aggressive interactions. The individuals were divided into one of the

following four categories, namely the alpha male (1), high ranking (2–3), middle

ranking (4–7), and low ranking (8–11).

Chimpanzees in the Kalinzu Forest Reserve

Field assistants and author S.S. observed the wild chimpanzees of the M group in

the Kalinzu Forest Reserve, Uganda, where long-term research has been conducted since

1992 (Hashimoto 1995; Hashimoto and Furuichi 2006). The observations were

conducted from February 3 to April 18, from June 23 to September 1, 2018, and from

March 11 to May 24 2019. During this study, the group consisted of approximately 100

individuals, including 15 adult males aged over 15 years (Goodall 1986), 29 adult

females, a few subadult males and females, and many juveniles and infants. All the

individuals in the M group were identified and habituated by the beginning of the study

period. The study subjects of the current study comprised ten adult males. The individual

names, dominance ranks, and estimated birth years of each adult male are shown in Table

2. The dominance rank of males was calculated using David's score (David 1987) based

on dyadic aggressive and submissive interactions. The individuals were divided into one

of the following four categories, namely the alpha male (1), high ranking (2–4), middle

ranking (5–7), and low ranking (8–10) (Table 2).

Data collection

Bonobos in Wamba were followed from approximately 0600 h from their

previous night's sleep site, and chimpanzees in Kalinzu were followed daily from

approximately 0700 h. We conducted observations using the same methods for both

bonobos and chimpanzees. We used the focal sampling method (Altmann 1974), in which

a focal animal was followed for as long as possible. We followed the first male we found

in the morning unless we had followed that individual on the previous day. When we lost

sight of the focal individual, we continued to observe if the individual was found again

within 30 min. However, if we could not find the individual for 30 min, we stopped the

focal observation for that day.

Party composition

Every 60 min during the focal following, we recorded the party composition that

the focal animal belonged to using the 1-h party method (Hashimoto et al. 2001;

Mulavwa et al. 2008; Tokuyama and Furuichi 2017). We recorded the individuals within

visible range at the beginning of each hour and included new individuals that joined the

party until the end of the hour. Each hour-long observation was included in the analysis

as a unit and described here as one observational hour unit (OHU) if we could observe

focal animals for more than 30 min within the hour of observation.

Aggressive interactions

We recorded all the observed intragroup aggressive interactions in each OHU

using ad libitum sampling, including cases where the focal animal was not involved. We

defined "aggressive interactions" as involving at least one aggressive behavior toward

one or more individuals: aggressive physical contact such as kicking, beating, and biting;

chasing; charging; directed aggressive displaying (including branch dragging and

provocative behaviors for bonobos); and non-vocal threatening. Undirected aggressive

displays, namely jumping in chimpanzees and branch dragging in bonobos, were only

recorded when other individuals responded by screaming or fleeing after that behavior.

All the aggressive interactions that involved subject males were included in the analyses.

To exclude the potential for overestimating the data on behaviors that were easily

observed, we excluded all observation data for days with less than three OHUs.

Definition of the presence of females with maximum swelling

The swelling status of the sexual skin of each female in each party was recorded

as one of the following two categories: non-swelling and maximum swelling (herein after

MS) (Hashimoto and Furuichi 2006). We defined all OHUs on days when we observed

females with MS at least once as "OHU under the presence of females with MS."

 Statistical analyses

In total, 217 OHUs were analyzed for all the subject bonobos with a mean of

19.7 ± 5.0 SD OHUs (range: 11–30) for each individual (Table 1). The mean duration of

OHU was 53.5 min (range: 30–60). For all the chimpanzee subjects, 677 OHUs were

analyzed with a mean of 67.7 ± 11.1 SD OHUs (range: 53–90) for each individual

(Table 2). The mean duration of OHU was 52.1 min (range: 30–60). We used R (version

4.2.2; R Foundation for Statistical Computing, Vienna, Austria, http://www.r-project.org

(accessed on 13 December 2022)) for all statistical analyses.

Comparison of party attendance of adult males

Among bonobos, females resume receptivity in the early stage of postpartum

infertility, which means that multiple females always show maximal swelling and

receptivity (Kano 1992; Furuichi and Hashimoto 2002; Hashimoto et al. 2022). In the

current study, more than 90% of OHUs (197 out of 217 units) on bonobos were defined

as "OHU under the presence of females with MS", and there were almost always

females showing MS. Therefore, we included all 217 OHUs on bonobos in the analyses

as a single dataset. Among chimpanzees, 376 out of 677 OHUs were defined as "OHU

under the presence of females with MS," and the remaining 301 OHUs were defined as

"OHU under the absence of females with MS." We inferred that females with MS were

present in the group when either research assistants or we observed one or more such

females on the same observation days. We compared the number of adult males

observed in a party between three different group conditions, namely bonobos,

chimpanzees with MS, and chimpanzees without MS. To determine whether the datasets

of the three conditions follow the same distribution, we first normalized each dataset

and then conducted the Kolmogorov–Smirnov two-sample tests for three combinations

by using the "ks.test()" function.

Intragroup aggressive interactions among adult males

In total, 92 aggressive interactions were recorded among the male bonobos. A

total of 76 cases were dyadic aggressive interactions between males, comprising 43

cases among adult males in 12 male–male dyads and 33 cases including adolescent

males in 10 male–male dyads. The remaining 16 cases were aggression from multiple

individuals, comprising 14 cases toward adult males and two cases toward adolescent

males. In total, 201 aggressive interactions among adult male chimpanzees including

177 dyadic aggressive interactions in 65 male–male dyads and 24 polyadic interactions

were recorded. Polyadic interactions include both cases of coalitions and cases when

there were multiple victims. We divided aggressive interactions into two conditions for

chimpanzees based on the presence/absence of females with MS. The aggressive

interactions were divided into four categories according to their intensity, namely

aggressive physical contact, chasing, directed aggressive display, and others including

non-vocal threatening. All the aggressive interactions among all the subject bonobos

(including adolescents), were summarized in the aggressive interaction matrix.

We calculated the frequencies of aggressive interactions in two different ways.

First, we calculated the frequency of overall aggressive interactions per OHU by

dividing the number of overall aggressive interactions by the number of OHUs.

Secondly, we calculated the frequencies of aggressive interactions per male in an OHU

by dividing the total number of males involved in aggressive interactions including both

aggressors and receivers by the total number of males observed during each OHU. To

determine whether the aggression frequencies in each group/condition were equal, we

conducted a chi-square test with a Bonferroni-corrected post hoc test using the

"pairwise.prop.test()" function.

Results
Grouping patterns of male bonobos at Wamba compared with chimpanzees at Kalinzu

Figure 1 shows the number of adult males observed in a party recorded using

the 1-hour party method in bonobos (Wamba E1 group) and chimpanzees (Kalinzu M

group). Among chimpanzees, the number of males in parties differed considerably

between days when receptive females showing MS (hereafter referred to as "females

with MS") were observed and days when they were not observed. In the absence of

females with MS, the modal value of the number of males was one, indicating that the

males were highly dispersed. This tendency of the high degree of dispersion was not

found in bonobos. The results of the Kolmogorov–Smirnov two-sample tests showed

that the party size distribution differed significantly in all three pairs (chimpanzees in

the absence of females with MS vs. chimpanzees in the presence of them, $D = 0.125$, P

$= 0.011$; bonobos vs. chimpanzees in the absence of females with MS, $D = 0.163$, $P =$

0.003; bonobos vs. chimpanzees in the presence of females with MS, $D = 0.145$, $P =$

0.006). Although the shapes of the party size distributions were similar between

bonobos and chimpanzees in the presence of females with MS and both showed the

highest frequency of the intermediate number of males, the null hypothesis that there is

no difference between the two distributions was statistically rejected.

Intragroup aggressive interactions among male bonobos at Wamba and male
chimpanzees at Kalinzu

Table 3 shows the numbers of dyadic aggression expressed and received among

11 male bonobos. Dyadic aggressive interactions were observed in 22 out of 55 dyads.

For 76 of the dyadic aggressive interactions, 37 cases were directed at two middle and

low-ranking males, Jiro and Jo. More than 80% of the interactions directed at them

comprising 30 cases out of 37 dyadic aggressions were from three high-ranking males

(Kitaro, Nobita, and Kiyota). These three males are the sons of the alpha female Kiku,

and Jiro and Jo are the sons of the second-ranking female, Jacky (Table 1). Out of 16

cases of aggression from multiple individuals, 15 were directed at Jiro and Jo. During

this observation period, the sons of Jacky were challenging the high status of the sons of

Kiku by provoking them (Shibata, personal observation), and aggressive interactions

among them frequently involved their mothers and brothers. A previous study has

shown examples of such behavior leading to the reversals of the dominance rank of

males (Furuichi 1997).

Table 4 shows the numbers of dyadic aggression expressed and received among

10 male chimpanzees. Dyadic aggressive interactions were observed in 33 out of 45

dyads. All the individuals except the alpha male were attacked multiple times by

higher-ranking males. For 100 of the dyadic aggressive interactions that occurred

among ten subject males, 47 cases were expressed by the alpha male, Goku.

Comparison of the frequency of aggressive interactions between chimpanzees and
bonobos

Table 5 shows the number of OHUs observed, the mean number of males in the

party, the total number of adult males observed in each OHU, and the number of

aggressive behaviors among adult male bonobos of the E1 group and chimpanzees of

the M group.

Among bonobos, approximately 50% of aggressive interactions were directed

display. Chasing behaviors were observed with similar frequency, but more than half

were polyadic aggression. Excluding them, the most frequent dyadic aggressive

interaction was directed display. Aggressive physical contact among adult males was not

observed.

Among chimpanzees, chasing was the most frequent aggressive interaction and

accounted for approximately 60% of the total aggressive interactions. Aggressive

physical contact accounted for 15% of all aggressive interactions, and more than 70% of

these occurred in the presence of females with MS.

The numbers of each type of polyadic aggressive interaction are shown in

Table 5 in parentheses. Among bonobos, all observed 14 polyadic aggressive

interactions involving adult males were chasing behaviors from multiple individuals

toward one individual, including at least one adult male as an aggressor. These

aggressive behaviors involved two to four aggressors. Females joined six cases of the

interactions. Kiku joined as an aggressor with her son in one of these six cases.

Out of all 24 polyadic aggressive interactions among chimpanzees, ten cases

were coalitionary aggression comprising nine chasing behaviors and one directed

aggressive display from two males. All of these coalitionary aggressions occurred in the

presence of females with MS, except for one case of chasing behavior. Female

chimpanzees did not join any of the coalitionary aggression as aggressors. Chimpanzees

also showed 14 aggressive interactions from one male towards multiple individuals.

These interactions include six cases of chasing directed at multiple individuals and eight

cases of aggressive displays after which other multiple males responded with screaming

or fleeing. These aggressive interactions from one individual to multiple individuals

were not observed in bonobos.

The frequency of overall aggressive interactions per OHU in bonobos (0.263)

was lower than that of chimpanzees (0.297). However, the difference was not

statistically significant (Chi-Square test: $\chi 2 = 0.778$, df = 1, p-value = 0.378). If the data

on chimpanzees was divided into two categories based on the presence/absence of

females with MS, the relationships in the frequencies of aggressive interactions were

chimpanzees in the presence of females with MS (0.404) > bonobos (0.263) >

chimpanzees in the absence of females with MS (0.163) and these differences were

statistically significant (Chi-Square test: $\chi2 = 48.41$, df = 2, p-value < 0.001). The

results of the Bonferroni-corrected post hoc tests showed that the differences were

significant in all three pairs: bonobos vs. chimpanzees in the presence of females with

MS (p-value < 0.01), bonobos vs. chimpanzees in the absence of females with MS

(p-value < 0. 001), chimpanzees in the presence of females with MS vs. chimpanzees in

the absence of them (p-value < 0.001) (Table 6).

However, when the difference in the mean number of males in the party was

considered, these differences between the species/conditions changed. The frequency of

aggressive interactions per male in bonobos (0.162) was higher than that of

chimpanzees (0.112), and the difference between the species was statistically significant

(Chi-Square test: $\chi2 = 4.821$, df = 1, p-value = 0.028). If the data on chimpanzees was

divided into two categories based on the presence/absence of females with MS, the

relationships in the frequencies of aggressive interactions per male were bonobos

(0.162) > chimpanzees in the presence of females with MS (0.136) > chimpanzees in

the absence of females with MS (0.072), and the differences were statistically

significant (Chi-Square test: $\chi2 = 20.471$, df = 2, p-value < 0.001). The results of the

Bonferroni-corrected post hoc tests showed that the differences were significant in

bonobos vs. chimpanzees in the absence of females with MS (p-value < 0. 001) and

chimpanzees in the presence of females with MS vs. chimpanzees in the absence of

them (p-value < 0.001), but not in bonobos vs. chimpanzees in the presence of females

with MS (p-value = 0.410) (Table 6).

Discussion

In this study, we have focused on the difference in the frequency and intensity

of male aggression between chimpanzees and bonobos (Idani 1990; Kano 1992; Mitani

and Watts 2005; Langergraber et al. 2017; Sakamaki et al. 2018; Tokuyama et al. 2019).

We have hypothesized that the patterns of association and behaviors of male bonobos

are strongly affected by the prolonged sexual receptivity of females and the constant

presence of multiple receptive females in parties, which may mitigate male mating

competition (Kano 1992; Furuichi and Hashimoto 2002; Hashimoto et al. 2022), and the

long-lasting mother–son relationships and strong support from mothers for mating

competition and acquisition of the alpha status (Furuichi 1989, 2011, 2019; Kano 1992;

Surbeck et al. 2011, 2019; Ishizuka et al. 2018). We have examined the four predictions

regarding those factors.

Among chimpanzees, the number of males in parties differed considerably

between days when receptive females showing MS were observed and days when they

were absent (Figure 1 b). In the absence of receptive females, the modal value of the

number of males in parties was one, indicating that the males were highly dispersed.

Low-ranking males were substantially more likely to range alone or in parties without

other males than high- and middle-ranking males in the absence of receptive females

(Shibata et al. 2022), which may have contributed to this trend. This tendency for

low-ranking males to avoid ranging with other males was likely to reduce the

probability of being attacked by other males (Shibata et al. 2022). However, in the

current study, similar tendencies of a high degree of dispersion were not found in

bonobos (Figure 1 a). In contrast, on the days when receptive females were observed,

the modal value of the number of chimpanzee males in parties was 6 or 7 (Figure 1 b).

Given that there were 15 adult males in this group, 40–47% of the males were found in

the observed party. In such a situation, there was no significant relationship between the

rate of solitary ranging and the rank of the males. This indicated that low-ranking males

actively participated in the party (Shibata et al. 2022). When we compared the

distribution of the number of adult males observed in the party between bonobos and

chimpanzees when females with MS were observed, a significant difference was

detected by the Kolmogorov–Smirnov two-sample test. However, both distributions

showed a similar tendency with unimodal distribution with a modal value of

approximately 40% of the maximum number of males. Therefore, bonobo males

seemed to show a similar tendency of attendance with chimpanzee males when females

with MS were present. This might be partially because male bonobos usually attend

mixed-sex parties that include most females including multiple receptive ones (Furuichi

and Hashimoto 2002; Furuichi et al. 2008; Hashimoto et al. 2022). Because of the long

receptive period of females in bonobos, receptive females are almost always present

(Furuichi and Hashimoto 2002; Furuichi 2011), and dispersal seen in the absence of

receptive females, as in chimpanzees, is rarely seen in bonobos. Considering this

difference in the periods of receptive female absence between chimpanzees and

bonobos, these results support the first prediction that male bonobos attend mixed-sex

parties where they find receptive females more frequently than male chimpanzees.

Comparison of aggressive interactions among adult males showed that the

frequencies of overall aggressive interactions per OHU were chimpanzees in the

presence of females with MS > bonobos > chimpanzees in the absence of females with

MS, and the differences were statistically significant in all pairs (Tables 5, 6). When the

difference in the number of males attending the party was considered, we obtained

different results. The frequency of aggressive interactions per male in bonobos was

higher than that of chimpanzees in the presence of females with MS. However, this

difference was not statistically significant (Table 6). The results in Table 5 seem to

indicate that male chimpanzees exhibit less aggressive behavior when females with MS

are absent. When females with MS are absent, low-ranking male chimpanzees

frequently range alone or in parties without other males. This tendency of low-ranking

males might make the frequency of aggressive interactions in the absence of MS

substantially lower because the calculations for the frequencies of aggression used all

recorded OHU including when focal males were observed without other males. When

the type of aggression is considered, aggressive interactions in bonobo males

predominantly comprised chasing or aggressive displays and aggressive physical

contact was not observed. Chimpanzee males showed 14 cases of aggressive behaviors

from one male toward multiple males. These interactions occurred when other males

were resting or grooming in proximity, and the alpha male was the aggressor in nine

cases of these interactions. Similar behavior has been reported in previous studies as

"separating intervention", where alpha males prevent other males from forming

affiliative relations and coalitions (de Waal 1982; Nishida and Hosaka 1996). This

highlights the complexity of the form of male competition among chimpanzees and the

strong influence of aggressive behavior from dominant individuals. These results do not

support the part of the second prediction that the frequency of aggressive interactions is

lower in bonobos than in chimpanzees. However, these results do support the second

part of this prediction that aggressive interactions between males involve less physical

contact in bonobos than in chimpanzees. This difference may explain why

lower-ranking bonobo males do not have to leave the party to avoid attacks by dominant

individuals.

436   Among chimpanzees, it has been reported that the presence of receptive

females increases male aggression (Watts 1998; Muller and Wrangham 2004). Similarly,

in the current study, the frequency of relatively severe aggressive interactions such as

aggressive physical contact and chasing behavior increased in chimpanzees when

females with MS were present (Table 5). In addition, nine cases out of ten coalitionary

aggressive interactions among male chimpanzees occurred when there were females

with MS. All these 10 cases of coalitionary attacks in chimpanzees were from two males

with no exception and the coalitionary attack formed by more than two males was not

observed. These results suggest that aggressive interactions among male chimpanzees

become more intense in the presence of receptive females, and males form coalitionary

attacks strategically for mating competition. Among bonobos, a previous study at

Wamba reported that the frequency of aggressive behavior that interferes with the

mating of subordinates is relatively low (2 of 36 cases of agonistic interactions observed

between August 1985 and January 1986, Furuichi, 1989). Similarly, out of 73 cases of

aggressive interactions in clear contexts observed in this study, seven cases occurred

over access to females with MS in this study (Shibata, unpublished data). These results

support the third prediction that aggressive interactions between males over access to

receptive females are less frequent in bonobos than in chimpanzees.

One characteristic of male aggression in bonobos observed here was that nearly

half of the male aggression occurred between the sons of the two highest-ranking

females (Table 3). During the observation, the sons of Jacky, the second-ranking female,

frequently challenged the sons of Kiku, the alpha female. If the interactions among

these males are excluded, the frequency of aggressive interactions among males

becomes substantially lower than in chimpanzees. This is likely to be the case in males

who are not competing over the dominance rank in the E1 group. Two old males and

one adolescent male showed only one or no aggressive behavior throughout the

observation. Therefore, these results partially support the second prediction that the

frequency of aggressive interactions between males is lower in bonobos than in

chimpanzees when we exclude the specific combinations of sons of two dominant

females. Among bonobos, all 14 cases of polyadic aggressive interactions were chasing

behaviors from multiple individuals towards one male. Three or four individuals joined

as aggressors in eight of the 14 cases, and females joined as aggressors in six cases of

all interactions. The victims of these interactions were a single male, Jiro, a son of the

second-ranking female, Jacky, in 13 of 14 cases. Kitaro or/and Nobita, the adult sons of

the alpha female, Kiku, joined in 12 of these 13 cases of aggression toward Jiro. This

also suggests that relatively large aggressive interactions involving multiple individuals

predominantly occur among males competing over dominance. However, only one case

of a mother supporting her son in aggressive interactions was observed during the study

period. Out of 27 cases of aggressive behaviors, including 16 dyadic and 12 polyadic

interactions, between Jiro and the adult sons of Kiku, Kiku only once joined aggressive

interaction with her son. In the PE group neighboring to the E1 group, the mother of one

of the males joined in approximately 6% of the aggressive interaction between males

(Tokuyama, personal communication). In LuiKotale, another study site, mothers joined

approximately 22% of aggressive interactions over access to receptive females (Surbeck

et al. 2011). The observations from the current study was lower than these. This might

be because Kiku had a newborn infant during the study period and could not join

aggressive interactions as before. Instead, Nao, another old female in a strong

relationship with Kiku, joined three cases of the remaining five interactions and

attacked Jiro with the sons of Kiku. Given that no polyadic aggressive interaction

involving females as the aggressor has been observed in chimpanzees in this study,

these observations indicate that bonobo females participate in aggressive interactions

more than female chimpanzees. Although such a tendency may differ depending on the

presence or absence of competing mother-son pairs, as far as this study is concerned,

these results support the fourth prediction that aggressive interactions between males

involve their mothers more frequently in bonobos than in chimpanzees. In addition, the

fact that one adolescent son of Kiku, Kiyota, was ranked as high-ranking also suggests a

strong influence from the dominance status of the mother on the dominance status of

sons in bonobos of the E1 group. Besides the constant presence of multiple receptive

females in the parties, such strong mother–son association might also promote high

attendance of males at mixed-sex parties addressed in the first prediction.

Among chimpanzees, dominance hierarchy is linear and aggressive behaviors

from dominant individuals to subordinate individuals are commonly observed in many

dyads (Nishida and Hosaka 1996; Newton-Fisher 2004). Similarly, in the present study,

all the individuals except the alpha male were attacked multiple times by higher-ranking

males (Table 4). In addition, the percentage of dyads in which aggressive interactions

occurred among all dyads was higher for chimpanzees than for bonobos. These results

suggest that aggressive interactions occur among a more limited number of individuals

in bonobos than in chimpanzees. Other previous studies suggest that when the

dominance relationships among chimpanzee males are stable, overt agonistic

interactions do not occur frequently (Kawanaka 1989; Takahata 1990). Like chimpanzee

males, aggressive interactions may be infrequent between bonobo males with a clear

dominance relationship or those who cannot participate in competition over dominance

status.

The results of the current study suggest that the mothers status among females

and their intervention in the conflict substantially impact the struggles over alpha status

among males in bonobos. A characteristic feature of struggles over alpha status among

bonobo males is that the mothers intervene in the conflict, and their status among

females has a substantial impact (Furuichi 2011, 2019; Surbeck et al. 2011). The history

of alpha status in the E1 group nearly the past 40 years from 1983–2020 has shown that

the status and support of the mother heavily influence alpha male status (Shibata and

Furuichi 2023). Such maternal support to males striving for alpha status in bonobos is

one of the most conspicuous aspects that differs from that of chimpanzees. Among

chimpanzees, age is one of the most important factors for acquiring alpha status in

males. Hasegawa and Kutsukake (2015) estimated the relationship between

competitiveness and age of males based on the data from three long-term study sites.

They reported that male chimpanzees have the highest potential to become alpha when

they are, on average, approximately 25 years old, except when males achieve alpha

status by coalition formation. In eastern chimpanzees in Mahale, alpha males acquire

their status at 19 to 37 years of age (Nishida 2008). In Gombe, middle-aged males reach

their highest rank and can dominate most, if not all, younger and older males by middle

age. Four cases have been reported in which males with a supportive elder brother

became alpha males (Bygott 1979). Although mothers sometimes supported their

offspring in fights, there is no evidence that such support influences the rank of adult

males (Bygott 1979).

These differences in maternal support between the two species might be caused

by the differences in mother–son relationships. The relationship between male offspring

and their mothers differs substantially between bonobos and chimpanzees (Kano 1992;

Schubert et al. 2013). Male bonobos maintain a close association with their mothers into

adulthood. In contrast, male chimpanzees significantly decrease their association with

their mothers when mothers resume cyclic receptivity, and eventually, they almost sever

their relationship with their mothers to associate with male adults. These differences

between bonobos and chimpanzees are also evident in a cluster analysis based on

association indices among all adult individuals in a group (Hashimoto and Furuichi

2015). Among bonobos, females gather and stay at the center of the party, and the males

remain peripheral (Kuroda 1979; Kano 1982; White 1988; Furuichi 1989). In contrast,

chimpanzee females are less gregarious than males and they spend a substantial amount

of time alone compared with males (Nishida 1968; Wrangham et al. 1992), although

different tendencies have been found between subspecies and populations (Lehmann

and Boesch 2004; Emery-Thompson and Wrangham, 2006). This tendency in bonobo

females makes the strong lifelong relationship between mothers and sons possible, and

maternal support has a substantial influence on the reproductive success of their sons.

The comparison of party attendance and aggressive interactions between

bonobos and chimpanzees using the same observation methods showed both similarities

and differences in the characteristics of male–male relationships. Unlike low-ranking

male chimpanzees in the absence of females with MS, male bonobos do not often range

alone. However, when the females with MS were present, both species show similar

tendencies in party attendance. Regarding aggression, the frequencies of aggressive

interactions were relatively similar between male bonobos and male chimpanzees in the

presence of females with MS. However, severe aggressive behaviors such as physical

contact aggression were not observed in bonobos. Previous studies and the study history

of Wamba suggested the strong impact of maternal support on the dominance status in

male bonobos (Shibata and Furuichi, 2023). In this study, nearly half of all aggressive

interactions among bonobos occurred between males with their mothers, which are the

two highest-ranking females in the group. The difference in male–male aggressive

interactions between the two species might also reflect considerable influences from the

difference in maternal support. Although the differences in male-male relationships

between chimpanzees and bonobos have been emphasized, this study has shown that

they share many similarities as members of the same male-philopatric society. In

addition, the current study has also shown that, despite the male-philopatric societies,

male-male relationships are greatly influenced by the mating behavior of females and

their social status. These results contribute to studies on the variation and evolution of

intermale relationships in the subfamily Homininae, including gorillas and humans,

which also show a similar male-philopatry tendency.

One limitation regarding the results of the current study is that the study period

for bonobos has been shortened because of the COVID-19 pandemic, and the study

periods for the two species were not equivalent. Moreover, because of bad visibility, we

could not clearly observe all aggressive interactions. Further studies addressing these

points will provide more accurate results. In addition, we need to note that the

behavioral tendencies of males found here cannot be generalized over populations of

bonobos and chimpanzees. Intermale relationships are likely to vary depending on

various factors, such as group size, the personality of the high-ranking male, the

presence or absence of supporting high-ranking mothers at the time, and differences

between the subspecies of chimpanzees. Studies using the same methods on other

populations or subspecies might yield different results. Therefore, further studies are

needed to understand better behavioral tendencies common to the species and the

divergence of inter–male relationships among male–philopatric societies.

Acknowledgments

We thank the National Forestry Authority of Uganda, the Uganda National Council for Science and Technology, and the Ministry of Scientific Research of the DRC. for their permission to conduct this research. We thank the local assistants for their help with field observations. We thank Nahoko Tokuyama, Tetsuya Sakamaki, Kazuya Toda, Chie Hashimoto, Hiroyuki Takemoto, Daisuke Shimizu, Akito Toge, and Shintaro Ishizuka for their help during the field research and their valuable discussions. This research was financially supported by the Leading Program in Primatology and Wildlife Science of Kyoto University, the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for JSPS Fellows, and JSPS Grants-in-aid for Scientific Research (to TF, 16H02753, 25304019 and 26257408). We also appreciate the anonymous reviewers for their helpful comments on this paper.

Author contributions

SS designed this study, collected and analyzed behavioral data, and wrote and

revised the manuscript. TF supervised SS in designing this study. All authors revised the

manuscript and gave final approval for publication.

Compliance with ethical standards

Conflict of interest

Ethical approval

This study was approved by the National Forestry Authority of Uganda and the Uganda

National Council for Science and Technology.

References

Altmann J (1974) Observational study of behavior: sampling methods. Behaviour 49:227–266

Aureli F, Schaffner CM, Verpooten J, et al (2006) Raiding parties of male spider monkeys: Insights

into human warfare? Am J Phys Anthropol 131:486–497. https://doi.org/10.1002/ajpa.20451

Bygott JD (1979) Agonistic behaviour, dominance, and social structure in wild chimpanzees of the

Gombe National Park. In: The great apes. pp 405–428

Campbell CJ (2006) Lethal Intragroup Aggression by Adult Male Spider Monkeys (*Ateles geoffroyi).*

Am J Primatol 68:1197–1201. https://doi.org/10.1002/ajp

David HA (1987) Ranking from Unbalanced Paired-Comparison Data. Biometrika 74:432–436

de Waal FBM (1982) Chimpanzee politics: power and sex among apes. Jonathan Cape, London

de Waal FBM (1989) Peacemaking among primates. Harvard University Press, Cambridge, MA

Duffy KG, Wrangham RW, Silk JB (2007) Male chimpanzees exchange political support for mating

opportunities. Curr Biol 17:586–587

Emery-Thompson, M., and Wrangham, R. W. (2006). Comparison of sex differences in

gregariousness in fission-fusion species: Reducing bias by standardizing for party size. In

Newton-Fisher, N. E., Notman, H., Reynolds, V., and Paterson, J. D. (eds.), Primates of

Western Uganda. Springer, New York.

Furuichi T (1997) Agonistic interactions and matrifocal dominance rank of wild bonobos (*Pan*

paniscus) at Wamba. Int J Primatol 18:855–875

Furuichi T (2011) Female contributions to the peaceful nature of bonobo society. Evol Anthropol

20:131–142. https://doi.org/10.1002/evan.20308

Furuichi T (1989) Social interactions and the life history of female Pan paniscus in Wamba, Zaire.

Int J Primatol 10:173–197. https://doi.org/10.1007/BF02735199

Furuichi T (2019) Bonobo and Chimpanzee: the lessons of social coexistence. Springer, New York

Furuichi T, Hashimoto C (2002) Why female bonobos have a lower copulation rate during estrus

631   than chimpanzees. In: Boesch C, Hohmann G, Marchant LF (eds) Behavioural diversity in

632   chimpanzees and bonobos. Cambridge University Press, Cambridge, UK, pp 156–167

633   Furuichi T, Idani G, Ihobe H, et al (2012) Long-term studies on wild bonobos at Wamba, Luo

634   Scientific Reserve, DR Congo: towards the understanding of female life history in a

635   male-philopatric species. In: Kappeler PM, Watts DP (eds) Long-term field studies of primates.

636   Springer, Berlin, pp 413–433

637   Furuichi T, Ihobe H (1994) Variation in Male Relationships in Bonobos and Chimpanzees.

638   Behaviour 130:211–228. https://doi.org/10.1163/156853994X00532

639   Furuichi T, Mulavwa, M;Yangozene K, Yamba-Yamba M, et al (2008) Relationships among fruit

640   abundance, ranging rate, and party size and composition of bonobos at Wamba. In: Furuichi T,

641   Thompson J (eds) The Bonobos: Behavior, Ecology, and Conservation. Springer, New York, pp

642   135–149

643   Gerloff U, Hartung B, Fruth B, et al (1999) Intracommunity relationships, dispersal pattern and

644   paternity success in a wild living community of Bonobos (*Pan paniscus*) determined from

645   DNA analysis of faecal samples. Proc R Soc B Biol Sci 266:1189–1195.

646   https://doi.org/10.1098/rspb.1999.0762

647 Goodall J (1986) The chimpanzees of Gombe: patterns of behavior. Cambridge University Press,

648  Cambrdige, UK

649 Hasegawa M, Kutsukake N (2015) Bayesian competitiveness estimation predicts dominance

650  turnover among wild male chimpanzees. Behav Ecol Sociobiol 69:89–99.

651  https://doi.org/10.1007/s00265-014-1821-9

652 Hashimoto C (1997) Context and Development of Sexual Behavior of Wild Bonobos (*Pan paniscus*)

653  at Wamba, Zaire. 18:

654 Hashimoto C (1995) Population census of the chimpanzees in the Kalinzu Forest, Uganda:

655  Comparison between methods with nest counts. Primates 36:477–488.

656  https://doi.org/10.1007/BF02382870

657 Hashimoto C, Furuichi T (2006) Comparison of behavioral sequence of copulation between

658  chimpanzees and bonobos. Primates 47:51–55. https://doi.org/10.1007/s10329-005-0144-x

659 Hashimoto C, Furuichi T (2015) Sex differences in ranging and association patterns in chimpanzees

660  in comparison with bonobos. In: Dispersing primate females. Springer, Tokyo

661 Hashimoto C, Furuichi T, Tashiro Y (2001) What factors affect the size of chimpanzee parties in the

662  Kalinzu Forest, Uganda? Examination of fruit abundance and number of estrous females. Int J

663  Primatol 22:947–959. https://doi.org/10.1023/A:1012061504420

Hashimoto C, Ryu H, Mouri K, et al (2022) Physical, behavioral, and hormonal changes in the

resumption of sexual receptivity during postpartum infertility in female bonobos at Wamba.

Primates 63:109–121. https://doi.org/10.1007/s10329-021-00968-w

Hashimoto C, Tashiro Y, Hibino E, et al (2008) Longitudinal structure of a unit-group of bonobos:

male philopatry and possible fusion of unit-groups. In: The Bonobos: Behavior, Ecology, and

Conservation. Springer, New York, pp 107–119

Idani G (1990) Relations Between Unit-Groups of Bonobos at Wamba, Zaire: Encounters and

Temporary Fusions. Afr Study Monogr 11:153–186

Ihobe H (1992) Male-male relationships among wild bonobos (*Pan paniscus*) at Wamba, Republic of

Zaire. Primates 33:163–179. https://doi.org/10.1007/BF02382747

Ishizuka S, Kawamoto Y, Sakamaki T, et al (2018) Paternity and kin structure among neighbouring

groups in wild bonobos at Wamba. R Soc Open Sci 5:171006.

https://doi.org/10.1098/rsos.171006

Ishizuka S, Kawamoto Y, Toda K, Furuichi T (2019) Bonobos' saliva remaining on the pith of

terrestrial herbaceous vegetation can serve as non-invasive wild genetic resources. Primates

60:7–13. https://doi.org/10.1007/s10329-018-00704-x

Kano T (1982) The social group of pygmy chimpanzees (*Pan paniscus*) of Wamba. Primates

23:171–188. https://doi.org/10.1007/BF02381159

Kano T (1992) The Last Ape: Pygmy Chimpanzee Behavior and Ecology. Stanford University Press,

Stanford, CA

Kawanaka K (1989) Age differences in social interactions of young males in a chimpanzee

unit-group at the Mahale Mountains National Park, Tanzania. Primates 30:285–305.

https://doi.org/10.1007/BF02381256

Kuroda S (1980) Social behavior of the pygmy chimpanzees. Primates 21:181–197

Kuroda S (1979) Grouping of the pygmy chimpanzees. Primates 20:161–183.

https://doi.org/10.1007/BF02373371

Langergraber KE, Mitani JC, Vigilant L (2007) The limited impact of kinship on cooperation in wild

chimpanzees. Proc Natl Acad Sci 104:7786–7790. https://doi.org/10.1073/pnas.0611449104

Langergraber KE, Watts DP, Vigilant L, et al (2017) Group augmentation, collective action, and

territorial boundary patrols by male chimpanzees. Proc Natl Acad Sci U S A 114:7337–7342.

https://doi.org/10.1073/pnas.1701582114

Lehmann J, Boesch C (2004) To fission or to fusion: effects of community size on wild chimpanzee

(*Pan troglodytesverus*) social organisation. Behav Ecol Sociobiol 56:207−216.

https://doi.org/10.1007/s00265-004-0781-x

Mitani JC (2009) Cooperation and competition in chimpanzees: Current understanding and future

challenges. Evol Anthropol 18:215–227. https://doi.org/10.1002/evan.20229

Mitani JC, Watts DP (2005) Correlates of territorial boundary patrol behaviour in wild chimpanzees.

Anim Behav 70:1079–1086. https://doi.org/10.1016/j.anbehav.2005.02.012

Mitani JC, Watts DP, Amsler SJ (2010) Lethal intergroup aggression leads to territorial expansion in

wild chimpanzees. Curr Biol 20:507–508. https://doi.org/10.1016/j.cub.2010.04.021

Mulavwa M, Furuichi T, Yangozene K (2008) Seasonal changes in fruit production and party size of

bonobos at Wamba. In: Furuichi T, Thompson J (Eds) The Bonobos. Springer, New York, pp

121–134

Muller MN, Wrangham RW (2004) Dominance, cortisol and stress in wild chimpanzees (*Pan

troglodytes schweinfurthii*). Behav Ecol Sociobiol 55:332–340.

https://doi.org/10.1007/s00265-003-0713-1

Newton-Fisher NE (2004) Hierarchy and social status in Budongo chimpanzees. Primates 45:81–87.

https://doi.org/10.1007/s10329-003-0064-6

Nishida T (1983) Alpha status and agonistic alliance in wild chimpanzees (*Pan troglodytes

schweinfurthii*). Primates 24:318–336. https://doi.org/10.1007/BF02381978

Nishida T (2012) Chimpanzees of the Lakeshore: Natural History and Culture at Mahale. Cambridge

University Press, Cambridge, UK

Nishida T (1970) Social behavior and relationship among wild chimpanzees of the Mahali mountains.

Primates 11:47–87. https://doi.org/10.1007/BF01730675

Nishida T (2008) Chimpanzee no shakai (the society of chimpanzees) (in Japanese). Toho Shuppan,

Osaka, Japan

Nishida T (1968) The social group of wild chimpanzees in the Mahali Mountains. Primates

9:167–224. https://doi.org/10.1007/BF01730971

Nishida T, Hosaka K (1996) Coalition strategies among adult male chimpanzees of the Mahale

Mountains, Tanzania. In: McGrew WC, Marchant LF, Nishida T (eds) Great Ape Societies.

ambridge University Press, Cambridge, UK, pp 114–134

Parish AR (1994) Sex and food control in the "uncommon chimpanzee": How Bonobo females

overcome a phylogenetic legacy of male dominance. Ethol Sociobiol 15:157–179.

https://doi.org/10.1016/0162-3095(94)90038-8

Sakamaki T, Ryu H, Toda K, et al (2018) Increased frequency of intergroup encounters in Wild

bonobos (Pan paniscus) around the yearly peak in fruit abundance at Wamba. Int J Primatol

39:685–704. https://doi.org/10.1007/s10764-018-0058-2

Santorelli CJ, Aureli F, Ramos-Fernández G, Schaffner CM (2013) Individual variation of whinnies

reflects differences in membership between spider monkey (Ateles geoffroyi) communities. Int

J Primatol 34:1172–1189. https://doi.org/10.1007/s10764-013-9736-2

Schubert G, Vigilant L, Boesch C, et al (2013) Co-residence between males and their mothers and

grandmothers is more frequent in bonobos than chimpanzees. PLoS One 8:1–11.

https://doi.org/10.1371/journal.pone.0083870

Schülke O, Bhagavatula J, Vigilant L, Ostner J (2010) Social bonds enhance reproductive success in

male macaques. Curr Biol 20:2207–2210. https://doi.org/10.1016/j.cub.2010.10.058

Shibata S, Furuichi T, Hashimoto C (2022) Adaptive Party Choice of Low-Ranking Males in

Fission–Fusion Dynamics of Chimpanzees in Kalinzu Forest Reserve, Uganda. Animals

12:2240. https://doi.org/10.3390/ani12172240

Shibata S, Furuichi T (in press) Intermale Relationships in Wild Bonobos at Wamba In Furuichi T,

Idani G, Kimura D, Ihobe H, Hashimoto C (eds) Bonobos and people at Wamba: 50 years and

counting.

Shimooka Y (2005) Sexual differences in ranging of *Ateles belzebuth belzebuth* at La Macarena,

Colombia. Int J Primatol 26:385–406. https://doi.org/10.1007/s10764-005-2930-0

Strier KB, Lee PC, Ives AR (2014) Behavioral flexibility and the evolution of primate social states.

PLoS One 9:1–14. https://doi.org/10.1371/journal.pone.0114099

Surbeck M, Boesch C, Crockford C, et al (2019) Males with a mother living in their group have

higher paternity success in bonobos but not chimpanzees. Curr Biol 29:R354–R355.

https://doi.org/10.1016/j.cub.2019.03.040

Surbeck M, Boesch C, Girard-Buttoz C, et al (2017) Comparison of male conflict behavior in

chimpanzees (*Pan troglodytes*) and bonobos (*Pan paniscus*), with specific regard to coalition

and post-conflict behavior. Am J Primatol 79:. https://doi.org/10.1002/ajp.22641

Surbeck M, Deschner T, Schubert G, et al (2012a) Mate competition, testosterone and intersexual

relationships in bonobos, Pan paniscus. Anim Behav 83:659–669.

https://doi.org/10.1016/j.anbehav.2011.12.010

Surbeck M, Deschner T, Weltring A, Hohmann G (2012b) Social correlates of variation in urinary

cortisol in wild male bonobos (*Pan paniscus*). Horm Behav 62:27–35.

https://doi.org/10.1016/j.yhbeh.2012.04.013

Surbeck M, Mundry R, Hohmann G (2011) Mothers matter! Maternal support, dominance status and

mating success in male bonobos (*Pan paniscus*). Proc R Soc B Biol Sci 278:590–598.

https://doi.org/10.1098/rspb.2010.1572

Takahata Y (1990) Adult males' social relations with adult females. In: The Chimpanzees of the

Mahale Mountains: Sexual and Life History Strategies. pp 133–148

Toda K, Furuichi T (2020) Low resistance of senior resident females toward female immigration in

bonobos (Pan paniscus) at Wamba, Democratic Republic of the Congo. Int J Primatol

41:415–427. https:// doi. org/ 10. 1007/ s10764- 020- 00151-w

Tokuyama N, Furuichi T (2017) Leadership of old females in collective departures in wild bonobos

(*Pan paniscus*) at Wamba. Behav Ecol Sociobiol 71:55.

https://doi.org/10.1007/s00265-017-2277-5

Tokuyama N, Sakamaki T, Furuichi T (2019) Inter-group aggressive interaction patterns indicate

male mate defense and female cooperation across bonobo groups at Wamba, Democratic

Republic of the Congo. Am J Phys Anthropol 170:535–550. https://doi.org/10.1002/ajpa.23929

van Hooff JARAM, van Schaik CP (1994) Male bonds: Afilliative relationships among nonhuman

primate males. Behaviour 130:309–337. https://doi.org/10.1163/156853994X00587

Watts DP (1998) Coalitionary mate guarding by male chimpanzees at Ngogo, Kibale National Park,

Uganda. Behav Ecol Sociobiol 44:43–55. https://doi.org/10.1007/s002650050513

White FJ (1988) Party composition and dynamics in *Pan paniscus*. Int J Primatol 9:179–193.

https://doi.org/10.1007/BF02737400

White FJ, Wood KD (2007) Female feeding priority in bonobos, *Pan paniscus*, and the question of

female dominance. Am J Primatol 69:837–850. https://doi.org/10.1002/ajp

Wilson ML, Hauser MD, Wrangham RW (2001) Does participation in intergroup conflict depend on

numerical assessment, range location, or rank for wild chimpanzees? Anim Behav

61:1203–1216. https://doi.org/10.1006/anbe.2000.1706

Wrangham RW (1999) Evolution of coalitionary killing. Am J Phys Anthropol Suppl 29:1–30.

https://doi.org/10.1002/(sici)1096-8644(1999)110:29+<1::aid-ajpa2>3.3.co;2-5

Wrangham RW, Clark AP, Isabirye-Basuta G (1992) Female social relationships and social

organization of Kibale Forest chimpanzees. In: Nishida T, McGrew W, Marler P, et al. (eds)

Topics in Primatology, Vol. 1: Human Origins. University of Tokyo Press, Tokyo, pp 81–98

Xiang ZF, Yang BH, Yu Y, et al (2014) Males collectively defend their one-male units against

bachelor males in a multi-level primate society. Am J Primatol 76:609–617.

https://doi.org/10.1002/ajp.22254

Figure 1. The number of OHUs in which the party included the respective number of adult males.

a) Male bonobos of the E1 group (N = 217). b) Male chimpanzees of the M group. Light blue and

orange bars represent the number of OHUs, when receptive females were absent (N = 301) and

present (N = 376), respectively.

Table 1. Subject males, dominance rank, birth year, and the number of observation hour units (OHU)

of bonobos of the E1 group.

Name (abbreviation)	Dominance rank	Birth year	Mother	OHU
Kitaro (KT)	Alpha	2004	Kiku	30
Nobita (NB)	High	1988	Kiku	18
Kiyota* (KY)	High	2009	Kiku	20
Loboko (LB)	Middle	1994**	-	11
Jiro (JR)	Middle	2002	Jackey	24
Dai (DI)	Middle	1975	-	23
Ten (TN)	Middle	1974	-	20
Tawashi (TW)	Low	1970	-	16
Gauche (GC)	Low	1988**	-	18
Hachiro* (HC)	Low	2009	Hoshi	16
Jo* (JO)	Low	2007	Jackey	21

In total, 217 OHUs were analyzed for all subject bonobos with a mean of 19.7 ± 5.0 SD OHUs

(range: 11–30) for each individual. The mean duration of OHU was 53.5 min (range: 30–60).

The dominance rank of males was calculated using David's score (David 1987) based on the dyadic

aggressive interactions.

* Adolescent male

** Year estimated

Table 2. Subject males, dominance rank, birth year, and the number of observation hour units (OHU)

of chimpanzees of the M group.

Name (abbreviation)	Dominance rank	Birth year	OHU
Goku (GK)	Alpha	1993*	90
Ponta (PO)	High	1995*	53
Ichiro (IC)	High	1980s*	76
Buru (BR)	High	1970s*	65
Prince (PR)	Middle	1997*	59
Taiki (TK)	Middle	1999	77
Deo (DO)	Middle	1970s*	62
Pietan (PT)	Low	2001	65
Black (BL)	Low	1998*	72
Jo (JO)	Low	2000*	58

In total, 677 OHUs were analyzed for all chimpanzee subjects with a mean of 67.7 ± 11.1 SD OHUs

(range: 53–90) for each individual. The mean duration of OHU was 52.1 min (range: 30–60). The

dominance rank of males was calculated using David's score (David 1987) based on the dyadic

aggressive interactions.

* Year estimated

Table 3. Aggressive interaction matrix with numbers of aggression expressed and received among 11

bonobos of the E1 group.

		Receiver									
		KT	NB	KY*	LB	JR	DI	TW	TN	GC	HC*	JO*
	KT	-	2	0	0	5	1	0	0	3	0	5
	NB	0	-	0	0	7	0	0	0	5	1	6
	KY*	0	0	-	1	1	0	0	0	0	0	6
	LB	0	0	0	-	0	3	0	0	0	0	0
Aggressor	JR	0	4	0	0	-	1	0	0	4	0	0
	DI	0	0	0	0	0	-	0	4	1	0	1
	TW	0	0	0	0	0	0	-	0	0	0	0
	TN	0	0	0	0	0	0	0	-	0	1	0
	GC	0	0	0	0	1	1	1	0	-	0	5
	HC*	0	0	0	0	0	0	0	1	0	-	0
	JO*	0	0	1	0	0	0	0	0	2	2	-

* Adolescent male

Dyadic aggressive interactions were observed in 22 out of 55 dyads.

37 cases of dyadic aggression were directed at two middle and low-ranking males, Jiro and Jo, and

30 cases of them were from three high-ranking males (Kitaro, Nobita, and Kiyota).

Table 4. Aggressive interaction matrix with numbers of aggression expressed and received among 10

adult chimpanzees of the M group.

		Receiver									
		GK	PO	BR	IC	PR	TK	DO	PT	BL	JO
Aggressor	GK	-	3	4	9	3	6	5	6	2	9
	PO	0	-	0	0	5	2	0	3	2	2
	BR	0	0	-	0	3	0	0	1	1	0
	IC	0	0	0	-	5	2	1	1	0	2
	PR	0	0	1	2	-	1	2	0	0	0
	TK	0	0	0	0	1	-	3	0	1	1
	DO	0	0	0	2	1	0	-	0	0	0
	PT	0	0	0	1	0	0	1	-	0	0
	BL	0	1	0	1	2	0	0	1	-	1
	JO	0	0	0	0	0	0	0	0	0	-

Dyadic aggressive interactions were observed in 33 out of 45 dyads.

47 cases out of 100 dyadic aggressive interactions were expressed by the alpha male, Goku.

Table 5. Type and frequency of aggressive interactions in bonobos of the E1 group and chimpanzees of the M group.

| Groups | Total number of OHUs observed | Mean number of males in the party | Total number of males observed | Number of male-male aggressive interactions | | | | | Total number of males involved in aggressive interactions | Frequency of aggressive interactions per OHU | Frequency of aggressive interaction per male |
				Physical contact	Chasing	Directed display	Others	Total			
Bonobos	217	3.60 ± 1.80	782	0 (0)	25 (14)	31 (0)	1 (0)	57 (14)	127	0.263	0.162
Chimpanzees	677	5.67 ± 3.09	3840	30 (0)	126 (15)	42 (9)	3 (0)	201 (24)	429	0.297	0.112
Chimpanzees (with MS females)	376	6.35 ± 2.98	2389	23 (0)	99 (13)	29 (6)	1 (0)	152 (19)	325	0.404	0.136
Chimpanzees (without MS females)	301	4.82 ± 3.03	1451	7 (0)	27 (2)	13 (3)	2 (0)	49 (5)	104	0.163	0.072

Numbers in parentheses represent the number of polyadic aggressive interactions. All 24 cases of polyadic aggression in chimpanzees involved two male aggressors. In six of the 14 cases polyadic aggression in bonobos, females joined the aggressive interactions as an aggressor.

Table 6. The results of the Bonferroni-corrected post hoc tests.

Group comparison	Adjusted p-value	
	Aggression per OHU	Aggression per male
Bonobos vs. Chimpanzees with MS	<0.01	NS
Bonobos vs. Chimpanzees without MS	<0.001	<0.001
Chimpanzees with MS vs. Chimpanzees without MS	<0.001	<0.001