Main Manuscript for

Reconfiguration of brain-wide neural activity after early life adversity.

Taylor W. Uselman[1], Russell E. Jacobs[2,3], Elaine L Bearer[1,3]

[1] University of New Mexico Health Sciences Center, Albuquerque, NM 87131
[2] Zilkha Neurogenetic Institute, Keck School of Medicine of University of Southern California, Los Angeles, CA 90033
[3] California Institute of Technology, Pasadena, CA 91125

Corresponding Author: Elaine L. Bearer
Email: elaine.bearer@gmail.com; ebearer@salud.unm.edu

Author Contributions: T.W.U. designed and performed research, contributed analytical tools, analyzed data and wrote the manuscript; R.E.J. designed and performed research, developed the imaging protocol, analyzed data and edited the manuscript; E.L.B. designed and performed research, analyzed data, wrote and edited the manuscript, and raised funding for the project.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences (Major), Neuroscience (Minor).

Keywords: manganese-enhanced magnetic resonance imaging, brain states, neuroimaging, early life adversity, acute threat

This PDF file includes:

      Main Text
      Figures 1 to 6

Abstract

Early life adversity (ELA) predisposes individuals to physical and mental disorders lifelong. How

ELA affects brain functions, leading to these vulnerabilities, is a mystery. To understand ELA's

impacts, investigations into neural activity affected by ELA must go beyond localized areas toward

simultaneous recordings from multiple widely distributed regions over time. Such studies will

expose relative activity between regions and discover shifts in regional activity in response to

different experiences. Here, we performed longitudinal manganese-enhanced magnetic resonance

imaging (MEMRI) to measure degrees of brain-wide neural activity in ELA-exposed mice across a

series of experiences in adulthood. To ascertain whether ELA resulted in atypical brain activity,

results were compared to those of the standard mouse (Std). MEMRI captured activity in the freely

moving home cage condition, and short- and long-term after exposure to TMT, a naturalistic

predator threat. Images were normalized and aligned then analyzed with statistical mapping and

automated segmentation. We found that neural activity in the home cage was greater in ELA

compared to Std in multiple striatal-pallidal and hypothalamic regions. Upon acute threat, neural

activity in Std increased in these regions to become more similar to that in ELA, while new

hyperactive responses in ELA emerged in the midbrain and hindbrain. Nine days after acute threat,

heightened neural activity in ELA persisted within locus coeruleus and increased within posterior

amygdala, ventral hippocampus, and dorsomedial and ventromedial hypothalamus. These results

reveal functional imbalances that arise between multiple brain-systems after ELA, which are

dependent upon context and cumulative experiences into adulthood.

Significance Statement

Early life adversity (ELA) is a crucial determinant of adult health. Yet the neurobiological basis for

this remains elusive. Localized brain regions display atypical neural activity in rodents who

experienced ELA, but how this contributes to overall brain state dynamics has been unknown. Here,

we used longitudinal manganese-enhanced MRI to detect brain-wide activities altered by ELA in

the adult across a series of conditions: freely moving, experiencing threat or its aftermath.

Computational analyses revealed that ELA produced widespread dynamic reconfiguration of brain-

wide neural activity in response to threat, shown here for the first time. Dynamic brain state

remodeling after ELA provides critical insights into human mental health vulnerabilities associated

with adverse childhood experiences (ACEs), and suggests plausible targets for intervention.

Main Text

Introduction

Early life adversity (ELA) poses risks for neuropsychiatric disorders, especially after a second

traumatic event, such as a life-threatening experience (1-5). ELA's long-term consequences on

global neural activity lasting into adulthood could serve as a biological basis for poor outcomes (6-

8). We are studying the brain states of adult mice who experienced ELA to test whether neural

activity in the home cage has been affected and whether their responses to threat are normal. To

ascertain how a potentially normal mouse is affected by ELA into adulthood, we compared ELA-

exposed mice to the standard mouse reared under generalized conditions (9, 10).

We invoked ELA in mice using a well-established paradigm: fragmented maternal care induced by

limited bedding (11, 12). Most studies using preclinical paradigms focus on localized brain regions

with invasive methods (13-27). Only a few have examined ELA's brain-wide consequences (28-

31). Lacking are any studies that examine the brain-wide effects of threat in the adult previously

exposed to ELA. Many questions remain that cannot be answered by localized techniques, such

as whether brain-wide activity is changed in an adult who has experienced ELA, and whether

activity in response to threat is also affected. To investigate these questions and overcome inherent

limitations of other techniques, we employed longitudinal manganese-enhanced MRI (MEMRI)

together with computational data-driven analysis. This provides quantitative information about

global brain activity, which sometimes is described with qualitative terms for directionality or to

generalize an aggregate of similar data.

Longitudinal MEMRI detects brain-wide activity before, during and after acute threat in awake freely

moving mice (32, 33). Predator odor is a widely accepted naturalistic paradigm to induce innate

threat responses in rodents (56-58). MEMRI does not require invasive or destructive procedures

(33). Mn(II) is safe – even continuous infusion of low doses has minimal effects on behavior (33,

34). We chose to deliver Mn(II) via two single bolus injections in order to avoid surgical procedures

required for continuous infusion and to allow for Mn(II) clearance between imaging sessions (32).

Systemic Mn(II) distributes throughout the brain parenchyma (35), enters active neurons through

voltage-gated calcium channels (36-38) and accumulates in neurons in an activity-dependent

manner (33, 39, 40). MR signal intensity is proportional to the frequency of action potentials (40).

Thus, MEMRI represents a quantitative readout of neural activity. Patterns of activity correspond

between MEMRI and IEG expression before and after challenge (32, 38, 41). Regional

accumulations of Mn(II) throughout the brain produce robust hyperintense MR signals with large

effect sizes at 100 µm^3 resolution (32, 33, 35, 41-43). These signals represent a composite picture
of regional neural activity occurring between injection and imaging timepoints. Alignment of multiple
MR images, collected repeatedly from large cohorts of individuals as they progress in parallel
through a series of experiences, produces a dataset that enables extraction of statistically
significant voxel-wise information about brain-wide activity over time, not possible with most other
techniques (32, 33, 44-55). Such processing is a well-accepted practice in human imaging that
yields much useful information and is routinely repurposed for rodent imaging (52, 56-59). To
ensure reliability and reproducibility, computational processing is verified and validated at each
step. Thus, longitudinal MEMRI can uniquely image enduring consequences of ELA on brain-wide
activity in the adult and its dynamics in response to threat.
80
Here, we examine the consequences of ELA on adult brain-wide activity during normal home cage
exploration in the adult and investigate how this activity changes in response to acute threat, short
and long term, with data-driven and hypothesis-based analyses of MEMRI images captured
longitudinally. Our alignments and computational processing pipeline were rigorously verified at
every step. We validated Mn(II) dose by comparing signal-to-noise ratios across all individuals. We
verified that Mn(II) signal replicated neural activity detected by other methods by measuring
intensities in circuits known to be involved in predator odor responses and by correlation of Mn(II)-
dependent enhancements with IEG (c-fos) expression. We used within- and between-group
statistical parametric mapping and automated segmentation with our new *InVivo* mouse brain atlas
to quantify relative activity within and between 102 segments brain-wide under each condition. We
found that volumes of activities in multiple segments throughout the brain differed significantly
between standard mice and those exposed to ELA undergoing the same experiential timeline.
93
94 **Results**
95
To witness neural activity in the home cage condition (HC) of adult mice who experienced ELA,
and their responses to predator stress short- and long-term, we adapted our longitudinal imaging
procedure (**Fig. 1A**) (32). We compared ELA mice (ELA, n = 12) to pool-weaned JAX-reared mice
of the same strain (Standard, Std, n = 12) to determine whether brain-wide activity of adult mice
who had experienced ELA differed from a phenotypically "normal" adult mouse and the degree of
difference.
102
Successful Replication of Behavioral Paradigms. To test whether our protocols for these two
paradigms, fragmented care (ELA) and predator stress (TMT), faithfully reproduced previously
reported behaviors (60-62), we compared videos of adult mice as they explored a custom arena

(**Fig. 1B-D**). For the effect of ELA experience, we measured the time ELA mice spent in the light at

baseline (BL) and compared that to Std mice. As expected, adult ELA mice spent 10% more time

in the light at BL than Std, demonstrating a persisting effect of ELA experience in an unprovoked

situation ($p < 0.05$ FDR, Cohen's d = 1.58) (**Fig. 1B**). For the effect of TMT, we measured time

spent in the light and compared between BL and TMT. As expected, TMT decreased time spent in

the light in both ELA and Std mice by more than 20%, demonstrating an acute effect of TMT ($p <$

0.05 FDR, Cohen's $d_{Std} = 2.29$, $d_{ELA} = 3.42$). To assess mouse recovery from MR anesthesia before

being presented with TMT, we measured distance traveled in 1 min epochs and tracked

movements from the beginning of a 10 min exposure to TMT (**Fig. 1C-D**). During the first minute of

TMT-exposure, distance traveled was equal to or greater than that at BL. Motility then decreased

by 50% or more over the 10 min TMT-exposure period ($p < 0.05$, FDR). These data support the

contention that our procedures faithfully reproduced ELA and TMT paradigms, giving us confidence

in assigning behavioral characteristics to unknown brain-wide activity.

Longitudinal Imaging. Our MR imaging protocol captures 8 sequential images across 12

individuals for each group (ELA and Std), yielding a total of 192 raw images with >2 million voxels

each (**Fig. 1E**). To ensure no differences between groups were introduced by processing, all

images were processed as a batch through our semi-automated processing pipeline (32, 33). This

pipeline yielded a combined dataset of skull-stripped, intensity normalized and spatially aligned

images ready for statistical evaluation. Quality assurance considerations included reviewing

imaging parameters, performing SNR measurements to confirm similar Mn(II) doses, visualizing

modal scaling output in voxel-by-intensity histograms to monitor equivalent intensity normalization,

calculating Jaccard Similarity and Mutual Information Indices for alignment accuracy, and

determining that smoothing does not significantly change regional signal intensities. All images

reported here met our quality assurance standards (*SI Appendix* **Fig. S1-2** and **Table S1**).

Brain States in Mice Exposed to ELA. To detect Mn(II)-enhancements in ELA mice across

conditions from the home cage (HC) to post-threat (TMT) to nine days after threat (D9), we

compared post-Mn(II) images for each condition to each mouse's respective pre-Mn(II) image using

voxel-wise statistical parametric mapping (SPM) (**Fig. 2**). SPM of images at HC revealed large

(16.9% of total) volumes of statistically significant signals in multiple brain regions identified by

anatomy, demonstrating that the unprovoked ELA brain was already highly active (T = 2.72, $p <$

0.01, cluster corr.) (**Fig. 2A**). Visual inspection of SPMs revealed larger volumes of signal within

regions of the olfactory bulb, striatum/pallidum, hypothalamus, hindbrain, and cerebellum, and

smaller volumes within subregions of hippocampus, amygdala, and hindbrain. Immediately after

TMT, volumes of statistically significant voxels increased by +1.4% overall. SPM volumes within

these regions were maintained or expanded, with sizable increases in the mid- and hindbrain (e.g.,
LC) (**Fig. 2B**). By D9, activity decreased in olfactory, hindbrain, and cerebellum, or increased in
hippocampus, amygdala, and midbrain (e.g., PAG), leading to an overall decrease of -2.9% from
HC (**Fig. 2C**). Other subregions also changed, such as a decrease of the volume of statistical signal
in the lateral septum (LS) and appearance of new signal in the paraventricular nucleus of the
thalamus (PVT). Direct SPM paired t-test comparisons between ELA post-Mn(II) images at each
condition added to these findings from activation volumes, to also identify statistical differences in
signal intensities within many regions across conditions (*SI Appendix* **Fig. S3**). These findings
indicate that ELA-exposed brains are already highly active before a second provocation in
adulthood, and that short- and long-term brain responses to TMT have unique patterns of activity,
with regions emerging or disappearing between conditions.

Magnitude of Signal Enhancements. To determine whether these statistical maps actually
represent changes in intensity values across conditions, we measured intensities within 3 x 3 x 3
voxel regions of interest (ROI) in TMT-responsive regions and compared between HC and TMT
(**Fig. 3A**, and *SI Appendix* **Tables S2** and **S3**) (63, 64). Our voxel-wise SPM of ELA images at HC
and TMT revealed significantly enhanced signal intensities after TMT in all four expected regions
($p < 0.05$ cluster corr.) (**Fig. 3A**, *top*). At HC, SPMs show that dlOB-DII and vlAON were already
enhanced and only minimally changed across conditions. Region of interest (ROI) measurements
within these locations revealed that SPM signals at HC represented large signal differences over
pre-Mn(II) (6.8%, dlOB-DII; 8.0%, vlAON) with standard deviations of similar magnitude, leading
to large Cohen's D effect sizes (d = 1.6, dlOB-DII; d = 1.5, vlAON; $p < 0.05$ FDR). Standard
deviations are shown in **Fig. 3A** and listed in *SI Appendix* **Table S3**. In aOC and aCEA, SPM
signals were minimal at HC and increased from HC to TMT (**Fig. 3A**, *top*). These SPM signal
changes were recapitulated in ROI comparisons, with aOC and aCEA showing small magnitudes
of signal enhancement over pre-Mn(II) at HC (1.4% and 3.0%), and larger magnitudes at TMT
(3.5% and 6.2%) (**Fig. 3A**, *bottom*). Thus, in this case SPMs represented signal intensity
differences. Signal intensity increased in all four regions expected to respond to TMT (+2.1-3.9%,
d = 0.85-1.20, $p < 0.05$ FDR), demonstrating robust TMT-induced enhancement.

Statistical mapping of ELA at HC (**Fig. 2** above) indicated signal enhancements within regions
normally only responsive after TMT in Std (32). To determine the degree of signal intensity
differences between ELA and Std at HC and test whether ELA signal truly exceeded Std in these
regions, we measured and compared signal intensities within 3 x 3 x 3 voxel ROIs (**Fig. 3B**, *SI*
Appendix **Tables S2** and **S4**). Magnitudes of signal enhancement at HC over pre-Mn(II)
corresponded to the presence or absence of significant SPM signal for both Std and ELA (non-

enhanced: SI = 2.0-3.6%, d = 0.5-0.7; enhanced: SI = 6.1-17.9%, d = 0.96-5.17). Direct

comparisons between groups revealed that ELA intensities were heightened at HC in all four

measured regions (SI = 4.1-6.6%, d > 1.00, p < 0.05 FDR). Standard deviations are shown in **Fig.**

3B and listed in ***SI Appendix* Table S4**. Thus, ELA signal intensities within threat-responsive

regions exceeded that of Std even before threat. ELA also displayed a lower variance than Std.

Together these observations support the conclusion that differences in SPM-identified activation

volumes, at p < 0.01 and cluster size ≥ 8 voxels, reliably represent intensity differences between

post-Mn(II) images. However, when SPM analysis reveals overlapping anatomy of activation

volumes between groups or conditions, direct measurements of post-Mn(II) intensities may be

necessary to uncover potential differences in activity levels that are not detected by activation

volumes.

Mn(II)-Enhanced Signal Corresponds to c-fos Expression. We next sought to test whether

differences in Mn(II)-enhancement corresponded quantitatively to differences in neural excitation,

as detected by c-fos expression (33, 41, 66, 67) (**Fig. 4**). We previously reported a correspondence

between c-fos and MEMRI before and after challenge (predator odor) and found similar changes

in activity between the two modalities (32). Here, we compare intensity values from *in vivo*

longitudinal MEMRI with positive nuclei counts from c-fos immunohistochemistry, which is an

endpoint where longitudinal dynamics are not feasible. A subset of the mice that underwent the

longitudinal MEMRI protocol were processed for c-fos immunohistochemistry (n_{Std} = 5 and n_{ELA} =

6). We selected pairs of regions in sections stained for c-fos where anatomy could be cross-

correlated with MR slices for each mouse: hypothalamus (HYP) and thalamus (THA); superior

central raphe (CS) and midbrain (MB) (**Fig. 4A-B**). First, we compared regions for each modality

separately (**Fig. 4C-D**). Statistical comparison of Mn(II)-enhanced signal intensities in HYP and CS

were statistically greater than in THA and MB, respectively (**Fig. 4C**, p < 0.001). Numbers of c-fos+

nuclei revealed similar differences between regions (**Fig. 4D**, p < 0.001). Robust effect sizes were

observed for both modalities (c-fos: MEMRI: d = 1.88-6.37; d = 3.75-5.03). Statistical differences

in MR signal intensities from this subset of mice were replicated in a second subset from each

group, demonstrating test-retest reliability (d = 2.94-8.43, p < 0.001, FDR) (***SI Appendix* Fig. S4**).

To test further whether intensities correlated linearly with c-fos counts, we calculated Pearson's

pairwise correlation coefficient for MEMRI versus c-fos data across the four regions with bootstrap

resampling and permutation testing. In both groups, correlations were statistically significant (Std:

r = 0.636, p < 0.0001; ELA: r = 0.582, p < 0.0013) (**Fig. 4E** and ***SI Appendix* Fig. S5**).

A between-group comparison of c-fos and MEMRI at D9 for each region found that both modalities
found increased activity in HYP ($d_{c\text{-}fos}$ = 0.38; d_{MEMRI} = 2.21) and CS ($d_{c\text{-}fos}$ = 1.07; d_{MEMRI} = 2.02) in
ELA relative to Std. While MEMRI differences were statistically significant in HYP and CS ($p < 0.05$,
FDR), c-fos differences were not, demonstrating that MEMRI could detect statistically significant
differences with only 5-6 mice possibly due to larger values and smaller variance in MEMRI data.
Comparison of c-fos at D9 with MEMRI from all three conditions in these regions revealed
statistically significant correlations ($r > 0.582$, $p < 0.0013$). This result shows that the relative levels
of activity were consistent in these regions across the longitudinal imaging procedure. Thus, our
data provide evidence that differences in the degree of MEMRI signal intensities reflect differences
in c-fos expression for the most part.

Deeper Dives into Brain States of Adult Mice Exposed to ELA. To assign neural activity in ELA
mice to specific anatomical regions, we aligned statistical maps of enhanced voxels with our *InVivo*
Atlas (***SI Appendix* Table S5**). Then we quantified activity in those anatomically defined segments
at each condition with a new approach, fractional activation volumes (FAV) (32) (**Fig. 5**). The FAV
is the fraction of activated to total voxels in each segment. Activated voxels were those found to be
statistically enhanced in SPM comparisons of post-Mn(II) to pre-Mn(II) images at each condition.
FAVs were visualized as column graphs. Comparing column graphs between conditions
demonstrated changes in FAVs as mice progressed through successive conditions. For
conciseness, we summarize key regional differences from this quantitative analysis using
qualitative terms for direction, magnitude, and aggregate patterns. Quantitative information
underlying these terms is available in ***SI Appendix* Tables S6-8**.

Overlays of statistical maps show that much of the global activity in ELA at HC (**Fig. 5A, blue**)
versus pre-Mn(II) coincides with that of TMT (**Fig. 5A, red**) in the same group (**Fig. 5A, overlap**
purple) ($p < 0.01$ cluster corr., $d = 0.79\text{-}8.19$) (***SI Appendix* Table. S6A**). Some regions expand
while a few others contract from HC to TMT. By FAV analysis of ELA, we see that at HC large FAVs
were prominent in various domains (OLF, STR/PAL, HYP, and CB) with some segments (MOBmi,
92%; MS, 100%; BST, 92%; and AHN, 97%) almost totally active. Immediately after TMT, FAV
values were retained or increased, with the largest increases in CEA (+9%) and LC (+42%). Several
STR/PAL and THA segments decreased in FAV after TMT, including LSc (-28%), LGd (-53%) and
MH (-34%). No relationships between segment volume and FAV were observed. Differences in
signal intensities between conditions in these segments were also statistically significant by SPM
comparisons between successive post-Mn(II) images, with the greatest differences in the LC ($p <$
0.05 cluster corr.) (***SI Appendix* Fig. S6** and **Table S6B**).

We then compared SPM overlays and segmental FAVs between TMT (**Fig. 5B, red**) and D9 (**Fig.**
5B, green; overlap, brown) of ELA mice to learn how TMT affected brain-wide activity long term
(**Fig. 5B**). Segmentation analysis revealed globally that FAV decreased in three domains (OLF,
HB, and CB; -14% on average) with the most substantial change in CB (-19%). Most other
segments either retained or further increased FAV, particularly within AMY, THA, HYP and MB
domains (+11% on average). In many segments where FAV had increased following TMT, FAV at
D9 either mostly persisted (LC, 34%) or increased (CEA, +29%), suggesting a longer lasting effect
of the threat experience. Segments where FAV decreased acutely after TMT typically remained
low or further decreased (LSc, -12%). Others increased at D9 (MH, +14%). These results suggest
that brain activity in some segments does not return to pre-threat levels.

To test whether activities return or not to pre-threat levels, we compared HC with D9 using a similar
strategy in both ELA and Std mice (**Fig. 5C and 5D**, also *SI Appendix* **Table S6A**). Both overlays
of SPMs (**Fig. 5C and 5D, overlap, turquoise**) and columns graphs of FAV show that at D9 after
threat, activity in both ELA and Std was increased substantially compared to HC in many domains
(AMY, STR/PAL, HYP, and MB). The magnitude of FAV increase in these domains was twice as
large on average in Std than in ELA (+28% vs +11%). This pattern also appeared in pairwise
comparisons of post-Mn(II) intensities between conditions (*SI Appendix* **Fig. S7** and **Table S6B**).
Neither the brain-wide activity of ELA nor Std returned to pre-threat levels at D9. These data
suggest regions in the brain remain hyperactive one week or more after acute threat.

Visual comparison of pairwise statistical maps within group at HC and D9 suggested statistical
differences between groups. At HC, before threat, FAVs of ELA were substantially greater than
those of Std across brain segments (31% vs 17% on average). These larger FAVs in ELA could
not be attributed to procedural differences, such as $MnCl_2$ dosing, MR parameters, imaging timing,
image processing, segmentation analysis (*SI Appendix* **Fig. S2** and **Table S1**), differences in pre-
Mn(II) signal intensities between groups (*SI Appendix* **Fig. S8**) or differences in anatomy between
groups (*SI Appendix* **Fig. S9**). While Std exhibited larger FAV increases from HC to D9 than ELA
in most segments (+12% vs +5% on average), ELA exhibited larger increases than Std in a subset
of segments, including RE (+35% vs +16%) and LC (+34% vs +4%). In other segments, ELA
displayed the opposite directional change from HC to D9 compared to Std, including PA (+19% vs
-4%), LSr (-18% vs +31%), IPN (-2% vs +33%), and PCG (+19% vs -5%). Except for IPN, these
segments also responded acutely to TMT in ELA (**Fig. 5B**). These findings suggest that brain-wide
activity and its evolution from threat-naïve (HC) to threat-experienced (D9) conditions differs
between groups and that ELA mice resemble a post-threat-like state already at HC. Furthermore,
those segments that respond to TMT in ELA appear to be major contributors to differences between

groups at D9. These observations prompted direct between-group statistical analyses for each
condition.

ELA Affected Many Segments Within Each Anatomical Domain. To determine whether signal
intensities between ELA and Std differed statistically between groups, and to map the anatomical
distribution of those differences, we performed direct SPM comparisons. First, we performed an
unpaired t-test comparing baseline pre-Mn(II) images between groups, which revealed small
intensity differences (*SI Appendix* **Fig. S7**). For subsequent comparisons, post-Mn(II) images were
adjusted to correct for these baseline differences. We then compared ELA and Std post-Mn(II)
images at each condition by an unpaired SPM t-test ($p < 0.05$, cluster corr., d = 0.70-5.30) (**Fig. 6**
and **Movie S1**). Segmentations of SPM difference maps at each condition were used to obtain the
number of statistically enhanced voxels (in either ELA > Std or Std > ELA) per segment, which
were then divided by the total number of voxels in each segment producing the fractional difference
volume (FDV) (**Fig. 6A-D**, and *SI Appendix* **Table S7**). At all three conditions, ELA exhibited
greater signal intensities than Std (i.e., heightened activity) in most segments but to a lesser extent
after TMT (HC, 17%; TMT 11%; D9, 12% on average), supporting observations from the within-
group SPMs shown in **Fig. 5**. From HC to D9, the total volume of enhanced voxels in ELA relative
to Std decreased by 4.4%.

Closer examination of FDVs within different anatomical domains identified heightened activity at
HC in ELA compared to Std within OLF (29% on average), STR/PAL (40% on average), and HYP
(26% on average) (**Fig. 6A**, yellow columns). At TMT, differences in these domains decreased (-
17% on average), although heightened HYP activity in ELA was somewhat preserved (-7%) and
new differences became noticeable in the CB (+16%) (**Fig. 6B**). At D9, heightened activity in the
HYP domain of ELA returned or even increased over HC (+13% on average) (**Fig. 6C**). While
heightened activity was present in most HYP segments for each condition (26% at HC vs 19% and
32% at TMT and D9), many other domains in ELA exhibited smaller volumes of heightened activity
over Std at TMT and D9, particularly segments within STR/PAL (40% at HC vs 14% at both TMT
and D9). In contrast, Std exhibited heightened activity compared to ELA within only a few segments
of CTX, THA, HYP, MB and CB domains (4.6% on average across all segments and conditions),
with the most differences occurring at TMT (6%) (**Fig. 6A-C**, blue columns). Thus, ELA in general
displayed heightened brain-wide activity over Std at HC, with FDV for ELA > Std of 10% or greater
in 53 of 102 total segments. Of the 53 segments, the degree of differences lessened upon
experiencing threat in most but not in all (TMT, 31; D9, 30). Next, we will take a deeper dive into
segmental activity dynamics.

Examining segment-wise FDV across conditions revealed at least five patterns of segmental
dynamics (**Fig. 6D**). To facilitate interpretation of activity differences across conditions, we plotted
FDVs indicated by arrows in **Fig. 6A-C** as line graphs (**Fig. 6D**) and FDV values listed in *SI*
Appendix **Table S7**. These graphs exemplify five dynamic patterns across conditions of
heightened activity in ELA > Std (yellow) or Std > ELA (blue). *One* pattern was heightened activity
in ELA at HC, which became more similar between groups at TMT (**Fig. 6D**, blue), as exemplified
by MS and VMH and observed across 26 segments. Some of these segments (18) continued to be
similar between groups at D9 (e.g., MS) while others returned (7) to heightened activity in ELA
(e.g., VMH). A *second* pattern followed an opposite dynamic in response to TMT. In these
segments (8), volumes of heightened ELA activity expanded acutely with TMT and returned to HC
levels at D9 (CEA and CS, red). A *third* pattern appeared in a few segments (8), where FDVs of
ELA rose only at D9 (PA and DMH, green). A *fourth* pattern emerged in 5 segments (e.g., DP and
ARH, black). Similar to pattern one or three, ELA activities in these segments were greater than of
Std at HC or D9. Dissimilar to pattern one or three, ELA activities decreased in these segments
below those of Std at TMT. A comparison between male and female found that ELA females might
contribute to between-group differences at D9 in VMH, ARH and PA more than ELA males, with
sex-differences in PA also present in pre-Mn(II) images (*SI Appendix* **Fig. S10**). A single segment
followed a *fifth* pattern (LC, purple), where activity was greater in Std at HC and then reversed after
TMT through D9 to become heightened in ELA. Together these data demonstrate that brain-wide
activity of ELA mice is reconfigured across conditions and that dynamics in response to threat, both
brain-wide and segment-wise, differ dramatically from Std.
342
343   **Discussion**
344
In this study we mapped the spatial-temporal transformations that ELA imposes on brain-wide
activity, as detected initially when in a home cage (HC) environment and then short and long-term
after challenge by an acute threat as compared to a Standard, phenotypically "normal," mouse.
Three different brain states (68), as defined here by widely distributed patterns of fractional
activation volumes among segments, emerged within each group. In all three conditions, brain
states of adult mice reared with the fragmented maternal care model for ELA differed from brain
states of adult Std mice, with greater overall neural activity after ELA (**Fig. 6E** and *SI Appendix*
Fig. S11). Even without provocation, ELA mice at HC had heightened fractional activation volumes
in regions known to be activated by predator odor or altered in reward processing: olfactory and
hypothalamic segments (67, 69-74), as well as striatum-pallidum (32, 75, 76). Distribution of
heightened activity expanded immediately after threat to engage subcortical structures as well as
monoaminergic nuclei and deeper brainstem segments in ELA, as also occurred in Std (32). Neither

group regained HC activity patterns nine days after threat. Across all conditions, activity in
hypothalamus of ELA was higher than Std. At D9, new differences emerged in the ventral
hippocampus and posterior amygdala, where activity was greater in ELA. Our longitudinal brain-
wide imaging strategy coupled with our computational processing and analytics empowered us to
follow these brain states in the same individuals as they progress through this series of experiences
in sample sizes sufficient for statistical significance. Our processing pipeline, aligning 192 whole-
brain images captured with $100\mu m^3$ resolution, allows precise localization of statistically significant
activity common across multiple individuals. This unique capability is not possible with most other
physiological approaches.

We found dynamic changes in segment-wise fractional activation volumes between conditions
throughout the brain in both cohorts. When a segment transitions from lower to higher FAV/FDV,
we predict that segment to have a more dominant functional role within its connectome. For
example in ELA, the increased volume of activity In CEA (from 43% to 53%) as the mouse
progresses from home cage to immediately after threat may contribute to heightened activity
throughout threat circuity (77). That ELA mice sustained heightened CEA activity at all three
conditions, suggests that ELA leads to persistent engagement of threat circuitry. This sustained
engagement of threat circuitry after ELA may sensitize CS and LC to threat, promoting
monoaminergic modulation of segmental activities toward the patterns observed at D9 (78-81).
Differing levels of activity among segments of a single network across different conditions could
produce different internal states (82, 83) and/or outcomes from that same network, be they actions,
decisions, emotions, thoughts, memories, or creative works, such as music, poetry, or art.

Results presented here should be considered in light of several limitations. 1) Theoretically, small
variations in rearing environment could contribute to differences between groups not attributable to
ELA. However, multi-site studies in mice find that different breeding/rearing facilities have no effect
on stress reactivity (84) and minimal explanatory value for at least 14 other physiological measures
(85). JAX-reared mice are the phenotypical standard and the most commonly used mouse by
researchers globally (9), making them an invaluable comparison group to generalize findings
across research facilities (86). 2) Although we found minimal sex differences between male and
female ELA mice, we cannot fully attribute heightened activity in HYP and vHIP/AMY segments to
ELA because we only had a small sample size and because our Std group was all male. Other
studies suggest a combinatorial effect of ELA and sex is likely within these threat responsive
regions (73, 87, 88). 3) Small anatomical variations could have influenced our results (89-92).
However, our Jacobian deformation analysis reported minimal alterations in anatomical volumes,
and our SPM analysis accounted for any signal differences by preserving intensity concentrations

within voxels (93, 94). 4) Atlas segmentation yields volumes that vary widely, ~10^2-10^4 voxels. This

may miss some large segments where small- to medium-sized activations only produce small

fractional volumes. 5) We recognize that statistical methods are limited due to the choice of

thresholding and pre-preprocessing, which is particularly difficult with MRI (33, 95-97). We

minimized these potential issues by rigorously examining each processing step for image quality,

by ensuring sample size was sufficient to detect moderate to large signal enhancements from

Mn(II), and by selecting statistical thresholds based on effect size and false positive rate

corrections. Fractional volumes, pioneered here, provide quantitative detail about the volumetric

extent of Mn(II)-enhanced signals, and are limited by their dependence on statistical parametric

mapping and segmentation, and could miss variability and discrete locations of signal intensities

within segments. Caveats of MEMRI technology include pre-existing tissue enhancement, intensity

normalization procedures, and differences in dosing and imaging parameters. We have dealt with

each of these in detail in this study, as described above.

To what degree does Mn(II) signal faithfully represent neural activity? Staining by c-fos shown here

and previous reports, including IEG and non-IEG methodology, contend that Mn(II) tags active

neurons (32, 36-41, 98, 99). Patterns of activity in response to challenge as seen by MEMRI

reproduce those detected by IEG expression (32, 98). Both MEMRI and c-fos are dependent on

calcium entry into neurons, though minor non-voltage-gated calcium channel dependent

mechanisms may exist for MEMRI (99, 100). MEMRI acts as a direct, quantitative reporter of

electrochemical flux and neural depolarization (40). Axonal transport from an active segment to

another inactive segment cannot explain the timing and pattern of accumulations across the brain.

Therefore, we believe that Mn(II)-enhanced signals are due to activity and not secondary to

transport or non-voltage dependent uptake. Spatiotemporal resolution of MEMRI is less granular

than some other methods. At 100 µm^3 voxels, our MEMRI has higher spatial resolution than current

BOLD imaging but still not at the level of single neurons studied with electrophysiology. MEMRI's

temporal resolution is less than electrophysiology or BOLD imaging. However, MEMRI's capacity

for repeated longitudinal brain-wide imaging across a long time span of neural activity in awake,

freely moving mice makes these tradeoffs worthwhile. Thus, Mn(II)-enhanced MR signal intensities

and statistical activation volumes are quantitative readouts of localized neural activity across the

brain. MEMRI pinpoints regions and conditions worthy of study at greater resolutions by other

techniques.

ELA is a naturalistic translationally relevant paradigm that provides an experimental platform to test

neuromodulatory interventions for human adverse childhood experiences (ACEs) and the impacts

of subsequent life-threatening events. That brain-wide destabilization exists in adult mice after ELA

provides insight into neurobiological observations of mental disorders arising after ACEs in humans
(101, 102), where similar longitudinal studies are not feasible. Regions highlighted as being
particularly affected, such as locus coeruleus, raphe nuclei, and hypothalamic nuclei, suggest
plausible neuro-chemical or -physiological targets.

In conclusion, ELA has a brain-wide impact on balance of neural activity among widely distributed
sub-regions and their dynamics across a series of experiences, from freely moving exploration in
the home cage to immediate- and post-threat conditions. We identify specific subcomponents of
this brain-wide network that are differentially activated depending upon condition in adult mice who
experienced ELA and compare these with the standard, commonly studied, adult mouse. Future
studies dissecting the functional connections suggested by these MEMRI results will provide
valuable information on coordination of coactivated regions. A next step will be to determine the
functional significance of these disparate segmental activities, how they interact in a serially
dependent manner to reconfigure future brain states, and whether these interactions contribute to
vulnerabilities for mental health disorders and/or compulsive drug use. Ultimately this longitudinal
MEMRI technique will provide insights into neurophysiological/biochemical mechanisms
contributing to the reconfiguration of brain states by ELA and inspire reparative/restorative
interventions to protect from the harm of ELA and secondary traumatic events later in adulthood.

Materials and Methods

Mice. Animal protocols were approved by Institutional Animal Care and Use Committees at both
California Institute of Technology (Caltech) and University of New Mexico. Numbers were
determined by statistical power analyses. Both Std and ELA reared groups of C57BL/6J mice
(Stock No: 000664) were bred at Jackson Labs (JAX). Adults from each group went through parallel
acute threat timelines, were handled by the same experimentalists, and imaged at Caltech (n = 24,
12 in each group), in a series of experiments performed over a period of 3.8 years. For ELA, we
followed a fragmented care paradigm (11, 12). Longitudinal MEMRI commenced at ages 11-13 wk.
This age was chosen because brain development is considered largely complete by 3 months
(103). Equal numbers of male (n = 6) and female (n = 6) ELA pups were used from 5 weaning
groups. Twelve male Std mice selected from 5 different weaning cages were subjected to the
longitudinal experimental protocol. No litter/cage effects were detected for either group.

Video Recordings and Acute Threat Exposure. We followed our previously published
longitudinal imaging protocol (32). Briefly, prior to imaging and during TMT exposure mice were
tracked in a custom arena using Noldus Ethovision (XT15) (32, 104, 105). Prior to TMT exposure,

mice were bright, alert and responsive, and displayed normal motility. Percentage of time spent in
the light side of the arena and distance travelled were calculated and statistically evaluated with
FDR correction in R v4.2.1 (*nlme* and *emmeans* packages) (106-111). Graphs were generated in
R (*ggplot2*) (112).

Longitudinal MEMRI. After capture of a non-enhanced image, mice were injected i.p. with 0.3
mmol/kg of $MnCl_2$-$4H_2O$ in sterile buffered saline and then imaged with contrast according to the
timeline previously reported (32). Between scanning sessions, video recordings, or $MnCl_2$
injections, mice were returned to their home cage. Mn(II) kinetics were followed by ROI analysis
across the timeline to test for expected intensity patterns. For optimization of MEMRI imaging
parameters, refer to our review (33).

MR Image Pre-Processing. MR images from both groups were pre-processed together in batch
according to standardized protocols (33, 50, 52). Briefly, raw MR image quality was visually
assessed and SNR quantified. Image artifacts were corrected when needed and outlier images
excluded from statistical analyses. Std and ELA pre- and post-Mn(II) images were then skull-
stripped, intensity normalized and anatomically aligned together in batch. Each processing step
was verified and validated (*SI Appendix* **Fig. S1-2**).

MR Image Analysis. *Comparison of Anatomy Between Groups.* For comparison of anatomy
between groups, we calculated Jacobian Determinants of the Deformation fields generated by the
affine-nonlinear alignment step. Average Jacobian determinant values were calculated using our
atlas-based segmentation process (see below). Two sample t-tests between Std and ELA values
were performed for each segment, FDR correction.

Measurement and Analysis of MR Intensities in ROIs. Measurements were made from pre- and
post-Mn(II)-enhanced MR images that had been skull-stripped, scaled, and aligned. Coordinates
(x/y/z) for regions of interest (3 x 3 x 3 voxels) were selected by projecting the annotated atlas
labels onto the aligned grayscale template in FSLeyes and placing the crosshairs within segments
that appeared different in SPM comparisons to pre-Mn(II) and that corresponded to previously
reported TMT-responsive regions (32, 77). Intensities beyond 3 standard deviations from the
sample average for each ROI were considered outliers and removed; none were found. Mean
intensities from ROI measurements were compared by Student's paired and two-sample t-tests, or
by Welch's test (if heteroscedastic), in R (*t.test*). Cohen's D effect sizes and FDR adjustments
(*p.adjust)* were calculated. Cohen's D for heteroscedastic data were calculated using Welch's non-
pooled standard deviation.

Statistical Parametric Mapping (SPM). For SPM, all images were smoothed to meet SPM assumptions with a 0.15 mm^3 Gaussian kernel, the minimum size allowable (57, 94). We adjusted for small baseline differences between groups in their pre-Mn(II) images by subtracting the average voxel-wise differences of intensities from each of the ELA images. After this adjustment, an unpaired t-test of pre-Mn(II) images yielded no voxels with significant differences between groups. Voxel-wise T-maps of statistically significant differences between images of different conditions within group (paired t-test) or between groups (unpaired t-test) were calculated using SPM12 (94). Statistical maps were cluster corrected (*cluster corr.*) in SPM12 with a minimum cluster size of 8 voxels (94). Fiber tracts and ventricles segments were masked from statistical maps using our *InVivo* Atlas. Setting statistical thresholds included consideration of minimal detectable effects and smallest cluster size needed to remove false positives while minimizing loss of signal.

Atlas Alignment, Segmentation, and Calculation of Fractional Volumes. Our updated *InVivo* Atlas grayscale image was inversely aligned to the dataset, and the warp field applied to the label image. This permitted automated segmentation of the images into 117 regions, including 102 neuronal clusters a.k.a. nuclei (32). Segment masks were created and applied using *fslmaths*. The number of statistically enhanced voxels were identified as those from SPM paired or unpaired t-tests exceeding thresholds set according to our criteria described above. The number of total voxels within each segment were counted using *fslstats* (24). The fraction of enhanced voxels from SPM paired (FAV) or unpaired (FDV) t-tests to total voxels within each segment were calculated in R and visualized as column graphs and line plots using *ggplot2* (111-113). Segments were categorized into patterns of FDV dynamics by the criteria of FDV > 0.1 for at least one condition and |ΔFDV| > 0.1 between conditions.

Immunohistochemistry, c-fos Quantification and Correlation to MEMRI. Brains from mice, euthanized at the conclusion of MR imaging (5 Std and 6 ELA), were prepared for histology according to our protocol (32, 114). We selected 1-3 sections also identifiable in MR slices from each mouse that had regions with both very high and very low numbers of c-fos stained nuclei. Nuclei were counted in a 3 x 3 grid of 100 μm^2 squares from 1-3 x 35 μm thick sections of each brain, and MR intensities (SI) from each condition measured from 100 μm^3 voxels in corresponding slices. Counts ranged from 0 to 27. MEMRI SI ranged from ~6,900-10,100. Counts and SI were visualized in quartile boxplots with R (*ggplot2*). A square root transformation linearized c-fos+ count data for determination of effect size and statistical significance. Regions counted in the same section or measured in the same slice were compared statistically by linear mixed models and by post-hoc paired t-tests with FDR correction. Regions were compared between groups for both

537 modalities by linear mixed models and post-hoc two-sample t-tests with FDR correction. Counts
538 and SI of each group were averaged by region and individual and then correlated, with p-value
539 determined by permutation testing.
540
541 Please see *SI Appendix* **Extended Methods** for additional methodological details.
542

Acknowledgments
We gratefully acknowledge support from the Biological Imaging Facility/Center and the Beckman
Institute at Caltech where imaging was performed, and funding from NIH R01MH096093 (ELB),
NIH NIA P20AG068077 (ELB), NIH NINDS F99NS139535 (TWU), Zilkha Neurogenetic Institute
(REJ) and Harvey Family Endowment (ELB). We appreciate staff, faculty and trainees at UNM,
USC and Caltech including Christopher S. Medina, Daniel Barto, Xiaowei Zhang and Sharon Wu
Lin (see *SI Appendix)*. We thank Chiara Padilla for proofreading support. We are grateful to Harry
B. Gray, Arnold O. Beckman Professor of Chemistry, at Caltech for insights on Mn(II)
metallobiology and advice on the manuscript. See *SI Appendix* **Additional Acknowledgements**.

 Figures and Tables

Figure 1.

Fig. 1. Longitudinal imaging procedures. *(A)* Schematic of the overall timeline (open arrows, top).

Boxes diagram events. *Left box*: ELA paradigm; *Middle box*: Home Cage and Acute Threat, timeline

of MR and video capture: Baseline, BL; Home Cage, HC; Acute Threat, TMT. *Right box*: Day 8

Baseline, D8-BL; Day 9 Post-Threat, D9-H. *(B-D)* Video tracking results: *(B)* Percent time spent in

the light (quartile boxplots, individuals indicated by dot plot overlays) (cyan, Std; yellow, ELA; BL

and TMT as per schematic). *(C)* Distance traveled after imaging during TMT exposure. Line graphs

of distance travelled per 1-min epochs with error bars representing standard error of the mean.

Repeated measures ANOVA with fixed effects of group and condition. For (B-C), paired and

unpaired t-tests with FDR correction for multiple comparisons; Cohen's D effect sizes (d); * $p <$

0.05, n = 22 (10 Std, 12 ELA). *(D)* Representative tracking of one mouse from each group, Std and

ELA, at BL (gray) and TMT (red). Shaded areas represent dark side of the arena. *(E)* Overview of

MR data processing and analysis used in this study (also see ***SI Appendix* Fig. S1-2**) (33, 50, 52).

Figure 2.

Fig. 2. Statistical parametric maps of MEMRI in ELA mice change as they progress from home cage to acute threat to post threat. *(A-C)* Shown are statistical maps of voxel-wise paired t-tests of post-Mn(II) images of ELA mice at each successive condition compared to their pre-Mn(II) images. Maps are overlaid on the grayscale dataset average of all pre-Mn(II) images (coronal slices, top rows; sagittal, bottom rows). Slice position is indicated on the grayscale image (top left, white lines). White arrows indicate notable differences between conditions. MOB, main olfactory bulb; LS, lateral septum; BST, bed nuclei of the stria terminalis; PVT, paraventricular thalamus; HIP, hippocampus; LC, locus coeruleus; CEA, central amygdala; and CB, cerebellum. For t-tests, n = 12, p < 0.01, T = 2.72, cluster corr. For comparisons between conditions of ELA mice see *SI Appendix,* **Fig. S3**.

Figure 3.

Fig. 3. Measurements of MEMRI signal intensities in candidate regions of interest in systems
known to be involved in predator odor-type threat responses. (A) Top: Representative slices from
two voxel-wise statistical maps (ELA HC & ELA TMT) of post-Mn(II) > pre-Mn(II) superimposed
onto the grayscale InVivo atlas. SPM paired t-tests: p < 0.01 cluster corr. ROI locations selected
where TMT signal did not overlap with HC (yellow crosshairs). Bottom: Quartile boxplots of signal
intensity as a percent over pre-Mn(II) measured in each ROI. Error bars: Mean ± standard deviation.
Effect sizes of the differences in signal between conditions at each ROI (Cohen's D, d), average
signal intensity differences (ΔSI) between the two conditions, and standard deviations of ΔSI are
indicated. Paired t-test, * p < 0.05, ** p < 0.01, FDR. dlOB-DII, dorsal lateral DII olfactory
subdomain; vlAON, ventrolateral anterior olfactory nucleus; aOC, anterior olfactory cortex; and
aCEA, anterior central amygdala. (B) Top: Similar comparison but between groups at one condition,
HC. Slices of paired t-tests of HC greater than pre-Mn(II) from either Std or ELA are superimposed
on the grayscale atlas. Crosshairs indicate ROI locations, selected where ELA signal did not
overlap with Std. Bottom: Quartile boxplots of unpaired t-tests of ROI measurements. Error bars:
Mean ± standard deviation. Welch's t-test and non-pooled variance-based d* and s* were used for
mOB-DII. * p < 0.05, FDR. mOB-DII, medial DII olfactory subdomain; dmNDB, dorsomedial

diagonal band nucleus; aLS, anterior lateral septum; dmACB, dorsomedial nucleus accumbens.

See ROI coordinates and summary statistics in *SI Appendix,* **Tables S2-4**.

Figure 4.

Fig. 4. MEMRI and c-fos staining correlate. (*A*) Coronal slices of the average intensities of day-9

post-Mn(II) images (color code, bottom) (n = 6). Minimum (300 a.u.) based average noise and

maximum (11,000 a.u.) from intensity histograms. Small white boxes indicate regions (3 x 3 x 1

voxels) where signal intensities (SI) were measured. HYP, hypothalamus; THA, thalamus; CS,

superior central raphe; MB, midbrain. Mag bar = 500 µm. (*B*) Two representative histological

sections from one of the mice contributing to *(A)* stained for c-fos corresponding to anatomy of

slices in *(A)* (large white boxes). Purple boxes indicate locations of 3 x 3 grid of 100 µm^2 squares

where numbers of c-fos+ nuclei were counted. Mag bar = 500 µm. See ***SI Appendix*** **Fig. S5A-B**

for corresponding images from Std. *(C)* Statistical comparisons of MEMRI SI in four locations for

both groups. Shown are quartile boxplots with dot plot overlays of MEMRI SI from ELA (yellow, left)

and Std (cyan, right). Paired t-tests were performed between regions within the same slice. Note

large Cohen's D effect sizes (d) for paired t-tests between active and inactive regions. For evidence

of test-retest reliability of MEMRI SI, see ***SI Appendix*** **Fig. S4**. *(D)* c-fos counts paralleled MEMRI

SI with high or low numbers of c-fos+ nuclei. Shown are quartile boxplots with dot plot overlay of

raw c-fos+ counts from ELA (purple, left) and Std (pink, right). A square root transformation

linearized c-fos+ counts for statistics. Note similar large Cohen's D effect sizes (d) between active

and inactive regions. *(E)* Strong Pearson correlations between MEMRI SI and numbers of c-fos+

nuclei (linearized) when paired by individual. Scatter plots with linear regressions (black lines) of

MEMRI SI (vertical axes) by c-fos+ counts (horizontal axes). Points represent the average values

for each of the 4 regions across 5 Std (*top*, *cyan*) and 6 ELA (*bottom*, *yellow*) brains. Note strong

correlations in both groups (r$_{Std}$ = 0.636, r$_{ELA}$ = 0.582). For paired t-tests in *(C-D)*, *** p < 0.001,

FDR corrected. For *(E)*, p < 0.001 by permutation testing and bootstrapping. Also see ***SI Appendix***

Fig. S5C-D.

Figure 5.

Fig. 5. Statistical maps show progressive modifications of activity across sequential conditions. *(A-*
D) Voxel-wise paired t-tests of post-Mn(II) compared to pre-Mn(II) images at two conditions: in ELA
mice, *(A)* HC (blue) and TMT (red), *(B)* TMT (red) and D9 (green), and *(C)* HC (blue) and D9
(green); and in Std mice, *(D)* HC (blue) and D9 (green). *Top*: Pairs of statistical maps were projected
onto the grayscale *InVivo* Atlas with segments outlined to show the anatomical overlaps between
conditions. *Bottom*: Corresponding column graphs show segment-wise fractional activation
volumes (FAV, vertical axes) quantified from segmentations of statistical maps. Anatomical
domains (color-coded on horizontal axes) are divided into 102 segments. Arrows point to examples
of segments with FAV alterations between conditions and color-coded to indicate direction of
change. For paired t-tests, $p < 0.01$, cluster *corr.* for multiple comparisons. ELA, $T_{(11)} = 2.72$, d $\geq$
0.79, n = 12; Std, $T_{(10)} = 2.76$, d $\geq$ 0.83, n = 11. See *SI Appendix* **Table S5** for list of domains and
segments, and **Table S6A-B** for FAV values.

Figure 6.

Fig. 6. ELA brain states differ from Std at every condition. *(A-C)* Fractional Difference Volumes
(FDVs) from segmentation of unpaired SPM t-tests between signal intensities of ELA and Std at
(A) HC, *(B)* TMT and *(C)* D9 conditions. All ELA images were adjusted for the voxel-wise average
differences between groups' pre-Mn(II) images. FDV color-code represents where ELA signal
intensities were heightened (yellow) or diminished (blue) relative to Std. Arrows in column graphs
point to example segments with notable FDV dynamics as in *(D)*: blue, MS and VMH; red, CEA
and CS; green, PA and DMH; black, DP and ARH; and purple, LC. *(D)* Shown are examples of
various dynamics in FDV in 9 segments, which were highlighted by arrows in column graphs *(A-*
C). Line graphs show FDVs (vertical axes) of each segment across three successive conditions
(horizontal axes). Graphs are grouped and color-coded according to the dynamical pattern of ELA
and Std FDV values. *(A-D)* Unpaired t-tests, p < 0.05, cluster corr.; HC: $T_{(21)}$ = 1.72, n_{Std} = 11, n_{ELA}
= 12; TMT and D9: $T_{(22)}$ = 1.72, n = 12 per group. *(E)* Three schematics summarizing differences in
activity between ELA and Std across the 10 segmented anatomical domains at each condition: HC
(*top*), TMT (*middle*) and D9 (*bottom*). Each color gradient corresponds to magnitudes and direction
of segmental fractional difference volumes (FDV), and proportion of coloring indicating proportion
of segments following that pattern within each domain; See **SI Appendix Movie 1** for a slice-by-
slice walkthrough of between group SPMs at each condition projected on brain anatomy, **Table S8**
for FDV values and **Table S6** for list of domains and segments.

References

1. Felitti VJ*, et al.* (1998) Relationship of Childhood Abuse and Household Dysfunction to Many of the Leading Causes of Death in Adults. *American Journal of Preventive Medicine* 14(4):245-258.

2. Merrick M*, et al.* (2019) Vital Signs: Estimated Proportion of Adult Health Problems Attributable to Adverse Childhood Experiences and Implications for Prevention — 25 States, 2015–2017. *MMWR. Morbidity and Mortality Weekly Report* 68.

3. Zarse EM*, et al.* (2019) The adverse childhood experiences questionnaire: Two decades of research on childhood trauma as a primary cause of adult mental illness, addiction, and medical diseases. *Cogent Medicine* 6(1):1581447.

4. Spadoni AD*, et al.* (2022) Contribution of early-life unpredictability to neuropsychiatric symptom patterns in adulthood. *Depression and Anxiety* 39(10-11):706-717.

5. Yehuda R*, et al.* (2010) Putative biological mechanisms for the association between early life adversity and the subsequent development of PTSD. *Psychopharmacology* 212(3):405-417.

6. Chen Y & Baram TZ (2016) Toward Understanding How Early-Life Stress Reprograms Cognitive and Emotional Brain Networks. *Neuropsychopharmacology* 41(1):197-206.

7. Kraaijenvanger EJ*, et al.* (2020) Impact of early life adversities on human brain functioning: A coordinate-based meta-analysis. *Neuroscience & Biobehavioral Reviews* 113:62-76.

8. Malave L, Van Dijk MT, & Anacker C (2022) Early life adversity shapes neural circuit function during sensitive postnatal developmental periods. *Translational Psychiatry* 12(1):306.

9. Garcia-Arocena D (2015) More researchers are using B6J mice than ever before. (The Jackson Laboratory).

10. Laboratory TJ (2024) Mouse Room Conditions.

11. Molet J, Maras PM, Avishai-Eliner S, & Baram TZ (2014) Naturalistic rodent models of chronic early-life stress. *Developmental Psychobiology* 56(8):1675-1688.

12. Rice CJ, Sandman CA, Lenjavi MR, & Baram TZ (2008) A Novel Mouse Model for Acute and Long-Lasting Consequences of Early Life Stress. *Endocrinology* 149(10):4892-4900.

13. Bercum FM, Navarro Gomez MJ, & Saddoris MP (2021) Elevated fear responses to threatening cues in rats with early life stress is associated with greater excitability and loss of gamma oscillations in ventral-medial prefrontal cortex. *Neurobiol Learn Mem* 185:107541.

14.   Bolton JL, *et al.* (2018) Anhedonia Following Early-Life Adversity Involves Aberrant
Interaction of Reward and Anxiety Circuits and Is Reversed by Partial Silencing of
Amygdala Corticotropin-Releasing Hormone Gene. *Biological Psychiatry* 83(2):137-147.
15.   Bolton JL, *et al.* (2022) Early stress-induced impaired microglial pruning of excitatory
synapses on immature CRH-expressing neurons provokes aberrant adult stress
responses. *Cell Rep* 38(13):110600.
16.   Brosens N, Simon C, Kessels HW, Lucassen PJ, & Krugers HJ (2023) Early life stress
lastingly alters the function and AMPA-receptor composition of glutamatergic synapses in
the hippocampus of male mice. *J Neuroendocrinol* 35(12):e13346.
17.   de Carvalho G, Khoja S, Haile MT, & Chen LY (2023) Early life adversity impaired dorsal
striatal synaptic transmission and behavioral adaptability to appropriate action selection
in a sex-dependent manner. *Front Synaptic Neurosci* 15:1128640.
18.   Guadagno A, *et al.* (2018) Reduced resting-state functional connectivity of the basolateral
amygdala to the medial prefrontal cortex in preweaning rats exposed to chronic early-life
stress. *Brain Struct Funct* 223(8):3711-3729.
19.   Guadagno A, Verlezza S, Long H, Wong TP, & Walker CD (2020) It Is All in the Right
Amygdala: Increased Synaptic Plasticity and Perineuronal Nets in Male, But Not Female,
Juvenile Rat Pups after Exposure to Early-Life Stress. *The Journal of neuroscience : the*
official journal of the Society for Neuroscience 40(43):8276-8291.
20.   Guadagno A, Wong TP, & Walker CD (2018) Morphological and functional changes in
the preweaning basolateral amygdala induced by early chronic stress associate with
anxiety and fear behavior in adult male, but not female rats. *Prog Neuropsychopharmacol*
Biol Psychiatry 81:25-37.
21.   Gunn BG, *et al.* (2013) Dysfunctional astrocytic and synaptic regulation of hypothalamic
glutamatergic transmission in a mouse model of early-life adversity: relevance to
neurosteroids and programming of the stress response. *The Journal of neuroscience :*
the official journal of the Society for Neuroscience 33(50):19534-19554.
22.   Johnson FK, *et al.* (2018) Amygdala hyper-connectivity in a mouse model of
unpredictable early life stress. *Transl Psychiatry* 8(1):49.
23.   Kooiker CL, Chen Y, Birnie MT, & Baram TZ (2023) Genetic Tagging Uncovers a Robust,
Selective Activation of the Thalamic Paraventricular Nucleus by Adverse Experiences
Early in Life. *Biol Psychiatry Glob Open Sci* 3(4):746-755.
24.   Levis SC, *et al.* (2022) Enduring disruption of reward and stress circuit activities by early-
life adversity in male rats. *Transl Psychiatry* 12(1):251.

25.   Malter Cohen M*, et al.* (2013) Early-life stress has persistent effects on amygdala
733   function and development in mice and humans. *Proc Natl Acad Sci U S A* 110(45):18274-
734   18278.

735   26.   Raineki C, Cortes MR, Belnoue L, & Sullivan RM (2012) Effects of early-life abuse differ
736   across development: infant social behavior deficits are followed by adolescent
737   depressive-like behaviors mediated by the amygdala. *The Journal of neuroscience : the*
738   *official journal of the Society for Neuroscience* 32(22):7758-7765.

739   27.   Yan CG*, et al.* (2017) Aberrant development of intrinsic brain activity in a rat model of
740   caregiver maltreatment of offspring. *Transl Psychiatry* 7(1):e1005.

741   28.   Holschneider DP, Guo Y, Mayer EA, & Wang Z (2016) Early life stress elicits visceral
742   hyperalgesia and functional reorganization of pain circuits in adult rats. *Neurobiol Stress*
743   3:8-22.

744   29.   Islam R*, et al.* (2024) Early adversity causes sex-specific deficits in perforant pathway
745   connectivity and contextual memory in adolescent mice. *Biol Sex Differ* 15(1):39.

746   30.   White JD*, et al.* (2020) Early life stress causes sex-specific changes in adult fronto-limbic
747   connectivity that differentially drive learning. *Elife* 9.

748   31.   Kooiker CL*, et al.* (2024) Paraventricular Thalamus Neuronal Ensembles Encode Early-
749   life Adversity and Mediate the Consequent Sex-dependent Disruptions of Adult Reward
750   Behaviors. (Cold Spring Harbor Laboratory).

751   32.   Uselman TW, Barto DR, Jacobs RE, & Bearer EL (2020) Evolution of brain-wide activity
752   in the awake behaving mouse after acute fear by longitudinal manganese-enhanced MRI.
753   *Neuroimage* 222:116975.

754   33.   Uselman TW, Medina CS, Gray HB, Jacobs RE, & Bearer EL (2022) Longitudinal
755   manganese-enhanced magnetic resonance imaging of neural projections and
756   activity.e4675.

757   34.   Vousden DA*, et al.* (2018) Continuous manganese delivery via osmotic pumps for
758   manganese-enhanced mouse MRI does not impair spatial learning but leads to skin
759   ulceration. *NeuroImage* 173:411-420.

760   35.   Aoki I, Wu YJ, Silva AC, Lynch RM, & Koretsky AP (2004) In vivo detection of
761   neuroarchitecture in the rodent brain using manganese-enhanced MRI. *Neuroimage*
762   22(3):1046-1059.

763   36.   Bedenk BT*, et al.* (2018) Mn(2+) dynamics in manganese-enhanced MRI (MEMRI):
764   Cav1.2 channel-mediated uptake and preferential accumulation in projection terminals.
765   *Neuroimage* 169:374-382.

766   37.   Narita K, Kawasaki F, & Kita H (1990) Mn and Mg influxes through Ca channels of motor
767   nerve terminals are prevented by verapamil in frogs. *Brain research* 510(2):289-295.

38.   Hattori S*, et al.* (2013) In vivo evaluation of cellular activity in alphaCaMKII heterozygous
knockout mice using manganese-enhanced magnetic resonance imaging (MEMRI). *Front*
Integr Neurosci 7:76.

39.   Lin YJ & Koretsky AP (1997) Manganese ion enhances T1-weighted MRI during brain
activation: an approach to direct imaging of brain function. *Magnetic resonance in*
medicine 38(3):378-388.

40.   Svehla P, Bedecarrats A, Jahn C, Nargeot R, & Ciobanu L (2018) Intracellular
manganese enhanced MRI signals reflect the frequency of action potentials in Aplysia
neurons. *J Neurosci Methods* 295:121-128.

41.   Malkova NV, Gallagher JJ, Yu CZ, Jacobs RE, & Patterson PH (2014) Manganese-
enhanced magnetic resonance imaging reveals increased DOI-induced brain activity in a
mouse model of schizophrenia. *Proc Natl Acad Sci U S A* 111(24):E2492-2500.

42.   Yu X, Sanes DH, Aristizabal O, Wadghiri YZ, & Turnbull DH (2007) Large-scale
reorganization of the tonotopic map in mouse auditory midbrain revealed by MRI.
Proceedings of the National Academy of Sciences 104(29):12193-12198.

43.   Yu X, Wadghiri YZ, Sanes DH, & Turnbull DH (2005) In vivo auditory brain mapping in
mice with Mn-enhanced MRI. *Nature Neuroscience* 8(7):961-968.

44.   Bearer EL*, et al.* (2007) Role of neuronal activity and kinesin on tract tracing by
manganese-enhanced MRI (MEMRI). *Neuroimage* 37 Suppl 1:S37-46.

45.   Bearer EL, Zhang X, & Jacobs RE (2007) Live imaging of neuronal connections by
magnetic resonance: Robust transport in the hippocampal–septal memory circuit in a
mouse model of Down syndrome. *NeuroImage* 37(1):230-242.

46.   Bearer EL, Zhang X, Janvelyan D, Boulat B, & Jacobs RE (2009) Reward circuitry is
perturbed in the absence of the serotonin transporter. *Neuroimage* 46(4):1091-1104.

47.   Zhang X*, et al.* (2010) Altered neurocircuitry in the dopamine transporter knockout mouse
brain. *PLoS One* 5(7):e11506.

48.   Gallagher JJ, Zhang X, Ziomek GJ, Jacobs RE, & Bearer EL (2012) Deficits in axonal
transport in hippocampal-based circuitry and the visual pathway in APP knock-out
animals witnessed by manganese enhanced MRI. *Neuroimage* 60(3):1856-1866.

49.   Gallagher JJ*, et al.* (2013) Altered reward circuitry in the norepinephrine transporter
knockout mouse. *PLoS One* 8(3):e57597.

50.   Delora A*, et al.* (2016) A simple rapid process for semi-automated brain extraction from
magnetic resonance images of the whole mouse head. *J Neurosci Methods* 257:185-193.

51.   Medina CS*, et al.* (2017) Hippocampal to basal forebrain transport of Mn(2+) is impaired
by deletion of KLC1, a subunit of the conventional kinesin microtubule-based motor.
Neuroimage 145(Pt A):44-57.

52.   Medina CS, Manifold-Wheeler B, Gonzales A, & Bearer EL (2017) Automated
805   Computational Processing of 3-D MR Images of Mouse Brain for Phenotyping of Living
806   Animals. *Curr Protoc Mol Biol* 119:29A 25 21-29A 25 38.

807   53.   Bearer EL*, et al.* (2018) Alterations of functional circuitry in aging brain and the impact of
808   mutated APP expression. *Neurobiol Aging* 70:276-290.

809   54.   Bearer EL, Barto D, & Jacobs RE (2019) Imaging the evolution acute fear: Longitudinal
810   whole brain imaging in living mice of neural activity with MEMRI. *Proceedings of the*
811   *International Society for Magnetic Resonance in Medicine... Scientific Meeting and*
812   *Exhibition. International Society for Magnetic Resonance in Medicine. Scientific Meeting*
813   *and Exhibition*, (NIH Public Access).

814   55.   Medina CS*, et al.* (2019) Decoupling the Effects of the Amyloid Precursor Protein From
815   Amyloid-beta Plaques on Axonal Transport Dynamics in the Living Brain. *Front Cell*
816   *Neurosci* 13:501.

817   56.   Phinyomark A, Ibanez-Marcelo E, & Petri G (2017) Resting-State fMRI Functional
818   Connectivity: Big Data Preprocessing Pipelines and Topological Data Analysis. *IEEE*
819   *Transactions on Big Data* 3(4):415-428.

820   57.   Flandin G & Friston K (2008) Statistical parametric mapping (SPM). *Scholarpedia*
821   3(4):6232.

822   58.   Penny WD, Friston KJ, Ashburner JT, Kiebel SJ, & Nichols TE (2011) *Statistical*
823   *Parametric Mapping: The Analysis of Functional Brain Images* (Elsevier Science).

824   59.   Pallast N*, et al.* (2019) Processing Pipeline for Atlas-Based Imaging Data Analysis of
825   Structural and Functional Mouse Brain MRI (AIDAmri). *Frontiers in Neuroinformatics* 13.

826   60.   Fendt M, Endres T, Lowry CA, Apfelbach R, & McGregor IS (2005) TMT-induced
827   autonomic and behavioral changes and the neural basis of its processing. *Neuroscience*
828   *and biobehavioral reviews* 29(8):1145-1156.

829   61.   Rosen J, Asok A, & Chakraborty T (2015) The smell of fear: innate threat of 2,5-dihydro-
830   2,4,5-trimethylthiazoline, a single molecule component of a predator odor. 9.

831   62.   Gallo M*, et al.* (2019) Limited Bedding and Nesting Induces Maternal Behavior
832   Resembling Both Hypervigilance and Abuse. 13.

833   63.   Matsukawa M, Yoshikawa M, Katsuyama N, Aizawa S, & Sato T (2022) The Anterior
834   Piriform Cortex and Predator Odor Responses: Modulation by Inhibitory Circuits. *Front*
835   *Behav Neurosci* 16:896525.

836   64.   Matsumoto H*, et al.* (2010) Spatial Arrangement of Glomerular Molecular-Feature
837   Clusters in the Odorant-Receptor Class Domains of the Mouse Olfactory Bulb.
838   103(6):3490-3500.

839   65.   Cohen J (2013) Statistical Power Analysis for the Behavioral Sciences.

840   66.   Kimbrough A*, et al.* (2020) Brain-wide functional architecture remodeling by alcohol
841   dependence and abstinence. 117(4):2149-2159.

842   67.   Motta SC*, et al.* (2009) Dissecting the brain's fear system reveals the hypothalamus is
843   critical for responding in subordinate conspecific intruders. *Proceedings of the National*
844   *Academy of Sciences* 106(12):4870-4875.

845   68.   Greene AS, Horien C, Barson D, Scheinost D, & Constable RT (2023) Why is everyone
846   talking about brain state? *Trends Neurosci* 46(7):508-524.

847   69.   Ben-Shaul Y, Katz LC, Mooney R, & Dulac C (2010) In vivo vomeronasal stimulation
848   reveals sensory encoding of conspecific and allospecific cues by the mouse accessory
849   olfactory bulb. 107(11):5172-5177.

850   70.   Canteras NS (2002) The medial hypothalamic defensive system: Hodological
851   organization and functional implications. *Pharmacology Biochemistry and Behavior*
852   71(3):481-491.

853   71.   Canteras NS, Chiavegatto S, Ribeiro do Valle LE, & Swanson LW (1997) Severe
854   Reduction of Rat Defensive Behavior to a Predator by Discrete Hypothalamic Chemical
855   Lesions. *Brain Research Bulletin* 44(3):297-305.

856   72.   Janitzky K, D'Hanis W, Kröber A, & Schwegler H (2015) TMT predator odor activated
857   neural circuit in C57BL/6J mice indicates TMT-stress as a suitable model for
858   uncontrollable intense stress. *Brain research* 1599:1-8.

859   73.   Kunwar PS*, et al.* (2015) Ventromedial hypothalamic neurons control a defensive
860   emotion state. *eLife* 4:e06633.

861   74.   Pérez-Gómez A*, et al.* (2015) Innate Predator Odor Aversion Driven by Parallel Olfactory
862   Subsystems that Converge in the Ventromedial Hypothalamus. *Current Biology*
863   25(10):1340-1346.

864   75.   Koob GF & Volkow ND (2016) Neurobiology of addiction: a neurocircuitry analysis. *The*
865   *lancet. Psychiatry* 3(8):760-773.

866   76.   Russo SJ & Nestler EJ (2013) The brain reward circuitry in mood disorders. *Nature*
867   *Reviews Neuroscience* 14(9):609-625.

868   77.   Silva BA, Gross CT, & Graff J (2016) The neural circuits of innate fear: detection,
869   integration, action, and memorization. *Learn Mem* 23(10):544-555.

870   78.   Coletta L*, et al.* (2020) Network structure of the mouse brain connectome with voxel
871   resolution. 6(51):eabb7187.

872   79.   Salvan P*, et al.* (2023) Serotonin regulation of behavior via large-scale neuromodulation
873   of serotonin receptor networks. *Nature Neuroscience* 26(1):53-63.

874   80.   Taylor NL*, et al.* (2022) Structural connections between the noradrenergic and cholinergic
875   system shape the dynamics of functional brain networks. *NeuroImage* 260:119455.

81. Zerbi V*, et al.* (2019) Rapid Reconfiguration of the Functional Connectome after
Chemogenetic Locus Coeruleus Activation. *Neuron* 103(4):702-718.e705.

82. David & Adolphs R (2014) A Framework for Studying Emotions across Species. *Cell*
157(1):187-200.

83. Flavell SW, Gogolla N, Lovett-Barron M, & Zelikowsky M (2022) The emergence and
influence of internal states. *Neuron* 110(16):2545-2570.

84. Jaric I*, et al.* (2022) The rearing environment persistently modulates mouse phenotypes
from the molecular to the behavioural level. *PLoS Biol* 20(10):e3001837.

85. Jaric I*, et al.* (2024) Using mice from different breeding sites fails to improve replicability
of results from single-laboratory studies. *Lab Anim (NY)* 53(1):18-22.

86. Usui T, Macleod MR, McCann SK, Senior AM, & Nakagawa S (2021) Meta-analysis of
variation suggests that embracing variability improves both replicability and
generalizability in preclinical research. *PLoS Biol* 19(5):e3001009.

87. Chang C-H & Gean P-W (2019) The Ventral Hippocampus Controls Stress-Provoked
Impulsive Aggression through the Ventromedial Hypothalamus in Post-Weaning Social
Isolation Mice. *Cell Reports* 28(5):1195-1205.e1193.

88. Manzano Nieves G, Bravo M, & Bath KG (2023) Early life adversity ablates sex
differences in active versus passive threat responding in mice. *Stress*:1-32.

89. Spring S, Lerch JP, & Henkelman RM (2007) Sexual dimorphism revealed in the
structure of the mouse brain using three-dimensional magnetic resonance imaging.
NeuroImage 35(4):1424-1433.

90. Bruce MR*, et al.* (2021) Sexually dimorphic neuroanatomical differences relate to ASD-
relevant behavioral outcomes in a maternal autoantibody mouse model. *Molecular*
Psychiatry 26(12):7530-7537.

91. Holz NE*, et al.* (2023) A stable and replicable neural signature of lifespan adversity in the
adult brain. *Nature Neuroscience*.

92. Lerch JP*, et al.* (2011) Maze training in mice induces MRI-detectable brain shape
changes specific to the type of learning. *NeuroImage* 54(3):2086-2095.

93. Ashburner J*, et al.* (2012) SPM8 manual.

94. Ashburner J*, et al.* (2014) SPM12 manual.

95. Carter CS, Lesh TA, & Barch DM (2016) Thresholds, Power, and Sample Sizes in
Clinical Neuroimaging. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*
1(2):99-100.

96. Mumford JA (2012) A power calculation guide for fMRI studies. *Social Cognitive and*
Affective Neuroscience 7(6):738-742.

97.   Vandekar SN & Stephens J (2021) Improving the replicability of neuroimaging findings by
912   thresholding effect sizes instead of p-values. 42(8):2393-2398.

913   98.   Lehallier B*, et al.* (2012) Brain Processing of Biologically Relevant Odors in the Awake
914   Rat, as Revealed by Manganese-Enhanced MRI. *PLoS ONE* 7(10):e48491.

915   99.   Petrus E, Saar G, Daoust A, Dodd S, & Koretsky AP (2021) A hierarchy of manganese
916   competition and entry in organotypic hippocampal slice cultures. *NMR in Biomedicine*
917   34(4).

918   100.   Bartelle BB, Szulc KU, Suero-Abreu GA, Rodriguez JJ, & Turnbull DH (2013) Divalent
919   metal transporter, DMT1: A novel MRI reporter protein. *Magnetic resonance in medicine*
920   70(3):842-850.

921   101.   Duffy KA, McLaughlin KA, & Green PA (2018) Early life adversity and health-risk
922   behaviors: proposed psychological and neural mechanisms. 1428(1):151-169.

923   102.   Hakamata Y, Suzuki Y, Kobashikawa H, & Hori H (2022) Neurobiology of early life
924   adversity: A systematic review of meta-analyses towards an integrative account of its
925   neurobiological trajectories to mental disorders. *Frontiers in Neuroendocrinology*
926   65:100994.

927   103.   Hammelrath L*, et al.* (2016) Morphological maturation of the mouse brain: An in vivo MRI
928   and histology investigation. *NeuroImage* 125:144-152.

929   104.   Brigman JL, Mathur P, Lu L, Williams RW, & Holmes A (2009) Genetic relationship
930   between anxiety-related and fear-related behaviors in BXD recombinant inbred mice.
931   20(2):204-209.

932   105.   Noldus LPJJ, Spink AJ, & Tegelenbosch RAJ (2001) EthoVision: A versatile video
933   tracking system for automation of behavioral experiments. *Behavior Research Methods,*
934   *Instruments, & Computers* 33(3):398-414.

935   106.   Fox J & Weisberg S (2019) An R Companion to Applied Regression.

936   107.   Lenth RV (2022) emmeans: Estimated Marginal Means, aka Least-Squares Means.

937   108.   Pinheiro J, Bates D, & Team RC (2022) nlme: Linear and Nonlinear Mixed Effects
938   Models.

109.   Pinheiro JC & Bates DM (2000) Mixed-Effects Models in S and S-PLUS.

110.   Posit t (2022) RStudio: Integrated Development Environment for R.

111.   Team RC (2022) R: A Language and Environment for Statistical Computing.

112.   Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis.

113.   Wickham H*, et al.* (2019) Welcome to the Tidyverse. 4(43):1686.

114.   Tyszka JM, Readhead C, Bearer EL, Pautler RG, & Jacobs RE (2006) Statistical diffusion
945   tensor histology reveals regional dysmyelination effects in the shiverer mouse mutant.
946   *NeuroImage* 29(4):1058-1065.