Human Pannexin 1 Channel is NOT Phosphorylated by Src Tyrosine Kinase at Tyr199 and Tyr309

Zheng Ruan[1#], Junuk Lee[1], Yangyang Li[1], Juan Du[1]*, Wei Lü[1]*

1. Department of Structural Biology, Van Andel Institute, Grand Rapids, MI 49503, USA

* To whom correspondence should be addressed
# Current Affiliation: Department of Biochemistry & Molecular Biology, Thomas Jefferson University, Philadelphia, PA 19107, USA

For editorial correspondence:
Juan Du, phone: +1 (616) 234-5358; E-mail: juan.du@vai.org;
Wei Lü, phone: +1 (616) 234-5022; E-mail: wei.lu@vai.org.

Keywords: PANX1, tyrosine phosphorylation, ion channel, Y198, Y308

ABSTRACT

Protein phosphorylation is one of the major molecular mechanisms regulating protein activity and function throughout the cell. Pannexin 1 (PANX1) is a large-pore channel permeable to ATP and other cellular metabolites. Its tyrosine phosphorylation and subsequent activation have been found to play critical roles in diverse cellular conditions, including neuronal cell death, acute inflammation, and smooth muscle contraction. Specifically, the non-receptor kinase Src has been reported to phosphorylate Tyr198 and Tyr308 of mouse PANX1 (equivalent to Tyr199 and Tyr309 of human PANX1), resulting in channel opening and ATP release. Although the Src-dependent PANX1 activation mechanism has been widely discussed in the literature, independent validation of the tyrosine phosphorylation of PANX1 has been lacking. Here, we show that commercially available antibodies against the two phosphorylation sites mentioned above—which were used to identify endogenous PANX1 phosphorylation at these two sites— are nonspecific and should not be used to interpret results related to PANX1 phosphorylation. We further provide evidence that neither tyrosine residue is a major phosphorylation site for Src kinase in heterologous expression systems. We call on the field to re-examine the existing paradigm of tyrosine phosphorylation-dependent activation of the PANX1 channel.

INTRODUCTION

PANX1 is a large-pore channel widely associated with many physiological and pathological processes, including immune response, inflammation, and vasoconstriction(1–3). The activation of PANX1 allows the release of ATP and other cellular metabolites to the extracellular space, serving as signaling molecules for neighboring cells(4, 5). While the functional relevance of PANX1 in human physiology and diseases has been investigated for nearly two decades since the PANX family was cloned(6–8), the detailed structural and biophysical properties of the channel were only beginning to be elucidated. In 2020, we and others determined structures of PANX1 by single-particle cryo-electron microscopy (cryo-EM), revealing the channel stoichiometry, domain architecture, ion permeation pathways, and inhibition mechanisms(9–14). These structures have in turn motivated small-molecule docking analysis and the development of the first PANX1-selective channel blockers(15). Despite the remarkable progress made in the PANX1 field, the dynamic regulation of the channel opening and closing in various cellular contexts is not well understood. Which cellular stimuli PANX1 responds to and how they achieve channel activation, especially in a reversible manner, are unclear and largely under debate(16, 17). Deciphering the regulatory principle of the PANX1 is not only important to better characterize the function of the channel, but may also provide clues to the development of pharmaceutical agents to treat PANX1-associated diseases.

Post-translational modifications, such as protease cleavage, glycosylation, phosphorylation, and S-nitrosylation, have been shown to play important roles in PANX1 regulation(18). For example, caspase-dependent cleavage of the regulatory C-terminal tail of PANX1 is critically implicated for apoptotic cell clearance by the immune system(4, 5). The glycosylation at the second extracellular loop of PANX1 sterically prevents the docking of two PANX1 heptamer into a gap-junction-like assembly(19–21). Our recent cryo-EM study on PANX1 directly visualized both mechanisms and corroborates with prior studies(14). While the caspase-dependent cleavage irreversibly activates PANX1, studies from several groups have converged on the concept that protein phosphorylation may be a novel mechanism to reversibly regulate PANX1 channel activity. In particular, the non-receptor tyrosine kinase Src seems to be a critical component during this process. For example, prolonged activation of P2X7 receptor leads to activation of PANX1 in many cell lines, opening of a large pore(22, 23). Through the use of a mimetic peptide and kinase inhibitors, Src was later found to be involved in mediating the initial steps of this signal transduction, but it does not directly phosphorylate PANX1 per se(24). In a more recent study PANX1 was proposed to make functional crosstalk with α7 nicotinic acetylcholine receptor (α7 nAChR) in neuroblastoma cell line(25). Src kinase is also involved in this process; however, it is unclear if PANX1 is a direct substrate of Src(25). In addition to tyrosine phosphorylation, murine PANX1 has been reported to undergo serine phosphorylation at Ser205 and Ser206, although two studies have shown contradictory effects of the two serine phosphorylation sites on PANX1 channel activity(26, 27). Likewise, the S394 of rat PANX1 could be phosphorylated by CaMKII upon an increase in cytoplasmic Ca^{2+} concentration(28). However, this residue is not conserved in human PANX1 (hPANX1). More recently, T302 and S328 were reported to be phosphorylated by protein kinase A, which affects the mechanically induced chloride conductance of PANX1 through the lateral tunnels(29).

Direct evidence of PANX1 phosphorylation by Src was provided by two groups using western
blot analysis, with Tyr198 and Tyr308 of mouse/rat PANX1 (equivalent to Tyr199 and Tyr309 of
hPANX1) being the target residues. Interestingly, the underlying signaling events of PANX1
phosphorylation are non-canonical. For example, the phosphorylation of Tyr308 by Src kinase
depends on ligand binding to the metabotropic NMDA receptor (NMDAR) in CA1 pyramidal
neurons, but does not require its pore opening and Ca^{2+} conductance(30, 31). As a result, it was
suggested that NMDARs, Src kinase and PANX1 could form a signaling complex, in which the
NMDAR is the ligand-sensing module whereas the PANX1 is the pore-forming unit.
Phosphorylation at Tyr308 is a key step for PANX1 activation in the NMDAR/Src/PANX1
signalosomes that eventually leads to the neuronal cell death during ischemia or stroke.
Likewise, the phosphorylation of Tyr198 of mouse PANX1 (mPANX1) was found to be relevant
in several different cellular context. For example, Lohman et al. showed that in vascular
endothelial cells, Tyr198 phosphorylation—which depends on type-1 TNF receptor (TNFR1) and
Src kinase—opens PANX1 channel, leading to ATP release and promoting leukocyte adhesion
and emigration(32). Moreover, Tyr198 phosphorylation is linked to vasoconstriction and blood
pressure regulation. Specifically, DeLalio et al. found that Tyr198 is constitutively
phosphorylated by Src in vascular smooth muscle cells, possibly through an α1-adrenergic
receptor (α1-AR)-based vasoconstriction mechanism(33). In a more recent study, Nouri-Nejad
et al. identified Tyr150 of PANX1 as another phosphorylation site through electrospray
ionization mass spectrometry, but only to a small degree (in 2 out of 11 LC-MS/MS spectra) (34).
The functional consequence of Tyr150 phosphorylation to PANX1 is unclear, as phosphor-
ablating mutant Y150F does not alter the properties of PANX1 current(34).

The initial aim of our study was to investigate how tyrosine phosphorylation regulates the
channel activity of PANX1 through structural approaches. To this end, we first examined
whether we could detect PANX1 phosphorylation using two commercially available antibodies
claimed to be specific for phosphorylated PANX1 at the Y198 and Y308 sites, respectively.
Surprisingly, neither antibody recognized PANX1 co-expressed with a constitutively active Src
variant. This result raised concerns about the validity of the conclusion that Tyr198 and Tyr308
of PANX1 are phosphorylated, as identification of both sites relied on these antibodies. We
further provided evidence that PANX1 is not phosphorylated at Tyr198 and Tyr308 by Src
kinase, at least in heterologous expression systems. Therefore, the paradigm of tyrosine
phosphorylation-dependent activation of PANX1 channel by Src kinase needs to be
reconsidered.

RESULT

Structural location and conservation of putative tyrosine phosphorylation sites in PANX1
Studies reporting the Src-mediated phosphorylation of Tyr198 and Tyr308 in murine PANX1
occurred prior to the availability of PANX1 structures(9–14, 31, 33). We first mapped the
corresponding residues, Tyr199 and Tyr309, into the apo state structure of hPANX1 (Figure 1A).
Interestingly, both residues are located in positions close to the side tunnel of PANX1, an anion-
selective permeation pathway responsible for the PANX1 basal current(14). Tyr199 is part of the
intersubunit interface of PANX1 and is partially buried, which may make it less accessible to Src
kinase. Tyr309, on the other hand, is completely buried inside the protein. Therefore,

phosphorylation of Tyr199 and Tyr309, if occurs, would require at least conformational changes
or partial unfolding of the intracellular domain. Such structural alternations are unlikely to occur
spontaneously without specific cellular conditions that favor this process. Accordingly,
phosphorylation of either of these two residues is expected to cause major conformational
changes of the intracellular domain due to the negative charges and the bulky size of the
introduced phosphate group(s).

We next compared the sequences of PANX1 across different species (Figure 1B). Both Tyr199
and Tyr309, as well as several residues nearby, are conserved among different PANX1
orthologs (Figure 1B), suggesting that phosphotyrosine antibodies developed against one
species are likely to cross-react with the others. Tyr199 is also conserved in PANX2 and PANX3,
the other two members of the pannexin family. In contrast, Tyr309 is only conserved in PANX1
and PANX3, while PANX2 has a phenylalanine at this position and lacks a few residues
preceding this residue (Figure 1B). Given the critical location of the two tyrosine residues, we
decided to experimentally characterize their phosphorylation status in the presence of Src
kinase.

Commercial antibodies, anti-PANX1-pY198 and anti-PANX1-pY308, do not recognize
phosphorylated human PANX1 in HEK293T cells
To investigate the phosphorylation of hPANX1 by Src, we used two commercially available
phosphor-specific antibodies that were used in previous studies to identify Tyr198 and Tyr308
as phosphorylation sites of murine PANX1(31–33) and were later reported to react with
hPANX1(35). In addition to wild-type hPANX1, we tested phosphor-ablating mutants Y199F and
Y309F (equivalent to mPANX1 Y198F and Y308F) as negative controls, which are expected to
abrogate the antibody binding even in the presence of active Src kinase. As in the prior
studies(31, 33), we used mouse Src (mSrc), which has 99.4% pairwise sequence similarity to
human Src. To increase the chance of detecting phosphorylated PANX1 and to introduce a
negative control for mSrc, we generated two mSrc mutants, Y529F and K297M, which
enhances and abolishes the kinase activity, respectively (Figure 2A). For the ease of
fluorescence imaging, our hPANX1 and mSrc genes are tagged with GFP and mCherry tags,
respectively (See Materials and Methods for detailed construct information), which allowed us to
compare the in-gel fluorescence with the western blot signals.

From the non-transfected HEK293T cells, neither anti-PANX1-pY198 nor anti-PANX1-pY308
produced any signal, suggesting that the antibodies do not recognize endogenous proteins
when hPANX1 and mSrc are absent (Figure 2B, lane 1). Similarly, from cells transfected with
hPANX1 alone, including WT, Y199F, and Y309F, neither antibody produced any signal, despite
the presence of clear GFP signals in the fluorescent gel indicating PANX1 expression (Figure
2B, lane 4-6). This is consistent with the notion that both antibodies do not recognize non-
phosphorylated PANX1.

Unexpectedly, from another control sample of cells transfected with WT mSrc alone, anti-
PANX1-pY198 yielded two prominent bands close to 75 kDa; anti-PANX1-pY308 also produced
a band corresponding to the lower band of anti-PANX1-pY198, albeit much weaker (Figure 2B,
lane 2). While the lower band corresponded to the fluorescent signal of mSrc (Figure 2B, lane 2),

the upper band may represent endogenous proteins induced and/or phosphorylated by mSrc.
The same signal pattern was detected with anti-PANX1-pY198 antibody from cells expressing
the constitutively active mSrc-Y529F variant alone, although both bands were slightly upshifted
(Figure 2B, lane 3). This upward shift was also observed in the fluorescent signal of mSrc-
Y529F, suggesting that it may be due to a more extensive phosphorylation of mSrc-Y529F itself
(lower band) and other endogenous proteins (upper band) associated with mSrc-Y529F (Figure
2B, lane 3). Interestingly, in addition to the band corresponding to the mSrc fluorescence signal,
the anti-PANX1-pY308 antibody showed a major band at about 50 kDa (Figure 2B, lane 3),
which may represent other sets of endogenous proteins induced and/or phosphorylated by
mSrc. Taken together, our control experiments showed that both antibodies demonstrate
promiscuous activity against mSrc and/or other unknown endogenous proteins, raising concerns
as to whether these antibodies should be used to interpret the phosphorylation of PANX1.

Despite the caveats of the antibodies mentioned above, we proceeded to test whether the two
antibodies could at least produce additional signals that may represent phosphorylated hPANX1
(when hPANX1 and mSrc are co-expressed). We designed several co-transfection experiments
combining WT mSrc or its constitutively active variant (Y529F) with WT PANX1 or its phosphor-
ablating mutants (Y199F and Y309F). All combinations yielded the same signal pattern as when
mSrc was expressed alone (Figure 2B, lanes 7–9 vs 2, lanes 10–12 vs 3), although under some
conditions the signals were weaker, which we attribute to the lower protein expression level
indicated by the fluorescent signals (Figure 2B, lane 10). Even when the constitutively active
mSrc was present, both antibodies failed to detect any signals corresponding to the GFP
florescence signal of hPANX1 (Figure 2B, lanes 7–12). Furthermore, there was no difference in
the signal pattern of WT hPANX1 and its two phosphor-ablating mutants, which is unexpected if
these antibodies are indeed specific for phosphorylated PANX1.

Our hPANX1 construct contains a GFP tag at its very C-terminus, which is unlikely to affect
PANX1 phosphorylation due to the flexibility of the C-terminus and its considerable distance
from the proposed phosphorylation sites. Nevertheless, to thoroughly address this concern, we
co-expressed a GFP-free version of hPANX1 with the constitutively active mSrc-Y529F, and
performed western blot analysis. Our result showed that both anti-PANX1-pY198 and anti-
PANX1-pY308 failed to detect hPANX1, regardless of the presence of the C-terminal GFP tag
(Figure 2-supplement 1, lanes 3-6).

Anti-PANX1-pY198 and anti-PANX1-pY308 antibodies do not recognize phosphorylated
human PANX1 *in vitro*
We next tested whether anti-PANX1-pY198 and anti-PANX1-pY308 antibodies could detect
PANX1 phosphorylation *in vitro,* as demonstrated in the literature(33), using purified hPANX1
produced in-house and purified human Src with a C-terminal GST tag (Src-GST) purchased
from a commercial source.

We first wanted to know whether purified hPANX1 could be phosphorylated by Src in an in vitro
context. To this end, we used a pan-specific phosphotyrosine antibody, anti-pY100, which is
expected to recognize a wide range of phosphorylated tyrosine sites. Our result clearly showed
that anti-pY100 detected only Src (Figure 3A, lanes 7-12), which is known to be

phosphorylated(36), but not PANX1 (Figure 3A, lanes 1-9), indicating that PANX1 is unlikely to
be phosphorylated in our *in vitro* setting. Similar to anti-pY100, anti-PANX1-pY198 yielded a
single band of the same size as the Src protein only in the presence of Src (Figure 3A, lanes 10-
15). These data confirmed that anti-PANX1-pY198 recognizes only Src, but not PANX1.

Notably, anti-PANX1-pY308 appeared to detect both Src and hPANX1 (Figure 3A, lanes 16-18).
However, the hPANX1 band was present even in the absence of Src (Figure 3A, lane 17), and
was not detected by anti-pY100 (Figure 3A, lane 7-9). This led us to speculate that anti-PANX1-
pY308 may recognize non-phosphorylated PANX1. To test this hypothesis, we performed *in*
vitro phosphorylation experiments on hPANX1 mutants around Y309, including L306A, K307A,
V308A, Y309F/A, E310A, I311A, and L312A (Figure 1B; Figure 3B,C). Our results showed that
these mutations also abolished the detection of the hPANX1 band by anti-PANX1-pY308
(Figure 3B,C). Thus, our data indicate that the anti-PANX1-pY308 antibody recognizes a non-
phosphorylated peptide epitope around Y309 of hPANX1.

We also speculate that anti-PANX1-pY308 prefers Src over hPANX1 as an epitope, because it
recognized Src both in its purified form and in crude whole-cell lysate, whereas it only
recognized hPANX1 in its purified form, but not in the whole-cell lysate (Figure 2B; Figure 3A-C).
As a result, the Src signal appeared to dominate in the western blot using whole-cell lysate
(Figure 2B). To test this hypothesis, we blotted whole-cell lysates from HEK293T cells
expressing either the WT PANX1 or the Y308F variant. Anti-PANX1-pY308 indeed detect a
weak band for WT PANX1, but not for the Y308F variant (Figure 3D).

Taken together, by using both cultured HEK293T cells and *in vitro* phosphorylation experiments,
we demonstrated that the anti-PANX1-pY198 and anti-PANX1-pY308 antibodies are non-
specific and may not be suitable for studying endogenous PANX1 protein phosphorylation.

Human PANX1 is not phosphorylated by mouse Src in HEK293T cells
Because commercially available phospho-specific antibodies failed to recognize PANX1, we
decided to use the Phos-tag gel to gauge the phosphorylation status of PANX1. The Phos-tag
gel allowed for an effective separation of proteins based on their phosphorylation level (37, 38).
We transiently transfected plasmids encoding hPANX1 and mSrc genes into HEK293T cells and
analyzed the cell lysates by Phos-tag gel and western blot. Figure 4A clearly showed differential
migration patterns of mSrc WT, Y529F and K297M in Phos-tag gel. De-phosphorylating of mSrc
by lambda protein phosphatase (λ-PPase) could partially convert the phosphorylated species
into the non-phosphorylated protein (Figure 4-figure supplement 1), suggesting that Y529F and
K297M mutation could affect the autophosphorylation of mSrc kinase itself (Figure 4A, lanes 3–
5). The hPANX1, when expressed by itself, showed one prominent band in both Phos-tag gel
and SDS-PAGE gel (Figure 4A and 3B, lane 2), which likely represents the non-phosphorylated
hPANX1 due to the absence of mSrc. We next co-expressed WT hPANX1 with the three mSrc
variants, and analyzed the cell lysates with Phos-tag gel to separate phosphorylated and
unphosphorylated hPANX1. Contrary to our expectation, the GFP fluorescence only showed a
monomeric band irrespective of mSrc activity, suggesting that hPANX1 is insensitive to
phosphorylation by mSrc (Figure 4A, lanes 6–8 vs 2). To ensure that the expression levels of
hPANX1 and mSrc were each comparable across the experimental conditions, we also

analyzed the same cell lysate samples using regular SDS-PAGE gel followed by western blot
and in-gel fluorescence and found no significant difference in protein expression (Figure 4B,
lanes 2-8). Our PANX1 gene is tagged with GFP at its C-terminus. To investigate whether GFP
would negatively affect the accessibility of the two tyrosine residues to mSrc, we also co-
expressed WT hPANX1 without GFP with mSrc variants in HEK293T cells and analyze the cell
lysate by Phos-tag gel and western blot analysis. No significant difference was observed
between hPANX1 expressed alone or with mSrc variants in Phos-tag gel (Figure 4-figure
supplement 2, lane 7-9 vs 3). Therefore, our Phos-tag gel analysis indicates that mSrc has no
impact on hPANX1 phosphorylation.

However, the inability to detect phosphorylated hPANX1 by Phos-tag gel analysis may be due
to the low level of phosphorylation. To address this possibility, we purified hPANX1 protein
expressed alone or co-expressed with the constitutively active mSrc-Y529F variant, and
conducted liquid chromatography with tandem mass spectrometry (LC-MS/MS) analysis on the
samples, which offers high sensitivity for detecting phosphorylated species. While we
unambiguously identified MS/MS spectra covered both Tyr199 and Tyr309 sites, the tyrosine
residues are not phosphorylated even when hPANX1 is co-expressed with mSrc-Y529F (Figure
4-figure supplement 3-5). Interestingly, we observed hPANX1 Ser385 phosphorylation when
mSrc-Y529F is present (Figure 4-figure supplement 6). However, out of 67 MS/MS spectra only
5 spectra contain the phosphorylated Ser385 (~7.5%). We note that phosphorylated peptides
may display a very different fragmentation and ionization efficiency compared to the non-
phosphorylated counterparts(39). Therefore, the ratio of the MS/MS spectra does not represent
the true phosphorylation level of hPANX1 protein. Our Phos-tag gel analysis suggests that the
Ser385 phosphorylation could be at a very low level as we did not observe other major bands in
Phos-tag gel when hPANX1 is co-expressed with mSrc-Y529F (Figure 4A). Nevertheless, our
result clearly demonstrated that Tyr199 and Tyr309 of hPANX1 are not phosphorylated by mSrc,
at least not to a detectable level, when transiently expressed in HEK293T cells.

Phosphorylation status of human PANX1 is not altered by Src in Neuro2A cells
Since our results conducted in HEK293T cells contradicted to what was known in the literature,
we wanted to test whether our result was cell line dependent. To this end, we co-expressed
hPANX1 and mSrc in Neuro2A cells, which were previously used as a model system to access
the phosphorylation of Y308 by NMDAR activation(31). We noticed that hPANX1 protein
expressed in Neuro2A cells was heavily glycosylated as indicated by its heterogeneous
migration bands in regular SDS-PAGE gel (Figure 5A, lane 2). However, the multimeric bands
can be reduced to a single band after PNGase F treatment, confirming that the heterogeneity is
indeed due to N-linked glycosylation (Figure 5A, lane 3). Next, we tested the phosphorylation
status of the samples using Phos-tag gel. Our result showed that deglycosylated hPANX1
migrates to the same location independent of mSrc, suggesting that Src does not alter the
phosphorylation status of hPANX1 (Figure 5B, lanes 8, 10, 12 vs 3).

Mouse PANX1 is not phosphorylated by Src in HEK293T cells
To rule out the possibility that hPANX1 may not be recognized by mouse Src, we repeated the
transient transfection experiment using mPANX1 (Figure 6). Using Phos-tag gel, we failed to
observe any difference between mPANX1 expressed alone or co-expressed with mSrc variants

in HEK293T cells, indicating that mPANX1 behaves in the same way as hPANX1 and is not
phosphorylated by mSrc (Figure 6A, lanes 6–8 vs 2). The expression levels of mPANX1 and
mSrc are each comparable across different experimental conditions as judged by western blot
and in-gel fluorescence image (Figure 6B).

We noted that the PANX1 antibody from Abcam (Catalogue number: ab124131) seems to
detect other non-specific bands in our experiment, including a major band close to 100 kDa and
a minor band at approximately 50 kDa (Figure 4B and 6B). To sort out the identity of the two
bands, we carried out a de-glycosylation experiment on non-transfected HEK293T cells and
hPANX1-expressing cells (Figure 6-figure supplement 1). Our result suggests that the 50 kDa
band could be partially de-glycosylated and migrate to a location where overexpressed PANX1
protein is expected to stay, whereas the 100 kDa band is insensitive to PNGase F treatment
(Figure 6-figure supplement 1). Thus, the lower band (50 kDa) likely represents endogenous
PANX1 from HEK293T cells, while the 100 kDa band is a non-specific target of anti-PANX1
antibody. This result highlights the caveats/complications of solely relying on western blot to
interpret the band identity. When combined with the fluorescence signal, it is possible to
determine with more confidence which bands correspond to the full-length mPANX1 and mSrc
protein in the SDS-PAGE gel.

The PANX1 current remains unchanged in the presence of active mSrc

To investigate whether hPANX1 channel activity is affected by the presence of active mSrc, we
co-expressed GFP-tagged hPANX1 with the mCherry-tagged constitutively active mSrc-Y529F,
and performed whole-cell patch clamp analysis. We targeted cells showing both green and red
fluorescence, indicative of expression of both hPANX1 and mSrc proteins (Figure 7A). We did
not observe any significant difference in the CBX-sensitive currents compared to the currents
when hPANX1 was expressed alone (Figure 7B). This data strongly suggests that mSrc does
not affect the gating of hPANX1 channel in HEK293T cells, further reinforcing the notion that
PANX1 is not phosphorylated by Src.

CONCLUSION

Protein phosphorylation is a common mechanism to regulate the physiology of large-pore
channel function. For example, Cx43 phosphorylation by p34[cdc2] kinase is associated with
channel internalization during mitosis(40–42). The volume-regulated anion channel was also
recently found to be phosphorylated by MSK1 kinase, which in turn controls cellular chloride
efflux(43). Not unexpectedly, PANX1 phosphorylation is also implicated in several physiological
conditions. The common scheme of these processes is that PANX1 acts downstream of a
membrane receptor, including P2X7, NMDARs, α7 nAChR, TNFR1, and α1-AR, which senses
various extracellular signal and results in PANX1 activation via a yet uncharacterized
intracellular signaling cascade(22, 25, 30–33). The Src non-receptor tyrosine kinase appears to
be a key player during this process, as inhibiting Src activity by kinase inhibitors could quench
PANX1 opening(22, 25, 30, 32, 33). Several evidence to support the notion that Src could
directly phosphorylate PANX1 comes from western blot analysis, in which phosphor-tyrosine
antibodies against Y198 and Y308 of murine PANX1 were used. In the study by DeLalio et al.,
attempts have been made to compare the effectiveness of pY198 antibody against WT

mPANX1 and Y198F mutant in HEK293T cells(33). However, the signal by the pY198 antibody
does not seem to match where the mPANX1 protein is expected to appear, although the blot is
rather noisy (see Figure S2A of DeLalio et al)(33). In the study by Weilinger et al., pY308 signal
is present even for Y308F mutant, suggesting that the antibody may not be specific for pY308 of
rat PANX1 (see Figure 5b of Weilinger et al)(31). Unfortunately, these antibodies are taken for
granted and continued to be used for scientific research(35).

Here, we combined multiple techniques, including Phos-tag gel, western blot, LC-MS/MS, and
patch-clamp electrophysiology to investigate Src-dependent PANX1 phosphorylation. Our
results, in contrast to previous findings, indicate that Tyr199 and Tyr309 of hPANX1 are not
effective substrates for Src kinase, at least in HEK293T and Neuro2A cells or in *in vitro*
experiments using purified hPANX1. Nevertheless, it is still possible that these two tyrosine sites
are phosphorylated by other tyrosine kinases in other cell lines. For example, recent
electrospray ionization mass spectrometry analysis on hPANX1 identified Tyr150 as a minor
phosphorylation site in human triple-negative breast cancer Hs578T cells(34). Our mass
spectrometry data failed to detect any fragments corresponding to a phosphorylated Tyr150
even when an active mSrc kinase is coexpressed with hPANX1, suggesting that Tyr150 may
not be effectively recognized by mSrc. Consistent with this, a number of other tyrosine kinases
have been predicted as the candidates for Tyr150 phosphorylation(34). Likewise, our study also
does not rule out the possibility that Src may still be involved in the regulation of PANX1, given
the fact that Src kinase inhibitors affect PANX1 channel opening under these conditions (22, 25,
30, 32, 33). It's possible that additional proteins are involved to relay the signaling output to
PANX1. For example, the endoplasmic reticulum (ER)-resident stromal interaction molecules
(STIM1/2) have recently been noted to mediate events between NMDAR and PANX1(44).
Future studies are required to fully decipher the underlying mechanisms.

Given the fact that the commercially available antibodies against phospho-PANX1 are non-
specific, we encourage the PANX1 research community to re-examine results derived from the
Src-mediated PANX1 phosphorylation and the associated antibodies. The technological
development in the production of phospho-specific antibodies represents a breakthrough in the
field of protein kinase and phosphoprotein research, providing unprecedented detection
precision and accuracy. While phospho-specific antibodies have been successfully developed
for many protein targets, great caution still needs to be taken when analyzing novel
phosphorylation sites and control experiments (such as the use of phosphor-ablating mutants
and pan-specific phosphorylation antibodies) should be conducted. Ideally, other
complementary techniques such as mass spectrometry and Phos-tag gel analysis should be
sought to independently verify the conclusion.

MATERIALS AND METHODS

Full-length human PANX1 in the pEGC Bacmam vector from our previous study was used(14).
Mouse PANX1 (UniprotID: Q9JIP4) was synthesized by GenScript and subcloned into the
pEGC Bacmam vector(45). The translated product contains the human or mouse PANX1
protein, a thrombin digestion site (LVPRGS), an enhanced GFP protein, and an 8x His tag.

pCMV5 mouse Src was a gift from Joan Brugge & Peter Howley (Addgene plasmid # 13663;
http://n2t.net/addgene:13663 ; RRID:Addgene_13663). The mouse Src gene was subcloned
into the pEGN Bacmam vector(45). The translated product contains a mCherry protein, a
thrombin digestion site, and the Src protein. Primers for site-directed mutagenesis were
designed using QuikChange Primer Design website
(https://www.agilent.com/store/primerDesignProgram.jsp) and synthesized by Eurofins
Genomics. The QuikChange mutagenesis protocol was used to generate all the mutants of the
study. Sanger sequencing was performed to identify positive clones.

Adherent HEK293T (ECACC, Catalogue Number: 96121229) cells were grown in DMEM media
supplemented with 10% fetal bovine serum. Transient transfection was conducted using
lipofectamine-2000 by following the manufacturer's protocol. Specifically, the cells were cultured
in 60mm Petri dishes until 80% confluency. Transfection solution was made by mixing 500ng of
plasmid DNA, 4uL of lipofectamine-2000 reagent, and 100uL Opti-MEM media. After 10min
incubation at room temperature, the DNA-lipid complexes were added to the cell culture and
incubated at 37°C. The next day, 10mM sodium butyrate was added to the cells to boost protein
expression. The cell culture was then grown at 30°C for another day before harvesting. The cell
pellet was flash-frozen with liquid nitrogen and stored at -80°C. For co-transfection experiments,
equal amounts of hPANX1 (250 ng) and mSrc (250 ng) plasmids were used in the transfection
mixture. For Neuro2A cell culture (ATCC, Catalogue Number: CCL-131), EMEM media is used
instead of DMEM. All the other procedures described above are the same for expressing
proteins in Neuro2A cells.

For the western blot experiment, the cell pellet was lysed in TBS buffer (20 mM Tris pH 8.0, 150
mM NaCl) with protease an inhibitor cocktail (1 mM phenylmethylsulfonyl fluoride, 2 mM
pepstatin, 0.8 µM aprotinin and 2 µg/ml leupeptin), 1mM sodium orthovanadate and 1% glyco-
diosgenin detergent for 30 min on ice. The lysate was clarified by centrifugation at 21,000rpm
for 20 min and the soluble portions were mixed with 2x SDS loading buffer supplemented with 5%
2-Mercaptoethanol. The samples were resolved in precast gradient SDS gel (4-20%) or 7.5%
PhosTag gel. A Chemidoc instrument was used to directly image the in-gel fluorescence signal
after electrophoresis. Subsequently, the protein in the gel was transferred to the nitrocellulose
membrane using the semi-dry transfer buffer (48 mM Tris base, 39 mM glycine, 20% methanol).
The membrane is blocked in the TBST buffer (20 mM Tris pH8.0, 150 mM NaCl, 0.1% Tween
80) with 4% non-fat milk for 1 h at room temperature. Afterward, primary antibodies (1:2000
dilution) were incubated with the membrane overnight at 4°C. The next day, the membrane was
washed with TBST buffer for 4 times, 10 min each before goat anti-rabbit IgG secondary
antibodies were added (1:25000 dilution). After 1h, the secondary antibodies were discarded
and the membrane was washed again with TBST for 4 times, 10 min each. The western blot
signal was detected using ECL Pierce substrate and imaged using a Chemidoc instrument. A
brightfield image was overlaid with the illuminance signal to visualize the position of protein
markers in the membrane. The anti-PANX1 antibody is obtained from abcam (Catalogue
number: ab124131). The anti-Src (Catalogue number: 36D10) and anti-pY100 (Catalogue
number: 9411) antibodies are obtained from Cell Signaling. The anti-PANX1-pY198 (Catalogue
number: ABN1681) and anti-PANX1-pY308 (Catalogue number: ABN1680) antibodies were
obtained from Millipore Sigma.

For de-glycosylation experiment, Neuro2A/HEK293T cells expressing designated genes were
solubilized in TBS buffer with protease inhibitor cocktail (1 mM phenylmethylsulfonyl fluoride, 2
mM pepstatin, 0.8 µM aprotinin and 2 µg/ml leupeptin), 1mM sodium orthovanadate and 1%
glyco-diosgenin detergent for 30min on ice. The samples were clarified by centrifugation at
20,000g for 30 min. The de-glycosylation reaction was made by mixing 16 uL of the supernatant
with 2uL of PNGase F enzyme and 2 uL of GlycoBuffer 2 (10X). The control reaction replaced
the PNGase F enzyme with water. The reaction was allowed to occur at room temperature
overnight. The next day, the samples were mixed with 20 uL 2× SDS sample-loading buffer
(Sigma) supplemented with 5% βME and resolved by SDS-PAGE. The gel was imaged in the
ChemiDoc system by probing the GFP and mCherry fluorescence signal.

For the dephosphorylation experiment, mSrc-mCherry WT, Y529F, and K297M were expressed
in HEK293T cells. The cells were lysed in TBS buffer (20mM Tris pH8, 150mM NaCl) with 1%
glyco-diosgenin detergent for 30min on ice. After clarifying the non-soluble debris by
centrifugation at 21,000rpm for 20min, 20 µL supernatant was mixed with 2.5 µL of 10X
NEBuffer for Protein MetalloPhosphatases, 2.5 µL 10 mM $MnCl_2$ and 2 µL Lambda Protein
Phosphatase (NEB). The dephosphorylation reaction was allowed to occur at 30 °C overnight.
Control samples without adding Lambda Protein Phosphatase are placed on ice. The next day,
the samples were mixed with 2× SDS sample-loading buffer (Sigma) supplemented with 5%
βME and resolved by SDS-PAGE or Phos-tag gel. The gel was imaged in the ChemiDoc system
by probing the mCherry fluorescence signal or western blot analysis.

For *in vitro* phosphorylation and mass-spectrometry analysis, the hPANX1 WT or mutants were
expressed alone or co-expressed with mSrc Y529F mutant in HEK293T cells. Specifically, 50
µg of plasmid DNA (in the case of the co-expressing experiment, 25 µg of human PANX1
plasmid and 25 µg of mouse Src Y529F mutant plasmid was used) was mixed with 150ug of
PEI 25K (Polysciences) in a and incubated at room temperature for 30 min. The DNA-PEI
complex was then added to the suspension cell culture of HEK293T cells at a density of $2x10^6$
cells/ml. After growing at 37 °C for 8 h, 10 mM sodium butyrate was added, and the temperature
was changed to 30°C for 40 h. The cells were then harvested and stored at -80 °C until
purification. The protein purification procedure was described in our previous study(14).

The *in vitro* phosphorylation is conducted using the human Src-GST protein (Sigma-Aldrich,
Catalogue number: S1076) according to the manufacturer's protocol. Specifically, 1.5 µg of
purified PANX1 protein is mixed with kinase assay buffer (25 mm MOPS, pH 7.2, 20 mM $MgCl_2$,
12.5 mM $MnCl_2$, 5 mM EGTA, 2 mM EDTA, 0.25 mM DTT), diluted 1:5 with 50 ng/µl bovine
serum albumin (BSA). The reaction was supplemented with 0.25 mM ATP, and 0.3 µg
recombinant human (active) Src-GST kinase. Samples were incubated for 1 h at 30 °C using a
PCR thermocycler. After 1 h, 2x SDS buffer is mixed with the sample to stop the reaction. The
sample is then resolved by an SDS-PAGE gel for subsequent western blot analysis.

Purified hPANX1 protein w- or w/o- co-expressing mSrc Y529F mutant was resolved in SDS-
PAGE gel and the band corresponding to hPANX1 protein was cut and subjected to in-gel
digestion with trypsin. Half of each digested sample was analyzed by nano LC-MS/MS with a
Waters M-Class HPLC system interfaced with a ThermoFisher Fusion Lumos mass

spectrometer. Peptides were loaded on a trapping column and eluted over a 75μm analytical
column at 350nL/min; both columns were packed with Luna C18 resin (Phenomenex). The
mass spectrometer was operated in data-dependent mode, with the Orbitrap operating at
60,000 FWHM and 15,000 FWHM for MS and MS/MS respectively. The instrument was run with
a 3 s cycle for MS and MS/MS.

Data were searched using a local copy of Mascot (Matrix Science) with the following parameters.
Enzyme: Trypsin/P; Database: SwissProt Human (concatenated forward and reverse plus
common contaminants); Fixed modification: Carbamidomethyl (C) Variable modifications:
Oxidation (M), Acetyl (N-term), Pyro-Glu (N-term Q), Deamidation (N/Q); Mass values:
Monoisotopic; Peptide Mass Tolerance: 10 ppm; Fragment Mass Tolerance: 0.02 Da; Max
Missed Cleavages: 2. Mascot DAT files were parsed into Scaffold (Proteome Software) for
validation, filtering and to create a non-redundant list per sample. Data were filtered using 1%
protein and peptide FDR and requiring at least two unique peptides per protein.

TsA201 cells were transfected using Lipofectamine 2000 (Thermo Fisher) according to the
manufacturer's protocol. The transfected cells were incubated at 37°C for 18-24 hours before
electrophysiological measurements. Whole-cell recordings were performed using a Multiclamp
700B (Axon Instruments) and Clampex software, with pipettes of 3-5 MOhm resistance filled
with an internal solution containing 145 mM NaCl, 10 mM Hepes, 10 mM EGTA, pH adjusted to
7.4. The external bath solution contained 160 mM NaCl, 10 mM Hepes, 3 mM KCl, 2 mM $CaCl_2$,
and 1 mM $MgCl_2$, also adjusted to pH 7.4. During the recordings, voltage steps ranging from
–100 mV to +140 mV were applied, each lasting 100 ms with a 20 mV increment between steps,
and membrane currents were digitally recorded at 10 kHz and filtered at 2 kHz. To precisely
measure the PANX1 channel's current, carbenoxolone disodium salt (CBX), a blocker of the
channel, was added to the bath solution at a final concentration of 0.1 mM, and CBX-sensitive
currents were subsequently calculated by comparing the difference in current amplitude in a cell
with and without the presence of CBX.

ACKNOWLEDGEMENTS

We acknowledge Dr. Henriette Remmer at the Proteomics & Peptide Synthesis Core in the
University of Michigan for the helpful discussion with Mass spectrometry data. We thank
members of the Du & Lü Labs for thoughtful discussions. Z.R. is supported by an American
Heart Association (AHA) postdoctoral fellowship (grant 20POST35120556) and the National
Institute of Health (NIH) (grant K99NS128258). W.L. is supported by the NIH (grant
R35GM138321). J.D. is supported by a McKnight Scholar Award, a Klingenstein-Simon Scholar
Award, a Sloan Research Fellowship in Neuroscience, a Pew Scholar in the Biomedical
Sciences award and an NIH grant (R01NS111031).
487

DATA AVAILABILITY
489 The LC-MS/MS data for WT human PANX1 with or without co-expressing Src-Y529F mutant is
490 deposited in Dryad database under https://doi.org/10.5061/dryad.4tmpg4fh7. The raw images of
491 the gels/immunoblots are provided as the source data for the corresponding figures.
492

 FIGURES

Figure 1: Location and sequence conservation of Tyr199 and Tyr309 in human PANX1. A,
The Tyr199 and Tyr309 of hPANX1 are located close to the side tunnel. Two adjacent subunits
are shown in semi-transparent surface representation. The Tyr199 and Tyr309 are shown in
stick representation. The green spheres indicate the pathway of the side tunnel. **B**, Sequence
alignment of PANX1 orthologs, human PANX2 and PANX3. Tyr198 and Tyr309 are highlighted
using a red circle. Secondary structural features are shown on the top.

Figure 2: Commercially available phospho-PANX1 antibodies (anti-PANX1-pY198 and
anti-PANX1-pY308) do not recognize PANX1. A, A schematic figure showing different mSrc
mutants used in the study. Mouse Src WT undergoes a dynamic equilibrium between the
inactive and active states that depends on the phosphorylation status of Tyr529. The Y529F
mutation renders Src constitutively active due to the disruption of the autoinhibitory interaction
between the SH2 domain and the C-terminal tail(36). The K297M is catalytically incompetent as
the mutation abolishes the ATP-binding capability of mSrc(46). **B**, Human PANX1 (WT, Y199F
or Y309F) are co-expressed with mSrc (WT or Y529F) in HEK293T cells. The cell lysates are
analyzed by SDS gel and blotted with anti-PANX1-pY198 (top) and anti-PANX1-pY308 (middle)
antibodies. The in-gel fluorescence of GFP and mCherry signal are shown at the bottom. The
positions of the signals detected by anti-PANX1-pY198 and anti-PANX1-pY308 antibodies do
not match the position of the GFP fluorescence signal, indicating that the two antibodies are not
specific for PANX1.
513
514 **Figure 2—source data 1: Uncropped immunoblots for Figure 2B.**
515 **Figure 2—source data 2: Raw image for anti-PANX1-pY198 immunoblot in Figure 2B.**
516 **Figure 2—source data 3: Raw image for anti-PANX1-pY308 immunoblot in Figure 2B.**
517 **Figure 2—source data 4: Raw image for in-gel fluorescence (GFP and mCherry) in Figure**
518 **2B.**
519
Figure 2—figure supplement 1: The hPANX1 cannot be detected by anti-PANX1-pY198
and anti-PANX1-pY308 irrespective of C-terminal GFP tag. Human PANX1 w- or w/o C-
terminal GFP tag were co-expressed with mSrc-Y529F in HEK293T cells. The cell lysates were
analyzed by SDS gel and blotted with anti-PANX1-pY198, anti-PANX1-pY308, anti-PANX1, and
anti-Src antibodies. The in-gel fluorescence of GFP and mCherry signal were shown at the
bottom. The positions of the signals detected by anti-PANX1-pY198 and anti-PANX1-pY308
antibodies did not match the position of the anti-PANX1 western blot signal and GFP
fluorescence signal.
528
Figure 3: The anti-PANX1-pY198 and anti-PANX1-pY308 are not specific to hPANX1 *in*
vitro. **A**, *In vitro* phosphorylation of hPANX1-GFP WT by active human Src-GST protein. The
leftmost panel showed the in-gel fluorescence of GFP overlaid with protein marker. The other
panels (from left to right) represent the western blot signal of anti-PANX1, anti-pY100, anti-Src,
anti-PANX1-PANX1-pY198, and anti-PANX1-pY308 antibodies, respectively. **B**, *In vitro*
phosphorylation of hPANX1-GFP WT, L306A, K307A, V308A, Y309F, E310A, I311A, or L312A
mutants. The left panel showed the in-gel fluorescence of GFP overlaid with protein marker. The

right panel showed the western blot signal of anti-PANX1-pY308 antibody. Only hPANX1 WT
lane showed western blot signal. **C**, *In vitro* phosphorylation of hPANX1-GFP WT, Y309A, or
Y309F mutants. The left panel showed the in-gel fluorescence of GFP overlaid with protein
marker (red). The right panel showed the western blot signal of anti-PANX1-pY308 antibody.
The Y309A mutant rendered PANX1 prone to oligomerization in our SDS gel experiment, but
not for Y309F. **D**, Whole cell lysate of HEK293T cells transiently transfected with hPANX1 WT
or Y309F. Left panel showed the in-gel fluorescence of GFP overlaid with protein marker.
Middle and right panels represent western blot result using anti-PANX1 and anti-PANX1-pY308
antibody, respectively. The location where PANX1 is expected to migrate in the SDS-PAGE gel
is indicated. Of note, visualizing such a band requires a very long exposure, which resulted in a
large number of non-specific bands.
547
Figure 3—source data 1: Uncropped immunoblots for Figure 3.
Figure 3—source data 2: Raw image for in-gel fluorescence (GFP and mCherry) in Figure
3A.
Figure 3—source data 3: Raw image for anti-PANX1 immunoblot in Figure 3A.
Figure 3—source data 4: Raw image for anti-pY100 immunoblot in Figure 3A.
Figure 3—source data 5: Raw image for anti-Src immunoblot in Figure 3A.
Figure 3—source data 6: Raw image for anti-PANX1-pY198 immunoblot in Figure 3A.
Figure 3—source data 7: Raw image for anti-PANX1-pY308 immunoblot in Figure 3A.
Figure 3—source data 8: Raw image for in-gel fluorescence (GFP and mCherry) in Figure
3B,C.
Figure 3—source data 9: Raw image for anti-PANX1-pY308 immunoblot in Figure 3B,C.
Figure 3—source data 10: Raw image for in-gel fluorescence (GFP and mCherry) in
Figure 3D.
Figure 3—source data 11: Raw image for anti-PANX1 immunoblot in Figure 3D.
Figure 3—source data 12: Raw image for anti-PANX1-pY308 immunoblot in Figure 3D.
563
Figure 4: Human PANX1 is not phosphorylated by mSrc when expressed in HEK293T
cells. A, The Phos-tag gel result of human PANX1 co-expressed with different mSrc mutants.
The GFP and mCherry fluorescence signal were overlaid to generate the image. The mSrc
kinase of different phosphorylation status are labeled. In all conditions, hPANX1 migrated as a
single band. **B**, Analysis of the same cell lysate sample in B using regular SDS gel. Western blot
experiment was conducted using anti-PANX1 (top) and anti-Src (middle) antibodies. The anti-
PANX1 antibody detected a non-specific band at 100 kDa from the whole-cell lysate with
unknown identity. The in-gel fluorescence for GFP and mCherry is shown at the bottom where
the position of mSrc and PANX1 protein could be compared with the western blot signal.

Figure 4—source data 1: Uncropped immunoblots for Figure 4.
Figure 4—source data 2: Raw image for in-gel fluorescence (GFP and mCherry) in Figure
4A.
Figure 4—source data 3: Raw image for anti-PANX1 immunoblot in Figure 4B.
Figure 4—source data 4: Raw image for anti-Src immunoblot in Figure 4B.
Figure 4—source data 5: Raw image for in-gel fluorescence (GFP and mCherry) in Figure

4B.

Figure 4—figure supplement 1: Dephosphorylation of mSrc protein by lambda protein

phosphotase (λ-PP). Phos-tag gel showed that mSrc-mCherry WT, Y529F, and K297M protein

had different phosphorylation status. Phosphorylated mSrc could be converted to the non-

phosphorylated form by λ-PP.

Figure 4—figure supplement 1—source data 1: Uncropped immunoblots for Figure 4—

figure supplement 1.

Figure 4—figure supplement 1—source data 2: Raw image for in-gel fluorescence

(mCherry) in Figure 4—figure supplement 1 (upper panel).

Figure 4—figure supplement 1—source data 3: Raw image for in-gel fluorescence

(mCherry) in Figure 4—figure supplement 1 (lower panel).

Figure 4—supplement 2: Wild-type hPANX1 without C-terminal GFP tag is not

phosphorylated my mSrc. Human PANX1 with/without C-terminal GFP tag is co-expressed

with Src-mCherry WT, Y529F, or K297M in HEK293T cells. Cell lysate was subjected to

PNGase F de-glycosylation prior to analysis on SDS-PAGE gel (upper and middle panels) or

Phos-tag gel (bottom panel). Upper panel showed the GFP and mCherry fluorescence signal in

SDS-PAGE gel. Middle panel showed the western blot result of anti-PANX1 antibody blot in

SDS-PAGE gel. Bottom panel showed the western blot result of anti-PANX1 antibody blot in

Phos-tag gel.

Figure 4—figure supplement 2—source data 1: Uncropped immunoblots for Figure 4—

figure supplement 2.

Figure 4—figure supplement 2—source data 2: Raw image for in-gel fluorescence (GFP

and mCherry) in Figure 4—figure supplement 2 (upper panel).

Figure 4—figure supplement 2—source data 3: Raw image for anti-PANX1 immunoblot in

Figure 4—figure supplement 2 (middle panel).

Figure 4—figure supplement 2—source data 4: Raw image for anti-PANX1 immunoblot in

Figure 4—figure supplement 2 (lower panel).

Figure 4—supplement 3: Coverage map of LC-MS/MS analysis of purified hPANX1-GFP.

A, The sequence coverage of hPANX1 expressed alone. **B**, The sequence coverage of

hPANX1 expressed with the constitutively active mSrc Y529F. Of note, both Tyr199 and Tyr309

are covered by LC-MS/MS spectra.

Figure 4—figure supplement 3—source data 1: LC-MS/MS data for WT human PANX1

with or without co-expressing Src-Y529F mutant [Dataset]. Dryad.

https://doi.org/10.5061/dryad.4tmpg4fh7

Figure 4—supplement 4: Representative LC-MS/MS spectra containing the Tyr199 of

hPANX1. A, The raw spectrum of (K)YPIVEQYLKTK(K) peptide from hPANX1 expressed

without Src. B ions and Y ions are colored in red and blue, respectively. **B**, The fragmentation

table of the (K)YPIVEQYLKTK(K) peptide shown in panel A. Red and blue color highlighted the

B ions and Y ions for the detected amino acid, respectively. The non-phosphorylated Tyr199 is
supported by both B ions and Y ions (blue frame). **C**, The raw spectrum of (K)YPIVEQYLK(T)
peptide from human PANX1 expressed with mSrc Y529F mutant. B ions and Y ions are colored
in red and blue, respectively. **D**, The fragmentation table of the the (K)YPIVEQYLK(T) peptide
shown in panel C. Red and blue color highlighted the B ions and Y ions for the detected amino
acid. The non-phosphorylated Tyr199 is supported by Y ions (blue frame).

Figure 4—supplement 5: Representative LC-MS/MS spectrum covering the Tyr309 for
hPANX1. A, The raw spectrum of (K)VYEILPTFDVLHFK(S) peptide from hPANX1 expressed
without Src. B ions and Y ions are colored in red and blue, respectively. **B**, The fragmentation
table of the (K)VYEILPTFDVLHFK(S) peptide shown in A. Red and blue color highlighted the B
ions and Y ions for the detected amino acid, respectively. The non-phosphorylated Tyr308 is
supported B ions (blue frame). **C**, The raw spectrum of (K)VYEILPTFDVLHFK(S) peptide from
human PANX1 expressed with mSrcY529F mutant. B ions and Y ions are colored in red and
blue, respectively. **D**, The fragmentation table of the the (K)VYEILPTFDVLHFK(S) peptide
shown in C. Red and blue color highlighted the B ions and Y ions for the detected amino acid.
The non-phosphorylated Tyr309 is supported by both B ions and Y ions (blue frame).

Figure 4—supplement 6: Representative LC-MS/MS spectrum covering the Ser385 for
hPANX1. A, The raw spectrum of (K)TPMSAEMR(E) peptide from hPANX1 expressed without
Src. B ions and Y ions are colored in red and blue, respectively. **B**, The fragmentation table of
the (K)TPMSAEMR(E) peptide shown in A. Red and blue color highlighted the B ions and Y ions
for the detected amino acid, respectively. The non-phosphorylated Ser385 is supported by both
B ions and Y ions (blue frame). **C**, One raw spectrum of (K)TPMSAEMR(E) peptide from human
PANX1 expressed with Src-Y529F mutant. B ions and Y ions are colored in red and blue,
respectively. **D**, The fragmentation table of the (K)TPMSAEMR(E) peptide shown in C. Red and
blue color highlighted the B ions and Y ions for the detected amino acid. The phosphorylated
Ser385 is supported by Y ions (blue frame). **E**, One raw spectrum of (K)TPMSAEMR(E) peptide
from hPANX1 expressed with Src-Y529F mutant. B ions and Y ions are colored in red and blue,
respectively. **F**, The fragmentation table of the (K)TPMSAEMR(E) peptide shown in E. Red and
blue color highlighted the B ions and Y ions for the detected amino acid. The non-
phosphorylated Ser385 is supported by both B ions and Y ions (blue frame).

Figure 5: Human PANX1 is not phosphorylated by Src when expressed in Neuro2A cells.
A, A regular SDS gel showing the GFP and mCherry fluorescence signal of hPANX1 and mSrc
co-expressed in Neuro2A cells. For each condition containing PANX1, PNGase F treatment is
performed to de-glycosylate the protein. **B**, A PhosTag gel result using the same samples
analyzed in **A**.

Figure 5—source data 1: Uncropped immunoblots for Figure 5.
Figure 5—source data 2: Raw image for in-gel fluorescence (GFP and mCherry) in Figure
5A.
Figure 5—source data 3: Raw image for in-gel fluorescence (GFP and mCherry) in Figure
5B.
660

Figure 6: Mouse PANX1 is not phosphorylated by Src when expressed in HEK293T cells.
A, A Phos-tag gel result of mPANX1 co-expressed with different mSrc mutants. The GFP and mCherry fluorescence signal are overlaid to generate the image. The mSrc kinase of different phosphorylation status are labeled. In all the conditions, mouse PANX1 migrates to as a single band. **B**, Analysis of the same cell lysate sample in B using regular SDS-PAGE gel. Western blot experiment is conducted using anti-PANX1 (top) and anti-Src (middle) antibodies. The in-gel fluorescence for GFP and mCherry is shown at the bottom where the position of mSrc and mPANX1 protein could be compared with the western blot signal.

Figure 6—source data 1: Uncropped immunoblots for Figure 6.
Figure 6—source data 2: Raw image for in-gel fluorescence (GFP and mCherry) in Figure 6A.
Figure 6—source data 3: Raw image for anti-PANX1 immunoblot in Figure 6B.
Figure 6—source data 4: Raw image for anti-Src immunoblot in Figure 6B.
Figure 6—source data 5: Raw image for in-gel fluorescence (GFP and mCherry) in Figure 6B.

Figure 6—supplement 1: The anti-PANX1 antibody detects both hPANX1 and non-specific proteins. The non-transfected HEK293T cells or cells transfected with PANX1 (without the GFP tag) is digested by PNGase F and analyzed by western blot. The anti-PANX1 antibody produced two bands from the non-transfected HEK293T cell lysate at approximately 100 kDa and 50 kDa. After PNGase F treatment, the 50 kDa band (indicated by blue dotted lines) can be partially shifted to a location where monomeric PANX1 protein is located. The 100 kDa band (indicated by dotted magenta lines) is not sensitive to PNGase F treatment. The right panel showed a long exposure image of the same blot on the left.

Figure 6—figure supplement 1—source data 1: Uncropped immunoblots for Figure 6—figure supplement 1.
Figure 6—figure supplement 1—source data 2: Raw image for anti-PANX1 immunoblot in Figure 6—figure supplement 1 (left panel).
Figure 6—figure supplement 1—source data 3: Raw image for anti-PANX1 immunoblot in Figure 6—figure supplement 1 (right panel).

Figure 7: The PANX1 current remains unchanged in the presence of active mSrc. A, Representative fluorescent images of HEK293T cells transfected with hPANX1-GFP and mSrc-Y529F-mCherry. Cells that show both green (GFP) and red (mCherry) fluorescence are selected for whole-cell patch-clamp analysis. **B**, The CBX sensitive whole-cell current of hPANX1-GFP and hPANX1+mSrc-Y529F-mCherry. Two-way ANOVA for repeated measurements was conducted to compare the hPANX1-GFP and hPANX1+mSrc-Y529F-mCherry. The analysis revealed a non-significant difference between the two groups ($F_{(1, 182)}$= 0.07286, P = 0.7875). Data are presented as mean ± SEM.

REFERENCES

1.   Adamson, S. E., and Leitinger, N. (2014) The role of pannexin1 in the induction and
resolution of inflammation. *FEBS Lett.* **588**, 1416–1422
2.   Makarenkova, H., and Shestopalov, V. (2014) The role of pannexin hemichannels in
inflammation and regeneration. *Front. Physiol.* **5**, 63
3.   Good, M. E., Begandt, D., DeLalio, L. J., Keller, A. S., Billaud, M., and Isakson, B. E.
(2015) Emerging concepts regarding pannexin 1 in the vasculature. *Biochem. Soc. Trans.*
43, 495–501
4.   Chekeni, F. B., Elliott, M. R., Sandilos, J. K., Walk, S. F., Kinchen, J. M., Lazarowski, E.
R., Armstrong, A. J., Penuela, S., Laird, D. W., Salvesen, G. S., Isakson, B. E., Bayliss, D.
A., and Ravichandran, K. S. (2010) Pannexin 1 channels mediate "find-me" signal release
and membrane permeability during apoptosis. *Nature.* **467**, 863–867
5.   Medina, C. B., Mehrotra, P., Arandjelovic, S., Perry, J. S. A., Guo, Y., Morioka, S., Barron,
B., Walk, S. F., Ghesquière, B., Krupnick, A. S., Lorenz, U., and Ravichandran, K. S.
(2020) Metabolites released from apoptotic cells act as tissue messengers. *Nature.* **580**,
130–135
6.   Baranova, A., Ivanov, D., Petrash, N., Pestova, A., Skoblov, M., Kelmanson, I., Shagin,
D., Nazarenko, S., Geraymovych, E., Litvin, O., Tiunova, A., Born, T. L., Usman, N.,
Staroverov, D., Lukyanov, S., and Panchin, Y. (2004) The mammalian pannexin family is
homologous to the invertebrate innexin gap junction proteins. *Genomics.* **83**, 706–716
7.   Sandilos, J. K., and Bayliss, D. A. (2012) Physiological mechanisms for the modulation of
pannexin 1 channel activity. *J. Physiol.* **590**, 6257–6266
8.   Scemes, E., and Velíšková, J. (2019) Exciting and not so exciting roles of pannexins.
Neurosci. Lett. **695**, 25–31
9.   Michalski, K., Syrjanen, J. L., Henze, E., Kumpf, J., Furukawa, H., and Kawate, T. (2020)
The Cryo-EM structure of pannexin 1 reveals unique motifs for ion selection and inhibition.
Elife. **9**, e54670
10.   Deng, Z., He, Z., Maksaev, G., Bitter, R. M., Rau, M., Fitzpatrick, J. A. J., and Yuan, P.
(2020) Cryo-EM structures of the ATP release channel pannexin 1. *Nat. Struct. Mol. Biol.*
27, 373–381
11.   Qu, R., Dong, L., Zhang, J., Yu, X., Wang, L., and Zhu, S. (2020) Cryo-EM structure of
human heptameric Pannexin 1 channel. *Cell Res.* **30**, 446–448
12.   Jin, Q., Zhang, B., Zheng, X., Li, N., Xu, L., Xie, Y., Song, F., Bhat, E. A., Chen, Y., Gao,
N., Guo, J., Zhang, X., and Ye, S. (2020) Cryo-EM structures of human pannexin 1
channel . *Cell Res.* **30**, 449–451
13.   Mou, L., Ke, M., Song, M., Shan, Y., Xiao, Q., Liu, Q., Li, J., Sun, K., Pu, L., Guo, L.,
Geng, J., Wu, J., and Deng, D. (2020) Structural basis for gating mechanism of Pannexin
1 channel. *Cell Res.* **30**, 452–454
14.   Ruan, Z., Orozco, I. J., Du, J., and Lü, W. (2020) Structures of human pannexin 1 reveal
ion pathways and mechanism of gating. *Nature.* **584**, 646–651
15.   Crocetti, L., Guerrini, G., Puglioli, S., Giovannoni, M. P., Di Cesare Mannelli, L., Lucarini,
E., Ghelardini, C., Wang, J., and Dahl, G. (2021) Design and synthesis of the first indole-
based blockers of Panx-1 channel. *Eur. J. Med. Chem.* **223**, 113650
16.   Chiu, Y.-H., Schappe, M. S., Desai, B. N., and Bayliss, D. A. (2017) Revisiting multimodal
activation and channel properties of Pannexin 1. *J. Gen. Physiol.* **150**, 19–39
17.   Mim, C., Perkins, G., and Dahl, G. (2021) Structure versus function: Are new
conformations of pannexin 1 yet to be resolved? *J. Gen. Physiol.* 10.1085/jgp.202012754
18.   Boyce, A. K. J., Epp, A. L., Nagarajan, A., and Swayne, L. A. (2018) Transcriptional and
post-translational regulation of pannexins. *Biochim. Biophys. acta. Biomembr.* **1860**, 72–
82
19.   Sosinsky, G. E., Boassa, D., Dermietzel, R., Duffy, H. S., Laird, D. W., MacVicar, B.,
Naus, C. C., Penuela, S., Scemes, E., Spray, D. C., Thompson, R. J., Zhao, H.-B., and

Dahl, G. (2011) Pannexin channels are not gap junction hemichannels. *Channels (Austin).*
5, 193–197

20. Dahl, G., and Locovei, S. (2006) Pannexin: to gap or not to gap, is that a question?
IUBMB Life. **58**, 409–419

21. Huang, Y., Grinspan, J. B., Abrams, C. K., and Scherer, S. S. (2007) Pannexin1 is
expressed by neurons and glia but does not form functional gap junctions. *Glia.* **55**, 46–
56

22. Locovei, S., Scemes, E., Qiu, F., Spray, D. C., and Dahl, G. (2007) Pannexin1 is part of
the pore forming unit of the P2X(7) receptor death complex. *FEBS Lett.* **581**, 483–488

23. Pelegrin, P., and Surprenant, A. (2006) Pannexin-1 mediates large pore formation and
interleukin-1beta release by the ATP-gated P2X7 receptor. *EMBO J.* **25**, 5071–5082

24. Iglesias, R., Locovei, S., Roque, A., Alberto, A. P., Dahl, G., Spray, D. C., and Scemes, E.
(2008) P2X7 receptor-Pannexin1 complex: pharmacology and signaling. *Am. J. Physiol.*
Cell Physiol. **295**, C752-60

25. Maldifassi, M. C., Momboisse, F., Guerra, M. J., Vielma, A. H., Maripillán, J., Báez-Matus,
X., Flores-Muñoz, C., Cádiz, B., Schmachtenberg, O., Martínez, A. D., and Cárdenas, A.
M. (2021) The interplay between α7 nicotinic acetylcholine receptors, pannexin-1
channels and P2X7 receptors elicit exocytosis in chromaffin cells. *J. Neurochem.* **157**,
1789–1808

26. Poornima, V., Vallabhaneni, S., Mukhopadhyay, M., and Bera, A. K. (2015) Nitric oxide
inhibits the pannexin 1 channel through a cGMP-PKG dependent pathway. *Nitric oxide*
Biol. Chem. **47**, 77–84

27. Medina, C. B., Chiu, Y.-H., Stremska, M. E., Lucas, C. D., Poon, I., Tung, K. S., Elliott, M.
R., Desai, B., Lorenz, U. M., Bayliss, D. A., and Ravichandran, K. S. (2021) Pannexin 1
channels facilitate communication between T cells to restrict the severity of airway
inflammation. *Immunity.* **54**, 1715-1727.e7

28. López, X., Palacios-Prado, N., Güiza, J., Escamilla, R., Fernández, P., Vega, J. L., Rojas,
M., Marquez-Miranda, V., Chamorro, E., Cárdenas, A. M., Maldifassi, M. C., Martínez, A.
D., Duarte, Y., González-Nilo, F. D., and Sáez, J. C. (2021) A physiologic rise in
cytoplasmic calcium ion signal increases pannexin1 channel activity via a C-terminus
phosphorylation by CaMKII. *Proc. Natl. Acad. Sci. U. S. A.* 10.1073/pnas.2108967118

29. López, X., Escamilla, R., Fernández, P., Duarte, Y., González-Nilo, F., Palacios-Prado,
N., Martinez, A. D., and Sáez, J. C. (2020) Stretch-Induced Activation of Pannexin 1
Channels Can Be Prevented by PKA-Dependent Phosphorylation. *Int. J. Mol. Sci.*
10.3390/ijms21239180

30. Weilinger, N. L., Tang, P. L., and Thompson, R. J. (2012) Anoxia-induced NMDA receptor
activation opens pannexin channels via Src family kinases. *J. Neurosci. Off. J. Soc.*
Neurosci. **32**, 12579–12588

31. Weilinger, N. L., Lohman, A. W., Rakai, B. D., Ma, E. M. M., Bialecki, J., Maslieieva, V.,
Rilea, T., Bandet, M. V, Ikuta, N. T., Scott, L., Colicos, M. A., Teskey, G. C., Winship, I. R.,
and Thompson, R. J. (2016) Metabotropic NMDA receptor signaling couples Src family
kinases to pannexin-1 during excitotoxicity. *Nat. Neurosci.* **19**, 432–442

32. Lohman, A. W., Leskov, I. L., Butcher, J. T., Johnstone, S. R., Stokes, T. A., Begandt, D.,
DeLalio, L. J., Best, A. K., Penuela, S., Leitinger, N., Ravichandran, K. S., Stokes, K. Y.,
and Isakson, B. E. (2015) Pannexin 1 channels regulate leukocyte emigration through the
venous endothelium during acute inflammation. *Nat. Commun.* **6**, 7965

33. DeLalio, L. J., Billaud, M., Ruddiman, C. A., Johnstone, S. R., Butcher, J. T., Wolpe, A. G.,
Jin, X., Keller, T. C. S. 4th, Keller, A. S., Rivière, T., Good, M. E., Best, A. K., Lohman, A.
W., Swayne, L. A., Penuela, S., Thompson, R. J., Lampe, P. D., Yeager, M., and Isakson,
B. E. (2019) Constitutive SRC-mediated phosphorylation of pannexin 1 at tyrosine 198
occurs at the plasma membrane. *J. Biol. Chem.* **294**, 6940–6956

34. Nouri-Nejad, D., O'Donnell, B. L., Patil, C. S., Sanchez-Pupo, R. E., Johnston, D.,
Sayedyahossein, S., Jurcic, K., Lau, R., Gyenis, L., Litchfield, D. W., Jackson, M. F.,
Gloor, G. B., and Penuela, S. (2021) Pannexin 1 mutation found in melanoma tumor
reduces phosphorylation, glycosylation, and trafficking of the channel-forming protein.
Mol. Biol. Cell. **32**, 376–390

35. Metz, L. M., and Elvers, M. (2022) Pannexin-1 Activation by Phosphorylation Is Crucial
for Platelet Aggregation and Thrombus Formation. *Int. J. Mol. Sci.*
10.3390/ijms23095059

36. Nada, S., Okada, M., MacAuley, A., Cooper, J. A., and Nakagawa, H. (1991) Cloning of a
complementary DNA for a protein-tyrosine kinase that specifically phosphorylates a
negative regulatory site of p60c-src. *Nature.* **351**, 69–72

37. Nagy, Z., Comer, S., and Smolenski, A. (2018) Analysis of Protein Phosphorylation Using
Phos-Tag Gels. *Curr. Protoc. protein Sci.* **93**, e64

38. Nishioka, K., Kato, Y., Ozawa, S., Takahashi, Y., and Sakamoto, W. (2021) Phos-tag-
based approach to study protein phosphorylation in the thylakoid membrane. *Photosynth.*
Res. **147**, 107–124

39. Dephoure, N., Gould, K. L., Gygi, S. P., and Kellogg, D. R. (2013) Mapping and analysis
of phosphorylation sites: a quick guide for cell biologists. *Mol. Biol. Cell.* **24**, 535–542

40. Xie, H., Laird, D. W., Chang, T. H., and Hu, V. W. (1997) A mitosis-specific
phosphorylation of the gap junction protein connexin43 in human vascular cells:
biochemical characterization and localization. *J. Cell Biol.* **137**, 203–210

41. Kanemitsu, M. Y., Jiang, W., and Eckhart, W. (1998) Cdc2-mediated phosphorylation of
the gap junction protein, connexin43, during mitosis. *Cell growth Differ. Mol. Biol. J. Am.*
Assoc. Cancer Res. **9**, 13–21

42. Lampe, P. D., Kurata, W. E., Warn-Cramer, B. J., and Lau, A. F. (1998) Formation of a
distinct connexin43 phosphoisoform in mitotic cells is dependent upon p34cdc2 kinase. *J.*
Cell Sci. **111 (Pt 6)**, 833–841

43. Serra, S. A., Stojakovic, P., Amat, R., Rubio-Moscardo, F., Latorre, P., Seisenbacher, G.,
Canadell, D., Böttcher, R., Aregger, M., Moffat, J., de Nadal, E., Valverde, M. A., and
Posas, F. (2021) LRRC8A-containing chloride channel is crucial for cell volume recovery
and survival under hypertonic conditions. *Proc. Natl. Acad. Sci. U. S. A.*
10.1073/pnas.2025013118

44. Patil, C. S., Li, H., Lavine, N. E., Shi, R., Bodalia, A., Siddiqui, T. J., and Jackson, M. F.
(2022) ER-resident STIM1/2 couples Ca(2+) entry by NMDA receptors to pannexin-1
activation. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2112870119

45. Goehring, A., Lee, C.-H., Wang, K. H., Michel, J. C., Claxton, D. P., Baconguis, I., Althoff,
T., Fischer, S., Garcia, K. C., and Gouaux, E. (2014) Screening and large-scale
expression of membrane proteins in mammalian cells for structural studies. *Nat. Protoc.*
9, 2574–2585

46. Roskoski, R. J. (2015) Src protein-tyrosine kinase structure, mechanism, and small
molecule inhibitors. *Pharmacol. Res.* **94**, 9–25

Figure 1

Figure 2

Figure 2-figure supplement 1

Figure 3

Figure 4

A

B

SDS-PAGE gel

Figure 4-figure supplement 1

Figure 4-figure supplement 2

Figure 4-figure supplement 3

A

70 exclusive unique peptides, 162 exclusive unique spectra, 266 total spectra, 508/690 amino acids (74% coverage)

B

84 exclusive unique peptides, 243 exclusive unique spectra, 542 total spectra, 579/690 amino acids (84% coverage)

Figure 4-figure supplement 4

A

B

B	B Ions	B+2H	B–NH3	B–H2O	AA	Y Ions	Y+2H	Y–NH3	Y–H2O	Y
1	164.1			146.1	Y	1,381.8	691.4	1,364.7	1,363.8	11
2	261.1			243.1	P	1,218.7	609.9	1,201.7	1,200.7	10
3	374.2			356.2	I	1,121.7	561.3	1,104.6	1,103.6	9
4	473.3			455.3	V	1,008.6	504.8	991.5	990.6	8
5	602.3			584.3	E	909.5	455.3	892.5	891.5	7
6	730.4	365.7	713.4	712.4	Q	780.5	390.7	763.4	762.5	6
7	893.4	447.2	876.4	875.4	Y	652.4	326.7	635.4	634.4	5
8	1,006.5	503.8	989.5	988.5	L	489.3	245.2	472.3	471.3	4
9	1,134.6	567.8	1,117.6	1,116.6	K	376.3	188.6	359.2	358.2	3
10	1,235.7	618.3	1,218.6	1,217.7	T	248.2		231.1	230.1	2
11	1,381.8	691.4	1,364.7	1,363.8	K	147.1		130.1	129.1	1

C

D

B	B Ions	B+2H	B–NH3	B–H2O	AA	Y Ions	Y+2H	Y–NH3	Y–H2O	Y
1	164.1			146.1	Y	1,152.6	576.8	1,135.6	1,134.6	9
2	261.1			243.1	P	989.6	495.3	972.5	971.6	8
3	374.2			356.2	I	892.5	446.8	875.5	874.5	7
4	473.3			455.3	V	779.4	390.2	762.4	761.4	6
5	602.3			584.3	E	680.4		663.3	662.4	5
6	730.4	365.7	713.4	712.4	Q	551.3		534.3	533.3	4
7	893.4	447.2	876.4	875.4	Y	423.3		406.2	405.2	3
8	1,006.5	503.8	989.5	988.5	L	260.2		243.2	242.2	2
9	1,152.6	576.8	1,135.6	1,134.6	K	147.1		130.1	129.1	1

Figure 4-figure supplement 5

A

B

B	B Ions	B+2H	B–NH3	B–H2O	AA	Y Ions	Y+2H	Y–NH3	Y–H2O	Y
1	100.1				V	1,720.9	861.0	1,703.9	1,702.9	14
2	263.1			245.1	Y	1,621.9	811.4	1,604.8	1,603.9	13
3	392.2			374.2	E	1,458.8	729.9	1,441.8	1,440.8	12
4	505.3			487.3	I	1,329.8	665.4	1,312.7	1,311.7	11
5	618.3			600.3	L	1,216.7	608.8	1,199.6	1,198.7	10
6	715.4	358.2		697.4	P	1,103.6	552.3	1,086.6	1,085.6	9
7	816.5	408.7		798.4	T	1,006.5	503.8	989.5	988.5	8
8	963.5	482.3		945.5	F	905.5	453.2	888.5	887.5	7
9	1,078.5	539.8		1,060.5	D	758.4	379.7	741.4	740.4	6
10	1,177.6	589.3		1,159.6	V	643.4	322.2	626.6	625.4	5
11	1,290.7	645.9		1,272.7	L	544.3	272.7	527.3	526.3	4
12	1,427.8	714.4		1,409.7	H	431.2	216.1	414.2	413.2	3
13	1,574.8	787.9		1,556.8	F	294.2		277.2	276.2	2
14	1,720.9	861.0	1,703.9	1,702.9	K	147.1		130.1		1

C

D

B	B Ions	B+2H	B–NH3	B–H2O	AA	Y Ions	Y+2H	Y–NH3	Y–H2O	Y
1	100.1				V	1,720.9	861.0	1,703.9	1,702.9	14
2	263.1			245.1	Y	1,621.9	811.4	1,604.8	1,603.9	13
3	392.2			374.2	E	1,458.8	729.9	1,441.8	1,440.8	12
4	505.3			487.3	I	1,329.8	665.4	1,312.7	1,311.7	11
5	618.3			600.3	L	1,216.7	608.8	1,199.6	1,198.7	10
6	715.4	358.2		697.4	P	1,103.6	552.3	1,086.6	1,085.6	9
7	816.5	408.7		798.4	T	1,006.5	503.8	989.5	988.5	8
8	963.5	482.3		945.5	F	905.5	453.2	888.5	887.5	7
9	1,078.5	539.8		1,060.5	D	758.4	379.7	741.4	740.4	6
10	1,177.6	589.3		1,159.6	V	643.4	322.2	626.4	625.4	5
11	1,290.7	645.9		1,272.7	L	544.3	272.7	527.3	526.3	4
12	1,427.8	714.4		1,409.7	H	431.2	216.1	414.2	413.2	3
13	1,574.8	787.9		1,556.8	F	294.2		277.2	276.2	2
14	1,720.9	861.0	1,703.9	1,702.9	K	147.1		130.1		1

Figure 4-figure supplement 6

Figure 5

A

SDS-PAGE gel

B

Phos-tag gel

Figure 6

Figure 6-figure supplement 1

Figure 7