Title:

Identifying spawning timing and locations of jack mackerel in a semi-closed bay using environmental DNA

analysis

Running title: Identifying Jack Mackerel Spawning with eDNA

Authors: Satsuki Tsuji[1,2]*, Hiroaki Murakami[3,4], Reiji Masuda[3] (*corresponding author)

Affiliations:

[1] Biosphere Informatics Laboratory, Graduate School of Informatics, Kyoto University, Yoshidahonmachi,

Sakyo-ku, Kyoto 606-8501, Japan

[2] Graduate School of Science, Kyoto University, Kitashirakawa-Oiwakecho, Sakyo-ku, Kyoto 606–8502, Japan

[3] Maizuru Fisheries Research Station, Kyoto University, Nagahama, Maizuru, Kyoto 625-0086, Japan

[4] Graduate School of Agricultural Science, Tohoku University, Aoba-ku, Sendai, 980-8572, Japan

e-mail addresses:

S.T.; satsuki.may425@gmail.com (Tel&Fax, +81–(75)–753–3131)

H.M.; hiroaki.murakami.d2@tohoku.ac.jp

R.M.; masuda.reiji.3w@kyoto-u.ac.jp

Abstract

Understanding spawning ecology is vital for species/population management and conservation. However,

conventional surveys are generally time- and labour-demanding and invasive. To overcome these challenges,

environmental DNA (eDNA) analysis has emerged as a promising spawning survey method. This study aimed to

identify the spawning season, times, and locations of Japanese jack mackerel (*Trachurus japonicus*) through

eDNA analysis and to investigate the utility of eDNA analysis in spawning surveys. We expected that eDNA

concentration and nuclear/mitochondrial DNA ratio should change overnight due to sperm-derived eDNA as this

species is a nocturnal spawner. The eDNA concentrations at sunset and sunrise at three sites in Maizuru Bay were

compared monthly for one year. Our results showed significant increases in eDNA concentrations and ratios at all

sites in July, suggesting potential spawning occurrences. At the site with a particularly large increase in

concentration, temporal monitoring showed a diurnal eDNA peak (9:00 PM–12:00 midnight), likely reflecting the

jack mackerel spawning time window. Subsequently, our eDNA-based estimations were supported by the

successful capture of their eggs using a plankton net. This is the first report providing evidence of saltwater fish

spawning using eDNA analysis, underscoring the usefulness of this approach for spawning detection.

Keywords: spawning, eDNA, egg, sperm, saltwater fish, Maizuru Bay, qPCR

Introduction

Reproductive events are an important factor in the recruitment process of fish populations because the success and magnitude of spawning directly influence population reproduction and future growth (Mertz and Myers, 1994; Scott et al., 2006). Hence, comprehending the time of day and location of fish spawning has long been recognised as a critical challenge for effective species and/or population management and conservation (Danylchuk et al., 2011; Di Muri et al., 2022; Spear et al., 2015). Conventional methods employed by fishermen, researchers, and conservation managers to identify fish spawning involve collecting fish eggs, observing the gonads of parent fish or conducting acoustic telemetry surveys. However, these methods are generally time-consuming and labour-intensive as well as sensitive to monitoring biases and field conditions (ex. weather, time and turbidity) (Caswell et al., 2004; Ko et al., 2013; Vautier et al., 2023). Furthermore, additional mortality is inevitable with survey methods that require the collection of individuals and eggs (Engstedt et al., 2014; Lefort et al., 2015; Wei et al., 2009). Consequently, an efficient and non-invasive method that could solve these conventional challenges would be welcomed as a new approach in fish spawning surveys.

In recent years, environmental DNA (eDNA) analysis has emerged as a rapidly advancing biodiversity assessment method and has entered a new phase where it can estimate ecological events such as spawning as well as the presence of the target species (Ip et al., 2023; Sato et al., 2021; Tsuji et al., 2022b; Wu et al., 2023). Environmental DNA on macroorganisms refers to generic terms for DNA derived from epidermal tissue, faeces, mucus and other sources released from organisms into their habitat (Barnes and Turner, 2016; Merkes et al., 2014). In species that perform in-vitro fertilisation, the reproductive materials such as oocytes, ovarian fluid and sperm released during spawning are also an eDNA source. In eDNA-based spawning surveys, the occurrence of spawning can be identified by observing rapid and significant increases in eDNA concentration and/or changes in nuclear/mitochondrial DNA ratios caused by sperm-derived eDNA (Bylemans et al., 2017; Tsuji et al., 2022b). Since only environmental samples like water need to be collected in the field, there is no impact on the target species or their habitats, and monitoring bias resulting from the surveyor's experience and amount of knowledge

are minimised. These advantages would solve some of the challenges of conventional methods, leading to

increasingly high expectations for eDNA analysis as a novel approach in spawning surveys.

Spawning surveys based on eDNA analysis are still in the developing stage, and their application for fish

is currently limited to freshwater areas (ex. lake, Vautier et al., 2023; Wu et al., 2023; river, Bracken et al., 2019;

Di Muri et al., 2022; Erickson et al., 2016; Saito et al., 2022; Tsuji and Shibata, 2023). Previous studies

investigating fish which spawn in the sea, such as the Japanese jack mackerel (*Trachurus japonicus*) and the

Japanese eel (*Anguilla japonica*), have reported significant increases in eDNA concentrations before and after

spawning in both species when artificially spawned in laboratory conditions (Takeuchi et al., 2019; Tsuji et al.,

2022b). These findings suggest that spawning surveys based on eDNA could be a useful method for investigating

spawning in saltwater fish. The decline of marine biodiversity is a pressing concern, with fish populations

inhabiting coral reefs and seagrass beds estimated to have decreased by 34–70% in the past 40 years until 2012

due to overfishing, ecosystem destruction and climate change (WWF 2015). Considering the importance of

conserving marine biodiversity and managing fisheries resources sustainably, it would be of great value to

investigate the applicability of spawning surveys based on eDNA analysis in saltwater areas, which are affected

by more complex currents than freshwater areas.

The Japanese jack mackerel (*Trachurus japonicus*) is one of the most commercially important resources

in East Asia and dominates in Maizuru Bay, Japan, from spring to autumn (Masuda, 2008; Masuda et al., 2008).

They primarily spawn at the shelf break of the southern East China Sea (Sassa et al., 2006), although small-scale

regional spawning also scatters along the Japanese coast (Nishida, 2006; Xie et al., 2005). Due to the complex

ocean currents affecting the Japanese archipelago, Japanese jack mackerel spawn at different seasons and in

different spawning grounds, contributing to the seasonal stock recruitment in fisheries. For example, they spawn

from February to May in the main spawning ground of the East China Sea and from November to next August in

the coastal waters of Japan (the eastern regions being later) (Nishida, 2006). While the spawning of Japanese jack

mackerel in Maizuru Bay has not been demonstrated, the otolith microstructure of juveniles caught in Kunda Bay,

just west of Maizuru Bay, suggests the existence of a population that may have been spawned near Maizuru Bay

and the southern Sea of Japan (Kanaji et al., 2009). Understanding the spawning season and grounds of Japanese

jack mackerel in Maizuru Bay is essential for fishery resource managers to comprehend population dynamics and

effectively manage resources.

In the present study, three experiments were conducted in Maizuru Bay with the aim of identifying the

spawning season (seasonability), time window (diel spawning periodicity) and location of Japanese jack mackerel

and examining the usefulness of eDNA analysis on spawning surveys. Since they spawn at night, it is assumed

that eDNA concentrations are higher at sunrise than before sunset during the spawning season. Therefore, in

Experiment 1 (Exp. 1), eDNA concentrations in Japanese jack mackerel were investigated once a month, both

before sunset and at sunrise the following day at multiple sites to estimate the spawning season and locations. In

Experiment 2 (Exp. 2), diurnal changes in eDNA concentrations were investigated during the estimated spawning

season to identify the time window of spawning. Finally, in Experiment 3 (Exp. 3), fish eggs were collected using

a plankton net to demonstrate the spawning of Japanese jack mackerel during the estimated spawning season.

Based on the results, we discuss the potential of eDNA analysis as a novel approach for conducting spawning

surveys in the sea.

Materials and Methods

The overview of sampling designs is presented in Fig.1. All experiments were conducted in Maizuru Bay, Japan,

which has a surface area of ca. 11 km^2 (35.481°N, 135. 332°E).

Experiment 1: Estimation of the spawning season and spawning sites

The water sampling survey was conducted in three sites in Maizuru Bay once a month from January 2021 to

January of the following year, on the day of the new moon (Fig. 1, Table S1). We chose a new moon day on the

assumption that a darker environment would be less likely to be attacked by natural enemies and spawning would

be more likely to occur. As Japanese jack mackerel spawn at night; if spawning occurs, it is expected that higher

DNA concentrations are observed at sunrise than before sunset. In each site, 1 L of surface water was collected in

triplicate at two-time points, two hours before sunset and sunrise on the following day (Fig. 1b, Table S1). As

Maizuru Bay is surrounded by mountains and some areas are dark before sunset, the water sampling time was set

at two hours before sunset. The three duplicated samples were treated as independent samples. To preserve eDNA

in water samples, benzalkonium chloride (BAC) was added and agitated well (final concentration, 0.01%;

OSVAN S 10 w/v % benzalkonium chloride, Nihon Pharmaceutical). Collected water samples were promptly

transported to the laboratory and vacuum-filtered using the Millipore Express PLUS disc PES (Polyethersulfone)

philic filters (1 L per filter, pore size 0.45 µm, 47 mm; Merck Millipore) and filter holders (PP-47; ADVANTEC)

within one hour of collection. To avoid contamination, filter holders for the number of samples were

decontaminated by prior soaking in 10% hypochlorite for at least two hours. After filtration of the sunrise samples,

1 L of ultrapure water was filtered in the same way as the collected water samples and used as a filter negative

control (Filt-NC). All filter samples were folded in half, wrapped in aluminium foil and immediately frozen at

−20°C. DNA was extracted from the filter samples, and the nuclear internal transcribed spacer-1 region (ITS1) of

the ribosomal RNA gene of Japanese jack mackerel was quantified using quantitative real-time PCR and

species-specific primer-probe sets (detailed below). For July 2021 samples, as the spawning trend was indicated

from the increase of ITS1 concentration (see results), the mitochondrial cytochrome b gene (cyt b) region was also

quantified using a species-specific primer-probe set to examine the ratio of nuclear DNA (nu-DNA) to

mitochondrial DNA (mt-DNA).

Experiment 2: Estimation of the spawning time window

Time-series water sampling was conducted between 29 and 30 June 2022. It was preferred to conduct Exp. 1 on

the new moon of July when spawning was shown to be possible (see results), but in 2022 the new moon was on

28-29 July. Thus, Exp. 2 was conducted at the end of June, i.e. at a new moon closer to 10 July, the date of the

new moon in July 2021 when Exp. 1 was conducted. At site 2, which was suggested as a possible main spawning

location (see results), 1 L of surface water was collected in triplicate at nine-time points: 12:00 noon, 3:00, 6:00,

9:00 PM, 12:00 midnight, 3:00, 6:00, 9:00 AM, and 12:00 noon (Table S2). Each water sample was added with

BAC (final concentration, 0.01%) and filtered on-site using the same techniques and equipment as in Experiment

1. After filtration of 12:00 samples on 30 June, 1 L of ultrapure water was filtered as a filter negative control

(Filt-NC). All filter samples were folded in half, wrapped in aluminum foil and immediately frozen at −20°C.

DNA was extracted from the filter samples, and ITS1 region of Japanese jack mackerel was quantified using the

same way as in Experiment 1.

Experiment 3: Field collection of Japanese jack mackerel egg

Egg surveys were conducted on 29 (9:10–10:17 AM) and 30 (1:09–1:58 PM) June 2022 at three sites in Maizuru

Bay (dashed lines, L1 to L3, in Fig. 1A). Oblique tawing of a plankton net (diameter 450 mm, mesh 0.3 mm) was

performed three or four times for each site (average 3 min 10 sec trawl per trial, wire length 40 m, wire angle

63–78°). Tables S3 and S4 show the towing conditions for each trawling and water quality at each site. The

collections were pooled for each site and immediately stored on ice. In the laboratory, approximately 0.6–1.2 mm

sized spherical fish eggs were sorted under a stereomicroscope and then fixed with 70% ethanol. For each

collected egg, DNA was extracted using the spin column (EconoSpin, EP-31201; GeneDesign Inc.) and the buffer

reagents supplied in the DNeasy Blood and Tissue kit (Qiagen). Eggs were soaked in a mixture of 180 µL buffer

ATL and 20 µL Proteinase K and incubated at 56°C for 3 hours to dissolve completely. 200 µL of ethanol was

added and mixed well, after which the entire elute was transferred to a spin column. After centrifugation at 6,000

g for 1 min, DNA was purified according to the manufacturer's protocol of the DNeasy Blood and Tissue kit.

Finally, the DNA was eluted in 50 µL of Buffer AE and stored at −20°C until use in the PCR assay. The species

identification was conducted based on the sequencing of the mitochondrial 12S rRNA region with the MiFish

fish-universal primer set (detailed below). Based on current laws and guidelines of Japan relating to animal

experiments on fish, the collection of fish eggs and the use of DNA samples are allowed without any ethical

approvals from any authorities.

DNA extraction from filter samples

A PES filter sample was folded in quarters and placed in the lower part of the spin column (EconoSpin) with the

silica gel membrane preliminarily removed. A mixed extraction buffer comprising 200 µL ultrapure water, 200 µL

Buffer AL (Qiagen), and 20 µL Proteinase K was added to each filter, and the spin columns were incubated for 30

minutes at 56°C. The filter was moved to the top of the spin column and then centrifuged at 6,000 g for 1 min.

600 µL of ethanol was added to the collected liquid and mixed well by pipetting. The DNA in the mixture was

purified using a DNeasy Blood and Tissue kit following the manufacturer's protocol. Finally, the DNA was eluted

in 100 µL of Buffer AE and stored at −20°C until use in the quantitative real-time PCR assay.

Quantitative real-time PCR (qPCR) assays

Quantitative real-time PCRs were performed in triplicate using the Light Cycler 96 system (Roche, Basel,

Switzerland). The ITS and cyt b regions of Japanese jack mackerel were amplified with species-specific

primer-probe sets, which were developed by previous studies (Jo et al., 2019; Yamamoto et al., 2016). The

sequences of each primer-probe set were shown in Table 1. Each qPCR was conducted in a total volume of 15 µL,

containing 900 nM of forward and reverse primers, 125 nM TaqMan probe, 0.075 µL of AmpErase uracil

N-glycosylase (Thermo Fisher Scientific), 7.5 µL of 2 × TaqMan Environmental Master Mix 2.0 (Thermo Fisher

Scientific), and 2 µL of DNA template. A four-step dilution series containing 3×10^1 to 3×10^4 copies of each

target region was used for each qPCR run as quantification standards. As PCR-negative controls, three reactions

were included in all qPCR runs in which 2.0 µL of ultrapure water was added instead of the DNA template. The

qPCR thermal conditions for each target region were as follows: ITS1, 2 min at 50°C, 10 min at 95°C, then 55

cycles of 15 s at 95°C, and 75 s at 60°C; cyt b, 2 min at 50°C, 10 min at 95°C, then 55 cycles of 15 s at 95°C, and

60 s at 60°C. As the ITS1 primer-probe set had a lower amplification efficiency than the cyt b primer-probe set,

the annealing and extension time was increased by 15 s compared to cyt b. To avoid contamination, reagent

preparation and qPCR were carried out in separate rooms. The R^2 values for the standard curve of all qPCR for

ITS1 and cyt b were shown in Table S5. No amplification was observed in any of the Filt-NC and PCR-negative

controls.

Species identification of eggs

To identify fish species of each egg, approximately 170 base pair of 12S rRNA region was amplified by a

two-step tailed PCR approach with two MiFish fish universal primer sets (Table S6, Miya et al., 2015). The

first-round PCR (1stPCR) was performed in 12-µL reactions containing 6 µL of 2×KODone PCR Master Mix

(TOYOBO), 1.4 µL of each primer mixture (MiFish-U: MiFish-U2 = 10: 3; 5µM) and 2□µl of template DNA.

There was no PCR replication. The thermal conditions were as follows: 30□cycles of 10 sec at 98°C, 5 sec at

65°C and 5 sec at 68°C. Amplicons of the 1stPCR were purified using Sera-Mag SpeedBeads

Carboxylate-Modified magnetic particles (Hydrophobic) (Cytiva) (PCR product : Sera-Mag beads□=□1 : 0.8).

The purified amplicons were adjusted to 0.1 ng/µL and used as the template in second-round PCR (2ndPCR). The

2ndPCR was performed in 12-µL reactions containing 6.0 µL of 2□×□KAPA HiFi HotStart ReadyMix (KAPA

Biosystems), 2.0 µL of each primer with index (1.8 µM) and 2.0 µL of the purified 1stPCR amplicon. The

sequences of the 2ndPCR primers were shown in Table S6. The thermal conditions were as follows: 3 min at 95°C,

12□cycles of 20□s at 98°C, 15□s at 72°C and 5 min at 72°C. The indexed 2nd PCR products were pooled in

equal volumes, and the target band (approx. 370□bp) was excised using a 2% E-Gel SizeSelect Agarose Gels

(Thermo Fisher Scientific). The 150 bp paired-end sequencing by NovaSeq was outsourced to Novogene. All raw

sequences were deposited in the DDBJ Sequence Read Archive (accession number: DRA016903). The

bioinformatics analysis was performed using the PMiFish pipeline (https://github.com/rogotoh/PMiFish). The top

BLAST hit with a sequence identity□≥□98% was identified as the species of the egg.

Statistical analysis

In experiment 1, we performed a Wilcoxon rank sum test to compare the eDNA concentration between sunset and

sunrise samples for each DNA region in July 2021 by wilcox.exact function in exactRankTests package ver.

0.8.34. In experiment 2, to examine diurnal changes in eDNA concentration, the ITS1 concentration observed

during the daytime (12:00 noon, 3:00 PM, 6:00PM of 29 June 2022, 6:00 AM, 9:00 AM, 12:00 noon of 30 June

2022) was compared with those observed at each time point during the nighttime (9:00 PM, 12:00 midnight of 29

of June 2022, 3:00 AM of 30 June 2022) using the Kruskal–Wallis test followed by the Conover's test with Holm

adjustment by kwAllPairsConoverTest function in PMCMRplus package ver.1.9.3. All statistical analyses and

graphic illustrations for this study were conducted using R version 4.2.2 software (R Core Team, 2022). Statistical

values were evaluated at a significance level of $\alpha = 0.05$.

Results

Japanese jack mackerel eDNA was detected and successfully quantified at all sites throughout the year. A

comparison of eDNA concentrations at two hours before sunset (Hereafter "sunset") and sunrise in each month

showed significantly higher ITS1 concentrations at sunrise than at two hours before sunset at all sites in July (Fig.

2 and Fig. 3a). Additionally, in March and October, higher DNA concentrations were found at sunrise than at

sunset in 2/3 of the sites (Fig. 2). In July, both ITS1 and cyt b region showed a significant increase from sunset to

sunrise ($P < 0.001$ for both region, Wilcoxon rank sum test; Fig. 3a,b). The nu-DNA/mt-DNA ratio, known to

increase with the presence of sperm-derived eDNA, , although not significant when all sites were considered, read

an increasing trend from sunset to sunrise at site 2 (Fig. 3c).

Diurnal changes in eDNA concentrations were observed in the time-series water sampling at site 2 (Fig.

4). Environmental DNA concentrations were significantly higher at nighttime than during the daytime ($P < 0.01$

for 9:00 PM and 12:00 midnight, $P < 0.05$ for 3:00 AM, Conover's test), peaking between 21:00 and 24:00.

In the egg collection survey, out of a total of 260 fish eggs successfully identified to species (or taxon)

by sequencing, two eggs were determined to be Japanese jack mackerel eggs. The two Japanese jack mackerel

eggs were recovered from line L3 in Fig. 1a. All remaining eggs were also determined to belong to species

inhabiting Maizuru Bay (Fig. S1A). Ultimately, 25 species were identified, which corresponds to approximately

90% of the expected species diversity based on the accumulation curve of the number of analyzed eggs and

species (Fig. S1b and c, Table S7). Eggs of species with relatively low occurrence frequencies, including Japanese

jack mackerel, were detected from only one of the two days of surveys (30 June 2022).

Discussion

This study successfully identified the spawning season, time window and location of Japanese jack mackerel in

Maizuru Bay through eDNA analysis. The eDNA-based findings were supported by the capture of Japanese jack

mackerel eggs using trawling with a plankton net. This is the first report to provide evidence of saltwater fish

spawning using eDNA analysis, indicating the high potential of this method for spawning detection.

During monthly eDNA monitoring throughout the year, the increase in nighttime eDNA concentrations

observed at all three sites in July was considered to reflect the occurrence of Japanese jack mackerel spawning. At

site 2, both ITS1 and cyt b exhibited the most significant increases in eDNA concentrations between sunset and

sunrise, and the ITS1/cyt b ratio also showed an increasing trend, suggesting the presence of sperm-derived eDNA

due to spawning (Bylemans et al., 2017). In Wakasa Bay, the outer bay of Maizuru Bay, two groups of Japanese

jack mackerel with distinct morphologies and ecological characteristics coexist: the offshore, seasonally migrating

group (black-type) and the coastal, non-migratory group (yellow-type). Observations of body length composition

and otoliths of their juveniles caught in Wakasa Bay suggested that black-type and yellow-type are derived from

eggs spawned between January and March and between June to July, respectively (Azeta and Ochiai, 1962). The

presence of cohorts presumably hatched between mid-June and early August has also been observed in Kunda

Bay, located to the west of Maizuru Bay (Kanaji et al., 2009). Taken together, the July spawning estimated by

eDNA analysis is likely attributed to the coastal yellow-type. Furthermore, while the water depth range in the

main spawning grounds has been estimated between 100 to 200 m (Nishida 2006), Maizuru Bay has a maximum

depth of approximately 30 meters near the bay's mouth and north of Toshima Island, and the depth is generally

shallower towards the back of the bay (cf. Minami et al., 2018). Thus, the depths of Maizuru Bay are not quite

comparable to those of the main spawning grounds, but if Japanese jack mackerel spawn in the bay, they are

likely to prefer deeper sites. However, the tidal currents are fast from the mouth of the bay to Toshima. So,

assuming that they spawned around Toshima where the currents begin to slow down, the most noticeable increase

in eDNA concentration and ratio observed at site 2 seems reasonable. (Miwa and Ikeno, 2008).

Increases in eDNA concentrations between sunset and sunrise were also observed at 2 out of 3 sites in

March and October, but this could be migration-related noise given Japanese jack mackerel ecology. In March,

the quantified eDNA concentrations were relatively low throughout the locations, both at sunset and sunrise,

making it unlikely that spawning behaviour with the release of large amounts of sperm had occurred. On the other

hand, October is the time when juveniles that entered from the outer bay during summer are growing and actively

swimming in the bay (Masuda et al., 2008). Therefore, relatively high eDNA concentrations were observed at

sunset at all sites. Although there was an increase in concentrations at sites 2 and 3, the increase was only 1.4 to

1.9-fold, which is hardly a remarkable increase. The amount of faeces, one of the sources of eDNA, should

increase as fish actively feed in the early morning. Therefore, it is reasonable to assume that the changes in eDNA

concentrations observed in October were most likely caused by fish migration and increased activity.

The significant diurnal changes in eDNA concentrations observed in the time-series sampling suggested

that Japanese jack mackerel mainly spawn between 9:00 PM and 12:00 midnight. This study is the first to identify

their spawning time window in the sea. This result is consistent with the peak spawning window observed in a

previous study when Japanese jack mackerel were artificially spawned in tanks (Nyuji et al., 2013). It is also

consistent with information on other *Trachurus* species, *T. symmetricus* and *T. trachurus*, which spawn at night

(Karlou-Riga and Economidis, 1997; Macewicz and Hunter, 1993). The spawning time window of Japanese jack

mackerel under natural conditions has been predicted to be before dawn, but this has not been clearly reported and

remained unknown (Nishida, 2006; Yoda et al., 2006). It is hoped that intensive egg recovery surveys during the

estimated spawning time window will be carried out in future studies, providing direct observations of eggs

immediately after spawning. Quantifying mt-DNA (cyt b) concentrations in the same samples and observing

time-series changes in nu-DNA to mt-DNA ratios would provide further insight. However, as the observations in

Exp. 1 showed (Fig. 3b), the concentration of cyt b was about 1/10th of ITS1, and daytime samples were

considered difficult to quantify. In addition, due to a change in the author's affiliation (S.T.), it was difficult to

perform the analysis in the same experimental environment and the same qPCR machine as in Exp. 3. Therefore,

these studies remain as future work.

The occurrence of Japanese jack mackerel spawning in Maizuru Bay during July (end of June), as

estimated based on eDNA analysis, was confirmed by the successful direct recovery of their eggs.   The fish egg

recovery surveys conducted in this study were not of an experimental design that provided quantitative data.

However, considering that the Japanese jack mackerel is one of the dominant species in Maizuru Bay, the fact that

only two out of the analysed 260 eggs were recovered likely indicates a low spawning quantity. Based on the

water temperature during the survey (20–25°C, Table SX), the recovered Japanese jack mackerel eggs may have

been floating for a maximum of one day (Ochiai et al., 1982). So, the possibility that the eggs were introduced

from outside the bay cannot be completely ruled out, depending on the tidal current conditions. However, as it

would have taken more than a day for the eggs to flow from the outer bay to Maizuru Bay, we concluded that the

eggs recovered in this study were likely spawned within the bay or at least inshore near the bay mouth.

In conclusion, this study demonstrated the utility of eDNA analysis in saltwater fish spawning surveys

through the identification of the spawning of Japanese jack mackerel in Maizuru Bay. While only Japanese jack

mackerel was focused on in this study, future research could potentially estimate the spawning ecology of

multiple species within the target taxonomic group simultaneously by employing quantitative eDNA

metabarcoding methods (Tsuji et al., 2022a; Ushio et al., 2018). Although eDNA analysis currently does not

provide some of the information available through conventional methods (e.g., size, age, sex, and hybridization of

fish) (Minamoto, 2022), its advantage lies in the ability to efficiently and non-invasively narrow down spawning

seasons, time of day, and locations. These characteristics could significantly alleviate the challenges associated

with spawning surveys in marine areas and foster advancements in the research field.

Acknowledgements

We wish to thank Dr. Keita Suzuki, Ms. Misaki Shiomi, Mr. Hunter Godfrey and Mr. Yoshihito Ogura (Maizuru

Fisheries Research Station, Kyoto University) for their help with the collection of fish eggs in experiment 3. This

study was supported by JSPS KAKENHI Grant Number JP20K15578 and JP19H05641.

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

Authors' contributions

S.T.: Conceptualization, Funding acquisition, Methodology, Fieldwork (all experiments), Molecular analysis,

Visualization, Writing–original draft. H.M.: Methodology, Fieldwork (experiment 1), Writing–review & editing.

R.M.: Conceptualization, Funding acquisition, Methodology, Fieldwork (experiment 3), Writing–review & editing,

Supervision.

Data availability

Full details of the qPCR results for each experiment of the present study are available in the supporting

information (Table S8 and S9). All raw sequences were deposited in the DDBJ Sequence Read Archive (accession

number: DRA016903).

References

Azeta M., Ochiai A., 1962. A study on the race of Jack mackerel found in Wakasa Bay. NIPPON SUISAN
        GAKKAISHI 28, 967–978. https://doi.org/10.2331/suisan.28.967
Barnes, M.A., Turner, C.R., 2016. The ecology of environmental DNA and implications for conservation genetics.
        Conservation genetics 17, 1–17. https://doi.org/10.1007/s10592-015-0775-4

Bracken, F.S.A., Rooney, S.M., Kelly-Quinn, M., King, J.J., Carlsson, J., 2019. Identifying spawning sites and
other critical habitat in lotic systems using eDNA "snapshots": A case study using the sea lamprey
Petromyzon marinus L. Ecology and Evolution 9, 553–567. https://doi.org/10.1002/ece3.4777
Bylemans, J., Furlan, E.M., Hardy, C.M., McGuffie, P., Lintermans, M., Gleeson, D.M., 2017. An environmental
DNA-based method for monitoring spawning activity: A case study, using the endangered Macquarie
perch (*Macquaria australasica*). Methods in Ecology and Evolution 8, 646–655.
https://doi.org/10.1111/2041-210X.12709
Caswell, N.M., Peterson, D.L., Manny, B.A., Kennedy, G.W., 2004. Spawning by lake sturgeon (*Acipenser
fulvescens*) in the Detroit River. Journal of Applied Ichthyology 20, 1–6.
https://doi.org/10.1111/j.1439-0426.2004.00499.x
Danylchuk, A.J., Cooke, S.J., Goldberg, T.L., Suski, C.D., Murchie, K.J., Danylchuk, S.E., Shultz, A.D., Haak,
C.R., Brooks, E.J., Oronti, A., Koppelman, J.B., Philipp, D.P., 2011. Aggregations and offshore
movements as indicators of spawning activity of bonefish (*Albula vulpes*) in The Bahamas. Mar Biol 158,
1981–1999. https://doi.org/10.1007/s00227-011-1707-6
Di Muri, C., Lawson Handley, L., Bean, C.W., Benucci, M., Harper, L.R., James, B., Li, J., Winfield, I.J.,
Hänfling, B., 2022. Spatio-temporal monitoring of lake fish spawning activity using environmental DNA
metabarcoding. Environmental DNA 5, 849–860. https://doi.org/10.1002/edn3.343
Engstedt, O., Engkvist, R., Larsson, P., 2014. Elemental fingerprinting in otoliths reveals natal homing of
anadromous Baltic Sea pike (*Esox lucius* L.). Ecology of Freshwater Fish 23, 313–321.
https://doi.org/10.1111/eff.12082
Erickson, R.A., Rees, C.B., Coulter, A.A., Merkes, C.M., McCalla, S.G., Touzinsky, K.F., Walleser, L., Goforth,
R.R., Amberg, J.J., 2016. Detecting the movement and spawning activity of bigheaded carps with
environmental DNA. Molecular Ecology Resources 16, 957–965.
https://doi.org/10.1111/1755-0998.12533
Ip, Y.C.A., Chang, J.J.M., Tun, K.P.P., Meier, R., Huang, D., 2023. Multispecies environmental DNA
metabarcoding sheds light on annual coral spawning events. Molecular Ecology 32, 6474–6488.
https://doi.org/10.1111/mec.16621
Jo, T., Arimoto, M., Murakami, H., Masuda, R., Minamoto, T., 2019. Particle size distribution of environmental
DNA from the nuclei of marine fish. Environmental Science & Technology 53, 9947–9956.
https://doi.org/10.1021/acs.est.9b02833
Kanaji, Y., Watanabe, Y., Kawamura, T., Xie, S., Yamashita, Y., Sassa, C., Tsukamoto, Y., 2009. Multiple cohorts
of juvenile jack mackerel *Trachurus japonicus* in waters along the Tsushima Warm Current. Fisheries
Research 95, 139–145. https://doi.org/10.1016/j.fishres.2008.08.004
Karlou-Riga, C., Economidis, P.S., 1997. Spawning frequency and batch fecundity of horse mackerel, *Trachurus
trachurus* (L.), in the Saronikos Gulf (Greece). Journal of Applied Ichthyology 13, 97–104.
https://doi.org/10.1111/j.1439-0426.1997.tb00108.x
Ko, H.-L., Wang, Y.-T., Chiu, T.-S., Lee, M.-A., Leu, M.-Y., Chang, K.-Z., Chen, W.-Y., Shao, K.-T., 2013.
Evaluating the accuracy of morphological identification of larval fishes by applying DNA barcoding.
PLOS ONE 8, e53451. https://doi.org/10.1371/journal.pone.0053451
Lefort, M.-C., Boyer, S., Barun, A., Khoyi, A.E., Ridden, J., Smith, V.R., Sprague, R., Waterhouse, B.R.,
Cruickshank, R.H., 2015. Blood, sweat and tears: non-invasive vs. non-disruptive DNA sampling for
experimental biology (No. e1580). PeerJ Inc. https://doi.org/10.7287/peerj.preprints.655v3
Macewicz, B.J., Hunter, R.J., 1993. Spawning frequency and batch fecundity of jack mackerel, *Trachurus
symmetricus*, off California during 1991. Calif. Coop. Oceanic Fish. Invest. Rep. 34, 112–121.

Masuda, R., 2008. Seasonal and interannual variation of subtidal fish assemblages in Wakasa Bay with reference
to the warming trend in the Sea of Japan. Environmental Biology of Fishes 82, 387–399.
Masuda, R., Yamashita, Y., Matsuyama, M., 2008. Jack mackerel *Trachurus japonicus* juveniles use jellyfish for
predator avoidance and as a prey collector. Fisheries science 74, 276–284.
Merkes, C.M., McCalla, S.G., Jensen, N.R., Gaikowski, M.P., Amberg, J.J., 2014. Persistence of DNA in
carcasses, slime and avian feces may affect interpretation of environmental DNA data. PLoS One 9,
e113346.
Mertz, G., Myers, R.A., 1994. Match/mismatch predictions of spawning duration versus recruitment variability.
Fisheries Oceanography 3, 236–245. https://doi.org/10.1111/j.1365-2419.1994.tb00101.x
Minami, K., Sawada, H., Masuda, R., Takahashi, K., Shirakawa, H., Yamashita, Y., 2018. Stage-specific
distribution of Japanese sea cucumber *Apostichopus japonicus* in Maizuru Bay, Sea of Japan, in relation
to environmental factors. Fish Sci 84, 251–259. https://doi.org/10.1007/s12562-017-1174-1
Minamoto, T., 2022. Environmental DNA analysis for macro-organisms: species distribution and more. DNA
Research 29, dsac018. https://doi.org/10.1093/dnares/dsac018
Miwa H., Ikeno H., 2008. Numerical analysis of tidal current and water quality in stratified field on summer in
Maizuru Bay. Annual journal of Hydraulic Engineering 52, 1387–1392.
https://doi.org/10.2208/prohe.52.1387
Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J.Y., Sato, K., Minamoto, T., Yamamoto, S., Yamanaka, H.,
Araki, H., 2015. MiFish, a set of universal PCR primers for metabarcoding environmental DNA from
fishes: detection of more than 230 subtropical marine species. Royal Society open science 2, 150088.
Nishida H., 2006. Reproductive biology of Japanese Jack Mackerel *Trachurus japonicus* and Japanese Sardine
Sardinops melanostictus. Bulletin of Fisheries Research Agency. Supplement 113–118.
Nyuji, M., Fujisawa, K., Imanaga, Y., Kitano, H., Yamaguchi, A., Matsuyama, M., 2013. GnRHa-induced
spawning of wild-caught jack mackerel *Trachurus japonicus*. Fish Sci 79, 251–258.
https://doi.org/10.1007/s12562-013-0599-4
Ochiai, A., Mutsutani, K., Umeda, S., 1982. Development of eggs, larvae and juveniles of Jack Mackerel,
Trachurus japonicus. Japanese Journal of Ichthyology 29, 86–92. https://doi.org/10.11369/jji1950.29.86
R Core Team, 2022. A Language and Environment for Statistical Computing.
Saito, M., Tsuji, S., Nakao, R., Miyazono, S., Akamatsu, Y., 2022. Comparative study on nuclear and
mitochondrial DNA of Ayu *Plecoglossus altivelis* for environmental DNA-based spawning evaluation.
Landscape Ecol Eng. https://doi.org/10.1007/s11355-022-00519-5
Sassa, C., Konishi, Y., Mori, K., 2006. Distribution of jack mackerel (*Trachurus japonicus*) larvae and juveniles in
the East China Sea, with special reference to the larval transport by the Kuroshio Current. Fisheries
Oceanography 15, 508–518. https://doi.org/10.1111/j.1365-2419.2006.00417.x
Sato, M., Inoue, N., Nambu, R., Furuichi, N., Imaizumi, T., Ushio, M., 2021. Quantitative assessment of multiple
fish species around artificial reefs combining environmental DNA metabarcoding and acoustic survey.
Scientific reports 11, 1–14.
Scott, B.E., Marteinsdottir, G., Begg, G.A., Wright, P.J., Kjesbu, O.S., 2006. Effects of population size/age
structure, condition and temporal dynamics of spawning on reproductive output in Atlantic cod (*Gadus*
morhua). Ecological Modelling 191, 383–415. https://doi.org/10.1016/j.ecolmodel.2005.05.015
Spear, S.F., Groves, J.D., Williams, L.A., Waits, L.P., 2015. Using environmental DNA methods to improve
detectability in a hellbender (*Cryptobranchus alleganiensis*) monitoring program. Biological
Conservation 183, 38–45. https://doi.org/10.1016/j.biocon.2014.11.016

Takeuchi, A., Iijima, T., Kakuzen, W., Watanabe, S., Yamada, Y., Okamura, A., Horie, N., Mikawa, N., Miller,
M.J., Kojima, T., 2019. Release of eDNA by different life history stages and during spawning activities of
laboratory-reared Japanese eels for interpretation of oceanic survey data. Scientific reports 9, 1–9.
Tsuji, S., Inui, R., Nakao, R., Miyazono, S., Saito, M., Kono, T., Akamatsu, Y., 2022a. Quantitative environmental
DNA metabarcoding shows high potential as a novel approach to quantitatively assess fish community.
Sci Rep 12, 21524. https://doi.org/10.1038/s41598-022-25274-3
Tsuji, S., Murakami, H., Masuda, R., 2022b. Analysis of the persistence and particle size distributional shift of
sperm-derived environmental DNA to monitor jack mackerel spawning activity. Environ. Sci. Technol. 56,
10754–10763. https://doi.org/10.1021/acs.est.2c01904
Tsuji, S., Shibata, N., 2023. Identification of fish spawning based on size-selected sampling and diurnal
concentration changes of environmental DNA resulting from sperm-release.
https://doi.org/10.1101/2023.06.19.545639
Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., Yamanaka, H., Minamoto, T., Kondoh, M.,
2018. Quantitative monitoring of multispecies fish environmental DNA using high-throughput sequencing.
Metabarcoding and Metagenomics 2, e23297. https://doi.org/10.3897/mbmg.2.23297
Vautier, M., Chardon, C., Goulon, C., Guillard, J., Domaizon, I., 2023. A quantitative eDNA-based approach to
monitor fish spawning in lakes: Application to European perch and whitefish. Fisheries Research 264,
106708. https://doi.org/10.1016/j.fishres.2023.106708
Wei, Q.W., Kynard, B., Yang, D.G., Chen, X.H., Du, H., Shen, L., Zhang, H., 2009. Using drift nets to capture
early life stages and monitor spawning of the Yangtze River Chinese sturgeon (*Acipenser sinensis*).
Journal of Applied Ichthyology 25, 100–106. https://doi.org/10.1111/j.1439-0426.2009.01269.x
Wu, L., Wu, Q., Inagawa, T., Okitsu, J., Sakamoto, S., Minamoto, T., 2023. Estimating the spawning activity of
fish species using nuclear and mitochondrial environmental DNA concentrations and their ratios.
Freshwater Biology 68, 103–114. https://doi.org/10.1111/fwb.14012
Xie, S., Watanabe, Y., Saruwatari, T., Masuda, R., Yamashita, Y., Sassa, C., Konishi, Y., 2005. Growth and
morphological development of sagittal otoliths of larval and early juvenile *Trachurus japonicus*. Journal
of Fish Biology 66, 1704–1719.
Yamamoto, S., Minami, K., Fukaya, K., Takahashi, K., Sawada, H., Murakami, H., Tsuji, S., Hashizume, H.,
Kubonaga, S., Horiuchi, T., Hongo, M., Nishida, J., Okugawa, Y., Fujiwara, A., Fukuda, M., Hidaka, S.,
Suzuki, K.W., Miya, M., Araki, H., Yamanaka, H., Maruyama, A., Miyashita, K., Masuda, R., Minamoto,
T., Kondoh, M., 2016. Environmental DNA as a 'Snapshot' of Fish Distribution: A Case Study of
Japanese Jack Mackerel in Maizuru Bay, Sea of Japan. PLoS ONE 11, e0149786.
https://doi.org/10.1371/journal.pone.0149786
Yoda, M., Mizuta, K., Matsuyama, M., 2006. Induction of ovarian maturetion and ovulation in jack mackerel
Trachurus japonics by human chorionic gonadotropin. Bull. Fish. Res. Agen 16, 15–18.
471
472

Table

Table 1 Species-specific primer-probe set for each target region of Japanese jack mackerel. Each primer-probe set was developed by Yamamoto et al. (2016) (cyt b) and Jo et al. (2019) (ITS1), respectively.

Primers/probe	Target region	Sequences (5'-> 3')	Tm (℃)	Amplicon length (bp)
Tja_Cyt b_F	cyt b	CAG ATA TCG CAA CCG CCT TT	58.7	127
Tja_Cyt b_R	(mt-DNA)	CCG ATG TGA AGG TAA ATG CAA A	57.6	
Tja_Cyt b_Pr		[FAM]-TAT GCA CGC CAA CGG CGC CT-[BHQ1]	67.9	
Tja_ITS1_F	ITS1	GCG GGT ACC CAA CTC TCT TC	60.1	164
Tja_ITS1_R	(nu-DNA)	CCT GAG CGG CAC ATG AGA G	60.5	
Tja_ITS1_Pr		[FAM]-CTC TCG CTT CTC CGA CCC CGG TCG-[BHQ1]	70.8	

Figure legends

Figure 1. Overview of the sampling designs. (a) Study sites in Maizuru Bay, Japan, (b)-(d) Sampling design for each experiment. The water sampling in Exp.1 and the egg surveys in Exp.3 were performed at st. 1 to 3 and L1 to 3, respectively.

Figure 2. Changes in eDNA concentrations of Japanese jack mackerel two hours before sunset and at sunrise on the following day observed in each month. Points indicate mean values and error bars indicate standard deviations. The grey background colour indicates months when eDNA concentrations were higher at sunrise than at sunset at multiple sites.

Figure 3. Comparisons of Japanese jack mackerel (a) ITS1 concentration, (b) cyt b concentration and (c) nu-DNA/mt-DNA ratio between two hours before sunset and at sunrise on the following day in July 2021. Yellow, pink, and blue plots indicate a different site, respectively. P values are indicated by asterisks (Wilcoxon rank sum test; *** $P < 0.001$).

Figure 4. Diurnal changes in eDNA concentration of Japanese jack mackerel ITS1 at site 2 on 29–30 June 2022. Orange and blue box-plot colour indicates the daytime (12:00 noon, 3:00 PM, 6:00 PM, 6:00 AM, 9:00 AM and day2 12:00 noon) and nighttime (9:00 PM, 12:00 midnight and 3:00 AM), respectively. The ITS1 concentration observed during the daytime was compared with those observed at each time point during the nighttime. P values are indicated by asterisks (Conover's test; * $P < 0.05$, ** $P < 0.01$). Sunset and sunrise times are as follows: 7:18 PM and 4:46 AM.

Figure S1. (a) species and their frequency of the collected eggs by egg surveys, (b) the number of detected species per survey date; (c) the number of analysed eggs and species diversity.

(a) Study sites

st. 1

st. 2

Maizuru Bay

L2   L1

L3

Toshima

2km

Japan Sea

Maizuru Bay

Pacific Ocean

600 km

45°N

40°N

35°N

30°N

130°E   140°E

st. 3

(b) Exp. 1: All sites

Jan. 2021 ~ Jan. 2022

New moon night

2h   sunset   sunrise

1L×3

(c) Exp. 2: st. 2

29-30 Jun. 2022, New moon night   24   1 L×3 each time point

Day 1   Day 2

12 3 6 9 12 3 6 9 12

(d) Exp. 3: Three dashed 3 lines in panel (a)

29, 30 Jun. 2022   sorting

DNA extraction & Sequencing

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1