Autophagy adaptors mediate Parkin-dependent mitophagy by forming

sheet-like liquid condensates

Zi Yang[1], Saori R. Yoshii[1], Yuji Sakai[1,2,3], Haruka Chino[1], Roland L. Knorr[1,4,5], Noboru

Mizushima[1,6*]

[1]Department of Biochemistry and Molecular Biology, Graduate School of Medicine, The

University of Tokyo, Tokyo, Japan.

[2]Department of Biosystems Science, Institute for Life and Medical Sciences, Kyoto

University, Sakyo-ku, Kyoto 606-8507, Japan

[3]Interdisciplinary Theoretical and Mathematical Sciences (iTHEMS) Program, RIKEN,

Wako, Saitama 351-0198, Japan

[4]Institute of Biology, Humboldt-Universität zu Berlin, Berlin, Germany

[5]International Research Frontiers Initiative, Institute of Innovative Research, Tokyo Institute

of Technology

[6]International Research Center for Neurointelligence (WPI-IRCN), UTIAS, The University of

Tokyo, Tokyo 113-8655, Japan

*Corresponding author. Department of Biochemistry and Molecular Biology, Graduate School

of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan; Tel:

+81-3-5841-3440; Fax: +81-3-3815-1490; E-mail: nmizu@m.u-tokyo.ac.jp

Abstract

During PINK1 and Parkin-mediated mitophagy, autophagy adaptors are recruited to depolarized mitochondria to promote the selective degradation of mitochondria. Autophagy adaptors such as OPTN and NDP52 bridge mitochondria and autophagosomal membranes by binding to ubiquitinated mitochondrial proteins and autophagosomal ATG8 family proteins. Here, we demonstrate that OPTN and NDP52 form sheet-like phase-separated condensates with liquid-like properties on the surface of ubiquitinated mitochondria. The dynamic and liquid-like feature of OPTN condensates is important for mitophagy activity because reducing the liquidity of OPTN–ubiquitin condensates suppresses the recruitment of ATG9 vesicles and impairs mitophagy. Based on these results, we propose a dynamic liquid-like model of autophagy adaptors, in contrast to a stoichiometric model, to explain their interactions between autophagic membranes (i.e., ATG9 vesicles and isolation membranes) and mitochondrial membranes during Parkin-mediated mitophagy. This model underscores the importance of liquid–liquid phase separation in facilitating membrane–membrane contacts, likely through the generation of capillary forces.

Keywords

autophagy; mitophagy; liquid–liquid phase separation; optineurin; wetting

Introduction

Macroautophagy (hereafter referred to as autophagy) is an intracellular degradation process mediated by a double-membraned organelle termed the autophagosome (Nakatogawa, 2020; Mizushima & Levine, 2020). Cytosolic components such as proteins and organelles are segregated into autophagosomes, which subsequently fuse with lysosomes for degradation. Autophagy can be either non-selective or selective (Lamark & Johansen, 2021; Vargas *et al*, 2023). Mitophagy, which is a selective type of autophagy, specifically degrades mitochondria and can be induced by mitochondrial damage or cellular stresses (Onishi *et al*, 2021; Ganley & Simonsen, 2022). For example, after losing the mitochondrial membrane potential, numerous mitochondrial outer-membrane proteins are ubiquitinated by the PINK1-Parkin system, initiating the recruitment of autophagy adaptors, including optineurin (OPTN), calcium binding and coiled-coil domain 2 (CALCOCO2/NDP52), sequestosome 1 (SQSTM1/p62), Next to BRCA1 gene 1 protein (NBR1), and Tax1 binding protein 1 (TAX1BP1) (Lamark & Johansen, 2021; Vargas *et al*, 2023). These autophagy adaptors bridge the mitochondrial and autophagosomal membranes by interacting with mitochondrial ubiquitinated proteins via a ubiquitin-binding domain (UBD) and with autophagosomal ATG8 family proteins (i.e., LC3 and GABARAP family proteins in mammals) via the LC3-interacting region (LIR). In addition to their role in bridging mitochondria and autophagosomes, OTPN and NDP52 also play crucial roles in inducing mitophagy by recruiting upstream autophagy factors (Lazarou *et al*, 2015); OPTN recruits ATG9 vesicles, whereas NDP52 recruits the ULK1 complex through interaction with FIP200 (Yamano *et al*, 2020; Vargas *et al*, 2023).

Liquid–liquid phase separation (LLPS) has been described in various biological processes (Musacchio, 2022; Hirose *et al*, 2023). LLPS involves the formation of liquid-like condensates through dynamic, multivalent interactions between various molecules that often possess intrinsically disordered regions. Emerging evidence suggests connections between autophagy and LLPS at multiple steps, including autophagosome formation and degradation (Noda *et al*, 2020; Ma *et al*, 2023). Notably, autophagy adaptors, such as p62,

undergo LLPS through multivalent interaction with poly-ubiquitins and form cytosolic

condensates (Zaffagnini *et al*, 2018; Sun *et al*, 2018). These condensates can be degraded

by autophagy, which is termed "fluidophagy" (Agudo-Canalejo *et al*, 2021). Furthermore,

recent studies showed that p62 undergoes LLPS on mitochondria and lysosomes that are

subjected to selective autophagy (Peng *et al*, 2021; Gallagher & Holzbaur, 2023).

Generally, phase-separated condensates can deform when they contact a rigid surface

in order to minimize the overall energy of the system, a phenomenon known as "wetting"

(Kusumaatmaja *et al*, 2021b; Gouveia *et al*, 2022). The wetting behavior of intracellular

condensates is determined by the interfacial tensions between the condensate–membrane,

condensate–cytosol, and cytosol-membrane interfaces. Depending on the relative

strengths of these interfacial tensions, the condensates exhibit partial wetting, complete

wetting, or de-wetting. In contrast, elastic surfaces, including membranes to which phase-

separated condensates adhere, also deform (Kusumaatmaja *et al*, 2021a, 2021b; Gouveia

et al, 2022). This indeed happens during fluidophagy; autophagosomal membranes bend

along the surface of p62 condensates (Agudo-Canalejo *et al*, 2021). Deformation of a

membrane by a phase-separated condensate is determined by the energy balance

between membrane deformation and the surface tension of the condensate, which is

known as elastocapillarity (Style *et al*, 2017; Kusumaatmaja *et al*, 2021b).

By forming capillary bridges, wetting droplets can also induce adhesion of surfaces

(Wexler *et al*, 2014). Whether LLPS and capillary bridges also contribute to the adhesion of

cellular membranes is not presently known. Therefore, we hypothesized that phase-

separated condensates of autophagy adaptors can wet both mitochondria and autophagic

membranes, including ATG9 vesicles and isolation membranes (also called phagophores),

thereby promoting their contact during Parkin-mediated mitophagy. In this study, we employ

live-cell imaging and a mathematical model to provide evidence that the autophagy

adaptors form phase-separated condensates on the surface of the mitochondrial

membrane, resulting in sheet-like condensates that cover the surface of ubiquitinated

mitochondria upon mitophagy induction. These condensates accumulate between

mitochondria or between mitochondria and autophagic membranes, exhibiting a dynamic

nature with a liquid-like property. Furthermore, we demonstrate the essential role of these

dynamic condensates in the recruitment of ATG9 vesicles and the initiation of mitophagy.

Based on these results, we propose a dynamic liquid-like model, rather than a

stoichiometric model, to describe the roles of autophagy adaptors in Parkin-mediated

mitophagy.

Results

Autophagy adaptors show distinct distributions during Parkin-mediated mitophagy

To better understand the role of each autophagy adaptor, we analyzed their localization

during Parkin-mediated mitophagy. When mitophagy was induced by treatment with the

mitochondrial uncoupler carbonyl cyanide *m*-chlorophenylhydrazone (CCCP) in HeLa cells

expressing exogenous Parkin, the autophagy adaptors p62, OPTN, NBR1, NDP52, and

TAX1BP1 translocated to mitochondria as previously observed (Fig. 1A and B) (Geisler *et*

al, 2010; Wong & Holzbaur, 2014; Heo *et al*, 2015; Moore & Holzbaur, 2016; Gallagher &

Holzbaur, 2023). Ubiquitin and all of these autophagy adaptors were distributed evenly on

the surface of separate mitochondria (Mt; Fig. 1A). In contrast, these adaptors exhibited

distinct localization patterns during the formation of isolation membranes on mitochondria

(Mt–IM; Fig. 1B and C). The signals of OPTN and NDP52 were enriched in areas where

LC3B signals colocalized, while they were mostly absent from the LC3B-negative side of

the same mitochondria (Fig. 1B and C). Endogenous OPTN and NDP52 also showed clear

enrichment on the LC3B-positive areas in comparison to the LC3B-negative side,

confirming that this localization is not caused by the overexpression of adaptors (Fig. 1D).

However, this inhomogeneous distribution pattern was not observed for ubiquitin, p62,

NBR1, and TAX1BP1 (Fig. 1B and C). Unlike OPTN and NDP52, p62 showed enrichment

between clustered mitochondria (Mt-Mt), which is consistent with previous reports (Fig.

EV1A and B) (Wong & Holzbaur, 2014). These accumulation patterns cannot be explained

by the stoichiometric interaction of the autophagy adaptors with ubiquitinated proteins that

are evenly distributed on mitochondria.

The localization of autophagy adaptors may be affected by their interaction with other

adaptors (Turco *et al*, 2021; Gubas & Dikic, 2022). To examine the localization of each

adaptor on its own, we used HeLa cells lacking all five autophagy adaptors (penta KO cells)

to exclude the effect of other adaptors (Lazarou *et al*, 2015). In penta KO cells, OPTN and

NDP52 exhibited significant enrichment in LC3B-positive areas, similar to that observed in

wild-type cells, suggesting that this accumulation occurred independently of heterologous

interaction with other adaptors (Fig. 1E and F). Mitophagy was not restored by exogenous

expression of p62 or NBR1 in penta KO cells, and thus colocalization with LC3 could not be

tested for these two adaptors (Lazarou *et al*, 2015). The accumulation of p62 between

clustered mitochondria was also observed in penta KO cells (Fig. EV1C and D). Given the

uniform distribution of ubiquitinated proteins on mitochondrial surfaces, these data suggest

that autophagy adaptors exhibit non-stoichiometric enrichment between membranes.

Hereafter, we used penta KO cells to study each autophagy adapter individually

OPTN and NDP52 show a dynamic exchange between the mitochondrial surface and

cytosol

We hypothesized that the non-stoichiometric distribution of the autophagy adaptors on

ubiquitinated mitochondria could be explained by LLPS. Typical condensates generated by

LLPS exhibit rapid exchange of their components, which is often demonstrated by

fluorescence recovery after photobleaching (FRAP) analysis (Taylor *et al*, 2019;

McSwiggen *et al*, 2019). We first conducted FRAP experiments for separate (i.e.,

unclustered) mitochondria in CCCP-treated cells. As expected, ubiquitin showed virtually no

recovery because it was conjugated to mitochondrial membrane proteins (Fig. 2A and B).

Additionally, p62 recovered only slightly, suggesting a minute exchange between the

mitochondrial surface and cytosol (Fig. 2A and B). In contrast, OPTN and NDP52 showed a

rapid recovery, indicating a dynamic exchange of these adaptors between the mitochondrial

surface and the cytosol (Fig. 2A and B).

Next, we examined the dynamics of adaptors between mitochondria and isolation

membranes. OPTN and NDP52 showed partial recovery (Fig. 2C and D). Although this

result indicates some exchange of these adaptors between membranes, the incomplete

recovery suggests the existence of a gel-like or immobile fraction. Treatment with 1,6-

hexanediol, which can be used to dissolve condensates, dispersed OPTN from

mitochondria without isolation membranes, suggesting that OPTN accumulation is

supported by weak interactions (Fig. 2E). These results indicate that OPTN and NDP52 on

ubiquitinated mitochondria are mobile and can quickly exchange between the mitochondrial

surface and the cytosol, supporting the hypothesis that they form phase-separated

condensates on damaged mitochondria.

Mathematical models of condensate formation and localization of autophagy

adaptors

To simulate the behavior of autophagy adaptors in the context of phase-separated

condensates, we developed mathematical models (see Materials and Methods for details).

First, we considered the case of two mitochondria approaching each other (Fig. 3A). In

Parkin-mediated mitophagy, various proteins in the mitochondrial outer membrane are

ubiquitinated. We modeled the mitochondrial outer membrane as a circle with a diameter of

$D_{Mt} = 600$ nm. On these mitochondria, we assumed a ring-shaped ubiquitin layer with a

thickness $w_{Ub} = 20$ nm and area-fraction (hereafter referred to as concentration) $\phi_{Ub} = 0.1$

(Milo & Phillips, 2015; Berry *et al*, 2018) (Fig. 3A). The closest distance between the two

mitochondrial outer membranes was set to d_{Mt} (Fig. 3A). Here, we considered the

dynamics of concentration changes of the adapter proteins and solvent components. We

assumed that mitochondria were very large compared with proteins and did not move on a

similarly short time scale, and that ubiquitin was tightly bound to mitochondrial membranes

and immobile, as suggested by the FRAP experiment (Fig. 2B). The adaptor proteins were

assumed to diffuse freely in solution and to bind to ubiquitin and themselves (self-

interactions) with strengths χ_{Ub} and χ_{self}, respectively.

Using the mathematical model, we assessed how the condensates deformed when the

distance between two mitochondria d_{Mt} was changed, where $\chi_{Ub} = 12k_BT$ and $\chi_{self} =$

$3k_BT$, with the Boltzmann constant k_B and temperature T. When two mitochondria were far

apart ($d_{Mt} = 900$ nm), the model predicted that adaptors cover the entire mitochondrial

surface (Fig. 3B). When the mitochondria were close enough to come into contact ($d_{Mt} =$

60 nm), the adaptors moved to the area between the mitochondria and filled the cleft

between them (arrowheads, Fig. 3B).

Next, we considered the case in which a growing isolation membrane surrounded and

enclosed a single mitochondrion (Fig. 3C and D). A mitochondrion with ubiquitin was

modeled as mentioned above, and the isolation membrane was modeled as a region of

thickness $w_{IM} = 50$ nm bound by two semicircles with the same center as the

mitochondrion. The closest distance between the isolation membrane and the

mitochondrial outer membrane was $d_{IM} = 100$ nm, and the isolation membrane bent at an

angle α to surround the mitochondrion. The surface of the isolation membrane was

assumed to be covered by an LC3 region with a thickness $w_{LC3} = 20$ nm and concentration

$\phi_{LC3} = 0.1$. The adaptors interacted with both LC3 and ubiquitin with strengths χ_{LC3} and

χ_{Ub}, respectively.

Using this model, we determined the distribution of the adaptors when the isolation

membrane appeared and elongated (changing the angle α surrounding the mitochondrion).

The interaction strengths were set to $\chi_{Ub} = \chi_{LC3} = 12 k_B T$ and $\chi_{self} = 3 k_B T$. In the absence

of the isolation membrane, the adaptors wet the entire mitochondrial surface uniformly (Fig.

3D). However, with the appearance of the isolation membrane, the adaptors remobilized to

the area between the isolation membrane and mitochondrion, and this adaptor-enriched

region elongated together with the isolation membrane. These modeling results suggest

that the distribution of the adaptors on a mitochondrion can change upon contact with

another mitochondrion or an isolation membrane.

OPTN condensates redistribute upon membrane contact

Our mathematical models predict that the OPTN distribution changes when mitochondria

cluster or are engulfed by isolation membranes. We then validated these predictions by

using live-cell imaging. When two small mitochondria with uniform OPTN distributions on

their surface approached each other, OPTN redistributed and formed a smooth surface that

covered the two adjacent mitochondria (Fig. 4A–D, Movie EV1). Notably, strong OPTN

enrichment was observed in the cleft between the two mitochondria (arrowheads, Fig. 4A

and C), showing a pattern distinct from that of the outer membrane protein Omp25 (OPTN

signal peaks localized outside of Omp25 signal peaks) (Fig. 4B and D). This phenomenon

was apparent when the size of both (Fig. 4A) or one (Fig. 4C, Movie EV1) of the two

mitochondria was less than 1 μm. Although OPTN formed sheet-like rather than spherical

condensates, this phenomenon appeared to be similar to condensate coalescence, which

is one of the hallmarks of liquid-like condensates (Hyman *et al*, 2014).

Furthermore, when an isolation membrane started to engulf a mitochondrion, OPTN,

which was initially distributed uniformly on the mitochondrial surface, underwent

redistribution to the area contacting the isolation membrane (Fig. 4E, Movie EV2). OPTN

signals on the isolation membrane-negative side of the mitochondrion diminished during

this process (Fig. 4E and F). The OPTN-enriched region expanded together with the

isolation membrane thereafter (Fig. 4E). These live-cell observations are consistent with

the predictions of our mathematical models (Fig. 3). Taken together, our experimental data

support the hypothesis that OPTN forms sheet-like phase-separated condensates on the

surface of ubiquitinated mitochondria, exhibiting liquid-like properties.

Liquid-like property of OPTN condensates is required for mitophagy

To investigate the importance of the liquid-like nature of OPTN condensates on

mitochondria, we sought to reduce its liquidity by strengthening the interaction but reducing

the multivalency between ubiquitin and GFP–OPTN constructs. To achieve this, we utilized

the anti-green fluorescent protein (GFP) nanobody, which exhibits a high binding affinity to

GFP (Rothbauer *et al*, 2008). Fusing the anti-GFP nanobody to the N-terminus of ubiquitin

(nanobody–mRuby–Ub) (Fig. 5A, bottom panel) resulted in significantly enhanced binding

to GFP–OPTN (Fig. 5B). Deleting the ubiquitin-binding domain in ABINs and NEMO (UBAN

domain) from OPTN (GFP–OPTNΔUBAN) eliminated the interaction with ubiquitinated

proteins (Fig. 5B). Wild-type GFP–OPTN accumulated on ubiquitinated mitochondria, but

GFP–OPTNΔUBAN failed to do so (Fig. 5A, top and middle panels). In contrast, the

expression of nanobody–mRuby–Ub rescued the recruitment of GFP–OPTNΔUBAN to

mitochondria (Fig. 5A, lower panel). FRAP experiments revealed that, compared with wild-type GFP–OPTN (with mRuby–Ub, not nanobody-mRuby–Ub), GFP–OPTNΔUBAN showed a substantial decrease in recovery on mitochondria when co-expressed with nanobody–mRuby–Ub, indicating a reduction in its dynamic properties (Fig. 5C).

We then evaluated the effect of impaired OPTN liquidity on mitophagy using the recently developed HaloTag cleavage assay (Yim *et al*, 2022). This assay utilizes the ligand-dependent conformational change of HaloTag (Halo); ligand-free Halo is efficiently degraded in lysosomes, whereas ligand-bound Halo becomes resistant to lysosomal degradation. When Halo is expressed in the mitochondrial matrix by fusing to the mitochondrial presequence of Fo-ATPase subunit 9 and SNAP-tag (mtHalo–SNAP), we can measure the extent of mitochondrial degradation by quantifying the amount of processed free Halo out of the total amount of Halo (mtHalo–SNAP + processed Halo). Mitophagy activity, which was impaired in penta KO cells, was restored by the expression of GFP–OPTN (Fig. 5D and E). However, GFP–OPTNΔUBAN failed to restore mitophagy, even when it was co-expressed with nanobody–mRuby–Ub to rescue ubiquitin binding and mitochondrial localization (Fig. 5A, B, D, and E). These data suggest that the localization of OPTN on mitochondria and its binding to ubiquitin alone is insufficient for mitophagy, and the liquidity of OPTN is important for effective mitophagy.

Liquid-like property of OPTN condensates is required for the recruitment of ATG9 vesicles

ATG9 vesicles are considered to be one of the sources of autophagosomal membranes (Yamamoto *et al*, 2012). Upon mitophagy induction, ATG9 vesicles accumulate at the sites of autophagosome formation (Itakura *et al*, 2012). OPTN plays a crucial role in this process by recruiting ATG9 vesicles to induce mitophagy (Yamano *et al*, 2020). In general, liquid-like condensates can be involved in clustering small vesicles such as synaptic vesicles, which exemplify complete wetting (Milovanovic & De Camilli, 2017; Sansevrino *et al*, 2023). Indeed, ATG9 vesicles are incorporated into condensates containing a glaucoma-

associated OPTN mutant (O'Loughlin *et al*, 2020) and synapsin (Park *et al*, 2023). We,

therefore, hypothesized that the liquid-like properties of OPTN condensates on

mitochondria facilitate the recruitment of ATG9 vesicles. Consistent with previous reports

(Itakura *et al*, 2012; Yamano *et al*, 2020), ATG9A was recruited to depolarized mitochondria

in cells expressing GFP–OPTN (Fig. 6A and B). However, this ATG9A recruitment was

almost completely abolished in cells expressing nanobody–mRuby–Ub and GFP–

OPTNΔUBAN (Fig. 6A and B), even though GFP–OPTNΔUBAN still retains the ATG9A-

interacting domain and the ability to bind with the ATG9 vesicles (Fig. 6C and D). Taken

together, these findings suggest that the liquidity of sheet-like OPTN condensates on the

mitochondrial surface is crucial for recruiting ATG9 vesicles and executing mitophagy.

Discussion

LLPS by autophagy adaptors during Parkin-mediated mitophagy

In the present study, we discovered that autophagy adaptors accumulate on ubiquitinated mitochondria with non-stoichiometric distribution compared with ubiquitin signals. They displayed characteristics consistent with phase-separated structures, including dynamic exchange with cytosol and redistribution upon coalescence. Notably, we observed the enrichment of OPTN at the isolation membrane-positive region, which expanded along with the elongation of the isolation membrane. These findings suggest that phase-separated adaptors facilitate the effective engulfment of mitochondria by the isolation membrane through the wetting effect and capillary forces (Fig. 6E, right panel). Moreover, the liquidity of OPTN condensates is critical for the localization of ATG9 vesicles onto ubiquitinated mitochondria, indicating that partial or complete wetting of phase-separated adaptors to ATG9 vesicles may be crucial for their recruitment to and/or retention on mitochondria (Fig. 6E, left panel). Therefore, we propose that LLPS of autophagy adaptors plays a critical role in two distinct steps of mitophagy by producing capillary forces, first, at the initiation step by recruiting and retaining the ATG9 vesicles in the condensates and, second, at the autophagosome elongation step by facilitating the attachment between ubiquitinated mitochondria and the isolation membrane.

LLPS in bulk and selective autophagy

Recent reports have shed light on the importance of LLPS in selective autophagy (Noda *et al*, 2020). In lysophagy, p62 undergoes LLPS on lysosomes to facilitate efficient lysophagy together with HSP27, which maintains the liquidity of p62 and prevents its gelation (Gallagher & Holzbaur, 2023). This may also be mediated by the wetting between damaged lysosomes and the isolation membrane by phase-separated p62. Indeed, p62 itself forms phase-separated condensates along with ubiquitinated proteins, producing capillary forces by the wetting effect to facilitate the engulfment by the isolation membrane (Agudo-

Canalejo *et al*, 2021). Similarly, Ape1, the cargo of the cytosol-to-vacuole targeting (Cvt)

pathway in yeast, forms gel-like condensates with Atg19 on the surface of Ape1

condensates to be sequestered by the isolation membrane (Yamasaki *et al*, 2020). This

association could also be mediated by partial wetting. These examples together with our

observations highlight the importance of LLPS and the wetting phenomena in various

selective autophagy processes.

LLPS may also be important for non-selective bulk autophagy. In yeast, the Atg1

complex (the ULK1 complex in mammals) forms pre-autophagosomal structures (PASs),

which have been shown to be condensates driven by LLPS (Fujioka *et al*, 2020). Phase

separation in this context provides a high local concentration of the Atg1 complex to

support the autoactivation of Atg1 kinase (Yamamoto *et al*, 2016). Mammalian autophagy

factors also accumulate at the autophagosome formation site; FIP200, a subunit of the

autophagy-initiation complex, undergoes LLPS triggered by calcium transients (Zheng *et al*,

2022). The liquid-like nature of these autophagy-initiating structures may also be important

for the recruitment of Atg9/ATG9 vesicles through the wetting effect.

The importance of phase separation during mitophagy

In contrast to Parkin-mediated mitophagy, hypoxia-induced mitophagy does not involve

autophagy adaptors, and instead, the outer mitochondrial membrane proteins NIX (also

known as BNIP3L) and BNIP3 directly interact with ATG8 homologs on the isolation

membrane (Novak *et al*, 2010; Onishi *et al*, 2021). In this context, ATG8 proteins interact

with NIX or BNIP3, likely in a one-to-one manner. This raises the problem of why phase

separation of autophagy adaptors is essential for Parkin-mediated mitophagy. One

possibility is that the quick induction and execution of Parkin-mediated mitophagy requires

amplification of the reactions through increased local concentration of adaptors and

autophagy-related proteins facilitated by LLPS. This is consistent with the fact that

mitophagy-inducing signals are also amplified by a mechanism involving phosphorylated

ubiquitin (Kane *et al*, 2014; Kazlauskaite *et al*, 2014; Koyano *et al*, 2014). Alternatively,

Parkin-mediated mitophagy requires a high level of precision and selectivity; mitophagy

should be induced only for damaged mitochondria, which is ensured by the recruitment and

retention of ATG9 vesicles only on phase-separated adaptor-positive mitochondria. This

strict specificity would enable proper quality control of the mitochondria. Further study

would be required to elucidate the functions and physiological importance of LLPS by

autophagy adaptors in mitophagy and beyond.

346   **Materials and Methods**

347

348   **Cell culture and generation of stable cell lines**

349   HeLa cells, HeLa penta-knockout (penta KO) cells, and human embryonic kidney (HEK)

350   293T cells authenticated by RIKEN were cultured in Dulbecco's modified Eagle's medium

351   (DMEM; D6546; Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS; 173012;

352   Sigma-Aldrich) and 2 mM L-glutamine (25030-081; Gibco) in a 5% CO_2 incubator at 37°C.

353   For stable expression, retrovirus was produced with HEK293T cells transfected with pMRX-

354   IP-based, pMRX-IB-based, pMRX-IU-based, or pMRX-IN-based retroviral plasmids, pCG-

355   VSV-G, and pCG-gag-pol by using Lipofectamine 2000 (11668019; Thermo Fisher

356   Scientific). After transfection, cells were further incubated at 37°C for 24h. The viral

357   supernatant was collected by filtration through a 0.45-μm filter unit (Ultrafree-MC; Millipore)

358   and then used for infection. Cells were plated onto 6-cm dishes 18 h before infection, and

359   the medium was replaced with viral supernatant diluted 1.5-fold with 8 μg/mL polybrene

360   (H9268; Sigma-Aldrich). Two days later, cells were selected in a medium containing 2

361   μg/mL puromycin (P8833; Sigma-Aldrich), 4 μg/mL blasticidin S hydrochloride (022-18713;

362   FUJIFILM Wako Pure Chemical Corporation), 1.5 mg/mL neomycin, or 250 μg/mL zeocin

363   (R25005; Thermo Fisher Scientific).

364

365   **Plasmids**

366   The pMRX-IPU, pMRX-IBU and pMRX-INU plasmids were generated by modifying the

367   multi-cloning site of pMRX-IP (Saitoh *et al*, 2002; Kitamura *et al*, 2003), pMRX-IB, and

368   pMRX-IN, respectively. DNA fragments encoding ubiquitin, p62 (NM_003900.5), OPTN

369   (NM_001008211.1), NBR1 (NM_001291571.2), NDP52 (NM_001261391.2), TAX1BP1

370   (NM_001206901.1), ATG9A (NM_001077198.3), LC3A (NM_032514.4), and LC3B

371   (NM_022818.5) were inserted into pMRX-IPU, pMRX-INU, or pMRX-IBU. DNAs encoding

372   the HA epitope, monomeric enhanced GFP with the A206K mutation (mGFP), codon-

373   optimized ultra-stable GFP (muGFP) (Scott *et al*, 2018), codon-optimized mRuby3 (Bajar *et*

al, 2016), HaloTag7 (G1891; Promega), 3×FLAG, and SNAP-tag (New England BioLabs,

N9181S) were used for tagging. The mitochondrial presequence of *Neurospora crassa* Fo-

ATPase subunit 9 (residues 1–69) was added to HaloTag7-SNAP to deliver the reporter into

the mitochondrial matrix (mtHalo-SNAP) (Eura, 2003). Truncated OPTN (OPTNΔUBAN, aa

445–502) was prepared by PCR-mediated site-directed mutagenesis. The resulting

plasmids were sequenced.

Antibodies and reagents

The primary antibodies used in this study were rabbit polyclonal anti-OPTN (Proteintech,

10837-AP), rabbit polyclonal anti-ATG9A (MBL, PD042), mouse monoclonal anti-Halo

(Promega, G9211), mouse monoclonal anti-HSP90 (BD Transduction Laboratories,

610419), and rabbit polyclonal anti-GFP (Thermo Fisher Scientific, A6455) antibodies. The

secondary antibodies used were HRP-conjugated goat polyclonal anti-rabbit IgG (Jackson

ImmunoResearch Laboratories, 111-035-144) and HRP-conjugated goat polyclonal anti-

mouse IgG (Jackson ImmunoResearch Laboratories, 115-035-003) antibodies. To induce

mitophagy, HeLa cells were treated with 10 μM carbonyl cyanide *m*-chlorophenyl hydrazine

(CCCP; Sigma-Aldrich) for 45 min, or 10 μM oligomycin (Cabiochem, 495455-10MGCN)

and 4 μM antimycin A (Sigma-Aldrich, A8674) for 18 h. After cells had been treated with

oligomycin and antimycin A for >6 h, 10 μM Q-VD-OPH (SM Biochemicals, SMPH001) was

added to block apoptotic cell death. To inhibit the formation of isolation membranes in the

1,6-hexanediol (Sigma-Aldrich, 240117-50G) treatment experiment, 20 μM wortmannin

(Sigma-Aldrich, W1628-1MG) was added into the medium.

Immunoprecipitation and immunoblotting

Cell lysates were prepared in a lysis buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM

EDTA, 1% Triton X-100, EDTA-free protease inhibitor cocktail [19543200; Roche]). After

centrifugation at 17,700 × *g* for 10 min, the supernatants were subjected to

immunoprecipitation using anti-FLAG M2 magnetic beads (M8823-1ML; Sigma-Aldrich).

Precipitated immunocomplexes were washed three times with lysis buffer and boiled in

sample buffer (46.7 mM Tris-HCl, pH 6.8, 5% glycerol, 1.67% sodium dodecyl sulfate,

1.55% dithiothreitol, and 0.02% bromophenol blue). For immunoprecipation of ATG9

vesicles, cells were disrupted by Dounce homogenization with hypotonic lysis buffer (10

mM HEPES, pH 7.9, 1.5 mM $MgCl_2$ and 10 mM KCl, EDTA-free protease inhibitor cocktail

[19543200; Roche]). Disruption was carried out by applying 35 strokes while the cell

suspension was cooled on ice. The supernatants were subjected to immunoprecipitation

using anti-FLAG M2 magnetic beads (M8823-1ML; Sigma-Aldrich). For immunoblotting, the

samples were separated by SDS-PAGE and transferred to Immobilon-P polyvinylidene

difluoride membranes (Millipore, WBKLS0500) with the Trans-Blot Turbo Transfer System

(Bio-Rad). After incubation with the relevant antibody in 5% skim milk in 20 mM Tris-HCl,

150 mM NaCl, and 0.1% Tween 20 (02194841-CF; MP Biomedicals), the signals from

incubation with SuperSignal West Pico Chemiluminescent Substrate (Thermo Fisher

Scientific, 34579) were detected with the FUSION SOLO.7S.EDGE imaging system (Vilber-

Lourmat). Contrast and brightness adjustment and quantification were performed using the

image processing software Fiji (Schindelin *et al*, 2012).

Fluorescence recovery after photobleaching

In-cell fluorescence recovery after photobleaching (FRAP) analyses were performed with

an Olympus Fluoview FV3000 confocal microscope equipped with a 60× oil-immersion

objective lens (1.40 NA, Olympus). The chamber was maintained at 37°C and continuously

supplied with humidified 5% CO_2. Bleaching was performed using 80% laser power (488 or

561 nm laser), and images were captured every 5 s for 30 frames. Recovery curves and

fitting were analyzed using OriginPro 2022. Fluorescence intensity of the bleached spot, an

unbleached control spot, and the background was measured using the software package

Fiji. Background intensity was subtracted, and the intensity values of the region of interest

are reported relative to the pre-bleached images during image acquisition. Each data point

represents the mean and standard error of fluorescence intensities in more than three

unbleached (control) or bleached (experimental) spots.

Live-cell imaging

Living imaging was conducted with the Olympus SpinSR10 spinning-disk confocal super-

resolution microscope equipped with a Hamamatsu ORCA-Flash 4.0 camera, a UPLAPO

OHR 100 × (NA 1.50) lens, and the SORA disk in place. The microscope was operated with

Olympus cellSens Dimension 2.3 software. Cells were passaged onto a four-chamber

glass-bottom dish (Greiner Bio-One) more than 24 h before imaging. To induce mitophagy,

cells were incubated with 10 µM CCCP in the presence of 200 nM SF650-conjugated Halo

ligand (GoryoChemical, A308-02). Images were processed using the image processing

software package Fiji (Schindelin *et al*, 2012). Fluorescence intensity of indicated

fluorophores was measured by Fiji, and spline connected graphs were created using

OriginPro 2022 software.

Immunofluorescence imaging

Cells were fixed with 4% PFA in 0.1 M phosphate buffer (pH 7.3) for 15 min at room

temperature (RT) and washed with PBS. Fixed cells were incubated with 10 µg/mL digitonin

for 5 min at RT and washed with PBS. Then, cells were incubated with primary antibodies

diluted (1:1000) in blocking buffer (3% BSA, in PBS) for 1 h at RT, washed with PBS,

incubated with Alexa Fluor-conjugated secondary antibodies (1:1000) in blocking buffer for

60 min at RT, washed again, and mounted on coverslips. Colocalization was calculated by

Fiji plugins BIOP JACoP, with the threshold of the ATG9A channel fixed to 500. ATG9A-

positive areas out of GFP-positive areas were calculated to obtain the overlapping areas.

Mathematical model of autophagy adaptor condensate formation

455   We formulated a mathematical model of autophagy adaptor condensate formation. For the

456   case of two mitochondria approaching each other (Fig. 3A), the free energy, F, of the

457   system can be written as

$$F = \int (f_{int} + f_{ent})dV, \tag{1}$$

with the interaction energy, f_{int}, and the entropic energy, f_{ent}, expressed as

$$f_{int} = \frac{\sigma}{2}(\nabla\phi_{ad})^2 - \chi_{self}\phi_{ad}^2 - \chi_{Ub}\phi_{Ub}\phi_{ad} + \chi_{Mt}\phi_{Mt}\phi_{ad}, \tag{2}$$

$$f_{ent} = \phi_{ad}\ln\phi_{ad} + \phi_{sol}\ln\phi_{sol}, \tag{3}$$

where ϕ_{ad}, ϕ_{Ub}, and $\phi_{sol} = 1 - \phi_{ad} - \phi_{Ub} - \phi_{Mt}$ are the adaptor, ubiquitin, and solvent concentrations, respectively, χ_{self} and χ_{Ub} represent the strength of the adaptor self-interaction and adaptor–ubiquitin interaction, respectively, σ represents the surface tension of the condensates, and χ_{Mt} represents the mitochondrial area exclusion effect, introduced to prevent proteins from entering the mitochondrial interior (ϕ_{Mt}).

For the case in which an isolation membrane surrounded a mitochondrion (Fig. 3C), the interaction energy, f_{int}, is instead expressed as

$$f_{int} = \frac{\sigma}{2}(\nabla\phi_{ad})^2 - \chi_{self}\phi_{ad}^2 - \chi_{Ub}\phi_{Ub}\phi_{ad} - \chi_{LC3}\phi_{LC3}\phi_{ad} + \chi_{Mt}\phi_{Mt}\phi_{ad} + \chi_{IM}\phi_{IM}\phi_{ad}, \tag{5}$$

where ϕ_{LC3} and χ_{LC3} represent the LC3 concentrations and the strength of the adaptor–LC3 interaction, respectively, and χ_{IM} represents the mitochondrial area exclusion effect, introduced to prevent proteins from entering the lumen of the isolation membrane (ϕ_{IM}). The solvent concentration that appears in the entropic energy (3) was also modified to $\phi_{sol} = 1 - \phi_{ad} - \phi_{Ub} - \phi_{LC3} - \phi_{Mt} - \phi_{IM}$.

For the numerical simulation, the space was discretized by a lattice with 10 nm on each side, and a concentration field was assigned to each location. The time evolution of the system was assumed to follow the Cahn–Hilliard equation,

$$\frac{\partial\phi_{ad}}{\partial t} = \frac{\partial}{\partial\vec{x}}\left[M\frac{\partial}{\partial\vec{x}}\left(\frac{\delta F}{\delta\phi_{ad}}\right)\right], \tag{4}$$

which is commonly used in analyses of phase separation dynamics (Berry *et al*, 2018). Here, $M = D/k_BT$ is the mobility of the protein with the diffusion constant $D = 10$ μm^2/sec (Milo & Phillips, 2015) expressed in terms of the Boltzmann constant k_B and temperature T. We considered a 3000 nm × 3000 nm square region with periodic boundary conditions containing two mitochondria and surrounding cytosolic components as a system and assumed that adaptor proteins are uniformly distributed in the cytosol around mitochondria

485   at a concentration of $\phi = 0.1$ in the initial state.

Statistical analysis

Statistical analysis was performed using OriginPro 2022 software. The statistical methods

used for each analysis are specified in the figure legends.

Acknowledgments

We thank Michael Lazarou (Walter and Eliza Hall Institute of Medical Research) for

providing pentaKO HeLa cells, Toshio Kitamura for pMXs-IP (The University of Tokyo),

Shoji Yamaoka (Tokyo Medical and Dental University, Tokyo, Japan) for pMRXIP, and

Teruhito Yasui (National Institutes of Biomedical Innovation, Health and Nutrition

(NIBIOHN), Osaka, Japan) for pCG-VSV-G and pCG-gag-pol. pKanCMV-mClover3-

mRuby3 was a gift from Michael Lin (Addgene plasmid # 74252 ;

http://n2t.net/addgene:74252 ; RRID:Addgene_74252).

This work was supported by the Exploratory Research for Advanced Technology (ERATO)

research funding program of the Japan Science and Technology Agency (JST)

(JPMJER1702 to NM), a Grant-in-Aid for Specially Promoted Research (22H04919 to NM)

and a Grant-in-Aid for Scientific Research (C) (23K05715 to Y.S.) from the Japan Society for

the Promotion of Science (JSPS), and German Research Foundation (Project 460056461 to

RLK). The numerical computations have been performed with the RIKEN supercomputer

system (HOKUSAI).

Author contributions

Z.Y., S.Y., H.C. R.L.K., and N.M. designed the project. Z.Y. performed most of the cell

biology experiments. Y.S. developed the mathematical model. Z.Y., S.Y., Y.S., R.L.K., and

N.M. wrote the manuscript. All authors analyzed and discussed the results and commented

on the manuscript.

Disclosure and competing interests statement

The authors declare that they have no conflict of interest.

References

References

Agudo-Canalejo J, Schultz SW, Chino H, Migliano SM, Saito C, Koyama-Honda I,
Stenmark H, Brech A, May AI, Mizushima N, *et al* (2021) Wetting regulates autophagy
of phase-separated compartments and the cytosol. *Nature* 591: 142–146

Bajar BT, Wang ES, Lam AJ, Kim BB, Jacobs CL, Howe ES, Davidson MW, Lin MZ & Chu J
(2016) Improving brightness and photostability of green and red fluorescent proteins for
live cell imaging and FRET reporting. *Sci Rep* 6: 20889

Berry J, Brangwynne CP & Haataja M (2018) Physical principles of intracellular organization
via active and passive phase transitions. *Reports on Progress in Physics* 81: 046601

Eura Y (2003) Two Mitofusin Proteins, Mammalian Homologues of FZO, with Distinct
Functions Are Both Required for Mitochondrial Fusion. *J Biochem* 134: 333–344

Fujioka Y, Alam JM, Noshiro D, Mouri K, Ando T, Okada Y, May AI, Knorr RL, Suzuki K,
Ohsumi Y, *et al* (2020) Phase separation organizes the site of autophagosome
formation. *Nature* 578: 301–305

Gallagher ER & Holzbaur ELF (2023) The selective autophagy adaptor p62/SQSTM1 forms
phase condensates regulated by HSP27 that facilitate the clearance of damaged
lysosomes via lysophagy. *Cell Rep* 42: 112037

Ganley IG & Simonsen A (2022) Diversity of mitophagy pathways at a glance. *J Cell Sci* 135:
jcs259748

Geisler S, Holmström KM, Skujat D, Fiesel FC, Rothfuss OC, Kahle PJ & Springer W (2010)
PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat Cell*
Biol 12: 119–131

Gouveia B, Kim Y, Shaevitz JW, Petry S, Stone HA & Brangwynne CP (2022) Capillary forces
generated by biomolecular condensates. *Nature* 609: 255–264

Gubas A & Dikic I (2022) A guide to the regulation of selective autophagy receptors. *FEBS J*
289: 75–89

Heo J-M, Ordureau A, Paulo JA, Rinehart J & Harper JW (2015) The PINK1-PARKIN
Mitochondrial Ubiquitylation Pathway Drives a Program of OPTN/NDP52 Recruitment
and TBK1 Activation to Promote Mitophagy. *Mol Cell* 60: 7–20

Hirose T, Ninomiya K, Nakagawa S & Yamazaki T (2023) A guide to membraneless
organelles and their various roles in gene regulation. *Nat Rev Mol Cell Biol* 24: 288–
304

Hyman AA, Weber CA & Jülicher F (2014) Liquid-Liquid Phase Separation in Biology. *Annu*
Rev Cell Dev Biol 30: 39–58

Itakura E, Kishi-Itakura C, Koyama-Honda I & Mizushima N (2012) Structures containing
Atg9A and the ULK1 complex independently target depolarized mitochondria at initial
stages of Parkin-mediated mitophagy. *J Cell Sci* 125: 1488–1499

Kane LA, Lazarou M, Fogel AI, Li Y, Yamano K, Sarraf SA, Banerjee S & Youle RJ (2014)
PINK1 phosphorylates ubiquitin to activate Parkin E3 ubiquitin ligase activity. *Journal of*
Cell Biology 205: 143–153

Kazlauskaite A, Kondapalli C, Gourlay R, Campbell DG, Ritorto MS, Hofmann K, Alessi DR,

Knebel A, Trost M & Muqit MMK (2014) Parkin is activated by PINK1-dependent phosphorylation of ubiquitin at Ser65. *Biochemical Journal* 460: 127–141

Kitamura T, Koshino Y, Shibata F, Oki T, Nakajima H, Nosaka T & Kumagai H (2003) Retrovirus-mediated gene transfer and expression cloning: powerful tools in functional genomics. *Exp Hematol* 31: 1007–1014

Koyano F, Okatsu K, Kosako H, Tamura Y, Go E, Kimura M, Kimura Y, Tsuchiya H, Yoshihara H, Hirokawa T, *et al* (2014) Ubiquitin is phosphorylated by PINK1 to activate parkin. *Nature* 510: 162–166

Kusumaatmaja H, May AI, Feeney M, McKenna JF, Mizushima N, Frigerio L & Knorr RL (2021a) Wetting of phase-separated droplets on plant vacuole membranes leads to a competition between tonoplast budding and nanotube formation. *Proceedings of the National Academy of Sciences* 118: e2024109118

Kusumaatmaja H, May AI & Knorr RL (2021b) Intracellular wetting mediates contacts between liquid compartments and membrane-bound organelles. *J Cell Biol* 220: e202103175

Lamark T & Johansen T (2021) Mechanisms of Selective Autophagy. *Annu Rev Cell Dev Biol* 37: 143–169

Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, Sideris DP, Fogel AI & Youle RJ (2015) The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. *Nature* 524: 309–314

Ma X, Li P & Ge L (2023) Targeting of biomolecular condensates to the autophagy pathway. *Trends Cell Biol* 33: 505–516

McSwiggen DT, Mir M, Darzacq X & Tjian R (2019) Evaluating phase separation in live cells: diagnosis, caveats, and functional consequences. *Genes Dev* 33: 1619–1634

Milo R & Phillips R (2015) Cell Biology by the Numbers Garland Science

Milovanovic D & De Camilli P (2017) Synaptic Vesicle Clusters at Synapses: A Distinct Liquid Phase? *Neuron* 93: 995–1002

Mizushima N & Levine B (2020) Autophagy in Human Diseases. *New England Journal of Medicine* 383: 1564–1576

Moore AS & Holzbaur ELF (2016) Dynamic recruitment and activation of ALS-associated TBK1 with its target optineurin are required for efficient mitophagy. *Proceedings of the National Academy of Sciences* 113: E3349–E3358

Musacchio A (2022) On the role of phase separation in the biogenesis of membraneless compartments. *EMBO J* 41: e109952

Nakatogawa H (2020) Mechanisms governing autophagosome biogenesis. *Nat Rev Mol Cell Biol* 21: 439–458

Noda NN, Wang Z & Zhang H (2020) Liquid–liquid phase separation in autophagy. *Journal of Cell Biology* 219: e202004062

Novak I, Kirkin V, McEwan DG, Zhang J, Wild P, Rozenknop A, Rogov V, Löhr F, Popovic D, Occhipinti A, *et al* (2010) Nix is a selective autophagy receptor for mitochondrial clearance. *EMBO Rep* 11: 45–51

O'Loughlin T, Kruppa AJ, Ribeiro ALR, Edgar JR, Ghannam A, Smith AM & Buss F (2020)

OPTN recruitment to a Golgi-proximal compartment regulates immune signalling and
cytokine secretion. *J Cell Sci* 133: 1–15

Onishi M, Yamano K, Sato M, Matsuda N & Okamoto K (2021) Molecular mechanisms and
physiological functions of mitophagy. *EMBO J* 40: e104705

Park D, Wu Y, Wang X, Gowrishankar S, Baublis A & De Camilli P (2023) Synaptic vesicle
proteins and ATG9A self-organize in distinct vesicle phases within synapsin
condensates. *Nat Commun* 14: 455

Peng S, Chen X-H, Chen S, Zhang J, Wang C, Liu W, Zhang D, Su Y & Zhang X (2021)
Phase separation of Nur77 mediates celastrol-induced mitophagy by promoting the
liquidity of p62/SQSTM1 condensates. *Nat Commun* 12: 5989

Rothbauer U, Zolghadr K, Muyldermans S, Schepers A, Cardoso MC & Leonhardt H (2008)
A Versatile Nanotrap for Biochemical and Functional Studies with Fluorescent Fusion
Proteins. *Molecular & Cellular Proteomics* 7: 282–289

Saitoh T, Nakano H, Yamamoto N & Yamaoka S (2002) Lymphotoxin-beta receptor mediates
NEMO-independent NF-kappaB activation. *FEBS Lett* 532: 45–51

Sansevrino R, Hoffmann C & Milovanovic D (2023) Condensate biology of synaptic vesicle
clusters. *Trends Neurosci* 46: 293–306

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for
biological-image analysis. *Nat Methods* 9: 676–682

Scott DJ, Gunn NJ, Yong KJ, Wimmer VC, Veldhuis NA, Challis LM, Haidar M, Petrou S,
Bathgate RAD & Griffin MDW (2018) A Novel Ultra-Stable, Monomeric Green
Fluorescent Protein For Direct Volumetric Imaging of Whole Organs Using CLARITY.
Sci Rep 8: 667

Style RW, Jagota A, Hui C-Y & Dufresne ER (2017) Elastocapillarity: Surface Tension and
the Mechanics of Soft Solids. *Annu Rev Condens Matter Phys* 8: 99–118

Sun D, Wu R, Zheng J, Li P & Yu L (2018) Polyubiquitin chain-induced p62 phase separation
drives autophagic cargo segregation. *Cell Res* 28: 405–415

Taylor NO, Wei M-T, Stone HA & Brangwynne CP (2019) Quantifying Dynamics in Phase-
Separated Condensates Using Fluorescence Recovery after Photobleaching. *Biophys*
J 117: 1285–1300

Turco E, Savova A, Gere F, Ferrari L, Romanov J, Schuschnig M & Martens S (2021)
Reconstitution defines the roles of p62, NBR1 and TAX1BP1 in ubiquitin condensate
formation and autophagy initiation. *Nat Commun* 12: 5212

Vargas JNS, Hamasaki M, Kawabata T, Youle RJ & Yoshimori T (2023) The mechanisms and
roles of selective autophagy in mammals. *Nat Rev Mol Cell Biol* 24: 167–185

Wexler JS, Heard TM & Stone HA (2014) Capillary Bridges between Soft Substrates. *Phys*
Rev Lett 112: 066102

Wong YC & Holzbaur ELF (2014) Optineurin is an autophagy receptor for damaged
mitochondria in parkin-mediated mitophagy that is disrupted by an ALS-linked mutation.
Proceedings of the National Academy of Sciences 111: E4439–E4448

Yamamoto H, Fujioka Y, Suzuki SW, Noshiro D, Suzuki H, Kondo-Kakuta C, Kimura Y, Hirano

H, Ando T, Noda NN, *et al* (2016) The Intrinsically Disordered Protein Atg13 Mediates
Supramolecular Assembly of Autophagy Initiation Complexes. *Dev Cell* 38: 86–99

Yamamoto H, Kakuta S, Watanabe TM, Kitamura A, Sekito T, Kondo-Kakuta C, Ichikawa R,
Kinjo M & Ohsumi Y (2012) Atg9 vesicles are an important membrane source during
early steps of autophagosome formation. *Journal of Cell Biology* 198: 219–233

Yamano K, Kikuchi R, Kojima W, Hayashida R, Koyano F, Kawawaki J, Shoda T, Demizu Y,
Naito M, Tanaka K, *et al* (2020) Critical role of mitochondrial ubiquitination and the
OPTN-ATG9A axis in mitophagy. *J Cell Biol* 219: e201912144

Yamasaki A, Alam JM, Noshiro D, Hirata E, Fujioka Y, Suzuki K, Ohsumi Y & Noda NN (2020)
Liquidity Is a Critical Determinant for Selective Autophagy of Protein Condensates. *Mol
Cell* 77: 1163-1175.e9

Yim WW-Y, Yamamoto H & Mizushima N (2022) A pulse-chasable reporter processing assay
for mammalian autophagic flux with HaloTag. *Elife* 11: e78923

Zaffagnini G, Savova A, Danieli A, Romanov J, Tremel S, Ebner M, Peterbauer T, Sztacho M,
Trapannone R, Tarafder AK, *et al* (2018) p62 filaments capture and present
ubiquitinated cargos for autophagy. *EMBO J* 37: e98308

Zheng Q, Chen Y, Chen D, Zhao H, Feng Y, Meng Q, Zhao Y & Zhang H (2022) Calcium
transients on the ER surface trigger liquid-liquid phase separation of FIP200 to specify
autophagosome initiation sites. *Cell* 185: 4082-4098.e22

Figure legends

Figure 1. Autophagy adaptors show distinct distributions during Parkin-mediated

mitophagy.

(A, B, and E) Representative images (left) and spline graphs of the intensity profiles along

the indicated lines in the respective images (right) of wild-type HeLa cells (A and B) or

HeLa cells lacking all five autophagy adaptors (penta KO cells) (E) expressing one of the

GFP-tagged autophagy adaptors or ubiquitin, mRuby–Omp25, and HaloTag7–LC3B at 45

min after 20 μM CCCP treatment. Separate mitochondria without contacting other

mitochondria or isolation membranes (A; Mt) or mitochondria with an isolation membrane

(B and E; Mt–IM) are shown. The *y*-axis in each of the graphs indicates the fluorescence

intensity. Scale bars indicate 1 μm.

(C and F) The relative intensity of GFP-tagged adaptors or ubiquitin (a and b) in LC3B-

positive areas compared with LC3B-negative areas in panels B and E. GFP signals were

normalized to the mRuby–Omp25 signals (a′ and b′) using the formula Relative Intensity =

(a/b)/(a′/b′), where a and a′ represent the intensities in the LC3B-positive area, and b and b′

represent the intensities in the LC3B-negative area. Solid horizontal bars indicate the

means, and dots indicate the data from five structures. Differences were statistically

analyzed by one-way analysis of variance with Dunnett's post-hoc test.

(D) Immunostaining of endogenous OPTN or NDP52 in wild-type HeLa cells expressing

mRuby–Omp25 and HaloTag7–LC3B at 45 min after 20 μM CCCP treatment. Scale bars

indicate 4 μm and 1 μm (magnified images).

Figure 2. OPTN and NDP52 show a dynamic exchange between the mitochondrial

surface and cytosol.

(A–D) Representative images (A and C) and quantification (B and D) of GFP fluorescence

recovery after photobleaching (FRAP) on separate mitochondria (Mt) (A and B) or on

mitochondria with an isolation membrane (Mt–IM) (C and D) in HeLa cells lacking all five

autophagy adaptors (penta KO cells) expressing GFP-tagged adaptors or ubiquitin at 45

min after 20 µM CCCP treatment. The photobleached areas are indicated by dotted lines.

The magnified panels display time-lapse images of the photobleached areas. Scale bars

indicate 2 µm and 1 µm (magnified images). Data were collected from four structures and

were fitted to the equation $y = a*(1 − \exp(−b*x))$. The dark shaded areas represent the 95%

confidence intervals, and the light shaded areas represent the 95% prediction intervals.

(E) Penta KO cells expressing GFP–OPTN at 45 min after 20 µM CCCP and 20 µM

wortmannin treatment. Wortmannin was added to inhibit autophagosome formation so that

OPTN would not be sequestered in a closed compartment. Images of cells before (upper

panels) and 2 min after (lower panels) the addition of 10% 1,6-hexanediol are displayed.

Scale bars indicate 5 µm and 2 µm (magnified images).

Figure 3. Mathematical models of condensate formation and localization of

autophagy adaptors.

(A) Schematic diagram of two separated mitochondria.

(B) Sheet-like condensates of autophagy adaptors on mitochondria deform when the inter-

mitochondrial distance d_{Mt} was changed. The strength of adaptor self-interaction (χ_{self}),

adaptor–ubiquitin interaction (χ_{Ub}), mitochondrial area exclusion (χ_{Mt}) and surface tension

(σ) were set to $\chi_{self} = 3k_BT$, $\chi_{Ub} = 12k_BT$, $\chi_{Mt} = 5k_BT$, and $\sigma = k_BT$, respectively. The

color indicates the adaptor concentration (ϕ_{ad}). Arrowheads indicate autophagy adaptors

filling the cleft between the two mitochondria.

(C) Schematic diagram of a single mitochondrion with an isolation membrane.

(D) Sheet-like condensates of autophagy adaptors on mitochondria deformed when the

isolation membrane (arrowheads) appeared and the angle α surrounding the mitochondria

changed. The strength of adaptor self-interaction (χ_{self}), adaptor–ubiquitin interaction (χ_{Ub}),

adaptor–LC3 interaction (χ_{LC3}), mitochondrial area exclusion (χ_{Mt}), isolation membrane

area exclusion (χ_{IM}), and surface tension (σ) were set to $\chi_{self} = 3k_BT$, $\chi_{Ub} = \chi_{LC3} = 12k_BT$,

$\chi_{Mt} = \chi_{IM} = 5k_BT$, and $\sigma = k_BT$, respectively. The color indicates the adaptor concentration

(ϕ_{ad}).

Figure 4. OPTN condensates redistribute upon contact with membranes.

(A–D) Time-lapse images (A and C) and spline graphs (B and D) of two separate

mitochondria approaching and contacting each other. Events of the attachment of two

mitochondria with comparable sizes (A and B) and with differing sizes (C and D) were

observed. The line graphs represent the intensity profiles along the indicated lines in the

respective images shown (B and D). The arrowheads indicate redistributed OPTN signals

filling the cleft between two adjacent mitochondria. Scale bars indicate 1 μm. See also

Movie EV1 for the time-lapse video corresponding to (C).

(E and F) Time-lapse images (E) and a spline graph (F) of a mitochondrion with an isolation

membrane elongating on its surface. The arrowhead indicates the elongating isolation

membrane. The line graph represent the intensity profiles along the indicated lines in the

respective images shown. Scale bars indicate 1 μm. See also Movie EV2 for the time-lapse

video corresponding to (E).

Figure 5. Liquid-like property of OPTN condensates is required for mitophagy.

(A) HeLa cells lacking all five autophagy adaptors (penta KO cells) expressing both GPF–

OPTN and mRuby–Ub (top), both GFP–OPTNΔUBAN and mRuby–Ub (middle), or both

GFP–OPTNΔUBAN and anti-GFP nanobody–mRuby–Ub (bottom) at 45 min after 20 μM

CCCP treatment. Time-lapse images of GFP FRAP are shown. Photobleached areas are

circled by dotted lines. Scale bars indicate 4 μm and 2 μm (magnified images).

(B) Interaction between GFP–OPTN mutants and FLAG–Ub or FLAG–nanobody–Ub was

investigated by immunoprecipitation with an anti-FLAG antibody and immunoblotting with

an anti-GFP antibody.

(C) Quantification of GFP FRAP on separate mitochondria (Mt) in penta KO cells

expressing both GFP–OPTN and mRuby–Ub or both GFP–OPTNΔUBAN and anti-GFP

nanobody–mRuby–Ub at 45 min after 20 μM CCCP treatment. Data were collected from

four structures and were fitted to the equation $y = a*(1 − \exp(−b*x))$. The dark shading

represent the 95% confidence intervals, and the light shading represent the 95% prediction

intervals.

(D and E) Representative data (D) and quantification (E) of HaloTag (Halo) processing

assay using cells expressing the indicated OPTN and Ub constructs. Cells expressing the

mtHalo–SNAP mitophagy reporter were treated without (0 h) and with 1 µM oligomycin and

2 µM antimycin for 18 h. The amount of processed Halo (proc. Halo) indicates the relative

amount of mitochondria degraded by the lysosomes. Solid horizontal bars indicate the

means, and dots indicate the data from four independent experiments. Differences were

statistically analyzed by one-way analysis of variance with Dunnett's post-hoc test.

Figure 6. Liquid-like property of OPTN condensates is required for the recruitment of

ATG9 vesicles.

(A and B) Localization of ATG9 vesicles under normal (CCCP−) or mitophagy-inducing

conditions (CCCP+, 60 min) (A) and quantification (B) of ATG9A colocalization with

mitochondria. Endogenous ATG9 was immunostained in penta KO cells expressing both

GFP–OPTN and mRuby–Ub or both GFP–OPTNΔUBAN and mRuby–nanobody–Ub

together with mitochondrially targeted Halo–SNAP (mtHalo–SNAP). Scale bars indicate 4

µm and 2 µm (magnified images). Solid horizontal bars indicate the means, each dot

indicates the mean value from one field of view with ≥10 cells (n=15). Differences were

statistically analyzed by one-way analysis of variance with Dunnett's post-hoc test.

(C and D) Representative data (C) and quantification (D) of organelle immunoprecipitation

in cells expressing GFP–OPTN or GFP–OPTNΔUBAN in the presence or absence of

FLAG-ATG9A. ATG9 vesicles were immunoprecipitated by FLAG M2 beads and GFP was

blotted to detect the interaction between GFP–OPTN (WT or ΔUBAN) and ATG9 vesicles at

60 min after 20 µM CCCP treatment. Solid horizontal bars indicate the means, and dots

indicate the data from five independent experiments. Differences were statistically analyzed

by one-way analysis of variance with Dunnett's post-hoc test.

(E) The models of ATG9 vesicle recruitment (left) and isolation membrane elongation (right)

mediated by liquid-like OPTN condensates. Partial or complete wetting of phase-separated

OPTN enables the recruitment of ATG9 vesicles to ubiquitinated mitochondria (left). Wetting

of OPTN on mitochondria and isolation membranes also facilitates the engulfment of

mitochondria by the isolation membranes (right).

Figure EV1. Localization of autophagy adaptors between mitochondria during

Parkin-mediated mitophagy.

(A and C) Representative images (left) and spline graphs of the intensity profiles along the

indicated lines (right) of wild-type HeLa cells (A) or HeLa cells lacking all five autophagy

adaptors (penta KO cells) (C) expressing one of the GFP–tagged autophagy adaptors or

ubiquitin and mRuby–Omp25 at 45 min after 20 µM CCCP treatment. Mitochondrial

clusters (Mt–Mt) are shown. The *y*-axis in each of the graphs indicates the fluorescence

intensity.

(B and D) The relative intensity of each adaptor was calculated as in Figure 1C. Solid

horizontal bars indicate the means, and dots indicate the data from five structures.

Differences were statistically analyzed by one-way ANOVA with Dunnett's post-hoc test.

Movie EV1. OPTN condensates redistribute upon mitochondrial contact.

Time-lapse video of two separate mitochondria approaching and contacting each other (30s

per frame). Scale bars indicate 1 µm. See also Figure 4C and D.

Movie EV2. OPTN condensates redistribute upon mitochondria–isolation membrane

contact.

Time-lapse video of a mitochondrion with an isolation membrane elongating on its surface

(30s per frame). Scale bars indicate 1 µm. See also Figure 4E and F.

Figure 1

Figure 1. Autophagy adaptors show distinct distributions during Parkin-mediated mitophagy.
(A, B, and E) Representative images (left) and spline graphs of the intensity profiles along the indicated lines in the respective images (right) of wild-type HeLa cells (A and B) or penta KO cells (E) expressing one of the GFP-tagged autophagy adaptors or ubiquitin, mRuby–Omp25 and HaloTag7–LC3B at 45 min after 20 µM CCCP treatment. Single mitochondria without contacting other mitochondria or isolation membranes (A; Mt) or mitochondria with an isolation membrane (B and E; Mt–IM) are shown. The Y axis in the graphs indicates the fluorescence intensity. Scale bars, 1 µm.
(C and F) The relative intensity of GFP-tagged adaptors or ubiquitin (a and b) in LC3B-positive areas compared to LC3B-negative areas in panels B and E. GFP signals were normalized to the mRuby–Omp25 signals (a' and b') using the indicated formula: Relative Intensity = (a/b)/(a'/b'), where a and a' represent the intensities in the LC3B-positive area, and b and b' represent the intensities in the LC3B-negative area. Solid bars indicate the means and dots indicate the data from 5 structures. Differences were statistically analyzed by one-way ANOVA with Dunnett's post hoc test.
(D) Immunostaining of endogenous OPTN or NDP52 in wild-type HeLa cells expressing mRuby–Omp25 and HaloTag7–LC3B at 45 min after 20 uM CCCP treatment. Scale bars, 4 um and 1 um (magnified images).

Figure 2

Figure 2. OPTN and NDP52 show a dynamic exchange between the mitochondrial surface and cytosol.
(A–D) Representative images (A and C) and quantification (B and D) of GFP FRAP on single mitochondria (Mt) (A and B) or on mitochondria with an isolation membrane (Mt–IM) (C and D) in penta KO cells expressing GFP-tagged adaptors or ubiquitin at 45 min after 20 μM CCCP treatment. The photobleached areas are indicated by dotted lines. The magnified panels display time-lapse images of the photobleached areas. Scale bars, 2 μm and 1 μm (magnified images). Data were collected from 4 structures and were fitted to the equation y = a*(1 - exp(-b*x)). The dark shadows represent the 95% confidence intervals, and the light shadows represent the 95% prediction intervals.
(E) Penta KO cells expressing GFP–OPTN at 45 min after 20 μM CCCP and 20 μM wortmannin treatment. Wortmannin was added to inhibit autophagosome formation so that OPTN is not sequestered in a closed compartment. Images of cells before (upper panels) and 2 min after (lower panels) the addition of 10% 1,6-hexanediol are displayed. Scale bars, 5 μm and 2 μm (magnified images).

Figure 3

Figure 3. Mathematical models of condensate formation and localization of autophagy adaptors.
(A) Schematic diagram of two separated mitochondria.
(B) Sheet-like condensates of autophagy adaptors on mitochondria deform when the inter-mitochondrial distance d_{Mt} was changed. The strength of adaptor self-interaction (χ_{self}), adaptor-ubiquitin interaction (χ_{Ub}), mitochondrial area exclusion (χ_{Mt}) and surface tension (σ) were set to $\chi_{self}=3k_BT$, $\chi_{Ub}=12k_BT$, $\chi_{Mt}=5k_BT$, and $\sigma=k_BT$ respectively. The color indicated the adaptor concentration (ϕ_{ad}). Arrowheads indicate autophagy adaptors filling the cleft between the two mitochondria.
(C) Schematic diagram of a mitochondrion with an isolation membrane.
(D) Sheet-like condensates of autophagy adaptors on mitochondria deformed when the isolation membrane (arrowheads) appeared and the angle α surrounding the mitochondria changed. The strength of adaptor self-interaction (χ_{self}), adaptor-ubiquitin interaction (χ_{Ub}), adaptor-LC3 interaction (χ_{LC3}), mitochondrial area exclusion (χ_{Mt}), isolation membrane area exclusion (χ_{IM}), and surface tension (σ) were set to $\chi_{Ub}=\chi_{LC3}=12k_BT$ and $\chi_{self}=3k_BT$. $\sigma=k_BT$ and $\chi_{Mt}=\chi_{IM}=5k_BT$. The color indicated the adaptor concentration (ϕ_{ad}).

Figure 4

Figure 4. OPTN condensates redistribute upon contact with membranes.

(A–D) Time-lapse images (A and C) and spline graphs (B and D) of two separate mitochondria approaching and contacting each other. Events of the attachment of two mitochondria with comparable sizes (A and B) and with differing sizes (C and D) were observed. The line graphs represent the intensity profiles along the indicated lines in the respective images shown (B and D). The arrowheads indicate redistributed OPTN signals filling the cleft between two adjacent mitochondria. Scale bars indicate 1 μm. See also Movie EV1 for the time-lapse video corresponding to (C).

(E and F) Time-lapse images (E) and a spline graph (F) of a mitochondrion with an isolation membrane elongating on its surface. The arrowhead indicates the elongating isolation membrane. The line graph represent the intensity profiles along the indicated lines in the respective images shown. Scale bars indicate 1 μm. See also Movie EV2 for the time-lapse video corresponding to (E).

Figure 5

Figure 5. Liquid-like property of OPTN condensates is required for mitophagy.

(A) HeLa cells lacking all five autophagy adaptors (penta KO cells) expressing both GPF–OPTN and mRuby–Ub (top), both GFP–OPTNΔUBAN and mRuby–Ub (middle), or both GFP–OPTNΔUBAN and anti-GFP nanobody–mRuby–Ub (bottom) at 45 min after 20 µM CCCP treatment. Time-lapse images of GFP FRAP are shown. Photobleached areas are circled by dotted lines. Scale bars indicate 4 µm and 2 µm (magnified images).

(B) Interaction between GFP–OPTN mutants and FLAG–Ub or FLAG–nanobody–Ub was investigated by immunoprecipitation with an anti-FLAG antibody and immunoblotting with an anti-GFP antibody.

(C) Quantification of GFP FRAP on separate mitochondria (Mt) in penta KO cells expressing both GFP–OPTN and mRuby–Ub or both GFP–OPTNΔUBAN and anti-GFP nanobody–mRuby–Ub at 45 min after 20 µM CCCP treatment. Data were collected from four structures and were fitted to the equation $y = a*(1 − \exp(−b*x))$. The dark shading represent the 95% confidence intervals, and the light shading represent the 95% prediction intervals.

(D and E) Representative data (D) and quantification (E) of HaloTag (Halo) processing assay using cells expressing the indicated OPTN and Ub constructs. Cells expressing the mtHalo–SNAP mitophagy reporter were treated without (0 h) and with 1 µM oligomycin and 2 µM antimycin for 18 h. The amount of processed Halo (proc. Halo) indicates the relative amount of mitochondria degraded by the lysosomes. Solid horizontal bars indicate the means, and dots indicate the data from four independent experiments. Differences were statistically analyzed by one-way analysis of variance with Dunnett's post-hoc test.

Figure 6

Figure 6. Liquid-like property of OPTN condensates is required for the recruitment of ATG9A vesicles.
(A and B) Localization of ATG9 vesicles under normal (CCCP−) or mitophagy-inducing conditions (CCCP+, 60 min) (A) and quantification (B) of ATG9A colocalization with mitochondria. Endogenous ATG9 was immunostained in penta KO cells expressing both GFP–OPTN and mRuby–Ub or both GFP–OPTNΔUBAN and mRuby–nanobody–Ub together with mitochondrially targeted Halo–SNAP (mtHalo–SNAP). Scale bars indicate 4 μm and 2 μm (magnified images). Solid horizontal bars indicate the means, each dot indicates the mean value from one field of view with ≥10 cells (n=15). Differences were statistically analyzed by one-way analysis of variance with Dunnett's post-hoc test.
(C and D) Representative data (C) and quantification (D) of organelle immunoprecipitation in cells expressing GFP–OPTN or GFP–OPTNΔUBAN in the presence or absence of FLAG-ATG9A. ATG9 vesicles were immunoprecipitated by FLAG M2 beads and GFP was blotted to detect the interaction between GFP–OPTN (WT or ΔUBAN) and ATG9 vesicles at 60 min after 20 μM CCCP treatment. Solid horizontal bars indicate the means, and dots indicate the data from five independent experiments. Differences were statistically analyzed by one-way analysis of variance with Dunnett's post-hoc test.
(E) The models of ATG9 vesicle recruitment (left) and isolation membrane elongation (right) mediated by liquid-like OPTN condensates. Partial or complete wetting of phase-separated OPTN enables the recruitment of ATG9 vesicles to ubiquitinated mitochondria (left). Wetting of OPTN on mitochondria and isolation membranes also facilitates the engulfment of mitochondria by the isolation membranes (right).

Figure EV1

Figure EV1. Localization of autophagy adaptors between mitochondria during Parkin-mediated mitophagy.

(A and C) Representative images (left) and spline graphs of the intensity profiles along the indicated lines (right) of wild-type HeLa cells (A) or penta KO cells (C) expressing one of the GFP–tagged autophagy adaptors or ubiquitin and mRuby–Omp25 at 45 min after 20 µM CCCP treatment. The Y axis in the graphs indicates the fluorescence intensity.

(B and D) The relative intensity of each adaptor was calculated as in Figure 1C. Solid bars indicate the means and dots indicate the data from 5 structures. Differences were statistically analyzed by one-way ANOVA with Dunnett's post hoc test.