Suppression of progesterone by influenza A virus causes adverse maternal and fetal

outcomes in mice

Patrick S. Creisher[1], Maclaine A. Parish[1], Jun Lei [2], Jin Liu[3], Jamie L. Perry[1], Ariana D.

Campbell[1], Morgan L. Sherer[1], Irina Burd[2], and Sabra L. Klein[1,†]

[1]W. Harry Feinstone Department of Molecular Microbiology and Immunology, The Johns

Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

[2]Department of Obstetrics, Gynecology & Reproductive Sciences, University of Maryland School

of Medicine, Baltimore, Maryland, USA

[3]Integrated Research Center for Fetal Medicine, Department of Gynecology and Obstetrics,

Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

[†]To whom correspondence should be addressed: Sabra L. Klein (sklein2@jhu.edu)

Short title: influenza and pregnancy

Key words: cytokine storm, macrophages, placenta, pregnancy, prostaglandins, progestins

Manuscript statistics:

Abstract: 138
Text: 6563
Figures: 7
References: 89
20

Abstract:

Influenza A virus infection during pregnancy can cause adverse maternal and fetal

outcomes, but the mechanism responsible remains elusive. Infection of outbred mice with 2009

H1N1 at embryonic day (E) 10 resulted in significant maternal morbidity, placental tissue

damage and inflammation, fetal growth restriction, and developmental delays that lasted

through weaning. Restriction of pulmonary virus replication was not inhibited during pregnancy,

but infected dams had suppressed circulating and placental progesterone (P4) concentrations

that were caused by H1N1-induced upregulation of pulmonary cyclooxygenase (COX)-1, but not

COX-2-, dependent synthesis and secretion of prostaglandin (PG) F2α. Treatment with 17-α-

hydroxyprogesterone caproate (17-OHPC), a synthetic progestin that is safe to use in

pregnancy, ameliorated the adverse maternal and fetal outcomes from H1N1 infection and

prevented placental cell death and inflammation. These findings highlight the therapeutic

potential of progestin treatments for influenza during pregnancy.

1 Introduction

Pregnant individuals are at increased risk of severe disease and death following infection

with seasonal and pandemic influenza viruses (Cox et al., 2006; Klein et al., 2010; Mertz et al.,

2017; Vousden et al., 2021; Woolston and Conley, 1918). For example, pregnancy was

associated with an 18.1% increase in one hospital's fatality rate during the 1918 H1N1

pandemic and a disproportionally greater percentage of morbidity and death during the 2009

H1N1 pandemic (Klein et al., 2010; Woolston and Conley, 1918). Further, seasonal influenza

viruses are reported to annually infect between 483 and 1097 per 10,000 pregnant women (Katz

et al., 2017), with greater odds of adverse outcomes, including intensive care unit (ICU)

admission, acute cardiopulmonary morbidity, and increased hospital length of stay, among

pregnant than nonpregnant people (Cox et al., 2006; Neuzil et al., 1998; Vousden et al., 2021).

In the absence of vertical transmission, influenza virus infection during pregnancy is still

associated with adverse perinatal and fetal outcomes (Hartel et al., 2016; He et al., 2017; Omer

et al., 2011), as was seen during the 2009 H1N1 pandemic, where maternal influenza A virus

(IAV) infection increased the relative risk of low birth weights and stillbirths by 71% and 136%,

respectively (He et al., 2017). Moreover, infection with influenza viruses during pregnancy has

been associated with long-lasting consequences for offspring, such as increasing the risk of

development of neuropsychiatric and metabolic disorders in adulthood (Brown et al., 2004;

Parboosing et al., 2013; Song et al., 2020).

The 2009 H1N1 pandemic spurred development of mouse models of IAV infection during

pregnancy to uncover mechanisms of adverse maternal and fetal outcomes. Collectively, these

studies have revealed pregnancy-specific mechanisms of influenza pathogenesis. Notably,

pregnant mice infected with diverse influenza viruses consistently experience greater morbidity

and/or mortality than nonpregnant female mice, consistent with observations in humans (Chan

et al., 2010; Engels et al., 2017; Kim et al., 2012; Littauer et al., 2017; Marcelin et al., 2011;

Pazos et al., 2012). Pregnancy impacts pulmonary innate immune activity and tissue repair

following IAV infection, as pregnant dams have greater proinflammatory cytokine concentrations

and numbers of neutrophils and macrophages in lungs than nonpregnant females (Chan et al.,

2010; Engels et al., 2017; Lauzon-Joset et al., 2019; Marcelin et al., 2011; Pazos et al., 2012).

Moreover, IAV infection during pregnancy is associated with pulmonary tissue damage,

impaired epithelial regeneration, reduced progesterone (P4) concentrations, and a shift of

alveolar macrophages from a classical to an alternative activation state (Lauzon-Joset et al.,

2019; Marcelin et al., 2011; Swieboda et al., 2020; Vermillion et al., 2018). Pregnant dams

infected with IAV have reduced adaptive immune responses including reduced IAV-specific

antibody titers, numbers of circulating CD4$^+$ and CD8$^+$ T cells, and IAV-specific cytotoxic T cell

activity, with these effects associated with changes in circulating P4 and estrogens (Chan et al.,

2010; Engels et al., 2017; Peluso et al., 1980; Swieboda et al., 2020). Whether IAV-induced

changes in circulating concentrations of sex steroids are a cause or consequence of maternal

disease during infection has not been determined.

Although mice do not recapitulate all aspects of human pregnancy, mice do exhibit many of

the cardiovascular, hormonal, and immunological changes associated with pregnancy (Abu-

Raya et al., 2020; Ander et al., 2019; Arany et al., 2022; Krishnan et al., 2013; Malassine et al.,

2003). For example, concentrations of P4 increase dramatically during pregnancy and support

the growth of the placenta in both mice and humans (Di Renzo et al., 2016; Malassine et al.,

2003), with secretion exclusively by the corpus luteum in mice and secretion shifting from the

corpus luteum to the placenta mid-gestation in humans (Malassine et al., 2003; Terakawa et al.,

2012). Mouse models of pregnancy have limitations, including shorter gestation in mice and

distinct mechanisms of placentation (Arany et al., 2022; Krishnan et al., 2013; Malassine et al.,

2003). Furthermore, when using inbred mice, the absence of a semi-allogeneic fetus limits

tolerogenic immune mechanisms observed in humans and outbred animals (Moldenhauer et al.,

2010; Tafuri et al., 1995), which has been shown to contribute to IAV pathogenesis (Engels et

al., 2017; Jacobsen et al., 2021).

In the current study, using outbred mice, IAV infection during pregnancy caused significant

maternal morbidity and adverse fetal outcomes that were associated with transient placental

tissue damage and inflammation. Both maternal disease and adverse perinatal outcomes were

mediated by IAV-induced activation of cyclooxygenase-1 (COX-1), but not COX-2, causing

increased activation of prostaglandin F2α (PGF2α) and suppression of P4. Treatment of IAV-

infected pregnant female mice with a synthetic progestin that is safe for administration during

pregnancy reversed the adverse maternal, placental, and fetal outcomes from IAV infection.

This novel study highlights utilizing hormone modulating treatments for the treatment of

influenza during pregnancy.

2 Materials and Methods

2.1 Viruses and cells

Viral seed stocks of mouse adapted A/California/04/09 H1N1 (maH1N1), generated by

reverse genetics from a published sequence (Ye et al., 2010) were kindly provided by Dr. Andy

Pekosz (Johns Hopkins University). Working stocks were generated by infecting Madin-Darby

canine kidney (MDCK) cells at a multiplicity of infection (MOI) of 0.01 tissue culture infectious

dose 50 ($TCID_{50}$) per cell in infection media [DMEM (Sigma D5796) supplemented with 0.5%

filter-sterilized bovine serum albumin (BSA; Sigma A9647), 100 U/mL penicillin and 100 µg/mL

streptomycin (Gibco 15149–122), 2 mM l-glutamine (Gibco 25030–081), and 5 µg/mL of N-

acetyl trypsin (Sigma T6763)]. Approximately 72 h post infection, the supernatant fluids were

collected, clarified by centrifugation (500 g for 10 minutes), and stored in aliquots at –70°C.

2.2 Experimental mice

Timed-pregnant adult (8–12 weeks of age) CD-1 IGS mice were purchased from Charles

River Laboratories. Animals arrived at embryonic day 8 (E8) and were singly housed under

standard biosafety level 2 housing conditions with ad libitum food and water. Pregnant mice

were acclimated for at least 24 h prior to infections (Creisher et al., 2022a). Monitoring and

procedures for animal experiments were performed consistently at the same time of day.

2.3 IAV infections and monitoring

Pregnant dams were intranasally infected at E10 with 10^3 TCID$_{50}$ units of maH1N1 in 30

µl of media (DMEM; Sigma D5796), a sublethal dose in nonpregnant CD-1 female mice, or

mock inoculated with 30 µl of media. Prior to intranasal infection, mice were anesthetized via

intraperitoneal ketamine/xylazine cocktail (80 mg/kg ketamine, 5 mg/kg xylazine). Following

intranasal infections, body mass and rectal temperature were monitored once daily in the

morning for 14 days or until tissue collection.

2.4 COX inhibitor treatments

Experimental animals were administered vehicle alone [20% DMSO (Sigma D2650) v/v

diluted in PBS], the COX-1 specific inhibitor SC-560 (10 mg/kg; abcam ab120649), or the COX-

2 specific inhibitor NS-398 (10 mg/kg; MedChemExpress HY-13913) in 100 µl intraperitoneally

(Futaki et al., 1993; Tanigawa et al., 2004; Thomas and Kuhn, 2005). Mice were administered

treatment twice daily, starting 1 day prior to infection and until tissue collection at 3 dpi. Serum

thromboxane B2 was measured to confirm COX-1 inhibition (Wallace et al., 1998; Wallace et

al., 1999) by ELISA per the manufacturer's instructions (Cayman Chemical 19030). Inhibitors

were reconstituted and diluted fresh for each dose.

2.5 17α-hydroxyprogesterone caproate treatment

17α-hydroxyprogesterone caproate (17-OHPC, 250 mg/mL) was obtained from the

Johns Hopkins Hospital Pharmacy (NDC 71225-105-01) and diluted to 20 mg/mL in vehicle

[46% v/v benzyl benzoate (Sigma B6630) in castor oil (Sigma C9606)]. Experimental animals

were administered vehicle alone or 17-OHPC subcutaneously at the time of intranasal infection.

17-OHPC was administered at a dose of 2 mg/dam, which previously demonstrated efficacy in

preventing adverse neurological outcomes in a mouse model of intrauterine inflammation

(Novak et al., 2018), and was administered as a single injection due to the prolonged half-life of

17-OHPC when delivered in an oil-based solution [>7 days (Shaik et al., 2016)].

2.6 Offspring measurements and developmental assessments

Offspring from IAV- and mock-inoculated dams were measured at post-natal day (PND)

0, within 12 hours of birth. Body mass (g), length measured from nose to anus (mm), and head

diameter measured from ear to ear (mm) were recorded for each pup, using a caliper, and the

average for each independent litter was calculated to avoid confounding litter effects. After birth

and until weaning, pups were monitored daily for body mass (g) and the appearance of key

developmental milestones including fur development, the opening of the ear canals and eyes,

and teeth eruption (Heyser, 2004; Hill et al., 2008). At PND5, pups were subjected to the

developmental neurobehavioral assay of surface righting, as described (Creisher et al., 2023;

Feather-Schussler and Ferguson, 2016). For each test, 1–2 male and 1–2 female offspring from

at least 5 independent litters were used per condition to avoid confounding litter effects. Pups

were subjected to 3 attempts, with the time to complete the test recorded on a stopwatch. The

upper limit of time was 60 seconds, and a pups' best trial for each test was used for analysis.

2.7 Tissue and serum collection

Experimental mice were euthanized via ketamine/xylazine overdose (160 mg/kg

ketamine, 10 mg/kg xylazine) followed by cardiac exsanguination at 1, 3, 6, or 8 days post-

infection. Maternal lungs, spleen, and ovaries were collected and flash frozen on dry ice for

homogenization. Fetuses and placenta were flash frozen in dry ice or fixed in 4%

paraformaldehyde (Thermo Fisher Scientific J19943.K2) for 72 h at 4°C for

immunohistochemistry. Serum was separated from blood by centrifugation at 10,000 rpm for

30 min at 4°C.

2.8 Tissue homogenization and infectious virus quantification

Frozen right cranial lungs, spleen, ovaries, placentas, and fetuses were weighed and

homogenized in 1:10 weight per volume of PBS using Lysing Matrix D tubes (MP Biomedicals

#6913100) in a MP Fast-prep 24 5G instrument. Homogenate was clarified by centrifugation at

12,000 rpm and stored at -70°C until analysis. IAV infectious virus titers in tissue homogenate

was determined by $TCID_{50}$ assay, as described previously (Hall et al., 2017; Vermillion et al.,

2018). Briefly, tissue homogenates were serially diluted in infection media and plated onto a

monolayer of 100% confluent MDCK cells in replicates of 6. Cells were incubated for 7 days at

32°C and 5% CO2, fixed for 1 hour with 4% formaldehyde, and then stained with naphthol blue

black for 4 hours. After staining, plates were scored for cytopathic effects and the Reed-Muench

method was used to calculate the $TCID_{50}$ titer for each sample.

2.9 Multiplex cytokine analysis

Cytokines and chemokines in the right cranial lung, spleen, and placenta homogenate

were multiplexed using the Immune Monitoring 48-Plex Mouse Procartaplex Panel (Thermo

Fisher Scientific EPX480-20834-901) miniaturized with the Curiox DA-Bead DropArray platform

(Versteeg et al., 2017). Briefly, 96-well DA-Bead Plates (Curiox 96-CC-BD-05) were blocked

with 10 µL per well of 1% BSA (Sigma A9647) in PBS for 30 min in a humidified magnet box

(Curiox HMBXWT01) at room temperature on a rotator. The plate was washed with Curiox wash

buffer [0.1% BSA (Sigma A9647) and 0.05% Tween-20 (Sigma P1379)] on a DropArray LT210

(Curiox). Procartaplex beads (5 µL) were added to each well, followed by 5 µL of serially diluted

ProCartaplex standards or 10 µL tissue homogenate per well, with each sample assayed in

duplicate. The plate was incubated overnight in a humidified magnet box on a rotator at 4°C.

The following day, the plate was washed three times, and 5 µL of ProCartaPlex detection

antibody added per well. The plate was incubated for 1 h in a humidified magnet box on a

rotator. After incubation, 10 µL streptavidin-phycoerythrin was added to each well and incubated

for 30 min in a humidified magnet box on a rotator. The plate was washed three times, and the

beads were reconstituted in Curiox wash buffer and transferred to a skirted v-bottom 96 well

plate (Biorad HSP9601) for reading on a MagPix system (Luminex). Analysis was performed

using BioPlex Manager (version 6.2). The following analytes were quantified: G-CSF, GM-CSF,

M-CSF, CXCL1, CXCL2, CXCL5, CXCL10, CCL2, CCL3, CCL4, CCL5, CCL7, CCL11, IFNα,

IFNγ, IL-1β, IL-1α, IL-2, IL-3, IL-2R, IL-4, IL-5, IL-6, IL-7, IL-7Rα, IL-9, IL-10, IL-12p70, IL-13, IL-

15/IL-15R, IL-17A IL-18, IL-19, IL-22, IL-23, IL-25, IL-27, IL-28, IL-31, IL-33, IL-33R, Leukemia

inhibitory factor (LIF), TNFα, receptor activator of nuclear factor kappa-B ligand (RANKL), B-cell

activating factor (BAFF), Betacellulin, vascular endothelial growth factor (VEGF-A), and Leptin.

2.10 Placental histology, immunohistochemistry, and TUNEL staining

Placentas were fixed overnight at 4°C in 4% PFA and were washed five times with PBS

and immersed in 30% sucrose until saturation. Using a Leica CM1950 cryostat, the specimens

were cut at 20-µm thickness and mounted on positively charged slides (Fisher Scientific12-550-

15). Routine H&E staining was performed to evaluate the morphological change of the

placentas. For immunohistochemical staining, slides were washed with PBS, followed by

permeabilization in PBS solution containing 0.05% Triton X-100 and 10% normal goat serum

(ThermoFisher Scientific 50197Z) for 30 min. Placentas were incubated with rabbit anti-vimentin

(1:200, Abcam ab92547) or rabbit anti-cytokeratin (1:200, Dako Z0622) overnight at 4°C. The

next day, sections were rinsed with PBS and incubated with donkey anti-rabbit (ThermoFisher

Scientific R37119) fluorescent secondary antibodies diluted 1:500 for 3 h at room temperature.

DAPI (Roche 10236276001) was applied for counterstaining, followed by mounting with

Fluoromount-G (eBioscience 00-4958-02). Images were taken using a Zeiss Axioplan 2

microscope (Jena) under 5x or 20x magnification. Cell density of vimentin and cytokeratin

positive cell quantification was performed using Image J (1.47v). The 20x images were captured

from the same batch of experiments, utilizing identical imaging parameters, including exposure

time for quantification. After setting the appropriate scale and threshold for positive expression,

the percentage of positive expression relative to the entire area was calculated. For each

placenta, six random images in the labyrinth at the middle level (thickest) of placenta were

taken, and the average fluorescent area calculated for that placenta. One placenta per dam was

used and 5-10 dams per group were analyzed. Terminal deoxynucleotidyl transferase dUTP

nick end labeling (TUNEL) was performed with Click-iT™ Plus TUNEL Assay (ThermoFisher

C10617) per the manufacturer protocol. Placental slides were fixed with 4% paraformaldehyde

in PBS for 15 min at 37°C, rinsed with PBS, and permeabilized with proteinase K solution for 15

min at RT followed by staining with TUNEL working solution for 60 min at 37°C. Specimens

were mounted on glass slides using Fluoromount-G (eBioscience #00-4958-02). For each

placenta, 10 random images in labyrinth at the middle level (thickest) of placenta were taken,

and the average percentage TUNEL$^+$ (TUNEL$^+$ fluorescent signal divided by the DAPI

fluorescent signal) given for that placenta. One placenta per dam was used and 5-9 dams per

group were analyzed.

2.11 Flow cytometry

Placentas and maternal blood were collected from a subset of experimental dams at 6

dpi for flow cytometry as described previously (Liu et al., 2021; Liu et al., 2022). Single cells

were prepared from 3 pooled placentas per dam. Manual digestion was performed by mincing

placentas into 2-4 mm pieces using a scalpel blade on ice, followed by enzymatic digestion with

collagenase D (Roche 11088882001) in 6 mL of Hanks' Balanced Salt Solution with calcium

and magnesium (HBSS+; Gibco 14025-092) at 37℃. Collagenase D digestion was stopped

after 30 min with 6 mL of Hanks' Balanced Salt Solution without calcium and magnesium

(HBSS-; Gibco 14175-095), and cells were passed through a 70 μm nylon mesh cell strainer

and washed with HBSS- thrice. Red blood cell lysis with ACK lysing buffer (Quality Biological

118-156-101), Fc blocking [anti-CD16/32 (BD Biosciences 554142)], and total cell counts were

performed prior to resuspension in cell staining buffer [0.5% BSA (Sigma A9647) + 0.05%

sodium azide (Sigma S8032) + 1mM EDTA (Quality Biological 351-027-101) in PBS] for

staining. Maternal blood (1 mL) was added to 10 ml of 5 mM EDTA (Quality Biological 351-027-

101) in HBSS+ (Gibco 14175-095), followed by centrifugation at 300g for 5 min at 4℃, red blood

cell lysis with ACK lysing buffer, Fc blocking, total cell counts, and resuspension in cell staining

buffer for staining. Single cell suspensions were split in half and stained with one of two surface

marker antibody cocktails at 1:100 for 30 min on ice in the dark. Panel one contained anti-

mouse CD45 (clone 30-F11) Super Bright 600 (Thermo Fisher Scientific 63-0451-82), anti-

mouse CD11b (clone M1-70) PerCP-Cyanine5.5 (Thermo Fisher Scientific 45-0112-82), anti-

mouse F4/80 (clone BM8) eFluor 450 (Thermo Fisher Scientific 48-4801-82), anti-mouse

CD170 (clone 1RNM44N) PE-Cyanine7 (Thermo Fisher Scientific 25-1702-82), anti-mouse

CD163 (clone TNKUPJ) PE (Thermo Fisher Scientific 12-1631-82), and anti-mouse CD68

(clone FA-11) Alexa Fluor 488 (Thermo Fisher Scientific 53-0681-82). Panel two contained anti-

mouse CD45 (clone 30-F11) PE (Thermo Fisher Scientific 12-0451-82), anti-mouse NK1.1

(clone PK136) PE-Cyanine7 (Thermo Fisher Scientific 25-5941-82), anti-mouse CD3 (clone

17A2) eFluor 450 (Thermo Fisher Scientific 48-0032-82), anti-mouse CD8a (clone 53-6.7)

eFluor 506 (Thermo Fisher Scientific 69-0081-82), and anti-mouse CD19 (clone eBio1D3)

Super Bright 600 (Thermo Fisher Scientific 63-0193-82). After incubation, cells were washed

and acquired on acquired on an Attune™ NxT Acoustic Focusing Cytometer (ThermoFisher

Scientific) and analyzed with FlowJo (version 10.8.1). Compensation was performed with

OneComp eBeads (Thermo Fisher Scientific 01-1111-41). Gating was performed as described

previously [(Liu et al., 2021; Liu et al., 2022) and **Supplemental Fig. 3**]. Debris and doublets

were excluded by gating on side scatter height versus side scatter area to delaminate singlets,

and specific populations including leukocytes, ($CD45^+$), NK cells ($CD45^+$, $CD3^-$, $CD19^+$), B cells

($CD45^+$, $CD3^-$, $CD19^+$), $CD4^+$ T cells ($CD45^+$, $CD3^+$, $CD4^+$), $CD8^+$ T cells ($CD45^+$, $CD3^+$, $CD8a^+$),

neutrophils ($CD45^+$, $CD11b^+$, $F4/80^-$, $CD170^-$, side scatter area high), eosinophils ($CD45^+$,

CD11b$^+$, F4/80$^+$, CD170$^+$), CD163$^+$ macrophages (CD45$^+$, CD11b$^+$, F4/80$^+$, CD163$^+$), and

CD68$^+$ macrophages (CD45$^+$, CD11b$^+$, F4/80$^+$, CD68$^+$) were gated based on unstained and

fluorescent minus one (FMO) controls.

2.12 Progesterone and prostaglandin ELISA

To measure P4, steroids were extracted from 50 µL of serum or tissue homogenate via

diethyl ether extraction at a 1:5 sample to ether ratio. Extracted steroids were diluted 1:100

(serum) or 1:10 (tissue homogenate) and P4 was quantified by ELISA according to the

manufacturer's instructions (Enzo Life Sciences ADI-900-11). PGF2α (Alpco 74-PGFHU-E01)

and PGE2 (Alpco 74-PG2HU-E01) in serum or tissue homogenate was quantified by ELISA

according to the manufacturer's instructions.

2.13 Western Blot

Flash frozen right caudal lung lobes and right ovaries were homogenized in 1X Cell lysis

Buffer (Cell Signaling Technology 9803) with 1X Protease Inhibitor cocktail (Sigma-Aldrich

P8340) and sodium fluoride (Fisher Scientific S299; 20µL lysis buffer per mg tissue). Protein

lysates were stored at -80°C until analysis. Protein concentration of each lysate was measured

using the Pierce BCA Protein Assay Kit. For each sample, 20 µg of protein was subjected to

sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) on NuPAGE 4-12%

Bis-Tris gels. The gel was blotted onto Immobilon-FL PVDF Membrane (Millipore #IPFL00010)

and the membranes were blocked with a 1:1 mixture of 1XPBS/Tween-20 solution (Sigma-

Aldrich P3563) and Intercept blocking buffer (LI-COR Biosciences 927-70001) for 30 min at

room temperature. Membranes were treated with a primary antibody diluted in blocking solution

at 4°C overnight on a rocker. Membranes were then washed with PBS-Tween three times and

incubated in secondary antibody solutions for one hour at room temperature on a rocker.

Membranes were washed three times in PBS-Tween and then imaged on an Azure 600 imager

(Azure Biosystems). Individual bands were quantified using Image Studio software (LI-COR

Biosciences; version 3.1.4) The signal from each band was normalized against the GAPDH

signal and graphed as arbitrary units. Primary antibodies used were rabbit anti-COX-1 (abcam

ab109025), rabbit anti-COX-2 (abcam ab179800), and mouse anti-GAPDH (abcam ab82450).

Secondary antibodies included goat anti-mouse Alexa Fluor 488 (Thermo Fisher Scientific

A11001) and donkey anti-rabbit Alexa Fluor Plus 647 (Thermo Fisher Scientific A32795).

2.14 Statistical analyses

Body mass change and rectal temperature were plotted and the area under the curve

(AUC) was calculated to provide individual data points that captured change over time, with

AUCs compared with either two tailed unpaired t-test or two-way analysis of variance (ANOVA)

followed by *post-hoc* Bonferroni multiple comparisons test. Viral titers in lungs from infected

dams were analyzed using one-way or two-way ANOVA followed by *post-hoc* Bonferroni

multiple comparisons test. Multiplex cytokine quantifications were analyzed using multiple

unpaired t-tests with Holm-Šídák correction for multiple comparisons or two-way ANOVA

followed by *post-hoc* Bonferroni multiple comparisons test. Developmental milestones were

analyzed using Mann-Whitney test or Kruskal-Wallis test followed by *post-hoc* Dun's multiple

comparison test. Birth and neurodevelopment measurements, IHC quantification, and TUNEL

quantification were analyzed with two tailed unpaired t-test, two-way ANOVA followed by *post-*

hoc Bonferroni multiple comparisons test, or three-way ANOVA followed by *post-hoc* Bonferroni

multiple comparisons test. Concentrations P4, PGs, and TxB2 were analyzed with two-way

ANOVA followed by *post-hoc* Bonferroni multiple comparisons test. Western blot quantification

and flow cytometry quantification were analyzed with two tailed unpaired t-test. Mean or median

differences were considered statistically significant at $p < 0.05$. Statistical analyses were

performed using GraphPad Prism v10.0.2 (GraphPad Software).

2.15 Study approval

All animal procedures were approved by the Johns Hopkins University Animal Care and

Use Committee (MO21H246).

3 Results

3.1 Influenza A virus infection during pregnancy causes a pulmonary cytokine storm and

maternal morbidity in outbred mice

Mouse models of IAV infection during pregnancy primarily use inbred mice (Chan et al.,

2010; Kim et al., 2012; Littauer et al., 2017; Pazos et al., 2012; Vermillion et al., 2018) that are

devoid of fetal genetic heterogeneity (Engels et al., 2017; Jacobsen et al., 2021). Intranasal IAV

infection of pregnant outbred CD1 mice at E10, roughly corresponding to the human second

trimester (Sones and Davisson, 2016), resulted in significant morbidity, in which IAV-infected

dams gained less body mass (**Fig. 1A**) and experienced hypothermia (**Fig. 1B**) relative to mock-

inoculated dams. IAV-infected dams experienced greater hypothermia than nonpregnant

females (**Supplemental Fig. 1A)**. Subsets of dams were euthanized at 3, 6, and 8 days post

infection (dpi) to measure pulmonary infectious viral loads (**Fig. 1C**) and cytokine concentrations

in lungs and spleens (**Fig. 1D, Supplemental Table 1, Supplemental Table 2**) prior to and

during peak disease. Pulmonary viral load peaked at 3 dpi and was low to undetectable at 8 dpi

(**Fig. 1C**), which was consistent and not significantly different from IAV-infected nonpregnant

female mice (**Supplemental Fig. 1B**). Prototypical cytokines and chemokines associated with

influenza (Teijaro, 2015), including interferons (IFNs), tumor necrosis factor (TNF) α, interleukin

(IL)-6, and CCL2 were upregulated throughout infection in the lungs, but not the spleens of IAV-

infected dams (**Fig. 1D, Supplemental Table 1, Supplemental Table 2**). Together, these data

indicate that outbred pregnant mice infected with IAV experience morbidity to a greater degree

than nonpregnant females and a pulmonary cytokine storm, consistent with previous studies

(Chan et al., 2010; Kim et al., 2012; Vermillion et al., 2018).

3.2 Influenza A virus infection during pregnancy causes intrauterine growth restriction and

postnatal developmental delays

IAV infection during pregnancy is associated with adverse fetal outcomes including pre-

term delivery, fetal growth restriction, and low birthweights (Hartel et al., 2016; He et al., 2017;

Omer et al., 2012; Omer et al., 2011) as well as long-term health consequences in adulthood

(Mazumder et al., 2010; Song et al., 2020) in humans. In mice, IAV infection during pregnancy

did not impact litter size (**Fig. 2A**) or increase the rate of stillbirths (98.78% viability) versus

mock inoculated dams (98.82% viability). IAV infection of dams resulted in significant fetal

growth restriction, with pups of IAV-infected dams being smaller at birth based on measures of

mass, length, and head diameter, than offspring of mock-inoculated dams (**Fig. 2B-D**). Offspring

of IAV-infected dams remained smaller than offspring from mock-inoculated dams through

weaning (**Fig. 2E**). Offspring of IAV-infected dams also displayed developmental delays,

reaching key developmental milestones (Hill et al., 2008) on average 1-2 days later than

offspring of mock-inoculated dams (**Fig. 2F**). Impaired surface righting, a developmental reflex

(Feather-Schussler and Ferguson, 2016; Hill et al., 2008), also was apparent, with offspring,

particularly males, of IAV-infected dams taking significantly longer to right themselves (**Fig. 2G**).

Fetal growth restriction and developmental delays were not caused by pre-term birth as all

dams, regardless of maternal infection, delivered pups at approximately E20 (Elovitz et al.,

2003). These data indicate that IAV infection of outbred mice causes both short and long-term

adverse perinatal and fetal outcomes, consistent with humans (He et al., 2017; Mazumder et al.,

2010).

3.3 Influenza A virus infection during pregnancy causes transient but significant placental cell

death and trophoblast loss

While many models of IAV infection during pregnancy have focused on uncovering

mechanisms of maternal disease (Engels et al., 2017; Marcelin et al., 2011; Pazos et al., 2012;

Swieboda et al., 2020; Vermillion et al., 2018), the mechanisms of adverse fetal outcomes

remain less clear. Placental tissue damage, in the absence of direct viral infection, has been

reported in mice (Chan et al., 2010; Littauer et al., 2017; Swieboda et al., 2020) and in limited

observational human studies (Kanmaz et al., 2011; Meijer et al., 2014). Despite no infectious

virus in the placenta (data not shown), at 3 dpi, there was significant placental damage,

including mixing of maternal and fetal red blood cells (**Fig. 3A**), suggestive of disruption of the

placental trophoblast-endothelial cell barrier, as well as increased cell death as compared with

placentas from mock-inoculated dams (TUNEL[+] cells, **Fig. 3B** for representative image, **Fig. 3E**

for quantification). Staining for cytokeratin (trophoblasts, **Fig. 3C** for representative image, **Fig.**

F for quantification) and vimentin (endothelial cells, **Fig. 3D** for representative image, **Fig. 3G**

for quantification) revealed a loss of trophoblasts, but not endothelial cells, in placentas from

IAV-infected compared with mock-inoculated dams at 3 dpi. Concentrations of IL-13 (**Fig. 3H**),

granulocyte-macrophage colony-stimulating factor (GM-CSF, **Fig. 3I**), and IL-18 (**Fig. 3J**),

cytokines secreted by trophoblasts *in vitro* (Megli et al., 2021; Naruse et al., 2010), were lower

in placentas from IAV-infected than mock-inoculated dams at 3 dpi. By 6 and 8 dpi, neither

trophoblast nor endothelial cell numbers differed between placentas of IAV-infected and mock-

inoculated dams (**Supplemental Fig. 2**), indicating resolution of placental damage.

3.4 Proinflammatory cytokines peak in the placenta following placental damage during Influenza

A virus infection

Proinflammatory cytokines, such as IL-1β and TNFα, in the placenta are associated with

placental damage (Chudnovets et al., 2020; Creisher et al., 2022b; Li et al., 2018).

Concentrations of proinflammatory cytokines, including IL-1β (**Fig. 4A**), IL-6 (**Fig. 4B**), and

TNFα (**Fig. 4C**) exhibited a transient increase in placentas from IAV-infected compared with

mock-inoculated dams at 6dpi. The increased concentrations of proinflammatory cytokines

followed the placental damage and reductions in trophoblasts observed at 3dpi. The profile of

cytokine secretion in the placenta (**Supplemental Table 3**) differed from the patterns observed

in either the spleen (**Supplemental Table 2**) or lungs (**Supplemental Table 1**). Increased

secretion of proinflammatory cytokines in the placenta could be mediated by infiltration of

immune cells, as we have observed in models of intrauterine inflammation (Liu et al., 2021; Liu

et al., 2022). Flow cytometry was used to evaluate adaptive (i.e., B cells, T cells) and innate

(i.e., NK cells, neutrophils, eosinophils, macrophages) immune cells in the placenta at 6 dpi

(**Fig. 4D-L, Supplemental Fig. 3** for representative gating). Frequencies of CD45$^+$ leukocytes

were reduced in placentas of IAV-infected dams compared to mock-inoculated dams (**Fig 4D**),

which was further observed for NK cells (**Fig. 4E**), B cells (**Fig. 4F**), CD4$^+$ T cells (**Fig. 4G**), and

CD8$^+$ T cells (**Fig. 4H**) in placentas of IAV-infected dams. Frequencies of neutrophils (**Fig. 4I**),

eosinophils (**Fig. 4J**) and CD163$^+$ macrophages (**Fig. 4K**) were similar in placentas from IAV-

and mock-inoculated dams. CD68$^+$ macrophages were the only immune cell population that

exhibited a significant increase in placentas of IAV-infected compared with mock-inoculated

dams (**Fig. 4L**). The patterns of immune cell infiltrates observed in placentas were not observed

in peripheral blood cells obtained from the same dams (**Supplemental Fig. 4**). CD68$^+$

macrophages in the placenta are associated with intrauterine growth restriction in humans

(Berezhna et al., 2021) and secretion of proinflammatory cytokines (Ye et al., 2022). Taken

together, these results indicate that maternal respiratory infection with IAV can induce transient

but significant cell death and proinflammatory cytokine release at the maternal fetal interface, in

the absence of direct viral infection or broad leukocyte infiltration.

3.5 Influenza A virus infection suppresses the pregnancy-associated rise in progesterone via

COX-1 activity

The mechanism underlying placental damage and cytokine secretion during IAV

infection might involve sex hormone regulation. Placental development and maturation, which

occurs at E10-E14 in mice (Hemberger et al., 2020) are supported by P4 signaling in both

humans and mice (Di Renzo et al., 2016; Malassine et al., 2003; Terakawa et al., 2012; Wang

et al., 1995). As IAV infection during pregnancy has been associated with reduced circulating

P4 in mice (Kim et al., 2012; Littauer et al., 2017; Vermillion et al., 2018), we evaluated if IAV

infection of outbred pregnant mice altered circulating (**Fig. 5A**) and placental (**Fig. 5B**) P4

concentrations. Circulating and placental concentrations of P4 in mock-inoculated dams

increased throughout gestation and peaked at approximately E16 (**Fig. 5A-B**) (Murr et al.,

1974). In contrast, IAV infection significantly reduced circulating and placental P4

concentrations throughout gestation relative to mock-inoculated dams, although not to

circulating concentrations observed in nonpregnant females **(Fig. 5A**).

In mice, P4 is secreted by the corpus luteum of the ovary throughout the entire

pregnancy where it is canonically regulated by the prostaglandin (PG) F2α (Diaz et al., 2002;

Peluso et al., 1980). To evaluate if secreted PGF2α upregulation was associated with reduced

P4, PGF2α as well as PGE2, a prostaglandin that has been observed to be upregulated with

IAV infection in nonpregnant mice (Coulombe et al., 2014), were measured in the lungs (i.e., the

site of IAV replication), systemic circulation, and the ovaries (i.e., the site of P4 secretion) of

IAV-infected and mock inoculated dams at 1 and 3 dpi (**Fig. 5C-E**, **Supplemental Fig. 5**).

Concentrations of PGF2α, but not PGE2 (**Supplemental Fig.5A-C**), were significantly increased

in the lung (**Fig. 5C**), sera (**Fig. 5D**), and ovaries (**Fig. 5E**) of IAV-infected dams as compared

with mock-inoculated dams at 3 dpi. Elevated concentrations of PGFα in the ovary were

associated with reduced serum P4 in both IAV-infected and mock-inoculated dams

(**Supplemental Fig. 5D).**

Prostaglandins, including PGF2α are canonically synthesized by COX enzymes, with

COX-1 constitutively expressed in most cells and COX2 induced by inflammatory stimuli

(Ricciotti and FitzGerald, 2011). Active COX-1 (Otto et al., 1993), but not COX-2, expression

was elevated in the lungs of IAV-infected relative to mock-inoculated dams at 3dpi (**Fig. 5F-G**,

Supplemental Fig. 6). Neither COX-1 nor COX-2 expression were altered in the ovaries

(**Supplemental Fig. 7**), suggesting that PGF2α generation by COX-1 is local at the site of IAV

infection. To evaluate if COX-1 activity is not just sufficient but necessary for PGF2α secretion

and suppression of P4 during IAV-infection, subsets of IAV-infected or mock-inoculated dams at

E10 were pretreated with either a COX-1-specific inhibitor (SC560), a COX-2-specific inhibitor

(NS-398), or vehicle (**Fig. 5H-I**). COX-1 inhibition, as measured by inhibition of serum

thromboxane B2 (TXB2) (Wallace et al., 1998; Wallace et al., 1999), was specific to COX-1

inhibitor treatment and not observed with COX-2 inhibitor treatment (**Supplemental Fig. 8A**).

COX-1, but not COX-2, inhibition significantly reduced PGF2α and increased P4 secretion in

IAV-infected dams (**Fig. 5H-I**, **Supplemental Fig. 8B**). These data illustrate that IAV infection

disrupts activity along the arachidonic acid pathway in pregnant dams to cause suppression of

P4, which is required for a healthy pregnancy.

3.6 Synthetic progestin treatment improves IAV-induced maternal morbidity, placental damage,

and perinatal developmental delays

To establish whether IAV-infection induced P4 suppression was the cause of adverse

maternal and fetal outcomes, a subset of IAV-infected and mock inoculated dams were treated

with a single dose of either 17-α-hydroxyprogesterone caproate (17-OHPC), a synthetic

progestin that is safe to use during pregnancy (Sibai et al., 2021) or vehicle. The administration

of 17-OHPC to mock-inoculated dams had no adverse impact on pregnancy outcomes.

Administration of 17-OHPC to IAV-infected dams significantly reduced hypothermia as

compared with IAV-infected dams treated with vehicle (**Fig. 6A**). Treatment of IAV-infected

dams with 17-OHPC did not affect pulmonary infectious virus titers (**Fig. 6B**). These data

suggest that IAV-induced suppression of P4 causes maternal morbidity, independent of

controlling virus replication.

To determine if administration of 17-OHPC to IAV-infected dams could reverse adverse

perinatal outcomes, subsets of IAV-infected and mock-inoculated dams with and without 17-

OHPC treatment carried their pregnancies to term. Treatment of mock-inoculated dams with 17-

OHPC had no effect on the development of offspring relative to vehicle-treated, mock-infected

dams. In contrast, treatment of IAV-infected dams with 17-OHPC significantly increased birth

mass (**Fig. 6C**), mitigated developmental delays in key milestones (**Fig. 6D**), and improved

neurobehavioral development as measured by surface righting at PND5 (**Fig. 6E**) as compared

with offspring from IAV-infected dams that were treated with vehicle. These data demonstrate

that IAV-induced suppression of P4 during pregnancy causes intrauterine growth restriction and

postnatal developmental delays, which are reversed by treatment with a safe, synthetic

progestin.

We hypothesized that 17-OHPC treatment might prevent placental damage to improve

offspring outcomes among IAV-infected dams. Treatment of mock-inoculated dams with 17-

OHPC did not alter placental histology, cell death, or numbers of trophoblasts as compared with

vehicle-treated, mock-infected dams. In contrast, treatment of IAV-infected dams with 17-OHPC

significantly reduced placental cell death (**Fig. 7A-B**), increased proportions of trophoblasts

(**Fig. 7C-D**), and reduced placental concentrations of proinflammatory cytokines (**Supplemental**

Table 4), including IL-1β (**Fig. 7E**), IL-6 (**Fig. 7F**), and TNFα (**Fig. 7G**). These data illustrate that

suppressed P4 is the cause of placental tissue damage and adverse perinatal outcomes during

IAV-infection and treatment with 17-OHPC improved maternal, placental, and offspring

outcomes during infection.

4 Discussion

Both seasonal and pandemic influenza virus infection during pregnancy result in adverse

perinatal and long-term detrimental outcomes that have been reported since the 1918 pandemic

(He et al., 2017; Mazumder et al., 2010; Noymer and Garenne, 2000). Using an outbred mouse

model of IAV infection during pregnancy, 2009 H1N1 infection caused maternal morbidity,

transient placental damage and inflammation, and long-term offspring growth restriction and

developmental delays. The IAV-induced adverse maternal and fetal outcomes were mediated

by a COX-1-dependent reduction in P4, including at the maternal-fetal interface. Previous

studies report damage to tissue architecture, necrosis, and pro-inflammatory cytokines in the

placenta following maternal IAV infection (Antonson et al., 2021; Chan et al., 2010; Van

Campen et al., 2020), including associations between placental damage and reduced placental

P4 (Littauer et al., 2017; Swieboda et al., 2020). Our study expands upon this literature by

establishing a novel causal role for the arachidonic acid biosynthesis pathway, which is

activated by IAV infection and disrupts P4 during pregnancy. Our data provide a mechanistic

connection between a pulmonary virus infection, which causes elevated COX-1 enzymatic

activity, and placental tissue damage, caused by increased prostaglandin (PG) and reduced P4

synthesis. Our results are consistent with knockout studies where selective disruption of

progesterone receptor signaling in immune cells causes damage to the placenta, intrauterine

growth restriction, and fetal wasting (Severance et al., 2022; Thiele et al., 2019). By treating

dams with a synthetic progestin that is safe during pregnancy, the IAV-induced placental

damage, inflammation, and adverse fetal outcomes, were reversed, highlighting the translational

significance of these findings.

IAV-induced suppression of P4 was dependent on COX-1, but not COX-2, signaling and

PGF2α secretion. Progesterone secretion during pregnancy is canonically regulated by PGF2α,

which signals in the corpus luteum and induces luteolysis to inhibit P4 secretion (Diaz et al.,

2002). PGF2α is generated within the arachidonic acid biosynthesis pathway, where COX

enzymes, predominantly COX-1 or COX-2, convert arachidonic acid into prostanoids that are

then converted to PG subclasses (i.e., PGF2α) by specific PG synthases (Dubois et al., 1998;

Wang et al., 2021). Historically, COX-1 has been characterized as constitutively active and

homeostatic, while COX-2 has been characterized as inducible and associated with pain and

inflammation (Adelizzi, 1999; Dubois et al., 1998). COX-1, however, also may have functions in

inflammatory responses, as COX-1 inhibition can alleviate neuroinflammation in a mouse model

of retinitis pigmentosa (Yang et al., 2020). Further, aspirin given at COX-1-specific doses is

associated with reduced rates of preeclampsia (Rolnik et al., 2022). Our data add to this

growing body of literature by illustrating that IAV infection upregulates COX-1 in the lungs and

increased COX-1-dependent PGF2α in several tissues, including the ovaries.

Mouse models of pregnancy are powerful tools to identify mechanisms of disease

pathogenesis and evaluate the safety and efficacy of potential treatments (Vermillion and Klein,

2018); mice, however, have limitations. The full-term gestation for mice is approximately 20

days (about 3 weeks), but 40 weeks (about 9 months) for humans, with disproportionally more

gestational time spent in the first-trimester equivalent in mice than in humans (Sones and

Davisson, 2016). Further, placental anatomy differs between mice and humans [reviewed in

(Ander et al., 2019)], with, for example, the placental labyrinth structure found in mice but not

humans. Inbred mice further lack genetic heterogeneity at the maternal-fetal interface, and in

the current study, outbred mice were used to avoid this limitation (Reichenbach et al., 2013).

With the use of outbred mice, genetic manipulations were not available to test hypotheses;

instead, pharmacologic inhibitors were used, which could have off-target effects, despite

dosages optimized for target-specificity.

Despite these limitations, results from this study highlight the value of treatments that

mitigate disease in addition to antiviral treatments for influenza during pregnancy. In the current

study, we utilized 17-OHPC, a synthetic progestin that has been prescribed to pregnant women

with history of preterm birth (EPPPIC Group, 2021). While the US FDA has withdrawn approval

due to a lack of efficacy of 17-OHPC for preterm birth (U.S. Food & Drug Administration, 2023),

translational studies demonstrated that progesterone and 17-OHPC, specifically, may contribute

to maternal immunomodulation and prevention of adverse perinatal outcomes other than

preterm birth (Novak et al., 2018; Severance et al., 2022). Furthermore, while nonselective COX

inhibitors [i.e. non-steroidal anti-inflammatory drugs (NSAIDs)] are generally not recommend for

use during pregnancy (Ostensen and Skomsvoll, 2004; U.S. Food & Drug Administration, 2022),

we have identified COX-1-dependent P4 regulation as a targetable pathway that warrants

further investigation. Our results add to growing literature utilizing P4 to treat adverse pregnancy

and perinatal outcomes (Alawadhi et al., 2022; Novak et al., 2018; Pei et al., 2022; Severance

et al., 2022). With pregnant people considered an 'at risk' population for many infectious

diseases ranging from influenza viruses and SARS-CoV-2 to Zika viruses, greater consideration

should be given to identifying the mechanisms of adverse pregnancy outcomes and the off label

use of compounds (e.g., 17-OHPC) that are safe during pregnancy and could mitigate the

impact of infectious diseases at the maternal-fetal interface.

5 Acknowledgements

The authors would like to thank the Klein, Pekosz, Davis, and Baumgarth laboratories for

discussions about these data. We would also like to thank the expert animal care staff at the

Johns Hopkins School of Public Health for assistance with maintenance of pregnant dams.

6 Funding

Funding provided by NIH/NICHD R01HD097608 (I.B. and S.L.K), the NIH/ORWH/NIA

Specialized Center of Research Excellence in Sex Differences U54AG062333 (S.L.K),

NIH/NIAID training grant T32AI007417-26 (P.C. and M.S.), the Vivien Thomas Scholars

Initiative at the Johns Hopkins University (A.C.).

7 Contributions

SK, PC, MP, and IB conceptualized and designed the experiments. PC, MP, JP, AC, and MS

performed animal experiments. PC grew and quantified viruses and multiplexed cytokines. PC

and MP performed ELISA and Western blot. PC, JP, and JinL performed flow cytometry. JunL

stained, imaged, and analyzed placental tissue. PC statistically analyzed and graphed data. PC

and SK wrote the manuscript with input from all authors. All authors read and provided edits to

drafts and approved the final submission. The order of authors was determined based on

contributions to the overall design, experimentation, analyses, and writing.

8 References

Abu-Raya, B., C. Michalski, M. Sadarangani, and P.M. Lavoie. 2020. Maternal Immunological
565   Adaptation During Normal Pregnancy. *Front Immunol* 11:575197.
566   Adelizzi, R.A. 1999. COX-1 and COX-2 in health and disease. *J Am Osteopath Assoc* 99:S7-12.
567   Alawadhi, M., A. Mouihate, N. Kilarkaje, and M. Al-Bader. 2022. Progesterone partially recovers
568   placental glucose transporters in dexamethasone-induced intrauterine growth
569   restriction. *Reprod Biomed Online* 44:595-607.
570   Ander, S.E., M.S. Diamond, and C.B. Coyne. 2019. Immune responses at the maternal-fetal
571   interface. *Sci Immunol* 4:
572   Antonson, A.M., A.D. Kenney, H.J. Chen, K.N. Corps, J.S. Yount, and T.L. Gur. 2021. Moderately
573   pathogenic maternal influenza A virus infection disrupts placental integrity but spares
574   the fetal brain. *Brain Behav Immun* 96:28-39.
575   Arany, Z., D. Hilfiker-Kleiner, and S.A. Karumanchi. 2022. Animal Models of Cardiovascular
576   Complications of Pregnancy. *Circ Res* 130:1763-1779.
577   Berezhna, V.A., T.V. Mamontova, and A.M. Gromova. 2021. Cd68+ M1 Macrophages Is
578   Associated with Placental Insufficiency under Fetal Growth Restriction. *Wiad Lek* 74:213-
579   219.
580   Brown, A.S., M.D. Begg, S. Gravenstein, C.A. Schaefer, R.J. Wyatt, M. Bresnahan, V.P. Babulas,
581   and E.S. Susser. 2004. Serologic evidence of prenatal influenza in the etiology of
582   schizophrenia. *Arch Gen Psychiatry* 61:774-780.
583   Chan, K.H., A.J. Zhang, K.K. To, C.C. Chan, V.K. Poon, K. Guo, F. Ng, Q.W. Zhang, V.H. Leung, A.N.
584   Cheung, C.C. Lau, P.C. Woo, H. Tse, W. Wu, H. Chen, B.J. Zheng, and K.Y. Yuen. 2010.
585   Wild type and mutant 2009 pandemic influenza A (H1N1) viruses cause more severe
586   disease and higher mortality in pregnant BALB/c mice. *PLoS One* 5:e13757.
587   Chudnovets, A., J. Lei, Q. Na, J. Dong, H. Narasimhan, S.L. Klein, and I. Burd. 2020. Dose-
588   dependent structural and immunological changes in the placenta and fetal brain in
589   response to systemic inflammation during pregnancy. *Am J Reprod Immunol* 84:e13248.
590   Coulombe, F., J. Jaworska, M. Verway, F. Tzelepis, A. Massoud, J. Gillard, G. Wong, G. Kobinger,
591   Z. Xing, C. Couture, P. Joubert, J.H. Fritz, W.S. Powell, and M. Divangahi. 2014. Targeted
592   prostaglandin E2 inhibition enhances antiviral immunity through induction of type I
593   interferon and apoptosis in macrophages. *Immunity* 40:554-568.
594   Cox, S., S.F. Posner, M. McPheeters, D.J. Jamieson, A.P. Kourtis, and S. Meikle. 2006. Influenza
595   and pregnant women: hospitalization burden, United States, 1998-2002. *J Womens
596   Health (Larchmt)* 15:891-893.

Creisher, P.S., A.D. Campbell, J.L. Perry, K. Roznik, I. Burd, and S.L. Klein. 2022a. Influenza
subtype-specific maternal antibodies protect offspring against infection but inhibit
vaccine-induced immunity and protection in mice. *Vaccine* 40:6818-6829.
Creisher, P.S., J. Lei, M.L. Sherer, A. Dziedzic, A.E. Jedlicka, H. Narasimhan, A. Chudnovets, A.D.
Campbell, A. Liu, A. Pekosz, I. Burd, and S.L. Klein. 2022b. Downregulation of
transcriptional activity, increased inflammation, and damage in the placenta following in
utero Zika virus infection is associated with adverse pregnancy outcomes. *Front Virol* 2:
Creisher, P.S., J.L. Perry, W. Zhong, J. Lei, K.R. Mulka, W.H. Ryan, R. Zhou, E.H. Akin, A. Liu, W.
Mitzner, I. Burd, A. Pekosz, and S.L. Klein. 2023. Adverse outcomes in SARS-CoV-2
infected pregnant mice are gestational age-dependent and resolve with antiviral
treatment. *The Journal of Clinical Investigation*
Di Renzo, G.C., I. Giardina, G. Clerici, E. Brillo, and S. Gerli. 2016. Progesterone in normal and
pathological pregnancy. *Horm Mol Biol Clin Investig* 27:35-48.
Diaz, F.J., L.E. Anderson, Y.L. Wu, A. Rabot, S.J. Tsai, and M.C. Wiltbank. 2002. Regulation of
progesterone and prostaglandin F2alpha production in the CL. *Mol Cell Endocrinol*
191:65-80.
Dubois, R.N., S.B. Abramson, L. Crofford, R.A. Gupta, L.S. Simon, L.B. Van De Putte, and P.E.
Lipsky. 1998. Cyclooxygenase in biology and disease. *FASEB J* 12:1063-1073.
Elovitz, M.A., Z. Wang, E.K. Chien, D.F. Rychlik, and M. Phillippe. 2003. A new model for
inflammation-induced preterm birth: the role of platelet-activating factor and Toll-like
receptor-4. *Am J Pathol* 163:2103-2111.
Engels, G., A.M. Hierweger, J. Hoffmann, R. Thieme, S. Thiele, S. Bertram, C. Dreier, P. Resa-
Infante, H. Jacobsen, K. Thiele, M. Alawi, D. Indenbirken, A. Grundhoff, S. Siebels, N.
Fischer, V. Stojanovska, D. Muzzio, F. Jensen, K. Karimi, H.W. Mittrucker, P.C. Arck, and
G. Gabriel. 2017. Pregnancy-Related Immune Adaptation Promotes the Emergence of
Highly Virulent H1N1 Influenza Virus Strains in Allogenically Pregnant Mice. *Cell Host*
Microbe 21:321-333.
EPPPIC Group. 2021. Evaluating Progestogens for Preventing Preterm birth International
Collaborative (EPPPIC): meta-analysis of individual participant data from randomised
controlled trials. *Lancet* 397:1183-1194.
Feather-Schussler, D.N., and T.S. Ferguson. 2016. A Battery of Motor Tests in a Neonatal Mouse
Model of Cerebral Palsy. *J Vis Exp*
Futaki, N., K. Yoshikawa, Y. Hamasaka, I. Arai, S. Higuchi, H. Iizuka, and S. Otomo. 1993. NS-398,
a novel non-steroidal anti-inflammatory drug with potent analgesic and antipyretic
effects, which causes minimal stomach lesions. *Gen Pharmacol* 24:105-110.
Hall, O.J., R. Nachbagauer, M.S. Vermillion, A.L. Fink, V. Phuong, F. Krammer, and S.L. Klein.
2017. Progesterone-Based Contraceptives Reduce Adaptive Immune Responses and
Protection against Sequential Influenza A Virus Infections. *J Virol* 91:
Hartel, C., A. Humberg, D. Viemann, A. Stein, T. Orlikowsky, J. Rupp, M.V. Kopp, E. Herting, and
W. Gopel. 2016. Preterm Birth during Influenza Season Is Associated with Adverse
Outcome in Very Low Birth Weight Infants. *Front Pediatr* 4:130.
He, J., Z.W. Liu, Y.P. Lu, T.Y. Li, X.J. Liang, P.C. Arck, S.M. Huang, B. Hocher, and Y.P. Chen. 2017.
A Systematic Review and Meta-Analysis of Influenza A Virus Infection During Pregnancy

Associated with an Increased Risk for Stillbirth and Low Birth Weight. *Kidney Blood Press*
Res 42:232-243.

Hemberger, M., C.W. Hanna, and W. Dean. 2020. Mechanisms of early placental development
in mouse and humans. *Nat Rev Genet* 21:27-43.

Heyser, C.J. 2004. Assessment of developmental milestones in rodents. *Curr Protoc Neurosci*
Chapter 8:Unit 8 18.

Hill, J.M., M.A. Lim, and M.M. Stone. 2008. Developmental Milestones in the Newborn Mouse.
In Neuropeptide Techniques. I. Gozes, editor Humana Press, Totowa, NJ. 131-149.

Jacobsen, H., K. Walendy-Gnirss, N. Tekin-Bubenheim, N.M. Kouassi, I. Ben-Batalla, N.
Berenbrok, M. Wolff, V.P. Dos Reis, M. Zickler, L. Scholl, A. Gries, H. Jania, A. Kloetgen, A.
Dusedau, G. Pilnitz-Stolze, A. Jeridi, A.O. Yildirim, H. Fuchs, V. Gailus-Durner, C. Stoeger,
M.H. de Angelis, T. Manuylova, K. Klingel, F.J. Culley, J. Behrends, S. Loges, B. Schneider,
S. Krauss-Etschmann, P. Openshaw, and G. Gabriel. 2021. Offspring born to influenza A
virus infected pregnant mice have increased susceptibility to viral and bacterial
infections in early life. *Nat Commun* 12:4957.

Kanmaz, H.G., O. Erdeve, S.S. Ogz, N. Uras, S. Celen, G. Korukluoglu, S. Zergeroglu, A. Kara, and
U. Dilmen. 2011. Placental transmission of novel pandemic influenza a virus. *Fetal*
Pediatr Pathol 30:280-285.

Katz, M.A., B.D. Gessner, J. Johnson, B. Skidmore, M. Knight, N. Bhat, H. Marshall, D.J. Horne,
J.R. Ortiz, and D.B. Fell. 2017. Incidence of influenza virus infection among pregnant
women: a systematic review. *BMC Pregnancy Childbirth* 17:155.

Kim, H.M., Y.M. Kang, B.M. Song, H.S. Kim, and S.H. Seo. 2012. The 2009 pandemic H1N1
influenza virus is more pathogenic in pregnant mice than seasonal H1N1 influenza virus.
Viral Immunol 25:402-410.

Klein, S.L., C. Passaretti, M. Anker, P. Olukoya, and A. Pekosz. 2010. The impact of sex, gender
and pregnancy on 2009 H1N1 disease. *Biol Sex Differ* 1:5.

Krishnan, L., T. Nguyen, and S. McComb. 2013. From mice to women: the conundrum of
immunity to infection during pregnancy. *J Reprod Immunol* 97:62-73.

Lauzon-Joset, J.F., N.M. Scott, K.T. Mincham, P.A. Stumbles, P.G. Holt, and D.H. Strickland. 2019.
Pregnancy Induces a Steady-State Shift in Alveolar Macrophage M1/M2 Phenotype That
Is Associated With a Heightened Severity of Influenza Virus Infection: Mechanistic
Insight Using Mouse Models. *J Infect Dis* 219:1823-1831.

Li, Z.H., L.L. Wang, H. Liu, K.P. Muyayalo, X.B. Huang, G. Mor, and A.H. Liao. 2018. Galectin-9
Alleviates LPS-Induced Preeclampsia-Like Impairment in Rats via Switching Decidual
Macrophage Polarization to M2 Subtype. *Front Immunol* 9:3142.

Littauer, E.Q., E.S. Esser, O.Q. Antao, E.V. Vassilieva, R.W. Compans, and I. Skountzou. 2017.
H1N1 influenza virus infection results in adverse pregnancy outcomes by disrupting
tissue-specific hormonal regulation. *PLoS Pathog* 13:e1006757.

Liu, J., Y. Liu, S. Panda, A. Liu, J. Lei, and I. Burd. 2021. Type 1 Cytotoxic T Cells Increase in
Placenta after Intrauterine Inflammation. *Front Immunol* 12:718563.

Liu, Y., Q. Na, J. Liu, A. Liu, A. Oppong, J.Y. Lee, A. Chudnovets, J. Lei, R. Sharma, S. Kannan, R.M.
Kannan, and I. Burd. 2022. Dendrimer-Based N-Acetyl Cysteine Maternal Therapy

Ameliorates Placental Inflammation via Maintenance of M1/M2 Macrophage
Recruitment. *Front Bioeng Biotechnol* 10:819593.
Malassine, A., J.L. Frendo, and D. Evain-Brion. 2003. A comparison of placental development
and endocrine functions between the human and mouse model. *Hum Reprod Update*
9:531-539.
Marcelin, G., J.R. Aldridge, S. Duan, H.E. Ghoneim, J. Rehg, H. Marjuki, A.C. Boon, J.A. McCullers,
and R.J. Webby. 2011. Fatal outcome of pandemic H1N1 2009 influenza virus infection is
associated with immunopathology and impaired lung repair, not enhanced viral burden,
in pregnant mice. *J Virol* 85:11208-11219.
Mazumder, B., D. Almond, K. Park, E.M. Crimmins, and C.E. Finch. 2010. Lingering prenatal
effects of the 1918 influenza pandemic on cardiovascular disease. *J Dev Orig Health Dis*
1:26-34.
Megli, C., S. Morosky, D. Rajasundaram, and C.B. Coyne. 2021. Inflammasome signaling in
human placental trophoblasts regulates immune defense against Listeria
monocytogenes infection. *J Exp Med* 218:
Meijer, W.J., A.M. Wensing, H.W. Bruinse, and P.G. Nikkels. 2014. High rate of chronic villitis in
placentas of pregnancies complicated by influenza A/H1N1 infection. *Infect Dis Obstet*
Gynecol 2014:768380.
Mertz, D., J. Geraci, J. Winkup, B.D. Gessner, J.R. Ortiz, and M. Loeb. 2017. Pregnancy as a risk
factor for severe outcomes from influenza virus infection: A systematic review and
meta-analysis of observational studies. *Vaccine* 35:521-528.
Moldenhauer, L.M., J.D. Hayball, and S.A. Robertson. 2010. Utilising T cell receptor transgenic
mice to define mechanisms of maternal T cell tolerance in pregnancy. *J Reprod Immunol*
87:1-13.
Murr, S.M., G.H. Stabenfeldt, G.E. Bradford, and Geschwind, II. 1974. Plasma progesterone
during pregnancy in the mouse. *Endocrinology* 94:1209-1211.
Naruse, K., B.A. Innes, J.N. Bulmer, S.C. Robson, R.F. Searle, and G.E. Lash. 2010. Secretion of
cytokines by villous cytotrophoblast and extravillous trophoblast in the first trimester of
human pregnancy. *J Reprod Immunol* 86:148-150.
Neuzil, K.M., G.W. Reed, E.F. Mitchel, L. Simonsen, and M.R. Griffin. 1998. Impact of influenza
on acute cardiopulmonary hospitalizations in pregnant women. *Am J Epidemiol*
148:1094-1102.
Novak, C.M., M. Ozen, M. McLane, S. Alqutub, J.Y. Lee, J. Lei, and I. Burd. 2018. Progesterone
improves perinatal neuromotor outcomes in a mouse model of intrauterine
inflammation via immunomodulation of the placenta. *Am J Reprod Immunol* 79:e12842.
Noymer, A., and M. Garenne. 2000. The 1918 influenza epidemic's effects on sex differentials in
mortality in the United States. *Popul Dev Rev* 26:565-581.
Omer, S.B., R. Bednarczyk, S.A. Madhi, and K.P. Klugman. 2012. Benefits to mother and child of
influenza vaccination during pregnancy. *Hum Vaccin Immunother* 8:130-137.
Omer, S.B., D. Goodman, M.C. Steinhoff, R. Rochat, K.P. Klugman, B.J. Stoll, and U.
Ramakrishnan. 2011. Maternal influenza immunization and reduced likelihood of
prematurity and small for gestational age births: a retrospective cohort study. *PLoS Med*
8:e1000441.

Ostensen, M.E., and J.F. Skomsvoll. 2004. Anti-inflammatory pharmacotherapy during
pregnancy. *Expert Opin Pharmacother* 5:571-580.
Otto, J.C., D.L. DeWitt, and W.L. Smith. 1993. N-glycosylation of prostaglandin endoperoxide
synthases-1 and -2 and their orientations in the endoplasmic reticulum. *J Biol Chem*
268:18234-18242.
Parboosing, R., Y. Bao, L. Shen, C.A. Schaefer, and A.S. Brown. 2013. Gestational influenza and
bipolar disorder in adult offspring. *JAMA Psychiatry* 70:677-685.
Pazos, M.A., T.A. Kraus, C. Munoz-Fontela, and T.M. Moran. 2012. Estrogen mediates innate
and adaptive immune alterations to influenza infection in pregnant mice. *PLoS One*
7:e40502.
Pei, J., Z. Liu, C. Wang, N. Chu, L. Liu, Y. Tang, H. Liu, Q. Xiang, H. Cheng, M. Li, and W. Gu. 2022.
Progesterone Attenuates SIRT1-Deficiency-Mediated Pre-Eclampsia. *Biomolecules* 12:
Peluso, J.J., S. Leiper, and R.W. Steger. 1980. Effects of PGF2 alpha-induced luteolysis and
progesterone on follicular growth in pregnant mice. *Prostaglandins* 19:437-447.
Reichenbach, D.K., Q. Li, R.A. Hoffman, A.L. Williams, W.D. Shlomchik, D.M. Rothstein, A.J.
Demetris, and F.G. Lakkis. 2013. Allograft outcomes in outbred mice. *Am J Transplant*
13:580-588.
Ricciotti, E., and G.A. FitzGerald. 2011. Prostaglandins and inflammation. *Arterioscler Thromb*
Vasc Biol 31:986-1000.
Rolnik, D.L., K.H. Nicolaides, and L.C. Poon. 2022. Prevention of preeclampsia with aspirin. *Am J*
Obstet Gynecol 226:S1108-S1119.
Severance, A.L., J.M. Kinder, L. Xin, A.R. Burg, T.Y. Shao, G. Pham, T. Tilburgs, W.A. Goodman, S.
Mesiano, and S.S. Way. 2022. Maternal-fetal conflict averted by progesterone- induced
FOXP3+ regulatory T cells. *iScience* 25:104400.
Shaik, I.H., J.R. Bastian, Y. Zhao, S.N. Caritis, and R. Venkataramanan. 2016. Route of
administration and formulation dependent pharmacokinetics of 17-
hydroxyprogesterone caproate in rats. *Xenobiotica* 46:169-174.
Sibai, B., G.R. Saade, A.F. Das, and J. Gudeman. 2021. Safety review of hydroxyprogesterone
caproate in women with a history of spontaneous preterm birth. *J Perinatol* 41:718-725.
Sones, J.L., and R.L. Davisson. 2016. Preeclampsia, of mice and women. *Physiol Genomics*
48:565-572.
Song, J.Y., K.V. Park, S.W. Han, M.J. Choi, J.Y. Noh, H.J. Cheong, W.J. Kim, M.J. Oh, and G.J. Cho.
2020. Paradoxical long-term impact of maternal influenza infection on neonates and
infants. *BMC Infect Dis* 20:502.
Swieboda, D., E.Q. Littauer, J.T. Beaver, L.K. Mills, K.M. Bricker, E.S. Esser, O.Q. Antao, D.T.
Williams, and I. Skountzou. 2020. Pregnancy Downregulates Plasmablast Metabolic
Gene Expression Following Influenza Without Altering Long-Term Antibody Function.
Front Immunol 11:1785.
Tafuri, A., J. Alferink, P. Moller, G.J. Hammerling, and B. Arnold. 1995. T cell awareness of
paternal alloantigens during pregnancy. *Science* 270:630-633.
Tanigawa, T., T. Watanabe, M. Hamaguchi, E. Sasaki, K. Tominaga, Y. Fujiwara, N. Oshitani, T.
Matsumoto, K. Higuchi, and T. Arakawa. 2004. Anti-inflammatory effect of two isoforms

of COX in H. pylori-induced gastritis in mice: possible involvement of PGE2. *Am J Physiol*
Gastrointest Liver Physiol 286:G148-156.

Teijaro, J.R. 2015. The role of cytokine responses during influenza virus pathogenesis and
potential therapeutic options. *Curr Top Microbiol Immunol* 386:3-22.

Terakawa, J., T. Watanabe, R. Obara, M. Sugiyama, N. Inoue, Y. Ohmori, Y.Z. Hosaka, and E.
Hondo. 2012. The complete control of murine pregnancy from embryo implantation to
parturition. *Reproduction* 143:411-415.

Thiele, K., A.M. Hierweger, J.I.A. Riquelme, M.E. Solano, J.P. Lydon, and P.C. Arck. 2019.
Impaired Progesterone-Responsiveness of CD11c(+) Dendritic Cells Affects the
Generation of CD4(+) Regulatory T Cells and Is Associated With Intrauterine Growth
Restriction in Mice. *Front Endocrinol (Lausanne)* 10:96.

Thomas, D.M., and D.M. Kuhn. 2005. Cyclooxygenase-2 is an obligatory factor in
methamphetamine-induced neurotoxicity. *J Pharmacol Exp Ther* 313:870-876.

U.S. Food & Drug Administration. 2022. FDA recommends avoiding use of NSAIDs in pregnancy
at 20 weeks or later because they can result in low amniotic fluid. In.

U.S. Food & Drug Administration. 2023. FDA Commissioner and Chief Scientist Announce
Decision to Withdraw Approval of Makena. In.

Van Campen, H., J.V. Bishop, V.M. Abrahams, H. Bielefeldt-Ohmann, C.K. Mathiason, G.J.
Bouma, Q.A. Winger, C.E. Mayo, R.A. Bowen, and T.R. Hansen. 2020. Maternal Influenza
A Virus Infection Restricts Fetal and Placental Growth and Adversely Affects the Fetal
Thymic Transcriptome. *Viruses* 12:

Vermillion, M.S., and S.L. Klein. 2018. Pregnancy and infection: using disease pathogenesis to
inform vaccine strategy. *NPJ Vaccines* 3:6.

Vermillion, M.S., A. Nelson, L. Vom Steeg, J. Loube, W. Mitzner, and S.L. Klein. 2018. Pregnancy
preserves pulmonary function following influenza virus infection in C57BL/6 mice. *Am J*
Physiol Lung Cell Mol Physiol 315:L517-L525.

Versteeg, L., X. Le Guezennec, B. Zhan, Z. Liu, M. Angagaw, J.D. Woodhouse, S. Biswas, and C.M.
Beaumier. 2017. Transferring Luminex(R) cytokine assays to a wall-less plate technology:
Validation and comparison study with plasma and cell culture supernatants. *J Immunol*
Methods 440:74-82.

Vousden, N., K. Bunch, M. Knight, and U.I.C.-I. Group. 2021. Incidence, risk factors and impact
of seasonal influenza in pregnancy: A national cohort study. *PLoS One* 16:e0244986.

Wallace, J.L., A. Bak, W. McKnight, S. Asfaha, K.A. Sharkey, and W.K. MacNaughton. 1998.
Cyclooxygenase 1 contributes to inflammatory responses in rats and mice: implications
for gastrointestinal toxicity. *Gastroenterology* 115:101-109.

Wallace, J.L., K. Chapman, and W. McKnight. 1999. Limited anti-inflammatory efficacy of cyclo-
oxygenase-2 inhibition in carrageenan-airpouch inflammation. *Br J Pharmacol* 126:1200-
1204.

Wang, B., L. Wu, J. Chen, L. Dong, C. Chen, Z. Wen, J. Hu, I. Fleming, and D.W. Wang. 2021.
Metabolism pathways of arachidonic acids: mechanisms and potential therapeutic
targets. *Signal Transduct Target Ther* 6:94.

Wang, M.W., D.L. Crombie, J.S. Hayes, and R.B. Heap. 1995. Aberrant maternal behaviour in
mice treated with a progesterone receptor antagonist during pregnancy. *J Endocrinol*
145:371-377.
Woolston, W.J., and D.O. Conley. 1918. Epidemic Pneumonia (Spanish Influenza) in Pregnancy:
Effect in One Hundred and One Cases. *Journal of the American Medical Association*
71:1898-1899.
Yang, W., G. Xiong, and B. Lin. 2020. Cyclooxygenase-1 mediates neuroinflammation and
neurotoxicity in a mouse model of retinitis pigmentosa. *J Neuroinflammation* 17:306.
Ye, J., E.M. Sorrell, Y. Cai, H. Shao, K. Xu, L. Pena, D. Hickman, H. Song, M. Angel, R.A. Medina, B.
Manicassamy, A. Garcia-Sastre, and D.R. Perez. 2010. Variations in the hemagglutinin of
the 2009 H1N1 pandemic virus: potential for strains with altered virulence phenotype?
PLoS Pathog 6:e1001145.
Ye, Y., L. Peng, A. Chelariu-Raicu, C. Kuhn, X. Dong, U. Jeschke, and V. von Schonfeldt. 2022.
Prostaglandin E2 receptor 3 promotes M1 macrophages polarization in unexplained
recurrent pregnancy loss. *Biol Reprod* 106:910-918.
823
Figure Legends

Figure 1. Influenza A virus infection of pregnant outbred mice results in morbidity and a

pulmonary cytokine storm.

Dams at embryonic day (E) 10 were intranasally inoculated with 10^3 $TCID_{50}$ of 2009 H1N1 or

were mock inoculated with media. Daily maternal body mass (A) and rectal temperature (B)

were recorded as measures of maternal morbidity for 14 days. Subsets of dams were

euthanized to measure viral titers in lung tissue via $TCID_{50}$ assay (C) and cytokine

concentrations in spleen and lung tissue (D) at 3 (E13), 6 (E16), and 8 (E18) days post

infection. Individual symbols (A-B) or bars (C) represent the mean ± standard error of the mean

from two independent replications (n =12-13/group) with individual mice indicated by symbols

(C). Significant differences (p < 0.05) were determined by two tailed unpaired t-test of AUCs (A-

B) or one-way ANOVA with Bonferroni post hoc test (C) and are indicated by an asterisk (*).

Figure 2. Influenza A virus (IAV) infection during pregnancy causes offspring growth

restriction and developmental delays.

Dams at embryonic day (E) 10 were intranasally inoculated with 10^3 TCID$_{50}$ of 2009 H1N1 or

were mock inoculated with media and were followed through delivery. Offspring were followed

through postnatal day (PND) 21 to assess fetal outcomes. Litter size (A), mass (B), nose to

anus length (C), and ear to head diameter (D) were measured at PND 0. Average

measurements of each independent litter were graphed to account for litter effects. Average pup

mass for each litter was taken daily to track growth through PND 21 (E). Developmental

milestones including fur growth, ear canal opening, eye opening, and teeth eruption were

recorded (F). At PND 5, a subset of pups was sexed and assessed for their surface righting

ability to measure neurological development. One pup per sex per dam was given 3 trials given

for the test, and each pup's best trial was reported (G). Bars (A-D, G) represent the mean ±

standard error of the mean from two independent replications (n =13/group) with individual

litters (A-D) or mice (G) indicated by symbols. For developmental milestones, the center bar

indicates the median day a litter reached each milestone, with the range indicated by the edges

of the box (D). Significant differences (p < 0.05) were determined by two tailed unpaired t-test

(A-D), two tailed unpaired t-test of AUCs (E), Mann-Whitney nonparametric test (F), or two-way

ANOVA with Bonferroni post hoc test (G) and are indicated by an asterisk (*).

Figure 3. Influenza A virus infection during pregnancy causes placental damage and

disruption to the trophoblast-endothelial cell barrier.

Dams at embryonic day (E) 10 were intranasally inoculated with 10^3 TCID$_{50}$ of 2009 H1N1 or

were mock inoculated with media and were euthanized at 3 dpi (E13) to collect placentas.

Representative H&E images were taken at 5x magnification and 20x magnification with arrows

indicating red blood cells (A). Representative TUNEL$^+$ (green) stained placentas with DAPI to

label nuclei taken (blue) were taken at 20x magnification and 100x (B).

Placentas were immunostained for cytokeratin (C, red) to mark trophoblasts or vimentin (D, red)

to mark endothelial cells and DAPI (blue) to label nuclei. Representative images were taken at

20x magnification and further zoomed 2.90-fold. Quantification of the percentage positive area

for each marker is shown (E-G). Placentas were homogenized and analyzed for concentrations

(pg/ml) of IL-13 (H), GM-CSF (I), and IL-18 (J). Bars represent the mean ± standard error of the

mean [n =5-6/group (image quantification) and 11-14/group (cytokines)] with each symbol

indicating one placenta and, for analysis of images, is the mean quantification of 6 fields of view.

Significant differences ($p < 0.05$) were determined by unpaired two tailed t-test and are

indicated by an asterisk (*). Scale bar: 1 mm (A, left panel/group), 100 μm [(A, right

panels/group; B, left panel/group), 200 μm (C-D, left panel/group), or 50 μm (C-D, right

panel/group).

Figure 4. Placental concentrations of proinflammatory cytokines and immune cells peak

at 6 days-post infection.

Dams at embryonic day 10 were intranasally inoculated with 10^3 TCID$_{50}$ of 2009 H1N1 or were

mock inoculated with media and were euthanized at 3 (E13), 6 (E16), or 8 dpi (E18) to collect

placental tissue (A-C). Placentas were homogenized and analyzed for concentrations (pg/ml) of

proinflammatory cytokines, including IL-1β (A), IL-6 (B), and TNFα (C). Flow cytometry was

performed using live placental cells at 6 dpi to assess immune cell type frequencies in placental

tissue including leukocytes (CD45$^+$, D), NK cells (CD45$^+$, CD3$^-$,CD19$^+$, E), B cells (CD45$^+$, CD3$^-$

,CD19$^+$, F), CD4$^+$ T-cells (CD45$^+$, CD3$^+$, CD4$^+$, G), CD8$^+$ T cells (CD45$^+$, CD3$^+$, CD8a$^+$, H),

neutrophils (CD45$^+$, CD11b$^+$, F4/80$^-$, CD170$^-$, side scatter area high, I), eosinophils (CD45$^+$,

CD11b$^+$, F4/80$^+$, CD170$^+$, J), CD163$^+$ macrophages (CD45$^+$, CD11b$^+$, F4/80$^+$, CD163$^+$, K), and

CD68+ macrophages (CD45$^+$, CD11b$^+$, F4/80$^+$, CD68$^+$, L). Bars represent the mean ± standard

error of the mean from two independent replications (n =11-14/group for cytokines and 5/group

for flow cytometry) with individual mice indicated by symbols. Significant differences ($p < 0.05$)

were determined by two-way ANOVA with Bonferroni post hoc test (A-C) or two tailed unpaired

t-test (D-L) and are indicated by an asterisk (*).

Figure 5. Downregulation of circulating and placental P4 caused by upregulation of COX-

1 through PGF2α secretion. Dams at embryonic day (E) 10 were intranasally inoculated with

10^3 TCID$_{50}$ of 2009 H1N1 or were mock inoculated with media. Dams were euthanized for tissue

and serum collection at 1 (E11), 3 (E13), 6 (E16) or 8 (E18) dpi. Serum (A) and placental (B) P4

concentrations (ng.ml) were measured by ELISA. PGF2α concentrations (ng/ml) were

measured by ELISA in lung homogenates (C), serum (D), and ovarian homogenates (E). COX-1

and COX-2 protein expression in lung tissue was assessed by western blot at 3dpi (F) and

quantified as the fluorescence signal for each protein normalized to GAPDH. Active COX-1 is

also shown as a ratio to inactive COX-1 (F). Lungs from two independent experiments were

analyzed together to minimize interassay variability. Independent cohorts of dams were

intranasally inoculated with 10^3 TCID$_{50}$ of 2009 H1N1 or were mock inoculated with media and

treated twice daily beginning 1 day before infection with either COX-1 inhibitor (SC-560, 10

mg/kg), COX-2 inhibitor (NS-398, 10 mg/kg), or vehicle until dams were euthanized at 3 dpi for

tissue and serum collection. PGF2α concentrations (ng/ml) were measured by ELISA in ovarian

tissue (H) and P4 concentrations (ng/ml) were measured by ELISA in serum (I). Bars represent

the mean ± standard error of the mean from two independent replications (n =6-12/group for

ELISAs and 4/group for western blot) with individual mice indicated by symbols. Significant

differences ($p < 0.05$) were determined by two-way ANOVA with Bonferroni post hoc test (A-E,

H-I) or two tailed unpaired t-test (G) and are indicated by an asterisk (*).

Figure 6. Synthetic progestin treatment improves maternal morbidity and prevents

offspring growth restriction and developmental delays.

Dams at embryonic day (E) 10 were intranasally inoculated with 10^3 TCID$_{50}$ of 2009 H1N1 or

were mock inoculated with media. At the time of inoculation, dams were subcutaneously

administered 17α-hydroxyprogesterone caproate (17-OHPC, 2mg/dam). Dams were euthanized

for tissue collection at 3 (E13) or 6 (E16) days post infection (dpi) or carried their pregnancy

through delivery (E20). Daily rectal temperatures were recorded as a measurement of maternal

morbidity through 14dpi (A). Viral titers were measured in lung tissue via $TCID_{50}$ assay at 3

(E13) and 6 (E16) dpi (B). At postnatal day (PND) 0, pups were weighed, with the average

measurements of each independent litter graphed to account for litter effects (C), and pups

were followed through weaning to observe developmental milestones (D) including fur growth,

ear canal opening, eye opening, and teeth eruption. (E) At PND 5, a subset of pups was sexed

and assessed for their surface righting ability with one pup per sex per dam given 3 trials for the

test, and each pup's best trial reported (G). Individual symbols (A) or bars (B-C, E) represent the

mean ± standard error of the mean from two independent replications (n =10/group) with

individual mice (B, E) or litters (C) indicated by symbols. For developmental milestones, the

center bar indicates the median day a littler reached each milestone, with the range indicated by

the edges of the box (D). Significant differences (p < 0.05) were determined by two-way ANOVA

with Bonferroni post hoc test of AUCs (A), two-way ANOVA with Bonferroni post hoc test (B-C),

Kruskal-Wallis with Dun's post hoc test (D), or three-way ANOVA with Bonferroni post hoc test

(G) and are indicated by an asterisk (*).

Figure 7. Synthetic progestin treatment ameliorates placental damage and inflammation.

Dams at embryonic day (E) 10 were intranasally inoculated with 10^3 $TCID_{50}$ of 2009 H1N1 or

were mock inoculated with media. At the time of inoculation, dams were subcutaneously

administered 17α-hydroxyprogesterone caproate (17-OHPC, 2mg/dam). Placentas were

collected at 3 (E13) or 6 (E16) days post infection (dpi). Representative TUNEL[+] (green) stained

placentas with DAPI to label nuclei taken (blue) at 100x magnification (A), with quantification of

the percentage positive area (B). Placentas were immunostained for cytokeratin (red) to mark

trophoblasts and DAPI (blue) to label nuclei (C) Representative images were taken at 20x

magnification, with quantification of the percentage positive area (D). Placentas were

homogenized to measure concentrations (pg/ml) of IL-1β (E), IL-6 (F), and TNFα (G). Bars

represent the mean ± standard error of the mean (n =5-9/group) with each symbol indicating

one placenta and, for analysis of images, is the mean quantification of 6 fields of view.

Significant differences ($p < 0.05$) were determined by unpaired two tailed t-test and are

indicated by an asterisk (*). Scale bar: 50 µm

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

DAPI TUNEL   DAPI Cytokeratin