Numerical and analytical simulation of the growth of amyloid-β plaques

Andrey V. Kuznetsov

Department of Mechanical and Aerospace Engineering, North Carolina State University,

Raleigh, NC 27695-7910, USA; e-mail: avkuznet@ncsu.edu

Abstract

Numerical and analytical solutions were employed to calculate the radius of an amyloid-β (Aβ) plaque over time. To my knowledge, this study presents the first model simulating the growth of Aβ plaques. Findings indicate that the plaque can attain a diameter of 50 μm after 20 years of growth, provided the degradation machinery responsible for degradation of Aβ monomers is malfunctioning. A mathematical model incorporates nucleation and autocatalytic growth processes using the Finke-Watzky model. The resulting system of ordinary differential equations was solved numerically, and for the simplified case of infinitely long Aβ monomer half-life, an analytical solution was found. Assuming that Aβ aggregates stick together and using the distance between the plaques as an input parameter of the model, it was possible to calculate the plaque radius from the concentration of Aβ aggregates. This led to the "cube root hypothesis," positing that Aβ plaque size increases as the cube root of time. This hypothesis helps explain why larger plaques grow more slowly. Furthermore, the obtained results suggest that the plaque size is independent of the kinetic constants governing Aβ plaque agglomeration. This indicates that the kinetics of Aβ plaque agglomeration is not a limiting factor for plaque growth. Instead, the plaque growth rate is limited by the rates of Aβ monomer production and degradation.

Keywords neuron; Alzheimer's disease; Finke−Watzky model, mathematical modeling; polymerization; cube root hypothesis

Abbreviations

Aβ   amyloid-β

AD   Alzheimer's disease

APP   amyloid precursor protein

CV   control volume

F-W   Finke-Watzky

1. Introduction

Alzheimer's disease (AD) is a devastating neurodegenerative disorder, impacting nearly 50 million individuals globally. While recent FDA-approved therapies may decelerate its progression, there are currently no definitive cures for AD [1-5]. It is crucial to develop mechanistic models of processes involved in AD to pave the way for potential future treatments.

Senile plaques, primarily composed of accumulated amyloid-β (Aβ) peptides [6], represent a key characteristic of AD [7]. A significant surge of interest in Aβ aggregation arose from the amyloid cascade hypothesis, which posited that the formation of Aβ aggregates initiates all other pathological processes in AD. While this hypothesis has undergone extensive investigation, it has also faced recent criticism from some authors [8-11]. Whether senile plaques serve as a causative factor in AD or not remains a subject of ongoing investigation, but they undeniably stand as a distinctive hallmark of the disease [12-15].

Previous mechanistic models of AD [16-18] primarily focused on elucidating intraneuronal mechanisms contributing to the development of AD. However, these investigations did not simulate the gradual extracellular buildup of Aβ plaques. In contrast, the present study aims to replicate the gradual growth of an Aβ plaque, a process that can span decades in humans. This process involves the generation of amyloid precursor protein (APP) within neurons and its subsequent cleavage by β- and γ-secretases at lipid membranes [6,14,19,20]. The majority of these newly formed Aβ monomers are released into the extracellular environment, where they hold the potential to aggregate [1].

2. Materials and models

2.1. Model equations

The F-W model simplifies the intricate process of protein aggregation into two key steps: nucleation and autocatalysis. During nucleation, new aggregates are continuously generated, while in the autocatalysis step, these aggregates undergo rapid surface growth [21,22]. These two pseudo-elementary reaction steps can be summarized as follows:

$$A \xrightarrow{k_1} B \,, \tag{1}$$

$$A + B \xrightarrow{k_2} 2B \,, \tag{2}$$

where A refers to a monomeric protein, and B represents a protein that has undergone amyloid conversion. The kinetic constants, denoted as k_1 and k_2, correspond to the rates of nucleation and autocatalytic growth, respectively [21]. The primary nucleation process, as defined by Eq. (1), exclusively involves

monomers. Conversely, secondary nucleation, described by Eq. (2), involves both monomers and preexisting aggregates of the same peptide [23].

In [21], the Finke-Watzky (F-W) model was utilized to fit previously published data on Aβ aggregation [24,25]. Typically, this data is presented for two isoforms, Aβ40 and Aβ42, with the latter being more prone to aggregation. In this context, the F-W model is applied to simulate the conversion of monomers, whose concentration is denoted as C_A, into aggregates, whose concentration is denoted as C_B. These aggregates encompass various forms of Aβ oligomers, protofibrils, and fibrils [7]. Aβ aggregates formed through this process assemble in amyloid plaques. It is worth noting that the simplified nature of the F-W model does not allow for differentiation between various aggregate types and sizes. The current model does not simulate the process by which adhesive Aβ fibrils assemble into an Aβ plaque. It is assumed that this assembly occurs faster than the formation of Aβ fibrils.

If soluble Aβ were to diffuse at a slower rate than the pace of aggregation, it would lead to growth primarily occurring at the tips of aggregates, resulting in a branching, tree-like structure. However, this contradicts what has been observed in experiments [26]. The rapid diffusion of Aβ monomers suggests minimal variation in their concentration, C_A, between neurons. With this assumption, the control volume (CV) displayed in Fig. 1 with a volume L^3 can be treated as a lumped capacitance body. Expressing the conservation of Aβ monomers within this CV yields the following equation:

$$\frac{dC_A}{dt} = -k_1 C_A - k_2 C_A C_B - C_A \frac{\ln(2)}{T_{1/2,A}} + \frac{q_{A,0}}{L} . \tag{3}$$

The first term on the right-hand side of Eq. (3) simulates the rate of conversion of Aβ monomers into aggregates, while the second term simulates the rate of this conversion through autocatalytic growth. These expressions for the first and second terms arise from applying the law of mass action to the reactions described by Eqs. (1) and (2), respectively. The third term on the right-hand side of Eq. (3) accounts for the rate at which Aβ monomers degrade. The fourth term represents the rate of Aβ monomer production, as Aβ monomers are continuously generated through the cleavage of APP [27,28].

If the conservation of Aβ aggregates is written for the CV, the following equation is obtained:

$$\frac{dC_B}{dt} = k_1 C_A + k_2 C_A C_B . \tag{4}$$

The first term on the right-hand side of Eq. (4) represents the rate at which Aβ aggregates are produced through nucleation, while the second term simulates the rate of their production via autocatalytic growth.

These terms have equal magnitudes, but opposite signs when compared to the first and second terms in Eq. (3). This is because, in the F-W model, the rate of aggregate production equals the rate of monomer disappearance. Eqs. (3) and (4) only account for the finite half-life of Aβ monomers, as it is expected that the half-life of Aβ aggregates is significantly longer [29].

The initial conditions are as follows:

At $t=0$: $C_A=0$, $C_B=0$. (5a,b)

The sole independent variable in the model is time, t. The dependent variables are listed in Table 1, while the parameters employed in the model are listed in Table 2.

Table 1. Dependent variables utilized in the model.

Symbol	Definition	Units
C_A	Concentration of Aβ monomers	μM
C_B	Concentration of Aβ aggregates	μM
r_{ABP}	Radius of a growing Aβ plaque	μm

Table 2. Model parameters and their estimated values.

Symbol	Definition	Units	Value or range	Reference or estimation method	Value(s) used in computations
a_{21}	Conversion factor from $\frac{mol}{\mu m^3}$ to μM	$\frac{\mu M\ \mu m^3}{mol}$	10^{21}		10^{21}
D_{ABP}	Diameter of Aβ plaque	μm	16.0-23.9 [a]	[30]	
k_1	Rate constant for the first pseudo-elementary step of the F-W model,	s⁻¹	2.22×10^{-9} [b]	[21,24]	2.22×10^{-9}

	describing the nucleation of Aβ aggregates				
k_2	Rate constant for the second pseudo-elementary step of the F-W model, describing the autocatalytic growth of Aβ aggregates	μM^{-1} s^{-1}	9.44×10^{-6} [b]	[21,24]	9.44×10^{-6}
L	Half distance between senile plaques	μm	50	Estimated from Fig. 1a in [31]	50
MW	Molecular weight of an Aβ monomer (the sum of weights of every atom in the monomer)	g mol^{-1}	4.51×10^3 [c]	[32]	4.51×10^3
N_A	Avogadro's number	mol^{-1}	6.022×10^{23}		6.022×10^{23}
$q_{A,0}$	Rate of production of Aβ monomers per unit area of one of the CV's faces. Aβ production is a surface phenomenon, occurring at the lipid membrane. The model assumes that all Aβ is released from one face of the CV. Aβ production is	μM μm s^{-1}	1.1×10^{-4} [d]	[32]	1.1×10^{-4}

	calculated per unit area of that face. The rate of Aβ production per unit volume then is $q_{A,0}/L$.				
$T_{1/2,A}$	Half-life of Aβ monomers	s	4.61×10^{3} [e]	[33]	4.61×10^{3}, 4.61×10^{9} (to represent $T_{1/2,A}\to\infty$)
ρ	Density of the Aβ plaque	g/μm^3	1.35×10^{-12} [f]	[34]	1.35×10^{-12}

[a] The average plaque area visible in 2D images is between 400-450 μm², with the smallest plaques having areas of less than 200 μm² [30]. While this parameter is not directly employed in the model, it is included in the table for discussing model predictions concerning plaque radius.

[b] Values reported in [21] were obtained by curve-fitting data from [24]. $k_1 = 8\times10^{-6}$ h^{-1}, $k_2 = 3.4\times10^{-2}$ μM^{-1} h^{-1} [21].

[c] Molecular weight of Aβ42 [32].

[d] According to [32], the human brain produces Aβ42 monomers at a rate of 0.33×10^{-6} μmol s^{-1}. Considering the approximate volume of the brain as 1.5×10^{15} μm^3, the rate of Aβ42 production per unit volume is estimated to be 2.2×10^{-22} μmol μm^{-3} s^{-1}. In a small cube with a side length of L (representing the CV in Fig. 1), the rate of Aβ42 production is 2.75×10^{-17} μmol s^{-1}. The production of Aβ monomers occurs at the lipid membrane [35]. To account for this, the production rate of Aβ42 was normalized assuming that production only occurs on one face of the CV with an area of L^2. The production rate of Aβ42 per unit area on that face was determined by dividing 2.75×10^{-17} μmol s^{-1} by L^2, resulting in 1.1×10^{-20} μmol μm^{-2} s^{-1} $=1.1\times10^{-5}$ μM μm s^{-1}. Since Aβ40 is significantly more abundant than Aβ42 in the brain [36], the total production rate of Aβ isoforms per unit area on that face is estimated to be ten times greater than the rate of Aβ42 production alone. This yields an estimated value of 1.1×10^{-4} μM μm

s[-1] for $q_{A,0}$. This results in an estimated value of $q_{A,0}$ as 1.1×10^{-4} µM µm s[-1]. The rate of Aβ42

production per unit volume of the CV can then be calculated as $q_{A,0}/L$.

[e] According to [33], the value is 1.28 hours.

[f] A commonly used density value, which is independent of the protein type, is 1.35 g/cm³ [34].

The dimensionless form of Eqs. (3) and (4) can be expressed as follows:

$$\frac{dC_A^*}{dt^*} = -C_A^* - C_A^* C_B^* - C_A^* \frac{\ln(2)}{T_{1/2,A}^*} + q_{A,0}^*, \tag{6}$$

$$\frac{dC_B^*}{dt^*} = C_A^* + C_A^* C_B^*. \tag{7}$$

The dimensionless form of the initial conditions defined by Eq. (5) is as follows:

At $t^* = 0$: $C_A^* = 0$, $C_B^* = 0$. $\tag{8a,b}$

Table 3 presents the dimensionless independent variable, while Table 4 summarizes the dimensionless dependent variables. Table 5 provides the dimensionless parameters used in the model.

Eqs. (6) and (7) constitute a pair of ordinary differential equations. MATLAB's ODE45 solver (MATLAB R2020b, MathWorks, Natick, MA, USA) was employed to solve these equations subject to initial conditions given by Eq. (8). To ensure accuracy of the solution, the error tolerance parameters, RelTol and AbsTol, were set to 1e-10.

Table 3. Dimensionless independent variable in the model.

Symbol	Definition
t^*	$t k_1$

Table 4. Dimensionless dependent variables employed in the model.

Symbol	Definition

C_A^*	$C_A k_2 / k_1$
C_B^*	$C_B k_2 / k_1$
r_{ABP}^*	r_{ABP} / L

Table 5. Dimensionless parameters utilized in the model.

Symbol	Definition	Value(s) obtained using the dimensional parameter values listed in the last column of Table 2
$q_{A,0}^*$	$q_{A,0} k_2 / \left(k_1^2 L \right)$	4.21×10^6
$T_{1/2,A}^*$	$T_{1/2,A} k_1$	1.02×10^{-5}, 1.02×10^1 (to represent $T_{1/2,A}^* \to \infty$)
ρ^*	$\dfrac{\rho}{MW} a_{21} \dfrac{k_2}{k_1}$	1.27×10^9

When Eqs. (6) and (7) were added together, the following result was obtained:

$$\frac{dC_A^*}{dt^*} + \frac{dC_B^*}{dt^*} = -C_A^* \frac{\ln(2)}{T_{1/2,A}^*} + q_{A,0}^* . \tag{9}$$

By integrating Eq. (9) with respect to time under the condition of $T_{1/2,A} \to \infty$ and using the initial condition given by Eq. (8), the following result was obtained:

$$C_A^* + C_B^* = q_{A,0}^* t^* . \tag{10}$$

The increase in $C_A^* + C_B^*$ over time is attributed to the continuous generation of Aβ monomers.

By eliminating C_A^* from Eq. (7) using Eq. (10), the following was obtained:

$$\frac{dC_B^*}{dt^*} = q_{A,0}^* t^* - C_B^* + \left(q_{A,0}^* t^* - C_B^* \right) C_B^* . \tag{11}$$

Eq. (11) is similar to the one examined in [37]. To find the exact solution of Eq. (11) with the initial condition given by Eq. (8b), the DSolve function followed by the FullSimplify function in Mathematica 13.3 (Wolfram Research, Champaign, IL) is utilized. The resulting solution is presented below:

$$C_B^*\left(t^*\right) = -1 + \frac{\exp\left[t^* + q_{A,0}^* \frac{\left(t^*\right)^2}{2}\right]\sqrt{q_{A,0}^*}}{\sqrt{q_{A,0}^*} - \sqrt{2}F\left(\frac{1}{\sqrt{2q_{A,0}^*}}\right) + \sqrt{2}\exp\left[t^* + q_{A,0}^* \frac{\left(t^*\right)^2}{2}\right]F\left(\frac{1 + q_{A,0}^* t^*}{\sqrt{2q_{A,0}^*}}\right)}. \tag{12}$$

In this equation, $F(x)$ represents Dawson's integral:

$$F(x) = \frac{1}{2}\sqrt{\pi}\, e^{-x^2} \mathrm{erfi}(x), \tag{13}$$

where $\mathrm{erfi}(x)$ is the imaginary error function.

C_A^* can then be determined using Eq. (10) as:

$$C_A^* = q_{A,0}^* t^* - C_B^*. \tag{14}$$

The exact solution given by Eqs. (12) and (14) is quite complex. A more elegant approximate solution, which is also valid when $T_{1/2,A} \to \infty$, is as follows:

$$C_A^* = \frac{q_{A,0}^*}{1 + q_{A,0}^* t^*}, \tag{15}$$

$$C_B^* = q_{A,0}^* t^*. \tag{16}$$

As $t^* \to \infty$, Eq. (15) predicts that $C_A^* \to 0$. It is important to note that Eqs. (15) and (16) are useful for larger times. However, they are not applicable when $t^* \to 0$; for instance, Eq. (15) yields $C_A^*(0) = q_{A,0}^*$, which contradicts the initial condition specified in Eq. (8a).

The Aβ plaque's growth (Fig. 1) is simulated using the following approach. The total number of Aβ monomers, denoted as N, that are integrated into the Aβ plaque at time t, is determined using the following equation [38]:

$$N(t) = \frac{C_B(t)}{a_{21}} L^3 N_A. \tag{17}$$

Alternatively, $N(t)$ can be calculated as described in [38]:

$$N(t) = \frac{V_{ABP}(t)\rho}{MW}N_A,$$

(18)

where MW denotes the average molecular weight of an Aβ monomer.

By equating the expressions on the right-hand sides of Eqs. (17) and (18) and solving for the volume of an Aβ plaque, the following result is obtained:

$$V_{ABP}(t) = \frac{C_B(t)}{a_{21}}L^3\frac{MW}{\rho},$$

(19)

where $a_{21} = 10^{21}\frac{\mu M \, \mu m^3}{mol}$ denotes the conversion factor from $\frac{mol}{\mu m^3}$ to μM.

Assuming that the Aβ plaque is spherical in shape,

$$V_{ABP}(t) = \frac{4}{3}\pi r_{ABP}^3(t).$$

(20)

Setting the right-hand sides of Eqs. (19) and (20) equal and solving for the radius of the Aβ plaque, the following is obtained:

$$r_{ABP} = \left(\frac{3}{4\pi}\frac{C_B}{a_{21}}L^3\frac{MW}{\rho}\right)^{1/3}.$$

(21)

Expressed in dimensionless variables, Eq. (21) becomes:

$$r_{ABP}^* = \left(\frac{3}{4\pi}\frac{1}{\rho^*}\right)^{1/3}\left(C_B^*\right)^{1/3}.$$

(22)

Substituting Eq. (16) into Eq. (22) yields the following approximate solution for the dimensionless radius of the Aβ plaque:

$$r_{ABP}^* = \left(\frac{3}{4\pi}\frac{q_{A,0}^*}{\rho^*}\right)^{1/3}\left(t^*\right)^{1/3}.$$

(23)

Returning to the dimensional variables, the following equation is obtained:

$$r_{ABP} = \left(\frac{3}{4\pi}q_{A,0}\frac{MW}{\rho a_{21}}L^2\right)^{1/3}t^{1/3}.$$

(24)

Eq. (24) suggests that the size of Aβ plaques increases proportionally to the cube root of time. Intriguingly, Eq. (24) also predicts that the radius of an Aβ plaque remains independent of the kinetic constants that describe the rate of Aβ plaque agglomeration, k_1 and k_2. This observation aligns with findings from [39], which revealed that individual plaques can form relatively quickly, typically within approximately 24 hours. Hence, the kinetics of Aβ plaque agglomeration does not appear to be a limiting factor. Instead, the growth of these plaques is constrained by the rate of Aβ monomer production and, if the half-life of the monomers is finite, by the rate at which these monomers degrade.

Based on Eq. (24), one can deduce that the rate of Aβ plaque growth is

$$\frac{dr_{ABP}}{dt} = \frac{1}{6^{2/3}\pi^{1/3}}\left(\frac{L^2 MW q_{A,0}}{a_{21}\rho}\right)^{1/3} t^{-2/3}.$$
(25)

Eq. (25) offers an explanation for the deceleration of Aβ plaque growth over time. This implies that smaller (younger) plaques tend to undergo faster growth compared to larger plaques, a phenomenon observed in previous research [40].

2.2. Sensitivity analysis

The sensitivity of the radius of a growing Aβ plaque, r_{ABP}, to the two parameters in Eq. (24), L and $q_{A,0}$, as well as to $T_{1/2,A}$ was investigated. To do this, the local sensitivity coefficients [41-44] were computed. The sensitivity coefficients represent the first-order partial derivatives of the observable (in this case, the radius of the growing Aβ plaque) to L, $q_{A,0}$, and $T_{1/2,A}$, for example, $\frac{\partial r_{ABP}}{\partial L}$. The dimensionless relative sensitivity coefficients [42,45] were then determined using the following equation, for example:

$$S_L^{r_{ABP}} = \frac{L}{r_{ABP}}\frac{\partial r_{ABP}}{\partial L}.$$
(26)

Utilizing Eq. (24):

$$S_L^{r_{ABP}} = \frac{2}{3},$$
(27)

$$S_{q_{A,0}}^{r_{ABP}} = \frac{1}{3}.$$
(28)

The positive values of $S_L^{r_{ABP}}$ and $S_{q_{A,0}}^{r_{ABP}}$ indicate that an increase in L and $q_{A,0}$ will result in an increase in r_{ABP}.

Because Eq. (24) is applicable only for the infinite half-life of Aβ monomers, the sensitivity of the radius of a growing Aβ plaque, r_{ABP}, to $T_{1/2,A}$ was computed numerically:

$$\frac{\partial r_{ABP}}{\partial T_{1/2,A}} \approx \frac{r_{ABP}\left(T_{1/2,A} + \Delta T_{1/2,A}\right) - r_{ABP}\left(T_{1/2,A}\right)}{\Delta T_{1/2,A}}\Bigg|_{\text{other parmeters kept constant}}, \tag{29}$$

where $\Delta T_{1/2,A} = 10^{-3} T_{1/2,A}$ represents the step size. Computations were conducted with various step sizes to assess whether the sensitivity coefficients remained unaffected by changes in the step size.

3. Results

Figs. 2-6 were generated using the parameter values listed in Table 2, unless otherwise specified.

In the scenario where Aβ monomers have an infinite half-life, the concentration of Aβ aggregates, C_B, increases linearly with time, consistent with the approximate solution provided by Eq. (16) (Fig. 2a). However, it is worth noting that this approximate solution is not valid at small times within the range of [0 0.04 year] (Fig. 2b). When the machinery responsible for degrading Aβ monomers works properly ($T_{1/2,A} = 4.61\times10^3$ s), the computational results show an approximately 2-year delay before the linear increase in Aβ aggregate concentration begins (Fig. 2a). This suggests that functional proteolytic machinery, which degrades Aβ monomers, can postpone the formation of Aβ plaques by several years [46].

In the scenario where Aβ monomers have an infinite half-life, the concentration of Aβ monomers, C_A, decreases as time progresses (Fig. 3a,b). This decline is a consequence of the increasing concentration of Aβ aggregates (Fig. 2a), leading to faster autocatalytic conversion of monomers into aggregates as described by Eq. (4). The approximate solution given by Eq. (15) closely matches the numerical solution (Fig. 3a), but only for times greater than 0.04 years (Fig. 3b). In the scenario where the machinery responsible for degrading Aβ monomers functions normally ($T_{1/2,A} = 4.61\times10^3$ s), there is approximately a 2-year delay in the decrease of Aβ monomer concentration. This delay is due to the postponed production of Aβ aggregates (as shown in Fig. 2a). These aggregates play a role in the autocatalytic conversion of monomers into aggregates (refer to Eq. (3)), which accelerates the monomer decay.

In the scenario of infinite half-life for Aβ monomers, the radius of a growing Aβ plaque, r_{ABP}, increases in proportion to the cube root of time, $t^{1/3}$, in agreement with the approximate solution provided in Eq. (24) (Fig. 4a). The agreement between the approximate and numerical solutions is excellent, except for when t is within the smaller time range of [0 0.04 year] (Fig. 4b). However, if the machinery responsible for Aβ monomer degradation functions normally ($T_{1/2,A} = 4.61 \times 10^3$ s), the computational result predicts a slower increase rate for the radius of an Aβ plaque over time (Fig. 4a). This suggests that the cube root hypothesis is applicable only when at least 0.04 years have passed since the beginning of Aβ aggregation, and the Aβ monomer degradation mechanism is not functioning correctly.

Fig. 5a illustrates the relationship between the radius of Aβ plaques after 20 years of growth and the half-distance between these plaques, L. When $T_{1/2,A} \to \infty$, r_{ABP} increases with L almost linearly. This occurs because, for a uniform volumetric production rate of Aβ monomers, a smaller number of plaques (larger distances between them) allows each individual plaque to access a greater supply of Aβ, thus enabling it to grow to a larger size. If $L = 500$ μm, the plaque's diameter after 20 years of growth is approximately 50 μm, which agrees with the representative diameter of an Aβ plaque observed in [31]. In the case of infinite half-life for Aβ monomers, the approximate solution provided by Eq. (24) precisely matches the numerical solution (Fig. 5a). In the scenario where Aβ monomer degradation functions normally ($T_{1/2,A} = 4.61 \times 10^3$ s), an increase in L first results in an increase in r_{ABP}, and subsequently (after reaching approximately $L = 250$ μm), it leads to a decrease in r_{ABP} (as shown in Fig. 5a). This occurs because an increase in L leads to a proportional increase in the supply of Aβ monomers into the CV through one of its faces, scaling with L^2. The number of Aβ monomers degraded within the CV scales with L^3, signifying that, at a certain point, Aβ monomer degradation becomes the prevailing process within the CV.

Fig. 5b illustrates how the radius of Aβ plaques after 20 years of growth is influenced by the rate of Aβ monomer production. When $T_{1/2,A} \to \infty$, r_{ABP} is directly proportional to $q_{A,0}^{1/3}$ because a higher production rate of Aβ monomers results in a greater supply of Aβ available for plaque growth (Fig. 5b). In the scenario where Aβ monomers have an infinite half-life, the approximate solution given by Eq. (24) precisely agrees with the numerical solution (Fig. 5b). However, if the machinery responsible for Aβ monomer degradation is functioning normally ($T_{1/2,A} = 4.61 \times 10^3$ s), the plaque grows nearly at the same rate as it does for $T_{1/2,A} \to \infty$. This is because Aβ monomer production is continuous, and the monomers convert into Aβ plaques before they can be broken down.

Fig. 5c illustrates the relation between the Aβ plaque radius after 20 years of growth and the half-life of Aβ monomers. The numerical solution generated a sigmoidal curve. Initially, with a half-life of 10 s, the plaque radius is zero. It gradually increases as the half-life increases, at $T_{1/2,A} = 10^5$ s reaching a value corresponding to the infinite half-life of monomers. Beyond this point, it aligns with the plaque radius predicted by the approximate solution given by Eq. (24).

Fig. 6 illustrates the dimensionless sensitivities of the Aβ plaque radius after 20 years of growth to various model parameters. Positive sensitivity indicates that an increase in the parameter leads to an increase in the plaque radius, while negative sensitivity signifies that an increase in the parameter results in a decrease in the plaque radius.

When the Aβ monomer degradation machinery is functioning normally ($T_{1/2,A} = 4.61 \times 10^3$ s), the dimensionless sensitivity of the Aβ plaque's radius to L decreases over time, turning negative after 0.5 years of growth. It reaches its minimum value after approximately 2 years of growth, then becomes positive again after 4 years of growth (Fig. 6a). In the scenario of infinite monomer half-life, the sensitivity equals 2/3, consistent with Eq. (27).

The dimensionless sensitivity of the Aβ plaque's radius to $q_{A,0}$ is positive for $T_{1/2,A} = 4.61 \times 10^3$ s (Fig. 6b). When Aβ monomer half-life is infinitely large, the sensitivity equals 1/3, consistent with Eq. (28).

At $T_{1/2,A} = 4.61 \times 10^3$ s, the dimensionless sensitivity to $T_{1/2,A}$ is positive, whereas for the infinite half-life of Aβ monomers, the dimensionless sensitivity to $T_{1/2,A}$ is zero (Fig. 6c).

4. Discussion, model limitations, and future directions

The analytical solution predicting the radius of Aβ plaques over time suggests that these plaques increase in size following a cube root relationship with time. This result helps clarify why larger plaques exhibit slower growth. Moreover, the model indicates that the radius of Aβ plaques remains unaffected by variations in the kinetic constants governing Aβ plaques aggregation rate, k_1 and k_2. This observation implies that the kinetics of Aβ aggregation is not the primary limiting factor dictating Aβ plaque growth; instead, the rates of Aβ monomer production and degradation play more crucial roles.

There are several limitations associated with this model. One limitation is the use of the lumped capacitance model to simulate the buildup of Aβ monomer and aggregate concentrations. This model assumes that these concentrations vary with time but not with location within the CV. Given that Aβ

monomers are generated at lipid membranes, it is reasonable to expect concentration variations between neurons, with the highest concentration occurring at the lipid membrane. Future models should consider incorporating diffusion-driven transport of Aβ monomers between neurons, as explored in a preliminary manner in [47]. Another limitation arises from the utilization of the F-W model for Aβ aggregation, which does not distinguish between aggregates of varying sizes, such as dimers, oligomers, protofibrils, and fibrils [6]. Addressing this limitation in future research by employing more precise protein polymerization models will lead to more accurate characterization of Aβ plaque structure, particularly their heterogeneity.

Data accessibility. This article has no additional data.

Authors' contributions. AVK is the sole author of this paper.

Competing interests. The author declares no competing interests.

Funding statement. AVK acknowledges the support provided by the National Science Foundation (grant CBET-2042834) and the Alexander von Humboldt Foundation through the Humboldt Research Award.

References

[1] Rahman, M. M., and Lendel, C., 2021, "Extracellular Protein Components of Amyloid Plaques and their Roles in Alzheimer's Disease Pathology," Molecular Neurodegeneration, **16**(1), pp. 59.

[2] Cummings, J., Zhou, Y., Lee, G., 2023, "Alzheimer's Disease Drug Development Pipeline: 2023," Alzheimer's & Dementia: Translational Research & Clinical Interventions, **9**(2), pp. e12385.

[3] Hardy, J., 2006, "A Hundred Years of Alzheimer's Disease Research," Neuron, **52**(1), pp. 3-13.

[4] Maqbool, M., Mobashir, M., and Hoda, N., 2016, "Pivotal Role of Glycogen Synthase Kinase-3: A Therapeutic Target for Alzheimer's Disease," European Journal of Medicinal Chemistry, **107**, pp. 63-81.

[5] Hung, S., and Fu, W., 2017, "Drug Candidates in Clinical Trials for Alzheimer's Disease," Journal of Biomedical Science, **24**, pp. 47.

[6] Hampel, H., Hardy, J., Blennow, K., 2021, "The Amyloid-B Pathway in Alzheimer's Disease," Molecular Psychiatry, **26**(10), pp. 5481-5503.

[7] Chen, G., Xu, T., Yan, Y., 2017, "Amyloid Beta: Structure, Biology and Structure-Based Therapeutic Development," Acta Pharmacologica Sinica, **38**(9), pp. 1205-1235.

[8] Hardy, J. A., and Higgins, G. A., 1992, "Alzheimers-Disease - the Amyloid Cascade Hypothesis," Science, **256**(5054), pp. 184-185.

[9] Karran, E., Mercken, M., and De Strooper, B., 2011, "The Amyloid Cascade Hypothesis for Alzheimer's Disease: An Appraisal for the Development of Therapeutics," Nature Reviews Drug Discovery, **10**(9), pp. 698-712.

[10] Herrup, K., 2015, "The Case for Rejecting the Amyloid Cascade Hypothesis," Nature Neuroscience, **18**(6), pp. 794-799.

[11] Ricciarelli, R., and Fedele, E., 2017, "The Amyloid Cascade Hypothesis in Alzheimer's Disease: It's Time to Change our Mind," Current Neuropharmacology, **15**(6), pp. 926-935.

[12] Murphy, M. P., and LeVine, H. 3., 2010, "Alzheimer's Disease and the Amyloid-Beta Peptide," Journal of Alzheimer's Disease : JAD, **19**(1), pp. 311-323.

[13] Hardy, J., and Selkoe, D. J., 2002, "The Amyloid Hypothesis of Alzheimer's Disease: Progress and Problems on the Road to Therapeutics," Science (New York, N.Y.), **297**(5580), pp. 353-356.

[14] O'Brien, R. J., and Wong, P. C., 2011, "Amyloid Precursor Protein Processing and Alzheimer's Disease," Annual Review of Neuroscience, Vol 34, **34**, pp. 185-204.

[15] Selkoe, D. J., and Hardy, J., 2016, "The Amyloid Hypothesis of Alzheimer's Disease at 25years," EMBO Molecular Medicine, **8**(6), pp. 595-608.

[16] Kuznetsov, I. A., and Kuznetsov, A. V., 2018, "How the Formation of Amyloid Plaques and Neurofibrillary Tangles may be Related – A Mathematical Modelling Study," Proceedings of the Royal Society A, **474**, pp. 20170777.

[17] Kuznetsov, I. A., and Kuznetsov, A. V., 2018, "Simulating the Effect of Formation of Amyloid Plaques on Aggregation of Tau Protein," Proceedings of the Royal Society A-Mathematical Physical and Engineering Sciences, **474**(2220), pp. 20180511.

[18] Torok, J., Maia, P. D., Verma, P., 2021, "Emergence of Directional Bias in Tau Deposition from Axonal Transport Dynamics," PLOS Computational Biology, **17**(7), pp. e1009258.

[19] Chow, V. W., Mattson, M. P., Wong, P. C., 2010, "An Overview of APP Processing Enzymes and Products," Neuromolecular Medicine, **12**(1), pp. 1-12.

[20] MacLeod, R., Hillert, E., Cameron, R. T., 2015, "The Role and Therapeutic Targeting of A-, B- and Γ-Secretase in Alzheimer's Disease," Future Science OA, **1**(3), pp. FSO11.

[21] Morris, A. M., Watzky, M. A., Agar, J. N., 2008, "Fitting Neurological Protein Aggregation Kinetic Data Via a 2-Step, Minimal/"Ockham's Razor" Model: The Finke-Watzky Mechanism of Nucleation Followed by Autocatalytic Surface Growth," Biochemistry, **47**(8), pp. 2413-2427.

[22] Iashchishyn, I. A., Sulskis, D., Nguyen Ngoc, M., 2017, "Finke-Watzky Two-Step Nucleation-Autocatalysis Model of S100A9 Amyloid Formation: Protein Misfolding as "Nucleation" Event," ACS Chemical Neuroscience, **8**, pp. 2152-2158.

[23] Thacker, D., Barghouth, M., Bless, M., 2023, "Direct Observation of Secondary Nucleation Along the Fibril Surface of the Amyloid B 42 Peptide," Proceedings of the National Academy of Sciences, **120**(25), pp. e2220664120.

[24] Bieschke, J., Zhang, Q., Powers, E. T., 2005, "Oxidative Metabolites Accelerate Alzheimer's Amyloidogenesis by a Two-Step Mechanism, Eliminating the Requirement for Nucleation," Biochemistry, **44**(13), pp. 4977-4983.

[25] Vestergaard, M., Kerman, K., Saito, M., 2005, "A Rapid Label-Free Electrochemical Detection and Kinetic Study of Alzheimer's Amyloid Beta Aggregation," Journal of the American Chemical Society, **127**(34), pp. 11892-11893.

[26] Cruz, L., Urbanc, B., Buldyrev, S. V., 1997, "Aggregation and Disaggregation of Senile Plaques in Alzheimer Disease," Proceedings of the National Academy of Sciences of the United States of America, **94**(14), pp. 7612-7616.

[27] Haass, C., Kaether, C., Thinakaran, G., 2012, "Trafficking and Proteolytic Processing of APP," Cold Spring Harbor Perspectives in Medicine, **2**(5), pp. a006270.

[28] Masters, C. L., and Selkoe, D. J., 2012, "Biochemistry of Amyloid B-Protein and Amyloid Deposits in Alzheimer Disease," Cold Spring Harbor Perspectives in Medicine, **2**(6), pp. a006262.

[29] Madrasi, K., Das, R., Mohmmadabdul, H., 2021, "Systematic in Silico Analysis of Clinically Tested Drugs for Reducing Amyloid-Beta Plaque Accumulation in Alzheimer's Disease," Alzheimer's & Dementia, **17**(9), pp. 1487-1498.

[30] Tandan, A. G., 2022, "A Brief Note on Amyloid Plaque," Global Journal of Neurology and Neurosurgery, **10**(1), pp. 3-4.

[31] Querol-Vilaseca, M., Colom-Cadena, M., Pegueroles, J., 2019, "Nanoscale Structure of Amyloid-B Plaques in Alzheimer's Disease," Scientific Reports, **9**(1), pp. 5181.

[32] Raskatov, J. A., 2019, "What is the "Relevant" Amyloid $\beta 42$ Concentration?" ChemBioChem, **20**(13), pp. 1725-1726.

[33] Waters, J., 2010, "The Concentration of Soluble Extracellular Amyloid-Beta Protein in Acute Brain Slices from CRND8 Mice," PLOS One, **5**(12), pp. e15709.

[34] Cavicchi, R. E., King, J., and Ripple, D. C., 2018, "Measurement of Average Aggregate Density by Sedimentation and Brownian Motion Analysis," Journal of Pharmaceutical Sciences, **107**(5), pp. 1304-1312.

[35] Dawkins, E., Derks, R. J. E., Schifferer, M., 2023, "Membrane Lipid Remodeling Modulates Γ-Secretase Processivity," Journal of Biological Chemistry, **299**(4).

[36] Gu, L., and Guo, Z., 2021, "Alzheimer's $A\beta 42$ and $A\beta 40$ Form Mixed Oligomers with Direct Molecular Interactions," Biochemical and Biophysical Research Communications, **534**, pp. 292-296.

[37] Kuznetsov, I. A., and Kuznetsov, A. V., 2022, "An Analytical Solution Simulating Growth of Lewy Bodies," Mathematical Medicine and Biology : A Journal of the IMA, **39**(3), pp. 299-312.

[38] Watzky, M. A., Finney, E. E., and Finke, R. G., 2008, "Transition-Metal Nanocluster Size Vs Formation Time and the Catalytically Effective Nucleus Number: A Mechanism-Based Treatment," Journal of the American Chemical Society, **130**(36), pp. 11959-11969.

[39] Meyer-Luehmann, M., Spires-Jones, T. L., Prada, C., 2008, "Rapid Appearance and Local Toxicity of Amyloid-Beta Plaques in a Mouse Model of Alzheimer's Disease," Nature, **451**, pp. 720-724.

[40] Yan, P., Bero, A. W., Cirrito, J. R., 2009, "Characterizing the Appearance and Growth of Amyloid Plaques in APP/PS1 Mice," The Journal of Neuroscience : The Official Journal of the Society for Neuroscience, **29**(34), pp. 10706-10714.

[41] Beck, J. V., and Arnold, K. J., 1977, *Parameter Estimation in Science and Engineering,* Wiley, New York.

[42] Zadeh, K. S., and Montas, H. J., 2010, "A Class of Exact Solutions for Biomacromolecule Diffusion-Reaction in Live Cells," Journal of Theoretical Biology, **264**(3), pp. 914-933.

[43] Zi, Z., 2011, "Sensitivity Analysis Approaches Applied to Systems Biology Models," IET Systems Biology, **5**(6), pp. 336-346.

[44] Kuznetsov, I. A., and Kuznetsov, A. V., 2019, "Investigating Sensitivity Coefficients Characterizing the Response of a Model of Tau Protein Transport in an Axon to Model Parameters," Computer Methods in Biomechanics and Biomedical Engineering, **22**(1), pp. 71-83.

[45] Kacser, H., Burns, J. A., and Fell, D. A., 1995, "The Control of Flux," Biochemical Society Transactions, **23**(2), pp. 341-366.

[46] Saido, T., and Leissring, M. A., 2012, "Proteolytic Degradation of Amyloid B-Protein," Cold Spring Harbor Perspectives in Medicine, **2**(6), pp. a006379.

[47] Kuznetsov, A. V., 2023, "Effect of Diffusivity of Amyloid Beta Monomers on the Formation of Senile Plaques," bioRxiv, https://doi.org/10.1101/2023.07.31.551367.

Figure captions

Fig. 1. A cubic CV with a side length of L containing a single growing Aβ plaque.

Fig. 2. (a) Molar concentration of Aβ aggregates, C_B, over time (in years). (b) Similar to Fig. 2a but focusing on a specific time range of [0 0.2 year] on the x-axis. The numerical solution is obtained by solving Eqs. (3) and (4) numerically with the initial condition (5). The approximate solution is plotted using Eq. (16) and converting the dimensionless variables into dimensional ones. The approximate solution is valid only for $T_{1/2,A} \rightarrow \infty$ and is plotted exclusively for this case. L=50 μm and $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1}.

Fig. 3. (a) Molar concentration of Aβ monomers, C_A, over time (in years). (b) Similar to Fig. 3a but focusing on a specific time range of [0 0.2 year] on the x-axis. The numerical solution is obtained by numerically solving Eqs. (3) and (4) with the initial condition (5). The approximate solution is plotted using Eqs. (15) and converting the dimensionless variables into dimensional ones. The approximate solution is valid only for $T_{1/2,A} \rightarrow \infty$ and is plotted exclusively for this case. L=50 μm and $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1}.

Fig. 4. Radius of a growing Aβ plaque, r_{ABP}, over time (in years). (b) Similar to Fig. 4a but focusing on a specific time range of [0 0.2 year] on the x-axis. The numerical solution is obtained by numerically solving Eqs. (3) and (4) with the initial condition (5) and then using Eq. (21) to calculate the radius of the Aβ plaque. The dimensionless variables are then converted into dimensional ones. The approximate solution is plotted using Eq. (24). The approximate solution is valid only for $T_{1/2,A} \rightarrow \infty$ and is plotted exclusively for this case. L=50 μm and $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1}.

Fig. 5. Radius of a growing Aβ plaque after 20 years of growth vs (a) half distance between senile plaques, $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1}; (b) rate of production of Aβ monomers, L=50 μm; and (c) half-life of Aβ monomers, L=50 μm and $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1}.

Fig. 6. Dimensionless sensitivity of the Aβ plaque's radius, r_{ABP}, to half distance between senile plaques, L, $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1} (a); rate of production of Aβ monomers, $q_{A,0}$, L=50 μm (b); and half-life of Aβ monomers, $T_{1/2,A}$, $q_{A,0} = 1.1 \times 10^{-4}$ μM μm s^{-1} and L=50 μm (c).

Figure 1

Figure 2a,b

Figure 3a,b

Figure 4a,b

Figure 5a,b

(c)

Figure 5c

(a)

(b)

Figure 6a,b

Figure 6c