Temperature, phenology, and plant defenses predict fitness near colder range limit

Daniel N. Anstett[1-6]

1 Plant Resilience Institute, Michigan State University, East Lansing, MI 48824 USA

2 Department of Plant Biology, Michigan State University, East Lansing, MI 48824 USA

3 Department of Entomology, Michigan State University, East Lansing, MI 48824 USA

4 Program in Ecology, Evolution, and Behavior, Michigan State University, East Lansing, MI

48824 USA

5 Biodiversity Research Centre and Department of Botany, University of British Columbia,

Vancouver, British Columbia V6T 1Z4, Canada

6 anstett3@msu.edu

Abstract

The space for time substitution posits that warmer locations can provide a source of genetic

variation that could be adaptive for future climate change conditions. While this approximation

might be useful for planning assisted gene flow, it relies on the importance of abiotic adaptations

over biotic ones. Here I address this gap by assessing influence of anti-herbivore defenses,

phenology, and morphology on the seed production of 149 populations of *Oenothera biennis*

close to the plant's cold range limit. Genotypes from 2.1° South of the common garden produce

more seeds than most northern lineages. Adaptations across space are a suitable substitute for

climate change, but there is still substantial fitness variability. These differences were best

explained by bolt date, flowering time, and greater defenses against herbivores. Given the

impacts of climate change, plant defenses might already be of similar adaptive importance to

phenology close to northern rage limits.

Key Words

Local adaptation, climate gradient, seed production, flowering time, herbivory, ellagitannins,

Oenothera bennis.

Introduction

Climate change is exposing populations to climate extremes that are increasingly further from historical averages (IPCC 2014; Pörtner *et al.* 2022). These deviations from baseline climatic conditions are overwhelming abiotic and possibly biotic adaptations (Etterson 2004; Anderson *et al.* 2012; Franks *et al.* 2014) leading to mortality (Van Mantgem & Stephenson 2007), population decline, (Willis *et al.* 2008) and range contractions (Freeman *et al.* 2018). Many populations that were locally adapted to their abiotic environment prior to the onset of climate change (Leimu & Fischer 2008; Sanford & Kelly 2011) are becoming maladapted (Aitken *et al.* 2008; Wang *et al.* 2010; Wilczek *et al.* 2014; Anderson & Wadgymar 2020; Bontrager *et al.* 2020). These populations may have to undergo range shifts to more suitable conditions to survive (Parmesan & Yohe 2003; Parmesan 2006; Poloczanska *et al.* 2013; Fadrique *et al.* 2018), or rapidly evolve to reverse demographic decline (Bell & Gonzalez 2011; Bell 2017). Such rapid evolution can occur from standing genetic variation (Barrett & Schluter 2008), yet extreme climate change may require adaptations beyond locally available genetic variation.

The space-for-time substitution is a useful approximation in which spatial biodiversity patterns across abiotic gradients are used to predict future temporal changes (Pickett 1989; Blois *et al.* 2013a). Space-for-time has been used widely to study community composition (Walker *et al.* 2010; Blois *et al.* 2013a), phenology (Buyantuyev *et al.* 2012), range-shifts (Eskildsen *et al.* 2013), and landscape-level climate maladaptation (Fitzpatrick & Keller 2015; Exposito-Alonso *et al.* 2018; Dauphin *et al.* 2021). When invoking space-for-time in the context of climate change adaptation, missing adaptive genetic variation from higher latitudes are assumed to be present in warmer populations, such that space can substitute for adaptations needed across time (Wogan & Wang 2018). Indeed, climate change experiments suggest that genetic variation from warmer

localities can perform better in historically colder locations (Anderson & Wadgymar 2020;

Bontrager *et al.* 2020). However, space-for-time fails to consider non-analogue future climates

(Williams & Jackson 2007), unchanging landscape features such as photoperiod (Walker *et al.*

2019), and increasingly strong biotic interactions as higher latitude sites become warmer (Blois *et*

al. 2013b; HilleRisLambers *et al.* 2013).

How the space-for-time substitution is influenced by differences in biotic interactions

between populations remains largely untested (Bucharova 2017). Plant populations may be

particularly vulnerable to increases in biotic interactions such as herbivory, which have been

documented to increase at higher latitudes due to greater herbivore biomass and increased intensity

of herbivore outbreaks (de Sassi & Tylianakis 2012; DeLucia *et al.* 2012; Ju *et al.* 2015; Jactel *et*

al. 2019). Plants have evolved chemical and physical defenses to protect against herbivory (Ehrlich

& Raven 1964; Agrawal & Weber 2015) as well as herbivore avoidance strategies such as changes

in phenology (Carmona *et al.* 2011). However, biogeographic theory predicts that populations at

higher latitudes will have lower plant defenses (Dobzhansky 1950; MacArthur 1972; Schemske *et*

al. 2009; Baskett & Schemske 2018) (but see Moles *et al.* 2011; Anstett *et al.* 2016). This could

leave higher latitude populations poorly defended and vulnerable to more mobile insect herbivores

and may make plant defenses adaptive during climatic change.

Nevertheless, assuming genetic variation from lower latitudes will be effective in

mediating herbivory at higher latitude sites ignores the complex and variable nature of plant-

herbivore interactions. Latitudinal patterns in herbivory are notoriously variable across species

(Moles *et al.* 2011; Anstett *et al.* 2016), meaning that some systems might naturally have higher

herbivory in colder climates. Herbivore identity and importance can also vary latitudinally with

differences in herbivory varying across herbivore guild and specialization (Hiura & Nakamura

2013; Anstett *et al.* 2014). In these conditions, plant defenses that protect against one herbivore

species might be less useful on another (Krischik *et al.* 1991; Kursar *et al.* 2009; Ali & Agrawal

2012) or could even act as a feeding stimulant (Malcolm & Brower 1989; Holeski *et al.* 2013;

Singh *et al.* 2021). Thus, genetic variation from donor sites could introduce plant defense traits

which could be unhelpful, or even attract greater herbivore damage. Similar arguments could be

made about other biotic interactions, making it vital to test the space-for-time substitution using

common gardens that measure biotic interactions and traits that mediate them.

A common garden can establish what historical climate best predicts performance, and

what biotic and abiotic adaptations may underly this increased performance. Multiple common

garden experiments have been carried out in tree species, testing the performance of range-wide

genetic variation across the range of the species including close to colder range limits (Schreiber

et al. 2013; Aitken & Bemmels 2016; Risk *et al.* 2021). Yet less focus has been placed on testing

the performance of range-wide genetic variation in forbs (but see Anderson & Wadgymar 2020)

and comparing what traits provide the biotic and abiotic adaptations that cause performance

differences. Given their shorter generation time, forbs might be more able to rapidly evolve and

establish, yet their smaller size might make them more vulnerable to larger herbivore loads.

To address these gaps, I investigate the performance of genotypes from across the range of

Oenothera biennis in a northern common garden, extending prior work on genetically-based clines

in anti-herbivore defenses and plant phenology (Anstett *et al.* 2015). Specifically, I ask: (1) Does

geography and environment of origin of populations from across the range of *O. biennis* predict

seed production when grown close to the northern range limit? (2) Does increased herbivory

predict decreased plant performance? (3) What phenological, morphological, and defensive traits

predict performance? This approach establishes that plant defenses are as important as abiotic

adaptations in predicting plant performance at higher latitudes and in explaining the traits

underlying space-for-time substitution.

Methods

Study system

Common evening primrose, *Oenothera biennis* L., Onagraceae, is a facultatively biennial,

semelparous species found along disturbed habitats such as roadsides and along bodies of water

(Dietrich *et al.* 1997). This herbaceous forb is found throughout Eastern North America with

limited populations in Western North America (Fig. 1), spanning a subtropical to temperate climate

gradient. Due to permanent translocation heterozygosity (PTH), *O. biennis* is functionally asexual

(Cleland 1972) and has near 100% selfing (Godfrey 2014; Johnson and Godfrey in review). Thus

populations are usually dominated by a clonal single genotype, and produce genetically identical

seeds (Levin 1975). This mating system in conjunction with a semelparous life history make it

possible to measure total lifetime male and female fitness in terms of the number of seeds produced

by each plant.

Oenothera biennis is an interesting case study for considering biotic adaptation during

climate change. It has substantial heritable genetic variation in morphological, physiological, life

history and herbivore resistance traits (Johnson *et al.* 2009) and established latitudinal gradients

across many of these traits (Anstett *et al.* 2015). A diverse community of herbivores attacks *O.*

biennis, including chewing herbivores, xylem feeders, and multiple specialized seed predators

(Johnson & Agrawal 2005). Attacks by seed predators target the plant's direct link to seed

production and fitness and have been shown to drive rapid evolution (Agrawal *et al.* 2012). These

seed predators are known to impact plants more at higher latitudes, while leaf chewing herbivory

shows no association with latitude (Anstett *et al.* 2014). Phenolics and specifically the ellagitannin

compound oenothein A have been associated with decreased seed predation and increased seed

production (Johnson *et al.* 2009; Agrawal *et al.* 2012). Total phenolics from *O. biennis* and a

mixture of oenothein A and structurally similar polymers decrease survival of both generalist and

specialist insect herbivores of *O. biennis* (Anstett *et al.* 2019). Ellagitannins such as oenothein A

are highly reactive in alkaline conditions (Salminen & Karonen 2011), such as those found in

lepidopteran insect guts (Appel 1993), making this compound valuable to anti-herbivore defense.

Experimental design

Seeds from 149 populations of *O. biennis,* including 137 from the main range and 12 from Western

North America (Fig. 1), were sourced from established seed collections provided by Marc T.J.

Johnson. These seed lines were collected one to three decades before the start of this experiment

and kept in dry conditions in a -20 freezer until germination. Five replicates per population were

germinated and grown for three weeks, and then planted in a common garden at the Koffler

Scientific Reserve (New Market, Ontario, Canada) and grown during 2012 and 2013 as outlined

in Anstett *et al.* (2015). Because of the PTH genetic system, these replicates are a single clone

from each of the 149 populations. I previously characterized latitudinal gradients in physical and

chemical defense traits, resistance to local herbivores and phenological differences across 137

populations in the main range of *O. biennis*. Here I assess how geography and climate of origin of

O. biennis impacts seed production, and what traits explain performance when measured as total

lifetime fitness. The timing of the common garden captures the early effects of climate change in

North America with mean annual temperature being 1.6 °C warmer at the common garden across

the 2012 and 2013 period when compared to the historical baseline temperatures under which *O.*

biennis evolved.

Climate & geography

Historical climate data from 1960 to 1990 was downloaded from Climate NA for each population

(Wang *et al.* 2016), which is representative of pre-climate change conditions. I selected seven

temperature and four precipitation associated variables that could impact *O. biennis* fitness. Most

variables were highly correlated ($r > 0.7$; Table S1), hence only mean annual temperature (MAT),

mean summer precipitation (MSP), cumulative moisture deficit (CMD), and relative humidity

(RH) were included in downstream calculations. Monthly climate data was downloaded from

Climate NA for 2012 and 2013 for the Koffler Science Reserve (44.026479, -79.544686). MAT,

CMD, and RH was calculated as a combined metric for July 2012 to September 2013, by first

averaging July, August and September values across both years and then calculating a 12 month

mean. This calculation reflects the entire length of the common garden experiment. MSP for the

common garden was only included for 2013 since supplemental watering was provided in 2012

after seedling transplant. Climatic distance to the common garden was calculated for each climatic

variable by subtracting the common garden value from the 30-year mean.

Trait measurement

I measured 23 traits including phenological (flowering date, bolt date, growth rate); morphological

(leaf toughness, trichome number, specific leaf area (SLA); % water content), secondary chemistry

across leaf, flower and fruit tissues (total phenolics, oxidative capacity; concentration of

ellagitannin compounds oenothein A and oenothein B); and herbivore resistance traits (generalist

leaf herbivory, number of *Philaenus spumarius* L. Hemiptera: Aphrophoridae, fruits damaged by

specialist *Schinia florida* G. Lepidoptera: Noctuidae, fruits damaged by specialist *Mompha*

brevivitella C. Lepidoptera: Momphidae). The detailed methodology for the estimation of 22 of

these traits is found in Anstett *et al.* (2015). Additionally, I monitored bolt date throughout the

season, defined as the first day the stem length was greater than 5 cm. The total number of fruits

and the length and width of 5 fruits produced by each plant was recorded. Seed number was

estimated by the following formula: seeds number = 2.33*Average_Fruit_Length *

Average_Fruit_Width - 102.47 * Damaged_Fruits. This calculation has been established to

accurately estimate fruit number in prior work conducted on *O. biennis* (Agrawal *et al.* 2012). The

damaging effects of specialist seed predators were also taken into account in the final estimation

of seed production by subtracting the total number of *S. florida* and 0.18*M. brevivitella* damaged

fruits. The latter represents average known amount of seed loss per pod due to *M. brevivitella*

damage (M.T.J. Johnson personal communication).

Statistical analyses

All analyses were performed using R (R Development Core Team 2020). Means of each

population were used for all traits ensuring a comparable design across all traits, since dried,

ground plant issue was averaged across genotypes prior to chemical analysis. I predicted seed

number using geographic, climatic, herbivory and plant traits. Seed number had inflated zeros and

was overdispersed (P <0.001) using *dispertiontest* in the *AER* package in R (Kleiber *et al.* 2020).

Hence, all regressions predicting seed number were performed using a negative binomial

generalized linear model in the *MASS* package in R (Ripley *et al.* 2013). For each model, I tested

first order patterns, and quadratic patterns (if this pattern appeared possible in the raw data) (Table

S2). Quadratic predictors were assessed separately from first order predictors using the I(Trait^2)

function in R. Regression models were reduced by using stepwise AIC, selecting the model with

the lowest AIC score. Significance of this model was assessed using a type 2 anova using the

Anova command in the CAR package in R (Fox *et al.* 2012). Results were visualized using visreg

in the *visreg* package in R (Breheny *et al.* 2020), with scale= "response", which applies an inverse

log-link function scaling the plots in seed number units.

Seed number was modeled as a function of latitude and longitude, distance to common

garden, climate present at each population, and climate distance to the common garden (Table S2).

Then to determine what types of traits underly the geographical and climatic patterns, resistance

to herbivores, phenology, morphology, and plant secondary chemistry were each used to predict

seed number. I used multiple regression within each class of traits whenever possible (see Table

S2 for every tested model). Leaf toughness was highly correlated with other morphological

variables (r>0.7) and was not included in the morphology multiple regression. For resistance to

herbivores, I ran separate regressions for seed predators. For secondary compounds, I ran

regressions separately for each tissue. Total phenolics was assessed separately from individual

compounds. Oxidative capacity was highly correlated with total phenolics (r>0.7); oenothein B

was highly correlated with oenothein A. Hence, I selected only total phenolics and oenothein A

based on prior knowledge of what traits more strongly predict resistance to herbivores (Agrawal

et al. 2012; Anstett *et al.* 2015).

To understand the major axes of variation across these predictive variables, I also carried

out principal component analysis using *prcomp()* in R and associated the most important axes with

seed production. Biplots were visualized using *autoplot()* in the ggfortify package in R (Tang *et*

al. 2016). To further establish which variables are most important in predicting *O. biennis* seed

number, I compared the relative importance of all variables that had evidence of predictive seed

number (flowering date, bolt date, water content, growth rate, and number of trichomes as well as

leaf, flower and fruit total phenolics and oenothein A). I carried out this comparison with the

machine learning algorithm, *randomForest* in R (Liaw & Wiener 2002) using 1000 trees and

plotted relative variable importance. I further visualized the relationship between latitude, seed

number and six of the most important traits by plotting trait change over latitude, while color

coding points for seed number using ggplot2 in R (Wickham 2011).

All analyses were repeated using undamaged fruits as a response instead of the estimated

seed number. No major differences were detected in trends and only minor difference in

significance (data not shown); hence I present only the seed number results.

Results

Geography & climate

Seed production increased with latitude, peaking at 41.9° N, 2.1° south of the common garden

before declining (Fig. 2A, Table 1; linear P = 0.001, quadratic P = 0.002). Indeed, five of the top

six genotypes with the most fitness were from 41.9°N or further south. Models with longitude had

higher AIC and were not selected. Seed production decreased at distances further from the

common garden (P = 0.044; Fig. 2B), although the highest performing genotypes were from 100's

or 1000's of km from the common garden.

Temperature and relative humidity explained plant fitness. Seed production increased

towards genotypes from warmer temperatures, peaking at 9.1°C, 1.2°C warmer than the common

garden before declining at higher temperatures of origin (Fig. 2C; linear P = 0.004, quadratic P =

0.001). Genotypes from locations with greater relative humidity had higher seed production up to

62.4% relative humidity, 6.6% higher than the common garden, before declining at higher relative

humidity (Fig. 2D). However, this effect was only weakly supported (linear $P = 0.063$, quadratic

$P = 0.062$). Models with May to September precipitation and Hargreaves climatic moisture deficit

had higher AIC and were not selected. Temperature distance from common garden has a similar

effect with genotypes from locations with greater temperature distance peaking 1.4°C warmer from

the common garden before declining (Fig. S1, Table S1, $P < 0.001$). There was also weak support

for mean summer precipitation distance predicting seed number ($P = 0.083$; Table S1). Models

with relative humidity distance and Hargreaves climatic moisture deficit distance were not

selected.

Resistance to herbivory

Herbivory had variable relationships with seed production (Table 1). Seed production increased

with *Philaenus spumarius* number of to 14 individuals and then declined (Fig. 3A; linear $P = 0.004$,

quadratic $P = 0.063$). The model that included leaf herbivory had higher AIC and was not selected.

Seed predators had contrasting relationships with seed production. Seed number increased with

higher *S. florida* damage or *S. florida* simply attacked more fecund plants (Fig. 3B; $P = 0.004$).

Seed number decreased with greater *M. brevivitella* damage although there was a lot of variability

($P = 0.051$; Fig. 3C).

Plant phenotype

All phenology variables predicted *O. biennis* seed number in some way (Table 1). Plants with early

or late flowering date performed poorly, with peak flowering date occurring at an intermediate

flowering date (Fig. 3D; $P < 0.001$). Genotypes with later bolt date had low seed production, while

plants with earlier bolt date showed higher seed production (on average) but with higher variability

(Fig. 3E; P < 0.001). Plants with intermediate growth rate had higher seed production although

this relationship was weak (linear P = 0.05, quadratic P = 0.073; Fig. 4F). Most plant morphology

variables also predicted seed number (Table 1). Seed production increased with trichome number

up to 110 per 5 mm^2, before decreasing at higher trichome numbers (Fig. 3G; P < 0.001). Fitness

also peaked at intermediate water content (linear P = 0.029; P = 0.03), with seeds increasing with

water content up to 74.3% before decreasing (Fig. 3H). Finally, seed number was not associated

with SLA (Fig. 3I; P = 0.12).

Plant defense chemistry metrics also predicted seed production but only for some tissues

(Table 2). Higher leaf total phenolics weakly predicted greater seed production (Fig. S2A; P =

0.84), while leaf total oenothein A does not predict seed number (linear P = 0.44; quadratic P =

0.76; Fig. S2B). Higher flower total phenolics predicted greater seed number up to 55 mg/g and

then declined (Fig. 4A; linear P = 0.12, quadratic P =0.03). Seed number increased with flower

oenothein A (Fig. 4C; linear P = 0.01) and declined slightly at only very high values (P = 0.01).

Greater fruit total phenolics predicted greater seed production (Fig. 4B; P = 0.03) with poor

evidence of decline at high fruit total phenolics (P = 0.11). Seed number also increases with greater

fruit oenothein A (P = 0.002; Fig. 4D).

Principle component analyses revealed that most of *O. biennis* trait variation was predicted

by the first two principal components (Fig. S3; Table S3), with PC1 explaining 33.65% of the

variation and PC2 explaining 13.8%. PC1 is primarily associated with flowering date, fruit

oenothein A, fruit phenolics, flower oenothein A, leaf oenothein A, leaf phenolics, and bolt date,

while PC2 is primarily associated with growth rate, SLA, and water content (Fig. 5A, Table S4).

Smaller values of PCA 1, associated with later phenology and higher plant defenses predicted

greater seed number (P=0.02). Smaller values of PC2, associated with decreased SLA, growth,

rate, and water content predicted greater seed number (P=0.004), but with much variation and with

the highest seed producers having intermediate seed numbers.

Gradient forest analysis showed that, out of the variables predictive of seed number, bolt

date, flower oenothein A, fruit oenothein A, fruit total phenolics, and flowering date were the five

most important (Fig. 6). After this there is a larger gap in importance, with growth rate, flower

total phenolics, trichome number and water content being the least important. Latitude-trait plots

with color-coded seed production emphasized flowering time just south of the common garden

leads to increased seed production (Fig. S4A), while later bolt date from further south leads to

massively decreased seed production (Fig. S4B). Populations from directly south of the common

garden that have greater fruit defenses perform considerably better (Fig. S5). Flower defenses

south of the common garden are also somewhat predictive of greater seed production (Fig. S6).

Discussion

This study shows that anti-herbivore defenses can at least partially explain the space-for-time

substitution that genotypes from warmer latitudes are more adapted to climate change conditions.

A 1.6 degree increase from historical baseline temperatures has made *O. biennis* genotypes from

2.1° South of the common garden perform better than local more northern genotypes. This

performance difference is best explained by both phenology and plant phenolics, suggesting both

biotic and abiotic factors impact performance during the early effects of climate change. The

importance of phenology as a climate change adaptation is already well established (Hamann *et*

al. 2018; Wadgymar *et al.* 2018). Here I extend our understanding of climate adaptation by

implicating the benefit of increased plant chemical traits typically associated with plant defense

as an important part of climate change adaptation.

Climate change adaptation close to cooler range limit

Plant defenses can be an important adaptation at higher latitude sites and could even be more

important than abiotic adaptations. Here I show that genotypes from across the range of *O.*

biennis with greater fruit total phenolics, greater fruit oenothein A and intermediate values of

flower oenothein A (mostly higher than the local population) had greater seed production when

grown close to the cold range limit of *O. biennis* (Fig. 4-5, Fig. S5-6). Additionally, *O. biennis*

chemical traits ranked higher in variable importance than flowering time (Fig. 6), which is

usually more commonly associated with the impacts of climate change. This importance of plant

defenses at higher latitudes in *O. biennis* is supported by prior work showing that greater

oenothein A explains decreased seed predator damage in this common garden (Fig. S4A; Anstett

et al. 2015) and during experimental evolution (Agrawal *et al.* 2012). Seed predator damage,

especially by *M. brevivitella*, is also known to lead to decreased seed production (Fig. 3C;

Agrawal *et al.* 2012), pointing to the importance of greater plant defenses at higher latitudes in

this system.

320   Flowering time and bolt date were also both predictive of greater seed production,

with earlier bolt date and an intermediate date of first flower being optimal. Some plants failed to

bolt until much later in the season, thus producing many fewer seeds (Fig. S3B) and making bolt

date the most important predictor of success for this small subset of populations (Fig. 6).

Oenothera biennis populations from the southern part of the range of the plant showed a

suboptimal, much later flowering time when grown in the northern common garden (Anstett *et*

al. 2015) likely due to photoperiod and/or temperature cue issues, which are often raised when

considering phenotypic mismatch during latitudinal translocations (Wadgymar & Weis 2017).

Indeed, these plants would have less time to acquire resources and produce seeds before the end

of the growing season (Inouye 2008; Franks 2015). Thus, plants from higher latitudes that are

just South of the common garden were better able to produce more seeds because they likely had

a photoperiod closer to the garden.

This begs the question as to why an intermediate flowering date leads to greater seed

production in *O. biennis* rather than the earliest possible flowering date. One possibility is that

these early flowering plants developed too early and cooler weather damaged the plants. Another

explanation is that plants that flower earlier are available for longer in the season and are thus

more susceptible to herbivory. Indeed, prior work in this system suggests that genotypes with

earlier flowering have greater damage by generalist and specialist herbivores (Agrawal *et al.*

2013; Anstett *et al.* 2015). This pattern is not surprising given that life history traits have been

found to be consistent predictors of resistance to herbivores (Pilson 2000; Kawagoe & Kudoh

2010), and at times more reliable than traditional chemical or physical defenses (Carmona *et al.*

2011). Overall, this suggests that even phenological adaptations at northern range limits may be

adaptations to biotic interactions.

Implications for space-time substitution

Using space as a proxy for time during translocations takes into account the effects of

temperature adaptation on average, but ignores how genetics within a region may impact the

realized fitness of a randomly selected individual. On average, genotypes from south of the

common garden with slightly higher mean annual temperature perform better (Fig. 2A,C), while

other climate traits were less important. This matches previous findings pointing to temperature

as a key factor in field transplant experiments (Bontrager *et al.* 2020). However, average

temperature or latitude may hide considerable genetic variability within a particular temperature

band. Multiple genotypes from latitudes at or slightly above the common garden still had above

average seed number (Fig. 2A), while some genotypes from across the entire range had low

fitness. This variability points to the importance of testing on-the-ground fitness at future

introduction sites or at least considering the variety of abiotic and biotic adaptations that might

be needed and where this genetic variation might be sourced.

This system suggests how possible trade-offs between biotic and abiotic adaptation may

be an important consideration for assisted gene flow. The benefits of assisted gene flow as a way

to rescue populations from climate change has frequently been weighed against risks of

introducing maladapted genetic variation (Vitt *et al.* 2010; Aitken & Whitlock 2013). Space-for-

time substitutions might serve as a way to source genetic material for assisted gene flow while

mitigating the risks of introducing further maladaptation in an already vulnerable species. This

study suggests there are risks of introducing genotypes that may be well suited for the new

abiotic environment but not the biotic one (or vice versa). These trade-offs are possible in *O.*

biennis especially given that multiple traits explain fitness and that these types of traits are

known to covary (Johnson *et al.* 2009). Given these findings, studies focusing on species of

immediate conservation concern should investigate the possible greater importance of biotic

adaptations close to northern range limits and how these may interact with abiotic adaptation

(Blois *et al.* 2013b; HilleRisLambers *et al.* 2013). This may be particularly true for translocations

across latitudinal gradients (Fielding *et al.* 1999), where adaptation to photoperiod may lead to

poor match with phenology (Walker *et al.* 2019).

Implications for biogeographic hypotheses

Greater plant defenses at higher latitudes may provide needed adaptations to increased herbivory.

Herbivore outbreaks at higher latitude sites are being attributed directly or indirectly to climate

change (de Sassi & Tylianakis 2012; DeLucia *et al.* 2012; Ju *et al.* 2015; Jactel *et al.* 2019),

raising the possibility that selection for greater defenses at higher latitudes might already be

occurring. Indeed, I previously documented that damage by seed predators is greater at higher

latitudes in *O. biennis* (Anstett *et al.* 2014), possibly due to early effects of climate change. This

idea falls against the established pre-climate change paradigm that greater biotic interactions at

warmer, lower latitudes leads to selection for greater traits that mediate these interactions

(Dobzhansky 1950; MacArthur 1972; Coley & Barone 1996; Schemske *et al.* 2009). However,

selection for greater plant defenses at higher latitudes due to climate change might explain why

more studies are finding greater plant defenses at higher latitudes (Moles *et al.* 2011; Anstett *et*

al. 2016).

As the impacts of the Anthropocene become more acute, long-held biogeographic

hypotheses may require a more complex and updated view. The causes and drivers of range

limits has been a key question throughout the history of evolutionary biology as well as a driver

of biogeographic patterns (Darwin 1859; MacArthur 1972; Sexton *et al.* 2009). Biotic

interactions are viewed as being less important for determining colder range limits (Louthan *et*

al. 2015; Paquette & Hargreaves 2021). Biotic interactions and the traits that mediate them are

also expected to be less intense and important at higher latitude sites (MacArthur 1972; Coley &

Aide 1991; Coley & Barone 1996; Schemske *et al.* 2009; Johnson & Rasmann 2011; Anstett *et*

al. 2016). Both of these ideas share the common viewpoint of diminishing the importance of

biotic interactions in colder climates. Climate change is changing this reality and is likely going

to pose new biotic challenges on historically cooler sites that may lack adaptations for greater

mean interaction intensity (de Sassi & Tylianakis 2012; Ju *et al.* 2015) and greater variability in

that intensity (Wetzel *et al.* 2023). Greater biotic adaptations such as plant defenses against

herbivores may become an increasingly important part of plant adaptation at higher latitudes

leading to the weakening of latitudinal gradients in biotic interactions and the traits that mediate

them. Lack of these adaptations at northern range limits could even slow plant range expansions

that are so often evoked when studying climate change.

Data Availability

All code and data for this paper is available on GitHub https://github.com/anstettd/KSR_North.

Conflict of Interest

The author has no conflicts of interest.

Acknowledgements

I thank C. Fitzpatrick, M. Turcotte, N. Turley, J. Hollister, C. Thomsen, R. Godfrey, A. Lee, P.

Maharaj, L. Sequeira, T. Didiano, D. Filice, D. Vigmond, H. Sekhon, A. Longley, J. Anstett A.

Koivuniemi, and J. Kim for field and lab support. J.-P. Salminen and J. Ahern aided in analytical

chemistry. M. Johnson and S. Greiner provided seeds. M. Johnson, W. Wetzel, and A.

VanWallendael also provided advice and read over a draft of the paper. J. Anstett aided in

coding. Special thanks to A. Agrawal for advice on seed estimation in *O. biennis*. Funding was

provided by Sigma Xi Grant-in-Aid of Research Grant, NSERC CGS-D Vanier, NSERC PDF,

NSERC Banting and a MSU Plant Resilience Institute Postdoc Fellowship to D. Anstett.

References

Agrawal, A.A., Hastings, A.P., Johnson, M.T.J., Maron, J.L. & Salminen, J.-P. (2012). Insect

herbivores drive real-time ecological and evolutionary change in plant populations. Science 338:

113-116.

Agrawal, A.A., Johnson, M.T.J., Hastings, A.P. & Maron, J.L. (2013). A field experiment

demonstrating plant life-history evolution and its eco-evolutionary feedback to seed predator

populations. Am. Nat. 181: S1-S11.

Agrawal, A.A. & Weber, M.G. (2015). On the study of plant defence and herbivory using

comparative approaches: how important are secondary plant compounds. Ecol. Lett. 18: 985-

991.

Aitken, S.N. & Bemmels, J.B. (2016). Time to get moving: assisted gene flow of forest trees.

Evol. Appl. 9: 271-290.

Aitken, S.N. & Whitlock, M.C. (2013). Assisted gene flow to facilitate local adaptation to

climate change. Annu. Rev. Ecol. S. 44: 367-388.

Aitken, S.N., Yeaman, S., Holliday, J.A., Wang, T. & Curtis-McLane, S. (2008). Adaptation,

migration or extirpation: climate change outcomes for tree populations. Evol. Appl. 1: 95-111.

Ali, J.G. & Agrawal, A.A. (2012). Specialist versus generalist insect herbivores and plant

defense. Trends Plant Sci. 17: 293-302.

Anderson, J.T., Inouye, D.W., McKinney, A.M., Colautti, R.I. & Mitchell-Olds, T. (2012).

Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in

response to climate change. Proc. R. Soc. London, Ser. B 279: 3843-3852.

Anderson, J.T. & Wadgymar, S.M. (2020). Climate change disrupts local adaptation and favours

upslope migration. Ecol. Lett. 23: 181-192.

Anstett, D.N., Ahern, J.R., Glinos, J., Nawar, N., Salminen, J.P. & Johnson, M.T.J. (2015). Can

genetically based clines in plant defence explain greater herbivory at higher latitudes? Ecol. Lett.

18: 1376–1386.

Anstett, D.N., Cheval, I., D'Souza, C., Salminen, J.P. & Johnson, M.T.J. (2019). Ellagitannins

from the Onagraceae Decrease the Performance of Generalist and Specialist Herbivores. J. Chem

Ecol. 45: 86-94.

Anstett, D.N., Naujokaitis-Lewis, I. & Johnson, M.T.J. (2014). Latitudinal gradients in herbivory

on Oenothera biennis vary according to herbivore guild and specialization. Ecology 95: 2915-

2923.

Anstett, D.N., Nunes, K.A., Baskett, C. & Kotanen, P.M. (2016). Sources of controversy

surrounding latitudinal patterns in herbivory and defense. Trends Ecol. Evol. 31: 789–802.

Appel, H.M. (1993). Phenolics in ecological interactions: the importance of oxidation. J. Chem

Ecol. 19: 1521-1552.

Barrett, R.D. & Schluter, D. (2008). Adaptation from standing genetic variation. Trends Ecol.

Evol. 23: 38-44.

Baskett, C.A. & Schemske, D.W. (2018). Latitudinal patterns of herbivore pressure in a

temperate herb support the biotic interactions hypothesis. Ecol. Lett. 21: 578-587.

Bell, G. (2017). Evolutionary rescue. Annu. Rev. Ecol. S. 48: 605-627.

Bell, G. & Gonzalez, A. (2011). Adaptation and evolutionary rescue in metapopulations

experiencing environmental deterioration. Science 332: 1327-1330.

Blois, J.L., Williams, J.W., Fitzpatrick, M.C., Jackson, S.T. & Ferrier, S. (2013a). Space can

substitute for time in predicting climate-change effects on biodiversity. PNAS. 110: 9374-9379.

Blois, J.L., Zarnetske, P.L., Fitzpatrick, M.C. & Finnegan, S. (2013b). Climate change and the

past, present, and future of biotic interactions. Science 341: 499-504.

Bontrager, M., Muir, C., Mahony, C., Gamble, D., Germain, R., Hargreaves, A. et al. (2020).

Climate warming weakens local adaptation. bioRxiv

doi: https://doi.org/10.1101/2020.11.01.364349.

Breheny, P., Burchett, W. & Breheny, M.P. (2020). Package 'visreg'.

Bucharova, A. (2017). Assisted migration within species range ignores biotic interactions and

lacks evidence. Restor. Ecol. 25: 14-18.

Buyantuyev, A., Xu, P., Wu, J., Piao, S. & Wang, D. (2012). A space-for-time (SFT) substitution

approach to studying historical phenological changes in urban environment. PLoS One 7:

e51260.

Carmona, D., Lajeunesse, M.J. & Johnson, M.T.J. (2011). Plant traits that predict resistance to

herbivores. Funct. Ecol. 25: 358-367.

Cleland, R.E. (1972). Oenothera: cytogenetics and evolution. Academic Press, New York, New

York, USA.

Coley, P.D. & Aide, T.M. (1991). Comparison of herbivory and plant defenses in temperate and

tropical broad-leaved forests. In: Plant-Animal Interaction: Evolutionary Ecology in Tropical and

Temperate Regions (eds. Price, PW, Lewinsohn, TM, Fernandes, GW & Benson, WW). Wiley

New York, pp. 25-49.

Coley, P.D. & Barone, J.A. (1996). Herbivory and plant defenses in tropical forests. Annu. Rev.

Ecol. Syst. 27: 305-335.

Darwin, C. (1859). On the origin of species by means of natural selection. J. Murray.

Dauphin, B., Rellstab, C., Schmid, M., Zoller, S., Karger, D.N., Brodbeck, S. et al. (2021).

Genomic vulnerability to rapid climate warming in a tree species with a long generation time.

Global Change Biol. 27: 1181-1195.

de Sassi, C. & Tylianakis, J.M. (2012). Climate change disproportionately increases herbivore

over plant or parasitoid biomass. PLoS One 7: e40557.

DeLucia, E.H., Nabity, P.D., Zavala, J.A. & Berenbaum, M.R. (2012). Climate change: resetting

plant-insect interactions. Plant Physiol.160: 1677-1685.

Dietrich, W., Wagner, W.L. & Raven, P.H. (1997). Systematics of Oenothera section Oenothera

subsection Oenothera (Onagraceae). Sys. Bot. Mono. 50: 1-234.

Dobzhansky, T. (1950). Evolution in the tropics. Am. Sci. 38: 209-221.

Ehrlich, P.R. & Raven, P.H. (1964). Butterflies and plants: a study in coevolution. Evolution 18:

586-608.

Eskildsen, A., le Roux, P.C., Heikkinen, R.K., Høye, T.T., Kissling, W.D., Pöyry, J. et al.

(2013). Testing species distribution models across space and time: high latitude butterflies and

recent warming. Glob. Ecol. Biogeogr. 22: 1293-1303.

Etterson, J.R. (2004). Evolutionary potential of Chamaecrista fasciculata in relation to climate

change. I. Clinal patterns of selection along an environmental gradient in the Great Plains.

Evolution 58: 1446-1456.

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H.A. & Weigel, D. (2018).

Genomic basis and evolutionary potential for extreme drought adaptation in Arabidopsis

thaliana. Nat. Ecol. Evol. 2: 352-358.

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J. et al. (2018). Widespread

but heterogeneous responses of Andean forests to climate change. Nature 564: 207-212.

Fielding, C., Whittaker, J., Butterfield, J. & Coulson, J. (1999). Predicting responses to climate

change: the effect of altitude and latitude on the phenology of the Spittlebug Neophilaenus

lineatus. Funct. Ecol. 13: 65-73.

Fitzpatrick, M.C. & Keller, S.R. (2015). Ecological genomics meets community-level modelling

of biodiversity: mapping the genomic landscape of current and future environmental adaptation.

Ecol. Lett 18: 1-16.

Fox, J., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S. et al. (2012). Package

'car'. Vienna: R Foundation for Statistical Computing 16.

Franks, S.J. (2015). The unique and multifaceted importance of the timing of flowering. Bot.

Soc. Am. 102: 1401-1402.

Franks, S.J., Weber, J.J. & Aitken, S.N. (2014). Evolutionary and plastic responses to climate

change in terrestrial plant populations. Evol. Appl 7: 123-139.

Freeman, B.G., Scholer, M.N., Ruiz-Gutierrez, V. & Fitzpatrick, J.W. (2018). Climate change

causes upslope shifts and mountaintop extirpations in a tropical bird community. PNAS 115:

11982-11987.

Godfrey, R. (2014). The Effects of Losing Sex on Genetic Variation in Oenothera (Onagraceae).

University of Toronto.

Hamann, E., Weis, A.E. & Franks, S.J. (2018). Two decades of evolutionary changes in Brassica

rapa in response to fluctuations in precipitation and severe drought. Evolution 72: 2682-2696.

HilleRisLambers, J., Harsch, M.A., Ettinger, A.K., Ford, K.R. & Theobald, E.J. (2013). How

will biotic interactions influence climate change–induced range shifts? Ann. NY. Acad. Sci.

1297: 112-125.

Hiura, T. & Nakamura, M. (2013). Different mechanisms explain feeding type-specific patterns

of latitudinal variation in herbivore damage among diverse feeding types of herbivorous insects.

Basic Appl. Ecol. 14: 480-488.

Holeski, L.M., Keefover-Ring, K., Bowers, M.D., Harnenz, Z.T. & Lindroth, R.L. (2013).

Patterns of phytochemical variation in *Mimulus guttatus* (yellow monkeyflower). J. Chem. Ecol.

39: 525-536.

Inouye, D.W. (2008). Effects of climate change on phenology, frost damage, and floral

abundance of montane wildflowers. Ecology 89: 353-362.

IPCC (2014). Climate Change 2014: Impacts, Adaptation and Vulnerability: Regional Aspects.

Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Jactel, H., Koricheva, J. & Castagneyrol, B. (2019). Responses of forest insect pests to climate

change: not so simple. Curr. Opin. Insect Sci. 35:103-108.

Johnson, M.T.J. & Agrawal, A.A. (2005). Plant genotype and environment interact to shape a

diverse arthropod community on evening primrose (*Oenothera biennis*). Ecology 86: 874-885.

Johnson, M.T.J., Agrawal, A.A., Maron, J.L. & Salminen, J.P. (2009). Heritability, covariation

and natural selection on 24 traits of common evening primrose (*Oenothera biennis*) from a field

experiment. J. Evol. Biol. 22: 1295-1307.

Johnson, M.T.J. & Rasmann, S. (2011). The latitudinal herbivory-defence hypothesis takes a

detour on the map. New Phytol. 191: 589-592.

Ju, R.-T., Zhu, H.-Y., Gao, L., Zhou, X.-H. & Li, B. (2015). Increases in both temperature means

and extremes likely facilitate invasive herbivore outbreaks. Sci. Rep. 5: 1-10.

Kawagoe, T. & Kudoh, H. (2010). Escape from floral herbivory by early flowering in

Arabidopsis halleri subsp gemmifera. Oecologia 164: 713-720.

Kleiber, C., Zeileis, A. & Zeileis, M.A. (2020). Package 'aer'. R package version 12.

Krischik, V.A., Goth, R.W. & Barbosa, P. (1991). Generalized plant defense: effects on multiple

species. Oecologia 85: 562-571.

Kursar, T.A., Dexter, K.G., Lokvam, J., Pennington, R.T., Richardson, J.E., Weber, M.G. et al.

(2009). The evolution of antiherbivore defenses and their contribution to species coexistence in

the tropical tree genus Inga. PNAS 106: 18073-18078.

Leimu, R. & Fischer, M. (2008). A meta-analysis of local adaptation in plants. PloS one 3:

e4010.

Levin, D.A. (1975). Genic heterozygosity and protein polymorphism among local populations of

Oenothera biennis. Genetics 79: 477-491.

Liaw, A. & Wiener, M. (2002). Classification and regression by randomForest. R news 2: 18-22.

Louthan, A.M., Doak, D.F. & Angert, A.L. (2015). Where and when do species interactions set

range limits? Trends Ecol. Evol. 30: 780-792.

MacArthur, R.H. (1972). Geographical ecology: patterns in the distribution of species. Princeton

University Press, Princeton.

Malcolm, S. & Brower, L. (1989). Evolutionary and ecological implications of cardenolide

sequestration in the monarch butterfly. Experientia 45: 284-295.

Moles, A.T., Bonser, S.P., Poore, A.G.B., Wallis, I.R. & Foley, W.J. (2011). Assessing the

evidence for latitudinal gradients in plant defence and herbivory. Funct. Ecol. 25: 380-388.

Paquette, A. & Hargreaves, A.L. (2021). Biotic interactions are more often important at species'

warm versus cool range edges. Ecol. Lett. 24: 2427-2438.

Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. Annu.

Rev. Ecol. Evol. Syst. 37: 637-669.

Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. Nature 421: 37-42.

Pickett, S.T. (1989). Space-for-time substitution as an alternative to long-term studies. In: Long-

term studies in ecology: approaches and alternatives. Springer New York, NY, pp. 110-135.

Pilson, D. (2000). Herbivory and natural selection on flowering phenology in wild sunflower,

Helianthus annuus. Oecologia 122: 72-82.

Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J. et

al. (2013). Global imprint of climate change on marine life. Nat. Clim. Change 3: 919-925.

Pörtner, H.O., Roberts, D.C., Adams, H., Adler, C., Aldunce, P., Ali, E. et al. (2022). IPCC,

2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Cambridge University

Press Cambridge, UK and New York, NY.

R Development Core Team (2020). R: A language and environment for statistical computing.

Ripley, B., Venables, B., Bates, D.M., Hornik, K., Gebhardt, A., Firth, D. et al. (2013). Package

'MASS'. CRAN Repository. http://cran. R-project. org/web/packages/MASS/MASS. pdf.

Risk, C., McKenney, D.W., Pedlar, J. & Lu, P. (2021). A compilation of North American tree

provenance trials and relevant historical climate data for seven species. Sci. Dat. 8: 1-8.

Salminen, J.-P. & Karonen, M. (2011). Chemical ecology of tannins and other phenolics: we

need a change in approach. Funct. Ecol. 25: 325-338.

Sanford, E. & Kelly, M.W. (2011). Local adaptation in marine invertebrates. Ann. Rev. Mar. Sci.

3: 509-535.

Schemske, D.W., Mittelbach, G.G., Cornell, H.V., Sobel, J.M. & Roy, K. (2009). Is there a

latitudinal gradient in the importance of biotic interactions? Annu. Rev. Ecol. Evol. Syst. 40:

245-269.

Schreiber, S.G., Ding, C., Hamann, A., Hacke, U.G., Thomas, B.R. & Brouard, J.S. (2013). Frost

hardiness vs. growth performance in trembling aspen: an experimental test of assisted migration.

J. Appl. Ecol. 50: 939-949.

Sexton, J.P., McIntyre, P.J., Angert, A.L. & Rice, K.J. (2009). Evolution and ecology of species

range limits. Annu. Rev. Ecol. Evol. Syst. 40: 415-436.

Singh, S., Kaur, I. & Kariyat, R. (2021). The multifunctional roles of polyphenols in plant-

herbivore interactions. Int. J. Mol. Sci. 22: 1442.

Tang, Y., Horikoshi, M. & Li, W. (2016). ggfortify: unified interface to visualize statistical

results of popular R packages. R J. 8: 474.

Van Mantgem, P.J. & Stephenson, N.L. (2007). Apparent climatically induced increase of tree

mortality rates in a temperate forest. Ecol. Lett. 10: 909-916.

Vitt, P., Havens, K., Kramer, A.T., Sollenberger, D. & Yates, E. (2010). Assisted migration of

plants: changes in latitudes, changes in attitudes. Biol. Conserve. 143: 18-27.

Wadgymar, S.M., Ogilvie, J.E., Inouye, D.W., Weis, A.E. & Anderson, J.T. (2018).

Phenological responses to multiple environmental drivers under climate change: insights from a

long-term observational study and a manipulative field experiment. New Phytol. 218: 517-529.

Wadgymar, S.M. & Weis, A.E. (2017). Phenological mismatch and the effectiveness of assisted

gene flow. Conservation Biology 31: 547-558.

Walker, L.R., Wardle, D.A., Bardgett, R.D. & Clarkson, B.D. (2010). The use of

chronosequences in studies of ecological succession and soil development. J. Ecol. 98: 725-736.

Walker, W.H., Meléndez-Fernández, O.H., Nelson, R.J. & Reiter, R.J. (2019). Global climate

change and invariable photoperiods: A mismatch that jeopardizes animal fitness. Ecol. Evol. 9:

10044-10054.

Wang, T., Hamann, A., Spittlehouse, D. & Carroll, C. (2016). Locally downscaled and spatially

customizable climate data for historical and future periods for North America. PloS one 11.

Wang, T., O'Neill, G.A. & Aitken, S.N. (2010). Integrating environmental and genetic effects to

predict responses of tree populations to climate. Ecol. Appl. 20: 153-163.

Wetzel, W.C., Inouye, B.D., Hahn, P.G., Whitehead, S.R. & Underwood, N. (2023). Variability

in Plant–Herbivore Interactions. Ann. Rev. Ecol. Evol. S. 54.

Wickham, H. (2011). ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics 3: 180-

185.

Wilczek, A.M., Cooper, M.D., Korves, T.M. & Schmitt, J. (2014). Lagging adaptation to

warming climate in Arabidopsis thaliana. PNAS. 111: 7906-7913.

Williams, J.W. & Jackson, S.T. (2007). Novel climates, no-analog communities, and ecological

surprises. Front. Ecol. Environ. 5: 475-482.

Willis, C.G., Ruhfel, B., Primack, R.B., Miller-Rushing, A.J. & Davis, C.C. (2008). Phylogenetic

patterns of species loss in Thoreau's woods are driven by climate change. PNAS. 105: 17029-

17033.

Wogan, G.O.U. & Wang, I.J. (2018). The value of space-for-time substitution for studying fine-

scale microevolutionary processes. Ecography 41: 1456-1468.

Table 1. P-values and model coefficients for geographic, climatic, climate distance, resistance

to herbivores, phenological, and morphological models predicting seed number. All variables

within horizontal lines are within the same model. Bold indicates P < 0.05; underlined indicates P

< 0.10.

	Coefficient	P-value
Geography		
Latitude	12544.58	**0.002**
Latitude^2	-149.73	**0.002**
Distance to Common Garden	-0.0003	**0.04**
Climate		
Mean Annual Temperature	0.49	**0.004**
Mean Annual Temperature^2	-0.03	**0.001**
Relative Humidity	0.91	<u>0.063</u>
Relative Humidity^2	-0.01	<u>0.062</u>
Climate Distance		
Mean Summer Precipitation Distance	0.002	<u>0.083</u>
Mean Annual Temperature Distance^2	-0.02	**<0.001**
Resistance to Herbivores		
P. spumarius Number	0.29	**<0.001**
P. spumarius Number^2	-0.01	**0.01**
Fruits Impacted by *S. florida*	0.02	**0.003**
Fruits Impacted by M. brevivitella	-0.05	<u>0.051</u>
Phenology		
Flowering Date	0.28	**<0.001**
Flowering Date^2	-0.001	**<0.001**
Bolt Date	-0.04	**<0.001**
Growth Rate	64.39	<u>0.0503</u>
Growth Rate^2	-314.2	<u>0.073</u>
Morphology		
SLA	-0.02	0.12
Water Content	2.36	**0.029**
Water Content^2	-0.02	**0.032**
Trichome Number	0.04	**<0.001**

Trichome Number^2	-0.0002	**<0.001**

647 **Table 2. P-values and model coefficients for chemical defenses and principal components**

648 **predicting seed number.** Bold indicates P < 0.05; underlined indicates P < 0.10.

	Coefficient	P-value
Total Phenolics		
Leaf Total Phenolics	0.02	<u>0.084</u>
Flower Total Phenolics	0.05	0.12
Flower Total Phenolics^2	-0.0005	<u>0.079</u>
Fruit Total Phenolics	0.04	**0.03**
Fruit Total Phenolics^2	-0.0001	0.11
Ellagitannins		
Leaf Oenothein A	0.02	0.44
Leaf Oenothein A^2	-0.0002	0.76
Flower Oenothein A	0.07	**0.01**
Flower Oenothein A^2	-0.002	**0.01**
Fruit Oenothein A	0.01	**0.002**
Principal Components		
PC1	-0.09	**0.02**
PC2	0.2	**0.004**

652 Figure 1. Locations of 149 sampled populations of *O. biennis* across North America. Each black

653 dot represents one sampled population. The star represents the Koffler Scientific Reserve, the

654 location of the common garden.

Figure 2. Seed production of *O. biennis* in a Northern common garden as predicted by (A) latitude

657 of origin, (B) Euclidian distance to the common garden in km, (C) mean annual temperature at

658 collected location, and (D) relative humidity at collected location. The solid vertical line gives the

659 latitude or environmental condition at origin of population with the highest predicted seed

660 production. The dashed vertical line gives the latitude or environmental condition at the common

661 garden location. Each point represents one population.

Figure 3. Herbivory, phenology and plant morphology predicts seed production in *O. biennis* in a northern common garden. Plots are given for (A) number of *Philaenus spumarius*, (B) number of *Schinia florida* damaged fruits, (C) number of *Mompha brevivitella* damaged fruits, (D) flowering date (in Julian days), (E) bolt date (in Julian days), (F) growth rate (cm/d), (G) trichome number in a 5 mm diameter disk, (H) % water content, and (D) specific leaf area (SLA). Each point represents one population.

Figure 4. Fruit chemical defenses predict seed production of *O. biennis* in a common garden. Plots

are given for (A) flower total phenolics, (B) fruit total phenolics, (C) flower oenothein A, and (D)

fruit oenothein A. Units are in mg of given secondary metabolite(s) per g of dry mass. Each point

represents one population.

677   Figure 5 Principle component analyses of chemical, phenological and morphological traits of *O*.

678   *biennis*. (A) PCA bioplot showing first and second principle component. PC1 describes mostly

 chemical and phenological traits, while PC2 describes primarily morphological traits and growth

 rate. Each dot represents one population. Number of seeds per plant is color coded for each point.

 Seeds are predicted by (A) PC1 and (B) PC2). Each point represents one population.

684 Figure 6. Variable importance for *O. biennis* traits predicting seed production in a Northern

685 common garden. Values were generated using gradient forest.