Synaptic and peptidergic connectomes of the *Drosophila* circadian clock

Nils Reinhard [1,#], Ayumi Fukuda [2,#], Giulia Manoli[1], Emilia Derksen[1], Aika Saito[2], Gabriel Möller[1], Manabu Sekiguchi[2], Dirk Rieger[1], Charlotte Helfrich-Förster[1,4], Taishi Yoshii[2,5] and Meet Zandawala[1,3,6]

[1] Neurobiology and Genetics, Theodor-Boveri-Institute, Biocenter, Julius-Maximilians-University of Würzburg, Am Hubland, 97074 Würzburg, Germany
[2] Graduate School of Natural Science and Technology, Okayama University, Okayama 700-8530, Japan
[3] Department of Biochemistry and Molecular Biology, University of Nevada Reno, Reno, 89557, NV, USA
[4] Contact: charlotte.foerster@biozentrum.uni-wuerzburg.de
[5] Contact: yoshii@okayama-u.ac.jp
[6] Corresponding author. Contact: meet.zandawala@uni-wuerzburg.de
joint first authors

The circadian clock of animals regulates various physiological and behavioral processes in anticipation of, and adaptation to, daily environmental fluctuations. Consequently, the circadian clock and its output pathways play a pivotal role in maintaining homeostasis and optimizing daily functioning. To obtain novel insights into how diverse rhythmic physiology and behaviors are orchestrated, we have generated the first comprehensive connectivity map of an animal circadian clock using the *Drosophila* FlyWire brain connectome. We reveal hitherto unknown extensive contralateral synaptic connectivity between the clock neurons, which might contribute to the robustness of the clock by synchronizing clock neurons across the two hemispheres. In addition, we discover novel direct and indirect light input pathways to the clock neurons that could facilitate clock entrainment. Intriguingly, we observe sparse monosynaptic connectivity between clock neurons and downstream higher-order brain centers and neurosecretory cells known to regulate several behaviors and physiology. Therefore, we integrated single-cell transcriptomic analysis and receptor mapping to additionally decipher paracrine peptidergic signaling between clock neurons and with neurosecretory cells. Our analyses identified additional novel neuropeptides expressed in clock neurons and suggest that peptidergic signaling greatly enriches the interconnectivity within the clock network. Neuropeptides also form the basis of output pathways which regulate rhythmic hormonal signaling. The *Drosophila* circadian clock and neurosecretory center connectomes provide the framework to understand more complex clock and hormonal networks, respectively, as well as the rhythmic processes regulated by them.

Keywords: Connectomics, circadian rhythm, single-cell sequencing, endocrine signaling, neuropeptide, GPCR

Introduction

Almost all living organisms from humans to bacteria possess a circadian clock (Bell-Pedersen *et al.*, 2005, Saini *et al.*, 2019). This internal timekeeping system enables organisms to anticipate and adapt to the rhythmic environmental changes that occur over a 24-hour cycle. At their molecular core, these clocks are comprised of a cell-autonomous transcription-translation negative-feedback loop (Takahashi, 2017). In most animals, the master circadian clock in the brain receives light cues via the eyes which enable synchronization (or entrainment) with the external 24-hour light-dark cycles. The master clock sits at the top of the hierarchy and in turn modulates the activity of downstream neurons, as well as the peripheral clocks located in tissues throughout the body via endocrine and systemic signaling. In vertebrates, the master clock is located in the suprachiasmatic nucleus (SCN) of the hypothalamus and is comprised of approximately 20,000 neurons (Mohawk *et al.*, 2012). Extensive intercellular coupling between these neurons likely via neurotransmitters, neuropeptides, and gap junctions forms a neuronal network that is resilient to internal and environmental perturbations (Maywood *et al.*, 2006, Liu *et al.*, 2007). Systematic characterization of these diverse coupling mechanisms between clock neurons is thus crucial to

understanding circadian clock entrainment and the mutual coupling of the clock neurons. In addition, unraveling the clock output pathways which generate rhythmic behaviors and hormonal signaling can provide mechanistic insights into circadian regulation of organismal physiology and homeostasis.

These aims are particularly challenging to accomplish in vertebrates due to the large neuronal network size and resultant increase in complexity. However, the molecular clock architecture as well as the neuronal network motifs are highly conserved from humans to insects (Panda *et al.*, 2002, Helfrich-Förster, 2004). Hence, *Drosophila melanogaster* with its powerful genetic toolkit and a complete brain connectome represents an ideal system to decipher the clock network and its input and output pathways (Zheng *et al.*, 2018, Delventhal *et al.*, 2019, Johnstone *et al.*, 2022, Dorkenwald *et al.*, 2023, Schlegel *et al.*, 2023). Not surprisingly, the fly circadian clock, comprised of approximately 150 neurons, is extremely well characterized (Dubowy and Sehgal, 2017). These neurons have historically been classified into different groups of Lateral and Dorsal Neurons (LN and DN, respectively) based on their size, anatomy, location in the brain, and differences in gene expression (Helfrich-Förster, 2003, Abruzzi *et al.*, 2017). Further, these neuronal

subgroups are active at different times during the day and are consequently distinct functionally (Nitabach and Taghert, 2008, Liang *et al.*, 2016). Single-cell transcriptome sequencing analyses of clock neurons recently revealed additional heterogeneity within some of these clusters, which can largely be explained by neuronal signaling molecules that they express (Ma *et al.*, 2021, Ma *et al.*, 2023). Comprehensive mapping of the synaptic partners of clock neurons and comparing this connectivity can determine if the molecular heterogeneity based on gene expression also translates into heterogenous synaptic connectivity. Nonetheless, given the rich array of neuropeptides expressed in clock neurons, both synaptic and paracrine signaling appear crucial in mediating the connectivity between clock neurons as well as their output pathways. While recent work has begun to uncover some of this connectivity (Cavanaugh *et al.*, 2014, Nagy *et al.*, 2019, Barber *et al.*, 2021, Reinhard *et al.*, 2022a, Reinhard *et al.*, 2022b, Shafer *et al.*, 2022, Hidalgo *et al.*, 2023, Kurogi *et al.*, 2023), global analyses encompassing entire neuronal networks across both brain hemispheres are lacking.

Here, we harnessed the power of connectomics to generate the first complete connectivity map of the circadian clock. Our analyses revealed that light input from extrinsic photoreceptors to the clock neurons is largely indirect. Furthermore, we discovered novel ipsilateral synaptic connectivity between the clock neurons and further identified a subset of DN as an important hub that link the clock network across the two brain hemispheres via contralateral projections. We also elucidated the output pathways from the clock network that could affect general behavioral activity levels and organismal physiology. In particular, we characterized clock inputs to subsets of neurosecretory cells (NSC) in the brain which collectively represent a major neuroendocrine center that is functionally analogous to the vertebrate hypothalamus (Nässel and Zandawala, 2020). We observed sparse monosynaptic connectivity between clock neurons and NSC, suggesting that multi-synaptic connections and peptidergic signaling account for most of the connectivity, as is characteristic of the vertebrate clock output pathways. Hence, as a complementary approach, we also generated a putative peptidergic clock connectome by extensively mapping the expression of clock neuropeptides and their receptors. This peptidergic signaling greatly enriches the connectivity within the clock network and is the major output pathway to the neuroendocrine center which regulates systemic physiology.

Results
Identification of the circadian neuronal network in the FlyWire and hemibrain connectomes
The master clock in the *Drosophila* brain is comprised of approximately 150 neurons, a number which is derived from neuronal expression of different clock genes (Ma *et al.*, 2021). *Clk856-Gal4*, based on the *Clk* promoter, faithfully recapitulates expression in most of these clock neurons (Figure 1A). These neurons can be broadly classified into four

classes each of LN and DN (Figure 1B and Table 1). The LN comprise Lateral Posterior Neurons (LPN), dorsoLateral Neurons (LN_d), Ion Transport Peptide (ITP)-expressing LN (LN^{ITP}), and Pigment-Dispersing Factor (PDF)-expressing ventroLateral Neurons (LN_v^{PDF}). Conversely, the DN include anterior Dorsal Neurons 1 (DN_{1a}), posterior DN_1 (DN_{1p}), DN_2 and DN_3. These clock neuron classes can be further subdivided into different cell types (Figure 1C and Table 1). The estimated number of clock neurons in *Drosophila* is likely an underestimation as the precise number of DN_3 is unknown. This is partly due to the fact that drivers like *Clk856-Gal4* only include a small proportion of these cells (Ma *et al.*, 2021). Nevertheless, roughly 80 DN_3 are predicted within the clock network, and these have recently been classified into different cell types (Sun *et al.*, 2022). Most of the DN_3 have small somata and project to the central brain, and hence they are aptly called small Central Projecting DN_3 (s-CPDN₃) (Table 1) (Sun *et al.*, 2022). Similarly, a pair of DN_3 with large somata project to the central brain (large Central Projecting DN_3; l-CPDN₃), whereas about 6 neurons per brain hemisphere project to the anterior brain (Anterior Projecting DN_3; APDN₃) (Sun *et al.*, 2022). These latter cells also have larger somata than the s-CPDN₃ (Reinhard *et al.*, 2022b). Despite the large number of DN_3, previous studies suggest that they are less important for behavioral rhythmicity compared to some of the LN, which are regarded as the master pacemaker neurons (Rieger *et al.*, 2006, Yoshii *et al.*, 2012, Li *et al.*, 2018).

As a first step in determining the synaptic connectivity within the clock network, we identified most of the neurons from all the clock clusters in the FlyWire connectome generated in our companion papers (Dorkenwald *et al.*, 2023, Schlegel *et al.*, 2023). In total, we successfully identified 175 putative clock neurons based on morphology and previously determined connectivity (Figure 1C-D, Supplementary Videos 1-2) (Schubert *et al.*, 2018, Reinhard *et al.*, 2022a, Reinhard *et al.*, 2022b, Shafer *et al.*, 2022, Sun *et al.*, 2022). Occasionally, single clock neurons were not detected in one brain hemisphere, resulting in odd cell numbers for a given cluster (Figure 1C and Table 1).

The FlyWire connectome combines automatically detected chemical synapses with proofread neurons. These synapses represent an additional anatomical feature that could potentially distinguish neuronal groups. Consequently, we asked whether the classification of clock neurons based on differences in their synaptic connectivity aligns with the traditional anatomical and recent gene expression-based classification. To address this, we clustered clock neurons based on cosine similarity between their inputs from other neurons. Our clustering analysis successfully recovered most of the previously established subgroups (Figure S1A-B). Hence, synaptic connectivity-based classification of clock neurons aligns with the ones determined based on anatomical and gene expression differences. The exception to this includes the largest group of clock neurons, the DN_3, which

Figure 1: *Drosophila melanogaster* circadian clock network. (A) *Clk856-Gal4* drives GFP expression in most of the circadian clock neurons and some non-clock neurons (asterisk). Scale bar = 50μm. **(B)** *Drosophila* clock neurons can be divided into four classes each of Dorsal neurons (DN) and Lateral neurons (LN), which can be further subdivided into different cell types based on their morphology and gene expression. **(C)** Classification and numbers of clock neurons identified in the FlyWire connectome. Refer to Table 1 for details. **(D)** The morphology of identified clock neurons in the connectome largely resembles the morphology of clock neurons marked by *Clk856-Gal4*. **(E)** Broad synaptic interconnectivity between different cell types within the clock network. The direction of the arrow indicates the flow of information. Strong connectivity is observed from DN_{1a} to LN^{ITP}, from s-$CPDN_3$ to DN_{1p} cand., as well as from $DN_{1p}A$ to LN^{ITP}, LN_d^{CRY+} and s-$CPDN_3$. Note extensive contralateral connectivity between different cell types, most of which can be accounted for by the $DN_{1p}A$. Dashed line indicates brain midline. **(F)** Individual $DN_{1p}A$ in the right hemisphere form both ipsilateral and contralateral connections with s-$CPDN_3$, s-$CPDN_3$ cand., LN_d^{CRY+} and LN^{ITP}.

Table 1: Identification and classification of *Drosophila* clock neurons in the FlyWire brain connectome

Clock Super Class	Clock Class	Clock Cell Type	Expected	Observed
Dorsal neurons (DN)	anterior DN_1 (DN_{1a})	DN_{1a}	4	4
	posterior DN_1 (DN_{1p})	$DN_{1p}A$	~10	9
		$DN_{1p}B$	~4	4
		unclassified/DN_{1p} candidates (DN_{1p} cand.)	~14	13
	DN_2	DN_2	4	3
	DN_3	small Central Projecting DN_3 (s-CPDN$_3$)	~80	67
		small Central Projecting DN_3 candidates (s-CPDN$_3$ cand.)		25
		large Central Projecting DN_3 (l-CPDN$_3$)		2
		Anterior Projecting DN_3 (APDN$_3$)		12
Lateral neurons (LN)	Lateral Posterior Neurons (LPN)	LPN	6	6
	dorsoLateral Neurons (LN_d)	Cryptochrome-negative LN_d (LN_d^{CRY-})	6	6
		Cryptochrome-positive LN_d (LN_d^{CRY+})	4	4
	ITP-positive Lateral Neurons (LN^{ITP})	Cryptochrome- and ITP-positive LN_d ($LN_d^{CRY+ \& ITP}$)	2	2
		5th ventroLateral Neuron (5th-LN_v)	2	2
	PDF-positive ventroLateral Neurons (LN_v^{PDF})	large ventroLateral Neuron (l-LN_v)	8	8
		small ventroLateral Neuron (s-LN_v)	8	8
		Total	~152	175

segregated into at least nine distinct cell types (Figure S1A and C). Interestingly, the nine DN_3 cell types clustered with different LN and DN groups. These heterogenous clusters comprising clock neurons of different classes receive similar inputs from other neurons and likely play similar roles in the clock network (Figure S1A). In addition, we classified 25 DN_3 as s-CPDN3 candidates (s-CPDN$_3$ cand.) as they largely resembled the s-CPDN$_3$ (Figure 1C-D, S1C and Table 1). Lastly, in case of the DN_{1p}, we identified several new candidates which could not be classified into the existing DN_{1p} cell types (Figure S1D and Table 1).

Having identified all the clock neurons, we next sought to determine their synaptic interconnectivity which could facilitate intercellular coupling within the network. Generally, we regarded anything >4 common synapses per neuron as significant connections and >9 synapses as strong connections. We first wanted to validate our analysis by comparing it with previously reported connections. In agreement with previous reports (Reinhard *et al.*, 2022b, Shafer *et al.*, 2022), we observed strong synaptic connectivity from DN_{1a} to LN^{ITP}, from $DN_{1p}A$ to LN^{ITP} and LN_d, and from $DN_{1p}B$ to DN_2 clusters (Figure 1E and S2), highlighting the robustness of our approach. Importantly, our analysis also uncovered novel connections between the different subgroups. Specifically, we observed strong contralateral and ipsilateral connectivity from $DN_{1p}A$ to LN^{ITP}, as well as additional significant connections with s-CPDN$_3$ and LN_d^{CRY+} across both hemispheres. Similarly, LN^{ITP} also provide synaptic inputs to s-CPDN$_3$ clusters in both hemispheres (Figure 1E and S2). This raised the question whether $DN_{1p}A$ represents a heterogenous population where one subgroup forms ipsilateral connections and the other contralateral. To address this, we examined the connectivity at cellular resolution (Figure S3-S4) which revealed that individual $DN_{1p}A$ indeed form both ipsilateral and contralateral connections (Figure 1F). Our analysis thus identified $DN_{1p}A$ as an important center which links the clock network across the two brain hemispheres.

In contrast, there are virtually no synaptic connections between the s-LN_v and the LN_d neurons (Figure 1E and S2-S4), which control morning and evening activities, respectively.

To assess the extent of inter-individual differences in the numbers, neuronal projections, and synaptic connectivity of clock neurons, we next performed comparisons with the partial hemibrain connectome (Scheffer *et al.*, 2020). Several groups of clock neurons were previously identified in the hemibrain connectome including all s-LN_v, l-LN_v, LN_d, LN^{ITP}, LPN, DN_{1a}, and some DN_2, DN_{1p}, and DN_3 (Reinhard *et al.*, 2022a, Reinhard *et al.*, 2022b, Shafer *et al.*, 2022). In total, 44 clock neurons have been identified in the hemibrain connectome, with the majority of missing neurons belonging to the DN_3 subgroups (Figure S5A-D). Comparison of different subgroups revealed stable neuronal numbers across the two connectomes (Table 1 and Figure S5D). Similarly, there is a high degree of stereotypy in the connectivity between the clock clusters (Figure S5E-F). For instance, l-LN_v and DN_2 form the least synaptic contacts with other clock clusters. At the opposite end of the spectrum, s-CPDN$_3$ are connected to all the clock clusters with the exception of s-LN_v, l-LN_v, DN_{1a}, and DN_2. Given its partial nature, the hemibrain connectome lacks information about all contralateral connections, reiterating the significance of characterizing information flow across entire networks. Taken together, our analyses revealed hitherto unknown connectivity between the clock neurons which could contribute to the robustness of the master clock. Moreover, identification of the complete circadian neuronal network in the FlyWire connectome underscores the power of the fruit fly in pushing forward the frontier of our understanding of chronobiology.

Validating clock neuron connectivity using trans-synaptic tracing

While the FlyWire and hemibrain connectomes exhibit a high degree of stereotypy, we further used an independent approach to validate our connectivity analyses. We performed light microscopy-based trans-synaptic circuit

Figure 2: Validating connectivity within the clock network using *trans*-Tango. (A) Driving *trans*-Tango using *Clk4.1M-Gal4* revealed DN$_{1p}$, DN$_2$, DN$_3$, LPN, LN$_d$, 5th-LN$_v$ and s-LN$_v$ as post-synaptic partners of DN$_{1p}$. **(B)** Expressing *trans*-Tango in a subset of the DN$_3$ (including s-CPDN$_3$ and APDN$_3$) generated a post-synaptic signal in all dorsal clock neuron clusters, as well as LPN, LN$_d$ and l-LN$_v$. Scale bars = 50μm for overview and 20μm for higher magnification images. These two examples highlight a general agreement in synaptic connectivity observed with the connectomes and *trans*-Tango. Abbreviations: PER, Period; ITP, Ion transport peptide, RFP, Red fluorescent protein.

tracing using *trans*-Tango (Talay *et al.*, 2017). Highly specific driver lines for different populations of clock neurons (Figure S6-S7) were used to drive expression of a *trans*-Tango variant wherein nuclei of post-synaptic neurons are labeled with RFP (Snell *et al.*, 2022). These preparations were then stained with antibodies against the clock protein Period (a marker for clock neurons) or ITP (a marker for LN^{ITP}) to identify the different types of post-synaptic clock neurons. We used this approach to first validate the connections of DN$_{1p}$ since our connectome analyses identified them as a vital hub linking the two hemispheres.

Upon driving *trans*-Tango with *Clk4.1M-Gal4* which labels most DN$_{1p}$ (Figure S6), we observed post-synaptic signals in DN$_{1p}$, DN$_2$, DN$_3$, LPN, LN$_d$, 5th-LN$_v$, and s-LN$_v$ (Figure 2A). However, l-LN$_v$ and DN$_{1a}$ were not post-synaptic to DN$_{1p}$. While post-synaptic signal was not detected in most clock neurons of control flies, occasionally, a false post-synaptic signal was detected in two clock neurons (LN$_d$) from the entire network (Figure S8A). Hence, any potential synaptic output to LN$_d$ should be treated with caution. Nonetheless, our *trans*-Tango analysis of DN$_{1p}$ agrees with the connectivity of DN$_{1p}$ based on the connectomes (Figure S2

and S5F). Similarly, a split-Gal4 line targeting DN_3 drives post-synaptic signals in DN_{1a}, DN_{1p}, DN_2, DN_3, LPN, LN_d, and l-LN_v (Figure 2B), which mirrors the connectivity seen in the connectomes. Overall, we observed similar congruency between the two approaches with other Gal4-lines including those targeting DN_2 (Figure S8B), LPN (Figure S8C), and LN^{ITP} (Figure S8D).

Differences compared to the connectomes were observed when driving *trans*-Tango with *Pdf-Gal4* (for s-LN_v and l-LN_v) (Figure S8E), and with the DN_{1a}-specific split-Gal4 line (Figure S8F). In both cases, *trans*-Tango generated post-synaptic signals in more clock neurons than anticipated based on the connectomes. In the case of *Pdf-Gal4*, this discrepancy could be explained by the presence of PDF tritocerebrum (PDF-Tri) neurons (Selcho *et al.*, 2018). Although PDF-Tri neurons undergo apoptosis after adult eclosion, their post-synaptic partners could remain labeled and contribute to the additional connectivity observed with *trans*-Tango. In addition, differences may be caused by daily remodeling of neural circuits, as shown previously for s-LN_v and DN_{1a} (Fernandez *et al.*, 2008, Song *et al.*, 2021, Hofbauer *et al.*, 2023). The connectomes represent a single snapshot in time, whereas *trans*-Tango represents a cumulative output seen over the course of the experiment. Daily remodeling of s-LN_v and DN_{1a} terminals could thus alter their synaptic connectivity, resulting in additional post-synaptic partners as evident with *trans*-Tango. In support of this hypothesis, the hemibrain connectome, which is derived from a brain fixed at 1.5 hours after lights-on, confirmed the presence of synapses between DN_{1a} and LPN (Figure S5F), which was also evident in our *trans*-Tango analysis (Figure S8F). This connectivity, however, was not detected in the FlyWire connectome which represents brain connectivity at a different time of the day (Figure S2). In summary, our *trans*-Tango analysis is largely in agreement with the clock network generated using the connectomes.

Deciphering light input pathways via in-silico retrograde tracing of clock neurons

Following successful validation of our connectivity data, we next identified all the major classes of neurons providing inputs to the clock network. To this end, we utilized the annotation scheme of our companion paper (Schlegel *et al.*, 2023), which provides a hierarchical classification of all neurons in the connectome (Figure 3A). We found that neurons intrinsic to the brain provide the majority of the inputs to the clock network (Figure 3A-C). This includes visual centrifugal neurons projecting from the central brain to the optic lobes, visual projection neurons projecting from the optic lobes to the central brain, as well neurons intrinsic to the optic lobes and central brain (Figure 3C). Examining inputs to specific clock clusters, we observed differential inputs across all the subgroups (Figure 3B). As expected, s-LN_v and l-LN_v receive most of their input from optic lobe and visual centrifugal neurons as they have a large number of input sites in the optic lobes and the accessory medulla (aMe) (Figure S9). In contrast, $APDN_3$, l-$CPDN_3$, and LN^{ITP}

populations receive major inputs from visual projection neurons. The remaining clock clusters receive most of their inputs from central brain neurons (Figure 3B, S9, and S10). In some cases, such as DN_{1p} cand., DN_2, s-$CPDN_3$, and s-$CPDN_3$ cand., a large portion of these central neurons are clock neurons themselves, confirming prominent intercellular synaptic connectivity between some clock clusters. Interestingly, only 3 sensory neurons provide direct inputs to the clock network. These are antennal sensory neurons (Figure 3A and D) which may provide temperature inputs to LPN and DN_{1p} cand. (Alpert *et al.*, 2022).

Having broadly classified the inputs from different neuronal super classes to the clock network, we probed further and identified individual cells providing the strongest synaptic inputs to clock neurons. For this purpose, we used a stringent threshold of 80 synapses to obtain a narrow list of candidate inputs. Our analysis discovered 13 neurons, including 7 aMe neurons that are strongly connected to specific downstream clock neurons (Figure 3E). For example, individual aMe3 and aMe6a neurons can form more than 79 synapses with $APDN_3$, while aMe8 are similarly connected to $LN_d^{CRY+ \& ITP}$ clock neurons (Figure 3E). The unifying feature of these aMe neurons is their dense arborisation in the aMe and often additionally in the posterior lateral protocerebrum (Figure 3E), where they anatomically interact with clock neuron dendrites (Figure S9 and S10) (Reinhard *et al.*, 2022b, Tang *et al.*, 2022). Interestingly, the aMe neurons themselves receive strong inputs from the extraretinal photoreceptors. Specifically, aME3 and aME6a neurons receive strong inputs directly from the Hofbauer-Buchner (HB) eyelets (Figure 3E). Conversely, aME8 receive indirect inputs from ocellar retinula cells via the ocellar ganglion neurons (OCG) type 2c (OCG02c) (Figure 3E). Therefore, this analysis suggests that the clock receives strong light inputs from extrinsic photoreceptor cells, albeit indirectly. This is not surprising since light is the most important Zeitgeber for circadian clocks (Helfrich-Förster, 2020). Flies synchronize their circadian clocks with the light-dark cycles using these extrinsic photoreceptor cells as well as via the blue-light photoreceptor Cryptochrome (CRY), which is expressed in about half of the clock neurons (Figure 3F) (Helfrich-Förster, 2020). While CRY interacts with the core clock protein Timeless and can quickly reset the clock (Ceriani *et al.*, 1999, Emery *et al.*, 2000), the different photoreceptor cells are important for sensing dawn, dusk, high light-intensities and daylength, and for adapting morning and evening activities to the appropriate time of day (Rieger *et al.*, 2006, Schlichting *et al.*, 2019).

Regardless, we found little direct inputs from the photoreceptor cells and other sensory cells to the clock neurons (Figure 3A). This is consistent with previous findings which revealed that most of the light-input to the clock appears to be indirect (Alejevski *et al.*, 2019). In light of this and our *in-silico* circuit tracing analysis described

Figure 3: Light and other inputs to the clock network. (A) Input to the clock neurons grouped by the nine neuronal super classes annotated in the FlyWire connectome. Intrinsic neurons, specifically the central and visual projection neurons, are the largest groups providing inputs to the clock. **(B)** Breakdown of inputs to different types of clock neurons. Several clock neuron types receive inputs from other clock neurons. **(C)** Neurons from the four different super classes (central, visual projection, visual centrifugal and optic) which provide major inputs (based on number of neurons) to the clock. **(D)** Antennal sensory neurons (colored) providing potential temperature-related inputs to LPN and DN_{1p} cand. (grey). (legend continued on next page)

(E) aMe neurons provide the strongest inputs to clock neurons. Upstream partners of aMe neurons with strong connectivity include HB-eyelet and OCG02a. Numbers above the arrows indicate the number of synapses and numbers in circles indicate the number of neurons. Neuronal reconstructions of aMe3, aMe6a and aMe8 are shown in one hemisphere. **(F)** Cryptochrome (CRY)-positive clock neurons are shown in blue. **(G)** Disynaptic inputs to the clock from the three types of extrinsic photoreceptors (HB-eyelet, ocelli, photoreceptor cells of the compound eye) using a threshold of >4 synapses. **(H)** Disynaptic inputs to the clock from the three types of extrinsic photoreceptors using a threshold of >2 synapses. Note: HB eyelet was only identified in the left hemisphere. The lamina was only attached to the right hemisphere and was used for retinal inputs. R7 and R8 are not completely proofread and thus depicted using arrows. Abbreviations: FB, Fan-shaped body; HB, Hofbauer-Buchner; OC, Ocelli.

above, we comprehensively characterized indirect connectivity between photoreceptor cells and clock neurons. For this purpose, we traced all the disynaptic connections between them. Using the normal threshold of >4 synapses, we again recovered the strong connections from the H-B eyelets via the aMe3/ aMe6a to the APDN$_3$, and additional weaker connections to the s-CPDN$_3$, LN^{ITP}, l-LN$_v$, and LN$_d^{CRY+}$ (Figure 3G). Furthermore, we revealed connections from R7/R8 compound eye photoreceptors to several clock neurons via aMe12 (Kind *et al.*, 2021) and other interneurons. While we did not observe any disynaptic connections from the ocellar retinula cells to clock neurons using the normal threshold (Figure 3G), reducing the threshold to >2 synapses revealed connections from the ocelli to APDN$_3$ and l-CPDN$_3$ via OCG02c (Figure 3H). The synaptic connections from the ocelli to OCG and beyond are extensively characterized in our companion paper and demonstrate interesting details that may also be valid for the other photoreceptor inputs to clock neurons (Dorkenwald *et al.*, 2023). The majority of ocellar photoreceptors are synaptically connected to ocellar ganglion neurons with thick axons (OCG01a-f) or directly to descending neurons (DNp28). These connections likely enable fast behavioral responses. In contrast, axons of OCG02c that connect to the clock neurons are rather thin and not suited for fast neurotransmission. Instead, these neurons appear suited for collecting light information over time – a property needed for entraining the circadian clock. Further, collecting light information over larger time intervals may not require a high synapse density. Thus, 3 to 4 synapses between retinula cells and the relevant downstream OCG observed here could be sufficient for this purpose (Figure 3H). The same is also true for the photoreceptor cells of the compound eyes. Reducing the threshold of significant connections from 5 to 3 synapses revealed indirect clock input from additional photoreceptor cells, including those that project from the dorsal rim area of the eye (Figure 3H). These photoreceptor cells are involved in polarized vision and might contribute to time-compensated sun compass orientation (Homberg, 2004). Whether the connectivity observed with a lower threshold of >2 synapses is functional *in vivo* remains to be seen; however, this is very likely since there are usually many photoreceptor cells that synapse onto only a few aMe neurons. For example, theoretically, the ~300 pale R8 cells project to only 3 aMe12 neurons (Kind *et al.*, 2021), resulting in ~100 connections on average per aMe neuron. Even if each of these connections were mediated via only 3 synapses, each aMe neuron could potentially receive inputs from R8 cells via 300 synapses, which is quite substantial (Figure 3H).

In-silico anterograde tracing of clock neurons

Delineating the output pathways that translate daily 24-hour oscillations of the molecular clock into physiological and behavioral rhythms remains a major focus in chronobiology. Using the same strategy as above to identify the inputs, we systematically classified all the neurons downstream of the clock network. Most synaptic output from the clock network is directed to intrinsic brain neurons, and in particular, the central brain neurons (Figure 4A-C). Except for l-LN$_v$, all clock clusters have a majority of their output onto central brain neurons. l-LN$_v$ mostly provide inputs to Medullary intrinsic neurons in the optic lobe (Figure 4B-C), consistent with their role in adapting the sensitivity of the visual system to the time of day (Chen *et al.*, 1992, Pyza and Meinertzhagen, 1997). Further, the majority of the output from DN$_{1p}$A is onto visual projecting and central brain neurons that are part of the clock network (Figure 4B). After broadly classifying clock outputs, we next focused on specific cell types which receive the strongest synaptic inputs (>49 synapses) from clock neurons using an approach similar to the one used earlier for clock inputs. Our analysis identified the enigmatic Clamp neurons (Figure 4D), which receive strong synaptic inputs from APDN$_3$. While the functions of most of these clamp neurons are still unknown, some of them output onto descending neurons, while others promote sleep (Sun *et al.*, 2022). Moreover, DN$_{1p}$ provide strong inputs to Tubercle-innervating neurons (Figure 4E), which are part of the anterior visual pathway (Hulse *et al.*, 2021). Lastly, several clock neurons are highly-connected to diverse neurons from different neuropil regions (Figure 4F).

Next, we examined clock inputs to descending neurons which could influence locomotor and other behaviors regulated by neurons in the ventral nerve cord. Interestingly, clock neurons provide direct inputs to 16 descending neurons (Figure 4A and G). These descending neurons include those which have not yet been classified (Figure 4H), as well as Allatostatin-C (AstC) and SIFamine (SIFa) peptidergic neurons (Figure 4I), the latter of which modulate feeding, mating, and sleep (Nässel and Zandawala, 2022). Direct synaptic inputs to downstream descending neurons predominantly derive from s-CPDN$_3$, s-CPDN$_3$ cand. and LPN (Figure 4B). When considering disynaptic connections, the connectivity between the clock network and descending neurons increased drastically, with approximately 20% of all descending neurons receiving indirect inputs from most of the clock neurons (Figure 4G). In summary, our connectivity analysis indicates that the clock can have a major influence on diverse behaviors, including locomotion, via outputs to descending neurons.

Figure 4: Direct and indirect clock output pathways. (A) Clock output grouped by the nine neuronal super classes annotated in the FlyWire connectome. Intrinsic neurons, specifically the central neurons, are the largest group of neurons downstream of clock neurons. **(B)** Breakdown of outputs from different types of clock neurons. All clock neurons, except for l-LN$_v$, provide a majority of their output to central neurons. l-LN$_v$ mostly provide inputs to optic neurons. **(C)** Neurons from the four different super classes (central, optic, visual projection and visual centrifugal) which receive inputs from the clock. **(D)** APDN$_3$ form the most output synapses in the clock network and are highly connected to several types of Clamp neurons (CL). **(E)** DN$_{1p}$ provide strong outputs to Tubercle-innervating neurons. **(F)** Several clock neurons provide strong inputs to neurons from diverse neuropils. **(G)** Mono- and di-synaptic clock inputs to descending neurons. **(H)** Novel and **(I)** peptidergic descending neurons receiving direct clock input. Mono- and di-synaptic clock inputs to neurons associated with the **(J)** central complex and **(K)** mushroom bodies. Note that there are only a few direct connections from the clock to downstream higher brain centers. Numbers above the arrows indicate the number of synapses and numbers in circles indicate the number of neurons. Abbreviations: EB, Ellipsoid body; FB, Fan-shaped body; NO, Noduli; EPG, Ellipsoid body-protocerebral bridge-gall neuron; PFGs, Protocerebral bridge glomerulus-fan-shaped body-ventral gall surround; KC, Kenyon cell; DAN, Dopaminergic neuron; MBON, Mushroom body output neuron.

In addition, the circadian clock is also known to modulate behaviors such as activity/sleep, spatial orientation, and learning and memory (Chouhan *et al.*, 2015, Fropf *et al.*, 2018, Liang *et al.*, 2019, Mathejczyk and Wernet, 2019, Warren *et al.*, 2019, Flyer-Adams *et al.*, 2020). These behaviors are regulated by higher brain centers such as the central complex and mushroom bodies. Consistent with previous results (Reinhard *et al.*, 2022a, Reinhard *et al.*, 2022b), we found few direct connections from the clock to neurons associated with the central complex (Figure 4J) and mushroom bodies (Figure 4K). We predicted that the clock output to these higher coordination centers is either indirect or paracrine via neuropeptides. Consistent with this prediction, we found prominent disynaptic connectivity between clock neurons and central complex neurons (mainly fan-shaped body (FB) and ellipsoid body neurons (EB)), as well as between clock neurons and Kenyon cells (KC), dopaminergic neurons (DANs) and mushroom body output neurons (MBONs) (Figure 4J-K). In support of paracrine signaling, receptors for several clock neuropeptides are also enriched in higher brain centers (Nässel and Zandawala, 2019). Taken together, circadian modulation of neurons regulating diverse behaviors is largely indirect or paracrine. Similarly, we observed very few direct connections between clock neurons and endocrine cells which influence organismal physiology and systemic homeostasis (Figure 4A-B). We address this connectivity in more detail below.

Identification and characterization of the neuroendocrine center in the FlyWire connectome
While recent work has unraveled some clock output pathways to endocrine cells, our collective understanding of the circadian regulation of endocrine rhythms is poor (Nagy *et al.*, 2019, King and Sehgal, 2020, Barber *et al.*, 2021, Hidalgo *et al.*, 2023). To address this knowledge gap, we first identified and classified all endocrine or NSC in the brain which are a major source of circulating hormones. These endocrine cells can be broadly classified into lateral, medial, and subesophageal zone NSC (l-NSC, m-NSC, and SEZ-NSC,

respectively) based on their location in the brain. Their axons exit the brain via a pair of nerves (nervii corpora cardiaca, NCC), and depending on the cell-type, innervate the corpora cardiaca, corpora allata, hypocerebral ganglion, crop, aorta or the anterior midgut (Figure 5A). Their axon terminations form neurohemal sites through which hormones are released into the circulation or locally on peripheral targets such as the crop. Collectively, the NSC form a major, yet distributed, neuroendocrine center that is functionally analogous to the hypothalamus as the cells express homologs of vertebrate corticotropin-releasing hormone, gonadotropin-releasing hormone and prolactin-releasing peptide (Nässel and Zandawala, 2020). We identified all brain NSC in the FlyWire connectome by isolating the nerve bundle containing their axons (Figure 5B-C). In total, we independently identified 80 brain NSC (Supplementary Video 3), in agreement with our companion studies (Dorkenwald *et al.*, 2023, Schlegel *et al.*, 2023). This number is substantially larger compared to the larvae where only 56 NSC are present (Table 2) (Huckesfeld *et al.*, 2021). Hence, we compared different types of NSC across development to determine whether the additional NSC in adults represent an expansion of NSC groups found in larvae or if they belong to a yet unidentified class. In larvae, two groups of m-NSC express myosuppressin (DMS) and Diuretic Hormone 44 (DH44) (Huckesfeld *et al.*, 2021). A third group of m-NSC express *Drosophila* Insulin-Like Peptides (DILP) 2, 3, and 5, and are commonly referred to as Insulin-Producing Cells (IPC). In addition, there are five groups of l-NSC which express ITP, corazonin (CRZ), Diuretic Hormone 31 (DH31), prothoracicotropic hormone, and eclosion hormone. The latter two populations of NSC undergo apoptosis soon after adult eclosion and are thus not found in mature adults. Lastly, the SEZ-NSC include two groups which express CAPA and Hugin neuropeptides. With the exception of CRZ neurons where 8 additional neurons are present in adults, the proportions of the remaining groups are thought to remain constant across development. We propose and utilize a systematic

Table 2: Classification of *Drosophila* neurosecretory cells (NSC) based on their cell body position in the central brain

NSC Super Class	Neuropeptide	Other common names	NSC Class	# in larvae	Expected # in adults	Observed # in adults
Medial	Myosuppressin (DMS)	SP3	m-NSC^{DMS}	4	4	4
Medial	Diuretic Hormone 44 (DH44)	DH44-PI	m-NSC^{DH44}	6	6	6
Medial	*Drosophila* Insulin-Like Peptides (DILP) 2, 3 and 5	IPC	m-NSC^{DILP}	14	14	18*
Medial	Eclosion Hormone (EH)	V$_m$	NSC^{EH}	2	0	0
Lateral	Corazonin (CRZ)	DLP, CC-LP2	l-NSC^{CRZ}	6	14**	6**
Lateral	Ion-Transport Peptide (ITP)	ALK, ipc-1	l-NSC^{ITP}	8	8	8
Lateral	Diuretic Hormone 31 (DH31)	CA-LP	l-NSC^{DH31}	6	6	6
Lateral	ProThoracicoTropic Hormone (PTTH)	PG-LP	l-NSC^{PTTH}	4	0	0
Subesophageal	Capability (CAPA)	Large SEG neurons	SEZ-NSC^{Capa}	2	2	2
Subesophageal	Hugin	HugRG	SEZ-NSC^{Hugin}	4	4	4
Medial	Unknown		m-NSC			12
Lateral	Unknown		l-NSC			14
			Total	56	58	80

Notes: * It is unclear at this point if all 18 m-NSC^{DILP} express all three or different combinations of DILP2, 3 and 5.
** Contrary to expectation, there are only 6 l-NSC^{CRZ} in adults based on our analysis (see Figure S12).

Figure 5: *Drosophila* neuroendocrine center connectome. (A) Schematic drawing of the different types of neurosecretory cells (NSC) and their projections to different release sites within the fly. **(B)** Reconstructions of the 80 NSC within the connectome. **(C)** All NSC projections exit the brain via NCC. Electron micrograph showcasing a cross section of the NCC. Scale bar = 750nm **(D)** Identification of 80 NSC and their classification based on location and neuropeptide expression. Refer to Table 2 for further details. **(E)** Direct and multisynaptic inputs from clock neurons to NSC. Di- and multisynaptic connections increase the connectivity between clock neurons and NSC. Numbers in circles indicate the number of neurons. Asterisk denotes one DN₁ₐ which was not found on Codex (https://codex.flywire.ai/) and was thus not included in this analysis. **(F)** LN^ITP (both hemispheres) and DN₁ₚ cand. (right hemisphere only) are the only clock neurons directly connected to NSC (l-NSC^DH31). Based on the direct connection shown in **(E)**. The numbers next to arrows indicate the number of synapses. **(G)** Reconstructions of non-clock neurons which mediate disynaptic connections between the clock network and NSC. Abbreviations: CC, Corpora cardiaca; CA, Corpora allata; NCC, Nervii corpora cardiaca; VNC, Ventral nerve cord; SEZ, Subesophageal zone.

nomenclature for all brain NSC based on their location and neuropeptide identity (Table 2).

While all adult SEZ-NSC and some l-NSC can easily be classified based on their morphology and location (Figure 5B-C), this approach is not feasible for m-NSC since they are clustered together in the superior medial protocerebrum and appear similar based on gross morphology. Therefore, we asked whether cosine similarity-based clustering, such as the one used previously for clock neurons, can be used to

distinguish and identify different m-NSC clusters, as well as the l-NSC^CRZ cluster. As expected, SEZ-NSC^Hugin, SEZ-NSC^Capa, and l-NSC^DH31 form three separate clusters (Figure S11A-C). Most l-NSC^ITP (Figure S11B) do not have any input synapses in our dataset and were thus excluded from this analysis. Notably, this analysis resulted in two clusters of m-NSC comprising 4 and 6 neurons each. Hence, these clusters likely represent m-NSC^DMS and m-NSC^DH44, respectively (Figure S11A and D). We obtained two additional clusters of m-NSC comprising 18 and 12 neurons, with the latter having

low similarity between the neurons. The cluster comprising 18 m-NSC likely represents IPCs (m-NSC^{DILP}). Whether all cells in this cluster express DILP2,3 and 5 remains unknown; however, DILP2 is expressed in more than 14 neurons in adults (Ohhara *et al.*, 2018).

Interestingly, we could only reliably identify 6 out of the expected 14 l-NSC^{CRZ} (Figure S11B and Table 2). These 6 neurons cluster into two separate clades (Figure S11A) as they represent a heterogenous population both anatomically and functionally (Oh *et al.*, 2019, Zandawala *et al.*, 2021, Manoli *et al.*, 2023). Our inability to identify the remaining 8 CRZ neurons inspired us to examine if these adult-specific CRZ neurons are indeed neurosecretory. Using *Gr64a-Gal4* to label the adult-specific CRZ neurons (Fujii *et al.*, 2015), we showed that there are only 6 adult l-NSC^{CRZ} (Figure S12). Contrary to our expectation, the adult-specific CRZ neurons do not project via the NCC, and are thus not endocrine. Hence, our clustering analysis accounts for all the NSC that persist into adulthood. Additionally, it uncovered 12 putative m-NSC and 14 putative l-NSC in the adult brain (Figure 5D, S11E, Supplementary Video 4). These neurons have comparatively smaller somata compared to other identified NSC (Figure S11), and have relatively fewer dense core vesicles than neurons such as l-NSC^{ITP} (data not shown). Hence, the type (neuropeptide, biogenic amine or fast-acting neurotransmitter) and the identity of the signaling molecules within these neurons remains unknown.

Informed by these new insights on different subgroups of NSC, we first explored potential inputs from clock neurons. Intriguingly, we observed sparse direct inputs from clock neurons to most NSC (Figure 5E), despite the fact that clock neuron projections are closely associated with NSC dendrites in the superior medial and lateral protocerebrum. The only exceptions are l-NSC^{DH31} which receive inputs from LN^{ITP} and DN$_{1p}$ cand. (Figure 5F). This observation prompted us to examine other types of synaptic inputs to NSC to ensure that the lack of synaptic connectivity between the clock network and NSC was not due to false negatives. Consistent with the location of m-NSC dendrites in the tritocerebrum (King *et al.*, 2017, Zandawala *et al.*, 2018, Hadjieconomou *et al.*, 2020), a large portion of their inputs derive from the subesophageal zone hemi-lineages (data not shown). Therefore, a lack of direct clock output to the NSC is genuine, and this connectivity is likely indirect or paracrine in nature. To explore the extent of indirect connections, we examined di- and multi-synaptic connectivity between clock neurons and NSC. Approximately half of the clock neurons provide inputs to half of the NSC disynaptically. The interneurons which facilitate these connections mainly include central neurons (Figure 5G). Interestingly, almost the entire clock network is connected to most of the NSC when accounting for multisynaptic connections (Figure 5E). Taken together, these prominent indirect connections between the clock network and NSC could form the basis of circadian regulation of systemic physiology.

Identifying the molecular basis of paracrine clock output pathways

Given the large repertoire of neuropeptides previously shown to be expressed in clock neurons (Abruzzi *et al.*, 2017, Ma *et al.*, 2021, Ma *et al.*, 2023), we predict that peptidergic paracrine signaling is crucial in mediating intercellular coupling within the clock network as well as output to downstream neurons such as NSC. First, we determined if any additional neuropeptides are expressed in clock neurons. To address this, we used a publicly-available single-cell transcriptome dataset of clock neurons (Ma *et al.*, 2021) combined with immunohistochemical localization and T2A-Gal4 lines. Unsupervised clustering of all clock neuron transcriptomes using t-SNE analysis yields 32 independent clusters (data not shown), 16 of which have high expression of clock genes (*tim* and *Clk*) and can be reliably identified based on known markers (Figure 6A-B). Our analysis revealed that most clock clusters express at least one neuropeptide (Figure 6B). Consistent with previous studies, l-LN$_v$ express high levels of *Pdf*, whereas s-LN$_v$ express both *Pdf* and *short neuropeptide F* (*sNPF*) (Figure 6B). Similar coexpression of neuropeptides is also observed in other clusters including the "DN$_{1p}$$^{CNMa \& AstC}$" cluster which coexpresses *CNMamide* (*CNMa*), *AstC*, and *Dh31* neuropeptides. In total, at least 12 neuropeptides are highly expressed in the clock network (Figure 6B-C). Importantly, this includes novel clock-related neuropeptides, namely DH44 and Proctolin (Proc). *Dh44* is expressed in several clock clusters including DN$_{1a}$, DN$_{1p}$^{AstA}, DN$_{1p}$^{sNPF}, DN$_3$^{VGlut}, LPN and LN$_d$^{NPF} (Figure 6B-C). Hence, *Dh44* is coexpressed with *CCHamide1* (*CCHa1*) in DN$_{1a}$ and with *Allatostatin-A* (*AstA*) in the DN$_{1p}$^{AstA} cluster (Figure 6B-C). We independently confirmed the presence of DH44 peptide in these clusters using a combination of DH44 antibody or *DH44-T2A-Gal4* (Figure 6D, S13, and S14). *Proc*, on the other hand, is expressed in the DN$_{1p}$^{CNMa} and the DN$_2$ clusters which was verified by driving GFP using a *Proc-T2A-LexA* driver (Figure 6E and S13). Lastly, we also detected *AstC* expression in additional clock neurons. AstC immunoreactivity was previously localized to DN$_{1p}$, DN$_3$, and LPN (Diaz *et al.*, 2019, Reinhard *et al.*, 2022a), which is in agreement with *AstC* transcript expression in DN$_{1p}$^{Rh7} and DN$_{1p}$$^{CNMa \& AstC}$ clusters (Figure 6B-C). Here, we show that AstC is additionally expressed in DN$_2$, which were labeled using an antibody against the clock protein Vrille (VRI) (Figure 6B, 6F, S13, and S14). Our comprehensive expression analyses revealed the complete neuropeptide complement of clock neurons (Figure 6B-F, S13, and S14) and provides the basis to explore paracrine targets of clock neurons.

As a first step in this direction and to validate our approach, we focused on select NSC which have been extensively characterized previously. We predicted that NSC are

Figure 6: Circadian clock neuropeptidome. (A) Single-cell RNA sequencing of clock neurons reveals 16 distinct clock neuron clusters (shown in a t-SNE plot) that can be reliably identified based on known markers. **(B and C)** Clock neurons express at least 12 different neuropeptides. **(D)** DH44 is a new clock-related neuropeptide which is expressed in APDN3, DN1a, LNd and LPN (arrowheads). **(E)** Proctolin is a new clock-related neuropeptide which is expressed in one DN1p (open arrowhead) and two DN2 (filled arrowheads). **(F)** AstC is expressed in DN2 in addition to other clock neurons. Double arrowheads indicate DN3 labelled by anti-AstC, filled arrowhead indicate DN2, and open arrowheads indicate DN1p. Scale bars = 50μm for overview and 10μm for higher magnification images. Abbreviations: PER, Period; TIM, Timeless; VRI, Vrille; PDF, Pigment dispersing factor; DH44, Diuretic hormone 44; AstC, Allatostatin-C. **(G)** Single-cell transcriptomes of NSC express receptors for clock-related neuropeptides.

targeted by clock-related neuropeptides since they receive sparse monosynaptic inputs from clock neurons despite being closely associated with them anatomically. To investigate potential paracrine signaling between clock neurons and NSC, we again turned our attention to single-cell transcriptomics. We identified m-NSC^{DH44}, m-NSC^{DILP}, l-NSC^{CRZ}, and l-NSC^{ITP} in single-cell RNA transcriptomes of the *Drosophila* brain (Davie *et al.*, 2018) based on previously identified markers, and quantified the expression of clock peptide receptors (Figure 6G). We could not reliably mine l-NSC^{DH31} and m-NSC^{DMS} due to the lack of multiple molecular markers. We also disregarded SEZ-NSC^{Capa} and SEZ-NSC^{Hugin} from this analysis since they are much further away from clock neuron projections. Consistent with our prediction, our analysis indicated that multiple receptors for clock peptides are indeed expressed in NSC. Modulatory inputs to m-NSC^{DILP} have been examined extensively and our analysis based on single-cell transcriptome sequencing data is in agreement with previous anatomical and functional studies (Kapan *et al.*, 2012, Hentze *et al.*, 2015, Nässel and Vanden Broeck, 2016, Nagy *et al.*, 2019, Oh *et al.*, 2019). Together, this analysis uncovers the molecular substrates of paracrine signaling between clock neurons and NSC.

We next explored the magnitude of peptidergic paracrine signaling between clock neurons themselves. To accomplish this, we extensively mapped the expression of clock neuropeptide receptors within the clock network, consequently generating a putative clock network neuropeptide connectome. Surprisingly, *Pdf receptor* (*Pdfr*) transcript is highly enriched in all clock neurons (Figure 7A-B), which we verified independently by expressing GFP using *Pdfr-T2A-GAL4* (Figure 7C). Other receptors are more sparsely expressed within the clock network (Figure 7A-B). However, most clock clusters express at least two receptors, with DN$_{1p}$^{sNPF} expressing at least 7 receptors. We validated our single-cell transcriptome analysis by mapping the expression of select receptors using T2A-Gal4 knock-in lines. Our anatomical mapping of receptors (Figure S15) is largely in agreement with transcriptome data (Figure 7D). In some cases, however, receptor mapping can provide additional insights. For instance, there are four transcript variants (A, B, C, and D) encoding the *Drosophila* Neuropeptide F receptor (NPFR). Knocking in the Gal4 cassette in different exons can enable expression mapping of specific transcript variants. Using Gal4 lines specific for NPFR-A/C and NPFR-B/D isoforms, we showed that these isoforms are differentially expressed across the clock clusters, with A/C isoforms expressed more broadly than B/D isoforms (Figure 7D and S15).

Finally, we utilized our expression data of neuropeptides and their cognate receptors in clock neurons to delineate potential paracrine signaling pathways within the network. Taking s-LN$_v$ as an example, this cluster expresses both PDF and sNPF. At the same time, PDFR is expressed in all clock clusters, whereas sNPF receptor is only expressed in DN$_2$, DN$_3$, and l-LN$_v$. Thus, in addition to providing synaptic inputs to DN$_2$ and DN$_3$, s-LN$_v$ can potentially provide paracrine inputs to all clock clusters (Figure 7E). Expanding this analysis to other clock clusters allowed us to generate a putative clock network neuropeptide connectome (Figure 7E). This peptidergic connectome, however, can only be considered putative for three main reasons: (1) all clock neuropeptides are also expressed in other non-clock neurons (Nässel and Zandawala, 2019), suggesting likely inputs from neurons extrinsic to the clock network, (2) we don't account for peptide efficacies and receptor affinities since these values were independently determined in distinct systems thus making comparisons difficult (Larsen *et al.*, 2001, Johnson *et al.*, 2005, Mertens *et al.*, 2005, Ida *et al.*, 2011) and (3) distance of peptide diffusion may vary. Nonetheless, the presence of receptors for clock peptides in other clock neurons provides the molecular basis for potential paracrine signaling between them. Our putative peptidergic connectome highlights additional novel connections between clock clusters, albeit with some limitations. For instance, all clock clusters may be connected to s-LN$_v$ despite forming few to no synaptic contacts (Figure 7E). Connectivity to l-LN$_v$ is also enhanced by paracrine signaling, although not to the same extent as it is for s-LN$_v$. In summary, peptidergic signaling greatly enriches the connectivity between different subsets of clock neurons. Additional investigations are necessary to determine which of these putative connections are functional *in vivo*.

Discussion

Peptidergic signaling supplements synaptic connectivity within the circadian clock network

Using a multipronged approach centered around the Fly-Wire connectome (Dorkenwald *et al.*, 2023, Schlegel *et al.*, 2023), we describe the first whole-brain neural connectome of an animal circadian clock. Our search using the complete brain connectome eluded identification of some clock neurons including a few DN$_{1p}$, DN$_2$, and s-CPDN$_3$. Nevertheless, given the fact that we identified approximately 98% of the expected clock neurons as well as some additional DN$_3$ (Table 1), our clock neuron synaptic wiring diagram is complete enough to be regarded as a connectome. Our clock connectome is also a significant upgrade (7-fold larger numerically) compared to the partial connectivity diagram based on the hemibrain connectome reported earlier (Shafer *et al.*, 2022). The previous analysis was based on only 24 clock neurons and largely focused on LN clusters, while excluding l-LN$_v$, DN$_2$, and DN$_3$ clusters due to the incomplete nature of the dataset. However, as evident from our analysis here, the DN in fact represent an important hub in the clock network and display high synaptic connectivity. In particular, DN$_{1p}$ play a large role in clock cluster interconnectivity and DN$_3$ appear important for clock output pathways. Our analysis also sheds light on the precise number of DN$_3$ in the clock network. Although approximately 80 DN$_3$ were previously estimated in the clock network, there appear to be at least 106

Figure 7: Paracrine signaling within the clock network (A) Expression of clock-related neuropeptide receptors in different clock clusters identified using single-cell transcriptome analysis. **(B)** t-SNE plots showing expression of receptors in different clock clusters. Clustering is based on Figure 6A. **(C)** *Pdfr-T2A-Gal4* drives GFP expression in all clock cell types. Arrowheads indicate different types of clock neurons which express PDFR. Scale bars = 20μm. Abbreviations: PDF, Pigment dispersing factor; TIM, Timeless. **(D)** Schematics depicting expression of receptors in different clock neurons. The schematics are based on GFP expression using T2A-GAL4 lines for different receptors (see Supplementary Figure 14 for confocal images). **(E)** Chord diagrams showing synaptic and potential peptidergic paracrine connections between clock neurons. This figure is based on synaptic connectivity in Figure 1E, and peptide and receptor expression mapping reported in Figures 6B-F, 7A-D and S13-15.

DN₃ based on the connectome. It is likely that common global clock network Gal4 drivers such as *Clk856*, *per*, and *tim*-Gal4 drive weak or no expression in some of these DN₃, making it difficult to characterize them anatomically and functionally (Kaneko and Hall, 2000, Ma *et al.*, 2021). We anticipate that resources such as NeuronBridge (Meissner *et* *al.*, 2023), which allow for comparisons between electron and light microscopy datasets, will facilitate the identification of Gal4 drivers that target these elusive DN₃. In addition, we also characterize the molecular basis for neuropeptide connectivity between the clock neurons, consequently generating a putative peptidergic clock connectome. Similar to

vertebrates (Wen *et al.*, 2020, Morris *et al.*, 2021), the *Drosophila* clock network is highly peptidergic. All clock neuron clusters express at least one neuropeptide, for a total of 12 neuropeptides within the entire network. Notably, a majority of the clock clusters express two neuropeptides, and several express three, while the LPN express four neuropeptides. For example, we showed here that DH44 and Proc are found in clock neurons that already express two to three other neuropeptides. Similar neuropeptide coexpression is also evident in SCN neurons and thus appears to be a common feature of clock neurons (Wen *et al.*, 2020). Surprisingly, there is little to no overlap in the neuropeptide complement of the *Drosophila* and vertebrate clock neurons (Figure 8). Orthologs of vertebrate clock neuropeptides including vasoactive intestinal peptide, arginine vasopressin, neuromedin S, cholecystokinin, gastrin-releasing peptide and prokineticin 2 are either absent in the *Drosophila* genome or expressed outside the clock network. Hence, *Drosophila* and vertebrates have evolved to utilize different signaling molecules while still conserving the diversity of neuropeptide signaling within the clock networks. Remarkably, with the exception of PDF, there appears to be little conservation in neuropeptide identities of clock neurons across different insects (Stengl and Arendt, 2016, Beer and Helfrich-Förster, 2020). This suggests that it is more important to conserve the mode of communication (paracrine signaling) rather than the messenger (specific neuropeptide).

Contralateral connectivity within the network prevents decoupling of clock neurons across the hemispheres

Analysis of interconnectivity within the clock network revealed extensive contralateral synaptic connectivity between the clock neurons, which is largely mediated by $DN_{1p}A$ and to a lesser extent by $DN_{1p}B$, s-$CPDN_3$, l-$CPDN_3$ and LN^{ITP}. Furthermore, paracrine peptidergic signaling amongst clock neurons has the potential to further strengthen this contralateral connectivity. Such a strong bilateral coupling of clock neurons prevents the internal desynchronization of clock neuron oscillations between the two hemispheres – a phenomenon that can happen in other insects and even in mammals, but so far has not been observed in fruit flies (Helfrich-Förster, 2004). In addition, we observed minimal synaptic connectivity between the s-LN_v and the LN_d neurons, which control morning and evening activities, respectively. Consistent with this observation, the phase relationship between morning and evening oscillators is plastic, consequently facilitating seasonal adaptations as in mammals (Helfrich-Förster, 2009, Yoshii *et al.*, 2012). Our data may also explain how the morning and evening oscillators in flies internally desynchronize under certain conditions (Helfrich-Förster, 2014). One such relevant condition is increased PDF signaling during long days (Hidalgo *et al.*, 2023), which was shown to delay the evening oscillators (Liang *et al.*, 2016, Liang *et al.*, 2017, Vaze and Helfrich-Förster, 2021) and may lead to internal desynchronization (Wulbeck *et al.*, 2008). Here, we confirm the presence of PDFR in the evening neurons. Thus, enhanced PDF signaling could delay the

Figure 8: Clock input and output pathways in *Drosophila* and vertebrates. Direct and indirect light input pathways to the *Drosophila* clock (comprised of LN and DN) and vertebrate suprachiasmatic nucleus (SCN). *Drosophila* and vertebrate clocks utilize different neuropeptides (grey box). Output from the clock to downstream targets is either synaptic (direct in red and indirect in orange) or paracrine (grey arrow). Abbreviations: CRY, Cryptochrome; MPS, Melanopsin, RHT, Retinohypothalamic tract; ipRGC, Intrinsically-photosensitive retinal ganglion cells; aMe, Accessory medulla neurons; NSC, Neurosecretory cell; AVP, Arginine vasopressin; CCK, Cholecystokinin; ENK, Met-enkephalin; GRP, Gastrin-releasing peptide; NMS, Neuromedin S; PROK2, Prokineticin 2; SST, Somatostatin; VIP, Vasoactive intestinal peptide.

evening neurons and bring them out of phase with the morning neurons, especially because the two sets of neurons are not connected via synapses. Taken together, the interconnectivity between the clock neurons, or lack thereof, results in a robust network which can adapt to different seasons and contexts.

Light and other inputs to Drosophila and vertebrate circadian clocks

Our analyses reveal that extrinsic light input from the photoreceptor cells of the compound eyes, HB eyelets, and ocelli to the clock neurons is largely indirect, with the former two transmitting light inputs via aMe neurons. This situation may appear to be different from mammals where intrinsically-photosensitive retinal ganglion cells in the retina project directly through the retinohypothalamic tract onto the SCN neurons (Figure 8) (Starnes and Jones, 2023). However, even the mammalian clock receives indirect photoreceptor inputs from rods and cones via bipolar cells and retinal ganglion cells. Furthermore, indirect light inputs may reach the SCN also via the intergeniculate leaflets of the thalamus (Kim and Harrington, 2008). Altogether, this suggests that the fly and mammalian system are not fundamentally different. One apparent difference is the presence of CRY, a cell-autonomous circadian photoreceptor, in subsets of *Drosophila* clock neurons that is sufficient for light entrainment in eyeless mutants (Helfrich-Förster *et al.*, 2001). Mammals lack light-sensitive CRY. Instead, they possess light-sensitive melanopsin in retinal ganglion cells, and mice lacking rods and cones can still entrain to light/dark cycles due to melanopsin (Foster *et al.*,

2020). Thus, flies and mice possess several redundant and partly parallel light-input pathways to entrain their clocks. The similarity is even higher when comparing flies with vertebrates in general. For instance, fish, birds, and reptiles possess additional photoreceptors in the pineal gland, which is reminiscent of the extraretinal HB eyelets or even the ocelli of flies (Figure 8). Most importantly, vertebrates and flies use their eyes for both vision and entraining their circadian clocks, tasks that require completely different properties of light inputs. Vision requires image formation and fast neurotransmission, whereas circadian entrainment is dependent on integrating light collection over a longer time that can be at a slower rate. The connectome reveals that the number of synapses as well as the axon thickness of the neurons mediating this connectivity are very different between the two types of photoreception. Hence, they are aptly suited to perform their required functions.

Paracrine modulation of the neuroendocrine system by circadian clock

Since circadian control of organismal physiology is likely mediated via hormones, in parallel with the clock connectome, we also characterized the complete neurosecretory center of an adult *Drosophila* brain. This neurosecretory center is comprised of 80 endocrine cells, located in distinct regions of the brain and having unique neuropeptide identities. To our knowledge, this is the largest neurosecretory center connectome characterized to date. Neurosecretory connectomes of larval *Drosophila* and the marine annelid, *Platynereis dumerilii* have been established previously (Williams *et al.*, 2017, Huckesfeld *et al.*, 2021). Unfortunately, these studies did not examine the connectivity between the circadian clock and the neuroendocrine centers. Thus, it remains to be seen whether the largely indirect and paracrine signaling between the adult *Drosophila* circadian network and NSC is a phenomenon conserved across other animals. However, given the relatively slow timescales at which the circadian output needs to be propagated to downstream neurons, neuropeptides seem suited for this role.

Limitations of our approach

The synaptic connectivity reported here is likely an underestimation due to several factors: 1) some DN were not identified in the connectome, 2) only ~30% of the photoreceptors are manually-proofread, and 3) we did not explore connectivity via gap junctions 4) we generally used a connectivity threshold of >4 synapses. Preliminary expression analysis using the single-cell transcriptomes of clock neurons suggests that gap junction genes are enriched in the clock network (data not shown) and they can influence activity-rest rhythms (Ramakrishnan and Sheeba, 2021). It remains to be seen which clock neurons are additionally coupled via gap junctions and how this electrical connectivity complements synaptic and peptidergic connectivity detailed here. Moreover, as discussed earlier, fewer than 5 synapses could also represent functional connections which were largely disregarded in our analyses. While the FlyWire and hemibrain connectomes exhibit a high degree of stereotypy, they are both based on an adult female brain. The lack of a male brain connectome currently prevents any comparisons on sex-specific differences within the circadian network and its output pathways which could influence sexually-dimorphic behaviors and physiology. Lastly, the connectome provides a singular snapshot of connectivity which could change depending on the time of day, age of the animal as well as its internal state.

Conclusion

In conclusion, our circadian clock connectome, the first of its magnitude, is a significant milestone in chronobiology. Given the high conservation of circadian network motifs between *Drosophila* and vertebrates, this connectome provides the framework to systematically investigate circadian dysregulation which is linked to various health issues in humans including sleep, metabolic, and mood disorders. Moreover, it will also facilitate development and experimental validation of novel hypotheses on clock function.

References

Abruzzi, K. C., A. Zadina, W. Luo, E. Wiyanto, R. Rahman, F. Guo, O. Shafer and M. Rosbash (2017). RNA-seq analysis of *Drosophila* clock and non-clock neurons reveals neuron-specific cycling and novel candidate neuropeptides. *PLoS Genet* **13**(2): e1006613. PMC5325595.

Alejevski, F., A. Saint-Charles, C. Michard-Vanhee, B. Martin, S. Galant, D. Vasiliauskas and F. Rouyer (2019). The HisCl1 histamine receptor acts in photoreceptors to synchronize *Drosophila* behavioral rhythms with light-dark cycles. *Nat Commun* **10**(1): 252. PMC6335465.

Alpert, M. H., H. Gil, A. Para and M. Gallio (2022). A thermometer circuit for hot temperature adjusts *Drosophila* behavior to persistent heat. *Curr Biol* **32**(18): 4079-4087 e4074. PMC9529852.

Barber, A. F., S. Y. Fong, A. Kolesnik, M. Fetchko and A. Sehgal (2021). *Drosophila* clock cells use multiple mechanisms to transmit time-of-day signals in the brain. *Proc Natl Acad Sci U S A* **118**(10). PMC7958456.

Bates, A. S., J. D. Manton, S. R. Jagannathan, M. Costa, P. Schlegel, T. Rohlfing and G. S. Jefferis (2020). The natverse, a versatile toolbox for combining and analysing neuroanatomical data. *Elife* **9**. PMC7242028.

Beer, K. and C. Helfrich-Förster (2020). Model and Non-model Insects in Chronobiology. *Front Behav Neurosci* **14**: 601676. PMC7732648.

Bell-Pedersen, D., V. M. Cassone, D. J. Earnest, S. S. Golden, P. E. Hardin, T. L. Thomas and M. J. Zoran (2005). Circadian rhythms from multiple oscillators: lessons from diverse organisms. *Nat Rev Genet* **6**(7): 544-556. PMC2735866.

Cannell, E., A. J. Dornan, K. A. Halberg, S. Terhzaz, J. A. T. Dow and S. A. Davies (2016). The corticotropin-releasing factor-like diuretic hormone 44 (DH44) and kinin neuropeptides modulate desiccation and starvation tolerance in *Drosophila melanogaster*. *Peptides* **80**: 96-107. PMC4889782.

Cavanaugh, D. J., J. D. Geratowski, J. R. Wooltorton, J. M. Spaethling, C. E. Hector, X. Zheng, E. C. Johnson, J. H. Eberwine and A. Sehgal (2014). Identification of a circadian output circuit for rest:activity rhythms in *Drosophila*. *Cell* **157**(3): 689-701. PMC4003459.

Ceriani, M. F., T. K. Darlington, D. Staknis, P. Mas, A. A. Petti, C. J. Weitz and S. A. Kay (1999). Light-dependent sequestration of TIMELESS by CRYPTOCHROME. *Science* **285**(5427): 553-556.

Chen, D. M., J. S. Christianson, R. J. Sapp and W. S. Stark (1992). Visual receptor cycle in normal and period mutant *Drosophila*: microspectrophotometry, electrophysiology, and ultrastructural morphometry. *Vis Neurosci* **9**(2): 125-135.

Chouhan, N. S., R. Wolf, C. Helfrich-Förster and M. Heisenberg (2015). Flies remember the time of day. *Curr Biol* **25**(12): 1619-1624.

Davie, K., J. Janssens, D. Koldere, M. De Waegeneer, U. Pech, L. Kreft, S. Aibar, S. Makhzami, V. Christiaens, C. Bravo Gonzalez-Blas, S. Poovathingal, G. Hulselmans, K. I. Spanier, T. Moerman, B. Vanspauwen, S. Geurs, T. Voet, J. Lammertyn, B. Thienpont, S. Liu, N. Konstantinides, M. Fiers, P. Verstreken and S. Aerts (2018). A Single-Cell Transcriptome Atlas of the Aging *Drosophila* Brain. *Cell* **174**(4): 982-998 e920. PMC6086935.

Delventhal, R., R. M. O'Connor, M. M. Pantalia, M. Ulgherait, H. X. Kim, M. K. Basturk, J. C. Canman and M. Shirasu-Hiza (2019). Dissection of central clock function in *Drosophila* through cell-specific CRISPR-mediated clock gene disruption. *Elife* **8**. PMC6794090.

Deng, B., Q. Li, X. Liu, Y. Cao, B. Li, Y. Qian, R. Xu, R. Mao, E. Zhou, W. Zhang, J. Huang and Y. Rao (2019). Chemoconnectomics: Mapping Chemical Transmission in *Drosophila*. *Neuron* **101**(5): 876-893 e874.

Diaz, M. M., M. Schlichting, K. C. Abruzzi, X. Long and M. Rosbash (2019). Allatostatin-C/AstC-R2 Is a Novel Pathway to Modulate the Circadian Activity Pattern in *Drosophila*. *Curr Biol* **29**(1): 13-22 e13. PMC6325008.

Dorkenwald, S., A. Matsliah, A. R. Sterling, P. Schlegel, S.-c. Yu, C. E. McKellar, A. Lin, M. Costa, K. Eichler, Y. Yin, W. Silversmith, C. Schneider-Mizell, C. S. Jordan, D. Brittain, A. Halageri, K. Kuehner, O. Ogedengbe, R. Morey, J. Gager, K. Kruk, E. Perlman, R. Yang, D. Deutsch, D. Bland, M. Sorek, R. Lu, T. Macrina, K. Lee, J. A. Bae, S. Mu, B. Nehoran, E. Mitchell, S. Popovych, J. Wu, Z. Jia, M. Castro, N. Kemnitz, D. Ih, A. S. Bates, N. Eckstein, J. Funke, F. Collman, D. D. Bock, G. S. X. E. Jefferis, H. S. Seung, M. Murthy and t. F. Consortium (2023). Neuronal wiring diagram of an adult brain. *bioRxiv*: 2023.2006.2027.546656.

Dorkenwald, S., C. E. McKellar, T. Macrina, N. Kemnitz, K. Lee, R. Lu, J. Wu, S. Popovych, E. Mitchell, B. Nehoran, Z. Jia, J. A. Bae, S. Mu, D. Ih, M. Castro, O. Ogedengbe, A. Halageri, K. Kuehner, A. R. Sterling, Z. Ashwood, J. Zung, D. Brittain, F. Collman, C. Schneider-Mizell, C. Jordan, W. Silversmith, C. Baker, D. Deutsch, L. Encarnacion-Rivera, S. Kumar, A. Burke, D. Bland, J. Gager, J. Hebditch, S. Koolman, M. Moore, S. Morejohn, B. Silverman, K. Willie, R. Willie, S. C. Yu, M. Murthy and H. S. Seung (2022). FlyWire: online community for whole-brain connectomics. *Nat Methods* **19**(1): 119-128. PMC8903166.

Dubowy, C. and A. Sehgal (2017). Circadian Rhythms and Sleep in Drosophila melanogaster. *Genetics* **205**(4): 1373-1397. PMC5378101.

Emery, P., R. Stanewsky, C. Helfrich-Förster, M. Emery-Le, J. C. Hall and M. Rosbash (2000). *Drosophila* CRY is a deep brain circadian photoreceptor. *Neuron* **26**(2): 493-504.

Fernandez, M. P., J. Berni and M. F. Ceriani (2008). Circadian remodeling of neuronal circuits involved in rhythmic behavior. *PLoS Biol* **6**(3): e69. PMC2270325.

Flyer-Adams, J. G., E. J. Rivera-Rodriguez, J. Yu, J. D. Mardovin, M. L. Reed and L. C. Griffith (2020). Regulation of Olfactory Associative Memory by the Circadian Clock Output Signal Pigment-Dispersing Factor (PDF). *J Neurosci* **40**(47): 9066-9077. PMC7673005.

Foster, R. G., S. Hughes and S. N. Peirson (2020). Circadian Photoentrainment in Mice and Humans. *Biology (Basel)* **9**(7). PMC7408241.

Fropf, R., H. Zhou and J. C. P. Yin (2018). The clock gene period differentially regulates sleep and memory in Drosophila. *Neurobiol Learn Mem* **153**(Pt A): 2-12. PMC6064670.

Fujii, S., A. Yavuz, J. Slone, C. Jagge, X. Song and H. Amrein (2015). *Drosophila* sugar receptors in sweet taste perception, olfaction, and internal nutrient sensing. *Curr Biol* **25**(5): 621-627. PMC4711800.

Gu, Z., L. Gu, R. Eils, M. Schlesner and B. Brors (2014). circlize Implements and enhances circular visualization in R. *Bioinformatics* **30**(19): 2811-2812.

Hadjieconomou, D., G. King, P. Gaspar, A. Mineo, L. Blackie, T. Ameku, C. Studd, A. de Mendoza, F. Diao, B. H. White, A. E. X. Brown, P. Y. Placais, T. Preat and I. Miguel-Aliaga (2020). Enteric neurons increase maternal food intake during reproduction. *Nature* **587**(7834): 455-459. PMC7610780.

Hao, Y., S. Hao, E. Andersen-Nissen, W. M. Mauck, 3rd, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T. Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P. Smibert and R. Satija (2021). Integrated analysis of multimodal single-cell data. *Cell* **184**(13): 3573-3587 e3529. PMC8238499.

Helfrich-Förster, C. (2003). The neuroarchitecture of the circadian clock in the brain of *Drosophila melanogaster*. *Microsc Res Tech* **62**(2): 94-102.

Helfrich-Förster, C. (2004). The circadian clock in the brain: a structural and functional comparison between mammals and insects. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **190**(8): 601-613.

Helfrich-Förster, C. (2009). Does the morning and evening oscillator model fit better for flies or mice? *J Biol Rhythms* **24**(4): 259-270.

Helfrich-Förster, C. (2014). From neurogenetic studies in the fly brain to a concept in circadian biology. *J Neurogenet* **28**(3-4): 329-347.

Helfrich-Förster, C. (2020). Light input pathways to the circadian clock of insects with an emphasis on the fruit fly *Drosophila melanogaster*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **206**(2): 259-272. PMC7069913.

Helfrich-Förster, C., C. Winter, A. Hofbauer, J. C. Hall and R. Stanewsky (2001). The circadian clock of fruit flies is blind after elimination of all known photoreceptors. *Neuron* **30**(1): 249-261.

Hentze, J. L., M. A. Carlsson, S. Kondo, D. R. Nässel and K. F. Rewitz (2015). The Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in *Drosophila*. *Sci Rep* **5**: 11680. PMC4485031.

Hidalgo, S., M. Anguiano, C. A. Tabuloc and J. C. Chiu (2023). Seasonal cues act through the circadian clock and pigment-dispersing factor to control EYES ABSENT and downstream physiological changes. *Curr Biol* **33**(4): 675-687 e675. PMC9992282.

Hofbauer, B., M. Zandawala, N. Reinhard, D. Rieger, C. Werner, J.-F. Evers and C. Wegener (2023). PDF neuropeptide signals independently of Bruchpilot-labelled active zones in daily remodelled terminals of *Drosophila* clock neurons. *bioRxiv*: 2023.2006.2020.545701.

Homberg, U. (2004). In search of the sky compass in the insect brain. *Naturwissenschaften* **91**(5): 199-208.

Huckesfeld, S., P. Schlegel, A. Miroschnikow, A. Schoofs, I. Zinke, A. N. Haubrich, C. M. Schneider-Mizell, J. W. Truman, R. D. Fetter, A. Cardona and M. J. Pankratz (2021). Unveiling the sensory and interneuronal pathways of the neuroendocrine connectome in *Drosophila*. *Elife* **10**. PMC8177888.

Hulse, B. K., H. Haberkern, R. Franconville, D. Turner-Evans, S. Y. Takemura, T. Wolff, M. Noorman, M. Dreher, C. Dan, R. Parekh, A. M. Hermundstad, G. M. Rubin and V. Jayaraman (2021). A connectome of the *Drosophila* central complex reveals network motifs suitable for flexible navigation and context-dependent action selection. *Elife* **10**. PMC9477501.

Ida, T., T. Takahashi, H. Tominaga, T. Sato, K. Kume, K. Yoshizawa-Kumagaye, H. Nishio, J. Kato, N. Murakami, M. Miyazato, K. Kangawa and M. Kojima (2011). Identification of the endogenous cysteine-rich peptide trissin, a ligand for an orphan G protein-coupled receptor in *Drosophila*. *Biochem Biophys Res Commun* **414**(1): 44-48.

Johnson, E. C., O. T. Shafer, J. S. Trigg, J. Park, D. A. Schooley, J. A. Dow and P. H. Taghert (2005). A novel diuretic hormone receptor in *Drosophila*:

evidence for conservation of CGRP signaling. *J Exp Biol* **208**(Pt 7): 1239-1246.

Johnstone, P. S., M. Ogueta, O. Akay, I. Top, S. Syed, R. Stanewsky and D. Top (2022). Real time, in vivo measurement of neuronal and peripheral clocks in *Drosophila melanogaster*. *Elife* **11**. PMC9662830.

Kahsai, L., N. Kapan, H. Dircksen, A. M. Winther and D. R. Nässel (2010). Metabolic stress responses in *Drosophila* are modulated by brain neurosecretory cells that produce multiple neuropeptides. *PLoS One* **5**(7): e11480. PMC2900207.

Kaneko, M. and J. C. Hall (2000). Neuroanatomy of cells expressing clock genes in *Drosophila*: transgenic manipulation of the *period* and *timeless* genes to mark the perikarya of circadian pacemaker neurons and their projections. *J Comp Neurol* **422**(1): 66-94.

Kapan, N., O. V. Lushchak, J. Luo and D. R. Nässel (2012). Identified peptidergic neurons in the *Drosophila* brain regulate insulin-producing cells, stress responses and metabolism by coexpressed short neuropeptide F and corazonin. *Cell Mol Life Sci* **69**(23): 4051-4066.

Kim, H. J. and M. E. Harrington (2008). Neuropeptide Y-deficient mice show altered circadian response to simulated natural photoperiod. *Brain Res* **1246**: 96-100. PMC2673519.

Kind, E., K. D. Longden, A. Nern, A. Zhao, G. Sancer, M. A. Flynn, C. W. Laughland, B. Gezahegn, H. D. Ludwig, A. G. Thomson, T. Obrusnik, P. G. Alarcon, H. Dionne, D. D. Bock, G. M. Rubin, M. B. Reiser and M. F. Wernet (2021). Synaptic targets of photoreceptors specialized to detect color and skylight polarization in *Drosophila*. *Elife* **10**. PMC8789284.

King, A. N., A. F. Barber, A. E. Smith, A. P. Dreyer, D. Sitaraman, M. N. Nitabach, D. J. Cavanaugh and A. Sehgal (2017). A Peptidergic Circuit Links the Circadian Clock to Locomotor Activity. *Curr Biol* **27**(13): 1915-1927 e1915. PMC5698909.

King, A. N. and A. Sehgal (2020). Molecular and circuit mechanisms mediating circadian clock output in the *Drosophila* brain. *Eur J Neurosci* **51**(1): 268-281. PMC6353709.

Kondo, S., T. Takahashi, N. Yamagata, Y. Imanishi, H. Katow, S. Hiramatsu, K. Lynn, A. Abe, A. Kumaraswamy and H. Tanimoto (2020). Neurochemical Organization of the *Drosophila* Brain Visualized by Endogenously Tagged Neurotransmitter Receptors. *Cell Rep* **30**(1): 284-297 e285.

Kurogi, Y., E. Imura, Y. Mizuno, R. Hoshino, M. Nouzova, S. Matsuyama, A. Mizoguchi, S. Kondo, H. Tanimoto, F. G. Noriega and R. Niwa (2023). Female reproductive dormancy in *Drosophila* is regulated by DH31-producing neurons projecting into the corpus allatum. *Development* **150**(10). PMC10233717.

Lamaze, A., P. Kratschmer, K. F. Chen, S. Lowe and J. E. C. Jepson (2018). A Wake-Promoting Circadian Output Circuit in *Drosophila*. *Curr Biol* **28**(19): 3098-3105 e3093.

Larsen, M. J., K. J. Burton, M. R. Zantello, V. G. Smith, D. L. Lowery and T. M. Kubiak (2001). Type A allatostatins from *Drosophila melanogaster* and *Diplotera puncata* activate two Drosophila allatostatin receptors, DAR-1 and DAR-2, expressed in CHO cells. *Biochem Biophys Res Commun* **286**(5): 895-901.

Li, H., J. Janssens, M. De Waegeneer, S. S. Kolluru, K. Davie, V. Gardeux, W. Saelens, F. P. A. David, M. Brbic, K. Spanier, J. Leskovec, C. N. McLaughlin, Q. Xie, R. C. Jones, K. Brueckner, J. Shim, S. G. Tattikota, F. Schnorrer, K. Rust, T. G. Nystul, Z. Carvalho-Santos, C. Ribeiro, S. Pal, S. Mahadevaraju, T. M. Przytycka, A. M. Allen, S. F. Goodwin, C. W. Berry, M. T. Fuller, H. White-Cooper, E. L. Matunis, S. DiNardo, A. Galenza, L. E. O'Brien, J. A. T. Dow, F. C. A. C. s. sign, H. Jasper, B. Oliver, N. Perrimon, B. Deplancke, S. R. Quake, L. Luo, S. Aerts, D. Agarwal, Y. Ahmed-Braimah, M. Arbeitman, M. M. Ariss, J. Augsburger, K. Ayush, C. C. Baker, T. Banisch, K. Birker, R. Bodmer, B. Bolival, S. E. Brantley, J. A. Brill, N. C. Brown, N. A. Buehner, X. T. Cai, R. Cardoso-Figueiredo, F. Casares, A. Chang, T. R. Clandinin, S. Crasta, C. Desplan, A. M. Detweiler, D. B.

Dhakan, E. Dona, S. Engert, S. Floc'hlay, N. George, A. J. Gonzalez-Segarra, A. K. Groves, S. Gumbin, Y. Guo, D. E. Harris, Y. Heifetz, S. L. Holtz, F. Horns, B. Hudry, R. J. Hung, Y. N. Jan, J. S. Jaszczak, G. Jefferis, J. Karkanias, T. L. Karr, N. S. Katheder, J. Kezos, A. A. Kim, S. K. Kim, L. Kockel, N. Konstantinides, T. B. Kornberg, H. M. Krause, A. T. Labott, M. Laturney, R. Lehmann, S. Leinwand, J. Li, J. S. S. Li, K. Li, K. Li, L. Li, T. Li, M. Litovchenko, H. H. Liu, Y. Liu, T. C. Lu, J. Manning, A. Mase, M. Matera-Vatnick, N. R. Matias, C. E. McDonough-Goldstein, A. McGeever, A. D. McLachlan, P. Moreno-Roman, N. Neff, M. Neville, S. Ngo, T. Nielsen, C. E. O'Brien, D. Osumi-Sutherland, M. N. Ozel, I. Papatheodorou, M. Petkovic, C. Pilgrim, A. O. Pisco, C. Reisenman, E. N. Sanders, G. Dos Santos, K. Scott, A. Sherlekar, P. Shiu, D. Sims, R. V. Sit, M. Slaidina, H. E. Smith, G. Sterne, Y. H. Su, D. Sutton, M. Tamayo, M. Tan, I. Tastekin, C. Treiber, D. Vacek, G. Vogler, S. Waddell, W. Wang, R. I. Wilson, M. F. Wolfner, Y. E. Wong, A. Xie, J. Xu, S. Yamamoto, J. Yan, Z. Yao, K. Yoda, R. Zhu and R. P. Zinzen (2022). Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science* **375**(6584): eabk2432. PMC8944923.

Li, M. T., L. H. Cao, N. Xiao, M. Tang, B. Deng, T. Yang, T. Yoshii and D. G. Luo (2018). Hub-organized parallel circuits of central circadian pacemaker neurons for visual photoentrainment in *Drosophila*. *Nat Commun* **9**(1): 4247. PMC6185921.

Liang, X., M. C. W. Ho, Y. Zhang, Y. Li, M. N. Wu, T. E. Holy and P. H. Taghert (2019). Morning and Evening Circadian Pacemakers Independently Drive Premotor Centers via a Specific Dopamine Relay. *Neuron* **102**(4): 843-857 e844. PMC6533154.

Liang, X., T. E. Holy and P. H. Taghert (2016). Synchronous *Drosophila* circadian pacemakers display nonsynchronous Ca(2)(+) rhythms *in vivo*. *Science* **351**(6276): 976-981. PMC4836443.

Liang, X., T. E. Holy and P. H. Taghert (2017). A Series of Suppressive Signals within the *Drosophila* Circadian Neural Circuit Generates Sequential Daily Outputs. *Neuron* **94**(6): 1173-1189 e1174. PMC5502710.

Liu, A. C., D. K. Welsh, C. H. Ko, H. G. Tran, E. E. Zhang, A. A. Priest, E. D. Buhr, O. Singer, K. Meeker, I. M. Verma, F. J. Doyle, 3rd, J. S. Takahashi and S. A. Kay (2007). Intercellular coupling confers robustness against mutations in the SCN circadian clock network. *Cell* **129**(3): 605-616. PMC3749832.

Ma, D., N. Herndon, J. Q. Le, K. C. Abruzzi, K. Zinn and M. Rosbash (2023). Neural connectivity molecules best identify the heterogeneous clock and dopaminergic cell types in the *Drosophila* adult brain. *Sci Adv* **9**(8): eade8500. PMC9946362.

Ma, D., D. Przybylski, K. C. Abruzzi, M. Schlichting, Q. Li, X. Long and M. Rosbash (2021). A transcriptomic taxonomy of *Drosophila* circadian neurons around the clock. *Elife* **10**. PMC7837698.

Manoli, G., M. Zandawala, T. Yoshii and C. Helfrich-Förster (2023). Characterization of clock-related proteins and neuropeptides in *Drosophila littoralis* and their putative role in diapause. *J Comp Neurol*.

Mathejczyk, T. F. and M. F. Wernet (2019). Heading choices of flying *Drosophila* under changing angles of polarized light. *Sci Rep* **9**(1): 16773. PMC6856357.

Maywood, E. S., A. B. Reddy, G. K. Wong, J. S. O'Neill, J. A. O'Brien, D. G. McMahon, A. J. Harmar, H. Okamura and M. H. Hastings (2006). Synchronization and maintenance of timekeeping in suprachiasmatic circadian clock cells by neuropeptidergic signaling. *Curr Biol* **16**(6): 599-605.

McKinney, W. (2010). <u>Data Structures for Statistical Computing in Python</u>. SciPy.

Meissner, G. W., A. Nern, Z. Dorman, G. M. DePasquale, K. Forster, T. Gibney, J. H. Hausenfluck, Y. He, N. A. Iyer, J. Jeter, L. Johnson, R. M. Johnston, K. Lee, B. Melton, B. Yarbrough, C. T. Zugates, J. Clements, C. Goina, H. Otsuna, K. Rokicki, R. R. Svirskas, Y. Aso, G. M. Card, B. J. Dickson, E. Ehrhardt, J. Goldammer, M. Ito, D. Kainmueller, W. Korff, L. Mais, R. Minegishi, S. Namiki, G. M. Rubin, G. R. Sterne, T. Wolff, O. Malkesman and T. FlyLight Project (2023). A searchable image resource of

Drosophila GAL4 driver expression patterns with single neuron resolution. *Elife* **12**. PMC10030108.

Mertens, I., A. Vandingenen, E. C. Johnson, O. T. Shafer, W. Li, J. S. Trigg, A. De Loof, L. Schoofs and P. H. Taghert (2005). PDF receptor signaling in *Drosophila* contributes to both circadian and geotactic behaviors. *Neuron* **48**(2): 213-219.

Miyamoto, T. and H. Amrein (2014). Diverse roles for the *Drosophila* fructose sensor Gr43a. *Fly (Austin)* **8**(1): 19-25. PMC3974889.

Mohawk, J. A., C. B. Green and J. S. Takahashi (2012). Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci* **35**: 445-462. PMC3710582.

Morris, E. L., A. P. Patton, J. E. Chesham, A. Crisp, A. Adamson and M. H. Hastings (2021). Single-cell transcriptomics of suprachiasmatic nuclei reveal a Prokineticin-driven circadian network. *EMBO J* **40**(20): e108614. PMC8521297.

Nagy, D., P. Cusumano, G. Andreatta, A. M. Anduaga, C. Hermann-Luibl, N. Reinhard, J. Gesto, C. Wegener, G. Mazzotta, E. Rosato, C. P. Kyriacou, C. Helfrich-Förster and R. Costa (2019). Peptidergic signaling from clock neurons regulates reproductive dormancy in *Drosophila melanogaster*. *PLoS Genet* **15**(6): e1008158. PMC6592559.

Nässel, D. R. and J. Vanden Broeck (2016). Insulin/IGF signaling in *Drosophila* and other insects: factors that regulate production, release and post-release action of the insulin-like peptides. *Cell Mol Life Sci* **73**(2): 271-290.

Nässel, D. R. and M. Zandawala (2019). Recent advances in neuropeptide signaling in *Drosophila*, from genes to physiology and behavior. *Prog Neurobiol* **179**: 101607.

Nässel, D. R. and M. Zandawala (2020). Hormonal axes in *Drosophila*: regulation of hormone release and multiplicity of actions. *Cell Tissue Res* **382**(2): 233-266. PMC7584566.

Nässel, D. R. and M. Zandawala (2022). Endocrine cybernetics: neuropeptides as molecular switches in behavioural decisions. *Open Biol* **12**(7): 220174. PMC9326288.

Nitabach, M. N. and P. H. Taghert (2008). Organization of the *Drosophila* circadian control circuit. *Curr Biol* **18**(2): R84-93.

Oh, Y., J. S. Lai, H. J. Mills, H. Erdjument-Bromage, B. Giammarinaro, K. Saadipour, J. G. Wang, F. Abu, T. A. Neubert and G. S. B. Suh (2019). A glucose-sensing neuron pair regulates insulin and glucagon in *Drosophila*. *Nature* **574**(7779): 559-564. PMC6857815.

Ohhara, Y., S. Kobayashi, K. Yamakawa-Kobayashi and N. Yamanaka (2018). Adult-specific insulin-producing neurons in *Drosophila melanogaster*. *J Comp Neurol* **526**(8): 1351-1367.

Panda, S., J. B. Hogenesch and S. A. Kay (2002). Circadian rhythms from flies to human. *Nature* **417**(6886): 329-335.

Pyza, E. and I. A. Meinertzhagen (1997). Circadian rhythms in screening pigment and invaginating organelles in photoreceptor terminals of the housefly's first optic neuropile. *J Neurobiol* **32**(5): 517-529.

Ramakrishnan, A. and V. Sheeba (2021). Gap junction protein Innexin2 modulates the period of free-running rhythms in *Drosophila melanogaster*. *iScience* **24**(9): 103011. PMC8426565.

Reinhard, N., E. Bertolini, A. Saito, M. Sekiguchi, T. Yoshii, D. Rieger and C. Helfrich-Förster (2022a). The lateral posterior clock neurons of *Drosophila melanogaster* express three neuropeptides and have multiple connections within the circadian clock network and beyond. *J Comp Neurol* **530**(9): 1507-1529.

Reinhard, N., F. K. Schubert, E. Bertolini, N. Hagedorn, G. Manoli, M. Sekiguchi, T. Yoshii, D. Rieger and C. Helfrich-Förster (2022b). The Neuronal Circuit of the Dorsal Circadian Clock Neurons in *Drosophila melanogaster*. *Front Physiol* **13**: 886432. PMC9100938.

Rieger, D., O. T. Shafer, K. Tomioka and C. Helfrich-Förster (2006). Functional analysis of circadian pacemaker neurons in *Drosophila melanogaster*. *J Neurosci* **26**(9): 2531-2543. PMC6793667.

Saini, R., M. Jaskolski and S. J. Davis (2019). Circadian oscillator proteins across the kingdoms of life: structural aspects. *BMC Biol* **17**(1): 13. PMC6378743.

Scheffer, L. K., C. S. Xu, M. Januszewski, Z. Lu, S. Y. Takemura, K. J. Hayworth, G. B. Huang, K. Shinomiya, J. Maitlin-Shepard, S. Berg, J. Clements, P. M. Hubbard, W. T. Katz, L. Umayam, T. Zhao, D. Ackerman, T. Blakely, J. Bogovic, T. Dolafi, D. Kainmueller, T. Kawase, K. A. Khairy, L. Leavitt, P. H. Li, L. Lindsey, N. Neubarth, D. J. Olbris, H. Otsuna, E. T. Trautman, M. Ito, A. S. Bates, J. Goldammer, T. Wolff, R. Svirskas, P. Schlegel, E. Neace, C. J. Knecht, C. X. Alvarado, D. A. Bailey, S. Ballinger, J. A. Borycz, B. S. Canino, N. Cheatham, M. Cook, M. Dreher, O. Duclos, B. Eubanks, K. Fairbanks, S. Finley, N. Forknall, A. Francis, G. P. Hopkins, E. M. Joyce, S. Kim, N. A. Kirk, J. Kovalyak, S. A. Lauchie, A. Lohff, C. Maldonado, E. A. Manley, S. McLin, C. Mooney, M. Ndama, O. Ogundeyi, N. Okeoma, C. Ordish, N. Padilla, C. M. Patrick, T. Paterson, E. E. Phillips, E. M. Phillips, N. Rampally, C. Ribeiro, M. K. Robertson, J. T. Rymer, S. M. Ryan, M. Sammons, A. K. Scott, A. L. Scott, A. Shinomiya, C. Smith, K. Smith, N. L. Smith, M. A. Sobeski, A. Suleiman, J. Swift, S. Takemura, I. Talebi, D. Tarnogorska, E. Tenshaw, T. Tokhi, J. J. Walsh, T. Yang, J. A. Horne, F. Li, R. Parekh, P. K. Rivlin, V. Jayaraman, M. Costa, G. S. Jefferis, K. Ito, S. Saalfeld, R. George, I. A. Meinertzhagen, G. M. Rubin, H. F. Hess, V. Jain and S. M. Plaza (2020). A connectome and analysis of the adult *Drosophila* central brain. *Elife* **9**. PMC7546738.

Schlegel, P., Y. Yin, A. S. Bates, S. Dorkenwald, K. Eichler, P. Brooks, D. S. Han, M. Gkantia, M. d. Santos, E. J. Munnelly, G. Badalamente, L. S. Capdevila, V. A. Sane, M. W. Pleijzier, I. F. M. Tamimi, C. R. Dunne, I. Salgarella, A. Javier, S. Fang, E. Perlman, T. Kazimiers, S. R. Jagannathan, A. Matsliah, A. R. Sterling, S.-c. Yu, C. E. McKellar, F. Consortium, M. Costa, H. S. Seung, M. Murthy, V. Hartenstein, D. D. Bock and G. S. X. E. Jefferis (2023). Whole-brain annotation and multi-connectome cell typing quantifies circuit stereotypy in Drosophila. *bioRxiv*: 2023.2006.2027.546055.

Schlichting, M., P. Menegazzi, M. Rosbash and C. Helfrich-Förster (2019). A Distinct Visual Pathway Mediates High-Intensity Light Adaptation of the Circadian Clock in *Drosophila*. *J Neurosci* **39**(9): 1621-1630. PMC6391560.

Schubert, F. K., N. Hagedorn, T. Yoshii, C. Helfrich-Förster and D. Rieger (2018). Neuroanatomical details of the lateral neurons of *Drosophila melanogaster* support their functional role in the circadian system. *J Comp Neurol* **526**(7): 1209-1231. PMC5873451.

Selcho, M., B. Muhlbauer, R. Hensgen, S. Shiga, C. Wegener and K. Yasuyama (2018). Anatomical characterization of PDF-tri neurons and peptidergic neurons associated with eclosion behavior in *Drosophila*. *J Comp Neurol* **526**(8): 1307-1328.

Shafer, O. T., G. J. Gutierrez, K. Li, A. Mildenhall, D. Spira, J. Marty, A. A. Lazar and M. P. Fernandez (2022). Connectomic analysis of the *Drosophila* lateral neuron clock cells reveals the synaptic basis of functional pacemaker classes. *Elife* **11**. PMC9365390.

Snell, N. J., J. D. Fisher, G. G. Hartmann, B. Zolyomi, M. Talay and G. Barnea (2022). Complex representation of taste quality by second-order gustatory neurons in *Drosophila*. *Curr Biol* **32**(17): 3758-3772 e3754. PMC9474709.

Song, B. J., S. J. Sharp and D. Rogulja (2021). Daily rewiring of a neural circuit generates a predictive model of environmental light. *Sci Adv* **7**(13). PMC7990339.

Starnes, A. N. and J. R. Jones (2023). Inputs and Outputs of the Mammalian Circadian Clock. *Biology (Basel)* **12**(4). PMC10136320.

Stengl, M. and A. Arendt (2016). Peptidergic circadian clock circuits in the Madeira cockroach. *Curr Opin Neurobiol* **41**: 44-52.

Sun, L., R. H. Jiang, W. J. Ye, M. Rosbash and F. Guo (2022). Recurrent circadian circuitry regulates central brain activity to maintain sleep. *Neuron* **110**(13): 2139-2154 e2135.

Takahashi, J. S. (2017). Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet* **18**(3): 164-179. PMC5501165.

Talay, M., E. B. Richman, N. J. Snell, G. G. Hartmann, J. D. Fisher, A. Sorkac, J. F. Santoyo, C. Chou-Freed, N. Nair, M. Johnson, J. R. Szymanski and G. Barnea (2017). Transsynaptic Mapping of Second-Order Taste Neurons in Flies by *trans*-Tango. *Neuron* **96**(4): 783-795 e784. PMC5693608.

Tang, M., L. H. Cao, T. Yang, S. X. Ma, B. Y. Jing, N. Xiao, S. Xu, K. R. Leng, D. Yang, M. T. Li and D. G. Luo (2022). An extra-clock ultradian brain oscillator sustains circadian timekeeping. *Sci Adv* **8**(35): eabo5506.

Vaze, K. M. and C. Helfrich-Förster (2021). The Neuropeptide PDF Is Crucial for Delaying the Phase of *Drosophila's* Evening Neurons Under Long Zeitgeber Periods. *J Biol Rhythms* **36**(5): 442-460. PMC8442139.

Warren, T. L., Y. M. Giraldo and M. H. Dickinson (2019). Celestial navigation in *Drosophila*. *J Exp Biol* **222**(Pt Suppl 1). PMC7375828.

Wen, S., D. Ma, M. Zhao, L. Xie, Q. Wu, L. Gou, C. Zhu, Y. Fan, H. Wang and J. Yan (2020). Spatiotemporal single-cell analysis of gene expression in the mouse suprachiasmatic nucleus. *Nat Neurosci* **23**(3): 456-467.

Williams, E. A., C. Veraszto, S. Jasek, M. Conzelmann, R. Shahidi, P. Bauknecht, O. Mirabeau and G. Jekely (2017). Synaptic and peptidergic connectome of a neurosecretory center in the annelid brain. *Elife* **6**. PMC5747525.

Wulbeck, C., E. Grieshaber and C. Helfrich-Förster (2008). Pigment-dispersing factor (PDF) has different effects on *Drosophila's* circadian clocks in the accessory medulla and in the dorsal brain. *J Biol Rhythms* **23**(5): 409-424.

Yang, Z., R. Huang, X. Fu, G. Wang, W. Qi, D. Mao, Z. Shi, W. L. Shen and L. Wang (2018). A post-ingestive amino acid sensor promotes food consumption in *Drosophila*. *Cell Res* **28**(10): 1013-1025. PMC6170445.

Yoshii, T., C. Hermann-Luibl, C. Kistenpfennig, B. Schmid, K. Tomioka and C. Helfrich-Förster (2015). Cryptochrome-dependent and -independent circadian entrainment circuits in *Drosophila*. *J Neurosci* **35**(15): 6131-6141. PMC6605168.

Yoshii, T., D. Rieger and C. Helfrich-Förster (2012). Two clocks in the brain: an update of the morning and evening oscillator model in *Drosophila*. *Prog Brain Res* **199**: 59-82.

Zandawala, M., T. Nguyen, M. Balanya Segura, H. A. D. Johard, M. Amcoff, C. Wegener, J. P. Paluzzi and D. R. Nässel (2021). A neuroendocrine pathway modulating osmotic stress in *Drosophila*. *PLoS Genet* **17**(3): e1009425. PMC7971876.

Zandawala, M., M. E. Yurgel, M. J. Texada, S. Liao, K. F. Rewitz, A. C. Keene and D. R. Nässel (2018). Modulation of *Drosophila* post-feeding physiology and behavior by the neuropeptide leucokinin. *PLoS Genet* **14**(11): e1007767. PMC6245514.

Zerr, D. M., J. C. Hall, M. Rosbash and K. K. Siwicki (1990). Circadian fluctuations of period protein immunoreactivity in the CNS and the visual system of *Drosophila*. *J Neurosci* **10**(8): 2749-2762. PMC6570283.

Zheng, Z., J. S. Lauritzen, E. Perlman, C. G. Robinson, M. Nichols, D. Milkie, O. Torrens, J. Price, C. B. Fisher, N. Sharifi, S. A. Calle-Schuler, L. Kmecova, I. J. Ali, B. Karsh, E. T. Trautman, J. A. Bogovic, P. Hanslovsky, G. Jefferis, M. Kazhdan, K. Khairy, S. Saalfeld, R. D. Fetter and D. D. Bock (2018). A Complete Electron Microscopy Volume of the Brain of Adult *Drosophila melanogaster*. *Cell* **174**(3): 730-743 e722. PMC6063995.

Acknowledgements

We would like to thank Selina Hilpert and Barbara Mühlbauer for technical assistance, Maria Steigmeier for preliminary investigations on PDFR expression, and Tatsuya Yokosako for generating the LN[ITP] split-Gal4 line. We are also grateful to Drs. Fumika Hamada for *Clk.9M-Gal4;pdf-Gal80*, Shu Kondo for T2A-Gal4 lines, Paul E. Hardin for VRI antibody, Ralf Stanewsky for PER antibody, Heinrich Dircksen for ITP antibody, Susan Morton for RFP antibody, and Jan Veenstra for AstC, CRZ and DH44 antibodies. We thank the Princeton FlyWire team and members of the Murthy and Seung labs, as well as members of the Allen Institute for Brain Science, for development and maintenance of FlyWire (supported by BRAIN Initiative grants MH117815 and NS126935 to Murthy and Seung). We also acknowledge members of the Princeton FlyWire team and the FlyWire consortium, especially Drs. Gregory Jefferis, Sven Dorkenwald, and Philipp Schlegel for troubleshooting and guidance. Special thanks to Austin T Burke and Mareike Selcho from the FlyWire consortium for contributing >10% effort towards editing and proofreading at least 10% of clock neurons. We are also thankful to Drs. Theresa McKim and Dick Nässel for helpful feedback during preparation of this manuscript, as well as the Division of Instrumental Analysis, Okayama University for the laser scanning confocal microscopes (FV1200 and FV3000). M.Z. was supported by funding from the University of Würzburg and Deutsche Forschungsgemeinschaft (DFG; ZA1296/1-1). D.R. (DFG; RI 2411/1-1) and C.H.F. (DFG; FO 207/16-1) were supported by DFG. T.Y. was supported by JSPS (KAKENHI 19H03265). A.S. and M.S. were supported by the OU fellowship (JST SPRING, Grant Number JPMJSP2126). We also acknowledge funding from the DFG for the Leica TCS SP8 microscope (251610680, INST 93/809-1 FUGG). This manuscript was typeset with the bioRxiv word template by @Chrelli: www.github.com/chrelli/bioRxiv-word-template.

Author contributions

N.R., C.H.F., T.Y., and M.Z. conceived the study. N.R., D.R., C.H.F., T.Y., and M.Z. supervised the project. N.R., A.F., G.M., E.D., A.S., G.M., M.S., and M.Z. performed the experimental work and analyzed the data. N.R. and M.Z. performed computational analyses. M.Z. wrote the manuscript with input from C.H.F and N.R. All authors read, provided feedback, and approved the final manuscript.

Competing interest statement

We declare we have no competing interests.

Materials and Methods

Fly strains

Drosophila melanogaster strains used in this study are listed in Supplementary Table 1. T2A-GAL and T2A-LexA knock-in lines generated previously (Deng *et al.*, 2019, Kondo *et al.*, 2020) were obtained from the Bloomington *Drosophila* Stock Center (BDSC) and Dr. Shu Kondo. Flies were maintained under LD12:12. Flies for peptide and receptor mapping were reared at 25°C on *Drosophila* medium containing 0.7% agar, 8.0% glucose, 3.3% yeast, 4.0% cornmeal, 2.5% wheat embryo, and 0.25% propionic acid. Flies for *trans*-Tango and GAL4 verification were raised at 18°C and 25°C, respectively, on standard medium containing 8.0% malt extract, 8.0% corn flour, 2.2% sugar beet molasses, 1.8% yeast, 1.0% soy flour, 0.8% agar and 0.3% hydroxybenzoic acid.

Immunohistochemistry and confocal imaging

Neuropeptide and receptor mapping

Immunostainings were performed as described previously (Yoshii *et al.*, 2015). Briefly, male flies were entrained in LD12:12 at 25 °C for at least 3 days. Whole flies sampled at Zeitgeber time (ZT) 20 were fixed in 4% paraformaldehyde in phosphate-buffered saline (PBS) with 0.1% Triton X-100 (PBS-T) for 2.5 h at room temperature (RT). Fixed flies were washed three times with PBS, before dissecting their brains. Samples were washed with PBS-T for three times. The samples were blocked in PBS-T containing 5% normal donkey serum for 1 hour at RT and subsequently incubated in primary antibodies at 4°C for 48 h. Following six washes with PBS-T, the brains were incubated in secondary antibodies at RT for 3 h. Lastly, the samples were washed six times in PBS-T and mounted in Vectashield mounting medium (Vector Laboratories, Burlingame, CA, USA). At least five brains for each strain were used for immunostaining to characterize the clock cells stained by anti-GFP antibodies (GAL4 or LexA expression) in the first

experiment and the clock cells were briefly characterized with PDP1 and PDF antibodies. In the second experiment, we conducted the same immunostaining with anti-TIM antibodies, but only for positive strains, to confirm the prior results. Images were taken from at least three different brains using laser scanning confocal microscopes (Olympus FV1200 and FV3000, Olympus, Tokyo, Japan).

Mapping DH44 and AstC expression in clock neurons, verification of clock neuron GAL4 lines, and trans-Tango analysis

Flies for PER staining were fixed at ZT23 (1 hour before lights-on) since PER levels are maximal at this ZT (Zerr *et al.*, 1990). Flies for VRI staining were fixed at ZT 20. Immunohistochemistry was performed as described previously (Reinhard *et al.*, 2022a, Reinhard *et al.*, 2022b). Images were scanned with a Leica TCS SP8 confocal microscope equipped with a photon multiplier tube and hybrid detector. A white light laser (Leica Microsystems, Wetzlar, Germany) was used for excitation. We used a 20-fold glycerol immersion objective (HC PL APO, Leica Microsystems, Wetzlar Germany) for wholemount scans and obtained confocal stacks with 2048×1024 pixels with a maximal voxel size of 0.3 x 0.3 x 2 µm and an optical section thickness of 3.12 µm. For noise reduction, we used a frame average of 3. Images were analyzed using Fiji. All the primary and secondary antibodies are listed in Supplementary Table 2.

Connectome datasets and neuron identification

For all analyses, we used the v630 snapshot of the FlyWire whole brain connectome and its annotations (Dorkenwald *et al.*, 2023, Schlegel *et al.*, 2023). We also used the adult hemibrain connectome for comparisons (Scheffer et al. 2020). Neurons were identified based on morphological features described previously (Lamaze *et al.*, 2018, Schubert *et al.*, 2018, Reinhard *et al.*, 2022a, Reinhard *et al.*, 2022b, Sun *et al.*, 2022) or based on NBLAST similarity with identified neurons in the hemibrain (Schlegel *et al.*, 2023). FlyWire cell IDs of identified clock neurons and NSC are provided in Supplementary Tables 3 and 4, respectively.

Data visualization

Data was visualized using ggplot2 (v 3.4.2, Wickham, 2016) and circlize (v 0.4.15) for R (v 4.2.2) in RStudio (2022.12.0) (Gu *et al.*, 2014). Reconstructions were downloaded using the navis library (v 1.0.4, https://github.com/navis-org) and cloud-volume library (v 8.10.0, https://github.com/seung-lab/cloud-volume) for python (v 3.8.5), and visualized using blender (v 3.01, Community, B. O. 2018). For visualizing a large number of neurons, Codex (doi: 10.13140/RG.2.2.35928.67844) and FlyWire neuroglancer were used (Dorkenwald *et al.*, 2022).

Connectivity analyses

Connectivity data was analyzed using the natverse libraries (v 0.2.4) for R in RStudio (Bates *et al.*, 2020). Cosine similarity analyses were conducted with coconatfly (v 0.1.1.9; https://github.com/flyconnectome/coconatfly) for R. Neuron similarity was analyzed based on all theirs. Filtered synapses were retrieved in python using the navis and pandas (v 1.1.3, (McKinney, 2010)) libraries. Unless stated otherwise, a connection with more than 4 synapses was considered significant.

Single-cell transcriptome analyses

Expression of neuropeptides and their cognate receptors in clock neurons was mined using the single-cell RNA sequencing dataset and analysis pipeline established earlier (Ma *et al.*, 2021). NSC transcriptomes were identified from the brain and IPC transcriptomes generated previously (Davie *et al.*, 2018, Li *et al.*, 2022). The parameters used to identify the different cell types were based on previous studies and provided below (Kahsai *et al.*, 2010, Kapan *et al.*, 2012, Miyamoto and Amrein, 2014, Cannell *et al.*, 2016, Yang *et al.*, 2018, Nässel and Zandawala, 2019, Oh *et al.*, 2019).

DH44 (12 cells): Dh44 > 1 & CG13248 > 1 & CG13743 > 0 & Lkr > 0
CRZ (11 cells): Crz > 1 & sNPF > 1 & Gr43a > 0 & Gr64a > 0
ITP (7 cells): Tk > 1 & sNPF > 1 & ITP > 1 & ImpL2 > 1 & Crz = 0
IPC (392 cells): Ilp2 > 0 & Ilp3 > 0 & Ilp5 > 0

All analyses were performed in R-Studio (v2022.02.0) using the Seurat package (v4.1.1 (Hao *et al.*, 2021)).