Kang Xu[1], Haoran Zeng[1], Emi Yumoto[2], Masashi Asahina[2, 3], Ken-ichiro Hayashi[4], Hidehiro

Fukaki[5], Hisashi Ito[6], Masaaki K. Watahiki[1, 6]

[1] Graduate School of Life Science, Hokkaido University, Sapporo, Japan

[2] Advanced Instrumental Analysis Center. Teikyo University, Utsunomiya, Japan

[3] Department of Biosciences, Teikyo University, Utsunomiya, Japan

[4] Department of Bioscience, Okayama University of Science, Okayama, Japan

[5] Department of Biology, Graduate School of Science, Kobe University, Kobe, Japan

[6] Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan

[7] Division of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo, Japan

Full title:

Carotenoid metabolism negatively regulates auxin-mediated root

growth

One-sentence summary
The β-carotene-derived apocarotenoid retinal negatively regulates auxin levels and signaling
and controls root growth.

List of authors:

Kang Xu: [ORCID ID: 0000-0001-6875-8027] kyokou950817@gmail.com

Haoran Zeng: [ORCID ID: 0009-0000-6707-4221] zenghaoran1998@outlook.com

Emi Yumoto: [ORCID ID: 0009-0005-2348-8410] yumoto@nasu.bio.teikyo-u.ac.jp

Masashi Asahina: [ORCID ID: 0000-0002-1828-0966] asahina@nasu.bio.teikyo-u.ac.jp

Ken-ichiro Hayashi: [ORCID ID: 0000-0002-9812-2801] ken-hayashi@ous.ac.jp

Hidehiro Fukaki: [ORCID ID: 0000-0002-6251-7668] h-fukaki@port.kobe-u.ac.jp

Hisashi Ito: [ORCID ID: 0000-0002-2482-5870] ito98@lowtem.hokudai.ac.jp

Masaaki K. Watahiki: [ORCID ID: 0000-0001-6669-9692] watahiki@sci.hokudai.ac.jp

Address correspondence to watahiki@sci.hokudai.ac.jp

The author responsible for distribution of materials integral to the findings represented in this

article in accordance with the policy described in the Instruction for Authors

(www.plantphysiol.org) is Masaaki K. Watahiki (watahiki@sci.hokudai.ac.jp).

ResearcherID: GFY-7778-2022

Abstract

Carotenoids support plant developmental activities as photosynthesis and photoprotective pigments, they also provide precursors for production of phytohormone abscisic acid and strigolactones, or bioactive apocarotenoids. Herein, the role of carotenoids as sources of apocarotenoids, which control endogenous auxin levels and plant root architecture was examined. Inhibition of β-carotene biosynthesis with the phytoene desaturase (PDS) inhibitor fluridone suppresses root growth, consequently inducing lateral root growth. PDS was confirmed to be the target of fluridone via the expression of the fluridone-insensitive trait *35S::mHvPDS* in an *Arabidopsis* plant. Inhibiting β-carotene biosynthesis elevated endogenous auxin levels and activated auxin signaling, thereby suppressing root growth. In addition, the root growth of the auxin-deficient mutant *yucQ* was partially restored via fluridone treatment. Consequently, suppressing PDS activity increases endogenous auxin levels and suppresses root growth. Conversely, the carotenoid pool indirectly suppresses endogenous auxin levels. Exogenous application of retinal, an apocarotenoid derived from β-carotene, complemented fluridone-mediated root growth inhibition and partially recovered auxin-mediated growth inhibition. Moreover, retinal inhibited the induction of the auxin-inducible reporter *pIAA19::ElucPEST*, indicating an antagonistic effect of retinal on endogenous auxin levels. Interestingly, the auxin-deficient mutants *wei2 wei7* and *yucQ* accumulated more β-carotene compared with the wild type. The expression of *CCDs*, which converts β-carotene into apocarotenoids, is auxin-inducible and increases following fluridone treatment. These results indicate feedback regulation during apocarotenoid biosynthesis in plant tissue. Thus, we conclude carotenoid regulates auxin levels and response, and this regulation is mediated by apocarotenoid retinal.

Introduction

The root system plays a vital role in water and nutrient uptake, anchors plants to the soil, and is a source of signaling chemicals that travel from root to root and root to shoot. The phytohormone auxin participates in establishing the root system architecture, which controls

root growth and lateral root patterning (Fukaki and Tasaka, 2009; Moreno-Risueno et al.,

2010). Auxin signaling begins with the degradation of AUXIN/INDOLE-3-ACETIC ACID

(Aux/IAA) proteins through ubiquitination by the E3 ligase complex TRANSPORT

INHIBITOR RESPONSE 1/AUXIN SIGNALING F-BOX PROTEIN (SCF[TIR1/AFB])

(Dharmasiri et al., 2005). The functional repression of the transcription factor AUXIN

RESPONSE FACTOR (ARF) by Aux/IAA is inactivated incrementally by cellular auxin

levels (Leyser, 2002; Shimizu-Mitao and Kakimoto, 2014). A gain-of-function mutation of

typical *aux/iaa* exhibited insensitivity to exogenous auxin addition and was accompanied by a

lateral root (LR) defect (i.e., *massugu2/auxin/indole-3-acetic acid19* [*msg2/aux/iaa19*],

solitary root/auxin/indole-3-acetic acid14 [*slr/aux/iaa14*], and *suppressor of hy2 mutation*

2/auxin/indole-3-acetic acid3 [*shy2/aux/iaa3*]). The auxin influx carriers AUXIN

RESISTANT1 (AUX1) and LIKE AUX1 (LAX) (Swarup and Peret, 2012), auxin efflux

carrier PINFORMED (PIN) (Paponov et al., 2005), and the ATP-binding cassette B

/MULTIPLE DRUG RESISTANCE/P-GLYCOPROTEIN (ABCB/MDR/PGP) (Geisler and

Murphy, 2006) regulate the flow and cellular levels of auxin to generate an auxin gradient,

which also shapes the root architecture (Vieten et al., 2007).

The conventional pathway of IAA biosynthesis in *Arabidopsis* involves the conversion of

tryptophan to indole-3-pyruvic acid (IPyA) by TRYPTOPHAN AMINOTRANSFERASE OF

ARABIDOPSIS (TAA), followed by oxidation of IPyA to IAA by the activity of YUCCA

(YUC) flavin monooxygenases. YUCCA is the rate-limiting enzyme in this pathway and

comprises 11 gene families in *Arabidopsis* (Won et al., 2011; Zhao, 2012). *YUCCA* genes are

highly redundant (Cheng et al., 2006; Xu et al., 2017) and multiple knockouts of the gene

family members result in an auxin-deficient phenotype (Cheng et al., 2007; Chen et al., 2014).

However, there are several non-IpyA pathways of IAA biosynthesis, including the

indole-3-acetaldoxime (IAOx) pathway. The IAOx pathway generates indole-glucosinolate

(IG) that prevent herbivores and also generates IAA through indole-3-acetonitrile (IAN) or

indole-3-acetamide (IAM) (Sugawara et al., 2009). One of the cytochrome P450 gene family

members of the IAOx pathway, *SUPERROOT2/CYP83B1* (*SUR2*), oxidizes IAOx as a

substrate for IG biosynthesis. A loss-of-function mutant of *sur2* accumulates IAA and exhibits

a pleiotropic effect, including high LR density, adventitious root, long hypocotyl, and epinastic cotyledon (Delarue et al., 1998; Barlier et al., 2000; Goto et al., 2023). Chorismic acid is a precursor of tryptophan biosynthesis, where the first step is catalyzed by a heterodimer of anthranilate synthase, WEAK ETHYLENE INSENSITIVE 2/ANTHRANILATE SYNTHASE α1 (WEI2/ASA1), and WEI7/ANTHRANILATE SYNTHASE β1 (WEI7/ASB1). The loss-of-function mutant of the anthranilate synthase subunit, *wei2* or *wei7*, decreases endogenous auxin levels and moderately reduces root length (Stepanova et al., 2005).

Carotenoids are a group of plastidial isoprenoid pigments that mainly synthesized by plants, fungi, algae and bacteria (Sun et al., 2018). In flowering plants, the photopigment carotenoid is synthesized by the 2-C-methyl-D-erythritol 4-phosphate/1-deoxy-D-xylulose 5-phosphate (MEP/DOXP) pathway (Rodriguez-Concepcion and Boronat, 2002). Geranylgeranyl pyrophosphate (GGPP) is condensed to phytoene by PHYTOENE SYNTHASE (PSY), which undergoes desaturation by PHYTOENE DESATURASE (PDS) to form ζ-carotene and is subsequently converted to tetra-*cis*-lycopene by ZETA-CAROTENE DESATURASE (ZDS) (Breitenbach and Sandmann, 2005). Tetra-*cis*-lycopene is isomerized to all-*trans*-lycopene by CAROTENOID CHLOROPLAST REGULATOR2/CAROTENOID ISOMERASE (CCR2/CRTISO) and its expression is mediated by histone methyltransferase SET DOMAIN GROUP 8 (SDG8)/CCR1 (Park et al., 2002; Cazzonelli et al., 2010). Subsequently, the pathway branches to produce α- or β-carotene (**Fig. 1A**). Fluridone is a herbicide used for eliminating aquatic plants and inhibiting PDS activity, resulting in chlorosis in plants (Bartels and Watson, 1978). Carotenoids exhibit diverse functions. They act as a protective photopigment in the photosynthetic complex and membrane, add color to fruit, and are a source of bioactive apocarotenoids, which are the products of oxidative cleavage of carotenoids (Cazzonelli, 2011). Although the apocarotenoid phytohormone abscisic acid (ABA) as well as strigolactones (SLs) (Schwartz et al., 1997; Umehara et al., 2008; Zhang et al., 2010) are well characterized, there has been an increase in the number of new bioactive apocarotenoids discovered during this century (Auldridge et al., 2006; Zheng et al., 2021). For example, β-cyclocitral is recently identified as a root growth regulator and reportedly

exhibits a positive effect on primary root (PR) elongation and LR capacity (Dickinson et al.,

2019). Anchorene, a β-carotene-derived regulatory metabolite, promotes anchor root

formation by modulating auxin distribution (Jia et al., 2019). Another example is retinal, a

precursor of vitamin A and retinoic acid in vertebrates (Olson and Hayaishi, 1965) that it also

exists in higher plants (Koschmieder et al., 2021) and plays a role in the periodic branching of

the root (Dickinson et al., 2021). In addition to their role in root architecture, the

apocarotenoids ionone derivatives repel or attract plant herbivores. The apocarotenoids bixin

and safranal are also important in the food industry as color additives (Zheng et al., 2021).

The family of carotenoid cleavage dioxygenases (CCD) and 9-*cis* epoxycarotenoid

dioxygenase (NCED) are responsible for carotenoid catabolism and producing apocarotenoids.

CCD1 and CCD4 exhibit a wide range of cleavage specificities, ranging from phytoene to

neoxanthin, to generate apocarotenoids, such as bixin, crocin, safranal, α-ionone, β-ionone,

and β-cyclocitral. CCD7 and CCD8 are involved in the synthesis of SLs from β-carotene.

NCEDs are responsible for the production of ABA (Yuan et al., 2015).

Carotenoids function as photopigments working with chlorophylls in chloroplasts and are

the precursors of apocarotenoids (e.g., retinal) and phytohormones (e.g., ABA and SLs).

Auxin modulates chlorophyll content and chloroplast biogenesis in the root (Kobayashi et al.,

2012), and regulates carotenoid accumulation during fruit ripening in tomato (Su et al., 2015).

Auxin and apocarotenoid SLs can regulate the level of each other and contribute to shoot

branching (Hayward et al., 2009). Application of IAA increases ABA sensitivity to seed

germination (Liu et al., 2007). Consequently, auxin plays a vital role in regulating plastid

activities, controlling carotenoid contents and interacting with apocarotenoids for

developmental activities. However, the interaction between auxin and carotenoids remains to

be elucidated, especially how carotenoid and apocarotenoid contents affect auxin homeostasis

and control root architecture. In this study, we found that carotenoid biosynthesis inhibition

resulted in auxin level elevation and response induction in the root. Apocarotenoid retinal

rescued carotenoid-mediated and auxin-mediated root growth suppression, suggesting an

intermediate position of retinal in carotenoid and auxin interaction pathway. Herein, we

demonstrate that carotenoid biosynthesis negatively couples with auxin biosynthesis and that

159   the exogenous apocarotenoid retinal may be a negative regulator for endogenous auxin levels.

Results

Fluridone suppresses the growth of the primary root

Fluridone inhibited PR growth and induced LR density in a dose-dependent manner (**Fig. 1, B,**

D and E). The emerging true leaves of the seedlings were bleached following fluridone

treatment; however, the established leaves, such as the cotyledons, were bleached to a lesser

extent (**Fig. 1, C and F**). Other PDS inhibitors, including norflurazon, diflufenican, and

picolinafen, also suppressed PR growth and increased LR density comparable with the effects

of fluridone treatment (Supplemental Fig. S1). To identify the target of fluridone for root

growth inhibition, the fluridone-insensitive trait PDS (*35S::mHvPDS*) (Arias et al., 2006) was

introduced into the *Arabidopsis* plant. The *35S::mHvPDS* plants exhibited less bleached

leaves (**Fig. 1C**) and less reduced carotenoid content (**Fig. 1F**) following fluridone treatment,

confirming that mutated *PDS* is less sensitive to fluridone (Arias et al., 2006). The

35S::mHvPDS plants also exhibited partial resistance to root growth inhibition (**Fig. 1D**).

These results indicate that PDS is the target of fluridone for PR growth inhibition and

implicates the carotenoid biosynthesis pathway for the involvement to the root growth

regulation. PR growth was inhibited for 2 days following fluridone treatment, with the extent

of inhibition further increasing after the third day (Supplemental Fig. S3). LR density was

observed to increase with fluridone treatment in the wild type (WT) and to a lesser extent, in

the *35S::mHvPDS* plants (**Fig. 1E**). Since the LR initiation and emergence are under the

control of auxin signaling (Du and Scheres, 2018), increased LR density indicates the

activation of auxin signaling.

Fluridone increases endogenous auxin levels and activates auxin signaling

The fluridone-induced PR growth inhibition and LR density elevation prompted us to

measure the endogenous auxin levels. Although the WT exhibited increased IAA levels 8 h

and 24 h following fluridone treatment, and then auxin levels returned to normal range (**Fig. 2**;

Supplemental Fig. S2A). However, no significant change of auxin levels was observed in the

35S::mHvPDS plants (Supplemental Fig. S2B). To determine the kinetics of auxin signaling,

a time-lapse image of the artificial auxin reporter *DR5:Luciferase* (*DR5:LUC*)

(Moreno-Risueno et al., 2010) was captured and compared between the WT and

35S::mHvPDS plants. Fluridone-induced *DR5:LUC* expression in the root tip and root–shoot

junction was observed in the WT and to a lesser extent, in the *35S::mHvPDS* plants

(Supplemental Movie. S1). *DR5:LUC* expression in the root tip increased instantly,

irrespective of fluridone treatment, decreasing thereafter (**Fig. 3**). This transient activation of

DR5:LUC may occur due to the stress caused by the transfer of plants to another medium.

Following this artificial elevation, *DR5:LUC* expression remained higher in the fluridone

medium compared with that in the control medium after 14 h in WT plants (**Fig. 3A**);

however, the *35S::mHvPDS* plants did not exhibit *DR5:LUC* expression (**Fig. 3B**). These

results suggest that PDS inhibition elevated auxin levels and activated auxin signaling.

Auxin biosynthesis and signaling are required for carotenoid-mediated root growth regulation

We hypothesized that fluridone-induced PR growth inhibition is mediated by the elevation of

endogenous auxin and activation of auxin signaling. Therefore, mutants of auxin signaling or

auxin biosynthesis were examined for fluridone-mediated PR growth inhibition. The roots of

the *AUX/IAA* dominant mutants *aux/iaa19/msg2-1* and *aux/iaa14/slr-1* exhibited a moderate

but nonsignificant resistance to fluridone. The *ARF* double-mutant *arf7-1 arf19-1* also

exhibited a similar level of resistance to that of *aux/iaa19/msg2-1* and *aux/iaa14/slr-1*. An

auxin permease mutant *aux1-21* exhibited higher resistance to fluridone compared with the

above mutants. A mutant of ubiquitin-activating enzyme E1, *axr1-12* (Lincoln et al., 1990),

exhibited the highest resistance to fluridone; however, the resistance of *aux/iaa3/shy2-101*

was comparable with that of the WT (**Fig. 4A**), thus indicating that a dominant mutant of the

AUX/IAA gene does not confer resistance to fluridone.

Auxin biosynthesis and auxin signaling are necessary for PR growth (Mallory et al., 2005;

Chen et al., 2014). Anthranilate is a precursor to tryptophane biosynthesis, which is an

intermediate of IAA biosynthesis. Mutants of the anthranilate synthase subunit *wei2* and *wei7*

decrease endogenous auxin levels (Stepanova et al., 2005), exhibiting moderate and no

resistance to fluridone, respectively, whereas their double-mutant, *wei2 wei7*, was insensitive

to fluridone (**Fig. 4B**). *SHADE AVOIDANCE 3* (*SAV3*), which encodes one of five tryptophan

aminotransferases in *Arabidopsis*, converts tryptophan to indole-3-pyruvate (Stepanova et al.,

2008). A single mutant of *sav3-2* did not demonstrate resistance to fluridone (**Fig. 4B**).

YUCCA encodes a flavin monooxygenase that catalyzes the last step in auxin biosynthesis,

the conversion of IpyA to IAA (Mashiguchi et al., 2011; Won et al., 2011). The quintuple

YUCCA gene mutants *yucQ* (*yuc 3 5 7 8 9*) decreased the endogenous auxin levels and

exhibited severe dwarf shoots and root phenotypes. This dwarf phenotype could be rescued by

exposure to exogenous auxin (Chen et al., 2014). Applying fluridone to *yucQ* plants did not

inhibit PR growth, instead encouraging it (**Fig. 4C**; >100% of inhibition), suggesting that

increased endogenous auxin levels can complement the growth defect in *yucQ* plants.

However, a single *YUCCA* gene mutant did not exhibit significant defects in PR growth, such

as in the *yucQ* plant, and exhibited less or no resistance to fluridone (Supplemental Fig. S4;

Supplemental Table. S1), indicating the functional redundancy of the *YUC* and *SAV3* gene

families in this mechanism. These results indicate that carotenoid biosynthesis (i.e., PDS

activity) negatively regulates endogenous auxin levels and auxin signaling.

Next, we evaluated the pharmaceutical perturbation of auxin signaling and biosynthesis in

combination with PDS inhibition by fluridone. Auxinole targets the auxin binding site of the

auxin receptor TIR1/AFB (Hayashi et al., 2012) and inhibits auxin signaling. Yucasin

difluorinated analog (YDF) targets the enzymatic activity of YUCCA (Tsugafune et al., 2017).

Although auxinole and YDF can alone suppress PR growth, adding fluridone to either

inhibitor did not enhance growth inhibition (**Fig. 4, D and E**). These results support the

hypothesis that fluridone-mediated PR growth inhibition requires auxin signaling and a

certain level of endogenous auxin. The addition of exogenous auxin or an inhibitor of auxin

efflux transport, such as N-1-naphthylphthalamic acid (NPA) (Casimiro et al., 2001), also

inhibited PR growth in a dose-dependent manner. However, in contrast to auxinole and YDF,

fluridone exhibited an additional effect to both exogenous auxin and NPA on PR growth

(Supplemental Fig. S5). These results suggest that fluridone-induced PR growth suppression occurs solely by an elevation of auxin levels and an efflux transporter may not be the major target of PDS inhibition. Taken together, auxin biosynthesis, signaling, and catabolism may be required for carotenoid-mediated PR growth regulation.

Auxin negatively regulates carotenoid content in root

One of the agricultural uses of auxin is as a herbicide, which causes chlorosis in broad-leaf plants (Grossmann, 2010). The auxin-insensitive mutants *arf7/msg1* and *axr1* are resistant to 2,4-dichlorophenoxyacetic acid (2,4-D) and chlorosis (Watahiki and Yamamoto, 1997). Auxin signaling negatively regulates chloroplast development (Kobayashi et al., 2012) and it is reasonable to consider that carotenoid content is also negatively regulated by auxin signaling. To clarify this hypothesis, the level of β-carotene in the root tissue was quantified. Auxinole and YDF increased β-carotene level in the root and fluridone addition reduced β-carotene level regardless of the presence of auxinole or YDF (**Fig. 5, A and B**). The auxin biosynthesis mutants *wei2 wei7* and *yucQ* increased the level of β-carotene more than two-fold compared with that in the WT (**Fig. 5C**). In addition, a negative regulation of carotenoid accumulation by auxin in the shoot tissue was observed (Supplemental Fig. S6). Conversely, the auxin signaling mutants *aux/iaa19/msg2-1* and *axr1-12* exhibited a minor but nonsignificant increase in β-carotene levels compared with that in the WT (**Fig. 5C**). These data suggest that endogenous auxin levels and signaling negatively control β-carotene levels.

Expression of auxin- and carotenoid-related genes

To explore whether carotenoids regulate auxin biosynthesis and signaling at the gene expression level, the expression of auxin-related genes in root tissue was measured following fluridone treatment. The expression of *AUX/IAA19* increased by approximately four-fold at 4 and 8 h following fluridone treatment in the WT but not in the *35S::mHvPDS* plant (**Fig. 6A**). Consistently, the expression of the auxin biosynthesis gene *YUC9* increased by ~2.5 times at 4 and 8 h following fluridone treatment (**Fig. 6B**). The induction time of auxin biosynthesis and signaling gene expression was associated with the induction of endogenous auxin levels (**Fig.

2). These data suggest that *PDS* inhibition activates auxin biosynthesis and signaling.

Next, we examined the expression of genes downstream of carotenoid biosynthesis or

catabolism. *CCD7* and *CCD8* are carotenoid catabolism genes (i.e., apocarotenoid

biosynthesis genes) (**Fig. 1A**) (Yuan et al., 2015). *CCD7* and *CCD8* were primarily induced at

8 and 24 h in the WT plants, whereas the expression of these genes remained unchanged in

the *35S::mHvPDS* plants (**Fig. 6, C and D**). On the other hand, carotenoid biosynthesis genes

CCR1 and *CCR2*, and catabolism gene *CCD1* were also examined. The expression of these

genes were also somewhat induced (Supplemental Fig. S7). These data imply carotenoid

catabolism rather than biosynthesis is activated following PDS inhibition.

Retinal controls auxin content in fluridone-mediated PR growth

regulation

The target of fluridone was PDS for PR growth inhibition; however, the downstream product

of PDS activity responsible for this process remains unknown. The extremely hydrophobic

nature of β-carotene made it impossible for the complementation test by exogenous

application to the plant, which was inhibited by fluridone (data not shown). Recently, various

β-carotene-derived apocarotenoids were reported to be bioactive. Strigolactone is one such

apocarotenoids and mutants of strigolactone biosynthesis genes *max3* and *max4* as well as the

strigolactone signaling gene *max2* were examined for fluridone-induced PR growth inhibition.

However, these mutants exhibited the same response to fluridone as the WT (data not shown)

and may not be responsible for fluridone-induced PR growth inhibition. Then, two

β-carotene-derived apocarotenoids, β-cyclocitral and retinal, were employed for the

complementation test to fluridone. Retinal, but not β-cyclocitral, completely restored

fluridone-induced PR growth inhibition (**Fig. 7A**; Supplemental Fig. S8). An auxin-inducible

pIAA19::ELucPEST reporter (Yamamoto et al., 2017) was used for quantifying auxin

signaling. *IAA19* activity increased two-fold in the root tip area following treatment with 10

nM 1-naphthaleneacetic acid (NAA), a synthetic auxin. However, this induction was

counteracted by cotreatment with retinal (**Fig. 7E**). These data suggest that the reduction of

retinal levels enhances auxin levels. If true, retinal may complement the inhibitory effect of

auxin. Indeed, retinal restored NAA-induced PR growth inhibition (**Fig. 7B**). Although retinal

did not affect PR growth in the WT, PR growth in the *wei2 wei7* and *yucQ* mutants was

significantly decreased (**Fig. 7C**). From these data, it is speculated that retinal negatively

regulates endogenous auxin levels; however, no significant reduction in IAA levels was

detected in the root of the retinal-treated plants (Supplemental Fig. S9G). Auxin catabolism

inhibitor KKI (Fukui et al., 2022) inhibits PR growth in a dose-dependent manner (**Fig. 7D**)

like as treatment with exogenous auxin (**Fig. 7B**). Interestingly, retinal did not affect

KKI-induced growth inhibition nor did it rescue PR growth inhibition; however, retinal

restored exogenous auxin–induced growth inhibition (**Fig. 7, B and D**). Although the retinal

levels in fluridone-treated plants remain to be determined, these results indicate that

fluridone-induced retinal levels reduction suppresses endogenous auxin catabolism,

consequently increasing auxin levels (**Fig. 2**).

Discussion

Herein, we demonstrated the coupling of carotenoid biosynthesis with root growth regulation.

We investigated that inhibiting PDS with fluridone suppressed PR growth (**Fig. 1D;**

Supplemental Fig. S3). In addition, other PDS inhibitors, such as norflurazon, diflufenican,

and picolinafen, suppressed PR growth and increased LR density, comparable with the effects

of fluridone (Supplemental Fig. S1). This inhibition of PR growth was caused by PDS activity

suppression, as *35S::mHvPDS* plants are less sensitive to fluridone (**Fig. 1D**; Supplemental

Fig. S3). PDS activity suppression coincides with elevated endogenous IAA levels and

suppressed PR growth (**Fig. 1, D** and **Fig. 2**; Supplemental Fig. S2A). The *msg2-1/aux/iaa19*

(Tatematsu et al., 2004) and *slr-1/aux/iaa14* (Fukaki et al., 2002) plants were less sensitive to

fluridone compared with the WT (**Fig. 4A**). In fact, *arf7 arf19* (Okushima et al., 2007)

exhibited similar resistance to fluridone as that of *msg2-1* and *slr-1* (**Fig. 4A**). The *axr1-12*

and *aux1-21* plants exhibited higher resistance to root inhibition by fluridone compared with

the other auxin signaling mutants (**Fig. 4A**). The *axr1-12* and *aux1-7* plants were previously

reported to exhibit higher resistance to exogenous auxin compared with *slr-1*, *arf7 arf19*, and

msg2-1 (Timpte et al., 1995; Fukaki et al., 2002; Tatematsu et al., 2004; Okushima et al.,

2005), which is relevant to fluridone resistance (**Fig. 4A**). This supports the hypothesis that fluridone-mediated growth inhibition requires auxin signaling and transport. However, no correlation was observed between fluridone resistance and LR formation defects among the mutants, as *msg2-1/aux/iaa19* and *slr-1/aux/iaa14* were highly LR-defective; however, *axr1-12* and *aux1-7* exhibited LR formation defects to a lesser extent. The *shy2-101/aux/iaa3* plant is also an LR-defective mutant (Goh et al., 2012), however, it exhibited sensitivity to fluridone, similar to that of the WT (**Fig. 4A**). *MSG2/AUX/IAA19* and *SLR/AUX/IAA14* expression is regulated by ARF and AUX/IAA during auxin signaling (Tatematsu et al., 2004; Ito et al., 2016), whereas *AUX/IAA3* expression is under the control of cytokinin (Bishopp et al., 2011) and brassinosteroid (Li et al., 2020) signaling. Fluridone activated a higher degree of *DR5:LUC* activity in the root tip in WT, but not in *35S::mHvPDS* (**Fig. 3**). These findings suggest that auxin-mediated gene expression participates in carotenoid-mediated PR regulation. Finally, auxinole, which is an inhibitor of the auxin receptor (Hayashi et al., 2012), inhibited PR growth; however, fluridone addition did not further suppress PR growth (**Fig. 4D**). This result also supports the requirement of auxin signaling in fluridone-mediated PR inhibition. Conversely, the suppression effect of PR growth by NPA and fluridone was additive (Supplemental Fig. S5B). This may exclude the involvement of the auxin efflux pathway in carotenoid-mediated PR growth regulation.

PDS inhibition activity increases auxin levels

Fluridone-induced PDS activity suppression elevates auxin levels in the root tissue, which may exceed the optimal concentration for root growth, while inhibiting root growth in the WT plants (**Fig. 1, D and 2**). In the case of *yucQ* plant, an acceleration of PR growth in *yucQ* plant which has suboptimal levels of auxin and shorter roots compared with those of the WT (Chen et al., 2014), may be explained by fluridone-induced elevation of auxin levels (**Fig. 4C**). In fact, root growth in *yucQ* was enhanced by exogenous auxin (Chen et al., 2014). Single mutants of the *YUC* gene family exhibited less or no resistance to fluridone (Supplemental Fig. S4), suggesting a functional redundancy of the *YUC* gene family in this mechanism. The ectopic expression of *YUC3* in the shoot of *yucQ* plant (*FRO6::XVE::YUC3*)

resulted in elongated hypocotyl (Chen et al., 2014), which is a typical phenotype of auxin

overproduction (Boerjan et al., 1995; Delarue et al., 1998). However, the short root of *yucQ*

was not rescued by the expression of *FRO6::XVE::YUC3*, suggesting a bare contribution of

shoot-derived auxin for root growth (Chen et al., 2014). Therefore, the effect on root growth

of fluridone in the WT and *yucQ* is root-specific and not based on shoot-derived auxin.

Notably, fluridone-induced elevation of auxin levels was detected in the root but not in the

shoot (Supplemental Table 2).

The *wei2-1 wei7-1* plants exhibited a short-root phenotype complemented by exogenous

auxin, such as the *yucQ* plant (Stepanova et al., 2005). Conversely, its root growth was not

rescued; however, it exhibited total insensitivity to fluridone (**Fig. 4B**). Although the IPA

pathway is a major way of auxin biosynthesis, there is a diversification of the auxin

biosynthesis pathway after tryptophan (Zhao et al., 2010; Kasahara, 2016). The difference in

response to fluridone between the *wei2-1 wei7-1* double-mutant and *yucQ* indicates the role

for the alternative pathway of auxin biosynthesis in fluridone-mediated root inhibition.

Together with the induction of *IAA19* and *YUC9* following fluridone treatment, (**Fig. 6, A and**

B) results all above indicate that fluridone activates auxin signaling and biosynthesis.

The inhibition of auxin signaling and biosynthesis by auxinole and YDF, respectively,

inhibited PR growth; however, fluridone addition did not further enhance this suppression

(**Fig. 4, D and E**). These results also indicate that auxin biosynthesis and signaling are

required to inhibit root growth by fluridone. In contrast to auxinole and YDF, exogenous

auxin and fluridone exhibited an additive effect on PR growth suppression (Supplemental Fig.

S5A). These pharmaceutical approaches can indicate the optimal level of auxin and auxin

signaling required for PR growth as well as whether excessive or insufficient auxin levels can

reduce PR growth. Taken together, fluridone-induced PDS inhibition increases endogenous

auxin levels, activates auxin signaling, and thus inhibits PR growth.

Interaction between auxin homeostasis and carotenoid catabolism

One of the apocarotenoids, retinal, completely restored PR growth inhibition induced by

fluridone (**Fig. 7A**) and NAA (**Fig. 7B**). Retinal is reported to be responsible for periodic root

branching and regulate LR formation (Dickinson et al., 2021). Besides of retinal,

apocarotenoid β-cyclocitral is also a regulator for both PR and LR development (Dickinson et

al., 2019). Conversely, β-cyclocitral did not rescue fluridone-induced PR growth inhibition

(Supplemental Fig. S8). Retinal counteracted the LR induction caused by NAA and

tryptophan and mitigated the exaggerated LR formation of the *sur2* mutant (Supplemental Fig.

S9, C–E). These findings indicate a negative role of retinal in auxin signaling. Indeed, retinal

suppressed the activation of auxin-inducible *AUX/IAA19* promoter by auxin (**Fig. 7E**).

Although retinal restored the fluridone-induced PR suppression over the first 2 d (**Fig. 7A**),

partial recovery at a higher concentration of retinal occurred on the third day after treatment

(Supplemental Fig. S9A). Notably, the rate of PR growth suppression accelerated after 48 h of

fluridone treatment (Supplemental Fig. S3). It is hypothesized that the pool of carotenoids

(i.e., β-carotene) was highly depleted at this time, and accordingly, the requirement of retinal

raised a high level. KKI is an inhibitor of the auxin–amino acid conjugating enzyme

GRETCHEN HAGEN 3 (GH3). The KKI-induced auxin catabolism inhibition rapidly

elevated endogenous IAA levels, leading to typical phenotype of excessive auxin, including

induced LR and adventitious root formation, and reduced PR growth (Fukui et al., 2022). In

contrast to the antagonistic effects of exogenous auxin (**Fig. 7B**), retinal did not rescue either

of these phenotypes (**Fig. 7E**; Supplemental Fig. S9F), suggesting that retinal may regulate

IAA levels through IAA catabolism. Fluridone-activated *CCD7* and *CCD8* expression at 8

and 24 h in the WT plants (**Fig. 6, C and D**). The *CCD7* and *CCD8* genes, which convert

9-*cis*-β-carotene to 9-*cis*-β-apo-10'-carotenal and 9-*cis*-β-apo-10'-carotenal to carlactone,

respectively, are induced by auxin (Bainbridge et al., 2005; Hayward et al., 2009). These data

suggest that fluridone-induced auxin (**Fig. 2 and 3**) activates the catabolism of β-carotene and

suggest the feedback regulation of apocarotenoid biosynthesis. The GENEVESTIGATOR

(https://genevestigator.com) (Hruz et al., 2008) database also confirmed that *CCD7*

expression is induced between 0.5 and 1 h following exogenous IAA treatment (Supplemental

Fig. S10).

Conclusion

Herein, we established a model of carotenoid-mediated auxin regulation and response via the apocarotenoid retinal (**Fig. 8**). The inhibition of carotenoid biosynthesis elevated auxin levels and induced auxin response, which may have reduced retinal levels and promoted endogenous auxin levels and response. We hypothesized that retinal may accelerate IAA catabolism to regulate endogenous auxin levels. Moreover, auxin may regulate carotenoid accumulation through a feedback mechanism involving the expression of genes associated with carotenoid catabolism. This model reveals the hidden part of the carotenoid and auxin interaction to regulate root growth. It also provides insight into the role of the apocarotenoid retinal in root architecture. Finally, the quantitative analysis of retinal and its derivatives as well as their targets, which may regulate auxin homeostasis, is warranted in the future.

Materials and methods

Plant materials and growth conditions

The *Arabidopsis thaliana* mutants and WT plants as well as *PDS* transgenic plants were in the Columbia background, except for *yuc5*, which is in the Landsberg *erecta* background. The *arf7-1 arf19-1*, *sav3-2*, *yuc1*, *yuc2*, *yuc3*, *yuc4-1*, *yuc5*, *yuc6*, *yuc7*, *yuc8*, *yuc9*, *yuc10*, and *yuc11* plants were described previously (Xu et al., 2017). The *axr1-12* (Timpte et al., 1995), *aux1-21* (Roman et al., 1995), *wei2-1*, *wei7-1*, *wei2-1/wei7-1* and *sur2* plants (Stepanova et al., 2005) were obtained from the Arabidopsis Biological Resource Center. The *yucQ* mutant was a generous gift from Dr Hiroyuki Kasahara (Tokyo University of Agriculture and Technology) with an agreement from Dr Yunde Zhao (University of California San Diego). *DR5:LUC* (Moreno-Risueno et al., 2010) was kindly provided from Dr Philip N. Benfey (Duke University). In this study, *DR5:LUC* reporter in the *DR5:LUC/VENUS* line (Toyokura et al., 2019) was observed. The *slr-1* (Fukaki et al., 2002) and *shy2-101* (Goh et al., 2012) mutants were described previously. Fluridone, norflurazon, deflufenican, picolinafen, and retinal were purchased from Fujifilm Wako pure chemical. KKI, YDF, and auxinole were provided by Kei-ichiro Hayashi. Seeds were surface-sterilized using chlorine gas for at least 45 min, suspended in 0.38% agarose, and sown on half-strength Murashige and Skoog (MS) medium

449   (Duchefa Biochemie) supplemented with 1% (w/v) sucrose, 0.6% (w/v) gellan gum, and 0.5

450   mM MES (pH 5.8) if not specifically indicated. Stratification was performed at 4°C for 3 d in

451   the dark. Plants were grown on vertically oriented plates at 22°C under constant light if not

452   specifically indicated (80–110 µmol m^{-2} s^{-1}). Stock solutions of phytohormones and

453   inhibitors were prepared in dimethyl sulfoxide and filtered through a 0.45 µM disc filter.

PDS-resistant vector and plant transformation and screening

The plasmid pPDSer1303 was kindly provided by Renée S. Arias (USDA ARS, USA). The

pds mutation vector construction was described previously (Arias et al., 2006). The plasmids

were transformed into *Agrobacterium* strain C58C1 and used to transform *Arabidopsis*

thaliana via the dip floral method. *Agrobacterium* harboring the plasmid pCambia1303

without *pds* was used as a control for the transformations and later as a control for herbicide

screening. Seeds from T1, T2, and T3 generations were screened with hygromycin (50 mg/L)

and fluridone (25 nM) separately. Seedlings that were resistant to hygromycin in T1 were

selected in either hygromycin or fluridone (25 nM) screening in T2 and T3. T3 seeds were

screened for resistance to fluridone at 25, 50, and 200 nM.

Quantification of LR density and PR growth

467   Plants were grown on vertical plates for 5 d and then transferred to a new half-strength MS

468   medium with or without inhibitors or phytohormones. Plant images were captured using a

469   flatbed scanner (GT-X980 and DS-50000, EPSON) within 4 d after transfer. The density of

470   emerged LRs from the root–shoot junction beside an adventitious root and the PR growth rate

471   were analyzed with Fiji software (https://fiji.sc/).

Quantification of PR growth via time-lapse imaging

Plants were grown on vertical plates for 4 d and then transferred to a new half-strength MS

medium. Plant images were captured using a digital camera (Lumix G5, Panasonic). The

interval was 30 min. After 1 d of photography, the seedlings were transferred to a new

half-strength MS medium with or without 800 nM fluridone. The PR growth rate was

analyzed using Fiji software (https://fiji.sc/).

Quantification of IAA

Seeds were sown in sterilized 300 μm nylon mesh and transferred to control or fluridone

plates after a 5-d germination together with mesh. Shoot and root samples were harvested at

specific timepoints after transfer and frozen in liquid nitrogen. Hormonome analysis was

conducted as previously described (Xu et al., 2017). Briefly, samples of ~100 mg fresh weight

(Fw) were suspended in 80% (v/v) aqueous methanol with $[^2H_5]$-IAA as internal standards.

The samples were homogenized and the supernatant was loaded onto a Bond Elut C18

cartridge (100 mg, 3 ml; Agilent Technologies) and eluted with 80% (v/v) aqueous methanol.

The concentrated samples were subjected to liquid chromatography with electrospray

ionization tandem mass spectrometry comprising a quadrupole tandem mass spectrometer

(Agilent 6460 Triple Quadrupole mass spectrometer) with an electrospray ion source and an

Agilent 1200 separation module. The raw data were extracted from MassHunter software

(Agilent Technologies) and examined in Excel (Microsoft).

Bioluminescence imaging analysis

D-Luciferin was purchased from Gold Biotechnology. For *DR5:LUC* expression, seeds were

germinated on medium containing 100 μM D-luciferin. Five-day-old plants were transferred

to 100 μM D-luciferin medium with or without fluridone, following which time-lapse imaging

was performed using a camera (ASI294MMPro, ZWO) with control software (SharpCap,

v4.0). For the retinal treatment of *pIAA19::ELucPEST*, seeds were germinated on 100 μM

D-luciferin medium. Four-day-old plants were transferred to 100 μM D-luciferin medium for

preincubation. After 1 d, seedlings were transferred to 100 μM D-luciferin medium with NAA

or NAA and retinal and time-lapse imaging was continued. The images were captured at an

interval of 1 h, including 15 min of exposure and 45 min of red and blue light (100 μmol m^{-2}

s^{-1}) for 5 d and then saved in a TIFF format for further analysis using Fiji software

(https://fiji.sc/).

RNA isolation and quantitative reverse transcription polymerase chain reaction analysis

Seeds were sown on control medium with 300 μm nylon mesh. Roots were placed on plates containing no herbicide (mock) or 800 nM fluridone, root samples were harvested at 0, 4, 8, and 24 h and frozen in liquid nitrogen. Total RNA was extracted and purified using an RNA isolation kit (ISOSPIN Plant RNA, NIPPON GENE). Complementary DNA was synthesized from the total RNA according to the manufacturer's instructions (ReverTra Ace qPCR RT Master Mix, Toyobo). Quantitative reverse transcription polymerase chain reaction (qRT–PCR) was performed in optical 96-well plates via the LightCycler 480 II system (Roche Life Science) using KOD SYBR qPCR Mix (Toyobo). Primer pairs spanning exon–exon junctions were designed using the QuantPrime program to avoid genomic DNA amplification as listed in Supplemental Table S3 (Xu et al., 2017). The specificity of the reactions was verified via melt curve analysis and capillary electrophoresis (Multina, Shimadzu). Standard curve analysis was used to evaluate the efficiency of the reactions. ACTIN2 was used as an internal standard. The qRT–PCR program included one cycle at 98°C for 2 min, followed by 40 cycles of 98°C for 10 s, 60°C for 10 s, and 68°C for 30 s. The cycle time values were determined using the second derivative maximum method (Tichopad et al., 2003) via LightCycler software (version 1.5, Roche Life Science). The data were analyzed using the comparative threshold cycle (Ct) method ($2^{-\triangle\triangle Ct}$) (Schmittgen and Livak, 2008).

Quantification of carotenoid via high-performance liquid chromatography analysis

Seeds were sown on control medium with 300 μm nylon mesh. The seedlings were grown under the conditions described above. Five days after germination, the seedlings together with nylon mesh were transferred to plates with or without fluridone and with or without NAA or auxinole or YDF. Shoot tissues (~20 mg) or root tissues (~100 mg) were harvested at specific timepoints and stored at −80°C. Total carotenoids were extracted using acetone wako 1st grade (FUJIFILM Wako Chemicals). Acetone was added to the tissue samples that had been mixed with beads and silica gel and shaken for 2 min. Total carotenoids were separated via

centrifugation for 10 min at 14000 rpm at 4℃. High-performance liquid chromatography analysis for carotenoid quantification was performed as previously described (Zapata et al., 2000). An aliquot of the sample was subjected to a C8 column (4.6 × 150 mm, Waters Symmetry C8, Waters) equilibrated with A solution (methanol:acetonitrile:aqueous pyridine solution [0.25 M pyridine] = 2:1:1 [v:v:v]). Pigments were eluted with a linear gradient of 0%–40% B solution (methanol:acetonitrile:acetone = 1:3:1 [v:v:v]) for 2 min, 40%–95% for 6 min, an isocratic elution for 5 min, a linear gradient of 95%–0% for 2 min, and an isocratic elution for 7 min at a flow rate of 1 mL/min. Pigments were identified based on their retention time relative to known standards and quantified by peak area on the chromatogram.

Statistical analysis

In this study the statistical significance was determined by 2-tailed Student's *t*-test (*: $P < 0.05$, **: $P < 0.01$) or one-way ANOVA following Tukey-Kramer test (Different lowercase letters indicate significant differences, $P < 0.05$).

Acknowledgments

We thank Dr. Renée Arias (United States Department of Agriculture) and Dr. Franck Dayan (Colorado State University) for the supply of plasmid *35S::mHvPDS* and for the kind reviewing of the manuscript.

Author contributions

Author contributions: K.X. conceived the project and designed the experiments, which were mostly carried out and analyzed by K.X.; H.Z. supported retinal-related experiment design, process and data analysis; E.Y. and M.A. were responsible for IAA quantification and data collection; K.X. and M.K.W. wrote the article. K.X., H.Z., E.Y., M.A., K.H., H.F., H.I. and M.K.W. read, commented and approved the final manuscript.

Supplemental data

The following materials are available in the online version of this article.

Supplemental Figure S1. Other PDS-target inhibitors exhibit a fluridone-like effect on

PR suppression and LR induction.

Supplemental Figure S2. Long-period auxin level change by fluridone.

Supplemental Figure S3. Kinetic PR growth suppression of WT and *mPDS* plants by

fluridone.

Supplemental Figure S4. *yuc* single mutants have weak or no resistance to

fluridone.

Supplemental Figure S5. Exogenous auxin and NPA suppress PR in an additive manner

with fluridone.

Supplemental Figure S6. Carotenoid in shoot tissue is negatively regulated by

auxin.

Supplemental Figure S7. Relative expression level change of carotenoid

biosynthesis genes (*CCR1*, *CCR2*) and the metabolism gene *CCD1*.

Supplemental Figure S8. β-cyclocitral is not involved in carotenoid-mediated PR

regulation.

Supplemental Figure S9. Retinal is responsible for fluridone effect and regulates

downstream auxin content and response.

Supplemental Figure S10. *CCD7* expression is rapidly induced by exogenous IAA

treatment.

Supplemental Table S1. PR growth rate in no fluridone-treated auxin-related mutants.

Supplemental Table S2. Endogenous IAA concentration in fluridone treatment in shoot

tissue.

Supplemental Table S3. Gene-specific primers used for qRT–PCR analysis.

Supplemental Movie S1. *DR5:LUC* bioluminescence video under fluridone treatment in

WT and *mPDS*.

Funding

This research was funded by grants-in-aid for scientific research from the Japan Society for

the Promotion of Science (JSPS) (JP19H05673 to H. F. and M. K. W., JP21K19097 to K.H.) ,

Japan Science and Technology Agency Support for Pioneering Research Initiated by the Next

Generation (JST SPRING) (Grant Number JPMJSP2119 to K. Y.) and ACRO Research grant

of Teikyo University (TeTe20-01 to M.A.).

conflict of interest statement. The authors have no conflicts of interests to declare.

References

Arias R, Dayan F, Michel A, Howell J, Scheffler B (2006) Characterization of a higher plant herbicide-resistant phytoene desaturase and its use as a selectable marker. *Plant Biotechnology Journal* **4:** 263-273

Auldridge M, McCarty D, Klee H (2006) Plant carotenoid cleavage oxygenases and their apocarotenoid products. *Current Opinion in Plant Biology* **9:** 315-321

Bainbridge K, Sorefan K, Ward S, Leyser O (2005) Hormonally controlled expression of the Arabidopsis *MAX4* shoot branching regulatory gene. *Plant Journal* **44:** 569-580

Barlier I, Kowalczyk M, Marchant A, Ljung K, Bhalerao R, Bennett M, Sandberg G, Bellini C (2000) The *SUR2* gene of *Arabidopsis thaliana* encodes the cytochrome P450 CYP83B1, a modulator of auxin homeostasis. *Proceedings of the National Academy of Sciences, USA* **97:** 14819-14824

Bartels PG, Watson CW (1978) Inhibition of carotenoid synthesis by fluridone and norflurazon. *Weed Science* **26:** 198-203

Bishopp A, Benkova E, Helariutta Y (2011) Sending mixed messages: auxin-cytokinin crosstalk in roots. *Current Opinion in Plant Biology* **14:** 10-16

Boerjan W, Cervera M-T, Delarue M, Beeckman T, Dewitte W, Bellini C, Caboche M, Van Onckelen H, Van Montagu M, Inzé D (1995) *superroot*, a recessive mutation in Arabidopsis, confers auxin overproduction. *The Plant Cell* **7:** 1405-1419

Breitenbach J, Sandmann G (2005) ζ-Carotene *cis* isomers as products and substrates in the plant poly-*cis* carotenoid biosynthetic pathway to lycopene. *Planta* **220:** 785-793

Casimiro I, Marchant A, Bhalerao R, Beeckman T, Dhooge S, Swarup R, Graham N, Inze D,

Sandberg G, Casero P, Bennett M (2001) Auxin transport promotes Arabidopsis lateral root initiation. *The Plant Cell* **13:** 843-852

Cazzonelli C (2011) Carotenoids in nature: insights from plants and beyond. *Functional Plant Biology* **38:** 833-847

Cazzonelli C, Roberts A, Carmody M, Pogson B (2010) Transcriptional control of *SET DOMAIN GROUP 8* and *CAROTENOID ISOMERASE* during *Arabidopsis* development. *Molecular Plant* **3:** 174-191

Chen Q, Dai X, De-Paoli H, Cheng Y, Takebayashi Y, Kasahara H, Kamiya Y, Zhao Y (2014) Auxin overproduction in shoots cannot rescue auxin deficiencies in Arabidopsis roots. *Plant and Cell Physiology* **55:** 1072-1079

Cheng Y, Dai X, Zhao Y (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in *Arabidopsis*. *Genes & Development* **20:** 1790-1799

Cheng Y, Dai X, Zhao Y (2007) Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in *Arabidopsis*. *The Plant Cell* **19:** 2430-2439

Delarue M, Prinsen E, Van Onckelen H, Caboche M, Bellini C (1998) *Sur2* mutations of *Arabidopsis thaliana* define a new locus involved in the control of auxin homeostasis. *The Plant Journal* **14:** 603-611

Dharmasiri N, Dharmasiri S, Estelle M (2005) The F-box protein TIR1 is an auxin receptor. *Nature* **435:** 441-445

Dickinson A, Lehner K, Mi J, Jia K, Mijar M, Dinneny J, Al-Babili S, Benfey P (2019) β-Cyclocitral is a conserved root growth regulator. *Proceedings of the National Academy of Sciences, USA* **116:** 10563-10567

Dickinson A, Zhang J, Luciano M, Wachsman G, Sandoval E, Schnermann M, Dinneny J, Benfey P (2021) A plant lipocalin promotes retinal-mediated oscillatory lateral root initiation. *Science* **373:** 1532-1536

Du Y, Scheres B (2018) Lateral root formation and the multiple roles of auxin. *Journal of Experimental Botany* **69:** 155-167

Fukaki H, Tameda S, Masuda H, Tasaka M (2002) Lateral root formation is blocked by a

gain-of-function mutation in the *SOLITARY-ROOT/IAA14* gene of *Arabidopsis*. *The Plant*

Journal **29:** 153-168

Fukaki H, Tasaka M (2009) Hormone interactions during lateral root formation. *Plant Molecular*

Biology **69:** 437-449

Fukui K, Arai K, Tanaka Y, Aoi Y, Kukshal V, Jez JM, Kubes MF, Napier R, Zhao Y, Kasahara H

(2022) Chemical inhibition of the auxin inactivation pathway uncovers the roles of metabolic

turnover in auxin homeostasis. *Proceedings of the National Academy of Sciences, USA* **119:**

e2206869119

Geisler M, Murphy A (2006) The ABC of auxin transport: The role of p-glycoproteins in plant

development. *FEBS Letters* **580:** 1094-1102

Goh T, Kasahara H, Mimura T, Kamiya Y, Fukaki H (2012) Multiple AUX/IAA-ARF modules

regulate lateral root formation: the role of *Arabidopsis* SHY2/IAA3-mediated auxin signalling.

Philosophical Transactions of the Royal Society B: Biological Sciences **367:** 1461-1468

Goto C, Ikegami A, Goh T, Maruyama K, Kasahara H, Takebayashi Y, Kamiya Y, Toyokura K,

Kondo Y, Ishizaki K, Mimura T, Fukaki H (2023) Genetic interaction between Arabidopsis

SUR2/CYP83B1 and *GNOM* indicates the importance of stabilizing local auxin accumulation

in lateral root initiation. *Plant and Cell Physiology* pcad084 .

Grossmann K (2010) Auxin herbicides: current status of mechanism and mode of action. *Pest*

Management Science **66:** 113-120

Hayashi K, Neve J, Hirose M, Kuboki A, Shimada Y, Kepinski S, Nozaki H (2012) Rational design

of an auxin antagonist of the SCF^{TIR1} auxin receptor complex. *ACS chemical biology* **7:**

590-598

Hayward A, Stirnberg P, Beveridge C, Leyser O (2009) Interactions between auxin and strigolactone

in shoot branching control. *Plant Physiology* **151:** 400-412

Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W,

Zimmermann P (2008) Genevestigator v3: a reference expression database for the

meta-analysis of transcriptomes. *Advances in Bioinformatics* **2008:** 420747

Ito J, Fukaki H, Onoda M, Li L, Li C, Tasaka M, Furutani M (2016) Auxin-dependent

compositional change in mediator in ARF7-and ARF19-mediated transcription. *Proceedings*

of the National Academy of Sciences, USA **113:** 6562-6567

Jia K, Dickinson A, Mi J, Cui G, Xiao T, Kharbatia N, Guo X, Sugiono E, Aranda M, Blilou I,

Rueping M, Benfey P, Al-Babili S (2019) Anchorene is a carotenoid-derived regulatory

metabolite required for anchor root formation in *Arabidopsis*. *Science Advances* **5:** eaaw6787

Kasahara H (2016) Current aspects of auxin biosynthesis in plants. *Bioscience Biotechnology and*

Biochemistry **80:** 34-42

Kobayashi K, Baba S, Obayashi T, Sato M, Toyooka K, Keranen M, Aro E, Fukaki H, Ohta H,

Sugimoto K, Masuda T (2012) Regulation of root greening by light and auxin/cytokinin

signaling in *Arabidopsis*. *The Plant Cell* **24:** 1081-1095

Koschmieder J, Wust F, Schaub P, Alvarez D, Trautmann D, Krischke M, Rustenholz C, Mano J,

Mueller M, Bartels D, Hugueney P, Beyer P, Welsch R (2021) Plant apocarotenoid

metabolism utilizes defense mechanisms against reactive carbonyl species and xenobiotics.

Plant Physiology **185:** 331-351

Leyser O (2002) Molecular genetics of auxin signaling. *Annual Review of Plant Biology* **53:** 377-398

Li T, Kang X, Lei W, Yao X, Zou L, Zhang D, Lin H (2020) SHY2 as a node in the regulation of root

meristem development by auxin, brassinosteroids, and cytokinin. *Journal of Integrative Plant*

Biology **62:** 1500-1517

Lincoln C, Britton JH, Estelle M (1990) Growth and development of the *axr1* mutants of *Arabidopsis*.

The Plant Cell **2:** 1071-1080

Liu PP, Montgomery TA, Fahlgren N, Kasschau KD, Nonogaki H, Carrington JC (2007)

Repression of *AUXIN RESPONSE FACTOR10* by microRNA160 is critical for seed

germination and post-germination stages. *The Plant Journal* **52:** 133-146

Mallory A, Bartel D, Bartel B (2005) MicroRNA-directed regulation of Arabidopsis *AUXIN*

RESPONSE FACTOR17 is essential for proper development and modulates expression of early

auxin response genes. *The Plant Cell* **17:** 1360-1375

Mashiguchi K, Tanaka K, Sakai T, Sugawara S, Kawaide H, Natsume M, Hanada A, Yaeno T,

Shirasu K, Yao H, McSteen P, Zhao Y, Hayashi K, Kamiya Y, Kasahara H (2011) The

main auxin biosynthesis pathway in *Arabidopsis*. *Proceedings of the National Academy of*

Sciences, USA **108:** 18512-18517

Moreno-Risueno M, Van Norman J, Moreno A, Zhang J, Ahnert S, Benfey P (2010) Oscillating

gene expression determines competence for periodic *Arabidopsis* root branching. *Science* **329:**

1306-1311

Okushima Y, Fukaki H, Onoda M, Theologis A, Tasaka M (2007) ARF7 and ARF19 regulate lateral

root formation via direct activation of *LBD/ASL* genes in *Arabidopsis*. *The Plant Cell* **19:**

118-130

Okushima Y, Overvoorde P, Arima K, Alonso J, Chan A, Chang C, Ecker J, Hughes B, Lui A,

Nguyen D, Onodera C, Quach H, Smith A, Yu G, Theologis A (2005) Functional genomic

analysis of the *AUXIN RESPONSE FACTOR* gene family members in *Arabidopsis thaliana*:

unique and overlapping functions of *ARF7* and *ARF19*. *The Plant Cell* **17:** 444-463

Olson JA, Hayaishi O (1965) The enzymatic cleavage of β-carotene into vitamin A by soluble

enzymes of rat liver and intestine. *Proceedings of the National Academy of Sciences, USA* **54:**

1364-1370

Paponov I, Teale W, Trebar M, Blilou K, Palme K (2005) The PIN auxin efflux facilitators:

evolutionary and functional perspectives. *Trends in Plant Science* **10:** 170-177

Park H, Kreunen S, Cuttriss A, DellaPenna D, Pogson B (2002) Identification of the carotenoid

isomerase provides insight into carotenoid biosynthesis, prolamellar body formation, and

photomorphogenesis. *The Plant Cell* **14:** 321-332

Rodriguez-Concepcion M, Boronat A (2002) Elucidation of the methylerythritol phosphate pathway

for isoprenoid biosynthesis in bacteria and plastids. A metabolic milestone achieved through

genomics. *Plant Physiology* **130:** 1079-1089

Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene

signal transduction in *Arabidopsis thaliana*: five novel mutant loci integrated into a stress

response pathway. *Genetics* **139:** 1393-1409

Schmittgen T, Livak K (2008) Analyzing real-time PCR data by the comparative C_T method. Nature

Protocols **3:** 1101-1108

Schwartz S, Tan B, Gage D, Zeevaart J, McCarty D (1997) Specific oxidative cleavage of

carotenoids by VP14 of maize. *Science* **276:** 1872-1874

Shimizu-Mitao Y, Kakimoto T (2014) Auxin sensitivities of all Arabidopsis Aux/IAAs for

degradation in the presence of every TIR1/AFB. *Plant and Cell Physiology* **55:** 1450-1459

Stepanova A, Hoyt J, Hamilton A, Alonso J (2005) A link between ethylene and auxin uncovered by

the characterization of two root-specific ethylene-insensitive mutants in Arabidopsis. *The*

Plant Cell **17:** 2230-2242

Stepanova A, Robertson-Hoyt J, Yun J, Benavente L, Xie D, DoleZal K, Schlereth A, Jurgens G,

Alonso J (2008) *TAA1*-mediated auxin biosynthesis is essential for hormone crosstalk and

plant development. *Cell* **133:** 177-191

Su L, Diretto G, Purgatto E, Danoun S, Zouine M, Li Z, Roustan J-P, Bouzayen M, Giuliano G,

Chervin C (2015) Carotenoid accumulation during tomato fruit ripening is modulated by the

auxin-ethylene balance. *BMC Plant Biology* **15:** 1-12

Sugawara S, Hishiyama S, Jikumaru Y, Hanada A, Nishimura T, Koshiba T, Zhao Y, Kamiya Y,

Kasahara H (2009) Biochemical analyses of indole-3-acetaldoxime-dependent auxin

biosynthesis in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **106:**

5430-5435

Sun T, Yuan H, Cao H, Yazdani M, Tadmor Y, Li L (2018) Carotenoid metabolism in plants: the role

of plastids. *Molecular Plant* **11:** 58-74

Swarup R, Peret B (2012) AUX/LAX family of auxin influx carriers-an overview. *Frontiers in Plant*

Science **3:**225

Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki M, Harper R, Liscum E, Yamamoto K (2004)

MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the

transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl

and formation of lateral roots in *Arabidopsis thaliana*. *The Plant Cell* **16:** 379-393

Tichopad A, Dilger M, Schwarz G, Pfaffl MW (2003) Standardized determination of real-time PCR

efficiency from a single reaction set-up. *Nucleic Acids Research* **31:** e122

Timpte C, Lincoln C, Pickett FB, Turner J, Estelle M (1995) The *AXR1* and *AUX1* genes of

Arabidopsis function in separate auxin-response pathways. *The Plant Journal* **8:** 561-569

Toyokura K, Goh T, Shinohara H, Shinoda A, Kondo Y, Okamoto Y, Uehara T, Fujimoto K,

Okushima Y, Ikeyama Y, Nakajima K, Mimura T, Tasaka M, Matsubayashi Y, Fukaki H

(2019) Lateral inhibition by a peptide hormone-receptor cascade during *Arabidopsis* lateral

root founder cell formation. *Developmental Cell* **48:** 64-75

Tsugafune S, Mashiguchi K, Fukui K, Takebayashi Y, Nishimura T, Sakai T, Shimada Y,

Kasahara H, Koshiba T, Hayashi K (2017) Yucasin DF, a potent and persistent inhibitor of

auxin biosynthesis in plants. *Scientific Reports* **7:** 13992

Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya

Y, Shirasu K, Yoneyama K, Kyozuka J, Yamaguchi S (2008) Inhibition of shoot branching

by new terpenoid plant hormones. *Nature* **455:** 195-200

Vieten A, Sauer M, Brewer P, Friml J (2007) Molecular and cellular aspects of

auxin-transport-mediated development. *Trends in Plant Science* **12:** 160-168

Watahiki M, Yamamoto K (1997) The *massugu1* mutation of Arabidopsis identified with failure of

auxin-induced growth curvature of hypocotyl confers auxin insensitivity to hypocotyl and leaf.

Plant Physiology **115:** 419-426

Won C, Shen X, Mashiguchi K, Zheng Z, Dai X, Cheng Y, Kasahara H, Kamiya Y, Chory J, Zhao

Y (2011) Conversion of tryptophan to indole-3-acetic acid by TRYPTOPHAN

AMINOTRANSFERASES OF *ARABIDOPSIS* and YUCCAs in *Arabidopsis*. *Proceedings of*

the National Academy of Sciences, USA **108:** 18518-18523

Xu D, Miao J, Yumoto E, Yokota T, Asahina M, Watahiki M (2017) *YUCCA9*-mediated auxin

biosynthesis and polar auxin transport synergistically regulate regeneration of root systems

following root cutting. *Plant and Cell Physiology* **58:** 1710-1723

Yamamoto KT, Watahiki MK, Matsuzaki J, Satoh S, Shimizu H (2017) Space–time analysis of

gravitropism in etiolated Arabidopsis hypocotyls using bioluminescence imaging of the *IAA19*

promoter fusion with a destabilized luciferase reporter. *Journal of Plant Research* **130:**

765-777

Yuan H, Zhang J, Nageswaran D, Li L (2015) Carotenoid metabolism and regulation in horticultural

crops. *Horticulture Research* **2:** 15036

Zapata M, Rodriguez F, Garrido J (2000) Separation of chlorophylls and carotenoids from marine

phytoplankton: a new HPLC method using a reversed phase C_8 column and

pyridine-containing mobile phases. *Marine Ecology Progress Series* **195:** 29-45

Zhang H, Han W, De Smet I, Talboys P, Loya R, Hassan A, Rong H, Jürgens G, Paul Knox J,

Wang MH (2010) ABA promotes quiescence of the quiescent centre and suppresses stem cell

differentiation in the Arabidopsis primary root meristem. *The Plant Journal* **64:** 764-774

Zhao Y (2012) Auxin biosynthesis: a simple two-step pathway converts tryptophan to indole-3-acetic

acid in plants. *Molecular Plant* **5:** 334-338

Zhao Y, Merchant S, Briggs W, Ort D (2010) Auxin biosynthesis and its role in plant development.

Annual Review of Plant Biology **61:** 49-64

Zheng X, Yang Y, Al-Babili S (2021) Exploring the diversity and regulation of apocarotenoid

metabolic pathways in plants. *Frontiers in Plant Science* **12:** 787049

Figure legends

Figure 1. Inhibition of PR growth by fluridone mediates phytoene desaturase activity. A) Carotenoid biosynthesis pathway. PSY, phytoene synthase; PDS, phytoene desaturase; ZDS, ζ-carotene desaturase; CCR1/SDG8, carotenoid chloroplast regulatory1/set domain group 8; CCR2/CRTISO, carotenoid chloroplast regulatory2/crtiso; β-LCY, β-cyclase; CCD1/4/7/8, carotenoid cleavage dioxygenase 1/4/7/8. **B)** Five-day-old plants were transferred to medium with or without 12.8 μM fluridone. Photographs were captured 3 d after transfer. Scale bar = 5 mm. **C)** Leaf pigment bleaching effect of WT (left) and *mPDS* (right) by fluridone. Five-day-old plants were transferred to medium with or without 800 nM fluridone. Photographs were captured 3 d after transfer. Scale bar = 1 mm. **D–F)** Five-day-old plants were transferred to medium with or without fluridone. PR growth rate **(D)** was analyzed from 2 d after transfer (dat) to 3 dat. LR density **(E)** was counted at 3 dat. Total carotenoid (lutein, neoxanthin, violaxanthin, and β-carotene) **(F)** was quantified at 2 dat. Data represent the means ± standard error (SE) from 12 seedlings in **(D**, **E)** and from 3 biological replicates in **(F)**. *Significant differences compared to WT in **(D–F)** (*: $P < 0.05$, **: $P < 0.01$; Student's *t*-test).

Figure 2. Fluridone induces endogenous IAA levels. Five-day-old plants were transferred to mock or 800 nM fluridone medium. IAA concentration of WT was measured. Data represent the means ± SE from 3 biological replicates. *Significant differences compared to mock plants (*: $P < 0.05$; Student's *t*-test).

Figure 3. *DR5:LUC* expression in the root tip was increased following fluridone treatment. A, **B)** *DR5:LUC* activity by fluridone treatment. Five-day-old seedlings grown in medium containing 100 μM D-luciferin were transferred to plates containing 100 μM D-luciferin with or without 800 nM fluridone. *DR5:LUC* activity was measured with an interval of 1 h in the WT **(A)** and *mPDS* **(B)**. The hollow and solid black dotted lines indicate WT mock and fluridone-treated seedlings, and the hollow and solid red dotted lines indicate

mPDS mock and fluridone-treated seedlings, respectively. Data represent the means (relative

to the mean of time 0) ± SE from 5 seedlings.

Figure 4. Auxin biosynthesis, signaling, and influx transport are involved in the

downstream process that fluridone suppresses PDS. A–**C)** Fluridone resistance screening

of auxin-related mutants. Five-day-old plants were transferred to new medium with or without

800 nM fluridone. PR growth rate of auxin signaling and influx transport mutants **(A)** and

wei2, *wei7*, *wei2 wei7*, and *taa1/sav3* mutants **(B)** and *yucQ* mutant **(C)** was analyzed from 2

to 3 dat. The relative growth was compared with non fluridone-treated plants. The PR growth

rate non fluridone-treated plants is listed in Supplemental Table. S1. **D**, **E)** PR growth

inhibition cotreated with fluridone and auxinole **(D)** or YDF **(E)**. Five-day-old plants were

transferred to new medium with or without 800 nM fluridone and auxin inhibitors. PR growth

rate was analyzed from 2 to 3 dat. Data represent the means ± SE from 8 seedlings. Different

lowercase letters above the bars in **(A**, **B, D, E)** indicate significant differences at $P < 0.05$

(one-way Analysis of Variance [ANOVA] following Tukey-Kramer test). *Significant

differences compared to WT in **(C)** (**: $P < 0.01$; Student's *t*-test).

Figure 5. Carotenoid accumulation is negatively regulated by auxin production and

signaling. A, **B)** β-carotene quantitation by cotreatment of fluridone and auxin inhibitors in

root tissue. Five-day-old plants were transferred to new medium with or without 800 nM

fluridone and 10 µM auxinole **(A)** or 100 µM YDF **(B)**. The level of β-carotene in root tissue

was measured at 2 dat. The bars indicate mock (white), fluridone (black), auxinole or yucasin

DF (diagonal striped black) and auxinole+fluridone or YDF+fluridone (horizontal striped

black). **C)** β-carotene quantitation of the auxin-related mutants in root tissue. The level of

β-carotene in root tissue was measured at 7 d following germination. Data represent the

means ± SE from 3 biological replicates. n.d. indicates no detectable data. Different lowercase

letters above the bars in **(A**–**C)** indicate significant differences at $P < 0.05$ (one-way ANOVA

following Tukey-Kramer test).

 Figure 6. Fluridone activates auxin biosynthesis, response, and carotenoid catabolism

genes. Five-day-old WT and *mPDS* seedlings were transferred to new medium with or

without 800 nM fluridone. **A–D)** Relative expression of *IAA19*, *YUC9*, *CCD7*, and *CCD8* to

non fluridone-treated plants after fluridone treatment in the WT (open circle) and *mPDS* (open

square).   Data represent the means ± SE from 3 biological replicates.

Figure 7. Retinal restores fluridone-induced PR growth and cancels the exogenous auxin

effect. A) Five-day-old plants were transferred to new medium with or without 800 nM

fluridone and retinal. PR growth rate was analyzed from 1 to 2 dat. **B)** Five-day-old plants

were transferred to new medium with or without 20 µM retinal and NAA. **C)** Five-day-old

WT and auxin-related mutants were transferred to new medium with or without 20 µM retinal.

D) Five-day-old plants were transferred to new medium with or without 20 µM retinal and

KKI. The bars in **(B)** indicate 0 nM NAA (white), 10 nM NAA (diagonal striped black), and

20 nM NAA (black). PR growth rate was analyzed from 2 to 3 dat. The relative growth in **(C)**

was compared to non retinal-treated plants. **E)** Relative *pIAA19::ELucPEST* expression by

cotreatment of NAA and retinal. Four-day-old plants grown in 100 µM D-luciferin medium

were transferred to new medium containing 100 µM D-luciferin for a 1 d preincubation and

transferred to new medium containing 100 µM D-luciferin, 10 nM NAA and with or without

20 µM retinal. Time 0 indicates transfer to NAA-containing medium with or without retinal.

pIAA19 activity was measured with an interval of 1 h. Data represent the means ± SE from 8

seedlings in **(A–D)** and the means (relative to the means of time –1 h) ± SE from 3 biological

replicates in **(E)**. Different lowercase letters above the bars in **(B**, **C)** indicate significant

differences at $P < 0.05$ (one-way ANOVA following Tukey-Kramer test). *Significant

differences compared to non fluridone-treated plants in **(A)** (*: $P < 0.05$, **: $P < 0.01$;

Student's *t*-test). n.s. indicates no significant difference.

Figure 8. Putative model of carotenoid-mediated PR growth regulation. Fluridone

suppresses the activity of *PDS* and activates auxin content and response via the apocarotenoid

retinal. Retinal may trigger IAA catabolism pathway to control endogenous auxin levels.

893   Enhanced auxin response feedback for the carotenoid catabolism gene *CCD* activity in the

894   regulation of carotenoid content.

Figure 1. Inhibition of PR growth by fluridone mediates phytoene desaturase activity. A) Carotenoid biosynthesis pathway. PSY, phytoene synthase; PDS, phytoene desaturase; ZDS, ζ-carotene desaturase; CCR1/SDG8, carotenoid chloroplast regulatory1/set domain group 8; CCR2/CRTISO, carotenoid chloroplast regulatory2/crtiso; β-LCY, β-cyclase; CCD1/4/7/8, carotenoid cleavage dioxygenase 1/4/7/8. **B)** Five-day-old plants were transferred to medium with or without 12.8 μM fluridone. Photographs were captured 3 d after transfer. Scale bar = 5 mm. **C)** Leaf pigment bleaching effect of WT (left) and *mPDS* (right) by fluridone. Five-day-old plants were transferred to medium with or without 800 nM fluridone. Photographs were captured 3 d after transfer. Scale bar = 1 mm. **D–F)** Five-day-old plants were transferred to medium with or without fluridone. PR growth rate **(D)** was analyzed from 2 d after transfer (dat) to 3 dat. LR density **(E)** was counted at 3 dat. Total carotenoid (lutein, neoxanthin, violaxanthin, and β-carotene) **(F)** was quantified at 2 dat. Data represent the means ± standard error (SE) from 12 seedlings in **(D, E)** and from 3 biological replicates in **(F)**. *Significant differences compared to WT in **(D–F)** (*: $P < 0.05$, **: $P < 0.01$; Student's *t*-test).

Figure 2. Fluridone induces endogenous IAA levels. Five-day-old plants were transferred to mock or 800 nM fluridone medium. IAA concentration of WT was measured. Data represent the means ± SE from 3 biological replicates. *Significant differences compared to mock plants (*: $P < 0.05$; Student's t-test).

A

B

Figure 3. *DR5:LUC* expression in the root tip was increased following fluridone treatment. **A**, **B)** *DR5:LUC* activity by fluridone treatment. Five-day-old seedlings grown in medium containing 100 µM D-luciferin were transferred to plates containing 100 µM D-luciferin with or without 800 nM fluridone. *DR5:LUC* activity was measured with an interval of 1 h in the WT **(A)** and *mPDS* **(B)**. The hollow and solid black dotted lines indicate WT mock and fluridone-treated seedlings, and the hollow and solid red dotted lines indicate *mPDS* mock and fluridone-treated seedlings, respectively. Data represent the means (relative to the mean of time 0) ± SE from 5 seedlings.

Figure 4. Auxin biosynthesis, signaling, and influx transport are involved in the downstream process that fluridone suppresses PDS. A–C) Fluridone resistance screening of auxin-related mutants. Five-day-old plants were transferred to new medium with or without 800 nM fluridone. PR growth rate of auxin signaling and influx transport mutants **(A)** and *wei2*, *wei7*, *wei2 wei7*, and *taa1/sav3* mutants **(B)** and *yucQ* mutant **(C)** was analyzed from 2 to 3 dat. The relative growth was compared with non fluridone-treated plants. The PR growth rate non fluridone-treated plants is listed in Supplemental Table. S1. **D**, **E)** PR growth inhibition cotreated with fluridone and auxinole **(D)** or YDF **(E)**. Five-day-old plants were transferred to new medium with or without 800 nM fluridone and auxin inhibitors. PR growth rate was analyzed from 2 to 3 dat. Data represent the means ± SE from 8 seedlings. Different lowercase letters above the bars in **(A, B, D, E)** indicate significant differences at $P < 0.05$ (one-way Analysis of Variance [ANOVA] following Tukey-Kramer test). *Significant differences compared to WT in **(C)** (**: $P < 0.01$; Student's *t*-test).

A

B

C

Figure 5. Carotenoid accumulation is negatively regulated by auxin production and signaling. A, **B)** β-carotene quantitation by cotreatment of fluridone and auxin inhibitors in root tissue. Five-day-old plants were transferred to new medium with or without 800 nM fluridone and 10 µM auxinole **(A)** or 100 µM YDF **(B)**. The level of β-carotene in root tissue was measured at 2 dat. The bars indicate mock (white), fluridone (black), auxinole or yucasin DF (diagonal striped black) and auxinole+fluridone or YDF+fluridone (horizontal striped black). **C)** β-carotene quantitation of the auxin-related mutants in root tissue. The level of β-carotene in root tissue was measured at 7 d following germination. Data represent the means ± SE from 3 biological replicates. n.d. indicates no detectable data. Different lowercase letters above the bars in **(A–C)** indicate significant differences at *P* < 0.05 (one-way ANOVA following Tukey-Kramer test).

A

B

C

D

Figure 6. Fluridone activates auxin biosynthesis, response, and carotenoid catabolism genes. Five-day-old WT and *mPDS* seedlings were transferred to new medium with or without 800 nM fluridone. **A–D)** Relative expression of *IAA19*, *YUC9*, *CCD7*, and *CCD8* to non fluridone-treated plants after fluridone treatment in the WT (open circle) and *mPDS* (open square). Data represent the means ± SE from 3 biological replicates.

Figure 7. Retinal restores fluridone-induced PR growth and cancels the exogenous auxin effect. A) Five-day-old plants were transferred to new medium with or without 800 nM fluridone and retinal. PR growth rate was analyzed from 1 to 2 dat. **B)** Five-day-old plants were transferred to new medium with or without 20 µM retinal and NAA. **C)** Five-day-old WT and auxin-related mutants were transferred to new medium with or without 20 µM retinal. **D)** Five-day-old plants were transferred to new medium with or without 20 µM retinal and KKI. The bars in **(B)** indicate 0 nM NAA (white), 10 nM NAA (diagonal striped black), and 20 nM NAA (black). PR growth rate was analyzed from 2 to 3 dat. The relative growth in **(C)** was compared to non retinal-treated plants. **E)** Relative *pIAA19::ELucPEST* expression by cotreatment of NAA and retinal. Four-day-old plants grown in 100 µM D-luciferin medium were transferred to new medium containing 100 µM D-luciferin for a 1 d preincubation and transferred to new medium containing 100 µM D-luciferin, 10 nM NAA and with or without 20 µM retinal. Time 0 indicates transfer to NAA-containing medium with or without retinal. *pIAA19* activity was measured with an interval of 1 h. Data represent the means ± SE from 8 seedlings in **(A–D)** and the means (relative to the means of time –1 h) ± SE from 3 biological replicates in **(E)**. Different lowercase letters above the bars in **(B, C)** indicate significant differences at $P < 0.05$ (one-way ANOVA following Tukey-Kramer test). *Significant differences compared to non fluridone-treated plants in **(A)** (*: $P < 0.05$, **: $P < 0.01$; Student's *t*-test). n.s. indicates no significant difference.

Figure 8. Putative model of carotenoid-mediated PR growth regulation. Fluridone suppresses the activity of *PDS* and activates auxin content and response via the apocarotenoid retinal. Retinal may trigger IAA catabolism pathway to control endogenous auxin levels. Enhanced auxin response feedback for the carotenoid catabolism gene *CCD* activity in the regulation of carotenoid content.

Parsed Citations

Arias R, Dayan F, Michel A, Howell J, Scheffler B (2006) Characterization of a higher plant herbicide-resistant phytoene desaturase and its use as a selectable marker. Plant Biotechnology Journal 4: 263-273
Google Scholar: Author Only Title Only Author and Title

Auldridge M, McCarty D, Klee H (2006) Plant carotenoid cleavage oxygenases and their apocarotenoid products. Current Opinion in Plant Biology 9: 315-321
Google Scholar: Author Only Title Only Author and Title

Bainbridge K, Sorefan K, Ward S, Leyser O (2005) Hormonally controlled expression of the Arabidopsis MAX4 shoot branching regulatory gene. Plant Journal 44: 569-580
Google Scholar: Author Only Title Only Author and Title

Barlier I, Kowalczyk M, Marchant A, Ljung K, Bhalerao R, Bennett M, Sandberg G, Bellini C (2000) The SUR2 gene of Arabidopsis thaliana encodes the cytochrome P450 CYP83B1, a modulator of auxin homeostasis. Proceedings of the National Academy of Sciences, USA 97: 14819-14824
Google Scholar: Author Only Title Only Author and Title

Bartels PG, Watson CW (1978) Inhibition of carotenoid synthesis by fluridone and norflurazon. Weed Science 26: 198-203
Google Scholar: Author Only Title Only Author and Title

Bishopp A, Benkova E, Helariutta Y (2011) Sending mixed messages: auxin-cytokinin crosstalk in roots. Current Opinion in Plant Biology 14: 10-16
Google Scholar: Author Only Title Only Author and Title

Boerjan W, Cervera M-T, Delarue M, Beeckman T, Dewitte W, Bellini C, Caboche M, Van Onckelen H, Van Montagu M, Inzé D (1995) superroot, a recessive mutation in Arabidopsis, confers auxin overproduction. The Plant Cell 7: 1405-1419
Google Scholar: Author Only Title Only Author and Title

Breitenbach J, Sandmann G (2005) ζ-Carotene cis isomers as products and substrates in the plant poly-cis carotenoid biosynthetic pathway to lycopene. Planta 220: 785-793
Google Scholar: Author Only Title Only Author and Title

Casimiro I, Marchant A, Bhalerao R, Beeckman T, Dhooge S, Swarup R, Graham N, Inze D, Sandberg G, Casero P, Bennett M (2001) Auxin transport promotes Arabidopsis lateral root initiation. The Plant Cell 13: 843-852
Google Scholar: Author Only Title Only Author and Title

Cazzonelli C (2011) Carotenoids in nature: insights from plants and beyond. Functional Plant Biology 38: 833-847
Google Scholar: Author Only Title Only Author and Title

Cazzonelli C, Roberts A, Carmody M, Pogson B (2010) Transcriptional control of SET DOMAIN GROUP 8 and CAROTENOID ISOMERASE during Arabidopsis development. Molecular Plant 3: 174-191
Google Scholar: Author Only Title Only Author and Title

Chen Q, Dai X, De-Paoli H, Cheng Y, Takebayashi Y, Kasahara H, Kamiya Y, Zhao Y (2014) Auxin overproduction in shoots cannot rescue auxin deficiencies in Arabidopsis roots. Plant and Cell Physiology 55: 1072-1079
Google Scholar: Author Only Title Only Author and Title

Cheng Y, Dai X, Zhao Y (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in Arabidopsis. Genes & Development 20: 1790-1799
Google Scholar: Author Only Title Only Author and Title

Cheng Y, Dai X, Zhao Y (2007) Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in Arabidopsis. The Plant Cell 19: 2430-2439
Google Scholar: Author Only Title Only Author and Title

Delarue M, Prinsen E, Van Onckelen H, Caboche M, Bellini C (1998) Sur2 mutations of Arabidopsis thaliana define a new locus involved in the control of auxin homeostasis. The Plant Journal 14: 603-611
Google Scholar: Author Only Title Only Author and Title

Dharmasiri N, Dharmasiri S, Estelle M (2005) The F-box protein TIR1 is an auxin receptor. Nature 435: 441-445
Google Scholar: Author Only Title Only Author and Title

Dickinson A, Lehner K, Mi J, Jia K, Mijar M, Dinneny J, Al-Babili S, Benfey P (2019) β-Cyclocitral is a conserved root growth regulator. Proceedings of the National Academy of Sciences, USA 116: 10563-10567
Google Scholar: Author Only Title Only Author and Title

Dickinson A, Zhang J, Luciano M, Wachsman G, Sandoval E, Schnermann M, Dinneny J, Benfey P (2021) A plant lipocalin promotes retinal-mediated oscillatory lateral root initiation. Science 373: 1532-1536

Google Scholar: Author Only Title Only Author and Title

Du Y, Scheres B (2018) Lateral root formation and the multiple roles of auxin. Journal of Experimental Botany 69: 155-167
Google Scholar: Author Only Title Only Author and Title

Fukaki H, Tameda S, Masuda H, Tasaka M (2002) Lateral root formation is blocked by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of Arabidopsis. The Plant Journal 29: 153-168
Google Scholar: Author Only Title Only Author and Title

Fukaki H, Tasaka M (2009) Hormone interactions during lateral root formation. Plant Molecular Biology 69: 437-449
Google Scholar: Author Only Title Only Author and Title

Fukui K, Arai K, Tanaka Y, Aoi Y, Kukshal V, Jez JM, Kubes MF, Napier R, Zhao Y, Kasahara H (2022) Chemical inhibition of the auxin inactivation pathway uncovers the roles of metabolic turnover in auxin homeostasis. Proceedings of the National Academy of Sciences, USA 119: e2206869119
Google Scholar: Author Only Title Only Author and Title

Geisler M, Murphy A (2006) The ABC of auxin transport: The role of p-glycoproteins in plant development. FEBS Letters 580: 1094-1102
Google Scholar: Author Only Title Only Author and Title

Goh T, Kasahara H, Mimura T, Kamiya Y, Fukaki H (2012) Multiple AUX/IAA-ARF modules regulate lateral root formation: the role of Arabidopsis SHY2/IAA3-mediated auxin signalling. Philosophical Transactions of the Royal Society B: Biological Sciences 367: 1461-1468
Google Scholar: Author Only Title Only Author and Title

Goto C, Ikegami A, Goh T, Maruyama K, Kasahara H, Takebayashi Y, Kamiya Y, Toyokura K, Kondo Y, Ishizaki K, Mimura T, Fukaki H (2023) Genetic interaction between Arabidopsis SUR2/CYP83B1 and GNOM indicates the importance of stabilizing local auxin accumulation in lateral root initiation. Plant and Cell Physiology pcad084 .
Google Scholar: Author Only Title Only Author and Title

Grossmann K (2010) Auxin herbicides: current status of mechanism and mode of action. Pest Management Science 66: 113-120
Google Scholar: Author Only Title Only Author and Title

Hayashi K, Neve J, Hirose M, Kuboki A, Shimada Y, Kepinski S, Nozaki H (2012) Rational design of an auxin antagonist of the SCFTIR1 auxin receptor complex. ACS chemical biology 7: 590-598
Google Scholar: Author Only Title Only Author and Title

Hayward A, Stirnberg P, Beveridge C, Leyser O (2009) Interactions between auxin and strigolactone in shoot branching control. Plant Physiology 151: 400-412
Google Scholar: Author Only Title Only Author and Title

Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, Zimmermann P (2008) Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. Advances in Bioinformatics 2008: 420747
Google Scholar: Author Only Title Only Author and Title

Ito J, Fukaki H, Onoda M, Li L, Li C, Tasaka M, Furutani M (2016) Auxin-dependent compositional change in mediator in ARF7-and ARF19-mediated transcription. Proceedings of the National Academy of Sciences, USA 113: 6562-6567
Google Scholar: Author Only Title Only Author and Title

Jia K, Dickinson A, Mi J, Cui G, Xiao T, Kharbatia N, Guo X, Sugiono E, Aranda M, Blilou I, Rueping M, Benfey P, Al-Babili S (2019) Anchorene is a carotenoid-derived regulatory metabolite required for anchor root formation in Arabidopsis. Science Advances 5: eaaw6787
Google Scholar: Author Only Title Only Author and Title

Kasahara H (2016) Current aspects of auxin biosynthesis in plants. Bioscience Biotechnology and Biochemistry 80: 34-42
Google Scholar: Author Only Title Only Author and Title

Kobayashi K, Baba S, Obayashi T, Sato M, Toyooka K, Keranen M, Aro E, Fukaki H, Ohta H, Sugimoto K, Masuda T (2012) Regulation of root greening by light and auxin/cytokinin signaling in Arabidopsis. The Plant Cell 24: 1081-1095
Google Scholar: Author Only Title Only Author and Title

Koschmieder J, Wust F, Schaub P, Alvarez D, Trautmann D, Krischke M, Rustenholz C, Mano J, Mueller M, Bartels D, Hugueney P, Beyer P, Welsch R (2021) Plant apocarotenoid metabolism utilizes defense mechanisms against reactive carbonyl species and xenobiotics. Plant Physiology 185: 331-351
Google Scholar: Author Only Title Only Author and Title

Leyser O (2002) Molecular genetics of auxin signaling. Annual Review of Plant Biology 53: 377-398
Google Scholar: Author Only Title Only Author and Title

Li T, Kang X, Lei W, Yao X, Zou L, Zhang D, Lin H (2020) SHY2 as a node in the regulation of root meristem development by auxin,

brassinosteroids, and cytokinin. Journal of Integrative Plant Biology 62: 1500-1517
Google Scholar: Author Only Title Only Author and Title

Lincoln C, Britton JH, Estelle M (1990) Growth and development of the axr1 mutants of Arabidopsis. The Plant Cell 2: 1071-1080
Google Scholar: Author Only Title Only Author and Title

Liu PP, Montgomery TA, Fahlgren N, Kasschau KD, Nonogaki H, Carrington JC (2007) Repression of AUXIN RESPONSE
FACTOR10 by microRNA160 is critical for seed germination and post-germination stages. The Plant Journal 52: 133-146
Google Scholar: Author Only Title Only Author and Title

Mallory A, Bartel D, Bartel B (2005) MicroRNA-directed regulation of Arabidopsis AUXIN RESPONSE FACTOR17 is essential for
proper development and modulates expression of early auxin response genes. The Plant Cell 17: 1360-1375
Google Scholar: Author Only Title Only Author and Title

Mashiguchi K, Tanaka K, Sakai T, Sugawara S, Kawaide H, Natsume M, Hanada A, Yaeno T, Shirasu K, Yao H, McSteen P, Zhao Y,
Hayashi K, Kamiya Y, Kasahara H (2011) The main auxin biosynthesis pathway in Arabidopsis. Proceedings of the National
Academy of Sciences, USA 108: 18512-18517
Google Scholar: Author Only Title Only Author and Title

Moreno-Risueno M, Van Norman J, Moreno A, Zhang J, Ahnert S, Benfey P (2010) Oscillating gene expression determines
competence for periodic Arabidopsis root branching. Science 329: 1306-1311
Google Scholar: Author Only Title Only Author and Title

Okushima Y, Fukaki H, Onoda M, Theologis A, Tasaka M (2007) ARF7 and ARF19 regulate lateral root formation via direct
activation of LBD/ASL genes in Arabidopsis. The Plant Cell 19: 118-130
Google Scholar: Author Only Title Only Author and Title

Okushima Y, Overvoorde P, Arima K, Alonso J, Chan A, Chang C, Ecker J, Hughes B, Lui A, Nguyen D, Onodera C, Quach H,
Smith A, Yu G, Theologis A (2005) Functional genomic analysis of the AUXIN RESPONSE FACTOR gene family members in
Arabidopsis thaliana: unique and overlapping functions of ARF7 and ARF19. The Plant Cell 17: 444-463
Google Scholar: Author Only Title Only Author and Title

Olson JA, Hayaishi O (1965) The enzymatic cleavage of β-carotene into vitamin A by soluble enzymes of rat liver and intestine.
Proceedings of the National Academy of Sciences, USA 54: 1364-1370
Google Scholar: Author Only Title Only Author and Title

Paponov I, Teale W, Trebar M, Blilou K, Palme K (2005) The PIN auxin efflux facilitators: evolutionary and functional
perspectives. Trends in Plant Science 10: 170-177
Google Scholar: Author Only Title Only Author and Title

Park H, Kreunen S, Cuttriss A, DellaPenna D, Pogson B (2002) Identification of the carotenoid isomerase provides insight into
carotenoid biosynthesis, prolamellar body formation, and photomorphogenesis. The Plant Cell 14: 321-332
Google Scholar: Author Only Title Only Author and Title

Rodriguez-Concepcion M, Boronat A (2002) Elucidation of the methylerythritol phosphate pathway for isoprenoid biosynthesis in
bacteria and plastids. A metabolic milestone achieved through genomics. Plant Physiology 130: 1079-1089
Google Scholar: Author Only Title Only Author and Title

Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene signal transduction in Arabidopsis
thaliana: five novel mutant loci integrated into a stress response pathway. Genetics 139: 1393-1409
Google Scholar: Author Only Title Only Author and Title

Schmittgen T, Livak K (2008) Analyzing real-time PCR data by the comparative CT method. Nature Protocols 3: 1101-1108
Google Scholar: Author Only Title Only Author and Title

Schwartz S, Tan B, Gage D, Zeevaart J, McCarty D (1997) Specific oxidative cleavage of carotenoids by VP14 of maize. Science
276: 1872-1874
Google Scholar: Author Only Title Only Author and Title

Shimizu-Mitao Y, Kakimoto T (2014) Auxin sensitivities of all Arabidopsis Aux/IAAs for degradation in the presence of every
TIR1/AFB. Plant and Cell Physiology 55: 1450-1459
Google Scholar: Author Only Title Only Author and Title

Stepanova A, Hoyt J, Hamilton A, Alonso J (2005) A link between ethylene and auxin uncovered by the characterization of two
root-specific ethylene-insensitive mutants in Arabidopsis. The Plant Cell 17: 2230-2242
Google Scholar: Author Only Title Only Author and Title

Stepanova A, Robertson-Hoyt J, Yun J, Benavente L, Xie D, Dolezal K, Schlereth A, Jurgens G, Alonso J (2008) TAA1-mediated
auxin biosynthesis is essential for hormone crosstalk and plant development. Cell 133: 177-191
Google Scholar: Author Only Title Only Author and Title

Su L, Diretto G, Purgatto E, Danoun S, Zouine M, Li Z, Roustan J-P, Bouzayen M, Giuliano G, Chervin C (2015) Carotenoid accumulation during tomato fruit ripening is modulated by the auxin-ethylene balance. BMC Plant Biology 15: 1-12
Google Scholar: Author Only Title Only Author and Title

Sugawara S, Hishiyama S, Jikumaru Y, Hanada A, Nishimura T, Koshiba T, Zhao Y, Kamiya Y, Kasahara H (2009) Biochemical analyses of indole-3-acetaldoxime-dependent auxin biosynthesis in Arabidopsis. Proceedings of the National Academy of Sciences, USA 106: 5430-5435
Google Scholar: Author Only Title Only Author and Title

Sun T, Yuan H, Cao H, Yazdani M, Tadmor Y, Li L (2018) Carotenoid metabolism in plants: the role of plastids. Molecular Plant 11: 58-74
Google Scholar: Author Only Title Only Author and Title

Swarup R, Peret B (2012) AUX/LAX family of auxin influx carriers-an overview. Frontiers in Plant Science 3:225
Google Scholar: Author Only Title Only Author and Title

Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki M, Harper R, Liscum E, Yamamoto K (2004) MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. The Plant Cell 16: 379-393
Google Scholar: Author Only Title Only Author and Title

Tichopad A, Dilger M, Schwarz G, Pfaffl MW (2003) Standardized determination of real-time PCR efficiency from a single reaction set-up. Nucleic Acids Research 31: e122
Google Scholar: Author Only Title Only Author and Title

Timpte C, Lincoln C, Pickett FB, Turner J, Estelle M (1995) The AXR1 and AUX1 genes of Arabidopsis function in separate auxin-response pathways. The Plant Journal 8: 561-569
Google Scholar: Author Only Title Only Author and Title

Toyokura K, Goh T, Shinohara H, Shinoda A, Kondo Y, Okamoto Y, Uehara T, Fujimoto K, Okushima Y, Ikeyama Y, Nakajima K, Mimura T, Tasaka M, Matsubayashi Y, Fukaki H (2019) Lateral inhibition by a peptide hormone-receptor cascade during Arabidopsis lateral root founder cell formation. Developmental Cell 48: 64-75
Google Scholar: Author Only Title Only Author and Title

Tsugafune S, Mashiguchi K, Fukui K, Takebayashi Y, Nishimura T, Sakai T, Shimada Y, Kasahara H, Koshiba T, Hayashi K (2017) Yucasin DF, a potent and persistent inhibitor of auxin biosynthesis in plants. Scientific Reports 7: 13992
Google Scholar: Author Only Title Only Author and Title

Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya Y, Shirasu K, Yoneyama K, Kyozuka J, Yamaguchi S (2008) Inhibition of shoot branching by new terpenoid plant hormones. Nature 455: 195-200
Google Scholar: Author Only Title Only Author and Title

Vieten A, Sauer M, Brewer P, Friml J (2007) Molecular and cellular aspects of auxin-transport-mediated development. Trends in Plant Science 12: 160-168
Google Scholar: Author Only Title Only Author and Title

Watahiki M, Yamamoto K (1997) The massugu1 mutation of Arabidopsis identified with failure of auxin-induced growth curvature of hypocotyl confers auxin insensitivity to hypocotyl and leaf. Plant Physiology 115: 419-426
Google Scholar: Author Only Title Only Author and Title

Won C, Shen X, Mashiguchi K, Zheng Z, Dai X, Cheng Y, Kasahara H, Kamiya Y, Chory J, Zhao Y (2011) Conversion of tryptophan to indole-3-acetic acid by TRYPTOPHAN AMINOTRANSFERASES OF ARABIDOPSIS and YUCCAs in Arabidopsis. Proceedings of the National Academy of Sciences, USA 108: 18518-18523
Google Scholar: Author Only Title Only Author and Title

Xu D, Miao J, Yumoto E, Yokota T, Asahina M, Watahiki M (2017) YUCCA9-mediated auxin biosynthesis and polar auxin transport synergistically regulate regeneration of root systems following root cutting. Plant and Cell Physiology 58: 1710-1723
Google Scholar: Author Only Title Only Author and Title

Yamamoto KT, Watahiki MK, Matsuzaki J, Satoh S, Shimizu H (2017) Space–time analysis of gravitropism in etiolated Arabidopsis hypocotyls using bioluminescence imaging of the IAA19 promoter fusion with a destabilized luciferase reporter. Journal of Plant Research 130: 765-777
Google Scholar: Author Only Title Only Author and Title

Yuan H, Zhang J, Nageswaran D, Li L (2015) Carotenoid metabolism and regulation in horticultural crops. Horticulture Research 2: 15036
Google Scholar: Author Only Title Only Author and Title

Zapata M, Rodriguez F, Garrido J (2000) Separation of chlorophylls and carotenoids from marine phytoplankton: a new HPLC method using a reversed phase C8 column and pyridine-containing mobile phases. Marine Ecology Progress Series 195: 29-45

Google Scholar: Author Only Title Only Author and Title

Zhang H, Han W, De Smet I, Talboys P, Loya R, Hassan A, Rong H, Jürgens G, Paul Knox J, Wang MH (2010) ABA promotes quiescence of the quiescent centre and suppresses stem cell differentiation in the Arabidopsis primary root meristem. The Plant Journal 64: 764-774

Google Scholar: Author Only Title Only Author and Title

Zhao Y (2012) Auxin biosynthesis: a simple two-step pathway converts tryptophan to indole-3-acetic acid in plants. Molecular Plant 5: 334-338

Google Scholar: Author Only Title Only Author and Title

Zhao Y, Merchant S, Briggs W, Ort D (2010) Auxin biosynthesis and its role in plant development. Annual Review of Plant Biology 61: 49-64

Google Scholar: Author Only Title Only Author and Title

Zheng X, Yang Y, Al-Babili S (2021) Exploring the diversity and regulation of apocarotenoid metabolic pathways in plants. Frontiers in Plant Science 12: 787049

Google Scholar: Author Only Title Only Author and Title