Kang Xu[1], Haoran Zeng[1], Feiyang Lin[1], Emi Yumoto[2], Masashi Asahina[2, 3], Ken-ichiro

Hayashi[4], Hidehiro Fukaki[5], Hisashi Ito[6], Masaaki K. Watahiki[1, 7]

[1] Graduate School of Life Science, Hokkaido University, Sapporo, Japan

[2] Advanced Instrumental Analysis Center. Teikyo University, Utsunomiya, Japan

[3] Department of Biosciences, Teikyo University, Utsunomiya, Japan

[4] Department of Bioscience, Okayama University of Science, Okayama, Japan

[5] Department of Biology, Graduate School of Science, Kobe University, Kobe, Japan

[6] Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan

[7] Division of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo, Japan

Full title:

The exogenous application of the apocarotenoid retinaldehyde

negatively regulates auxin-mediated root growth

One-sentence summary

The short-term inhibition of carotenoid biosynthesis mediates the production of

β-carotene-derived apocarotenoid retinal which negatively regulates auxin levels and

signaling to control root growth.

List of authors:

Kang Xu: [ORCID ID: 0000-0001-6875-8027] kyokou950817@gmail.com

Haoran Zeng: [ORCID ID: 0009-0000-6707-4221] zenghaoran1998@outlook.com

Feiyang Lin: [ORCID ID: 0000-0003-4401-508X] feiyanglin05@gmail.com

Emi Yumoto: [ORCID ID: 0009-0005-2348-8410] yumoto@nasu.bio.teikyo-u.ac.jp

Masashi Asahina: [ORCID ID: 0000-0002-1828-0966] asahina@nasu.bio.teikyo-u.ac.jp

Ken-ichiro Hayashi: [ORCID ID: 0000-0002-9812-2801] ken-hayashi@ous.ac.jp

Hidehiro Fukaki: [ORCID ID: 0000-0002-6251-7668] h-fukaki@port.kobe-u.ac.jp

Hisashi Ito: [ORCID ID: 0000-0002-2482-5870] ito98@lowtem.hokudai.ac.jp

Masaaki K. Watahiki: [ORCID ID: 0000-0001-6669-9692] watahiki@sci.hokudai.ac.jp

Address correspondence to watahiki@sci.hokudai.ac.jp

The author responsible for distribution of materials integral to the findings represented in this

article in accordance with the policy described in the Instruction for Authors

(www.plantphysiol.org) is Masaaki K. Watahiki (watahiki@sci.hokudai.ac.jp).

ResearcherID: GFY-7778-2022

Abstract

Root development is essential for plant survival. The lack of carotenoid biosynthesis in the *phytoene desaturase 3* (*pds3*) mutant results in short primary roots (PR) and reduced lateral root (LR) formation. In this study, we show that short-term inhibition of PDS by fluridone suppressed PR growth in WT, but to a lesser extent in auxin mutants. Such an inhibition of PDS activity increased endogenous indole-3-acetic acid (IAA) levels, promoted auxin signaling, and partially complemented the PR growth of auxin deficient mutant, *YUCCA 3 5 7 8 9* quadruple mutant (*yucQ*). The exogenous application of retinaldehyde (retinal), an apocarotenoid derived from β-carotene, complemented the fluridone-induced suppression of root growth, as well as the short roots of the *pds3* mutant. Retinal also partially complemented the auxin-induced suppression of root growth. These results suggest that retinal may play a role in regulating root growth by modulating endogenous auxin levels.

Introduction

The root system plays a crucial role in water and nutrient uptake, anchoring plants to the soil, and serving as a source of signaling chemicals. Primary root (PR) elongation and lateral root (LR) branching are controlled by environmental cues and determine the architecture of the root system, allowing plants to adapt to the environment. The phytohormone auxin controls root growth and LR patterning (Moreno-Risueno et al., 2010; Saini et al., 2013; Chen et al., 2014). An early auxin signaling requires a family of repressor genes known as *AUXIN/INDOLE-3-ACETIC ACID* (*Aux/IAA*s). *Aux/IAA* dimerizes with the transcription factor *AUXIN RESPONSE FACTOR* (*ARF*) and suppresses the transcriptional function of *ARF*s (Leyser, 2002; De Smet et al., 2010). Application of exogenous auxin suppresses PR length and induces LR formation (Hobbie and Estelle 1994; Okumura et al., 2013). Overexpression of *ARF19* reduces PR length, suggesting a negative role for auxin signaling in PR elongation (Okushima et al., 2005). Polar auxin transport (PAT), which controls the flow of auxin to create an auxin gradient, is also involved in root growth and LR formation (Reed et al., 1998; Vieten et al., 2007). The auxin permease mutant, *auxin resistant 1* (*aux1*) exhibits both agravitropism and resistance to PR growth suppression by exogenous IAA (Yamamoto

and Yamamoto, 1998). The longer PRs of *aux1* compared to wild-type (WT) also suggest a negative role of endogenous auxin in PR growth (Roman et al., 1995).

YUCCA (*YUC*) encodes flavin monooxygenases which converts indole-3-pyruvic acid (IPyA) to indole-3-acetic acid (IAA) (Won et al., 2011; Zhao, 2012). *YUCCA* consists of a family of 11 genes and is highly redundant among them (Cheng et al., 2006; Xu et al., 2017). *YUC3*, *5*, *7*, *8*, and *9* are predominantly expressed in roots, and their quintuple mutant *yucQ* has a shorter PR and about half the amount of free IAA in roots compared to WT (Chen et al., 2014). The root meristem of *yucQ* is smaller and partially differentiated (Chen et al., 2014), suggesting that an appropriate level of auxin is required to maintain root meristem activity. Anthranilate is a precursor of tryptophan biosynthesis and contributes to IAA biosynthesis. The *WEAK ETHYLENE INSENSITIVE 2/ANTHRANILATE SYNTHASE α1* (*WEI2/ASA1*) and *WEI7/ANTHRANILATE SYNTHASE β1* (*WEI7/ASB1*) genes encode the subunits of anthranilate synthase, and their double mutant, *wei2 wei7* exhibits a dwarf phenotype with severely reduced PR length and IAA levels (Stepanova et al., 2005; Doyle et al., 2019).

Auxin regulates not only root development but also chloroplast development and chlorophyll accumulation. The *auxin resistant 1* (*axr1*) and *massugu1* (*msg1*)/*arf7* mutant, which are insensitive to auxin, are resistant to leaf chlorosis caused by 2,4-dichlorophenoxyacetic acid (2,4-D) (Watahiki and Yamamoto, 1997). The auxin signaling genes, *SOLITARY-ROOT/AUXIN/INDOLE-3-ACETIC ACID14* (*SLR/AUX/IAA14*), *ARF7*, and *ARF19*, are coupled with light and cytokinin signaling to regulate the gene sets associated with chloroplast development in the root (Kobayashi et al., 2012). Carotenoids are a diverse group of plastidial isoprenoid pigments that are essential for photosynthesis (Ruiz-Sola and Rodríguez-Concepción, 2012). In flowering plants, carotenoids are synthesized in the 2-C-methyl-D-erythritol 4-phosphate/1-deoxy-D-xylulose 5-phosphate (MEP/DOXP) pathway (Ruiz-Sola and Rodríguez-Concepción, 2012). PHYTOENE SYNTHASE (PSY) and PHYTOENE DESATURASE (PDS) are enzymes that convert geranylgeranyl pyrophosphate (GGPP) into phytoene and phytoene into ζ-carotene, respectively (Breitenbach and Sandmann, 2005; Rodríguez-Villalón et al., 2009). ζ-carotene then undergoes desaturation, isomerization and cyclization to generate carotenes like α-carotene and β-carotene (Ruiz-Sola and

Rodríguez-Concepción, 2012). Carotenoids act not only as photopigments but also as precursors of phytohormones, abscisic acid (ABA) and strigolactones (SL) (Nambara and Marion-Poll, 2005; Umehara et al., 2008). Besides, the discovery of new bioactive apocarotenoids, which are the products of the oxidative cleavage of carotenoids, has increased during this century (Auldridge et al., 2006; Zheng et al., 2021). β-cyclocitral is formed by the oxidative cleavage of the C7', C8' double bond position of β-carotene and promotes PR growth and LR formation (Rodrigo et al., 2013; Dickinson et al., 2019). Anchorene, formed by cutting the C11, C12 and C11', C12' bonds in almost all plant carotenoids, promotes the adventitious root formation by modulating auxin distribution (Jia et al., 2019). Impairment of apocarotenoid biosynthesis shows shorter roots and reduced LR formation compared to WT (Van Norman et al., 2014). This suggests a role for carotenoids and apocarotenoids in root growth regulation. The apocarotenoid retinaldehyde (retinal), a precursor of vitamin A and retinoic acid in vertebrates formed by the oxidative cleavage of the C15, C15' bond of β-carotene (Olson and Hayaishi, 1965), is mysteriously formed in flowering plants (Koschmieder et al., 2021), and is found to play a role in the periodic formation of LR (Dickinson et al., 2021).

In this study, we found that the herbicide fluridone (Bartels and Watson, 1978), which inhibits the enzymatic activity of PDS, suppressed root growth and partially mimicked the dwarf phenotype of the *pds3* mutant (Qin et al., 2007). This mechanism of root growth suppression remains to be elucidated. Here, we show that fluridone rapidly induces endogenous auxin levels and activates the auxin response. Genetic analysis revealed that this rapid response is dependent on PDS activity. Complementary approaches revealed that the apocarotenoid retinal negatively regulates endogenous auxin levels. Our results suggest that PDS activity positively regulates auxin catabolism to maintain auxin homeostasis via retinal.

Results

Fluridone suppresses the growth of the primary root

After transferring 5-day-old seedlings to fluridone medium or mock medium, the position of the root tip was marked and incubated for another 4 days. Fluridone suppressed PR growth in

a dose-dependent manner (**Fig. 1, B and C**), and bleached the newly emerged true leaves,

while the established cotyledons exhibited chlorosis (Supplemental Fig. S4A). The LR

number and density in the newly grown areas of PRs after transfer to fluridone medium and

the pre-grown areas of PRs were measured. The LR number in the pre-grown areas was

induced by fluridone. In contrast, fluridone reduced the number of LRs in the newly grown

areas (Supplemental Fig. S3). Overall, LR density was increased by fluridone (**Fig. 1D**).

Other PDS inhibitors, such as norflurazon, diflufenican and picolinafen, also suppressed PR

growth and increased LR density (Supplemental Fig. S2). To investigate the target specificity

of fluridone, a fluridone-insensitive trait of PDS (*35S::mHvPDS*) (Arias et al., 2006) was then

introduced into the *Arabidopsis* plants. The *35S::mHvPDS* plants were confirmed to have

resistance to fluridone in terms of chlorophyll and carotenoid contents (Supplemental Fig. S4,

B–G) (Arias et al., 2006). *35S::mHvPDS* also showed resistance to fluridone in the

suppression of PR growth and induction of LR formation (**Fig. 1, C and D**). These results

indicate that PDS is the target of fluridone for the suppression of PR growth and induction of

LR formation. Subsequently, carotenoid-deficient mutants were examined for fluridone

sensitivity. The phytoene synthase mutant *psy* showed reduced PR growth and LR formation

(Van Norman et al., 2014) (Supplemental Fig. S1). PDS is encoded by a single gene, *PDS3*,

in *Arabidopsis thaliana* (Qin et al., 2007), and its T-DNA insertion mutant, *pds3-1* (McElver

et al., 2001; Meinke, 2019), also showed reduced PR growth rate and LR density (**Fig. 1, E–**

G). The *pds3-1* plants were insensitive to fluridone for both PR growth suppression and LR

density induction (**Fig. 1, E–G**), further indicating that PDS is a target for fluridone. Unlike

the *pds3-1* plants, the *psy* plants showed weak resistance to fluridone compared to *pds3-1*, and

their LR formation was induced by fluridone (Supplemental Fig. S1, A and B). These results

indicate that the change in root architecture induced by fluridone is specifically dependent on

the activity of PDS. Fluridone suppressed PR growth immediately after application, and the

rate of suppression increased after 60 h of treatment (Supplemental Fig. S6).

Fluridone inhibits the biosynthesis of ABA and SLs by inhibiting their precursor β-carotene

(Yoshioka et al., 1998; López-Ráez et al., 2008). ABA and SLs promote root growth at low

doses (Zhang et al., 2010; Ruyter-Spira et al., 2011). Therefore, we investigated whether

exogenous ABA or the SL analog *racemic*-GR24 (*rac*-GR24) could complement

fluridone-induced PR growth inhibition. The growth suppression of PR was not reversed by

neither ABA nor *rac*-GR24 (Supplemental Fig. S5, G and H). The biosynthesis and signaling

mutants of ABA or SLs were also tested for fluridone sensitivity. All the mutants examined

showed comparable responses to fluridone compared to WT (Supplemental Fig. S5, A–C).

These results suggest that ABA and SLs play a minor role in fluridone-induced PR growth

suppression. The suppression of PR growth suggests an activation of auxin signaling by

fluridone, as the increase in LR formation is under the control of auxin signaling (Du and

Scheres, 2018).

Fluridone increases endogenous auxin levels and activates auxin signaling

The levels of endogenous auxin in roots were measured during the temporary inhibition of

PDS activity. The WT plants showed an increase in IAA levels after 8 h and 24 h of fluridone

treatment (**Fig. 2**; Supplemental Fig. S7A). In contrast, the *35S::mHvPDS* plants showed no

significant change of it (Supplemental Fig. S7B). The expression of the artificial auxin

reporter *DR5::luciferase* (*DR5:LUC*) (Moreno-Risueno et al., 2010) was then measured. The

DR5:LUC expression was transiently activated at the root tip regardless of the presence or

absence of fluridone, which may be a response to plant stress (**Fig. 3**). After this transient

activation, the *DR5:LUC* expression in fluridone-treated WT plants remained higher than in

mock-treated plants after 4 h (**Fig. 3A**), while no such induction was detected in

35S::mHvPDS plants (**Fig. 3B**). These results indicate that the fluridone-activated auxin

response, like the induction of endogenous auxin, depends on PDS activity. The expression of

DR5:LUC in fluridone-treated WT plants, but not in *35S::mHvPDS* plants, was reduced to

below the mock level after 140 h (**Fig. 3**). This data suggests that the temporary inhibition of

PDS activity leads to an increase in auxin levels and activation of the auxin signaling pathway,

whereas long-term inhibition of PDS activity reduces the auxin response, possibly due to a

carotenoid deficiency.

Carotenoid-mediated regulation of root growth requires auxin biosynthesis and signaling

It is hypothesized that the temporary inhibition of PDS activity suppresses PR growth by increasing endogenous auxin levels and activating auxin signaling. To test this hypothesis, mutants in auxin signaling or auxin biosynthesis were examined. Indeed, the auxin permease mutant, *aux1-21*, and the ubiquitin-activating enzyme E1 mutant, *axr1-12* (Lincoln et al., 1990), were resistant to fluridone. *AUX/IAA* gain-of-function mutants *aux/iaa19/msg2-1* (Tatematsu et al., 2004) and *aux/iaa14/slr-1* (Fukaki et al., 2002), and the downstream transcription factor mutant *arf7-1 arf19-1* (Okushima et al., 2005) were weakly resistant to fluridone (**Fig. 4A**). *axr1-12* and *aux1-7* are reported to show a high level of auxin resistance on root elongation, whereas *msg2-1*, *slr-1*, and *arf7-1 arf19-1* exhibit relatively low or no resistance to auxin on root growth (Timpte et al., 1995; Fukaki et al., 2002; Tatematsu et al., 2004; Okushima et al., 2005). These results suggest that there is a correlation between fluridone resistance and auxin resistance in the auxin mutants.

Optimal levels of auxin are required for root growth, and both supraoptimal and suboptimal levels of auxin reduce root growth (Zhao, 2018). The root growth of the *wei2-1 wei7-1* double mutant was insensitive to fluridone, while its single mutant showed moderate or no resistance to fluridone, indicating the redundancy of these genes (**Fig. 4B**). The quintuple *YUCCA* mutant, *yucQ*, exhibited a severe root defect due to a reduction in endogenous auxin levels. Surprisingly, fluridone tended not to suppress the PR growth in the *yucQ* plants but did promote it (**Fig. 4C**; >100% of inhibition). This suggests that the inhibition of PDS activity increases endogenous auxin levels (**Fig. 2**), which compensates for the growth defect in the roots of *yucQ* (Chen et al., 2014). None of the *yuc* single mutants showed resistance to fluridone, indicating the redundancy of *YUCCA* genes (Supplemental Fig. S8, A and B). These results suggest that PDS activity negatively regulates endogenous auxin levels.

Pharmaceutical interference with auxin signaling and biosynthesis was then combined with the effect of fluridone. Auxinole is an antagonist of the auxin receptor TRANSPORT INHIBITOR RESPONSE 1/AUXIN SIGNALING F-BOX PROTEIN (TIR1/AFB) (Hayashi et al., 2012). Yucasin difluorinated analog (YDF) is a competitive inhibitor of YUCCA flavin

monooxygenase (Tsugafune et al., 2017). Both auxinole and YDF suppressed PR growth, but

the addition of fluridone to either inhibitor did not enhance the suppression of PR growth (**Fig.**

4, D and E). These results support the hypothesis that fluridone-induced suppression of PR

growth requires an increase in endogenous auxin and proper auxin signaling. An inhibitor of

auxin efflux transport, *N*-1-naphthylphthalamic acid (NPA), also suppressed PR growth in a

dose-dependent manner (Reed et al., 1998). In contrast to auxinole and YDF, NPA showed an

additive effect to that of fluridone in suppressing PR growth (Supplemental Fig. S9B),

suggesting that the efflux transporter may not be the primary target of PDS inhibition.

Negative control of carotenoid levels by auxin

The albino phenotype in the *pds3-1* and *psy* plants (**Fig. 1E**; Supplemental Fig. S1C) indicates

the essential role of carotenoid biosynthesis in chloroplast development (Qin et al., 2007). The

development of such chloroplasts is repressed by auxin signaling (Kobayashi et al., 2012).

arf7/msg1 and *axr1* are resistant to 2,4-D-induced chlorosis (Watahiki and Yamamoto, 1997),

suggesting that carotenoid contents are also negatively regulated by auxin. Auxinole and YDF

increased β-carotene levels in roots, whereas the application of fluridone abolished the

β-carotene levels regardless of the presence of auxinole or YDF (**Fig. 5, A and B**). The

auxin-deficient mutants, *wei2-1 wei7-1* and *yucQ* showed doubled β-carotene levels compared

to WT (**Fig. 5C**), while auxin signaling mutants showed insignificant increased β-carotene

levels (Supplemental Fig. S10C) which may be due to the redundant functions in these genes.

These data suggest that endogenous auxin levels and signaling negatively control β-carotene

levels. In addition, this negative regulation of carotenoid levels by auxin was also observed in

the shoots (Supplemental Fig. S10, A and B).

Expression of auxin- and carotenoid-related genes

To investigate whether PDS activity regulates auxin biosynthesis and signaling at the gene

expression level, the expression of auxin-related genes in roots was measured after fluridone

treatment. The expression of an early auxin-induced gene, *AUX/IAA19* or auxin biosynthesis

gene, *YUCCA9*, increased approximately 4 and 2.5 times, respectively, after 4 h of fluridone

treatment in WT, but not in *35S::mHvPDS* (**Fig. 6, A and B**). The induced expression of these genes was associated with the induction time of the *DR5:LUC* expression (**Fig. 3A**). These data suggest that the temporary inhibition of PDS activity activates auxin biosynthesis and signaling.

The expression of genes in carotenoid biosynthesis (*CAROTENOID CHLOROPLAST REGULATOR2* [*CCR2*]), *CCR1* which regulates the expression of *CCR2* (Ruiz-Sola and Rodríguez-Concepción, 2012), and the carotenoid catabolism pathway (*CAROTENOID CLEAVAGE DIOXYGENASE 1* [*CCD1*], *CCD7* and *CCD8*) (Yuan et al., 2015) (**Fig. 1A**) was examined following fluridone treatment. The expression of *CCD7* and *CCD8* was strongly induced at 8 and 24 h in WT plants (**Fig. 6, C and D**). In contrast, the expression of *CCR1*, *CCR2* and *CCD1* showed little induction in comparison to *CCD7* and *CCD8* (Supplemental Fig. S11). These data suggest that after the temporary inhibition of PDS activity, carotenoid catabolism is activated rather than biosynthesis. *CCD7* and *CCD8* are reported to be induced by auxin (Bainbridge et al., 2005; Hayward et al., 2009). The GENEVESTIGATOR database (https://genevestigator.com) (Hruz et al., 2008) also shows that *CCD7* expression is induced within 0.5 to 1 h following exogenous IAA treatment (Supplemental Fig. S15). These suggest that the increased auxin levels by the inhibition of PDS activity likely feedback to upregulate carotenoid catabolism.

Fluridone-mediated PR growth regulation in relation to retinal and auxin levels

Recent evidence has shown that the apocarotenoids β-cyclocitral and retinal, both derived from the oxidation of β-carotene, regulate root architecture (Olson and Hayaishi, 1965; Rodrigo et al., 2013; Dickinson et al., 2019; Dickinson et al., 2021). Fluridone potentially blocks the biosynthesis of such apocarotenoids and may result in the suppression of PR growth. Therefore, both β-cyclocitral and retinal were investigated to complement the suppression of PR growth by fluridone application. Retinal, but not β-cyclocitral, fully restored the PR growth suppression induced by fluridone (**Fig. 7A**; Supplemental Fig. S12). In addition, retinal also promoted the growth of PRs in the *pds3-1* mutant (**Fig. 7B**),

suggesting that retinal is responsible for the short root phenotype of *pds3-1*. This also supports the hypothesis that the retinal is responsible for the suppression of root growth by fluridone. To further investigate the relationship between auxin signaling and retinal, we examined the expression of the auxin reporter *pIAA19::emerald luciferase-PEST* (*pIAA19::Eluc-PEST*) (Yamamoto et al., 2017), which was induced two-fold at the root tip by 10 nM 1-naphthalene acetic acid (NAA), a synthetic auxin analog. However, this induction was suppressed by the cotreatment with retinal (**Fig. 7E**). This result indicates that retinal suppresses the exogenous auxin response and raises the hypothesis that retinal may complement the inhibitory effect of auxin on root growth. Indeed, retinal restored the NAA-induced PR growth suppression in WT (**Fig. 7B**). Conversely, the short PR phenotype in *wei2-1 wei7-1* and *yucQ* was enhanced by the application of retinal (**Fig. 7C**). This suggests a further suppression of auxin levels by retinal in these mutants. In addition, retinal also showed an antagonistic effect to auxin on the LR formation induced by NAA and tryptophan (Supplemental Fig. S13, D and E). *SUPERROOT2/FEWER ROOT SUPPRESSOR1/CYP83B1* (*SUR2/FSP1*) is one of the cytochrome P450 gene family members of the indole-3-acetaldoxime (IAOx) pathway, which represents a non-IPyA pathway for IAA biosynthesis (Barlier et al., 2000; Sugawara et al., 2009). *sur2* mutant and the weaker allele *fsp1* mutant exhibit high LR density due to an overaccumulation of IAA (Delarue et al., 1998; Goto et al., 2023). Such a high LR density of *sur2* was also reduced by retinal (Supplemental Fig. S13G). These data are consistent with a negative regulation of endogenous auxin levels by retinal. However, no significant reduction in IAA levels was detected in the roots of retinal-treated WT plants (Supplemental Fig. S13I). Kakeimide (KKI), an inhibitor of the IAA-amino acid conjugating enzyme GRETCHEN HAGEN 3 (GH3s), accumulates endogenous IAA rapidly (Fukui et al., 2022) and suppressed PR growth in a dose-dependent manner (**Fig. 7D**), similar to the treatment with exogenous auxin (Supplemental Fig. S13C). However, unlike the case of exogenous auxin (**Fig. 7B**; Supplemental Fig. S13D), retinal did not counteract the KKI-induced suppression of PR growth and induction of LR formation (**Fig. 7D**; Supplemental Fig. S13H). Although the levels of retinal in fluridone-treated plants remain to be determined, these results suggest that

307 retinal may target the degradation of auxin, thereby exerting an antagonistic effect on auxin.

Discussion

Inhibition of PDS activity increases the levels of auxin in roots

The regulation of root growth by temporary inhibition of carotenoid biosynthesis was the

subject of this study. Here we showed that a temporary application of fluridone increased

endogenous auxin levels in the roots but not in the shoots (**Fig. 2**; Supplemental Table 2).

Such a temporary inhibition of PDS activity may exceed the optimal concentration of auxin

levels for the root growth, thereby resulting in the suppression of PR growth and the

promotion of LR formation (**Fig. 1, C and D**). The endogenous auxin levels in *yucQ* were

suboptimal, and the short roots of *yucQ* could be rescued by a low dose application of

exogenous IAA (5 nM) (Chen et al., 2014). The partial recovery of root growth in the short

roots of *yucQ* correlates with the root-specific increase in auxin levels by the inhibition of

PDS activity (**Fig. 2 and 4C**). The *sur2/fsp1* and *yuc1D* plants accumulate endogenous auxin

and increase the growth of hypocotyls (Delarue et al., 1998; Zhao et al., 2001). However, the

length of hypocotyls was similar between fluridone-treated and mock-treated plants (data not

shown), confirming a root-specific induction of auxin by fluridone. Like the *yucQ* plants, the

wei2 wei7 (*asa1 asb1*) plants showed a reduction in endogenous IAA levels and a short root

phenotype. Although the root of *wei2 wei7* showed resistance to fluridone, it did not restore

the root growth of *wei2-1 wei7-1*, in contrast to that of *yucQ* (**Fig. 4, B and C**). A low dose

application of anthranilic acid, a precursor of IAA and tryptophan, partially recovered the root

growth in *wei2-1 wei7-1* without rescuing its endogenous IAA levels (Doyle et al., 2019).

This suggests that the short roots of the *wei2-1 wei7-1* plants require anthranilic acid rather

than increasing the suboptimal concentration of IAA to compensate their growth. In fact, a

higher concentration of anthranilic acid significantly but not completely rescued root growth

of *wei2-1 wei7-1* (Doyle et al., 2019). Low dose application of exogenous NAA did not

rescue or inhibit root growth in *wei2-1 wei7-1* (Supplemental Fig. S8C), further supporting

that endogenous auxin is suboptimal and it is not a primary requirement for rescuing the PR

growth in *wei2-1 wei7-1*. The inhibition of auxin signaling and biosynthesis by auxinole and

YDF, respectively, suppressed PR growth; however, fluridone addition did not further

enhance this suppression (**Fig. 4, D and E**). In contrast to auxinole and YDF, exogenous

auxin and fluridone exhibited an additive effect on PR growth suppression (Supplemental Fig.

S9A). This suggests that the optimal level of auxin and auxin signaling are required for PR

growth, and that excessive or insufficient levels of auxin can reduce PR growth. Taken

together, fluridone-induced PDS inhibition increases root auxin levels, activates auxin

signaling, and thus suppresses PR growth.

The roles of apocarotenoids on fluridone-mediated root growth

The temporary inhibition of PDS activity triggers the auxin-mediated suppression of PR

growth. However, the downstream product of PDS activity responsible for this process has

yet to be identified. The extremely hydrophobic nature of β-carotene made it impossible for

the complementation test by exogenous application to the plants (data not shown), which is

inhibited by fluridone. Although ABA and SLs are the candidates for their close interactions

with auxin in the regulation of root architecture (Ruyter-Spira et al., 2011; Emenecker and

Strader, 2020), this study provides no evidence that they are responsible for the temporary

inhibition of PDS activity on the root architecture regulation (Supplemental Fig. S5).

Consistently, the LR formation of the excised tomato roots is also induced by fluridone, even

in the presence of ABA (Hooker and Thorpe, 1998).

Dickinson et al. (2021) report that retinal (1 μM) induces LR capacity. In this study, the

application of a higher dose of retinal (20 μM) showed antagonistic effects to auxin on the PR

growth and the LR formation (**Fig. 7, B and E**; Supplemental Fig. S13, C, D, E and G).

Although retinal partially complemented the short roots of the *pds3-1* plants (**Fig. 7B**), it did

not alter LR density (Supplemental Fig. S13D). 20 μM retinal also did not alter PR growth or

LR density in the WT plants (**Fig. 7B**; Supplemental Fig. S13D). Instead, the sites of LR

emergence in retinal-treated WT plants were shifted away from the root tip compared to those

of mock-treated WT plants (Supplemental Fig. S13F). Therefore, it is speculated that retinal

suppressed the auxin response during the development of the LR primordia and delayed the

emergence of LRs, resulting in longer zones of unbranched PR. It should be noted that

Dickinson et al. (2021) measure the LR capacity by excising the root tips to encourage LR

formation which evokes the root cut response (Xu et al., 2017), which differs from this study.

The severe short root phenotype in *yucQ* is further repressed by YDF due to a further

inhibition of auxin biosynthesis in *yucQ* (Tsugafune et al., 2017). Although retinal did not

consistently alter endogenous free IAA levels (Supplemental Fig. S13I), it further suppressed

PR growth in the auxin-deficient mutants (**Fig. 7C**) and had an additive effect on the

YDF-induced suppression of root growth (data not shown). Instead, the antagonistic effect of

retinal on auxin was abolished by the auxin catabolism inhibitor KKI (**Fig. 7D**; Supplemental

Fig. S13H), suggesting that retinal may target downstream substrates of GH3s in the auxin

catabolic pathway. The study of mutants in the auxin catabolic pathway on fluridone and

retinal may be required for this confirmation. Although the expression of *YUC9* was enhanced

by fluridone treatment (**Fig. 6B**), a single *yuc9* mutant did not show resistance to fluridone

(Supplemental Fig. S8A). This suggests that the fluridone-induced accumulation of

endogenous IAA may be primarily contributed by the suppression of the auxin catabolic

pathway rather than by the activation of the auxin biosynthesis pathway.

The difference between continuous carotenoid deficiency and temporary inhibition of PDS activity

Van Norman et al. (2014) and this study showed that both the *psy* and *pds3-1* plants had

shorter PR lengths compared to WT (**Fig. 1, E and F**; Supplemental Fig. S1, A and C).

Similarly, temporary exposure to fluridone reduced PR growth. Short-term fluridone

treatment led to rapid and modest suppression of root growth, whereas long-term exposure

(>60 h) resulted in accelerated suppression of root growth (Supplemental Fig. S6), which may

correspond to the short PRs in *psy* and *pds3-1*. The root growth of fluridone-treated WT

plants was fully restored by retinal on the second day after treatment, but the extent of root

growth recovery in *pds3-1* was less than that of WT (Supplemental Fig. S14). By the third day

after treatment (>60 h), such retinal-mediated recovery of root growth became ineffective in

WT, whereas it remained at the same level in *pds3-1* (Supplemental Fig. S13A and S14).

Temporary application of fluridone resulted in the degradation of chlorophylls, leading to

chlorosis in established cotyledons and bleaching in newly emerged leaves (Supplemental Fig.

S4A) (Kim et al., 2004), while the albino *pds3* mutant lost the ability to differentiate plastids

(Qin et al., 2007). The difference in root growth recovery between the fluridone-exposed

plants and the *pds3-1* plants may reflect the process of plastid degradation by fluridone and

undifferentiated proplastids, respectively, suggesting a further role of proplastid

differentiation for root growth.

Although both *psy* and *pds3-1* reduced the number of LRs (**Fig. 1G**; Supplemental Fig.

S1B), fluridone increased the number of LRs in the pre-grown areas of PRs and decreased it

in the newly grown areas of PRs in WT (Supplemental Fig. S3), which corresponds to the

activation of the *DR5:LUC* expression in the upper part of PRs (Supplemental Movie. S1). An

induction of LR formation by fluridone has been reported in the excised tomato roots (Hooker

and Thorpe, 1998). It is speculated that the process of plastid degradation increases auxin

levels in pre-grown roots, while newly grown roots remain undifferentiated proplastids and

reduce LRs, as in the *psy* and *pds3-1* plants (Supplemental Fig. S3) (Van Norman et al., 2014).

This hypothesis is consistent with the result that the *DR5:LUC* expression at the root tips was

increased upon fluridone application, and eventually decreased upon long-term incubation

(**Fig. 3A**).

Conclusion

In this study, we found that the apocarotenoid retinal controls the growth of PRs through

endogenous auxin levels (**Fig. 8**). Temporary inhibition of PDS activity blocks retinal

biosynthesis and then leads to auxin accumulation to suppress root growth. Long-term

exposure to fluridone further suppresses root growth, the underlying mechanisms of which

remain to be elucidated. The role of retinal here may be different from that of promoting LR

capacity (Dickinson et al., 2021). Retinal may regulate auxin catabolism and auxin

homeostasis. In addition, auxin may further regulate carotenoid accumulation through a

feedback mechanism involving the expression of genes associated with carotenoid catabolism.

This model provides insight into the role of the apocarotenoid retinal in root growth

regulation. The quantitative analysis of retinal and its derivatives, as well as their targets that

may regulate auxin homeostasis, is warranted in the future.

Materials and methods

Plant materials and growth conditions

The *Arabidopsis thaliana* mutants and WT plants as well as *PDS* transgenic plants were in the

Columbia background, except for *yuc5*, which is in the Landsberg *erecta* background. *pds3-1*

used in this work refers to *pigment defective 226* (*pde226-1*). The *arf7-1 arf19-1*, *slr-1*, *yuc1*,

yuc2, *yuc3*, *yuc4-1*, *yuc5*, *yuc6*, *yuc7*, *yuc8*, *yuc9*, *yuc10*, and *yuc11* plants were described

previously (Xu et al., 2017). The *axr1-12* (Lincoln et al., 1990), *aux1-21* (Roman et al., 1995),

aba1-6 (Niyogi et al., 1998), *aba2-1* (González-Guzmán et al., 2002), *psy* (Alonso et al., 2003)

and *pde226-1* (McElver et al., 2001) plants were obtained from the Arabidopsis Biological

Resource Center. *wei2-1*, *wei7-1*, *wei2-1 wei7-1* (Stepanova et al., 2005) and *sur2* (Delarue et

al., 1998) plants were provided from Dr Hidehiro Fukaki (Kobe University). *aba3-2*

(Koornneef et al., 1982), *abi1c* (Koornneef et al., 1984), *abi3-9*, *abi4-11*, *abi5-7* (Nambara et

al., 2002), *ost1-1* (Mustilli et al., 2002) and *pyrQ* (Park et al., 2009) plants were provided

from Dr Masanori Okamoto (RIKEN Center for Sustainable Resource Science). *max1-1*,

max2-1 (Stirnberg et al., 2002) and *max3-9* (Booker et al., 2004) plants were provided from Dr

Kiyoshi Tatematsu (National Institute for Basic Biology). The *yucQ* mutant was a generous

gift from Dr Hiroyuki Kasahara (Tokyo University of Agriculture and Technology) with an

agreement from Dr Yunde Zhao (University of California San Diego). *DR5:LUC*

(Moreno-Risueno et al., 2010) was kindly provided from Dr Philip N. Benfey (Duke

University). In this study, *DR5:LUC* reporter in the *DR5:LUC/VENUS* line (Toyokura et al.,

2019) was observed. Fluridone, norflurazon, deflufenican, picolinafen, and retinal were

purchased from Fujifilm Wako Pure Chemical. KKI, YDF, and auxinole were provided by

Kei-ichiro Hayashi. Seeds were surface-sterilized using chlorine gas for at least 45 min,

suspended in 0.38% agarose, and sown on half-strength Murashige and Skoog (MS) medium

(Duchefa Biochemie) supplemented with 1% (w/v) sucrose, 0.6% (w/v) gellan gum, and 0.5

mM MES (pH 5.8). Stratification was performed at 4°C for 3 d in the dark. Plants were grown

on vertically oriented plates at 22°C under constant light if not specifically indicated (80–110

μmol m^{-2} s^{-1}). Stock solutions of phytohormones and inhibitors were prepared in dimethyl

sulfoxide and filtered through a 0.45 μM disc filter.

PDS-resistant vector and plant transformation and screening

The plasmid pPDSer1303 was kindly provided by Renée S. Arias (USDA ARS, USA). The

pds mutation vector construction was described previously (Arias et al., 2006). The plasmids

were transformed into *Agrobacterium* strain C58C1 and used to transform *Arabidopsis*

thaliana via the dip floral method. *Agrobacterium* harboring the plasmid pCambia1303

without *pds* was used as a control for the transformations and later as a control for herbicide

screening. Seeds from T1, T2, and T3 generations were screened with hygromycin (50 mg/L)

and fluridone (25 nM) separately. Seedlings that were resistant to hygromycin in T1 were

selected in either hygromycin or fluridone (25 nM) screening in T2 and T3. T3 seeds were

screened for resistance to fluridone at 25, 50, and 200 nM.

Quantification of LR density and PR growth

Plants were grown on vertical plates for 5 d and then transferred to a new half-strength MS

medium with or without chemicals indicated. Plant images were captured using a flatbed

scanner (GT-X980 and DS-50000, EPSON) within 4 d after transfer. The LR density (the

number of emerged LRs besides an adventitious root/the root length from the root–shoot

junction to the youngest LR toward the root tip) (Dubrovsky and Forde, 2012) and the PR

growth rate were analyzed with Fiji software (https://fiji.sc/).

Quantification of PR growth via time-lapse imaging

475 Plants were grown on vertical plates for 4 d and then transferred to a new half-strength MS

476 medium. Plant images were captured using a digital camera (Lumix G5, Panasonic). The

477 interval was 30 min. After 1 d of photography, the seedlings were transferred to a new

478 half-strength MS medium with or without 800 nM fluridone. The PR growth rate was

479 analyzed using Fiji software (https://fiji.sc/).

Quantification of IAA

Seeds were sown in sterilized 300 μm nylon mesh and transferred to control or fluridone plates after 5-d germination together with mesh. Shoot and root samples were harvested at specific time points after transfer and frozen in liquid nitrogen. Hormonome analysis was conducted as previously described (Xu et al., 2017). Briefly, samples of ~100 mg fresh weight (Fw) were suspended in 80% (v/v) aqueous methanol with $[^2H_5]$-IAA as internal standards. The samples were homogenized and the supernatant was loaded onto a Bond Elut C18 cartridge (100 mg, 3 ml; Agilent Technologies) and eluted with 80% (v/v) aqueous methanol. The concentrated samples were subjected to liquid chromatography with electrospray ionization tandem mass spectrometry comprising a quadrupole tandem mass spectrometer (Agilent 6460 Triple Quadrupole mass spectrometer) with an electrospray ion source and an Agilent 1200 separation module. The raw data were extracted from MassHunter software (Agilent Technologies) and examined in Excel (Microsoft).

Bioluminescence imaging analysis

D-Luciferin was purchased from Gold Biotechnology. Four-day-old plants grown on 100 μM D-luciferin medium were transferred to new D-luciferin medium for a 1 d preincubation and transferred to new D-luciferin medium with or without chemicals indicated. Time-lapse imaging was performed using a camera (ASI294MMPro, ZWO) with control software (SharpCap, v4.0). The images were captured at an interval of 1 h, including 15 min of exposure and 45 min of red and blue light (100 μmol m^{-2} s^{-1}) and then saved in a TIFF format for further analysis using Fiji software (https://fiji.sc/).

RNA isolation and quantitative reverse transcription polymerase chain reaction analysis

Seeds were sown on control medium with 300 μm nylon mesh. Roots were placed on plates containing no herbicide (mock) or 800 nM fluridone, root samples were harvested at 0, 4, 8, and 24 h and frozen in liquid nitrogen. Total RNA was extracted and purified using an RNA isolation kit (ISOSPIN Plant RNA, NIPPON GENE). Complementary DNA was synthesized

from the total RNA according to the manufacturer's instructions (ReverTra Ace qPCR RT Master Mix, Toyobo). Quantitative reverse transcription polymerase chain reaction (qRT–PCR) was performed in optical 96-well plates via the LightCycler 480 II system (Roche Life Science) using KOD SYBR qPCR Mix (Toyobo). Primer pairs spanning exon–exon junctions were designed using the QuantPrime program to avoid genomic DNA amplification as listed in Supplemental Table S3 (Xu et al., 2017). The specificity of the reactions was verified via melt curve analysis and capillary electrophoresis (Multina, Shimadzu). Standard curve analysis was used to evaluate the efficiency of the reactions. ACTIN2 was used as an internal standard. The qRT–PCR program included one cycle at 98°C for 2 min, followed by 40 cycles of 98°C for 10 s, 60°C for 10 s, and 68°C for 30 s. The cycle time values were determined using the second derivative maximum method (Tichopad et al., 2003) via LightCycler software (version 1.5, Roche Life Science). The data were analyzed using the comparative threshold cycle (Ct) method ($2^{-\triangle\triangle Ct}$) (Schmittgen and Livak, 2008).

Quantification of carotenoid via high-performance liquid chromatography (HPLC) analysis

Seeds were sown on control medium with 300 μm nylon mesh. The seedlings were grown under the conditions described above. Five days after germination, the seedlings together with nylon mesh were transferred to plates with or without chemicals indicated. Shoot tissues (~20 mg) or root tissues (~100 mg) were harvested at specific time points and stored at −80°C. Total carotenoids were extracted using acetone wako 1[st] grade (FUJIFILM Wako Chemicals). Acetone was added to the tissue samples that had been mixed with beads and silica gel and shaken for 2 min. Total carotenoids were separated via centrifugation for 10 min at 14000 rpm at 4°C. High-performance liquid chromatography analysis for carotenoid quantification was performed as previously described (Zapata et al., 2000). An aliquot of the sample was subjected to a C8 column (4.6 × 150 mm, Waters Symmetry C8, Waters) equilibrated with A solution (methanol:acetonitrile:aqueous pyridine solution [0.25 M pyridine] = 2:1:1 [v:v:v]). Pigments were eluted with a linear gradient of 0%–40% B solution (methanol:acetonitrile:acetone = 1:3:1 [v:v:v]) for 2 min, 40%–95% for 6 min, an isocratic

539 elution for 5 min, a linear gradient of 95%–0% for 2 min, and an isocratic elution for 7 min at

540 a flow rate of 1 mL/min. Pigments were identified based on their retention time relative to

541 known standards and quantified by peak area on the chromatogram.

Statistical analysis

544 All data shown in graphs and tables are represented as the means ± SE from 3 replicates,

545 except for Supplemental Fig. S2, which is performed once. The statistical significance is

546 determined by 2-tailed Student's *t*-test (*: $P < 0.05$, **: $P < 0.01$) or one-way ANOVA

547 following Tukey-Kramer test (Different lowercase letters indicate significant differences, $P <$

548 0.05).

Author contributions

551 Author contributions: K.X. conceived the project and designed the experiments, which were

552 mostly carried out and analyzed by K.X.; H.Z. and F.L. supported retinal-related experiment

553 design, process and data analysis; E.Y. and M.A. were responsible for IAA quantification and

554 data collection; K.X. and M.K.W. wrote the article. K.X., H.Z., F.L., E.Y., M.A., K.H., H.F.,

555 H.I. and M.K.W. read, commented and approved the final manuscript.

Supplemental data

The following materials are available in the online version of this article.

Supplemental Figure S1. The phenotypes and fluridone resistance of the *psy* mutant.

Supplemental Figure S2. PDS inhibitors exhibit a fluridone-like effect on PR suppression

and LR induction.

Supplemental Figure S3. Fluridone induces LR formation in the pre-grown areas of PR

but severely reduces it in the newly grown areas of PR.

Supplemental Figure S4. Fluridone reduces chlorophyll and carotenoid contents, and

mPDS plants are resistant to it.

Supplemental Figure S5. ABA and SLs are not involved in fluridone-mediated regulation

of root architecture.

Supplemental Figure S6. The kinetic growth rate of PRs in WT and *mPDS* after fluridone

treatment.

Supplemental Figure S7. Long-period change of auxin levels by fluridone.

Supplemental Figure S8. The *yuc* single mutants have weak or no resistance to fluridone,

and *wei2-1 wei7-1* shows resistance to low dose of exogenous NAA.

Supplemental Figure S9. Exogenous auxin and NPA suppress PR growth in an additive

manner with fluridone.

Supplemental Figure S10. Carotenoid contents in the shoots are negatively regulated by

auxin, and auxin signaling mutants insignificantly increase carotenoid contents in the roots.

Supplemental Figure S11. The expression of *CCR1*, *CCR2* and *CCD1* is somewhat

induced by fluridone.

Supplemental Figure S12. β-cyclocitral is not involved in carotenoid-mediated regulation

of PR growth.

Supplemental Figure S13. Retinal is responsible for the fluridone effect and regulates

auxin levels and response.

Supplemental Figure S14. The different effect of retinal to fluridone-treated WT plants

and *pds3-1* plants in root growth recovery.

Supplemental Figure S15. *CCD7* expression is rapidly induced by exogenous IAA

treatment.

Supplemental Table S1. PR growth rate in mock-treated auxin-related mutants.

Supplemental Table S2. Endogenous IAA concentration in fluridone treatment in the

shoots.

Supplemental Table S3. Gene-specific primers used for qRT–PCR analysis.

Supplemental Movie S1. The *DR5:LUC* bioluminescence video under fluridone

treatment in WT and *mPDS*.

Acknowledgments

We thank Dr. Renée Arias (United States Department of Agriculture) and Dr. Franck Dayan

(Colorado State University) for the supply of plasmid *35S::mHvPDS* and for the kind

reviewing of the manuscript. This research was funded by grants-in-aid for scientific research

from the Japan Society for the Promotion of Science (JSPS) (JP19H05673 to H. F. and M. K.

W., JP21K19097 to K.H.), Japan Science and Technology Agency Support for Pioneering

Research Initiated by the Next Generation (JST SPRING) (Grant Number JPMJSP2119 to K.

Y.) and ACRO Research grant of Teikyo University (TeTe20-01 to M.A.).

conflict of interest statement. The authors have no conflicts of interests to declare.

References

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen HM, Shinn P, Stevenson DK, Zimmerman J,

Barajas P, Cheuk R, Gadrinab C, Heller C, Jeske A, Koesema E, Meyers CC, Parker H,

Prednis L, Ansari Y, Choy N, Deen H, Geralt M, Hazari N, Hom E, Karnes M,

Mulholland C, Ndubaku R, Schmidt I, Guzman P, Aguilar-Henonin L, Schmid M,

Weigel D, Carter DE, Marchand T, Risseeuw E, Brogden D, Zeko A, Crosby WL, Berry

CC, Ecker JR (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science*

301: 653-657

Arias R, Dayan F, Michel A, Howell J, Scheffler B (2006) Characterization of a higher plant

herbicide-resistant phytoene desaturase and its use as a selectable marker. *Plant Biotechnology*

Journal **4:** 263-273

Auldridge M, McCarty D, Klee H (2006) Plant carotenoid cleavage oxygenases and their

apocarotenoid products. *Current Opinion in Plant Biology* **9:** 315-321

Bainbridge K, Sorefan K, Ward S, Leyser O (2005) Hormonally controlled expression of the

Arabidopsis *MAX4* shoot branching regulatory gene. *Plant Journal* **44:** 569-580

Barlier I, Kowalczyk M, Marchant A, Ljung K, Bhalerao R, Bennett M, Sandberg G, Bellini C

(2000) The *SUR2* gene of *Arabidopsis thaliana* encodes the cytochrome P450 CYP83B1, a

modulator of auxin homeostasis. *Proceedings of the National Academy of Sciences, USA* **97:**

14819-14824

Bartels PG, Watson CW (1978) Inhibition of carotenoid synthesis by fluridone and norflurazon. *Weed*

Science **26:** 198-203

Booker J, Auldridge M, Wills S, McCarty D, Klee H, Leyser O (2004) MAX3/CCD7 is a carotenoid

cleavage dioxygenase required for the synthesis of a novel plant signaling molecule. *Current*

Biology **14:** 1232-1238

Breitenbach J, Sandmann G (2005) ζ-Carotene *cis* isomers as products and substrates in the plant

poly-*cis* carotenoid biosynthetic pathway to lycopene. *Planta* **220:** 785-793

Chen Q, Dai X, De-Paoli H, Cheng Y, Takebayashi Y, Kasahara H, Kamiya Y, Zhao Y (2014)

Auxin overproduction in shoots cannot rescue auxin deficiencies in Arabidopsis roots. *Plant*

and Cell Physiology **55:** 1072-1079

Cheng Y, Dai X, Zhao Y (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls

the formation of floral organs and vascular tissues in *Arabidopsis*. *Genes & Development* **20:**

1790-1799

De Smet I, Lau S, Voss U, Vanneste S, Benjamins R, Rademacher EH, Schlereth A, De Rybel B,

Vassileva V, Grunewald W, Naudts M, Levesque MP, Ehrismann JS, Inzé D, Luschnig C,

Benfey PN, Weijers D, Van Montagu MCE, Bennett MJ, Jürgens G, Beeckman T (2010)

Bimodular auxin response controls organogenesis in *Arabidopsis*. *Proceedings of the National*

Academy of Sciences, USA **107:** 2705-2710

Delarue M, Prinsen E, Van Onckelen H, Caboche M, Bellini C (1998) *Sur2* mutations of

Arabidopsis thaliana define a new locus involved in the control of auxin homeostasis. *The*

Plant Journal **14:** 603-611

Dickinson AJ, Lehner K, Mi J, Jia K, Mijar M, Dinneny JR, Al-Babili S, Benfey P (2019)

β-Cyclocitral is a conserved root growth regulator. *Proceedings of the National Academy of*

Sciences, USA **116:** 10563-10567

Dickinson AJ, Zhang J, Luciano M, Wachsman G, Sandoval E, Schnermann M, Dinneny JR,

Benfey P (2021) A plant lipocalin promotes retinal-mediated oscillatory lateral root initiation.

Science **373:** 1532-1536

Doyle SM, Rigal A, Grones P, Karady M, Barange DK, Majda M, Pařízková B, Karampelias M,

Zwiewka M, Pěnčík A, Almqvist F, Ljung K, Novák O, Robert S (2019) A role for the

auxin precursor anthranilic acid in root gravitropism via regulation of PIN-FORMED protein

polarity and relocalisation in *Arabidopsis*. *New Phytologist* **223:** 1420-1432

Du Y, Scheres B (2018) Lateral root formation and the multiple roles of auxin. *Journal of*
Experimental Botany **69:** 155-167

Dubrovsky JG, Forde BG (2012) Quantitative analysis of lateral root development: pitfalls and how
to avoid them. *Plant Cell* **24:** 4-14

Emenecker RJ, Strader LC (2020). Auxin-abscisic acid interactions in plant growth and development.
Biomolecules **10:** 281

Fukaki H, Tameda S, Masuda H, Tasaka M (2002) Lateral root formation is blocked by a
gain-of-function mutation in the *SOLITARY-ROOT/IAA14* gene of *Arabidopsis*. *The Plant*
Journal **29:** 153-168

Fukui K, Arai K, Tanaka Y, Aoi Y, Kukshal V, Jez JM, Kubes MF, Napier R, Zhao Y, Kasahara H
(2022) Chemical inhibition of the auxin inactivation pathway uncovers the roles of metabolic
turnover in auxin homeostasis. *Proceedings of the National Academy of Sciences, USA* **119:**
e2206869119

González-Guzmán M, Apostolova N, Bellés JM, Barrero JM, Piqueras P, Ponce MR, Micol JL,
Serrano R, Rodríguez PL (2002) The short-chain alcohol dehydrogenase ABA2 catalyzes
the conversion of xanthoxin to abscisic aldehyde. *Plant Cell* **14:** 1833-1846

Goto C, Ikegami A, Goh T, Maruyama K, Kasahara H, Takebayashi Y, Kamiya Y, Toyokura K,
Kondo Y, Ishizaki K, Mimura T, Fukaki H (2023) Genetic interaction between Arabidopsis
SUR2/CYP83B1 and *GNOM* indicates the importance of stabilizing local auxin accumulation
in lateral root initiation. *Plant and Cell Physiology* pcad084

Hayashi K, Neve J, Hirose M, Kuboki A, Shimada Y, Kepinski S, Nozaki H (2012) Rational design
of an auxin antagonist of the SCF^{TIR1} auxin receptor complex. *ACS Chemical Biology* **7:**
590-598

Hayward A, Stirnberg P, Beveridge C, Leyser O (2009) Interactions between auxin and strigolactone
in shoot branching control. *Plant Physiology* **151:** 400-412

Hobbie L, Estelle M (1994) Genetic approaches to auxin action. *Plant, Cell & Environment* **17:**
525-540

Hooker TS, Thorpe TA (1998) Effects of fluridone and abscisic acid on lateral root initiation and root

elongation of excised tomato roots cultured *in vitro*. *Plant Cell, Tissue and Organ Culture* **52:**

199-203

Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W,

Zimmermann P (2008) Genevestigator v3: a reference expression database for the

meta-analysis of transcriptomes. *Advances in Bioinformatics* **2008:** 420747

Jia K, Dickinson A, Mi J, Cui G, Xiao T, Kharbatia N, Guo X, Sugiono E, Aranda M, Blilou I,

Rueping M, Benfey P, Al-Babili S (2019) Anchorene is a carotenoid-derived regulatory

metabolite required for anchor root formation in *Arabidopsis*. *Science Advances* **5:** eaaw6787

Kim JS, Yun BW, Choi JS, Kim TJ, Kwak SS, Cho KY (2004) Death mechanisms caused by

carotenoid biosynthesis inhibitors in green and in undeveloped plant tissues. *Pesticide*

Biochemistry and Physiology **78:** 127-139

Kobayashi K, Baba S, Obayashi T, Sato M, Toyooka K, Keranen M, Aro E, Fukaki H, Ohta H,

Sugimoto K, Masuda T (2012) Regulation of root greening by light and auxin/cytokinin

signaling in *Arabidopsis*. *The Plant Cell* **24:** 1081-1095

Koornneef M, Jorna ML, Brinkhorst-van der Swan DLC, Karssen CM (1982) The isolation of

abscisic acid (ABA) deficient mutants by selection of induced revertants in non-germinating

gibberellin sensitive lines of *Arabidopsis thaliana* (L.) Heynh. *Theoretical and Applied*

Genetics **61:** 385-393

Koornneef M, Reuling G, Karssen CM (1984) The isolation and characterization of abscisic

acid-insensitive mutants of *Arabidopsis thaliana*. *Physiologia Plantarum* **61:** 377-383

Koschmieder J, Wust F, Schaub P, Alvarez D, Trautmann D, Krischke M, Rustenholz C, Mano J,

Mueller M, Bartels D, Hugueney P, Beyer P, Welsch R (2021) Plant apocarotenoid

metabolism utilizes defense mechanisms against reactive carbonyl species and xenobiotics.

Plant Physiology **185:** 331-351

Leyser O (2002) Molecular genetics of auxin signaling. *Annual Review of Plant Biology* **53:** 377-398

Lincoln C, Britton JH, Estelle M (1990) Growth and development of the *axr1* mutants of *Arabidopsis*.

The Plant Cell **2:** 1071-1080

López-Ráez JA, Charnikhova T, Gómez-Roldán V, Matusova R, Kohlen W, De Vos R, Verstappen

F, Puech-Pages V, Bécard G, Mulder P, Bouwmeester H (2008) Tomato strigolactones are

derived from carotenoids and their biosynthesis is promoted by phosphate starvation. *New*

Phytologist **178:** 863-874

McElver J, Tzafrir I, Aux G, Rogers R, Ashby C, Smith K, Thomas C, Schetter A, Zhou Q,

Cushman MA, Tossberg J, Nickle T, Levin JZ, Law M, Meinke D, Patton D (2001)

Insertional mutagenesis of genes required for seed development in *Arabidopsis thaliana*.

Genetics **159:** 1751-1763

Meinke DW (2019) Genome-wide identification of *EMBRYO-DEFECTIVE* (*EMB*) genes required for

growth and development in *Arabidopsis. New Phytologist* **226:** 306-325

Moreno-Risueno M, Van Norman J, Moreno A, Zhang J, Ahnert S, Benfey P (2010) Oscillating

gene expression determines competence for periodic *Arabidopsis* root branching. *Science* **329:**

1306-1311

Mustilli A-C, Merlot S, Vavasseur A, Fenzi F, Giraudat J (2002) Arabidopsis OST1 protein kinase

mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive

oxygen species production. *Plant Cell* **14:** 3089-3099

Nambara E, Marion-Poll A (2005) Abscisic acid biosynthesis and catabolism. *Annual Review of Plant*

Biology **56:** 165-185

Nambara E, Suzuki M, Abrams S, McCarty DR, Kamiya Y, McCourt P (2002) A screen for genes

that function in abscisic acid signaling in *Arabidopsis thaliana*. *Genetics* **161:** 1247-1255

Niyogi KK, Grossman AR, Björkman O (1998) Arabidopsis mutants define a central role for the

xanthophyll cycle in the regulation of photosynthetic energy conversion. *Plant Cell* **10:**

1121-1134

Okumura K, Goh T, Toyokura K, Kasahara H, Takebayashi Y, Mimura T, Kamiya Y, Fukaki H

(2013) GNOM/FEWER ROOTS is required for the establishment of an auxin response

maximum for Arabidopsis lateral root initiation. *Plant and Cell Physiology* **54:** 406-417

Okushima Y, Overvoorde P, Arima K, Alonso J, Chan A, Chang C, Ecker J, Hughes B, Lui A,

Nguyen D, Onodera C, Quach H, Smith A, Yu G, Theologis A (2005) Functional genomic

analysis of the *AUXIN RESPONSE FACTOR* gene family members in *Arabidopsis thaliana*:

unique and overlapping functions of *ARF7* and *ARF19*. *The Plant Cell* **17:** 444-463

Olson JA, Hayaishi O (1965) The enzymatic cleavage of β-carotene into vitamin A by soluble

enzymes of rat liver and intestine. *Proceedings of the National Academy of Sciences, USA* **54:**

1364-1370

Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A,

Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ, Desveaux D, Rodriguez PL,

Mccourt P, Zhu JK, Schroeder JI, Volkman BF, Cutler SR (2009) Abscisic acid inhibits

type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* **324:**

1068-1071

Qin G, Gu H, Ma L, Peng Y, Deng XW, Chen Z, Qu LJ (2007) Disruption of phytoene desaturase

gene results in albino and dwarf phenotypes in *Arabidopsis* by impairing chlorophyll,

carotenoid, and gibberellin biosynthesis. *Cell Research* **17:** 471-482

Reed RC, Brady SR, Muday G (1998) Inhibition of auxin movement from the shoot into the root

inhibits lateral root development in Arabidopsis. *Plant Physiology* **118:** 1369-1378

Rodrigo MJ, Alquézar B, Alós E, Medina V, Carmona L Bruno M, Al-Babili S, Zacarías L (2013)

A novel carotenoid cleavage activity involved in the biosynthesis of *Citrus* fruit-specific

apocarotenoid pigments. *Journal of Experimental Botany* **64:** 4461-4478

Rodríguez-Villalón A, Gas E, Rodríguez-Concepción M (2009) Phytoene synthase activity controls

the biosynthesis of carotenoids and the supply of their metabolic precursors in dark-grown

Arabidopsis seedlings. *Plant Journal* **60:**424-435

Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene

signal transduction in *Arabidopsis thaliana*: five novel mutant loci integrated into a stress

response pathway. *Genetics* **139:** 1393-1409

Ruiz-Sola MÁ, Rodríguez-Concepción M (2012) Carotenoid biosynthesis in Arabidopsis: a colorful

pathway. *Arabidopsis Book* **10:** e0158

Ruyter-Spira C, Kohlen W, Charnikhova T, van Zeijl A, van Bezouwen L, de Ruijter N, Cardoso

C, Lopez-Raez JA, Matusova R, Bours R, Verstappen F, Bouwmeester H (2011)

Physiological effects of the synthetic strigolactone analog GR24 on root system architecture in

Arabidopsis: another belowground role for strigolactones? *Plant Physiology* **155:** 721-734

Saini S, Sharma I, Kaur N, Pati PK (2013) Auxin: a master regulator in plant root development.

Plant Cell Report **32:** 741-757

Schmittgen T, Livak K (2008) Analyzing real-time PCR data by the comparative C_T method. Nature

Protocols **3:** 1101-1108

Stepanova A, Hoyt J, Hamilton A, Alonso J (2005) A link between ethylene and auxin uncovered by

the characterization of two root-specific ethylene-insensitive mutants in Arabidopsis. *The*

Plant Cell **17:** 2230-2242

Stirnberg P, van de Sande K, Leyser HMO (2002) MAX1 and MAX2 control shoot lateral branching

in *Arabidopsis*. *Development* **129:** 1131-1141

Sugawara S, Hishiyama S, Jikumaru Y, Hanada A, Nishimura T, Koshiba T, Zhao Y, Kamiya Y,

Kasahara H (2009) Biochemical analyses of indole-3-acetaldoxime-dependent auxin

biosynthesis in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **106:**

5430-5435

Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki M, Harper R, Liscum E, Yamamoto K (2004)

MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the

transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl

and formation of lateral roots in *Arabidopsis thaliana*. *The Plant Cell* **16:** 379-393

Tichopad A, Dilger M, Schwarz G, Pfaffl MW (2003) Standardized determination of real-time PCR

efficiency from a single reaction set-up. *Nucleic Acids Research* **31:** e122

Timpte C, Lincoln C, Pickett FB, Turner J, Estelle M (1995) The *AXR1* and *AUX1* genes of

Arabidopsis function in separate auxin-response pathways. *The Plant Journal* **8:** 561-569

Toyokura K, Goh T, Shinohara H, Shinoda A, Kondo Y, Okamoto Y, Uehara T, Fujimoto K,

Okushima Y, Ikeyama Y, Nakajima K, Mimura T, Tasaka M, Matsubayashi Y, Fukaki H

(2019) Lateral inhibition by a peptide hormone-receptor cascade during *Arabidopsis* lateral

root founder cell formation. *Developmental Cell* **48:** 64-75

Tsugafune S, Mashiguchi K, Fukui K, Takebayashi Y, Nishimura T, Sakai T, Shimada Y,

Kasahara H, Koshiba T, Hayashi K (2017) Yucasin DF, a potent and persistent inhibitor of

auxin biosynthesis in plants. *Scientific Reports* **7:** 13992

Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya

Y, Shirasu K, Yoneyama K, Kyozuka J, Yamaguchi S (2008) Inhibition of shoot branching

by new terpenoid plant hormones. *Nature* **455:** 195-200

Van Norman JM, Zhang J, Cazzonelli CI, Pogson BJ, Harrison PJ, Bugg TD, Chan KX,
Thompson AJ, Benfey PN (2014) Periodic root branching in *Arabidopsis* requires synthesis
of an uncharacterized carotenoid derivative. *Proceedings of the National Academy of Sciences,*
USA **111:** E1300-E1309

Vieten A, Sauer M, Brewer P, Friml J (2007) Molecular and cellular aspects of
auxin-transport-mediated development. *Trends in Plant Science* **12:** 160-168

Watahiki M, Yamamoto K (1997) The *massugu1* mutation of Arabidopsis identified with failure of
auxin-induced growth curvature of hypocotyl confers auxin insensitivity to hypocotyl and leaf.
Plant Physiology **115:** 419-426

Won C, Shen X, Mashiguchi K, Zheng Z, Dai X, Cheng Y, Kasahara H, Kamiya Y, Chory J, Zhao
Y (2011) Conversion of tryptophan to indole-3-acetic acid by TRYPTOPHAN
AMINOTRANSFERASES OF *ARABIDOPSIS* and YUCCAs in *Arabidopsis*. *Proceedings of*
the National Academy of Sciences, USA **108:** 18518-18523

Xu D, Miao J, Yumoto E, Yokota T, Asahina M, Watahiki M (2017) *YUCCA9*-mediated auxin
biosynthesis and polar auxin transport synergistically regulate regeneration of root systems
following root cutting. *Plant and Cell Physiology* **58:** 1710-1723

Yamamoto KT, Watahiki MK, Matsuzaki J, Satoh S, Shimizu H (2017) Space–time analysis of
gravitropism in etiolated Arabidopsis hypocotyls using bioluminescence imaging of the *IAA19*
promoter fusion with a destabilized luciferase reporter. *Journal of Plant Research* **130:**
765-777

Yamamoto M, Yamamoto KT (1998) Differential effects of 1-naphthaleneacetic acid, indole-3-acetic
acid and 2,4-dichlorophenoxyacetic acid on the gravitropic response of roots in an
auxin-resistant mutant of Arabidopsis, *auxl*. *Plant and Cell Physiology* **39:** 660-664

Yoshioka T, Endo T, Satoh S (1998) Restoration of seed germination at supraoptimal temperatures by
fluridone, an inhibitor of abscisic acid biosynthesis. *Plant and Cell Physiology* **39:** 307-312

Yuan H, Zhang J, Nageswaran D, Li L (2015) Carotenoid metabolism and regulation in horticultural
crops. *Horticulture Research* **2:** 15036

Zapata M, Rodriguez F, Garrido J (2000) Separation of chlorophylls and carotenoids from marine
phytoplankton: a new HPLC method using a reversed phase C_8 column and

pyridine-containing mobile phases. *Marine Ecology Progress Series* **195:** 29-45

Zhang H, Han W, De Smet I, Talboys P, Loya R, Hassan A, Rong H, Jürgens G, Paul Knox J,

Wang MH (2010) ABA promotes quiescence of the quiescent centre and suppresses stem cell

differentiation in the Arabidopsis primary root meristem. *The Plant Journal* **64:** 764-774

Zhao Y (2012) Auxin biosynthesis: a simple two-step pathway converts tryptophan to indole-3-acetic

acid in plants. *Molecular Plant* **5:** 334-338

Zhao Y (2018) Essential roles of local auxin biosynthesis in plant development and in adaptation to

environmental changes. *Annual Review of Plant Biology* **69:** 417-435

Zhao Y, Christensen SK, Fankhauser C, Cashman JR, Cohen JD, Weigel D, Chory J (2001) A role

for flavin monooxygenase-like enzymes in auxin biosynthesis. *Science* **291:** 306-309

Zheng X, Yang Y, Al-Babili S (2021) Exploring the diversity and regulation of apocarotenoid

metabolic pathways in plants. *Frontiers in Plant Science* **12:** 787049

Figure legends

Figure 1. Fluridone inhibits phytoene desaturase activity to suppress PR growth. A) The carotenoid biosynthesis pathway. PSY, phytoene synthase; PDS, phytoene desaturase; CCR1/SDG8, carotenoid chloroplast regulatory1/set domain group 8; CCR2/CRTISO, carotenoid chloroplast regulatory2/crtiso; β-LCY, β-cyclase; CCD1/4/7/8, carotenoid cleavage dioxygenase 1/4/7/8. **B)** Five-day-old plants were transferred to medium with or without 12.8 μM fluridone. Photographs were captured at 3 d after transfer (dat). Scale bar = 5 mm. **C, D)** Five-day-old plants of WT and *mPDS* were transferred to medium with or without fluridone. **E–G)** Five-day-old plants of WT and *pds3-1* were transferred to medium with or without 800 nM fluridone. Photographs were captured at 3 dat. Scale bar = 5 mm. Red lines indicate the root tips at the time point of transfer. The PR growth rate in **(C, F)** was analyzed from 2 to 3 dat. The LR density in **(D, G)** was counted at 3 dat. Data represent the means ± standard error (SE) from 10-12 seedlings. *Significant differences compared to WT in **(C, D)** (*: $P < 0.05$, **: $P < 0.01$; Student's *t*-test). Different lowercase letters above the bars in **(F, G)** indicate significant differences at $P < 0.05$ (one-way Analysis of Variance [ANOVA] following Tukey-Kramer test).

Figure 2. Fluridone induces endogenous IAA levels. Five-day-old plants were transferred to medium with or without 800 nM fluridone at time 0. The IAA contents in the roots were measured at the indicated time points. Data represent the means ± SE from 3 biological replicates. *Significant differences compared to mock-treated plants (*: $P < 0.05$; Student's *t*-test).

Figure 3. The expression of *DR5:LUC* at the root tips was increased following fluridone treatment. Four-day-old plants grown on 100 μM D-luciferin medium were transferred to D-luciferin-containing medium for a 1 d preincubation. Then plants were transferred to new D-luciferin-containing medium with or without 800 nM fluridone at time 0. The *DR5:LUC* expression in WT **(A)** and *mPDS* **(B)** was measured at an interval of 1 h. Data represent the means (relative to the mean of time –1 h) ± SE from 6 seedlings.

Figure 4. Auxin biosynthesis, signaling, and influx transport are required for fluridone-mediated root growth regulation. A–C) The resistance to fluridone in auxin-related mutants. Five-day-old plants were transferred to medium with or without 800 nM fluridone. **(A)** auxin signaling and influx transport mutants. **(B)** *wei2*, *wei7* and *wei2 wei7* mutants. **(C)** *yucQ* mutant. **D**, **E)** Five-day-old plants were transferred to medium with or without 800 nM fluridone and 2 μM auxinole **(D)** or 100 μM YDF **(E)**. The PR growth rate was analyzed from 2 to 3 dat. The relative PR growth rate in (A–C) was compared to mock-treated plants whose root growth rate is listed in Supplemental Table S1. Data represent the means ± SE from 20 seedlings in **(A)** and from 8 seedlings in **(B–E)**. Different lowercase letters above the bars in **(A, B, D, E)** indicate significant differences at $P < 0.05$ (one-way ANOVA following Tukey-Kramer test). *Significant differences compared to WT in **(C)** (**: $P < 0.01$; Student's *t*-test).

Figure 5. The carotenoid accumulation in the roots is negatively regulated by auxin levels and signaling. A, **B)** Five-day-old plants were transferred to medium with or without 800 nM fluridone and 10 μM auxinole **(A)** or 100 μM YDF **(B)**. The levels of β-carotene in the roots were measured at 2 dat. The bars indicate mock (white), fluridone (diagonal striped black), auxinole **(A)** or YDF **(B)** (horizontal striped black), and auxinole + fluridone **(A)** or YDF + fluridone **(B)** (black). **C)** The β-carotene levels of the auxin-deficient mutants in the roots were measured at 7 d after germination. Data represent the means ± SE from 3 biological replicates. Different lowercase letters above the bars in **(A–C)** indicate significant differences at $P < 0.05$ (one-way ANOVA following Tukey-Kramer test). n.d. indicates no detectable data.

Figure 6. Fluridone activates auxin biosynthesis, response, and carotenoid catabolism genes. Five-day-old plants of WT and *mPDS* were transferred to medium with or without 800 nM fluridone at time 0. **A–D)** Relative expression of *IAA19*, *YUC9*, *CCD7*, and *CCD8* after fluridone treatment in the WT (open circle) and *mPDS* (open square). Data represent the

means ± SE from 3 biological replicates.

Figure 7. Retinal restores fluridone- and auxin-induced suppression of PR growth. A)

Five-day-old plants were transferred to new medium with or without 800 nM fluridone and

retinal. **B)** Five-day-old plants of WT and *pds3-1* were transferred to medium with or without

20 μM retinal and 20 nM NAA. **C)** Five-day-old plants of WT and the auxin-deficient

mutants were transferred to medium with or without 20 μM retinal. **D)** Five-day-old plants

were transferred to medium with or without 20 μM retinal and KKI. The PR growth rate was

analyzed from 1 to 2 dat in **(A)** and 2 to 3 dat in **(B–D)**. The bars in **(B)** indicate mock (white),

20 nM NAA (diagonal stripe black), 20 μM retinal (horizontal striped black) and NAA +

retinal (black). The relative growth in **(C)** was compared to mock-treated plants. **E)** The

relative *pIAA19::Eluc-PEST* expression by cotreatment of NAA and retinal. Four-day-old

plants grown on 100 μM D-luciferin medium were transferred to new D-luciferin-containing

medium for a 1 d preincubation. Then plants were transferred to new medium containing 100

μM D-luciferin, 10 nM NAA and with or without 20 μM retinal at time 0. The *pIAA19*

expression was measured at an interval of 1 h. Data represent the means ± SE from 8-10

seedlings in **(A–D)** and the means (relative to the means of time –1 h) ± SE from 6 seedlings

in **(E)**. Different lowercase letters above the bars in **(A–C)** indicate significant differences at

$P < 0.05$ (one-way ANOVA following Tukey-Kramer test).

Figure 8. Putative model of carotenoid-mediated PR growth regulation. Under normal

conditions, PR growth is controlled by auxin homeostasis. The short-term fluridone treatment

inhibits PDS activity and blocks consequent retinal biosynthesis to increase endogenous auxin

levels. The increase of auxin levels may exceed its optimal levels for root growth. Retinal

may trigger the IAA catabolism pathway to control endogenous auxin levels. Under long-term

fluridone incubation, the carotenoid deficiency leads to an acceleration of root growth

suppression via unknown factors. The activated auxin response feedback for the carotenoid

catabolism gene *CCD* activity in the regulation of carotenoid content. Black dashed lines

indicate unknown or predicted mechanisms.

Figure 1. Fluridone inhibits phytoene desaturase activity to suppress PR growth. A) The carotenoid biosynthesis pathway. PSY, phytoene synthase; PDS, phytoene desaturase; CCR1/SDG8, carotenoid chloroplast regulatory1/set domain group 8; CCR2/CRTISO, carotenoid chloroplast regulatory2/crtiso; β-LCY, β-cyclase; CCD1/4/7/8, carotenoid cleavage dioxygenase 1/4/7/8. **B)** Five-day-old plants were transferred to medium with or without 12.8 µM fluridone. Photographs were captured at 3 d after transfer (dat). Scale bar = 5 mm. **C, D)** Five-day-old plants of WT and *mPDS* were transferred to medium with or without fluridone. **E–G)** Five-day-old plants of WT and *pds3-1* were transferred to medium with or without 800 nM fluridone. Photographs were captured at 3 dat. Scale bar = 5 mm. Red lines indicate the root tips at the time point of transfer. The PR growth rate in **(C, F)** was analyzed from 2 to 3 dat. The LR density in **(D, G)** was counted at 3 dat. Data represent the means $\pm$ standard error (SE) from 10-12 seedlings. *Significant differences compared to WT in **(C, D)** (*: $P < 0.05$, **: $P < 0.01$; Student's *t*-test). Different lowercase letters above the bars in **(F, G)** indicate significant differences at $P < 0.05$ (one-way Analysis of Variance [ANOVA] following Tukey-Kramer test).

Figure 2. Fluridone induces endogenous IAA levels. Five-day-old plants were transferred to medium with or without 800 nM fluridone at time 0. The IAA contents in the roots were measured at the indicated time points. Data represent the means ± SE from 3 biological replicates. *Significant differences compared to mock-treated plants (*: $P < 0.05$; Student's *t*-test).

A

B

Figure 3. The expression of *DR5:LUC* at the root tips was increased following fluridone treatment. Four-day-old plants grown on 100 µM D-luciferin medium were transferred to D-luciferin-containing medium for a 1 d preincubation. Then plants were transferred to new D-luciferin-containing medium with or without 800 nM fluridone at time 0. The *DR5:LUC* expression in WT **(A)** and *mPDS* **(B)** was measured at an interval of 1 h. Data represent the means (relative to the mean of time –1 h) $\pm$ SE from 6 seedlings.

Figure 4. Auxin biosynthesis, signaling, and influx transport are required for fluridone-mediated root growth regulation. A–**C)** The resistance to fluridone in auxin-related mutants. Five-day-old plants were transferred to medium with or without 800 nM fluridone. **(A)** auxin signaling and influx transport mutants. **(B)** *wei2*, *wei7* and *wei2 wei7* mutants. **(C)** *yucQ* mutant. **D**, **E)** Five-day-old plants were transferred to medium with or without 800 nM fluridone and 2 µM auxinole **(D)** or 100 µM YDF **(E)**. The PR growth rate was analyzed from 2 to 3 dat. The relative PR growth rate in (A–C) was compared to mock-treated plants whose root growth rate is listed in Supplemental Table S1. Data represent the means ± SE from 20 seedlings in **(A)** and from 8 seedlings in **(B–E)**. Different lowercase letters above the bars in **(A, B, D, E)** indicate significant differences at *P* < 0.05 (one-way ANOVA following Tukey-Kramer test). *Significant differences compared to WT in **(C)** (**: *P* < 0.01; Student's *t*-test).

A **B**

C

Figure 5. The carotenoid accumulation in the roots is negatively regulated by auxin levels and signaling. A, **B**) Five-day-old plants were transferred to medium with or without 800 nM fluridone and 10 µM auxinole **(A)** or 100 µM YDF **(B)**. The levels of β-carotene in the roots were measured at 2 dat. The bars indicate mock (white), fluridone (diagonal striped black), auxinole **(A)** or YDF **(B)** (horizontal striped black), and auxinole + fluridone **(A)** or YDF + fluridone **(B)** (black). **C)** The β-carotene levels of the auxin-deficient mutants in the roots were measured at 7 d after germination. Data represent the means $\pm$ SE from 3 biological replicates. Different lowercase letters above the bars in **(A–C)** indicate significant differences at $P < 0.05$ (one-way ANOVA following Tukey-Kramer test). n.d. indicates no detectable data.

Figure 6. Fluridone activates auxin biosynthesis, response, and carotenoid catabolism genes. Five-day-old plants of WT and *mPDS* were transferred to medium with or without 800 nM fluridone at time 0. **A**–**D)** Relative expression of *IAA19*, *YUC9*, *CCD7*, and *CCD8* after fluridone treatment in the WT (open circle) and *mPDS* (open square). Data represent the means ± SE from 3 biological replicates.

Figure 7. Retinal restores fluridone- and auxin-induced suppression of PR growth. A) Five-day-old plants were transferred to medium with or without 800 nM fluridone and retinal. **B)** Five-day-old plants of WT and *pds3-1* were transferred to medium with or without 20 µM retinal and 20 nM NAA. **C)** Five-day-old plants of WT and the auxin-deficient mutants were transferred to medium with or without 20 µM retinal. **D)** Five-day-old plants were transferred to medium with or without 20 µM retinal and KKI. The PR growth rate was analyzed from 1 to 2 dat in **(A)** and 2 to 3 dat in **(B–D)**. The bars in **(B)** indicate mock (white), 20 nM NAA (diagonal stripe black), 20 µM retinal (horizontal striped black) and NAA + retinal (black). The relative growth in **(C)** was compared to mock-treated plants. **E)** The relative *pIAA19::Eluc-PEST* expression by cotreatment of NAA and retinal. Four-day-old plants grown on 100 µM D-luciferin medium were transferred to new D-luciferin-containing medium for a 1 d preincubation. Then plants were transferred to new medium containing 100 µM D-luciferin, 10 nM NAA and with or without 20 µM retinal at time 0. The *pIAA19* expression was measured at an interval of 1 h. Data represent the means $\pm$ SE from 8-10 seedlings in **(A–D)** and the means (relative to the means of time –1 h) $\pm$ SE from 6 seedlings in **(E)**. Different lowercase letters above the bars in **(A–C)** indicate significant differences at $P < 0.05$ (one-way ANOVA following Tukey-Kramer test).

Figure 8. Putative model of carotenoid-mediated PR growth regulation. Under normal conditions, PR growth is controlled by auxin homeostasis. The short-term fluridone treatment inhibits PDS activity and blocks consequent retinal biosynthesis to increase endogenous auxin levels. The increase of auxin levels may exceed its optimal levels for root growth. Retinal may trigger the IAA catabolism pathway to control endogenous auxin levels. Under long-term fluridone incubation, the carotenoid deficiency leads to an acceleration of root growth suppression via unknown factors. The activated auxin response feedback for the carotenoid catabolism gene CCD activity in the regulation of carotenoid content. Black dashed lines indicate unknown or predicted mechanisms.

Parsed Citations

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen HM, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, Gadrinab C, Heller C, Jeske A, Koesema E, Meyers CC, Parker H, Prednis L, Ansari Y, Choy N, Deen H, Geralt M, Hazari N, Hom E, Karnes M, Mulholland C, Ndubaku R, Schmidt I, Guzman P, Aguilar-Henonin L, Schmid M, Weigel D, Carter DE, Marchand T, Risseeuw E, Brogden D, Zeko A, Crosby WL, Berry CC, Ecker JR (2003) Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science 301: 653-657
  Google Scholar: Author Only Title Only Author and Title

Arias R, Dayan F, Michel A, Howell J, Scheffler B (2006) Characterization of a higher plant herbicide-resistant phytoene desaturase and its use as a selectable marker. Plant Biotechnology Journal 4: 263-273
  Google Scholar: Author Only Title Only Author and Title

Auldridge M, McCarty D, Klee H (2006) Plant carotenoid cleavage oxygenases and their apocarotenoid products. Current Opinion in Plant Biology 9: 315-321
  Google Scholar: Author Only Title Only Author and Title

Bainbridge K, Sorefan K, Ward S, Leyser O (2005) Hormonally controlled expression of the Arabidopsis MAX4 shoot branching regulatory gene. Plant Journal 44: 569-580
  Google Scholar: Author Only Title Only Author and Title

Barlier I, Kowalczyk M, Marchant A, Ljung K, Bhalerao R, Bennett M, Sandberg G, Bellini C (2000) The SUR2 gene of Arabidopsis thaliana encodes the cytochrome P450 CYP83B1, a modulator of auxin homeostasis. Proceedings of the National Academy of Sciences, USA 97: 14819-14824
  Google Scholar: Author Only Title Only Author and Title

Bartels PG, Watson CW (1978) Inhibition of carotenoid synthesis by fluridone and norflurazon. Weed Science 26: 198-203
  Google Scholar: Author Only Title Only Author and Title

Booker J, Auldridge M, Wills S, McCarty D, Klee H, Leyser O (2004) MAX3/CCD7 is a carotenoid cleavage dioxygenase required for the synthesis of a novel plant signaling molecule. Current Biology 14: 1232-1238
  Google Scholar: Author Only Title Only Author and Title

Breitenbach J, Sandmann G (2005) ζ-Carotene cis isomers as products and substrates in the plant poly-cis carotenoid biosynthetic pathway to lycopene. Planta 220: 785-793
  Google Scholar: Author Only Title Only Author and Title

Chen Q, Dai X, De-Paoli H, Cheng Y, Takebayashi Y, Kasahara H, Kamiya Y, Zhao Y (2014) Auxin overproduction in shoots cannot rescue auxin deficiencies in Arabidopsis roots. Plant and Cell Physiology 55: 1072-1079
  Google Scholar: Author Only Title Only Author and Title

Cheng Y, Dai X, Zhao Y (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in Arabidopsis. Genes & Development 20: 1790-1799
  Google Scholar: Author Only Title Only Author and Title

De Smet I, Lau S, Voss U, Vanneste S, Benjamins R, Rademacher EH, Schlereth A, De Rybel B, Vassileva V, Grunewald W, Naudts M, Levesque MP, Ehrismann JS, Inzé D, Luschnig C, Benfey PN, Weijers D, Van Montagu MCE, Bennett MJ, Jürgens G, Beeckman T (2010) Bimodular auxin response controls organogenesis in Arabidopsis. Proceedings of the National Academy of Sciences, USA 107: 2705-2710
  Google Scholar: Author Only Title Only Author and Title

Delarue M, Prinsen E, Van Onckelen H, Caboche M, Bellini C (1998) Sur2 mutations of Arabidopsis thaliana define a new locus involved in the control of auxin homeostasis. The Plant Journal 14: 603-611
  Google Scholar: Author Only Title Only Author and Title

Dickinson AJ, Lehner K, Mi J, Jia K, Mijar M, Dinneny JR, Al-Babili S, Benfey P (2019) β-Cyclocitral is a conserved root growth regulator. Proceedings of the National Academy of Sciences, USA 116: 10563-10567
  Google Scholar: Author Only Title Only Author and Title

Dickinson AJ, Zhang J, Luciano M, Wachsman G, Sandoval E, Schnermann M, Dinneny JR, Benfey P (2021) A plant lipocalin promotes retinal-mediated oscillatory lateral root initiation. Science 373: 1532-1536
  Google Scholar: Author Only Title Only Author and Title

Doyle SM, Rigal A, Grones P, Karady M, Barange DK, Majda M, Pařízková B, Karampelias M, Zwiewka M, Pěnčík A, Almqvist F, Ljung K, Novák O, Robert S (2019) A role for the auxin precursor anthranilic acid in root gravitropism via regulation of PIN-FORMED protein polarity and relocalisation in Arabidopsis. New Phytologist 223: 1420-1432
  Google Scholar: Author Only Title Only Author and Title

Du Y, Scheres B (2018) Lateral root formation and the multiple roles of auxin. Journal of Experimental Botany 69: 155-167
  Google Scholar: Author Only Title Only Author and Title

Dubrovsky JG, Forde BG (2012) Quantitative analysis of lateral root development: pitfalls and how to avoid them. Plant Cell 24: 4-14

Google Scholar: Author Only Title Only Author and Title

Emenecker RJ, Strader LC (2020). Auxin-abscisic acid interactions in plant growth and development. Biomolecules 10: 281

Google Scholar: Author Only Title Only Author and Title

Fukaki H, Tameda S, Masuda H, Tasaka M (2002) Lateral root formation is blocked by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of Arabidopsis. The Plant Journal 29: 153-168

Google Scholar: Author Only Title Only Author and Title

Fukui K, Arai K, Tanaka Y, Aoi Y, Kukshal V, Jez JM, Kubes MF, Napier R, Zhao Y, Kasahara H (2022) Chemical inhibition of the auxin inactivation pathway uncovers the roles of metabolic turnover in auxin homeostasis. Proceedings of the National Academy of Sciences, USA 119: e2206869119

Google Scholar: Author Only Title Only Author and Title

González-Guzmán M, Apostolova N, Bellés JM, Barrero JM, Piqueras P, Ponce MR, Micol JL, Serrano R, Rodríguez PL (2002) The short-chain alcohol dehydrogenase ABA2 catalyzes the conversion of xanthoxin to abscisic aldehyde. Plant Cell 14: 1833-1846

Google Scholar: Author Only Title Only Author and Title

Goto C, Ikegami A, Goh T, Maruyama K, Kasahara H, Takebayashi Y, Kamiya Y, Toyokura K, Kondo Y, Ishizaki K, Mimura T, Fukaki H (2023) Genetic interaction between Arabidopsis SUR2/CYP83B1 and GNOM indicates the importance of stabilizing local auxin accumulation in lateral root initiation. Plant and Cell Physiology pcad084

Google Scholar: Author Only Title Only Author and Title

Hayashi K, Neve J, Hirose M, Kuboki A, Shimada Y, Kepinski S, Nozaki H (2012) Rational design of an auxin antagonist of the SCFTIR1 auxin receptor complex. ACS Chemical Biology 7: 590-598

Google Scholar: Author Only Title Only Author and Title

Hayward A, Stirnberg P, Beveridge C, Leyser O (2009) Interactions between auxin and strigolactone in shoot branching control. Plant Physiology 151: 400-412

Google Scholar: Author Only Title Only Author and Title

Hobbie L, Estelle M (1994) Genetic approaches to auxin action. Plant, Cell & Environment 17: 525-540

Google Scholar: Author Only Title Only Author and Title

Hooker TS, Thorpe TA (1998) Effects of fluridone and abscisic acid on lateral root initiation and root elongation of excised tomato roots cultured in vitro. Plant Cell, Tissue and Organ Culture 52: 199-203

Google Scholar: Author Only Title Only Author and Title

Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, Zimmermann P (2008) Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. Advances in Bioinformatics 2008: 420747

Google Scholar: Author Only Title Only Author and Title

Jia K, Dickinson A, Mi J, Cui G, Xiao T, Kharbatia N, Guo X, Sugiono E, Aranda M, Blilou I, Rueping M, Benfey P, Al-Babili S (2019) Anchorene is a carotenoid-derived regulatory metabolite required for anchor root formation in Arabidopsis. Science Advances 5: eaaw6787

Google Scholar: Author Only Title Only Author and Title

Kim JS, Yun BW, Choi JS, Kim TJ, Kwak SS, Cho KY (2004) Death mechanisms caused by carotenoid biosynthesis inhibitors in green and in undeveloped plant tissues. Pesticide Biochemistry and Physiology 78: 127-139

Google Scholar: Author Only Title Only Author and Title

Kobayashi K, Baba S, Obayashi T, Sato M, Toyooka K, Keranen M, Aro E, Fukaki H, Ohta H, Sugimoto K, Masuda T (2012) Regulation of root greening by light and auxin/cytokinin signaling in Arabidopsis. The Plant Cell 24: 1081-1095

Google Scholar: Author Only Title Only Author and Title

Koornneef M, Jorna ML, Brinkhorst-van der Swan DLC, Karssen CM (1982) The isolation of abscisic acid (ABA) deficient mutants by selection of induced revertants in non-germinating gibberellin sensitive lines of Arabidopsis thaliana (L.) Heynh. Theoretical and Applied Genetics 61: 385-393

Google Scholar: Author Only Title Only Author and Title

Koornneef M, Reuling G, Karssen CM (1984) The isolation and characterization of abscisic acid-insensitive mutants of Arabidopsis thaliana. Physiologia Plantarum 61: 377-383

Google Scholar: Author Only Title Only Author and Title

Koschmieder J, Wust F, Schaub P, Alvarez D, Trautmann D, Krischke M, Rustenholz C, Mano J, Mueller M, Bartels D, Hugueney P, Beyer P, Welsch R (2021) Plant apocarotenoid metabolism utilizes defense mechanisms against reactive carbonyl species and xenobiotics. Plant Physiology 185: 331-351

Google Scholar: Author Only Title Only Author and Title

Leyser O (2002) Molecular genetics of auxin signaling. Annual Review of Plant Biology 53: 377-398
Google Scholar: Author Only Title Only Author and Title

Lincoln C, Britton JH, Estelle M (1990) Growth and development of the axr1 mutants of Arabidopsis. The Plant Cell 2: 1071-1080
Google Scholar: Author Only Title Only Author and Title

López-Ráez JA, Charnikhova T, Gómez-Roldán V, Matusova R, Kohlen W, De Vos R, Verstappen F, Puech-Pages V, Bécard G, Mulder P, Bouwmeester H (2008) Tomato strigolactones are derived from carotenoids and their biosynthesis is promoted by phosphate starvation. New Phytologist 178: 863-874
Google Scholar: Author Only Title Only Author and Title

McElver J, Tzafrir I, Aux G, Rogers R, Ashby C, Smith K, Thomas C, Schetter A, Zhou Q, Cushman MA, Tossberg J, Nickle T, Levin JZ, Law M, Meinke D, Patton D (2001) Insertional mutagenesis of genes required for seed development in Arabidopsis thaliana. Genetics 159: 1751-1763
Google Scholar: Author Only Title Only Author and Title

Meinke DW (2019) Genome-wide identification of EMBRYO-DEFECTIVE (EMB) genes required for growth and development in Arabidopsis. New Phytologist 226: 306-325
Google Scholar: Author Only Title Only Author and Title

Moreno-Risueno M, Van Norman J, Moreno A, Zhang J, Ahnert S, Benfey P (2010) Oscillating gene expression determines competence for periodic Arabidopsis root branching. Science 329: 1306-1311
Google Scholar: Author Only Title Only Author and Title

Mustilli A-C, Merlot S, Vavasseur A, Fenzi F, Giraudat J (2002) Arabidopsis OST1 protein kinase mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive oxygen species production. Plant Cell 14: 3089-3099
Google Scholar: Author Only Title Only Author and Title

Nambara E, Marion-Poll A (2005) Abscisic acid biosynthesis and catabolism. Annual Review of Plant Biology 56: 165-185
Google Scholar: Author Only Title Only Author and Title

Nambara E, Suzuki M, Abrams S, McCarty DR, Kamiya Y, McCourt P (2002) A screen for genes that function in abscisic acid signaling in Arabidopsis thaliana. Genetics 161: 1247-1255
Google Scholar: Author Only Title Only Author and Title

Niyogi KK, Grossman AR, Björkman O (1998) Arabidopsis mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion. Plant Cell 10: 1121-1134
Google Scholar: Author Only Title Only Author and Title

Okumura K, Goh T, Toyokura K, Kasahara H, Takebayashi Y, Mimura T, Kamiya Y, Fukaki H (2013) GNOM/FEWER ROOTS is required for the establishment of an auxin response maximum for Arabidopsis lateral root initiation. Plant and Cell Physiology 54: 406-417
Google Scholar: Author Only Title Only Author and Title

Okushima Y, Overvoorde P, Arima K, Alonso J, Chan A, Chang C, Ecker J, Hughes B, Lui A, Nguyen D, Onodera C, Quach H, Smith A, Yu G, Theologis A (2005) Functional genomic analysis of the AUXIN RESPONSE FACTOR gene family members in Arabidopsis thaliana: unique and overlapping functions of ARF7 and ARF19. The Plant Cell 17: 444-463
Google Scholar: Author Only Title Only Author and Title

Olson JA, Hayaishi O (1965) The enzymatic cleavage of β-carotene into vitamin A by soluble enzymes of rat liver and intestine. Proceedings of the National Academy of Sciences, USA 54: 1364-1370
Google Scholar: Author Only Title Only Author and Title

Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ, Desveaux D, Rodriguez PL, Mccourt P, Zhu JK, Schroeder JI, Volkman BF, Cutler SR (2009) Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. Science 324: 1068-1071
Google Scholar: Author Only Title Only Author and Title

Qin G, Gu H, Ma L, Peng Y, Deng XW, Chen Z, Qu LJ (2007) Disruption of phytoene desaturase gene results in albino and dwarf phenotypes in Arabidopsis by impairing chlorophyll, carotenoid, and gibberellin biosynthesis. Cell Research 17: 471-482
Google Scholar: Author Only Title Only Author and Title

Reed RC, Brady SR, Muday G (1998) Inhibition of auxin movement from the shoot into the root inhibits lateral root development in Arabidopsis. Plant Physiology 118: 1369-1378
Google Scholar: Author Only Title Only Author and Title

Rodrigo MJ, Alquézar B, Alós E, Medina V, Carmona L Bruno M, Al-Babili S, Zacarías L (2013) A novel carotenoid cleavage activity involved in the biosynthesis of Citrus fruit-specific apocarotenoid pigments. Journal of Experimental Botany 64: 4461-4478
Google Scholar: Author Only Title Only Author and Title

Rodríguez-Villalón A, Gas E, Rodríguez-Concepción M (2009) Phytoene synthase activity controls the biosynthesis of carotenoids and the supply of their metabolic precursors in dark-grown Arabidopsis seedlings. Plant Journal 60:424-435
Google Scholar: Author Only Title Only Author and Title

Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene signal transduction in Arabidopsis thaliana: five novel mutant loci integrated into a stress response pathway. Genetics 139: 1393-1409
Google Scholar: Author Only Title Only Author and Title

Ruiz-Sola MÁ, Rodríguez-Concepción M (2012) Carotenoid biosynthesis in Arabidopsis: a colorful pathway. Arabidopsis Book 10: e0158
Google Scholar: Author Only Title Only Author and Title

Ruyter-Spira C, Kohlen W, Charnikhova T, van Zeijl A, van Bezouwen L, de Ruijter N, Cardoso C, Lopez-Raez JA, Matusova R, Bours R, Verstappen F, Bouwmeester H (2011) Physiological effects of the synthetic strigolactone analog GR24 on root system architecture in Arabidopsis: another belowground role for strigolactones? Plant Physiology 155: 721-734
Google Scholar: Author Only Title Only Author and Title

Saini S, Sharma I, Kaur N, Pati PK (2013) Auxin: a master regulator in plant root development. Plant Cell Report 32: 741-757
Google Scholar: Author Only Title Only Author and Title

Schmittgen T, Livak K (2008) Analyzing real-time PCR data by the comparative CT method. Nature Protocols 3: 1101-1108
Google Scholar: Author Only Title Only Author and Title

Stepanova A, Hoyt J, Hamilton A, Alonso J (2005) A link between ethylene and auxin uncovered by the characterization of two root-specific ethylene-insensitive mutants in Arabidopsis. The Plant Cell 17: 2230-2242
Google Scholar: Author Only Title Only Author and Title

Stirnberg P, van de Sande K, Leyser HMO (2002) MAX1 and MAX2 control shoot lateral branching in Arabidopsis. Development 129: 1131-1141
Google Scholar: Author Only Title Only Author and Title

Sugawara S, Hishiyama S, Jikumaru Y, Hanada A, Nishimura T, Koshiba T, Zhao Y, Kamiya Y, Kasahara H (2009) Biochemical analyses of indole-3-acetaldoxime-dependent auxin biosynthesis in Arabidopsis. Proceedings of the National Academy of Sciences, USA 106: 5430-5435
Google Scholar: Author Only Title Only Author and Title

Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki M, Harper R, Liscum E, Yamamoto K (2004) MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. The Plant Cell 16: 379-393
Google Scholar: Author Only Title Only Author and Title

Tichopad A, Dilger M, Schwarz G, Pfaffl MW (2003) Standardized determination of real-time PCR efficiency from a single reaction set-up. Nucleic Acids Research 31: e122
Google Scholar: Author Only Title Only Author and Title

Timpte C, Lincoln C, Pickett FB, Turner J, Estelle M (1995) The AXR1 and AUX1 genes of Arabidopsis function in separate auxin-response pathways. The Plant Journal 8: 561-569
Google Scholar: Author Only Title Only Author and Title

Toyokura K, Goh T, Shinohara H, Shinoda A, Kondo Y, Okamoto Y, Uehara T, Fujimoto K, Okushima Y, Ikeyama Y, Nakajima K, Mimura T, Tasaka M, Matsubayashi Y, Fukaki H (2019) Lateral inhibition by a peptide hormone-receptor cascade during Arabidopsis lateral root founder cell formation. Developmental Cell 48: 64-75
Google Scholar: Author Only Title Only Author and Title

Tsugafune S, Mashiguchi K, Fukui K, Takebayashi Y, Nishimura T, Sakai T, Shimada Y, Kasahara H, Koshiba T, Hayashi K (2017) Yucasin DF, a potent and persistent inhibitor of auxin biosynthesis in plants. Scientific Reports 7: 13992
Google Scholar: Author Only Title Only Author and Title

Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya Y, Shirasu K, Yoneyama K, Kyozuka J, Yamaguchi S (2008) Inhibition of shoot branching by new terpenoid plant hormones. Nature 455: 195-200
Google Scholar: Author Only Title Only Author and Title

Van Norman JM, Zhang J, Cazzonelli CI, Pogson BJ, Harrison PJ, Bugg TD, Chan KX, Thompson AJ, Benfey PN (2014) Periodic root branching in Arabidopsis requires synthesis of an uncharacterized carotenoid derivative. Proceedings of the National Academy of Sciences, USA 111: E1300-E1309
Google Scholar: Author Only Title Only Author and Title

Vieten A, Sauer M, Brewer P, Friml J (2007) Molecular and cellular aspects of auxin-transport-mediated development. Trends in Plant Science 12: 160-168
Google Scholar: Author Only Title Only Author and Title

Watahiki M, Yamamoto K (1997) The massugu1 mutation of Arabidopsis identified with failure of auxin-induced growth curvature of hypocotyl confers auxin insensitivity to hypocotyl and leaf. Plant Physiology 115: 419-426

Google Scholar: Author Only Title Only Author and Title

Won C, Shen X, Mashiguchi K, Zheng Z, Dai X, Cheng Y, Kasahara H, Kamiya Y, Chory J, Zhao Y (2011) Conversion of tryptophan to indole-3-acetic acid by TRYPTOPHAN AMINOTRANSFERASES OF ARABIDOPSIS and YUCCAs in Arabidopsis. Proceedings of the National Academy of Sciences, USA 108: 18518-18523

Google Scholar: Author Only Title Only Author and Title

Xu D, Miao J, Yumoto E, Yokota T, Asahina M, Watahiki M (2017) YUCCA9-mediated auxin biosynthesis and polar auxin transport synergistically regulate regeneration of root systems following root cutting. Plant and Cell Physiology 58: 1710-1723

Google Scholar: Author Only Title Only Author and Title

Yamamoto KT, Watahiki MK, Matsuzaki J, Satoh S, Shimizu H (2017) Space–time analysis of gravitropism in etiolated Arabidopsis hypocotyls using bioluminescence imaging of the IAA19 promoter fusion with a destabilized luciferase reporter. Journal of Plant Research 130: 765-777

Google Scholar: Author Only Title Only Author and Title

Yamamoto M, Yamamoto KT (1998) Differential effects of 1-naphthaleneacetic acid, indole-3-acetic acid and 2,4-dichlorophenoxyacetic acid on the gravitropic response of roots in an auxin-resistant mutant of Arabidopsis, auxl. Plant and Cell Physiology 39: 660-664

Google Scholar: Author Only Title Only Author and Title

Yoshioka T, Endo T, Satoh S (1998) Restoration of seed germination at supraoptimal temperatures by fluridone, an inhibitor of abscisic acid biosynthesis. Plant and Cell Physiology 39: 307-312

Google Scholar: Author Only Title Only Author and Title

Yuan H, Zhang J, Nageswaran D, Li L (2015) Carotenoid metabolism and regulation in horticultural crops. Horticulture Research 2: 15036

Google Scholar: Author Only Title Only Author and Title

Zapata M, Rodriguez F, Garrido J (2000) Separation of chlorophylls and carotenoids from marine phytoplankton: a new HPLC method using a reversed phase C8 column and pyridine-containing mobile phases. Marine Ecology Progress Series 195: 29-45

Google Scholar: Author Only Title Only Author and Title

Zhang H, Han W, De Smet I, Talboys P, Loya R, Hassan A, Rong H, Jürgens G, Paul Knox J, Wang MH (2010) ABA promotes quiescence of the quiescent centre and suppresses stem cell differentiation in the Arabidopsis primary root meristem. The Plant Journal 64: 764-774

Google Scholar: Author Only Title Only Author and Title

Zhao Y (2012) Auxin biosynthesis: a simple two-step pathway converts tryptophan to indole-3-acetic acid in plants. Molecular Plant 5: 334-338

Google Scholar: Author Only Title Only Author and Title

Zhao Y (2018) Essential roles of local auxin biosynthesis in plant development and in adaptation to environmental changes. Annual Review of Plant Biology 69: 417-435

Google Scholar: Author Only Title Only Author and Title

Zhao Y, Christensen SK, Fankhauser C, Cashman JR, Cohen JD, Weigel D, Chory J (2001) A role for flavin monooxygenase-like enzymes in auxin biosynthesis. Science 291: 306-309

Google Scholar: Author Only Title Only Author and Title

Zheng X, Yang Y, Al-Babili S (2021) Exploring the diversity and regulation of apocarotenoid metabolic pathways in plants. Frontiers in Plant Science 12: 787049

Google Scholar: Author Only Title Only Author and Title