Unravelling species demographic responses to local and global past ecological disruptions using whole-genome sequencing in volcanic oceanic island organisms

Maëva Gabrielli[1,2], Thibault Leroy[3], Jordi Salmona[1], Benoit Nabholz[4], Borja Milá[5*], Christophe Thébaud[1*]

1 Laboratoire Évolution et Diversité Biologique (EDB), UMR 5174 (Université Paul Sabatier, CNRS, IRD), Toulouse, France.

2 Department of Life Sciences and Biotechnology, University of Ferrara, Ferrara, Italy

3 GenPhySE, INRAE, INP, ENVT, Université de Toulouse, Castanet-Tolosan, France.

4 Institut des Sciences de l'Evolution de Montpellier, UMR 5554 (Université de Montpellier, CNRS, IRD, EPHE), Montpellier, France.

5 National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid, Spain.

* Equal authorship contributions

Running title: Demographic responses to global and local factors

Keywords: demographic inferences, effective population size, global change, population genomics, Mascarene archipelago, *Zosterops*

Abstract

Disentangling the effects of ecological disruptions operating at different spatial and temporal scales in shaping past species' demography is particularly important in the current context of rapid environmental changes driven by both local and global factors. We argue that volcanic oceanic islands provide useful settings to study the influence of past ecological disruptions operating at local and global scales on population demographic histories. We investigate

potential drivers of past population dynamics for three closely related species of passerine
birds from two volcanic oceanic islands, Reunion and Mauritius (Mascarene archipelago), with
distinct volcanic history. Using ABC and PSMC inferences from complete genomes, we
reconstructed the demographic history of the Reunion grey white-eye (*Zosterops borbonicus*),
the Reunion olive white-eye (*Z. olivaceus*), and the Mauritius grey white-eye (*Z. mauritianus*),
and searched for possible causes underlying similarities or differences between species living
on the same or different islands. Both demographic inferences strongly support ancient and
long-term expansions in all species. They also reveal different trajectories between species
inhabiting different islands, but consistent demographic trajectories in species or populations
from the same island. Species from Reunion appear to have experienced synchronous
reductions in population size during the Last Glacial Maximum, a trend not seen in the
Mauritian species. Overall, this study suggests that local events have played an important role
in shaping population trajectories of these island species. It also highlights the potential of our
conceptual framework to disentangle the effects of local and global drivers on past species'
demography and long-term population processes.

43   **Introduction**

Ecological disruptions due to variations in environmental conditions can have profound effects
on local population size by directly affecting individual physiology or indirectly modifying
population niche limits. These disruptions can be triggered by major environmental events,
such as global climatic changes, that can lead to drastic reductions in population size. Such
bottlenecks have important consequences for gene pools owing to their effect on effective
population sizes (N_e) (Maruyama & Fuerst, 1985; Nei et al., 1975; Tajima, 1989; Watterson,
1984). Such effects have been documented across different taxa in the case of past global
climatic events, implying some synchrony in population fluctuations at a very large spatial
scale in response to climatic changes (Chattopadhyay et al., 2019; Germain et al., 2023). In
turn, severe ecological disruptions at the local level can have more temporary and localized
effects on populations. These include explosive volcanic eruptions, forest fires, and direct or
indirect human impacts on the environment through predation or habitat modification
(Dornelas, 2010). Distinguishing the effects of local-scale drivers relative to large-scale
processes in shaping the demography of populations and species over time, especially when
both occurred in the past before human impact, might be essential in the context of human-
driven climate change which challenges the global persistence of many species and their
conservation (Nogués-Bravo et al., 2018; Urban et al., 2016). Yet, evaluating the relative

importance of local-based and global-based influences remains challenging since past local
effects are generally poorly understood at the scale of a species' range and can be
confounded by global effects.

Volcanic oceanic islands possess a number of features that could help to solve
problems associated with the difficulty of distinguishing the effects of past ecological
disruptions operating at local versus global scales on population demographic histories
(Warren et al., 2015). First, they have usually undergone a series of localized processes, such
as major geodynamic events (Whittaker et al., 2008) or human impact, but have also
experienced the effects of past changes in the global climate (Weigelt & Kreft, 2013). Second,
most volcanic oceanic islands are configured into groups of islands that are closely spaced,
and have similar origins, geophysical settings, and climates. Yet, each island within such
volcanic archipelagos has its own geological life cycle (Whittaker et al., 2008), and different
islands have experienced independent bouts of ecological disruptions, mainly owing to
volcanic episodes and related geodynamic processes that are often asynchronous across
islands. Therefore, studying volcanic archipelagos with islands at different stages of their life
cycle could allow for assessing the relative importance of local (*i.e.*, island-specific) and global
(*i.e.*, at the archipelago scale) disruption events on population size trajectories by comparing
long-term demographic responses of species living on the same or different islands within an
archipelago (Figure 1). In addition, the isolated nature of these islands is an asset for inferring
demographic histories of insular lineages as most species are island endemics and
interference from external source populations that can generate complex demographic
scenarios, as is often found in continental settings, is expected to be minimal (Losos &
Ricklefs, 2009). Furthermore, because different islands often host allopatric sister species,
archipelagos allow for the comparison of demographic histories from immediately related
species inhabiting islands with the same versus different geographical features, including age,
size, and maximum elevation, while minimizing possible higher-level confounding effects (J.
T. Weir & Lawson, 2015).

Some studies have inferred the demographic histories of species from volcanic
archipelagos (for example*,* (Lamichhaney et al., 2015; Martin et al., 2021), but did not attempt
to compare the population histories of species living on the same and different islands. Here,
we propose to do so using, as a study system, the Mascarene archipelago, a remote island
group in the southwestern Indian Ocean composed of two main islands, Mauritius and
Reunion. The two islands are separated by 140 km of deep water, were never connected to
each other (Thébaud et al., 2009). They are comparable in size while being at different stages
of island ontogeny (see Fernández-Palacios et al., 2011), and therefore exhibit very different

topographies: Reunion, a younger island (2 my) is in the erosion and landslide phase and has

a very rugged landscape, reaching an elevation of 3,071 m, while Mauritius, an older island (8

my) in the late basal plain phase, has experienced the long-term effects of erosion and

culminates at just 828 m (Duncan, 2009). Global climatic events, including Late Quaternary

events, appear to have affected species' demographic histories on both islands (de Boer et

al., 2015; Garot et al., 2019; Heller et al., 2008; Salmona et al., 2012; Salmona, Heller,

Quéméré, et al., 2017). However, the intensity of ecological disruption as experienced by

animal and plant populations may have been very different on the two islands since the

persistence of possible refuges during extreme cooling events is positively related to

mountainous topography (Harter et al., 2015; Mastretta-Yanes et al., 2015; but see Salces-

Castellano et al., 2021). Localized and temporary, island-specific events likely to have affected

many species' population trajectories have also occurred on both islands (A. Cheke & Hume,

2010). In particular a series of large explosive volcanic eruptions that took place around

200,000 ya (198,800 ya ± 2,500 years) on Reunion (Castellanos Melendez et al., 2023), and

a series of basaltic lava flows that have occurred regularly (on average every 21,000 years)

in the last 500,000 years on Mauritius, which currently cover nearly 75% of the island (Moore

et al., 2011).

To illustrate potential effects of local and global disruption events on population

demographic trajectories, we focused on three species of white-eyes (Zosteropidae), all

single-island endemics in the Mascarene Archipelago (Milá et al., 2010; Warren et al., 2006).

These small passerine birds show limited dispersal capacity (Bertrand et al., 2014; Chris

Smith, pers. comm.), a feature that could make them sensitive to major climatic and

environmental changes (Germain et al., 2023). We hypothesized that 1) if island-specific

events had major effects on species demography, we should recover island-specific

demographic signals shared by species living on a particular island, and similarly, 2) if regional

or global climatic events had severe impacts on species at the scale of the entire archipelago,

we should find similar signals of demographic events between species living on different

islands (Figure 1). Taking advantage of the recent public release of a high-quality reference

genome for the Reunion grey white-eye (Leroy, Anselmetti, et al., 2021), we used shotgun

sequencing to generate whole-genome sequence data from the Reunion grey white eye

(*Zosterops borbonicus*), the Reunion olive white-eye (*Zosterops olivaceus*) and the Mauritius

grey white-eye (*Zosterops mauritianus*). The first two species are endemic to Reunion

whereas the third species is endemic to Mauritius. All three species are immediate relatives

(Milá et al., 2010; Warren et al., 2006) and are common and widely distributed in their

respective islands. Therefore, it seems reasonable to assume that their long-term population

dynamics have been little affected by anthropogenic activities since humans colonized these

islands (A. Cheke & Hume, 2010). In addition, the Reunion grey white-eye is represented by
multiple geographic forms that originated and diversified within Reunion (Gabrielli et al., 2020;
Gill, 1973). Three forms are restricted to the lowlands, whereas a fourth form is only found in
the highlands, primarily between 1,400 and 3,000 m (J. Bertrand et al., 2016; Cornuault et al.,
2015). These forms can therefore be considered as replicates of lineages from the same island
but also as indicators of within-island environmental differences. To reconstruct the long-term
trajectory of effective population sizes (Emerson et al., 2001), we first used a series of
Markovian coalescent analyses at the individual species level. Although these approaches
have been intensively used in the past decade (Li & Durbin, 2011; Schiffels & Durbin, 2014;
Terhorst et al., 2017), a potentially serious limitation to using Markovian coalescent methods
is that they assume that the focal species behaves as a panmictic unit (Mazet et al., 2016).
Therefore, we also used demographic inferences based on Approximate Bayesian
Computation (ABC; Beaumont et al., 2002), since they allow the evaluation of various
scenarios of population size changes accounting for periods of gene exchange and estimate
the timing of demographic events. Our results strongly support ancient expansions in the three
species, likely associated with evolutionary processes that took place on the islands
themselves since they diverged from an ancestral stock. Furthermore, our results indicate that
the different forms of the Reunion grey white-eye exhibit similar demographic histories, distinct
from the demographic history of the Mauritius grey white-eye. These findings suggest that
local events play a crucial role in shaping population trajectories in Mascarene white-eyes.
Our conceptual framework and its working hypotheses appear to provide an improved basis
for disentangling the effects of local and global drivers on past demography of species and
better understanding population processes that are of paramount conservation importance in
the current context of rapid environmental change driven by both local and global factors of
ecological disruption.

Materials and Methods

a) Population sampling and DNA sequence data

Blood samples from a total of 42 individuals, captured in mistnets and then released, were
collected between 2007 and 2017 throughout the entire distribution of the Reunion grey white-
eye (22 individuals from 18 localities), the Reunion olive white-eye (11 individuals from 11
localities) and the Mauritius grey white-eye (nine individuals from nine localities) (Figure 2;

Table S1). Approximately 1 µg of high-quality DNA was extracted using a QIAGEN DNeasy
Blood & Tissue kit following the manufacturer's instructions, with an extra pre-digestion
grinding step. Genomic DNA extractions were sent to Novogene Bioinformatics Technology
(Beijing, China; 25 individuals) and GeT-PlaGe core facility (INRA, Toulouse; nine individuals)
for shotgun whole-genome resequencing with an Illumina HiSeq2500 sequencer. Our dataset
includes data from eight individuals sequenced using similar procedures in previous studies
(Bourgeois et al., 2017; Leroy, Anselmetti, et al., 2021).

b) Reference-based whole-genome processing

To genotype individuals, sequencing reads of all three species were first mapped against the
high-quality genome assembly (scaffold N50 exceeding one megabase) of a Reunion grey
white-eye (Leroy, Anselmetti, et al., 2021) using the bwa-mem algorithm (Li & Durbin, 2009)
with default parameters. Single nucleotide polymorphisms (SNPs) were then called for each
species independently using GATK v. 3.7 (McKenna et al., 2010) using base quality score
recalibration and indel realignment. Both variant and invariant sites were called (option "--
includeNonVariantSites"). Variant filtration was performed using a custom script to speed up
computations, but following the same procedures than under GATK, using the following
thresholds: QD>2.0, FS<60.0, MQ>40.0, MQRANKSUM>-2.0, READPOSRANKSUM>-2.0
and RAW_MQ > 45,000. The three species were also genotyped together for multi-population
demographic inferences, using the same variant filtration procedure. To check for the general
consistency of the obtained genomic dataset, we performed a Principal Components Analysis
with PLINK v. 1.90b5.3 (Purcell et al., 2007) and plotted the results in R version 3.5 (R core
team, 2020) with adegenet v. 2.1.1 (Jombart, 2008) (Figure S1).

c) Mutation rate and generation time estimates

Mutation rate and generation time are two key parameters in population genetic inference. In
this study, we used the same set of parameter values for the three focal species, considering
that they diverged from a common ancestor 1.2 million years ago at most (Martins et al., 2020;
Warren et al., 2006), they apparently have similar ecologies, and genomic features are very
conserved in birds (Kawakami et al., 2014; Singhal et al., 2015). We used a mutation rate of
$4.6e^{-9}$ mutations per site per generation (95% confidence interval: $3.4e^{-9}$ - $5.9e^{-9}$) based on a
direct estimate for the collared flycatcher (*Ficedula albicollis*) (Smeds et al., 2016). A few
estimates of generation times are available for *Zosterops* species, suggesting short generation

times, possibly less than a year (Cornetti et al., 2015; Moyle et al., 2009), but generation time

is between two and three years in the Heron Island population of silvereye (*Zosterops lateralis*

chlorocephalus) (Clegg et al., 2002; Estoup & Clegg, 2003). Thus, we used a generation time

of one year but also run our analyses using values ranging between 0.5 and 2 years to account

for uncertainty.

d) Demographic inferences

We used two independent approaches to reconstruct the demographic histories of the three

Mascarene white-eyes.

Markovian coalescent analyses: In order to estimate the historical population size trajectory in

each species, we first performed inferences of N_e using the Pairwise Sequentially Markovian

Coalescent (PSMC, Li and Durbin, 2011) and whole-genome sequence data. Briefly, the

PSMC model infers the local time to the most recent common ancestor (TMRCA) on the basis

of the local density of heterozygotes. We considered two different thresholds for the minimum

sequence length, either 1,000 bp or 100,000 bp. For each individual, a custom script was used

to convert genotypes into PSMC input format, *i.e.* converting each 100-base-pair window into

K if the window contains at least one heterozygote site and T otherwise, using the following

filters: QUAL > 100, 10<depth<150, N_ratio (ratio of missing data in the window) < 0.9. Sites

outside these thresholds were considered to be missing (N). We set the upper limit of the

TMRCA to 5 and the initial θ/ρ value to 1 and we used the atomic interval scheme

"4+30*2+4+6+10", as in Nadachowska-Brzyska et al. (2016).

Approximate Bayesian Computation inferences: We performed ABC inferences from whole-

genome sequence data using DILS (Demographic Inferences with Linked Selection; Fraïsse

et al., 2021). This pipeline gathers ABC demographic inferences of population size changes

and timing of gene flow during divergence using either single population models (Figure S2A)

or two population models Figure S2B). As DILS needs sequences as input, we performed a

whole-genome sequence reconstruction at the individual level (two sequences per

chromosome considering diploidy) for all Zosterops individuals as published in Leroy,

Rousselle, et al. (2021), and then generated DNA segments of 10 kb for all the scaffolds with

a scaffold size exceeding 10 kb and kept scaffolds between 1 and 10 kb as one DNA segment.

In addition, reconstructed sequences with an N content exceeding 50% (missing information)

were discarded. Eight summary statistics were used for single population models: number of

biallelic sites, pairwise nucleotide diversity (π) (Tajima, 1983), Watterson's *θ* (Watterson,

1975) and Tajima's D (Tajima, 1989), considering the average and standard deviation of each
measure. Thirty-seven summary statistics were used for two population models, including the
same estimates of genetic diversity for each population, and measures of differentiation and
divergence between populations (F_{ST}; B. S. Weir & Cockerham, 1984; Wright, 1943); net
divergence; Nei and Li, 1979) (see Table S2 for a list of all summary statistics). For all
inferences, model comparison was performed using a random forest algorithm. Two million
simulations were run under the best model to perform parameter estimations, and four rounds
of goodness-of-fit were run to refine parameter estimation (Fraïsse et al., 2021).

We first used single population models for each of the three species to compare three
scenarios involving either a constant population size (i) or a change of population size over
time, with either an expansion (ii) or a contraction (ii) (Figure S2A). Models including
homogeneous or heterogeneous N_e across the genome were compared. Three parameters
were estimated for single population models, namely population size prior to the population
size change event (N_{ANC}), population size after this event (N_{POP}), and time since this event in
number of generations (T_{dem}). The priors used for each parameter are detailed in Table S3.

Since single population models assume closed systems, we also used models of two
population divergence with different timing of gene flow, with two models assuming current
isolation, *i.e.* absence of ongoing gene flow, (SI: Strict Isolation; AM: Ancestral Migration), and
two models assuming ongoing gene flow (IM: Isolation with Migration; SC: Secondary Contact)
(Figure S2B). These four models assume the instantaneous split of an ancestral population
into two daughter populations that diverge either without any migration (SI), with continuous
introgression since the time of split (IM), with a period of migration in the early times of lineage
divergence only (AM), or with a period of migration occurring upon secondary contact after an
initial period of strict isolation (SC). The models also included a possible change in population
size in the daughter populations after the ancestral split. Eight parameters were estimated for
all two-population models, namely the population size of the ancestral population (N_{ANC}), the
time of the split between the two populations (T_{split}), the population sizes of the daughter
populations after the split (N_{ANC1} and N_{ANC2}), the time of the potential demographic change
(T_{dem1} and T_{dem2}), and the population sizes of the daughter populations after the demographic
change (N_{POP1} and N_{POP2}). The time of the secondary contact (T_{SC}) was estimated in the SC
model, and the time of ancestral migration cessation (T_{AM}) was estimated in the AM model.
Daughter population migration rates were estimated for the three models involving migration
(IM, AM, SC). The priors used for each parameter are detailed in Table S3. We run these
models for the three possible pairs of species (*Z. borbonicus- Z. mauritianus*; *Z. borbonicus-*
Z. olivaceus and *Z. mauritianus-Z. olivaceus*).

Results

Population size history reconstruction though PSMC analyses support contrasting population size trajectories between species living on different islands (Figure 3). While cycles of expansions and contractions are observed in the Reunion grey and olive white-eyes, Mauritius grey white-eyes present a clear signal of long-term expansion. Specifically, in the Reunion grey white-eye, a population decline was detected that could coincide with the Last Glacial Maximum (LGM; 21,000 ya), while, over the same period of time, only a slight reduction or a population size stasis was detected in the Reunion olive white-eye. In contrast, PSMC analyses indicate that the Mauritius grey white-eye underwent an expansion in the corresponding time window, suggesting that the LGM might have had a lower impact on this species than on Reunion species. Similar demographic histories were reconstructed for the different geographic forms of the Reunion grey white-eye (Figure S3), with a possibly stronger reduction in population size around the time of the LGM in the highland form. Even after considering substantial variation of the mutation rate and generation time (Figure S4), our main conclusions still hold. Similarly, restricting the analysis to the longest sequences with a minimal length of 100,000 bp instead of 1,000 bp led to similar results (Figure S5), suggesting that our results are robust to a certain extent to methodological limits and parameter uncertainties.

In general agreement with the PSMC analyses, ABC demographic reconstruction highlighted a strong signal of expansion for all species, in single population models as well as in two-population models. For single population inferences, models including an expansion phase had a very high posterior probability (≥ 0.95) compared to models including a bottleneck or constant population size (Table S4). More specifically, ancestral N_e values were estimated to be around 35,000, 65,000 and 130,000 individuals for the Reunion grey white-eye, Mauritius grey white-eye and Reunion olive white-eye, respectively, and current N_e values were estimated to be around 100,000, 490,000 and 700,000, respectively, representing 2.8, 7.5 and 5.4 fold increases (Table S5). Based on these models, the onsets of expansion were very different between Reunion and Mauritius grey white-eyes (around 20,000 generations ago and 45,000 generations ago for the Reunion and the Mauritius grey white-eyes, respectively) and the Reunion olive white-eye (around 250,000 generations ago).

Two-population models of divergence inferred an Ancestral Migration scenario as the most likely (posterior probability higher than 0.81 for current isolation models and higher than 0.67 for Ancestral Migration versus Strict Isolation; Figure S6), with a brief period of ancestral gene flow in all species comparisons (median values for the T_{AM}/T_{SPLIT} ratio: 0.72, 0.94 and

0.73 for *Z. borbonicus-Z. mauritianus*; *Z. borbonicus-Z. olivaceus* and *Z. mauritianus-Z.*
olivaceus, respectively). Evidence for current isolation between species was expected,
especially for species living in different islands and estimates of divergence times that were
obtained in our models are consistent with previous estimates (Cai et al., 2019; Warren et al.,
2006), suggesting that the temporal framework of our demographic inferences using ABC may
be quite robust. Parameter estimations obtained using these models suggested 10-fold
expansions from ancestral N_e of between 35,000 and 75,000 individuals to current N_e between
400,000 and 800,000 individuals across the three species (Table 1). These expansions
seemed to be rather ancient for the Reunion grey and olive white-eyes, with an expansion
estimated around 70-90% of the time of divergence, while the expansion inferred for the
Mauritius white-eye seemed more recent (30-50% of the time of divergence), highlighting
again the importance of island-specific events on population size trajectories.

Results from ABC demographic inferences for one of the four geographic forms of the
Reunion grey white-eye, the Grey-Headed Brown form (GHB), are consistent with those
considering all Reunion grey white-eye geographic forms together (Tables S4 and S5, Table
1), suggesting that the different geographic forms may have experienced similar population
trajectories.

Discussion

Reconstructing long-term population dynamics using whole-genome data has been key in
revealing how past global climatic events influenced changes in population size through time
in many different taxa (*e.g.* Chattopadhyay et al., 2019; Germain et al., 2023; Nadachowska-
Brzyska et al., 2015). Yet populations and species face ecological disruptions over time that
can vary from being temporary and localized to regional or even global with no necessary
synchrony between the two types of events. Therefore, estimates of the changes in effective
population sizes as visualized in PSMC profiles may reflect demographic responses to a
number of distinct events acting at different temporal and spatial scales and individual effects
may be difficult to distinguish from one another. In addition, the combined effects of these
different sources of variation on population demographic histories may also be accompanied
by variation in population structure, to which current approaches have been shown to be very
sensistive (Chikhi et al., 2018; Mazet et al., 2016; Wakeley, 1999). In an attempt to alleviate
this difficulty and distinguish the effects of local and global disruption events on population
demographic trajectories, our study used volcanic oceanic islands as a spatial and temporal
framework, whole-genome data obtained from closely related species, and ABC demographic

inference in addition to PSMC to uncover population size changes that could be attributable
to the different types of events. We obtained two key findings. First, we found that species
from the same island (Reunion) showed similarities in their population size trajectories that
differed from that reconstructed for the species from the other island (Mauritius); and second,
that this pattern also applied to within-island geographic forms of the Reunion grey white-eye
that exhibited similar demographic histories, distinct from the demographic history of the sister
taxon, the Mauritius grey white-eye.

These findings suggest that local events have played a major role in shaping
population trajectories in these island bird species. They also highlight the potential of our
conceptual framework and its working hypotheses in disentangling the effects of local and
global drivers on past demography of species. This is crucial to obtain a better understanding
of long-term population processes that are of paramount conservation importance in the
current context of rapid environmental change driven by both local and global factors of
ecological disruption (Leung et al., 2020; Nogués-Bravo et al., 2018; Urban et al., 2016).

What could be the factors explaining shared response to local (island-specific)
ecological disruptions in our study system? In the two Reunion species of white-eyes, the
period of lowest N_e corresponds approximately to the LGM, while in the Mauritius grey white-
eye this episode seems to have had no effect on the demographic history. Thus, global drivers
like LGM seem to have some effects but response to these effects may vary depending on
local conditions and the ecology of the species, although this last factor is probably
unimportant here given how closely related and ecologically similar white-eye species are in
both islands (A. S. Cheke, 1987). A similar signal of bottleneck associated to the LGM timing
has been observed in the endemic Reunion coffee tree (Garot et al., 2019) and was interpreted
as being the result of range contraction due to major drops in both precipitation (Yan et al.,
2016) and temperature (Barrows & Juggins, 2005), especially on the leeward side of the
island. Since the Reunion coffee tree is a widespread forest species, this may imply that forest
extent may have been greatly reduced during LGM, which might have impacted bird
populations, especially at lower elevations, according to a well-known scenario of distribution
shifts in response to Quaternary climatic fluctuations (Hewitt, 2000). Reduction of forest area
at the LGM and restriction in refugia have been reconstructed in many instances (*e.g.*
Carnaval and Moritz, 2008) with possible demographic consequences on the fauna the forest
remnants sheltered (*e.g.* Maldonado-Coelho, 2012). Elevational downward shifts of
ecosystems towards the coast can be expected during glacial events (Fernández-Palacios et
al., 2016), especially in high-elevation islands like Reunion (see Salces-Castellano et al.,
2021). The highland habitat might have become less suitable and the species' range may

have contracted towards the coast. In line with this, our PSMC results indicate that the
highland form of the Reunion grey white-eye may have experienced a stronger bottleneck
associated with the LGM than the other forms restricted to lowlands. By contrast, our Mauritius
grey white-eye PSMC analyses suggest that the population kept expanding during the LGM.
As Mauritius is a low elevation island, it is possible that during the LGM Mauritius offered more
suitable lowland habitats compared to Reunion. Furthermore, the globally low sea level
associated with the LGM translated into an increase in land that was more substantial for
Mauritius (50%) than for Reunion (10%) (Norder et al., 2018). The 4,200 BP megadrought that
has been documented in East Africa (Marchant & Hooghiemstra, 2004; Thompson, 2002),
North America (Booth et al., 2016), India (Staubwasser et al., 2003), Madagascar (Gasse &
Van Campo, 2001) and Mauritius (de Boer et al., 2015; Rijsdijk et al., 2011) and has been
associated with massive mortality and population bottlenecks of Western Indian ocean and
African species (Heller et al., 2008; Rijsdijk et al., 2011; Salmona et al., 2012; Salmona, Heller,
Lascoux, et al., 2017) left no detectable signature in the population trajectories that we
recovered. This was not unexpected since PSMC analysis lacks resolution at such recent
dates, as shown in Figure 3, and spurious signals of bottlenecks are often observed in the
lower time limit of PSMC efficiency (1,000 - 10,000 ya) limiting the use of that approach for
recent periods (Li & Durbin, 2011). Lastly, we did not detect bottlenecks associated with major
dated volcanic eruptions that occurred in either Reunion or Mauritius, as it has been observed
in other island bird systems (Bemmels et al., 2022). This could indicate that the volcanic
eruptions in the Mascarenes did not affect the totality of the islands in spite of the magnitude
of some events, in particular a series that occurred around 200 kya on Reunion and are
thought to have covered a large area of the island (Castellanos Melendez et al., 2023;
Fretzdorff et al., 2000) and/or that the white-eyes were able to find habitat refugia in which
they maintained relatively high population sizes in spite of the possible devastation of most of
the island. This view is supported by our results from both PSMC and ABC demographic
inferences that strongly support ancient and long-term expansions in the three study species,
likely associated with evolutionary processes which took place on the islands themselves
since the species diverged from an ancestral stock and reflecting the radiation of the
Mascarene white-eyes (Warren et al., 2006).

While our study recovered potentially important information about how the
demographic history of island species may have been influenced by ecological disruption
acting at different spatial and temporal scales, it also highlights a series of difficulties and
limitations inherent to this kind of investigations. First, population structure was not explicitly
taken into account in our study, yet population structure can severely bias demographic
inferences. For instance, it has been shown that a PSMC signal of population size changes

can be obtained in a population without any change of population in its history but with just changes of connectivity between subpopulations (Grusea et al., 2019; Mazet et al., 2016; Rodríguez et al., 2018). All inferences for one particular geographic form of the Reunion grey white-eye produced similar results that those obtained using all geographic forms together, making us confident that our conclusions are fairly robust to deviations due to population structure, at least for that species. Second, we only considered models with up to two populations since the framework that we used (DILS; Fraïsse et al., 2021), while fast and malleable, only works with two populations at most. We tried using the Fastsimcoal modelling approach (Excoffier et al., 2013) as an alternative but when compared to DILS, it yielded unrealistic results with very poor fits to the data (results not shown). In addition, Fastsimcoal does not allow for N_e and N_m heterogeneity along the genome, which can lead to inaccurate demographic inferences while DILS parametrizes this heterogeneity. Our ABC analyses indeed unconditionally support N_e heterogeneity along the genome (posterior probabilities of 0.99 for all the species, Table S4) indicating that accounting for this factor may be critical in demographic inference based on whole-genome sequence data. While strong signals of expansion were detected in our ABC demographic inferences, we cannot really rule out the possibility that bottlenecks occurred over the course of the species' population histories. More complex ABC models including several waves of demographic changes would be necessary to properly address this, but current ABC approaches may not be powerful enough to detect ancient and subtle population size changes. However, PSMC analyses, which account for the possibility of multiple population size changes across time, do not suggest many demographic changes, but are rather consistent with a general strong expansion.

To conclude, the shared population histories in the Reunion grey and olive white-eyes and their contrast with the Mauritius grey white-eye population history reveal that, in this system and at a relatively small spatial scale, local events may have been the most important drivers of population trajectories. Using more species from Reunion and Mauritius could further validate whether the differences in demographic trajectories that we recovered are general. As white-eyes could be more resilient to environmental changes than other species groups, the use of other species, with different ecologies or being less behaviourally plastic, could yield different population trajectories in response to both local and global events. More importantly, investigating the demographic history of multiple species from a range of taxa in several archipelagos by comparing species living on the same or different islands and controlling for their evolutionary relatedness, as recently advocated by Cerca et al. (2023) for another purpose, would greatly help understanding how species respond to local and global events of ecological disruption and would provide an useful tool for assessing the impacts and risks of plausible future human-driven changes in climate.

Acknowledgments

We thank Thomas Duval, Jennifer Devillechabrolle, Guillaume Gélinaud, Marie Manceau, Juli
Broggi, Josselin Cornuault, Yann Bourgeois, Joris Bertrand, Boris Delahaie, Dominique
Strasberg, Philipp Heeb, René-Claude Billot, Ben Warren and Jean-Michel Probst for their
assistance in the field in Reunion and Mauritius; Reunion National Park and Mauritius National
Parks and Conservation Service for granting us permission to conduct fieldwork in protected
areas of Reunion and Mauritius, respectively; Benoit Lequette (Reunion National Park) and
Vikash Tatayah (Mauritius Wildlife Foundation) for facilitating permit collection. We thank
Hélène Holota and Amaya Iribar-Pelozuelo for help with the laboratory work; Lounès Chikhi
for stimulating discussions on population structure and demographic inferences; Camille Roux
for his valuable advice on the use of DILS for demographic inferences; the IFB cluster and
Genotoul bioinformatics platform for the computational resources they provided and in
particular Marie-Stéphane Trotard for bioinformatics support.

Funding

This work was supported by the Agence Française pour le Développement, the National
Geographic Society, the Fondation pour la Recherche sur la Biodiversité, the 'Laboratoire
d'Excellence' TULIP (ANR-10-LABX-41), ANR grant (BirdIslandGenomic project, ANR-14-
CE02-0002) to B.N., a PhD studentship from the Ministère de l'Enseignement Supérieur et de
la Recherche and a grant from the Fonds Inkermann and François Lacoste to M.G. Logistic
support on Reunion was provided by the Marelongue field station run by the University of
Reunion and the Observatoire des Sciences de l'Univers de La Réunion.

Data Accessibility and Benefit-Sharing

Raw sequence reads of whole genomes are deposited in the SRA (BioProject PRJNA661201).

Author Contributions

B.M. and C.T. initiated and coordinated the project. M.G., B.N., B.M. and C.T. conceived the
study and designed the experiments. B.M. and C.T. conducted the fieldwork, with the
assistance of M.G., T.L. and B.N. M.G. performed the analyses. M.G., T.L., J.S., B.N., B.M.
and C.T. interpreted the analyses and wrote the manuscript.

References

Barrows, T. T., & Juggins, S. (2005). Sea-surface temperatures around the Australian margin and Indian Ocean during the Last Glacial Maximum. *Quaternary Science Reviews*, *24*(7), Article 7. https://doi.org/10.1016/j.quascirev.2004.07.020

Beaumont, M. A., Zhang, W., & Balding, D. J. (2002). Approximate Bayesian Computation in Population Genetics. *Genetics*, *162*(4), Article 4.

Bemmels, J. B., Haddrath, O., Colbourne, R. M., Robertson, H. A., & Weir, J. T. (2022). Legacy of supervolcanic eruptions on population genetic structure of brown kiwi. *Current Biology*, *32*(15), 3389-3397.e8. https://doi.org/10.1016/j.cub.2022.05.064

Bertrand, J. A. M., Bourgeois, Y. X. C., Delahaie, B., Duval, T., García-Jiménez, R., Cornuault, J., Heeb, P., Milá, B., Pujol, B., & Thébaud, C. (2014). Extremely reduced dispersal and gene flow in an island bird. *Heredity*, *112*(2), 190–196. https://doi.org/10.1038/hdy.2013.91

Bertrand, J., Delahaie, B., Bourgeois, Y. X. C., Duval, T., García-Jiménez, R., Cornuault, J., Pujol, B., Thébaud, C., & Milá, B. (2016). The role of selection and historical factors in driving population differentiation along an elevational gradient in an island bird. *Journal of Evolutionary Biology*, *29*(4), 824–836. https://doi.org/10.1111/jeb.12829

Booth, R. K., Jackson, S. T., Forman, S. L., Kutzbach, J. E., E. A. Bettis, I. I. I., Kreigs, J., & Wright, D. K. (2016). A severe centennial-scale drought in midcontinental North America 4200 years ago and apparent global linkages: *The Holocene*. https://doi.org/10.1191/0959683605hl825ft

Bourgeois, Y. X. C., Delahaie, B., Gautier, M., Lhuillier, E., Malé, P.-J. G., Bertrand, J. A. M., Cornuault, J., Wakamatsu, K., Bouchez, O., Mould, C., Bruxaux, J., Holota, H., Milá, B., & Thébaud, C. (2017). A novel locus on chromosome 1 underlies the evolution of a melanic plumage polymorphism in a wild songbird. *Royal Society Open Science*, *4*(2), 160805. https://doi.org/10.1098/rsos.160805

Cai, T., Cibois, A., Alström, P., Moyle, R. G., Kennedy, J. D., Shao, S., Zhang, R., Irestedt, M., Ericson, P. G. P., Gelang, M., Qu, Y., Lei, F., & Fjeldså, J. (2019). Near-complete phylogeny and taxonomic revision of the world's babblers (Aves: Passeriformes). *Molecular Phylogenetics and Evolution*, *130*, 346–356. https://doi.org/10.1016/j.ympev.2018.10.010

Carnaval, A. C., & Moritz, C. (2008). Historical climate modelling predicts patterns of current biodiversity in the Brazilian Atlantic forest. *Journal of Biogeography*, *35*(7), Article 7. https://doi.org/10.1111/j.1365-2699.2007.01870.x

Castellanos Melendez, M. P., Di Muro, A., Laurent, O., Kuiper, K., Wijbrans, J. R., & Bachmann, O. (2023). Explosive volcanism of Piton des Neiges (Reunion Island) and excess age dispersion in sanidine: Insights into magma chamber processes in a hotspot setting. *Chemical Geology*, *632*, 121539. https://doi.org/10.1016/j.chemgeo.2023.121539

Cerca, J., Cotoras, D. D., Bieker, V. C., De-Kayne, R., Vargas, P., Fernández-Mazuecos, M., López-Delgado, J., White, O., Stervander, M., Geneva, A. J., Andino, J. E. G., Meier, J. I., Roeble, L., Brée, B., Patiño, J., Guayasamin, J. M., Torres, M. de L., Valdebenito, H., Castañeda, M. del R., … Martin, M. D. (2023). Evolutionary genomics of oceanic island radiations. *Trends in Ecology & Evolution*, *38*(7), 631–642. https://doi.org/10.1016/j.tree.2023.02.003

Chattopadhyay, B., Garg, K. M., Ray, R., & Rheindt, F. E. (2019). Fluctuating fortunes: Genomes and habitat reconstructions reveal global climate-mediated changes in bats' genetic diversity. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1911), 20190304. https://doi.org/10.1098/rspb.2019.0304

Cheke, A., & Hume, J. P. (2010). *Lost Land of the Dodo: The Ecological History of Mauritius, Réunion and Rodrigues*. Bloomsbury Publishing.

Cheke, A. S. (1987). The ecology of the surviving native land-birds of Réunion. In A. W. Diamond (Ed.), *Studies of Mascarene Island Birds* (pp. 301–358). Cambridge University Press. https://doi.org/10.1017/CBO9780511735769

Chikhi, L., Rodríguez, W., Grusea, S., Santos, P., Boitard, S., & Mazet, O. (2018). The IICR (inverse instantaneous coalescence rate) as a summary of genomic diversity: Insights into demographic inference and model choice. *Heredity, 120*(1), 13–24. https://doi.org/10.1038/s41437-017-0005-6

Clegg, S. M., Degnan, S. M., Kikkawa, J., Moritz, C., Estoup, A., & Owens, I. P. F. (2002). Genetic consequences of sequential founder events by an island-colonizing bird. *Proceedings of the National Academy of Sciences, 99*(12), Article 12. https://doi.org/10.1073/pnas.102583399

Cornetti, L., Valente, L. M., Dunning, L. T., Quan, X., Black, R. A., Hebert, O., & Savolainen, V. (2015). The genome of the" great speciator" provides insights into bird diversification. *Genome Biology and Evolution, 7*(9), 2680–2691. https://doi.org/10.1093/gbe/evv168

Cornuault, J., Delahaie, B., Bertrand, J. A., Bourgeois, Y. X., Milá, B., Heeb, P., & Thébaud, C. (2015). Morphological and plumage colour variation in the Réunion grey white-eye (Aves: *Zosterops borbonicus*): assessing the role of selection. *Biological Journal of the Linnean Society, 114*(2), 459–473. https://doi.org/10.1111/bij.12428

de Boer, E. J., Vélez, M. I., Rijsdijk, K. F., de Louw, P. G., Vernimmen, T. J., Visser, P. M., Tjallingii, R., & Hooghiemstra, H. (2015). A deadly cocktail: How a drought around 4200 cal. yr BP caused mass mortality events at the infamous 'dodo swamp' in Mauritius. *The Holocene, 25*(5), 758–771. https://doi.org/10.1177/0959683614567886

Dornelas, M. (2010). Disturbance and change in biodiversity. *Philosophical Transactions of the Royal Society B: Biological Sciences, 365*(1558), 3719–3727. https://doi.org/10.1098/rstb.2010.0295

Duncan, R. A. (2009). Mascarene Islands, Geology. In *Encyclopedia of Islands* (pp. 620–622). University of California Press.

Emerson, B. C., Paradis, E., & Thébaud, C. (2001). Revealing the demographic histories of species using DNA sequences. *Trends in Ecology & Evolution, 16*(12), Article 12. https://doi.org/10.1016/S0169-5347(01)02305-9

Estoup, A., & Clegg, S. M. (2003). Bayesian inferences on the recent island colonization history by the bird Zosterops lateralis lateralis. *Molecular Ecology, 12*(3), 657–674. https://doi.org/10.1046/j.1365-294X.2003.01761.x

Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V. C., & Foll, M. (2013). Robust Demographic Inference from Genomic and SNP Data. *PLOS Genetics, 9*(10), e1003905. https://doi.org/10.1371/journal.pgen.1003905

Fernández-Palacios, J. M., Nascimento, L. de, Otto, R., Delgado, J. D., García-del-Rey, E., Arévalo, J. R., & Whittaker, R. J. (2011). A reconstruction of Palaeo-Macaronesia, with particular reference to the long-term biogeography of the Atlantic island laurel forests. *Journal of Biogeography, 38*(2), Article 2. https://doi.org/10.1111/j.1365-2699.2010.02427.x

Fernández-Palacios, J. M., Rijsdijk, K. F., Norder, S. J., Otto, R., Nascimento, L., Fernández-Lugo, S., Tjørve, E., Whittaker, R. J., & Santos, A. (2016). Towards a glacial-sensitive model of island biogeography. *Global Ecology and Biogeography, 25*(7), Article 7. https://doi.org/10.1111/geb.12320

Fraïsse, C., Popovic, I., Mazoyer, C., Spataro, B., Delmotte, S., Romiguier, J., Loire, É., Simon, A., Galtier, N., Duret, L., Bierne, N., Vekemans, X., & Roux, C. (2021). DILS: Demographic inferences with linked selection by using ABC. *Molecular Ecology Resources, 21*(8), 2629–2644. https://doi.org/10.1111/1755-0998.13323

Fretzdorff, S., Paterne, M., Stoffers, P., & Ivanova, E. (2000). Explosive activity of the Reunion Island volcanoes through the past 260,000 years as recorded in deep-sea sediments. *Bulletin of Volcanology, 62*(4), 266–277. https://doi.org/10.1007/s004450000095

Gabrielli, M., Nabholz, B., Leroy, T., Milá, B., & Thébaud, C. (2020). Within-island diversification in a passerine bird. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1923), Article 1923. https://doi.org/10.1098/rspb.2019.2999

Garot, E., Joët, T., Combes, M., Severac, D., & Lashermes, P. (2019). Plant population dynamics on oceanic islands during the Late Quaternary climate changes: Genetic evidence from a tree species (*Coffea mauritiana*) in Reunion Island. *New Phytologist*, *224*(2), 974–986. https://doi.org/10.1111/nph.16052

Gasse, F., & Van Campo, E. (2001). Late Quaternary environmental changes from a pollen and diatom record in the southern tropics (Lake Tritrivakely, Madagascar). *Palaeogeography, Palaeoclimatology, Palaeoecology*, *167*(3), Article 3. https://doi.org/10.1016/S0031-0182(00)00242-X

Germain, R. R., Feng, S., Chen, G., Graves, G. R., Tobias, J. A., Rahbek, C., Lei, F., Fjeldså, J., Hosner, P. A., Gilbert, M. T. P., Zhang, G., & Nogués-Bravo, D. (2023). Species-specific traits mediate avian demographic responses under past climate change. *Nature Ecology & Evolution*, *7*(6), Article 6. https://doi.org/10.1038/s41559-023-02055-3

Gill, F. (1973). Intra-island variation in the Mascarene white-eye *Zosterops Borbonica*. *Ornithological Monographs*.

Grusea, S., Rodríguez, W., Pinchon, D., Chikhi, L., Boitard, S., & Mazet, O. (2019). Coalescence times for three genes provide sufficient information to distinguish population structure from population size changes. *Journal of Mathematical Biology*, *78*(1), 189–224. https://doi.org/10.1007/s00285-018-1272-4

Harter, D. E. V., Irl, S. D. H., Seo, B., Steinbauer, M. J., Gillespie, R., Triantis, K. A., Fernández-Palacios, J.-M., & Beierkuhnlein, C. (2015). Impacts of global climate change on the floras of oceanic islands – Projections, implications and current knowledge. *Perspectives in Plant Ecology, Evolution and Systematics*, *17*(2), 160–183. https://doi.org/10.1016/j.ppees.2015.01.003

Heller, R., Lorenzen, E. D., Okello, J. B. A., Masembe, C., & Siegismund, H. R. (2008). Mid-Holocene decline in African buffalos inferred from Bayesian coalescent-based analyses of microsatellites and mitochondrial DNA. *Molecular Ecology*, *17*(22), Article 22. https://doi.org/10.1111/j.1365-294X.2008.03961.x

Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*(6789), Article 6789. https://doi.org/10.1038/35016000

Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers. *Bioinformatics (Oxford, England)*, *24*(11), 1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Kawakami, T., Smeds, L., Backström, N., Husby, A., Qvarnström, A., Mugal, C. F., Olason, P., & Ellegren, H. (2014). A high-density linkage map enables a second-generation collared flycatcher genome assembly and reveals the patterns of avian recombination rate variation and chromosomal evolution. *Molecular Ecology*, *23*(16), Article 16. https://doi.org/10.1111/mec.12810

Lamichhaney, S., Berglund, J., Almén, M. S., Maqbool, K., Grabherr, M., Martinez-Barrio, A., Promerová, M., Rubin, C.-J., Wang, C., Zamani, N., Grant, B. R., Grant, P. R., Webster, M. T., & Andersson, L. (2015). Evolution of Darwin's finches and their beaks revealed by genome sequencing. *Nature*, *518*(7539), 371–375. https://doi.org/10.1038/nature14181

Leroy, T., Anselmetti, Y., Tilak, M.-K., Bérard, S., Csukonyi, L., Gabrielli, M., Scornavacca, C., Milá, B., Thébaud, C., & Nabholz, B. (2021). A bird's white-eye view on avian sex chromosome evolution. *Peer Community Journal*, *1*. https://doi.org/10.24072/pcjournal.70

Leroy, T., Rousselle, M., Tilak, M.-K., Caizergues, A. E., Scornavacca, C., Recuerda, M., Fuchs, J., Illera, J. C., De Swardt, D. H., Blanco, G., Thébaud, C., Milá, B., & Nabholz, B. (2021). Island songbirds as windows into evolution in small populations. *Current Biology*, *31*(6), 1303-1310.e4. https://doi.org/10.1016/j.cub.2020.12.040

Leung, B., Hargreaves, A. L., Greenberg, D. A., McGill, B., Dornelas, M., & Freeman, R. (2020). Clustered versus catastrophic global vertebrate declines. *Nature*, *588*(7837), Article 7837. https://doi.org/10.1038/s41586-020-2920-6

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, *25*(14), 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li, H., & Durbin, R. (2011). Inference of human population history from individual whole-genome sequences. *Nature*, *475*(7357), 493–496. https://doi.org/10.1038/nature10231

Losos, J. B., & Ricklefs, R. E. (2009). Adaptation and diversification on islands. *Nature*, *457*(7231), Article 7231. https://doi.org/10.1038/nature07893

Maldonado-Coelho, M. (2012). Climatic oscillations shape the phylogeographical structure of Atlantic Forest fire-eye antbirds (Aves: Thamnophilidae). *Biological Journal of the Linnean Society*, *105*(4), Article 4. https://doi.org/10.1111/j.1095-8312.2011.01823.x

Marchant, R., & Hooghiemstra, H. (2004). Rapid environmental change in African and South American tropics around 4000 years before present: A review. *Earth-Science Reviews*, *66*(3), Article 3. https://doi.org/10.1016/j.earscirev.2004.01.003

Martin, C. A., Armstrong, C., Illera, J. C., Emerson, B. C., Richardson, D. S., & Spurgin, L. G. (2021). Genomic variation, population history and within-archipelago adaptation between island bird populations. *Royal Society Open Science*, *8*(2), 201146. https://doi.org/10.1098/rsos.201146

Martins, F. C., Cox, S. C., Irestedt, M., Prŷs-Jones nvestigation, R. P., & Day, J. J. (2020). A comprehensive molecular phylogeny of Afrotropical white-eyes (Aves: Zosteropidae) highlights prior underestimation of mainland diversity and complex colonisation history. *Molecular Phylogenetics and Evolution*, 106843. https://doi.org/10.1016/j.ympev.2020.106843

Maruyama, T., & Fuerst, P. A. (1985). Population bottlenecks and nonequilibrium models in population genetics. III. Genic homozygosity in populations which experience periodic bottlenecks. *Genetics*, *111*(3), 691–703. https://doi.org/10.1093/genetics/111.3.691

Mastretta-Yanes, A., Moreno-Letelier, A., Piñero, D., Jorgensen, T. H., & Emerson, B. C. (2015). Biodiversity in the Mexican highlands and the interaction of geology, geography and climate within the Trans-Mexican Volcanic Belt. *Journal of Biogeography*, *42*(9), 1586–1600. https://doi.org/10.1111/jbi.12546

Mazet, O., Rodríguez, W., Grusea, S., Boitard, S., & Chikhi, L. (2016). On the importance of being structured: Instantaneous coalescence rates and human evolution—lessons for ancestral population size inference? *Heredity*, *116*(4), 362–371. https://doi.org/10.1038/hdy.2015.104

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Milá, B., Warren, B. H., Heeb, P., & Thébaud, C. (2010). The geographic scale of diversification on islands: Genetic and morphological divergence at a very small spatial scale in the Mascarene grey white-eye (Aves: *Zosterops borbonicus*). *BMC Evolutionary Biology*, *10*, 158. https://doi.org/10.1186/1471-2148-10-158

Moore, J., White, W., Paul, D., Duncan, R., Abouchami, W., & Galer, S. (2011). Evolution of shield-building and rejuvenescent volcanism of Mauritius. *Journal of Volcanology and Geothermal Research*, *207*, 47–66. https://doi.org/10.1016/j.jvolgeores.2011.07.005

Moyle, R. G., Filardi, C. E., Smith, C. E., & Diamond, J. (2009). Explosive Pleistocene diversification and hemispheric expansion of a "great speciator". *Proceedings of the National Academy of Sciences*, *106*(6), 1863–1868. https://doi.org/10.1073/pnas.0809861105

Nadachowska-Brzyska, K., Burri, R., Smeds, L., & Ellegren, H. (2016). PSMC analysis of effective population sizes in molecular ecology and its application to black-and-white Ficedula flycatchers. *Molecular Ecology*, *25*(5), 1058–1072. https://doi.org/10.1111/mec.13540

Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., & Ellegren, H. (2015). Temporal Dynamics of Avian Populations during Pleistocene Revealed by Whole-Genome Sequences. *Current Biology*, *25*(10), 1375–1380. https://doi.org/10.1016/j.cub.2015.03.047

Nei, M., & Li, W.-H. (1979). Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences of the United States of America*, *76*(10), 5269–5273. https://doi.org/10.1073/pnas.76.10.5269

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The Bottleneck Effect and Genetic Variability in Populations. *Evolution*, *29*(1), 1–10. https://doi.org/10.2307/2407137

Nogués-Bravo, D., Rodríguez-Sánchez, F., Orsini, L., de Boer, E., Jansson, R., Morlon, H., Fordham, D. A., & Jackson, S. T. (2018). Cracking the code of biodiversity responses to past climate change. *Trends in Ecology & Evolution*, *33*(10), 765–776. https://doi.org/10.1016/j.tree.2018.07.005

Norder, S. J., Baumgartner, J. B., Borges, P. A. V., Hengl, T., Kissling, W. D., van Loon, E. E., & Rijsdijk, K. F. (2018). A global spatially explicit database of changes in island palaeo-area and archipelago configuration during the late Quaternary. *Global Ecology and Biogeography*, *27*(5), Article 5. https://doi.org/10.1111/geb.12715

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics*, *81*(3), 559–575. https://doi.org/10.1086/519795

R core team. (2020). R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria.*

Rijsdijk, K. F., Zinke, J., de Louw, P. G. B., Hume, J. P., van der Plicht, H. J., Hooghiemstra, H., Meijer, H. J. M., Vonhof, H. B., Porch, N., Florens, F. B. V., Baider, C., van Geel, B., Brinkkemper, J., Vernimmen, T., & Janoo, A. (2011). Mid-Holocene (4200 kyr BP) mass mortalities in Mauritius (Mascarenes): Insular vertebrates resilient to climatic extremes but vulnerable to human impact. *The Holocene*, *21*(8), 1179–1194. https://doi.org/10.1177/0959683611405236

Rodríguez, W., Mazet, O., Grusea, S., Arredondo, A., Corujo, J. M., Boitard, S., & Chikhi, L. (2018). The IICR and the non-stationary structured coalescent: Towards demographic inference with arbitrary changes in population structure. *Heredity*, *121*(6), 663–678. https://doi.org/10.1038/s41437-018-0148-0

Salces-Castellano, A., Stankowski, S., Arribas, P., Patiño, J., Karger, D. N., Butlin, R., & Emerson, B. C. (2021). Long-term cloud forest response to climate warming revealed by insect speciation history. *Evolution*, *75*(2), 231–244. https://doi.org/10.1111/evo.14111

Salmona, J., Heller, R., Lascoux, M., & Shafer, A. B. A. (2017). Inferring demographic history using genomic data. In *Population Genomics: Concepts, Approaches and Applications* (pp. 511–537). Springer.

Salmona, J., Heller, R., Quéméré, E., & Chikhi, L. (2017). Climate change and human colonization triggered habitat loss and fragmentation in Madagascar. *Molecular Ecology*, *26*(19), Article 19. https://doi.org/10.1111/mec.14173

Salmona, J., Salamolard, M., Fouillot, D., Ghestemme, T., Larose, J., Centon, J.-F., Sousa, V., Dawson, D. A., Thebaud, C., & Chikhi, L. (2012). Signature of a Pre-Human Population Decline in the Critically Endangered Reunion Island Endemic Forest Bird Coracina newtoni. *PLOS ONE, 7*(8), e43524. https://doi.org/10.1371/journal.pone.0043524

Schiffels, S., & Durbin, R. (2014). Inferring human population size and separation history from multiple genome sequences. *Nature Genetics*, *46*(8), 919–925. https://doi.org/10.1038/ng.3015

Singhal, S., Leffler, E. M., Sannareddy, K., Turner, I., Venn, O., Hooper, D. M., Strand, A. I., Li, Q., Raney, B., Balakrishnan, C. N., Griffith, S. C., McVean, G., & Przeworski, M. (2015). Stable recombination hotspots in birds. *Science*, *350*(6263), Article 6263. https://doi.org/10.1126/science.aad0843

Smeds, L., Qvarnström, A., & Ellegren, H. (2016). Direct estimate of the rate of germline mutation in a bird. *Genome Research*, *26*(9), 1211–1218. https://doi.org/10.1101/gr.204669.116

Staubwasser, M., Sirocko, F., Grootes, P. M., & Segl, M. (2003). Climate change at the 4.2 ka BP termination of the Indus valley civilization and Holocene south Asian monsoon variability. *Geophysical Research Letters*, *30*(8), Article 8. https://doi.org/10.1029/2002GL016822

Tajima, F. (1983). Evolutionary relationship of DNA sequences in finite populations. *Genetics*, *105*(2), 437–460. https://doi.org/10.1093/genetics/105.2.437

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, *123*(3), 585–595. https://doi.org/10.1093/genetics/123.3.585

Terhorst, J., Kamm, J. A., & Song, Y. S. (2017). Robust and scalable inference of population history from hundreds of unphased whole-genomes. *Nature Genetics*, *49*(2), 303–309. https://doi.org/10.1038/ng.3748

Thébaud, C., Warren, B. H., Cheke, A., & Strasberg, D. (2009). Mascarene Islands, Biology. In *Encyclopedia of Islands* (pp. 612–619). University of California Press.

Thompson, L. G. (2002). Kilimanjaro Ice Core Records: Evidence of Holocene Climate Change in Tropical Africa. *Science*, *298*(5593), Article 5593. https://doi.org/10.1126/science.1073198

Urban, M. C., Bocedi, G., Hendry, A. P., Mihoub, J.-B., Pe'er, G., Singer, A., Bridle, J. R., Crozier, L. G., De Meester, L., Godsoe, W., Gonzalez, A., Hellmann, J. J., Holt, R. D., Huth, A., Johst, K., Krug, C. B., Leadley, P. W., Palmer, S. C. F., Pantel, J. H., … Travis, J. M. J. (2016). Improving the forecast for biodiversity under climate change. *Science*, *353*(6304), aad8466. https://doi.org/10.1126/science.aad8466

Wakeley, J. (1999). Nonequilibrium migration in human history. *Genetics*, *153*(4), Article 4.

Warren, B. H., Bermingham, E., Prys-Jones, R. P., & Thébaud, C. (2006). Immigration, species radiation and extinction in a highly diverse songbird lineage: White-eyes on Indian Ocean islands. *Molecular Ecology*, *15*(12), 3769–3786. https://doi.org/10.1111/j.1365-294X.2006.03058.x

Warren, B. H., Simberloff, D., Ricklefs, R. E., Aguilée, R., Condamine, F. L., Gravel, D., Morlon, H., Mouquet, N., Rosindell, J., Casquet, J., Conti, E., Cornuault, J., Fernández-Palacios, J. M., Hengl, T., Norder, S. J., Rijsdijk, K. F., Sanmartín, I., Strasberg, D., Triantis, K. A., … Thébaud, C. (2015). Islands as model systems in ecology and evolution: Prospects fifty years after MacArthur-Wilson. *Ecology Letters*, *18*(2), 200–217. https://doi.org/10.1111/ele.12398

Watterson, G. A. (1975). On the number of segregating sites in genetical models without recombination. *Theoretical Population Biology*, *7*(2), 256–276. https://doi.org/10.1016/0040-5809(75)90020-9

Watterson, G. A. (1984). Allele frequencies after a bottleneck. *Theoretical Population Biology*, *26*(3), 387–407. https://doi.org/10.1016/0040-5809(84)90042-X

Weigelt, P., & Kreft, H. (2013). Quantifying island isolation – insights from global patterns of insular plant species richness. *Ecography*, *36*(4), 417–429. https://doi.org/10.1111/j.1600-0587.2012.07669.x

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-Statistics for the Analysis of Population Structure. *Evolution*, *38*(6), Article 6.

Weir, J. T., & Lawson, A. (2015). Evolutionary rates across gradients. *Methods in Ecology and Evolution*, *6*(11), 1278–1286. https://doi.org/10.1111/2041-210X.12419

822   Whittaker, R. J., Triantis, K. A., & Ladle, R. J. (2008). A general dynamic theory of oceanic
823   island biogeography. *Journal of Biogeography*, *35*(6), 977–994.
824   https://doi.org/10.1111/j.1365-2699.2008.01892.x
825   Wright, S. (1943). Isolation by Distance. *Genetics*, *28*(2), Article 2.
826   Yan, M., Wang, B., & Liu, J. (2016). Global monsoon change during the Last Glacial
827   Maximum: A multi-model study. *Climate Dynamics*, *47*(1), Article 1.
828   https://doi.org/10.1007/s00382-015-2841-5

830 <u>Table 1</u>: Parameter estimation for two population demographic scenarios in DILS. Numbers
831 outside brackets indicate median values while numbers inside brackets indicate confidence
832 intervals (2.5%-97.5%).

	ABC model parameters								
	N_{ANC}	T_{split}	N_{ANC1}	N_{ANC2}	T_{dem1}	N_{POP1}	T_{dem2}	N_{POP2}	T_{AM}
Z. borbonicus - Z. mauritianus	115,681 (103,039-130,232)	128,291 (95,638-168,209)	46,070 (24,675-76,275)	48,886 (30,636-77,610)	98,879 (80,019-121,191)	631,186 (487,328-742,422)	69,215 (55,562-86,687)	537,411 (396,932-661,929)	92,045 (75,532-110,466)
GHB – Z. mauritianus	84,075 (73,819-97,160)	115,323 (86,075-152,274)	66,184 (36,219-109,992)	47,120 (27,039-74,713)	91,004 (69,428-116,346)	443,404 (338,203-581,931)	60,632 (46,932-80,056)	405,894 (284,879-536,721)	93,817 (74,518-112,903)
Z. borbonicus - Z. olivaceus	234,347 (200,173-275,027)	318,314 (255,817-398,370)	67,133 (22,956-120,907)	55,254 (24,862-96,435)	234,291 (188,145-290,008)	797,268 (604,018-915,561)	293,743 (234,835-364,791)	816,811 (651,688-945,258)	299,302 (241,213-350,777)
Z. mauritianus - Z. olivaceus	237,298 (192,706-277,644)	431,242 (299,381-559,243)	76,860 (41,129-120,390)	34,639 (16,827-58,066)	118,964 (91,229-141,605)	745,166 (556,712-916,230)	291,374 (218,052-356,087)	407,800 (322,593-453,945)	316,104 (258,245-385,771)

<u>Figure 1</u>: Proposed strategy to compare the effects of local events and global events on
genetic diversity by comparing the demographic histories of closely related populations or
species inhabiting different islands from the same volcanic oceanic archipelago. Plots
represent effective population size (N_e) through time for a population or species, and numbers
indicate different events of ecological disruption associated with bottlenecks: event 1 is a
volcanic eruption on island A while event 2 corresponds to a global drop in temperatures, as
observed during a glacial maximum, that affects demographic histories on islands A and B
synchronously, all else being equal. Contemporary bottlenecks suggest the importance of
events at a large geographic scale while bottlenecks associated with island specific events
suggest the importance of local events as drivers of population histories.

<u>Figure 2</u>: Sampling localities for each of the three species used in this study: (A) Reunion
island grey white-eye, (B) Reunion island olive white-eye, and (C) Mauritius island grey white-
eye. Localities are labelled with numbers (Table S1). Grey colours represent elevation classes;
for Reunion: light grey = elevation below 1,000 m, medium grey = elevation between 1,000
and 2,000 m, dark grey = elevation above 2,000 m; for Mauritius island: light grey = elevation
below 400 m, medium grey = elevation between 400 and 600 m, dark grey = elevation above
600 m. Dotted lines represent distribution ranges of the four geographic forms of the Reunion
grey white-eye.

854
855

Figure 3: PSMC inferences of population size histories over time. Colours correspond to the
three white-eye species. Vertical dotted lines and grey vertical bars indicate (from left to right):
the 4,200 BP megadrought, the Last Glacial Maximum 21,000 ya, a catastrophic period of
large explosive eruptions in Reunion around 200,000 ya. In Mauritius, volcanic activity has
been quite intense throughout the last 500,000 years but was never explosive during that
period. Please note that the plots are in log-scales.

862