Demographic responses of oceanic island birds to local and regional ecological disruptions revealed by whole-genome sequencing

Maëva Gabrielli[1,2], Thibault Leroy[3], Jordi Salmona[1], Benoit Nabholz[4], Borja Milá[5*], Christophe Thébaud[1*]

1 Laboratoire Évolution et Diversité Biologique (EDB), UMR 5174 (Université Paul Sabatier, CNRS, IRD), Toulouse, France

2 Department of Life Sciences and Biotechnology, University of Ferrara, Ferrara, Italy

3 GenPhySE, INRAE, INP, ENVT, Université de Toulouse, Castanet-Tolosan, France

4 Institut des Sciences de l'Evolution de Montpellier, UMR 5554 (Université de Montpellier, CNRS, IRD, EPHE), Montpellier, France

5 National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid, Spain

* Equal authorship contributions

Running title: Demographic responses of oceanic island birds

Keywords: demographic inferences, effective population size, global change, population genomics, Mascarene archipelago, *Zosterops*

Abstract

Disentangling the effects of ecological disruptions operating at different spatial and temporal scales in shaping past species' demography is particularly important in the current context of rapid environmental changes driven by both local and regional factors. We argue that volcanic oceanic islands provide useful settings to study the influence of past ecological disruptions operating at local and regional scales on population demographic histories. We investigate

potential drivers of past population dynamics for three closely related species of passerine
birds from two volcanic oceanic islands, Reunion and Mauritius (Mascarene archipelago), with
distinct volcanic history. Using ABC and PSMC inferences from complete genomes, we
reconstructed the demographic history of the Reunion Grey White-eye (*Zosterops borbonicus*
(Pennant, 1781)), the Reunion Olive White-eye (*Z. olivaceus* (Linnaeus, 1766)), and the
Mauritius Grey White-eye (*Z. mauritianus* (Gmelin, 1789)), and searched for possible causes
underlying similarities or differences between species living on the same or different islands.
Both demographic inferences strongly support ancient and long-term expansions in all
species. They also reveal different trajectories between species inhabiting different islands,
but consistent demographic trajectories in species or populations from the same island.
Species from Reunion appear to have experienced synchronous reductions in population size
during the Last Glacial Maximum, a trend not seen in the Mauritian species. Overall, this study
suggests that local events may have played a role in shaping population trajectories of these
island species. It also highlights the potential of our conceptual framework to disentangle the
effects of local and regional drivers on past species' demography and long-term population
processes.

Introduction

Ecological disruptions due to variations in environmental conditions can have profound effects
on local population size by directly affecting individual physiology or indirectly modifying
population niche limits. These disruptions can be triggered by major environmental events,
such as global climatic changes, that can lead to drastic reductions in population size. Such
bottlenecks have important consequences for gene pools owing to their effect on effective
population sizes (N_e) (Maruyama and Fuerst 1985; Nei, Maruyama, and Chakraborty 1975;
Tajima 1989; Watterson 1984). Such effects have been documented across different taxa in
the case of past global climatic events, implying some synchrony in population fluctuations at
a very large spatial scale in response to climatic changes (Chattopadhyay et al. 2019; Germain
et al. 2023). In turn, severe ecological disruptions at the local level can have more temporary
and localized effects on populations. These include volcanic eruptions, forest fires, and direct
or indirect human impacts on the environment through predation or habitat modification, all of
which have the capacity to radically transform local landscapes (Dornelas 2010).
Distinguishing the effects of local-scale drivers relative to large-scale processes in shaping
the demography of populations and species over time, especially when both occurred in the
past before human impact, might be essential if we want to predict accurately how human-

driven climate change will challenge the global persistence of many species and their
conservation (Nogués-Bravo et al. 2018; Urban et al. 2016). Yet, evaluating the relative
importance of local-based and global-based influences remains challenging since past local
effects are generally poorly understood at the scale of a species' range and can be
confounded by global effects.

Volcanic oceanic islands possess a number of features that could help to solve
problems associated with the difficulty of distinguishing the effects of past ecological
disruptions operating at local versus regional scales on population demographic histories
(Warren et al. 2015). First, they have usually undergone a series of localized processes, such
as major geodynamic events (Whittaker, Triantis, and Ladle 2008) or human impact, but have
also experienced the effects of past changes in the global climate (Weigelt and Kreft 2013).
Second, most volcanic oceanic islands are configured into groups of islands that are closely
spaced and have similar origins, geophysical settings and climates. Yet, each island within
such volcanic archipelagos has its own geological life cycle (Whittaker et al. 2008), and
different islands have experienced independent bouts of ecological disruptions, mainly owing
to volcanic episodes and related geodynamic processes that are often asynchronous across
islands. Therefore, studying volcanic archipelagos with islands at different stages of their life
cycle could allow for assessing the relative importance of local (*i.e.*, island-specific) and
regional (*i.e.*, at the archipelago scale) disruption events on population size trajectories by
comparing long-term demographic responses of species living on the same or different islands
within an archipelago (Figure 1). In addition, the isolated nature of these islands is an asset
for inferring demographic histories of insular lineages as most species are island endemics
and interference from external source populations that can generate complex demographic
scenarios, as is often found in continental settings, is expected to be minimal (Losos and
Ricklefs 2009). Furthermore, because different islands often host allopatric sister species,
archipelagos allow for the comparison of demographic histories from immediately related
species, minimizing possible higher-level confounding effects (Weir and Lawson 2015).

Some studies have inferred the demographic histories of species from volcanic
archipelagos (*e.g.* (Lamichhaney et al. 2015; Martin et al. 2021), but did not attempt to
compare the population histories of species living on the same and different islands. Here, we
propose to do so using, as a study system, the Mascarene archipelago, a remote island group
in the southwestern Indian Ocean composed of two main islands, Mauritius and Reunion. The
two islands are separated by 140 km of deep water and were never connected to each other
(Thébaud et al. 2009). They are comparable in size while being at different stages of island
ontogeny (see Fernández-Palacios et al., 2011), and therefore exhibit very different

topographies: Reunion, a younger island (2 my) is in the erosion and landslide phase and has
a very rugged landscape, reaching an elevation of 3,071 m, while Mauritius, an older island (8
my) in the late basal plain phase, has experienced the long-term effects of erosion and
culminates at just 828 m (Duncan 2009). Global climatic events, including Late Quaternary
events, appear to have affected species' demographic histories on both islands (de Boer et al.
2015; Garot et al. 2019; Heller et al. 2008; Salmona et al. 2012; Salmona, Heller, Quéméré,
et al. 2017). However, the intensity of ecological disruption as experienced by animal and plant
populations may have been very different on the two islands since the persistence of possible
refugia during extreme cooling events is positively related to mountainous topography (Harter
et al., 2015; Mastretta-Yanes et al., 2015; but see Salces-Castellano et al., 2021). Localized
and temporary, island-specific events likely to have affected many species' population
trajectories have also occurred on both islands (Cheke and Hume 2010). In particular a series
of large explosive volcanic eruptions that took place around 200,000 ya (198,800 ya ± 2,500
years) on Reunion (Castellanos Melendez et al. 2023), and a series of basaltic lava flows that
have occurred regularly (on average every 21,000 years) in the last 500,000 years on
Mauritius, which currently cover nearly 75% of the island (Moore et al. 2011).

To illustrate potential effects of local and regional disruption events on population
demographic trajectories, we focused on three species of white-eyes (Zosteropidae), all
single-island endemics in the Mascarene Archipelago (Milá et al., 2010; Warren et al., 2006;
Figure 2). These small passerine birds show limited dispersal capacity (Bertrand et al., 2014;
Chris Smith, pers. comm.), a feature that could make them sensitive to major climatic and
environmental changes (Germain et al. 2023). We hypothesized that (1) if island-specific
events had major effects on species' demography, we should recover island-specific
demographic signals shared by species living on a particular island, and similarly, (2) if
regional climatic events had severe impacts on species at the scale of the entire archipelago,
we should find similar signals of demographic events between species living on different
islands (Figure 1). Taking advantage of the recent public release of a high-quality reference
genome for the Reunion Grey White-eye (*Zosterops borbonicus* (Pennant, 1781)) (Leroy,
Anselmetti, et al. 2021), we used shotgun sequencing to generate whole-genome sequence
data from the Reunion Grey White-eye, the Reunion Olive White-eye (*Zosterops olivaceus*
(Linnaeus, 1766)) and the Mauritius Grey White-eye (*Zosterops mauritianus* (Gmelin, 1789)).
The first two species are endemic to Reunion whereas the third species is endemic to
Mauritius. All three species are immediate relatives (Milá et al., 2010; Warren et al., 2006;
Figure 2) and are common and widely distributed in their respective islands. Therefore, it
seems reasonable to assume that their long-term population dynamics have been little
affected by anthropogenic activities since humans colonized these islands (Cheke and Hume

2010). In addition, the Reunion Grey White-eye is represented by multiple geographic forms

that originated and diversified within Reunion (Gabrielli et al. 2020; Gill 1973). Three forms are

restricted to the lowlands, whereas a fourth form is only found in the highlands, primarily

between 1,400 and 3,000 m (Bertrand et al. 2016; Cornuault et al. 2015). These forms can

therefore be considered as replicates of lineages from the same island but also as indicators

of within-island environmental differences. To reconstruct the long-term trajectory of effective

population sizes (Emerson, Paradis, and Thébaud 2001), we first used a series of Markovian

coalescent analyses at the individual level. Although these approaches have been intensively

used in the past decade (Li and Durbin 2011; Schiffels and Durbin 2014; Terhorst, Kamm, and

Song 2017), a potentially serious limitation to using Markovian coalescent methods is that they

assume that the focal species behaves as a panmictic unit (Mazet et al. 2016). Therefore, we

also used demographic inferences based on Approximate Bayesian Computation (ABC;

Beaumont et al., 2002), since they allow the evaluation of various scenarios of population size

changes accounting for periods of gene exchange and estimate the timing of demographic

events. Our results strongly support ancient expansions in the three species, likely associated

with evolutionary processes that took place on the islands themselves since they diverged

from an ancestral stock. Furthermore, our results indicate that the two species of Reunion

white-eyes similar demographic histories, distinct from the demographic history of the

Mauritius Grey White-eye. These findings suggest that local events have played a rather

significant role in shaping population trajectories in Mascarene white-eyes. Our conceptual

framework and its working hypotheses appear to provide an improved basis for disentangling

the effects of local and regional drivers on past demography of species and a better

understanding of population processes that are of paramount conservation importance in the

current context of rapid environmental change driven by both local and regional factors of

ecological disruption.

158   **Materials and Methods**

a) Population sampling and DNA sequence data

Blood samples from a total of 42 individuals, captured in mistnets and then released, were

collected between 2007 and 2017 throughout the entire distribution of the Reunion Grey

White-eye (22 individuals from 18 localities), the Reunion Olive White-eye (11 individuals from

11 localities), and the Mauritius Grey White-eye (nine individuals from nine localities) (Figure

2; Table S1). Approximately 1 µg of high-quality DNA was extracted using a QIAGEN DNeasy

Blood & Tissue kit following the manufacturer's instructions, with an extra pre-digestion

grinding step. Genomic DNA extractions were sent to Novogene Bioinformatics Technology

(Beijing, China; 25 individuals) and GeT-PlaGe core facility (INRA, Toulouse; nine individuals)

for shotgun whole-genome resequencing with an Illumina HiSeq2500 sequencer. Our dataset

includes data from eight individuals sequenced using similar procedures in previous studies

(Bourgeois et al. 2017; Leroy, Anselmetti, et al. 2021).

<u>b) Reference-based whole-genome processing</u>

To genotype individuals, sequencing reads of all three species were first mapped against the

high-quality genome assembly (scaffold N50 exceeding one megabase) of a Reunion Grey

White-eye (Leroy, Anselmetti, et al. 2021) using the bwa-mem algorithm (Li and Durbin 2009)

with default parameters. Single nucleotide polymorphisms (SNPs) were then called for each

species independently using GATK v. 3.7 (McKenna et al. 2010) using base quality score

recalibration and indel realignment. Both variant and invariant sites were called (option "--

includeNonVariantSites"). Variant filtration was performed using a custom script to speed up

computations, but following the same procedures than under GATK, using the following

thresholds: QD>2.0, FS<60.0, MQ>40.0, MQRANKSUM>-2.0, READPOSRANKSUM>-2.0

and RAW_MQ > 45,000. The three species were also genotyped together for multi-population

demographic inferences, using the same variant filtration procedure. After filtering, the mean

individual coverage was 15x (see Table S2 for details about the samples used). To check for

the general consistency of the obtained genomic dataset, we performed a Principal

Component Analysis with PLINK v. 1.90b5.3 (Purcell et al. 2007) and plotted the results in R

version 3.5 (R core team 2020) with adegenet v. 2.1.1 (Jombart 2008) (Figure S1).

<u>c) Mutation rate and generation time estimates</u>

Mutation rate and generation time are two key parameters in population genetic inference. In

this study, we used the same set of parameter values for the three focal species, considering

that they diverged from a common ancestor 1.2 million years ago at most (Martins et al. 2020;

Warren et al. 2006), and genomic features tend to be conserved in birds (Kawakami et al.

2014; Singhal et al. 2015). We used a mutation rate of $4.6e^{-9}$ mutations per site per generation

(95% confidence interval: $3.4e^{-9}$ - $5.9e^{-9}$) based on a direct estimate for the Collared Flycatcher

(*Ficedula albicollis* (Temminck, 1815)) (Smeds, Qvarnström, and Ellegren 2016). A few

estimates of generation times are available for *Zosterops* species, suggesting short generation

times, possibly less than a year (Cornetti et al. 2015; Moyle et al. 2009), but generation time

is between two and three years in the Heron Island population of Silvereye (*Zosterops lateralis*

chlorocephalus Gould, 1841*)* (Clegg et al. 2002; Estoup and Clegg 2003). Thus, we used a

generation time of one year but also run our analyses using values ranging between 0.5 and

2 years to account for uncertainty.

d) Demographic inferences

We used two independent approaches to reconstruct the demographic histories of the three

Mascarene white-eyes.

Markovian coalescent analyses: In order to estimate the historical population size trajectory in

each species, we first performed inferences of N_e using the Pairwise Sequentially Markovian

Coalescent (PSMC, Li and Durbin, 2011) since this method can be readily applied to unphased

whole-genome sequence data. Briefly, the PSMC model infers the local time to the most

recent common ancestor (TMRCA) on the basis of the local density of heterozygotes and

estimates the inverse instantaneous rate (IICR) of a sample which is equivalent to population

size in panmictic models (Mazet et al. 2016). We considered two different thresholds for the

minimum sequence length, either 1,000 bp or 100,000 bp. For each individual, a custom script

was used to convert genotypes into PSMC input format, *i.e.* converting each 100-base-pair

window into K if the window contains at least one heterozygote site and T otherwise, using the

following filters: QUAL > 100, 10<depth<150, N_ratio (ratio of missing data in the window) <

0.9. Sites outside these thresholds were considered to be missing (N). We set the upper limit

of the TMRCA to 5 and the initial θ/ρ value to 1 and we used the atomic interval scheme

"4+30*2+4+6+10", as in Nadachowska-Brzyska et al. (2016).

Approximate Bayesian Computation inferences: We performed ABC inferences from whole-

genome sequence data using DILS (Demographic Inferences with Linked Selection; Fraïsse

et al., 2021). This pipeline gathers ABC demographic inferences of population size changes

and timing of gene flow during divergence using either single-population models (Figure S2A)

or two-population models (Figure S2B). As DILS needs sequences as input, we performed a

whole-genome sequence reconstruction at the individual level (two sequences per

chromosome considering diploidy) for all individuals as published in Leroy, Rousselle, et al.

(2021), and then generated DNA segments of 10 kb for all the scaffolds with a scaffold size

exceeding 10 kb and kept scaffolds between 1 and 10 kb as one DNA segment. In addition,

reconstructed sequences with an N content exceeding 50% (missing information) were
discarded. Eight summary statistics were used for single-population models: number of
biallelic sites, pairwise nucleotide diversity (π) (Tajima 1983), Watterson's θ (Watterson 1975)
and Tajima's D (Tajima 1989), considering the average and standard deviation of each
measure. Thirty-seven summary statistics were used for two-population models, including the
same estimates of genetic diversity for each population, and measures of differentiation and
divergence between populations (F_{ST}; B. S. Weir & Cockerham, 1984; Wright, 1943); net
divergence; Nei and Li, 1979) (see Table S3 for a list of all summary statistics). For all
inferences, model comparison was performed using a random forest algorithm. Two million
simulations were run under the best model to perform parameter estimations, and four rounds
of goodness-of-fit were run to refine parameter estimation (Fraïsse et al. 2021).

We first used single-population models for each of the three species to compare three
scenarios involving either a constant population size (i) or a change of population size over
time, with either an expansion (ii) or a contraction (iii) (Figure S2A). Models including
homogeneous or heterogeneous N_e across the genome were compared. Three parameters
were estimated for single-population models, namely population size prior to the population
size change event (N_{ANC}), population size after this event (N_{POP}), and time since this event in
number of generations (T_{dem}). The priors used for each parameter are detailed in Table S4.

Since single-population models assume closed systems, we also used models of two-
population divergence with different timing of gene flow, with two models assuming current
isolation, *i.e.* absence of ongoing gene flow, (SI: Strict Isolation; AM: Ancestral Migration), and
two models assuming ongoing gene flow (IM: Isolation with Migration; SC: Secondary Contact)
(Figure S2B). These four models assume the instantaneous split of an ancestral population
into two daughter populations that diverge either without any migration (SI), with continuous
introgression since the time of split (IM), with a period of migration in the early times of lineage
divergence only (AM), or with a period of migration occurring upon secondary contact after an
initial period of strict isolation (SC). The models also included a possible change in population
size in the daughter populations after the ancestral split. Eight parameters were estimated for
all two-population models, namely the population size of the ancestral population (N_{ANC}), the
time of the split between the two populations (T_{split}), the population sizes of the daughter
populations after the split (N_{ANC1} and N_{ANC2}), the time of the potential demographic change
(T_{dem1} and T_{dem2}) and the population sizes of the daughter populations after the demographic
change (N_{POP1} and N_{POP2}). The time of the secondary contact (T_{SC}) was estimated in the SC
model, and the time of ancestral migration cessation (T_{AM}) was estimated in the AM model.
Daughter population migration rates were estimated for the three models involving migration

(IM, AM, SC). The priors used for each parameter are detailed in Table S4. We run these
models for the three possible pairs of species (Reunion Grey White-eye - Mauritius Grey
White-eye; Reunion Grey White-eye - Reunion Olive White-eye and Mauritius Grey White-eye
- Reunion Olive White-eye).

271 **Results**

Population size history reconstructions though PSMC analyses support contrasting population
size trajectories between species living on different islands (Figure 3). While cycles of
expansions and contractions are observed in the Reunion Grey and Olive White-eyes,
Mauritius Grey White-eyes present a clear signal of long-term expansion. Specifically, in the
Reunion Grey White-eye, a population decline was detected that could coincide with the Last
Glacial Maximum (LGM; 21,000 ya), while, over the same period of time, only a slight reduction
or population size stasis was detected in the Reunion Olive White-eye. In contrast, PSMC
analyses indicate that the Mauritius Grey White-eye underwent an expansion in the
corresponding time window, suggesting that the LGM might have had a lower impact on this
species than on the Reunion species. Similar demographic histories were reconstructed for
the different geographic forms of the Reunion Grey White-eye (Figure S3), with a possibly
stronger reduction in population size around the time of the LGM in the highland form. Even
after considering substantial variation in the mutation rate and generation time (Figure S4),
our main conclusions still hold. Similarly, restricting the analysis to the longest sequences with
a minimal length of 100,000 bp instead of 1,000 bp led to similar results (Figure S5),
suggesting that our results are robust to a certain extent to methodological limits and
parameter uncertainties.

In general agreement with the PSMC analyses, ABC demographic reconstruction
highlighted a strong signal of expansion for all species, in single-population models as well as
in two-population models. For single-population inferences, models including an expansion
phase had a very high posterior probability (≥ 0.95) compared to models including a bottleneck
or constant population size (Table S5). More specifically, ancestral N_e values were estimated
to be around 35,000, 65,000 and 130,000 individuals for the Reunion Grey White-eye,
Mauritius Grey White-eye and Reunion Olive White-eye, respectively, and current N_e values
were estimated to be around 100,000, 490,000 and 700,000, respectively, representing 2.8-,
7.5- and 5.4-fold increases (Table S6). Based on these models, the onsets of expansion were
very different between Reunion and Mauritius Grey White-eyes (around 20,000 generations

ago and 45,000 generations ago, respectively) and the Reunion Olive White-eye (around
250,000 generations ago).

Two-population models of divergence inferred an Ancestral Migration scenario as the
most likely (posterior probability higher than 0.81 for current isolation models and higher than
0.67 for Ancestral Migration versus Strict Isolation; Figure S6), with a brief period of ancestral
gene flow in all species comparisons (median values for the T_{AM}/T_{SPLIT} ratio: 0.72, 0.94 and
0.73 for Reunion Grey White-eye - Mauritius Grey White-eye; Reunion Grey White-eye -
Reunion Olive White-eye and Mauritius Grey White-eye - Reunion Olive White-eye,
respectively). Evidence for current isolation between species was expected, especially for
species living in different islands and estimates of divergence times that were obtained in our
models are consistent with previous estimates (Cai et al. 2019; Warren et al. 2006), suggesting
that the temporal framework of our demographic inferences using ABC may be quite robust.
Parameter estimations obtained using these models suggested 10-fold expansions from
ancestral N_e of between 35,000 and 75,000 individuals to current N_e between 400,000 and
800,000 individuals across the three species (Table 1). These expansions seemed to be rather
ancient for the Reunion Grey and Olive White-eyes, with an expansion estimated around 70-
90% of the time of divergence, while the expansion inferred for the Mauritius Grey White-eye
seemed more recent (30-50% of the time of divergence), highlighting again the importance of
island-specific events on population size trajectories.

Results from ABC demographic inferences for one of the four geographic forms of the
Reunion Grey White-eye, the Grey-Headed Brown form (GHB), are consistent with those
considering all Reunion Grey White-eye geographic forms together (Tables S5 and S6, Table
1), suggesting that the different geographic forms may have experienced similar population
trajectories.

Discussion

Reconstructing long-term population dynamics using whole-genome data has been key in
revealing how past regional climatic events influenced changes in population size through time
in many different taxa (*e.g.* Chattopadhyay et al., 2019; Germain et al., 2023; Nadachowska-
Brzyska et al., 2015). Yet, populations and species face ecological disruptions over time that
can vary from being temporary and localized to regional or even global with no necessary
synchrony between the two types of events. Therefore, estimates of the changes in effective
population sizes as visualized in PSMC profiles may reflect demographic responses to a

number of distinct events acting at different temporal and spatial scales and individual effects
that may be difficult to distinguish from one another. In addition, the combined effects of these
different sources of variation on population demographic histories may also be accompanied
by variation in population structure, to which current approaches have been shown to be very
sensistive (Chikhi et al. 2018; Mazet et al. 2016; Wakeley 1999). In an attempt to alleviate this
difficulty and distinguish the effects of local and regional disruption events on population
demographic trajectories, our study used volcanic oceanic islands as a spatial and temporal
framework, whole-genome data obtained from closely related species, and ABC demographic
inference in addition to PSMC to uncover population size changes that could be attributable
to the different types of events. We obtained two key findings. First, we found that species
from the same island (Reunion) showed similarities in their population size trajectories that
differed from the one reconstructed for the species from the other island (Mauritius); and
second, that this pattern also applied to within-island geographic forms of the Reunion Grey
White-eye that exhibited similar demographic histories, distinct from the demographic history
of the sister taxon, the Mauritius Grey White-eye.

These findings suggest that local events have played a significant role in shaping
population trajectories in these island bird species. They also highlight the potential of our
conceptual framework and its working hypotheses in disentangling the effects of local and
regional drivers on past demography of species. This is crucial to obtain a better
understanding of long-term population processes that are of paramount conservation
importance in the current context of rapid environmental change driven by both local and
global factors of ecological disruption (Leung et al. 2020; Nogués-Bravo et al. 2018; Urban et
al. 2016).

What could be the factors explaining shared response to local (island-specific)
ecological disruptions in our study system? In the two Reunion species of white-eyes, the
period of lowest N_e corresponds approximately to the LGM, while in the Mauritius Grey White-
eye, this episode seems to have had no effect on the demographic history. Thus, global drivers
like LGM seem to have some effects but the response to these effects may vary depending
on local conditions and the ecology of the species, although this last factor is probably
unimportant here given how closely related and ecologically similar white-eye species are in
both islands (Cheke 1987). Similar evidence of a bottleneck associated to the LGM timing has
been observed in the Reunion coffee tree (*Coffea mauritiana* Lam., 1783) (Garot et al. 2019)
and was interpreted as being the result of a range contraction due to major drops in both
precipitation (Yan, Wang, and Liu 2016) and temperature (Barrows and Juggins 2005),
especially on the leeward side of the island. Since the Reunion coffee tree is a widespread

forest species, this may imply that forest extent may have been greatly reduced during the
LGM, which might have impacted bird populations, especially at lower elevations, according
to a well-known scenario of distribution shifts in response to Quaternary climatic fluctuations
(Hewitt 2000). Reduction of forest area at the LGM and restriction in refugia have been
reconstructed in many instances (*e.g.* Carnaval and Moritz, 2008) with possible demographic
consequences on the fauna the forest remnants sheltered (*e.g.* Maldonado-Coelho, 2012).
Elevational downward shifts of ecosystems towards the coast can be expected during glacial
events (Fernández-Palacios et al. 2016), especially in high-elevation islands like Reunion (see
Salces-Castellano et al., 2021). The highland habitat might have become less suitable and
the species' range may have contracted towards the coast. In line with this, our PSMC results
indicate that the highland form of the Reunion Grey White-eye may have experienced a
stronger bottleneck associated with the LGM than the other forms restricted to the lowlands.
By contrast, our Mauritius Grey White-eye PSMC analyses suggest that the population kept
expanding during the LGM. As Mauritius is a low elevation island, it is possible that during the
LGM Mauritius offered more suitable lowland habitats compared to Reunion. Furthermore, the
globally low sea level associated with the LGM translated into an increase in land area that
was more substantial for Mauritius (50%) than for Reunion (10%) (Norder et al. 2018). The
4,200 BP megadrought that has been documented in East Africa (Marchant and Hooghiemstra
2004; Thompson 2002), North America (Booth et al. 2005), India (Staubwasser et al. 2003),
Madagascar (Gasse and Van Campo 2001) and Mauritius (de Boer et al. 2015; Rijsdijk et al.
2011) and has been associated with massive mortality and population bottlenecks of Western
Indian ocean and African species (Heller et al. 2008; Rijsdijk et al. 2011; Salmona et al. 2012;
Salmona, Heller, Lascoux, et al. 2017) must have impacted populations of our study species,
but it left no detectable signature in the population trajectories that we recovered. However,
PSMC analysis lacks resolution at such recent dates, as shown in Figure 3, and spurious
signals of bottlenecks are often observed in the lower time limit of PSMC efficiency (1,000 -
10,000 ya) limiting the use of that approach for recent periods (Li and Durbin 2011). While
some new methods have become available to reconstruct demographic histories close to
present (Terhorst et al. 2017), understanding the impact on population size of very recent
events still remains a challenge and requires larger datasets with multiple resequenced
individuals per species (Patton et al. 2019; Wilder et al. 2023). Lastly, we did not detect
bottlenecks associated with major dated volcanic eruptions that occurred in either Reunion or
Mauritius, as it has been observed in other island bird systems (Bemmels et al. 2022). This
could indicate that the volcanic eruptions in the Mascarenes did not affect the totality of the
islands in spite of the magnitude of some events, in particular a series that occurred around
200 kya on Reunion and are thought to have wreaked havoc on a large area of the island
(Castellanos Melendez et al. 2023; Fretzdorff et al. 2000), and/or that the white-eyes were

able to find habitat refugia in which they maintained relatively high population sizes in spite of

the possible devastation of most of the island. This view is supported by our results from both

PSMC and ABC demographic inferences that strongly support ancient and long-term

expansions in the three study species, likely associated with evolutionary processes which

took place on the islands themselves since the species diverged from an ancestral stock and

reflecting the radiation of the Mascarene white-eyes (Warren et al. 2006).

410   While our study recovered potentially important information about how the

demographic history of island species may have been influenced by ecological disruption

acting at different spatial and temporal scales, it also highlights a series of difficulties and

limitations inherent to this kind of investigations. First, population structure was not explicitly

taken into account in our study; yet, it can severely bias demographic inferences. For instance,

it has been shown that a PSMC signal consistent with population size changes can be

obtained in a population without any change of population in its history but with just changes

in connectivity between subpopulations (Grusea et al. 2019; Mazet et al. 2016; Rodríguez et

al. 2018). Our PSMC reconstructions were similar for all four geographic forms of the Reunion

Grey White-eye, and all ABC inferences for one particular geographic form (the GHB form)

produced similar results to those obtained using all geographic forms together, making us

confident that our conclusions are fairly robust to deviations from panmixia due to population

structure, at least for that species. Second, we assumed similar generation times for the three

species, while these have different social structure and population densities, that may

influence this parameter. For example, cooperative breeding has been reported in Reunion

and Mauritius Grey White-eyes (Gill 1973; Hansen, Olesen, and Jones 2002) that may delay

first reproduction and overall increase generation time (Kreider et al. 2022). However, we note

that the results from PSMC analyses did not change substantially after varying both mutation

rate and generation time, suggesting that our results are also robust, to some extent, to

variations in these parameters. Third, we only considered models with up to two populations

since the framework that we used (DILS; Fraïsse et al., 2021), while fast and malleable, only

works with two populations at most. We tried using the Fastsimcoal modelling approach

(Excoffier et al. 2013) as an alternative but when compared to DILS, it yielded unrealistic

results with very poor fits to the data (results not shown). In addition, Fastsimcoal does not

allow for N_e and N_m heterogeneity along the genome, which can lead to inaccurate

demographic inferences while DILS parametrizes this heterogeneity. Our ABC analyses

indeed unconditionally support N_e heterogeneity along the genome (posterior probabilities of

0.99 for all the species, Table S4) indicating that accounting for this factor may be critical in

demographic inference based on whole-genome sequence data. While strong signals of

expansion were detected in our ABC demographic inferences, we cannot really rule out the

possibility that bottlenecks occurred over the course of the species' population histories. More complex ABC models including several waves of demographic changes would be necessary to properly address this, but current ABC approaches may not be powerful enough to detect ancient and subtle population size changes. However, PSMC analyses, which account for the possibility of multiple population size changes across time, do not suggest many demographic changes, but are rather consistent with a general strong expansion.

To conclude, the shared population histories in the Reunion Grey and Olive White-eyes and their contrast with the Mauritius Grey White-eye population history suggest that, in this system and at a relatively small spatial scale, local events may have been important drivers of population trajectories. Using more species from Reunion and Mauritius will be necessary to further validate whether the differences in demographic trajectories that we recovered are general. As white-eyes could be more resilient to environmental changes than other species groups, the use of other species, with different ecologies or being less behaviourally plastic, could yield different population trajectories in response to both local and regional events. More importantly, investigating the demographic history of multiple species from a range of taxa in several archipelagos by comparing species living on the same or different islands and controlling for their evolutionary relatedness, as recently advocated by Cerca et al. (2023) for another purpose, would greatly help understanding how species respond to local and regional events of ecological disruption and would provide an useful tool for assessing the impacts and risks of plausible future human-driven changes in climate.

Acknowledgments:

We thank Thomas Duval, Jennifer Devillechabrolle, Guillaume Gélinaud, Marie Manceau, Juli Broggi, Josselin Cornuault, Yann Bourgeois, Joris Bertrand, Boris Delahaie, Dominique Strasberg, Philipp Heeb, René-Claude Billot, Ben Warren and Jean-Michel Probst for their assistance in the field in Reunion and Mauritius; Reunion National Park and Mauritius National Parks and Conservation Service for granting us permission to conduct fieldwork in protected areas of Reunion and Mauritius, respectively; Benoit Lequette (Reunion National Park) and Vikash Tatayah (Mauritius Wildlife Foundation) for facilitating permit collection. We thank Hélène Holota, Uxue Suescun, and Amaya Iribar-Pelozuelo for help with the laboratory work; Lounès Chikhi for stimulating discussions on population structure and demographic inferences; Camille Roux for his valuable advice on the use of DILS for demographic inferences; the IFB cluster and Genotoul bioinformatics platform for the computational resources they provided and in particular Marie-Stéphane Trotard for bioinformatics support. We are also grateful to Stanislav Harvančík, Simon Colenutt, and Hilary Jones for allowing us to reproduce their photos of white-eyes.

Funding:

This work was supported by the Agence Française pour le Développement, The National
Geographic Society, the Fondation pour la Recherche sur la Biodiversité, the 'Laboratoire
d'Excellence' TULIP (ANR-10-LABX-41), an ANR grant (BirdIslandGenomic project, ANR-14-
CE02-0002) to B.N., a PEPS grant from CNRS, a PhD studentship from the Ministère de
l'Enseignement Supérieur et de la Recherche and a grant from the Fonds Inkermann and
François Lacoste to M.G. Logistic support on Reunion was provided by the Marelongue field
station run by the University of Reunion and the Observatoire des Sciences de l'Univers de
La Réunion.

Data Accessibility and Benefit-Sharing

Raw sequence reads of whole genomes are deposited in the SRA (BioProject PRJNA661201
and PRJEB18566). See Table S2 for accession codes of all the samples used in this study.

Author Contributions

B.M. and C.T. initiated and coordinated the project. M.G., B.N., B.M. and C.T. conceived the
study and designed the experiments. B.M. and C.T. conducted the fieldwork, with the
assistance of M.G., T.L. and B.N. M.G. performed the analyses. M.G., T.L., J.S., B.N., B.M.
and C.T. interpreted the analyses and wrote the manuscript.

References

Barrows, Timothy T., and Steve Juggins. 2005. 'Sea-Surface Temperatures around the Australian Margin and Indian Ocean during the Last Glacial Maximum'. *Quaternary Science Reviews* 24(7):1017–47. doi: 10.1016/j.quascirev.2004.07.020.

Beaumont, Mark A., Wenyang Zhang, and David J. Balding. 2002. 'Approximate Bayesian Computation in Population Genetics'. *Genetics* 162(4):2025–35.

Bemmels, Jordan B., Oliver Haddrath, Rogan M. Colbourne, Hugh A. Robertson, and Jason T. Weir. 2022. 'Legacy of Supervolcanic Eruptions on Population Genetic Structure of Brown Kiwi'. *Current Biology* 32(15):3389-3397.e8. doi: 10.1016/j.cub.2022.05.064.

Bertrand, J., Y. X. C. Bourgeois, B. Delahaie, T. Duval, R. García-Jiménez, J. Cornuault, P. Heeb, B. Milá, B. Pujol, and C. Thébaud. 2014. 'Extremely Reduced Dispersal and Gene Flow in an Island Bird'. *Heredity* 112(2):190–96. doi: 10.1038/hdy.2013.91.

Bertrand, Joris, Boris Delahaie, Y. X. C. Bourgeois, T. Duval, R. García-Jiménez, J. Cornuault, B. Pujol, C. Thébaud, and B. Milá. 2016. 'The Role of Selection and Historical Factors in Driving Population Differentiation along an Elevational Gradient in an Island Bird'. *Journal of Evolutionary Biology* 29(4):824–36. doi: 10.1111/jeb.12829.

de Boer, Erik J., Maria I. Vélez, Kenneth F. Rijsdijk, Perry GB de Louw, Tamara JJ Vernimmen, Petra M. Visser, Rik Tjallingii, and Henry Hooghiemstra. 2015. 'A Deadly Cocktail: How a Drought around 4200 Cal. Yr BP Caused Mass Mortality Events at the Infamous "Dodo Swamp" in Mauritius'. *The Holocene* 25(5):758–71. doi: 10.1177/0959683614567886.

Booth, Robert K., Stephen T. Jackson, Steven L. Forman, John E. Kutzbach, E. A. Bettis, Joseph Kreigs, and David K. Wright. 2005. 'A Severe Centennial-Scale Drought in Midcontinental North America 4200 Years Ago and Apparent Global Linkages'. *The Holocene* 15(3):321–28. doi: 10.1191/0959683605hl825ft.

Bourgeois, Yann X. C., Boris Delahaie, Mathieu Gautier, Emeline Lhuillier, Pierre-Jean G. Malé, Joris A. M. Bertrand, Josselin Cornuault, Kazumasa Wakamatsu, Olivier Bouchez, Claire Mould, Jade Bruxaux, Hélène Holota, Borja Milá, and Christophe Thébaud. 2017. 'A Novel Locus on Chromosome 1 Underlies the Evolution of a Melanic Plumage Polymorphism in a Wild Songbird'. *Royal Society Open Science* 4(2):160805. doi: 10.1098/rsos.160805.

Cai, Tianlong, Alice Cibois, Per Alström, Robert G. Moyle, Jonathan D. Kennedy, Shimiao Shao, Ruiying Zhang, Martin Irestedt, Per G. P. Ericson, Magnus Gelang, Yanhua Qu, Fumin Lei, and Jon Fjeldså. 2019. 'Near-Complete Phylogeny and Taxonomic Revision of the World's Babblers (Aves: Passeriformes)'. *Molecular Phylogenetics and Evolution* 130:346–56. doi: 10.1016/j.ympev.2018.10.010.

Carnaval, Ana Carolina, and Craig Moritz. 2008. 'Historical Climate Modelling Predicts Patterns of Current Biodiversity in the Brazilian Atlantic Forest'. *Journal of Biogeography* 35(7):1187–1201. doi: 10.1111/j.1365-2699.2007.01870.x.

Castellanos Melendez, Maria Paula, Andrea Di Muro, Oscar Laurent, Klaudia Kuiper, Jan R. Wijbrans, and Olivier Bachmann. 2023. 'Explosive Volcanism of Piton Des Neiges (Reunion Island) and Excess Age Dispersion in Sanidine: Insights into Magma Chamber Processes in a Hotspot Setting'. *Chemical Geology* 632:121539. doi: 10.1016/j.chemgeo.2023.121539.

Cerca, José, Darko D. Cotoras, Vanessa C. Bieker, Rishi De-Kayne, Pablo Vargas, Mario Fernández-Mazuecos, Julia López-Delgado, Oliver White, Martin Stervander, Anthony J. Geneva, Juan Ernesto Guevara Andino, Joana Isabel Meier, Lizzie Roeble, Baptiste Brée, Jairo Patiño, Juan M. Guayasamin, María de Lourdes Torres, Hugo Valdebenito, María del Rosario Castañeda, Jaime A. Chaves, Patricia Jaramillo Díaz, Luis Valente, Matthew L. Knope, Jonathan P. Price, Loren H. Rieseberg, Bruce G. Baldwin, Brent C. Emerson, Gonzalo Rivas-Torres, Rosemary Gillespie, and Michael D. Martin. 2023. 'Evolutionary Genomics of Oceanic Island Radiations'. *Trends in Ecology & Evolution* 38(7):631–42. doi: 10.1016/j.tree.2023.02.003.

Chattopadhyay, Balaji, Kritika M. Garg, Rajasri Ray, and Frank E. Rheindt. 2019. 'Fluctuating Fortunes: Genomes and Habitat Reconstructions Reveal Global Climate-Mediated Changes in Bats' Genetic Diversity'. *Proceedings of the Royal Society B: Biological Sciences* 286(1911):20190304. doi: 10.1098/rspb.2019.0304.

Cheke, Anthony, and Julian P. Hume. 2010. *Lost Land of the Dodo: The Ecological History of Mauritius, Réunion and Rodrigues*. Bloomsbury Publishing.

Cheke, Anthony S. 1987. 'The Ecology of the Surviving Native Land-Birds of Réunion'. Pp. 301–58 in *Studies of Mascarene Island Birds*, edited by A. W. Diamond. Cambridge: Cambridge University Press.

Chikhi, Lounès, Willy Rodríguez, Simona Grusea, Patrícia Santos, Simon Boitard, and Olivier Mazet. 2018. 'The IICR (Inverse Instantaneous Coalescence Rate) as a Summary of Genomic Diversity: Insights into Demographic Inference and Model Choice'. *Heredity* 120(1):13–24. doi: 10.1038/s41437-017-0005-6.

Clegg, S. M., S. M. Degnan, J. Kikkawa, C. Moritz, A. Estoup, and I. P. F. Owens. 2002. 'Genetic Consequences of Sequential Founder Events by an Island-Colonizing Bird'. *Proceedings of the National Academy of Sciences* 99(12):8127–32. doi: 10.1073/pnas.102583399.

Cornetti, L., L. M. Valente, L. T. Dunning, X. Quan, R. A. Black, O. Hebert, and V. Savolainen. 2015. 'The Genome of the" Great Speciator" Provides Insights into Bird Diversification'. *Genome Biology and Evolution* 7(9):2680–91. doi: 10.1093/gbe/evv168.

Cornuault, Josselin, Boris Delahaie, Joris AM Bertrand, Yann XC Bourgeois, Borja Milá, Philipp Heeb, and Christophe Thébaud. 2015. 'Morphological and Plumage Colour Variation in the Réunion Grey White-Eye (Aves: *Zosterops Borbonicus*): Assessing the Role of Selection'. *Biological Journal of the Linnean Society* 114(2):459–73. doi: 10.1111/bij.12428.

Dornelas, Maria. 2010. 'Disturbance and Change in Biodiversity'. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1558):3719–27. doi: 10.1098/rstb.2010.0295.

Duncan, R. A. 2009. 'Mascarene Islands, Geology'. Pp. 620–22 in *Enclyclopedia of Islands*.
University of California Press.

Emerson, Brent C., Emmanuel Paradis, and Christophe Thébaud. 2001. 'Revealing the
Demographic Histories of Species Using DNA Sequences'. *Trends in Ecology &*
Evolution 16(12):707–16. doi: 10.1016/S0169-5347(01)02305-9.

Estoup, A., and S. M. Clegg. 2003. 'Bayesian Inferences on the Recent Island Colonization
History by the Bird Zosterops Lateralis Lateralis'. *Molecular Ecology* 12(3):657–74.
doi: 10.1046/j.1365-294X.2003.01761.x.

Excoffier, Laurent, Isabelle Dupanloup, Emilia Huerta-Sánchez, Vitor C. Sousa, and
Matthieu Foll. 2013. 'Robust Demographic Inference from Genomic and SNP Data'.
PLOS Genetics 9(10):e1003905. doi: 10.1371/journal.pgen.1003905.

Fernández-Palacios, José María, Lea de Nascimento, Rüdiger Otto, Juan D. Delgado,
Eduardo García-del-Rey, José Ramón Arévalo, and Robert J. Whittaker. 2011. 'A
Reconstruction of Palaeo-Macaronesia, with Particular Reference to the Long-Term
Biogeography of the Atlantic Island Laurel Forests'. *Journal of Biogeography*
38(2):226–46. doi: 10.1111/j.1365-2699.2010.02427.x.

Fernández-Palacios, José María, Kenneth F. Rijsdijk, Sietze J. Norder, Rüdiger Otto, Lea
Nascimento, Silvia Fernández-Lugo, Even Tjørve, Robert J. Whittaker, and Ana
Santos. 2016. 'Towards a Glacial-sensitive Model of Island Biogeography'. *Global*
Ecology and Biogeography 25(7):817–30. doi: 10.1111/geb.12320.

Fraïsse, Christelle, Iva Popovic, Clément Mazoyer, Bruno Spataro, Stéphane Delmotte,
Jonathan Romiguier, Étienne Loire, Alexis Simon, Nicolas Galtier, Laurent Duret,
Nicolas Bierne, Xavier Vekemans, and Camille Roux. 2021. 'DILS: Demographic
Inferences with Linked Selection by Using ABC'. *Molecular Ecology Resources*
21(8):2629–44. doi: 10.1111/1755-0998.13323.

Fretzdorff, S., M. Paterne, P. Stoffers, and E. Ivanova. 2000. 'Explosive Activity of the
Reunion Island Volcanoes through the Past 260,000 Years as Recorded in Deep-
Sea Sediments'. *Bulletin of Volcanology* 62(4):266–77. doi: 10.1007/s004450000095.

Gabrielli, Maëva, Benoit Nabholz, Thibault Leroy, Borja Milá, and Christophe Thébaud.
2020. 'Within-Island Diversification in a Passerine Bird'. *Proceedings of the Royal*
Society B: Biological Sciences 287(1923):20192999. doi: 10.1098/rspb.2019.2999.

Garot, Edith, Thierry Joët, Marie-Christine Combes, Dany Severac, and Philippe Lashermes.
2019. 'Plant Population Dynamics on Oceanic Islands during the Late Quaternary
Climate Changes: Genetic Evidence from a Tree Species (*Coffea Mauritiana*) in
Reunion Island'. *New Phytologist* 224(2):974–86. doi: 10.1111/nph.16052.

Gasse, F., and E. Van Campo. 2001. 'Late Quaternary Environmental Changes from a
Pollen and Diatom Record in the Southern Tropics (Lake Tritrivakely, Madagascar)'.
Palaeogeography, Palaeoclimatology, Palaeoecology 167(3):287–308. doi:
10.1016/S0031-0182(00)00242-X.

Germain, Ryan R., Shaohong Feng, Guangji Chen, Gary R. Graves, Joseph A. Tobias,
Carsten Rahbek, Fumin Lei, Jon Fjeldså, Peter A. Hosner, M. Thomas P. Gilbert,
Guojie Zhang, and David Nogués-Bravo. 2023. 'Species-Specific Traits Mediate
Avian Demographic Responses under Past Climate Change'. *Nature Ecology &*
Evolution 7(6):862–72. doi: 10.1038/s41559-023-02055-3.

Gill, Frank. 1973. 'Intra-Island Variation in the Mascarene White-Eye *Zosterops Borbonica*'. *Ornithological Monographs*.

Grusea, Simona, Willy Rodríguez, Didier Pinchon, Lounès Chikhi, Simon Boitard, and Olivier Mazet. 2019. 'Coalescence Times for Three Genes Provide Sufficient Information to Distinguish Population Structure from Population Size Changes'. *Journal of Mathematical Biology* 78(1):189–224. doi: 10.1007/s00285-018-1272-4.

Hansen, Dennis M., Jens M. Olesen, and Carl G. Jones. 2002. 'Trees, Birds and Bees in Mauritius: Exploitative Competition between Introduced Honey Bees and Endemic Nectarivorous Birds?' *Journal of Biogeography* 29(5–6):721–34. doi: 10.1046/j.1365-2699.2002.00720.x.

Harter, David E. V., Severin D. H. Irl, Bumsuk Seo, Manuel J. Steinbauer, Rosemary Gillespie, Kostas A. Triantis, José-María Fernández-Palacios, and Carl Beierkuhnlein. 2015. 'Impacts of Global Climate Change on the Floras of Oceanic Islands – Projections, Implications and Current Knowledge'. *Perspectives in Plant Ecology, Evolution and Systematics* 17(2):160–83. doi: 10.1016/j.ppees.2015.01.003.

Heller, R., E. D. Lorenzen, J. B. A. Okello, C. Masembe, and H. R. Siegismund. 2008. 'Mid-Holocene Decline in African Buffalos Inferred from Bayesian Coalescent-Based Analyses of Microsatellites and Mitochondrial DNA'. *Molecular Ecology* 17(22):4845–58. doi: 10.1111/j.1365-294X.2008.03961.x.

Hewitt, Godfrey. 2000. 'The Genetic Legacy of the Quaternary Ice Ages'. *Nature* 405(6789):907–13. doi: 10.1038/35016000.

Jombart, Thibaut. 2008. 'Adegenet: A R Package for the Multivariate Analysis of Genetic Markers'. *Bioinformatics (Oxford, England)* 24(11):1403–5. doi: 10.1093/bioinformatics/btn129.

Kawakami, Takeshi, Linnéa Smeds, Niclas Backström, Arild Husby, Anna Qvarnström, Carina F. Mugal, Pall Olason, and Hans Ellegren. 2014. 'A High-Density Linkage Map Enables a Second-Generation Collared Flycatcher Genome Assembly and Reveals the Patterns of Avian Recombination Rate Variation and Chromosomal Evolution'. *Molecular Ecology* 23(16):4035–58. doi: 10.1111/mec.12810.

Kreider, Jan J., Boris H. Kramer, Jan Komdeur, and Ido Pen. 2022. 'The Evolution of Ageing in Cooperative Breeders'. *Evolution Letters* 6(6):450–59. doi: 10.1002/evl3.307.

Lamichhaney, Sangeet, Jonas Berglund, Markus Sällman Almén, Khurram Maqbool, Manfred Grabherr, Alvaro Martinez-Barrio, Marta Promerová, Carl-Johan Rubin, Chao Wang, Neda Zamani, B. Rosemary Grant, Peter R. Grant, Matthew T. Webster, and Leif Andersson. 2015. 'Evolution of Darwin's Finches and Their Beaks Revealed by Genome Sequencing'. *Nature* 518(7539):371–75. doi: 10.1038/nature14181.

Leroy, Thibault, Yoann Anselmetti, Marie-Ka Tilak, Sèverine Bérard, Laura Csukonyi, Maëva Gabrielli, Céline Scornavacca, Borja Milá, Christophe Thébaud, and Benoit Nabholz. 2021. 'A bird's white-eye view on avian sex chromosome evolution'. *Peer Community Journal* 1(e63). doi: 10.24072/pcjournal.70.

Leroy, Thibault, Marjolaine Rousselle, Marie-Ka Tilak, Aude E. Caizergues, Céline Scornavacca, María Recuerda, Jérôme Fuchs, Juan Carlos Illera, Dawie H. De Swardt, Guillermo Blanco, Christophe Thébaud, Borja Milá, and Benoit Nabholz.

2021. 'Island Songbirds as Windows into Evolution in Small Populations'. *Current*
Biology 31(6):1303-1310.e4. doi: 10.1016/j.cub.2020.12.040.

Leung, Brian, Anna L. Hargreaves, Dan A. Greenberg, Brian McGill, Maria Dornelas, and
Robin Freeman. 2020. 'Clustered versus Catastrophic Global Vertebrate Declines'.
Nature 588(7837):267–71. doi: 10.1038/s41586-020-2920-6.

Li, Heng, and Richard Durbin. 2009. 'Fast and Accurate Short Read Alignment with
Burrows–Wheeler Transform'. *Bioinformatics* 25(14):1754–60. doi:
10.1093/bioinformatics/btp324.

Li, Heng, and Richard Durbin. 2011. 'Inference of Human Population History from Individual
Whole-Genome Sequences'. *Nature* 475(7357):493–96. doi: 10.1038/nature10231.

Losos, Jonathan B., and Robert E. Ricklefs. 2009. 'Adaptation and Diversification on
Islands'. *Nature* 457(7231):830–36. doi: 10.1038/nature07893.

Maldonado-Coelho, Marcos. 2012. 'Climatic Oscillations Shape the Phylogeographical
Structure of Atlantic Forest Fire-Eye Antbirds (Aves: Thamnophilidae)'. *Biological*
Journal of the Linnean Society 105(4):900–924. doi: 10.1111/j.1095-
8312.2011.01823.x.

Marchant, Robert, and Henry Hooghiemstra. 2004. 'Rapid Environmental Change in African
and South American Tropics around 4000 Years before Present: A Review'. *Earth-*
Science Reviews 66(3):217–60. doi: 10.1016/j.earscirev.2004.01.003.

Martin, Claudia A., Claire Armstrong, Juan Carlos Illera, Brent C. Emerson, David S.
Richardson, and Lewis G. Spurgin. 2021. 'Genomic Variation, Population History and
within-Archipelago Adaptation between Island Bird Populations'. *Royal Society Open*
Science 8(2):201146. doi: 10.1098/rsos.201146.

Martins, Frederico C., Siobhan C. Cox, Martin Irestedt, Robert P. Prŷs-Jones nvestigation,
and Julia J. Day. 2020. 'A Comprehensive Molecular Phylogeny of Afrotropical
White-Eyes (Aves: Zosteropidae) Highlights Prior Underestimation of Mainland
Diversity and Complex Colonisation History'. *Molecular Phylogenetics and Evolution*
106843. doi: 10.1016/j.ympev.2020.106843.

Maruyama, T., and P. A. Fuerst. 1985. 'Population Bottlenecks and Nonequilibrium Models
in Population Genetics. III. Genic Homozygosity in Populations Which Experience
Periodic Bottlenecks'. *Genetics* 111(3):691–703. doi: 10.1093/genetics/111.3.691.

Mastretta-Yanes, Alicia, Alejandra Moreno-Letelier, Daniel Piñero, Tove H. Jorgensen, and
Brent C. Emerson. 2015. 'Biodiversity in the Mexican Highlands and the Interaction of
Geology, Geography and Climate within the Trans-Mexican Volcanic Belt'. *Journal of*
Biogeography 42(9):1586–1600. doi: 10.1111/jbi.12546.

Mazet, O., W. Rodríguez, S. Grusea, S. Boitard, and L. Chikhi. 2016. 'On the Importance of
Being Structured: Instantaneous Coalescence Rates and Human Evolution—
Lessons for Ancestral Population Size Inference?' *Heredity* 116(4):362–71. doi:
10.1038/hdy.2015.104.

McKenna, Aaron, Matthew Hanna, Eric Banks, Andrey Sivachenko, Kristian Cibulskis,
Andrew Kernytsky, Kiran Garimella, David Altshuler, Stacey Gabriel, Mark Daly, and
Mark A. DePristo. 2010. 'The Genome Analysis Toolkit: A MapReduce Framework

for Analyzing next-Generation DNA Sequencing Data'. *Genome Research*
20(9):1297–1303. doi: 10.1101/gr.107524.110.

Milá, Borja, Ben H. Warren, Philipp Heeb, and Christophe Thébaud. 2010. 'The Geographic
Scale of Diversification on Islands: Genetic and Morphological Divergence at a Very
Small Spatial Scale in the Mascarene Grey White-Eye (Aves: *Zosterops*
Borbonicus)'. *BMC Evolutionary Biology* 10:158. doi: 10.1186/1471-2148-10-158.

Moore, Jacob, William White, Debajyoti Paul, Robert Duncan, Wafa Abouchami, and
Stephen Galer. 2011. 'Evolution of Shield-Building and Rejuvenescent Volcanism of
Mauritius'. *Journal of Volcanology and Geothermal Research* 207:47–66. doi:
10.1016/j.jvolgeores.2011.07.005.

Moyle, Robert G., Christopher E. Filardi, Catherine E. Smith, and Jared Diamond. 2009.
'Explosive Pleistocene Diversification and Hemispheric Expansion of a "Great
Speciator"'. *Proceedings of the National Academy of Sciences* 106(6):1863–68. doi:
10.1073/pnas.0809861105.

Nadachowska-Brzyska, Krystyna, Reto Burri, Linnéa Smeds, and Hans Ellegren. 2016.
'PSMC Analysis of Effective Population Sizes in Molecular Ecology and Its
Application to Black-and-White Ficedula Flycatchers'. *Molecular Ecology* 25(5):1058–
72. doi: 10.1111/mec.13540.

Nadachowska-Brzyska, Krystyna, Cai Li, Linnea Smeds, Guojie Zhang, and Hans Ellegren.
2015. 'Temporal Dynamics of Avian Populations during Pleistocene Revealed by
Whole-Genome Sequences'. *Current Biology* 25(10):1375–80. doi:
10.1016/j.cub.2015.03.047.

Nei, Masatoshi, and Wen-Hsiung Li. 1979. 'Mathematical Model for Studying Genetic
Variation in Terms of Restriction Endonucleases'. *Proceedings of the National
Academy of Sciences of the United States of America* 76(10):5269–73. doi:
10.1073/pnas.76.10.5269.

Nei, Masatoshi, Takeo Maruyama, and Ranajit Chakraborty. 1975. 'The Bottleneck Effect
and Genetic Variability in Populations'. *Evolution* 29(1):1–10. doi: 10.2307/2407137.

Nogués-Bravo, David, Francisco Rodríguez-Sánchez, Luisa Orsini, Erik de Boer, Roland
Jansson, Helene Morlon, Damien A. Fordham, and Stephen T. Jackson. 2018.
'Cracking the Code of Biodiversity Responses to Past Climate Change'. *Trends in
Ecology & Evolution* 33(10):765–76. doi: 10.1016/j.tree.2018.07.005.

Norder, Sietze J., John B. Baumgartner, Paulo A. V. Borges, Tomislav Hengl, W. Daniel
Kissling, E. Emiel van Loon, and Kenneth F. Rijsdijk. 2018. 'A Global Spatially
Explicit Database of Changes in Island Palaeo-Area and Archipelago Configuration
during the Late Quaternary'. *Global Ecology and Biogeography* 27(5):500–505. doi:
10.1111/geb.12715.

Patton, Austin H., Mark J. Margres, Amanda R. Stahlke, Sarah Hendricks, Kevin Lewallen,
Rodrigo K. Hamede, Manuel Ruiz-Aravena, Oliver Ryder, Hamish I. McCallum,
Menna E. Jones, Paul A. Hohenlohe, and Andrew Storfer. 2019. 'Contemporary
Demographic Reconstruction Methods Are Robust to Genome Assembly Quality: A
Case Study in Tasmanian Devils'. *Molecular Biology and Evolution* 36(12):2906–21.
doi: 10.1093/molbev/msz191.

Purcell, Shaun, Benjamin Neale, Kathe Todd-Brown, Lori Thomas, Manuel A. R. Ferreira,
David Bender, Julian Maller, Pamela Sklar, Paul I. W. de Bakker, Mark J. Daly, and
Pak C. Sham. 2007. 'PLINK: A Tool Set for Whole-Genome Association and
Population-Based Linkage Analyses'. *The American Journal of Human Genetics*
81(3):559–75. doi: 10.1086/519795.

R core team. 2020. 'R: A Language and Environment for Statistical Computing'. *R*
Foundation for Statistical Computing, Vienna, Austria.

Rijsdijk, Kenneth F., Jens Zinke, Perry G. B. de Louw, Julian P. Hume, Hans J. van der
Plicht, Henry Hooghiemstra, Hanneke J. M. Meijer, Hubert B. Vonhof, Nick Porch, F.
B. Vincent Florens, Claudia Baider, Bas van Geel, Joost Brinkkemper, Tamara
Vernimmen, and Anwar Janoo. 2011. 'Mid-Holocene (4200 Kyr BP) Mass Mortalities
in Mauritius (Mascarenes): Insular Vertebrates Resilient to Climatic Extremes but
Vulnerable to Human Impact'. *The Holocene* 21(8):1179–94. doi:
10.1177/0959683611405236.

Rodríguez, Willy, Olivier Mazet, Simona Grusea, Armando Arredondo, Josué M. Corujo,
Simon Boitard, and Lounès Chikhi. 2018. 'The IICR and the Non-Stationary
Structured Coalescent: Towards Demographic Inference with Arbitrary Changes in
Population Structure'. *Heredity* 121(6):663–78. doi: 10.1038/s41437-018-0148-0.

Salces-Castellano, Antonia, Sean Stankowski, Paula Arribas, Jairo Patiño, Dirk N. Karger,
Roger Butlin, and Brent C. Emerson. 2021. 'Long-Term Cloud Forest Response to
Climate Warming Revealed by Insect Speciation History'. *Evolution* 75(2):231–44.
doi: 10.1111/evo.14111.

Salmona, Jordi, Rasmus Heller, Martin Lascoux, and Aaron B. A. Shafer. 2017. 'Inferring
Demographic History Using Genomic Data'. Pp. 511–37 in *Population Genomics:*
Concepts, Approaches and Applications. Springer.

Salmona, Jordi, Rasmus Heller, Erwan Quéméré, and Lounès Chikhi. 2017. 'Climate
Change and Human Colonization Triggered Habitat Loss and Fragmentation in
Madagascar'. *Molecular Ecology* 26(19):5203–22. doi: 10.1111/mec.14173.

Salmona, Jordi, Marc Salamolard, Damien Fouillot, Thomas Ghestemme, Jerry Larose,
Jean-François Centon, Vitor Sousa, Deborah A. Dawson, Christophe Thebaud, and
Lounès Chikhi. 2012. 'Signature of a Pre-Human Population Decline in the Critically
Endangered Reunion Island Endemic Forest Bird Coracina Newtoni'. *PLOS ONE*
7(8):e43524. doi: 10.1371/journal.pone.0043524.

Schiffels, Stephan, and Richard Durbin. 2014. 'Inferring Human Population Size and
Separation History from Multiple Genome Sequences'. *Nature Genetics* 46(8):919–
25. doi: 10.1038/ng.3015.

Singhal, Sonal, Ellen M. Leffler, Keerthi Sannareddy, Isaac Turner, Oliver Venn, Daniel M.
Hooper, Alva I. Strand, Qiye Li, Brian Raney, Christopher N. Balakrishnan, Simon C.
Griffith, Gil McVean, and Molly Przeworski. 2015. 'Stable Recombination Hotspots in
Birds'. *Science* 350(6263):928–32. doi: 10.1126/science.aad0843.

Smeds, Linnéa, Anna Qvarnström, and Hans Ellegren. 2016. 'Direct Estimate of the Rate of
Germline Mutation in a Bird'. *Genome Research* 26(9):1211–18. doi:
10.1101/gr.204669.116.

Staubwasser, M., F. Sirocko, P. M. Grootes, and M. Segl. 2003. 'Climate Change at the 4.2
835   Ka BP Termination of the Indus Valley Civilization and Holocene South Asian
836   Monsoon Variability'. *Geophysical Research Letters* 30(8):1425–32. doi:
837   10.1029/2002GL016822.

838   Tajima, F. 1983. 'Evolutionary Relationship of DNA Sequences in Finite Populations'.
839   *Genetics* 105(2):437–60. doi: 10.1093/genetics/105.2.437.

840   Tajima, F. 1989. 'Statistical Method for Testing the Neutral Mutation Hypothesis by DNA
841   Polymorphism'. *Genetics* 123(3):585–95. doi: 10.1093/genetics/123.3.585.

842   Terhorst, Jonathan, John A. Kamm, and Yun S. Song. 2017. 'Robust and Scalable Inference
843   of Population History from Hundreds of Unphased Whole-Genomes'. *Nature*
844   *Genetics* 49(2):303–9. doi: 10.1038/ng.3748.

845   Thébaud, Christophe, Ben H. Warren, A. Cheke, and Dominique Strasberg. 2009.
846   'Mascarene Islands, Biology'. Pp. 612–19 in *Encyclopedia of Islands*. Berkeley:
847   University of California Press.

848   Thompson, L. G. 2002. 'Kilimanjaro Ice Core Records: Evidence of Holocene Climate
849   Change in Tropical Africa'. *Science* 298(5593):589–93. doi:
850   10.1126/science.1073198.

851   Urban, M. C., G. Bocedi, A. P. Hendry, J. B. Mihoub, G. Pe'er, A. Singer, J. R. Bridle, L. G.
852   Crozier, L. De Meester, W. Godsoe, A. Gonzalez, J. J. Hellmann, R. D. Holt, A. Huth,
853   K. Johst, C. B. Krug, P. W. Leadley, S. C. F. Palmer, J. H. Pantel, A. Schmitz, P. A.
854   Zollner, and J. M. J. Travis. 2016. 'Improving the Forecast for Biodiversity under
855   Climate Change'. *Science* 353(6304):aad8466. doi: 10.1126/science.aad8466.

856   Wakeley, J. 1999. 'Nonequilibrium Migration in Human History'. *Genetics* 153(4):1863–71.

857   Warren, Ben H., Eldredge Bermingham, Robert P. Prys-Jones, and Christophe Thébaud.
858   2006. 'Immigration, Species Radiation and Extinction in a Highly Diverse Songbird
859   Lineage: White-Eyes on Indian Ocean Islands'. *Molecular Ecology* 15(12):3769–86.
860   doi: 10.1111/j.1365-294X.2006.03058.x.

861   Warren, Ben H., Daniel Simberloff, Robert E. Ricklefs, Robin Aguilée, Fabien L. Condamine,
862   Dominique Gravel, Hélène Morlon, Nicolas Mouquet, James Rosindell, Juliane
863   Casquet, Elena Conti, Josselin Cornuault, José María Fernández-Palacios, Tomislav
864   Hengl, Sietze J. Norder, Kenneth F. Rijsdijk, Isabel Sanmartín, Dominique Strasberg,
865   Kostas A. Triantis, Luis M. Valente, Robert J. Whittaker, Rosemary G. Gillespie,
866   Brent C. Emerson, and Christophe Thébaud. 2015. 'Islands as Model Systems in
867   Ecology and Evolution: Prospects Fifty Years after MacArthur-Wilson'. *Ecology*
868   *Letters* 18(2):200–217. doi: 10.1111/ele.12398.

869   Watterson, G. A. 1975. 'On the Number of Segregating Sites in Genetical Models without
870   Recombination'. *Theoretical Population Biology* 7(2):256–76. doi: 10.1016/0040-
871   5809(75)90020-9.

872   Watterson, G. A. 1984. 'Allele Frequencies after a Bottleneck'. *Theoretical Population*
873   *Biology* 26(3):387–407. doi: 10.1016/0040-5809(84)90042-X.

874   Weigelt, Patrick, and Holger Kreft. 2013. 'Quantifying Island Isolation – Insights from Global
875   Patterns of Insular Plant Species Richness'. *Ecography* 36(4):417–29. doi:
876   10.1111/j.1600-0587.2012.07669.x.

Weir, B. S., and C. Clark Cockerham. 1984. 'Estimating F-Statistics for the Analysis of
Population Structure'. *Evolution* 38(6):1358–70.

Weir, Jason T., and Adam Lawson. 2015. 'Evolutionary Rates across Gradients'. *Methods in
Ecology and Evolution* 6(11):1278–86. doi: 10.1111/2041-210X.12419.

Whittaker, Robert J., Kostas A. Triantis, and Richard J. Ladle. 2008. 'A General Dynamic
Theory of Oceanic Island Biogeography'. *Journal of Biogeography* 35(6):977–94. doi:
10.1111/j.1365-2699.2008.01892.x.

Wilder, Aryn P., Megan A. Supple, Ayshwarya Subramanian, Anish Mudide, Ross Swofford,
Aitor Serres-Armero, Cynthia Steiner, Klaus-Peter Koepfli, Diane P. Genereux, Elinor
K. Karlsson, Kerstin Lindblad-Toh, Tomas Marques-Bonet, Violeta Munoz Fuentes,
Kathleen Foley, Wynn K. Meyer, Zoonomia Consortium, Oliver A. Ryder, and Beth
Shapiro. 2023. 'The Contribution of Historical Processes to Contemporary Extinction
Risk in Placental Mammals'. *Science* 380(6643):eabn5856. doi:
10.1126/science.abn5856.

Wright, Sewall. 1943. 'Isolation by Distance'. *Genetics* 28(2):114–38.

Yan, Mi, Bin Wang, and Jian Liu. 2016. 'Global Monsoon Change during the Last Glacial
Maximum: A Multi-Model Study'. *Climate Dynamics* 47(1):359–74. doi:
10.1007/s00382-015-2841-5.

895
896

Table 1: Parameter estimation for two-population demographic scenarios in DILS. Numbers
outside brackets indicate median values while numbers inside brackets indicate confidence
intervals (2.5%-97.5%). Nine parameters were estimated, namely the population size of the
ancestral population (N_{ANC}), the time of the split between the two populations (T_{split}), the
population sizes of the daughter populations after the split (N_{ANC1} and N_{ANC2}), the time of the
potential demographic change (T_{dem1} and T_{dem2}), the population sizes of the daughter
populations after the demographic change (N_{POP1} and N_{POP2}), and the time of ancestral
migration cessation (T_{AM}). For each pair, populations 1 and 2 are the first and second
population of the pair, respectively. See Figure S2B for a representation of this scenario of
divergence.

	ABC model parameters								
	N_{ANC}	T_{split}	N_{ANC1}	N_{ANC2}	T_{dem1}	N_{POP1}	T_{dem2}	N_{POP2}	T_{AM}
Reunion Grey White-eye - Mauritius Grey White-eye	115,681 (103,039-130,232)	128,291 (95,638-168,209)	46,070 (24,675-76,275)	48,886 (30,636-77,610)	98,879 (80,019-121,191)	631,186 (487,328-742,422)	69,215 (55,562-86,687)	537,411 (396,932-661,929)	92,045 (75,532-110,466)
Reunion Grey White-eye (GHB form) - Mauritius Grey White-eye	84,075 (73,819-97,160)	115,323 (86,075-152,274)	66,184 (36,219-109,992)	47,120 (27,039-74,713)	91,004 (69,428-116,346)	443,404 (338,203-581,931)	60,632 (46,932-80,056)	405,894 (284,879-536,721)	93,817 (74,518-112,903)
Reunion Grey White-eye - Reunion Olive White-eye	234,347 (200,173-275,027)	318,314 (255,817-398,370)	67,133 (22,956-120,907)	55,254 (24,862-96,435)	234,291 (188,145-290,008)	797,268 (604,018-915,561)	293,743 (234,835-364,791)	816,811 (651,688-945,258)	299,302 (241,213-350,777)
Mauritius Grey White-eye - Reunion Olive White-eye	237,298 (192,706-277,644)	431,242 (299,381-559,243)	76,860 (41,129-120,390)	34,639 (16,827-58,066)	118,964 (91,229-141,605)	745,166 (556,712-916,230)	291,374 (218,052-356,087)	407,800 (322,593-453,945)	316,104 (258,245-385,771)

<u>Figure 1</u>: Proposed strategy to compare the effects of local events and regional events on
genetic diversity by comparing the demographic histories of closely related populations or
species inhabiting different islands from the same volcanic oceanic archipelago. Plots
represent effective population size (N_e) through time for a population or species, and numbers
indicate different events of ecological disruption associated with bottlenecks: event 1 is a
volcanic eruption on island A while event 2 corresponds to a regional drop in temperatures,
as observed during a glacial maximum, that affects demographic histories on islands A and B
synchronously, all else being equal. Contemporary bottlenecks suggest the importance of
events at a large geographic scale while bottlenecks associated with island-specific events
suggest the importance of local events as drivers of population histories.
918

Figure 2: Distribution range and sampling localities. Localities are labelled with numbers (see
Table S1 for names and coordinates) and circles with different colours represent samples from
Reunion Grey White-eye (purple), Reunion Olive White-eye (green), and Mauritius Grey
White-eye (orange). Grey colours represent elevation classes; for Reunion: light grey =
elevation below 1,000 m, medium grey = elevation between 1,000 and 2,000 m, dark grey =
elevation above 2,000 m; for Mauritius: light grey = elevation below 400 m, medium grey =
elevation between 400 and 600 m, dark grey = elevation above 600 m. Dotted lines separate
the distribution ranges of the four Reunion Grey White-eye geographic forms. Phylogenetic
relationships between the four endemic Mascarene white-eyes are adapted from Warren et
al. (2006), with the asterisk denoting the divergence time between the two sister lineages,
estimated at 1.2 Mya. Illustrations from Birds of the World reproduced with permission from
Lynx edicions. Photo credits: Stanislas Harvančík (Reunion Grey White-eye), Simon Colenutt
(Reunion Olive White-eye), and Hilary Jones (Mauritius Grey White-eye).

<u>Figure 3</u>: PSMC inferences of population size histories over time. Colours correspond to the
three white-eye species, and each line represents a different individual. Vertical dotted lines
and grey vertical bars indicate (from left to right): the Last Glacial Maximum 21,000 ya and a
catastrophic period of large explosive eruptions in Reunion around 200,000 ya. In Mauritius,
volcanic activity has been quite intense throughout the last 500,000 years but was never
explosive during that period. Please note that the plots are in log-scales. Illustrations from
Birds of the World reproduced with permission from Lynx edicions.

942