Title: Functional connectome through the human life span

Authors: Lianglong Sun[1,2,3], Tengda Zhao[1,2,3,#], Xinyuan Liang[1,2,3,#], Mingrui Xia[1,2,3,#],
Qiongling Li[1,2,3], Xuhong Liao[4], Gaolang Gong[1,2,3,5], Qian Wang[1,2,3], Chenxuan Pang[1,2,3], Qian
Yu[1,2,3], Yanchao Bi[1,2,3,5], Pindong Chen[6], Rui Chen[1], Yuan Chen[7], Taolin Chen[8], Jingliang Cheng[7],
Yuqi Cheng[9], Zaixu Cui[5], Zhengjia Dai[1,2,3], Yao Deng[1], Yuyin Ding[1], Qi Dong[1], Dingna Duan[1,2,3],
Jia-Hong Gao[10,11,12], Qiyong Gong[8,13], Ying Han[14], Zaizhu Han[1,3], Chu-Chung Huang[15], Ruiwang
Huang[1,3], Ran Huo[16], Lingjiang Li[17,18], Ching-Po Lin[19,20,21], Qixiang Lin[1,2,3], Bangshan
Liu[17,18],Chao Liu[1,3], Ningyu Liu[1], Ying Liu[16], Yong Liu[22], Jing Lu[1], Leilei Ma[1], Weiwei Men[10,11],
Shaozheng Qin[1,2,3,5], Jiang Qiu[23,24], Shijun Qiu[25], Tianmei Si[26], Shuping Tan[27], Yanqing Tang[28],
Sha Tao[1], Dawei Wang[29], Fei Wang[28], Jiali Wang[1], Pan Wang[30], Xiaoqin Wang[23,24], Yanpei Wang[1],
Dongtao Wei[23,24], Yankun Wu[26], Peng Xie[31,32], Xiufeng Xu[9], Yuehua Xu[1,2,3], Zhilei Xu[1,2,3],
Liyuan Yang[1,2,3], Huishu Yuan[16], Zilong Zeng[1,2,3], Haibo Zhang[1], Xi Zhang[33], Gai Zhao[1], Yanting
Zheng[25], Suyu Zhong[22], Alzheimer's Disease Neuroimaging Initiative, Cam-CAN, Developing
Human Connectome Project, DIDA-MDD Working Group, MCADI, NSPN, and Yong He[1,2,3,5,*]

Affiliations:

[1] State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing, China;
[2] Beijing Key Laboratory of Brain Imaging and Connectomics, Beijing Normal University, Beijing, China;
[3] IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing, China;
[4] School of Systems Science, Beijing Normal University, Beijing, China;
[5] Chinese Institute for Brain Research, Beijing, China;
[6] Brainnetome Center & National Laboratory of Pattern Recognition, Institute of Automation, Chinese
Academy of Sciences, Beijing, China;
[7] Department of Magnetic Resonance Imaging, The First Affiliated Hospital of Zhengzhou University,
Zhengzhou, China;
[8] Huaxi MR Research Center (HMRRC), Department of Radiology, West China Hospital, Sichuan University,
Chengdu, China;
[9] Department of Psychiatry, First Affiliated Hospital of Kunming Medical University, Kunming, China;
[10] Center for MRI Research, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing,
China;
[11] Beijing City Key Laboratory for Medical Physics and Engineering, Institute of Heavy Ion Physics, School of
Physics, Peking University, Beijing, China;
[12] IDG/McGovern Institute for Brain Research, Peking University, Beijing, China;
[13] Research Unit of Psychoradiology, Chinese Academy of Medical Sciences, Chengdu, China;
[14] Department of Neurology, Xuanwu Hospital of Capital Medical University, Beijing, China;
[15] Key Laboratory of Brain Functional Genomics (Ministry of Education), Affiliated Mental Health Center
(ECNU), School of Psychology and Cognitive Science, East China Normal University, Shanghai, China;
[16] Department of Radiology, Peking University Third Hospital, Beijing, China;
[17] Department of Psychiatry, and National Clinical Research Center for Mental Disorders, The Second Xiangya
Hospital of Central South University, Changsha, China;
[18] Mental Health Institute of Central South University, China National Technology Institute on Mental
Disorders, Hunan Technology Institute of Psychiatry, Hunan Key Laboratory of Psychiatry and Mental Health,
Hunan Medical Center for Mental Health, Changsha, China;
[19] Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai, China;
[20] Institute of Neuroscience, National Yang Ming Chiao Tung University, Taipei, China;
[21] Department of Education and Research, Taipei City Hospital, Taipei, China;
[22] Center for Artificial Intelligence in Medical Imaging, School of Artificial Intelligence, Beijing University of
Posts and Telecommunications, Beijing, China;
[23] Key Laboratory of Cognition and Personality (SWU), Ministry of Education, Chongqing, China;

[24] Department of Psychology, Southwest University, Chongqing, China;
[25] Department of Radiology, The First Affiliated Hospital of Guangzhou University of Chinese Medicine,
Guangzhou, China;
[26] Peking University Sixth Hospital, Peking University Institute of Mental Health, NHC Key Laboratory of
Mental Health (Peking University), National Clinical Research Center for Mental Disorders (Peking
University Sixth Hospital), Peking University, Beijing, China;
[27] Beijing Huilongguan Hospital, Peking University Huilongguan Clinical Medical School, Beijing, China;
[28] Department of Psychiatry, The First Affiliated Hospital of China Medical University, Shenyang, China;
[29] Department of Radiology, Qilu Hospital of Shandong University, Ji'nan, China;
[30] Department of Neurology, Tianjin Huanhu Hospital, Tianjin University, Tianjin, China;
[31] Chongqing Key Laboratory of Neurobiology, Chongqing, China;
[32] Department of Neurology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China;
[33] Department of Neurology, the Second Medical Centre, National Clinical Research Centre for Geriatric
Diseases, Chinese PLA General Hospital, Beijing, China.
63

64 [#] These authors contributed equally

65 [*] Correspondence to: yong.he@bnu.edu.cn

66 **Keywords**: brain chart, brain atlas, connectomics, fMRI

Abstract

The lifespan growth of the functional connectome remains unknown. Here, we assemble task-free functional and structural magnetic resonance imaging data from 33,250 individuals aged 32 postmenstrual weeks to 80 years from 132 global sites. We report critical inflection points in the nonlinear growth curves of the global mean and variance of the connectome, peaking in the late fourth and late third decades of life, respectively. After constructing a fine-grained, lifespan-wide suite of system-level brain atlases, we show distinct maturation timelines for functional segregation within different systems. Lifespan growth of regional connectivity is organized along a primary-to-association cortical axis. These connectome-based normative models reveal substantial individual heterogeneities in functional brain networks in patients with autism spectrum disorder, major depressive disorder, and Alzheimer's disease. These findings elucidate the lifespan evolution of the functional connectome and can serve as a normative reference for quantifying individual variation in development, aging, and neuropsychiatric disorders.

Introduction

The resting human brain, characterized by intrinsic or spontaneous brain activities, has been increasingly understood from a connectome perspective over the past two decades [1-5]. The emergence, development, and aging of the intrinsic connectome architecture enables the dynamic reorganization of functional specialization and integration throughout the lifespan, contributing to continuous changes in human cognition and behavior [6-9]. Understanding the spatiotemporal growth process of the typical functional connectome is critical for elucidating network-level developmental principles in healthy individuals and for pinpointing periods of heightened vulnerability or potential. Disruption of these normative connectome patterns, especially during specific time windows, can predispose individuals to a spectrum of neurodevelopmental [10-12], neurodegenerative [13], and psychiatric disorders [14-16]. The growth chart framework provides an invaluable tool for charting normative reference curves in the human brain [17-20]. Recently, Bethlehem et al. [18] delineated the life-cycle growth curves of brain morphometry by aggregating the largest multisite structural magnetic resonance imaging (MRI) dataset to date (101,457 individuals between 115 days post-conception to 100 years of age), marking a significant step toward reproducible and generalizable brain charts. However, the normative growth charts of the functional brain connectome across the human lifespan remain unknown.

Previous studies using task-free functional MRI (fMRI) data have reported age-related characteristics of the functional connectome [21-23]. However, most of these studies were limited to specific periods of growth with narrow age intervals. For example, data from the perinatal and early postnatal period (e.g., 0-6 years) are rarely included in studies spanning childhood, adolescence, and adulthood; thus missing the opportunity to depict a continuous life-cycle dynamic evolution from gestation to old age. Although a few studies have attempted to include a broader age range from childhood to late adulthood, they have suffered from challenges in robustly estimating normative growth curves due to limited sample sizes (typically < 1,000) [24-29]. More recently, Rutherford et al. [30] have made great strides in establishing a lifespan normative model of the functional connectome using a large sample dataset (~22,000 individuals aged 2-100 years). However, this work primarily focused on intersystem functional connectivity using population-based system-level atlas. Furthermore, there are large inconsistencies in the literature regarding functional connectivity trajectories, with no consensus emerging on the developmental directions and growth milestones. In particular, Cao et al. [25] reported that global functional connectivity in the whole brain peaks at around 30 years of age, whereas other studies suggest earlier peaks [24] or show a continuous decline across the lifespan [31]. Different trends have been observed for sensorimotor regions, with reports of ascending [32], descending [33], and stable [34] developmental trajectories from childhood to adolescence. Similarly, connectivity patterns between the default and frontoparietal networks have been reported to both increase [35] and decrease [36, 37] during this period. Such discrepancies between studies are likely due to the high sensitivity of high-dimensional fMRI data to variations in scanner platforms and sequences, image quality, data processing, and statistical methods, as well as the population heterogeneity of cohorts [6]. This underscores the paramount importance of large sample sizes, rigorous data quality control procedures, consistent data processing protocols, and standardized statistical modeling frameworks to accurately characterize growth curves of the functional connectome across the lifespan.

To address this gap, we assembled a large multimodal neuroimaging dataset with rigorous

quality control, consisting of cross-sectional task-free fMRI and structural MRI data from 33,250 individuals ranging in age from 32 postmenstrual weeks to 80 years, collected from 132 global sites (Fig. 1a). We conducted a comprehensive network modeling analysis to delineate the nonlinear growth patterns of the functional connectome across multiple scales. We began by characterizing lifespan growth in the overall patterns of the global functional connectome, revealing important life-course milestones. We then constructed continuous age-related, system-level atlases across the lifespan and further provided a previously unreported portrayal of the distinct growth patterns across brain systems. Next, we sought to elucidate the spatiotemporal principles governing connectome growth at a finer regional scale. Finally, we investigated the potential clinical value of the established connectome-based normative models. We selected autism spectrum disorder (ASD, N = 414), major depressive disorder (MDD, N = 622), and Alzheimer's disease (AD, N = 180) as representative conditions characterized by network dysfunction across different life stages. These conditions typically manifest in childhood, adolescence/adulthood, and older adulthood, respectively [16, 38-40]. Using individual deviation scores relative to the 50th percentile, we presented a multiscale characterization to quantify the individual heterogeneity of patients with ASD, MDD, or AD.

Results

We initially aggregated 44,030 participants with multimodal structural MRI and task-free fMRI data. After a rigorous quality control process (for details, see the Methods and Supplementary Figs 1 and 2), we obtained a final sample of 34,466 participants with high-quality imaging data, including 33,250 healthy individuals (Fig. 1a) and 1,216 patients. The detailed demographics and acquisition parameters of the datasets are provided in Supplementary Tables 1 and 2, respectively. Using the standardized and highly uniform processing pipeline (Methods and Supplementary Fig. 3), we obtained the surface-based preprocessed blood oxygenation level-dependent (BOLD) signals in fsaverage4 space for each participant (4,609 vertices in total). We then constructed a vertexwise $4,609 \times 4,609$ functional connectome matrix by calculating Pearson's correlation coefficient between the time courses of each vertex. Figure 1b shows the functional connectome matrices of representative participants at different ages. Next, we examined the individual connectome at the global, system, and vertex levels. In accordance with the recommendations of the World Health Organization recommendation [41], the age-related nonlinear growth patterns were described using the generalized additive model for location, scale, and shape (GAMLSS) [41, 42], based on cross-sectional data from healthy populations (N = 33,250). Sex and in-scanner head motion (mean framewise displacement) were included as fixed effect covariates, and the scanner site was included as a random effect covariate. GAMLSS provides a robust framework for modeling nonlinear growth curves and has been widely used in neurodevelopmental studies [18, 43-45]. To assess the rate of growth (velocity) and inflection points, we calculated the first derivatives of the lifespan growth curves. The GAMLSS specifications, model estimations, and model evaluations are detailed in the Methods section.

Mapping the normative growth of the global functional connectome across the lifespan

To provide basic developmental and aging insights into the global functional connectome, we first characterized the normative growth patterns of the global mean and variance (estimated by standard deviation) of the functional connectome. The lifespan curve of the global mean of functional connectome (Fig. 1c) exhibited a nonlinear increase from 32 postmenstrual weeks

onward, peaking in the late fourth decade of life (38.2 years, 95% bootstrap confidence interval
(CI) 36.0-40.1), followed by a nonlinear decline. This growth curve is primarily driven by age-
related changes of middle- and long-range connections (Supplementary Fig. 4). The global
variance of functional connectome (Fig. 1d) also exhibited a nonlinear growth pattern, reaching
its peak in the late third decade of life (28.2 years, 95% bootstrap CI 26.4-30.1). The utilization
of the GAMLSS enabled the delineation of normative growth curves for interindividual
variability [18] in the two global measures (Supplementary Result 1 and Supplementary Fig. 5a).
The curves demonstrated a slight decline in inter-individual variability during the initial stages of
early development, a gradual increase until the late sixth decade of life (peaking at 55.2 years, 95%
CI [53.9, 56.0] for the global mean; peaking at 56.8 years, 95% CI [55.1, 58.1] for the global
variance), and then a rapid decline. These nonlinear growth patterns in the global connectome
measures indicated a temporally coordinated manner across the lifespan.

181
Fig. 1 | Normative growth patterns of the functional connectome at a global level over the lifespan. a,
Quality-controlled MRI data from 132 scanning sites comprising 33,250 healthy participants who collectively
spanned the age range from 32 postmenstrual weeks to 80 years. Box plots show the age distribution of
participants at each site of data acquisition. The detailed participant demographics and acquisition parameters
of each site are provided in Supplementary Tables 1 and 2, respectively. **b,** The functional connectome matrices
of representative participants at different ages. **c,** Normative growth curve (left panel) and growth rate (right
panel) of the global mean of the connectome as estimated by GAMLSS. The median (50th) centile is
represented by a solid line, while the 5th, 25th, 75th, and 95th centiles are indicated by dotted lines. The
growth rate is characterized by the first derivative of the median centile line. The gray shaded areas represent
the 95% confidence interval, which was estimated by bootstrapping 1,000 times (see Methods for details). **d,**
Normative growth curve (left panel) and growth rate (right panel) of global variance of the connectome. wk,
week; yr, year.

Lifespan growth of system-specific organization in the functional connectome

Functional segregation and integration are two fundamental organizational principles of the human brain connectome [1]. To understand the lifespan growth patterns of functional segregation and integration, we established the normative models of the functional connectome at the systems level. The first step was to parcellate the cortex into distinct functional systems for each participant. Converging evidence has shown that relying on population-level atlases for individual analysis overlooks crucial intersubject variability in functional topography organization [46-49]. This oversight leads to the misinterpretation of spatial distribution differences as system-level disparities [47, 50], thereby increasing the risk of inaccuracies in mapping both intra- and intersystem connectivity. Moreover, although previous studies of fetal and infant brains have elucidated the early emergence of basic forms of large-scale functional systems, including the visual [51-54], somatomotor [51-54], dorsal attention [55, 56], ventral attention [51], frontoparietal [52, 54, 56], and default mode networks [51-54, 56], the functional architecture of an individual's system undergoes dramatic refinement and reorganization over the protracted life course [21, 57]. To increase the precision of the construction of individual-specific functional networks, it is essential to establish a set of continuous growth atlases with accurate system correspondences across the life course.

To address this issue, we proposed a Gaussian-weighted iterative age-specific group atlas (GIAGA) generation approach (see Methods and Supplementary Fig. 6a). The iterative refinement process is central to this approach. Briefly, we first divided all participants aged 32 postmenstrual weeks to 80 years into 26 distinct age groups. Yeo's adult atlas [58] was then used as a prior to generate a personalized parcellation for each participant in a given age group. These personalized parcellations were further aggregated to construct an age-specific population-level atlas, where the contribution of participants was weighted according to their age position within a Gaussian probability distribution. This process was repeated until the age-specific population-level atlas converged, resulting in a set of age-specific brain atlases across the lifespan (Fig. 2a, Supplementary Figs 7 and 8). Validation analysis revealed greater global homogeneity when using these age-specific group atlases than using the adult-based group atlas across all age groups (all $p < 10^{-9}$, Bonferroni-corrected, Supplementary Fig. 9), particularly evident during early development. Notably, each of the 26 brain atlases was parcellated into seven canonical functional networks. For each network, we calculated the network size ratio, measured by the proportion of vertices, and the distribution score, defined by the number of spatially discontinuous subregions (Fig. 2b). We found that the default mode (DM), frontoparietal (FP), and ventral attention (VA) networks showed a slight expansion in network size during the first month of life, while their distribution scores developed until early childhood (4-6 years). In contrast, the somatomotor (SM), visual (VIS), and dorsal attention (DA) networks showed a relatively stable pattern of network size and network discretization throughout the lifespan. A hierarchical clustering analysis of these system-level brain atlases revealed three overarching patterns. Cluster I covered atlases from 34 postmenstrual weeks to 1 month, cluster II covered atlases from 3 months to 24 months, and cluster III covered atlases from 4 years to 80 years of age (Supplementary Fig. 10). To further quantify the growth patterns of the whole-cortical atlas and the system-specific atlases, we computed their network similarity to the designated reference atlas using both the overlay index and the Dice coefficient (Methods). The reference atlas was derived from the average of eight adult-like atlases, identified as a homogeneous cluster of 18- to 80-year-old atlases (Supplementary Fig. 10). We found that the overall similarity of the whole-

cortical atlas exhibited a rapid increase during the first two decades of life, followed by a plateau, and a subsequent slight decrease with age (Fig. 2c). At the system level, we observed that both the VIS and SM networks exhibited adult-like patterns (80% similarity) in the perinatal period, whereas the DM, FP, DA, and VA networks developed adult-like patterns (80% similarity) at 4-6 years of age (Fig. 2d and 2e).

Fig. 2 | Population-level and individual-level functional atlases throughout the lifespan. a, Employing the Gaussian-weighted iterative group atlas generation approach (for details, see Methods and Supplementary Fig. 6a), the lifespan set of seven-network functional atlases from 32 postmenstrual weeks to 80 years was established (26 atlases in total). Only the left hemisphere is displayed here; for the whole-cortical atlases, refer to Supplementary Figs 7 and 8. Labels of each system were mapped onto the HCP fs_LR_32k surface and visualized using BrainNet Viewer [59]. **b**, Network size ratio and network distribution score of each system in all age-specific group atlases. The network size ratio was calculated as the vertex number of the system divided by the total cortical vertex number. The network distribution score was measured by the number of spatially discontinuous subregions (≥ 5 vertices) in the system. **c**, Global similarity of each age-specific group atlas with the reference atlas across the lifespan. The degree of global similarity was defined as the number of vertices with the same label in the two atlases divided by the total number of vertices in both atlases. **d**, System similarity of each age-specific group atlas with the corresponding system in the reference atlas across the lifespan. System similarity was quantified using the Dice coefficient. **e**, The ages at which the system similarity of each age-specific group atlas reached 0.8 and 0.98. **f-g**, Normative growth curve and growth rate of global atlas similarity with the reference atlas when using personalized functional atlas for each participant. The gray shaded areas represent the 95% confidence interval, which was estimated by bootstrapping 1,000 times. VIS, visual; SM, somatomotor; DA, dorsal attention; VA, ventral attention; LIM, limbic; FP, frontoparietal; DM, default mode. wk, week; mon, month; yr, year.

Based on the age-specific group atlases established above, we proceeded to map individual-level functional systems for each participant. Specifically, we used an iterative parcellation procedure (see Methods and Supplementary Fig. 6b), as proposed by Wang et al. [60], which has been demonstrated to accurately identify personalized functional networks in both healthy [47, 60] and diseased individuals [61-63]. As expected, the individual-level atlases exhibited significantly greater global homogeneity than both the age-specific group atlases (all $p < 10^{-9}$, Bonferroni-corrected) and the adult-based group atlas (all $p < 10^{-8}$, Bonferroni-corrected), regardless of the age groups considered (Supplementary Fig. 9). Consistent with the growth pattern observed in the age-specific group atlas (Fig. 2c), the global similarity of the individualized atlas to the reference increased from 32 postmenstrual weeks and reached a peak in adulthood (31.6 years, 95% bootstrap CI 30.5-32.9) (Fig. 2f and 2g).

Using the person-specific network mapping approach, which integrates individual-level iterative processes with the age-specific group atlases, we characterized the lifespan growth patterns of within-system connectivity (functional segregation) and between-system connectivity (functional integration) (Supplementary Result 2, Supplementary Figs 11 and 12). To further quantify the differences in within-system connectivity relative to between-system connectivity, we calculated the system segregation index for each brain system [64]. This index measures the difference between mean within-system connectivity and mean between-system connectivity as a proportion of mean within-system connectivity [64] (Methods). Interestingly, global segregation across all systems peaked in the third decade of life (25.7 years, 95% bootstrap CI 24.8-26.8) (Fig. 3a). At the system level, different networks manifested distinct nonlinear growth patterns (Fig. 3b-3d). The primary VIS network consistently showed the greatest segregation across all ages (Fig. 3b and 3c), suggesting that the VIS network is more functionally specialized and relatively less integrated in inter-network communication compared to other systems. The DA and VIS networks exhibited similar trends in life-cycle growth patterns, peaking in early childhood and pre-adolescence, respectively (Fig. 3b and 3c). The DM and FP networks showed the lowest levels of segregation in the early stages of neurodevelopment (Fig. 3b and 3c). However, segregation increased rapidly with age peaks at the end of the third decade and decreased rapidly in the late stages of senescence (Fig. 3b-3d). Finally, the SM and VA networks showed similar growth patterns of system segregation, increasing and decreasing moderately over the lifetime (Fig. 3b-3d).

Fig. 3 | Lifespan normative growth patterns of brain system segregation. a, Normative growth curve and growth rate of global system segregation. The peak occurred in the third decade of life (25.7 years, 95% bootstrap confidence interval 24.8-26.8). The gray shaded areas represent the 95% confidence interval, which was estimated by bootstrapping 1,000 times. **b-c**, Normative growth curves and growth rate of system segregation for each network. The median (50th) centile is represented by a solid line, while the 5th, 25th, 75th, and 95th centiles are indicated by dotted lines. The key inflection points are marked in blue font. **d**, Growth rate of system-specific segregation visualized in the cortex, with black lines depicting system boundaries. The values of each system are mapped and visualized on the HCP fs_LR_32k surface. VIS, visual; SM, somatomotor; DA, dorsal attention; VA, ventral attention; LIM, limbic; FP, frontoparietal; DM, default mode. wk, week; yr, year.

Lifespan growth of functional connectivity at the regional level reveals a spatial gradient pattern

Having identified distinct growth patterns in different brain systems, we further explored the more nuanced spatiotemporal growth patterns of the functional connectome at the regional level. First, we plotted the normative growth curves of each vertex's functional connectivity strength (FCS) by calculating the average connectivity with all other vertices. Figure 4a shows the curves for several vertices located in different brain regions, and Figure 4b shows the fitted FCS and its growth rate across the cortex. Notably, the most pronounced changes in functional connectivity at the regional level occurred within the first decade of life. We then sought to elucidate how the overall growth patterns varied spatially across the cortex by mapping the primary spatial axis of FCS development. To this end, we used a principal component analysis (PCA) on the zero-centered 50th centiles of the growth curves. The first PC, accounting for 60.4% of the variance, was identified as the dominant axis of regional functional connectivity growth (Fig. 4c). This axis captured a hierarchical spatial transition, starting from primary sensorimotor and visual

cortices and culminating in higher-order association regions, including the angular gyrus, precuneus, temporal, and prefrontal cortices. To better illustrate the spatiotemporal pattern of growth curves throughout the cortex, we segmented the main growth axis into 20 equal bins and averaged the curves for vertices within each bin. A continuous spectrum of curves along the lifespan axis is shown in Fig. 4d.

The cortical landscape of the human brain is organized by a fundamental gradient known as the sensorimotor-association (S-A) axis [65]. This axis spans from primary cortices critical for sensory and motor functions to advanced transmodal regions responsible for complex cognitive and socioemotional tasks. It has been shown to play an important role in shaping neurodevelopmental processes [66-68]. Here, we sought to investigate the extent to which our defined growth axis aligns with the classic S-A axis as formulated by Sydnor et al. [66]. (Fig. 4e). Using a spin-based spatial permutation test [69], we found a significant association between the main growth axis and the S-A axis (r = 0.72, $p_{spin} < 0.0001$) (Fig. 4f). This finding suggests that the spatiotemporal growth of the functional connectome throughout the human lifespan follows the canonical sensorimotor-association organization.

Fig. 4 | Lifespan normative growth patterns of regional functional connectivity strength. a, Normative growth curves of example vertices from different regions. **b**, The fitted 50th centiles (top panel) and their growth rates (bottom panel) for all vertices at representative ages. **c**, The lifespan growth axis of brain functional connectivity, represented by the first principal component from a PCA on regional level FCS curves. **d**, Based on the lifespan principal axis, all vertices across the brain were equally divided into 20 bins. The zero-centered curves of all vertices within each bin were averaged. The first vigintile (depicted in darkest yellow) represents one pole of the axis, while the twentieth vigintile represents the opposite pole (depicted in darkest blue). The patterns of growth curves vary continuously along the axis, with the greatest differences observed between the two poles. **e**, The sensorimotor-association (S-A) axis, as formulated by Sydnor et al. [66], represents a cortical continuum that transitions from primary regions to transmodal areas. **f**, A strong correlation was observed between the lifespan principal growth axis and the S-A axis (r = 0.72, $p_{spin} < 0.0001$) (linear association shown with 95% confidence interval). All brain maps were mapped to the HCP fs_LR_32k surface for visualization. FCS, functional connectivity strength; wk, week; yr, year.

Sex differences in lifespan growth patterns

It is becoming increasingly evident that sex differences exert a significant influence on brain development and aging [70, 71]. In GAMLSS modeling, we included a sex effect as an additional variable to establish lifespan normative growth curves. We characterized the sex-stratified growth curves and interindividual variability curves of the functional connectome (Supplementary Result 3, Supplementary Figs 13 and 14). Specifically, we observed that the global mean of the functional connectome was significantly greater in males than in females (p_{FDR} = 0.0002), thereby confirming and extending conclusions from previous studies [72, 73]. Conversely, the global variance of the connectome was greater in females than in males (p_{FDR} = 0.0009). Furthermore, females showed greater global system segregation ($p_{FDR} = 10^{-24}$) and system-specific segregation in the VIS, VA, FP, and DM networks (all $p_{FDR} < 0.01$), but lower system-specific segregation in the SM and limbic (LIM) networks (all $p_{FDR} < 10^{-32}$) than males. At the regional level, the lateral and medial parietal cortex and lateral prefrontal cortex showed greater FCS in females, whereas the sensorimotor cortex, medial prefrontal cortex, and superior temporal gyrus showed greater FCS in males ($p_{FDR} < 0.05$). These results are compatible with a previous study employing seed-based and independent component analysis (ICA)-based functional connectivity analysis [23]. Additionally, in a recent study, Zhang et al. [74] used a large data set (36,531 participants from the UK Biobank, mean age 69) to report that females had lower functional connectivity in somatosensory/premotor regions and greater functional connectivity in the inferior parietal and posterior cingulate cortex, which aligns with our findings. The detailed statistical values of the sex variable within each normative model are presented in Supplementary Tables 3 and 4. The sex differences in the interindividual variability curves are detailed in the Supplementary Result 3.

Identifying individual heterogeneity in brain disorders using connectome-based normative models

Recent studies have highlighted the potential of normative models to disentangle the inherent heterogeneity in clinical cohorts by enabling statistical inference at the individual level [18, 75-81]. This approach enables the quantification of individual deviations of brain phenotypes from normative expectations, thereby providing unique insights into the typicality or atypicality of individual's brain structure or function. To validate the clinical value of our connectome-based normative models, we selected three representative brain disorders characterized by connectome dysfunction, each manifesting at distinct life stages. These were ASD, which mainly presents in early development; MDD, which mainly presents in adolescence and adulthood; and AD, which mainly presents in older adulthood.

We characterized the individual deviation z-scores (age- and sex-specific) of the functional metrics at the global, system, and regional levels in patients with ASD (N_{ASD} = 414, aged 5-59 years), MDD (N_{MDD} = 622, aged 11-77 years), or AD (N_{AD} = 180, aged 51-80 years), and their matched healthy controls (HCs). The standard protocol for normative modeling [82] emphasizes the importance of incorporating some control samples from the same imaging sites as the patients into the testing set. This approach verifies that the observed case☐control differences are not due to the analysis with controls in the training set and cases in the testing set [78, 82]. This approach also allows for the estimation of site effects within the case☐control datasets. In the present study,

we reconstructed the connectome-based normative models for all three disorders using the same
set of healthy participants. Specifically, we randomly divided the HCs of all case–control
datasets ($N_{HC} = 591$ in ASD datasets, $N_{HC} = 535$ in MDD datasets, and $N_{HC} = 187$ in AD datasets)
in half, stratified by age, sex, and site. The training set ($N = 32,591$), which was used to construct
the normative model, consisted of half of the HCs ($N_{train} = 654$) and all samples from other
datasets ($N = 31,937$). The testing set, comprising remaining half of the HCs ($N_{test} = 659$) and the
patient cases, was used as a completely independent set to determine their deviation scores. This
process was repeated 100 times, generating 100 new normative models and 100 sets of deviation
scores. We observed a high degree of stability of both normative curves and patient's deviation
scores across the 100 repetitions (average $r > 0.95$ and average mean square error [MSE] < 0.2
for all functional metrics, see Supplementary Fig. 15, Supplementary Tables 5 and 6). We then
averaged the 100 sets of deviation scores for patients in each disease group, and then assessed
the extreme deviations ($z > |2.6|$) for each metric. Among the ASD patients, 92% had at least one
metric with an extreme negative deviation, and 32% had at least one metric with an extreme
positive deviation (Supplementary Fig. 16). For MDD patients, the percentages were 89% and
39%, respectively, and for the AD patients, they were 61% and 25%, respectively. Furthermore,
we calculated the proportion of patients with extreme deviations in each metric and found that no
more than 10% of the patients had extreme deviations in any single metric (Fig. 5a,
Supplementary Fig. 16). These results highlight the considerable individual heterogeneity within
each disease group.

Using the k-means clustering algorithm, we identified two subtypes for ASD ($N_{ASD1} = 238$ for
subtype 1, $N_{ASD2} = 176$ for subtype 2), two for MDD ($N_{MDD1} = 375$ for subtype 1, $N_{MDD2} = 247$
for subtype 2), and two for AD ($N_{AD1} = 67$ for subtype 1, $N_{AD1} = 113$ for subtype 2) (Fig. 5b).
For each disorder, different subtypes showed distinct patterns of deviation and case–control
differences in the functional connectome (Fig. 5c-f, Supplementary Figs 17 and 18). Specifically,
ASD subtype 1 showed greater positive deviations in the bilateral ventral prefrontal cortex and
negative deviations primarily in the sensorimotor and insular cortices in comparison with HCs
($p_{FDR} < 0.05$, Fig. 5c-d). In contrast, ASD subtype 2 exhibited greater positive deviations in the
sensorimotor and insular cortices ($p_{FDR} < 0.05$, Fig. 5e-f). For MDD, subtype 1 patients showed
greater positive deviations in the lateral frontal and parietal regions, and insular cortices, and
greater negative deviations in the visual and sensorimotor cortices ($p_{FDR} < 0.05$, Fig. 5c-d). MDD
subtype 2 patients showed greater positive deviations in the visual and sensorimotor cortices, and
greater negative deviations in the lateral and medial prefrontal and parietal regions, and the
insula ($p_{FDR} < 0.05$, Fig. 5e-f). For AD, subtype 1 showed very few positive or negative
deviations (Fig. 5c-d), but subtype 2 showed greater positive deviations in the visual and
sensorimotor cortices, and negative deviations in the lateral and medial parietal regions and the
insula ($p_{FDR} < 0.05$, Fig. 5e-f).

We further investigated the classification performance of each disorder with and without
subtyping, characterized by the mean area under the curves (AUCs) (Fig. 5g, Supplementary Fig.
19). Specifically, the mean AUCs for ASD subtypes 1 and 2 were 0.76 and 0.67, respectively (all
$p_{FDR} < 0.001$, permutation tests), but 0.55 without subtyping ($p_{FDR} < 0.05$, permutation tests).
The mean AUCs for MDD subtypes 1 and 2 were 0.71 and 0.73, respectively (all $p_{FDR} < 0.001$,
permutation tests), but 0.58 without subtyping ($p_{FDR} < 0.05$, permutation tests). The mean AUCs
for AD subtypes 1 and 2 were 0.71 ($p_{FDR} < 0.05$, permutation tests) and 0.82 ($p_{FDR} < 0.001$,
permutation tests), respectively, but 0.71 without subtyping ($p_{FDR} < 0.001$, permutation tests).

Furthermore, we investigated the potential of the connectome deviations to predict clinical scores (Fig. 5h, Supplementary Fig. 20). For ASD, the patterns of connectome-based deviations in

Fig. 5 | Clinical relevance of connectome-based deviation patterns in three brain disorders. a, Percentage of patients with extreme deviations. Subplots from left to right display the percentage of patients with extreme positive and negative deviations in ASD, MDD, and AD. The bar plot shows the percentage in global mean of the connectome (G1), global variance of the connectome (G2), global system segregation (G3), and system-specific segregation. The brain map shows the percentage of regional-level FCS. Orange yellow represents extreme positive deviations, while blue represents extreme negative deviations. **b**, The optimal number of subtypes (left panel) and the similarity matrix of deviation patterns across patients (right panel) for each disorder. **c**, Mean deviation patterns in patients in subtype 1 of each disorder. **d**, Individual deviation scores of patients in subtype 1 were compared to the median of healthy controls (HCs). For each metric, the significance of the median differences between the case group and HCs was assessed using the Mann Whitney U test. P-values were adjusted for multiple comparisons using FDR correction across all possible pairwise tests (p < 0.05). The color bar represents the proportion of tests that passed the significance threshold in 100 comparisons. **e**, Mean deviations pattern in patients in subtype 2 of each disorder. **f**, Individual deviation scores of patients in subtype 2 were compared to the median of HCs. **g**, Disease classification performance based on individual deviation patterns using support vector machine analysis. **h**, Prediction accuracy of clinical scores based on individual deviation patterns using support vector regression analysis. All brain maps were mapped to the HCP fs_LR_32k surface and are shown in the left hemisphere. For whole-cortex visualizations, refer to

patients predicted total Repetitive Restrictive Behavior (RRB) scores for each subtype ($r = 0.22$,
$p_{perm} = 0.04$ for subtype 1; $r = 0.24$, $p_{perm} = 0.03$ for subtype 2). However, the prediction was not
significant without ASD subtyping ($r = 0.04$, $p_{perm} = 0.3$). For MDD, the connectome-based
deviation patterns showed significant predictive accuracy for total Hamilton Depression Rating
Scale (HDRS) score, both with and without subtyping ($r = 0.19$, $p_{perm} < 0.001$ for all patients; $r =$
0.27, $p_{perm} < 0.001$ for subtype 1; $r = 0.18$, $p_{perm} = 0.002$ for subtype 2). For AD, the prediction of
the Mini-Mental State Examination (MMSE) score was significant without subtyping ($r = 0.33$,
$p_{perm} < 0.001$), but only for AD subtype 2 ($r = 0.02$, $p_{perm} = 0.5$ for subtype 1; $r = 0.34$, $p_{perm} =$
0.002 for subtype 2). These results demonstrate the clinical relevance of connectome-based
normative modeling in identifying disease subtypes, as evidenced by enhanced performance in
both disease classification and prediction of symptomatic scores.

Sensitivity analyses

The lifespan growth patterns of functional connectomes were validated at the global, system, and
regional levels using various analysis strategies (for details, see Methods). Each validation
strategy yielded growth patterns that highly matched the main results (Supplementary Tables 7-
12 and Supplementary Fig. 21). (*i*) To validate the potential effects of head motion, the analyses
were reperformed using data from 24,494 participants with a stricter quality control threshold for
head motion (mean framewise displacement (FD) < 0.2 mm) (Supplementary Fig. 22). (*ii*) To
mitigate the impact of uneven sample and site distributions across ages, a balanced sampling
strategy was employed to ensure uniformity in participant and site numbers (N = 6,770,
resampling 1,000 times) (Supplementary Fig. 23). (*iii*) To validate reproducibility of our results,
a split half approach was adopted (Supplementary Fig. 24). (i*v*) To examine the potential effects
of data samples, a bootstrap resampling analysis was performed (1,000 times, Supplementary Fig.
25). (*v*) To examine the potential effects of specific sites, a leave-one-site-out (LOSO) analysis
was conducted (Supplementary Fig. 26). The results of these sensitive analyses were
quantitatively assessed in comparison to the main results (Supplementary Tables 7-12).
Specifically, a series of 80 points at one-year intervals was sampled for each curve, and
Pearson's correlation coefficients were then calculated between the corresponding curves
(Supplementary Table 7). At both global and system levels, all growth curves in the sensitivity
analyses exhibited a high degree of correlations with those shown in the main results ($r = 0.97$-
1.0 for global mean of the connectome; $r = 0.98$-1.0 for global variance of the connectome; $r =$
0.99-1.0 for global system segregation; $r = 0.98$-1.0 for system segregation of VIS, DA, VA, FP,
and DM networks; $r = 0.91$-1.0 for system segregation of SM networks; $r = 0.8$-1.0 for system
segregation of LIM networks, except for $r = 0.51$ of the balanced resampling analysis; all $p_{FDR} <$
10^{-5}). At the regional level, the lifespan growth axes in the sensitivity analyses were highly
spatially associated with that shown in the main results (all $r = 0.94$-1.0, $p_{spin} < 0.0001$). The
similar results of growth rate are shown in Supplementary Table 8. We observed consistent
results when the sampling was obtained with six-month intervals (160 points) and monthly
intervals (1,000 points) (Supplementary Tables 9-12).

Discussion

Using a large multimodal structural and task-free fMRI dataset from 33,250 individuals aged 32
postmenstrual weeks to 80 years, we mapped the growth patterns of the functional connectome
across the human lifespan at the global, system, and regional levels. We charted the multiscale,
nonlinear growth curves of the functional connectome and revealed previously unidentified key
growth milestones. To provide a lifespan characterization of functional brain systems, we created
age-specific atlases spanning 32 postmenstrual weeks to 80 years of age to serve as a
foundational resource for future research. Using three representative disease datasets of ASD,
MDD, and AD, we explored the utility of the connectome-based normative model in capturing
individual heterogeneity, identifying disease biotypes, and performing classification and
prediction analyses within these clinical populations, highlighting their potential to advance our
understanding of neuropsychiatric disorders.

At the global level, we observed continuous nonlinear changes in the global mean and variance
of functional connectivity across the life cycle, peaking in the late fourth and late third decades,
respectively. Similarly, the growth curve of global brain structure shows a pattern of increase
followed by decline, albeit peaking earlier [18]. Taken together, these functional and anatomical
findings suggest that the human brain remains in a state of dynamic adaptation throughout the
lifespan. At the systems level, an intriguing observation is that the DM and FP networks, relative
to other networks, undergo more rapid development of system segregation during infancy,
childhood, and adolescence, peak later, and decline precipitously during aging. The accelerated
early development of these networks can be attributed to their initially less organized functional
architecture in utero [53, 83] and the subsequent need for rapid postnatal development to support the
emergence and development of advanced cognitive functions [8, 84, 85]. Moreover, the increased
susceptibility of these networks to accelerated decline during aging may be exacerbated by their
increased sensitivity to environmental, genetic, and lifestyle factors, as well as neurodegenerative
agents such as amyloid-β and tau [86-89]. At the regional level, our results validate and extend the
replicable findings of Luo and colleagues [32], who, using four independent datasets, observed an
increase in FCS in primary regions and a decrease in higher-order regions from childhood to
adolescence. Furthermore, the life-cycle growth curves of regional FCS are constrained by their
positions along the S-A axis, highlighting the role of the S-A axis as a key organizational
principle that influences cortical development and aging [66].

Emerging evidence increasingly implicates abnormal interregional brain communication and
global network dysfunction as critical factors in the pathogenesis of various neuropsychiatric
disorders [13, 15, 16]. After establishing lifespan growth curves, we focused on characterizing the
degree to which individual functional metrics deviated from established population norms. This
analysis provided preliminary insights into the clinical utility of our connectome-based
normative models. Using age- and sex-normalized metrics, we first elucidated individual
heterogeneity in functional brain deviations at the global, system, and regional levels across three
clinically relevant populations, namely ASD, MDD, and AD. Through subtype analysis based on
individual deviation scores, we validated the potential of the connectome-based normative
models to parse complex intragroup heterogeneity and enhance the prediction of disease
discrimination and clinical symptoms. A biological exploration of the underlying causes of
positive and negative deviations in individual functional brain connectomes would provide
valuable insights into the similarities and differences between disparate clinical disorders [78].
Furthermore, future studies could include more disease cohorts with large sample sizes to allow
transdiagnostic comparisons between disorders. It is important to note that considerable work is

still needed to effectively translate growth charts and their derived heterogeneity metrics into
clinical utility [18, 90, 91]. Therapeutically, the incorporation of individual functional deviations
along with finely stratified subtypes may improve the efficacy of interventions using
connectome-guided transcranial magnetic stimulation [92]. In summary, the integration of the
connectomic framework with normative growth curves provides an unprecedented opportunity to
study brain network dysfunction in clinical populations.

A promising avenue to explore for future research is the interaction between lifespan growth
curves of brain networks under different modalities. This interaction could be investigated by
examining how different structural and functional connectivity metrics coevolve across the
lifespan and whether there are similar or variable temporal key points within these curves. It
would be valuable to determine whether milestones of the structural connectome precede those
of the functional connectome, thereby providing an anatomical scaffold for the dynamic
maturation of functional communication. Furthermore, identifying the critical physiological
factors that shape growth patterns across the lifespan is a complex but essential endeavor. Recent
evidence suggests that population-based life-cycle trajectories of cortical thickness align with
patterns of molecular and cellular organization, with varying degrees of biological explanation at
different life stages [93]. A genome-wide association meta-analysis by Brouwer et al. [94] identified
common genetic variants that influence the growth rates in cortical morphology development or
atrophy across the lifespan. These findings underscore the necessity of a multifaceted approach
encompassing anatomical, genetic, molecular, and metabolic methodologies to elucidate the
complex factors that regulate typical and atypical alterations in the human brain connectome.

A number of challenges warrant further consideration. First, the data used to delineate lifespan
growth patterns in the current study were aggregated from existing neuroimaging datasets, which
are disproportionately derived from European, North American, Asian, and Australian
populations. This geographic bias has also been found in other neuroimaging normative
references or big data studies, such as those involving cortical morphology growth maps [18] and
genome-wide association studies of brain structure across the lifespan [94]. Future research should
include more neuroimaging cohort studies designed to achieve a balanced representation of
diverse ethnic populations [95]. In addition, it is critical to consider the diversity of environmental
factors, such as socioeconomic status, education level, industrialization, and regional culture,
which pose potential challenges to the application of lifespan trajectories. Second, as previously
outlined by Bethlehem et al. [18], we also encountered challenges related to the uneven age
distribution of the neuroimaging sample, particularly with the underrepresentation of the infant
and middle-aged (30-40 years) populations. It is evident that functional changes in the uterus are
dramatic, however, the paucity of available fetal fMRI data limits our understanding of this
critical period. Future research should complement the current models with more neuroimaging
data, especially from the fetal stages. Third, the presence of artifacts and low signal-to-noise
ratios in fMRI images of the orbitofrontal cortex, partly due to head movement and magnetic
field inhomogeneity, represents a significant challenge [96, 97]. The development of advanced
imaging techniques and algorithms will be crucial for addressing this issue. Fourth, adjusting for
multisite effects in retrospective data represents another significant challenge. Studies have
shown that incorporating site variables as random effects in models, rather than the use of
ComBat, is a more effective approach in normative modeling [18, 98, 99]. Therefore, we adopted a
conservative analytical approach by modeling site effects as random effects (for a comparison of
results using different methods, see Supplementary Result 4 and Supplementary Fig. 27). Future

research may benefit from integrating prospective cohort designs, phantom scans, and scans of
traveling subjects. Fifth, due to the ambiguity in interpreting negative functional connectivity, we
focused on positive connectivity in our main results. Nonetheless, we also analyzed the
normative growth patterns of negative connectivity across the lifespan at global, system, and
regional levels (Supplementary Result 5 and Supplementary Fig. 28). Sixth, considering the
methodological challenges of surface-based analyses in integrating cortical and subcortical
structures, we focused on cortical connectomes in our main results. In light of the significance of
subcortical structures, we also presented lifespan growth curves of subcortical connectomes
using volume-based analysis (Supplementary Result 6 and Supplementary Fig. 29). Seventh, the
data used in this study are cross-sectional, which may result in an underestimation of age-related
changes in the functional connectome [100]. Therefore, integrating more densely collected
longitudinal data across all ages is essential to accurately characterize lifespan trajectories.
Finally, it is anticipated that the connectome-based growth charts established here will serve as a
dynamic resource. As more high-quality, multimodal connectome datasets become available, the
lifespan normative growth model will be updated accordingly.

Methods

Datasets and participants

To delineate the normative growth of the functional connectome in the human brain, we
aggregated the available multisite neuroimaging datasets, each containing both 3T structural and
task-free fMRI data. For participants with multiple test-retest scans, only the first session was
included. The total number of imaging scans collected was 46,178 with 44,030 participants
ranging in age from 32 postmenstrual weeks to 80 years. These scans were obtained from 172
sites in 28 datasets. Participant demographics and imaging scan parameters for each site were
presented in Supplementary Table 1 and 2, respectively. Written informed consent was obtained
from participants or their legal guardians, and the recruitment procedures were approved by the
local ethics committees for each dataset.

Image quality control process

The implementation of a rigorous and standardized quality control procedure is essential to
ensure the authenticity of neuroimaging data, thereby enhancing the credibility of growth curves.
Previous research has shown that inadequate quality control of MRI scans can diminish the
benefits of large sample sizes in detecting meaningful associations [101]. In this study, we
employed a comprehensive four-step data quality control framework that combined automated
assessment approaches and expert manual review to assess both structural and functional images
across all 46,178 imaging scans from 44,030 participants (Supplementary Figs 1 and 2). This
rigorous framework effectively identified imaging artifacts or errors, thereby ensuring the
accuracy and reliability of our neuroimaging data.

Step 1: Quality control of the raw images. First, we performed a preliminary quality control to
filter out low-quality scans with problematic acquisitions. For several publicly available datasets
(dHCP, HCP-Development, HCP-Aging, HCP-Young Adult, and ABCD) that provide
information on image quality, we performed initial quality control according to their
recommended inclusion criteria. For the BCP dataset, each scan was visually reviewed by two
neuroradiologists experienced in pediatric MRI. For the other datasets, we conducted automated

quality assessment using the MRI Quality Control (MRIQC) tool [102], which extracted non-
reference quality metrics for each structural (T1w and T2w) and fMRI image. In each dataset,
structural images were excluded if they were marked as outliers (more than 1.5 times the
interquartile range (IQR) in the adverse direction) in at least three of the following quality
metrics: entropy-focus criterion (EFC), foreground-background energy ratio (FBER), coefficient
of joint variation (CJV), contrast-to-noise ratio (CNR), signal-to-noise ratio (SNR), and
Dietrich's SNR (SNRd). Similarly, functional images were excluded if they were marked as
outliers in three or more of the following quality metrics: AFNI's outlier ratio (AOR), AFNI's
quality index (AQI), DVARS_std, DVARS_vstd, SNR, and temporal signal-to-noise ratio (tSNR).
This step resulted in the exclusion of 838 structural and 963 functional images.

Step 2: Determination of whether to pass the entire processing pipeline. Following the initial
quality control step, the images were submitted to the pre- and post-processing pipelines. A
detailed description of the latter is provided in the "**Data processing pipeline**" section. Any scan
that could not pass the entire data processing pipeline was excluded, resulting in the removal of
2,910 structural and 2,969 functional images.

Step 3: Surface quality control and head motion control. For structural images, the Euler
number was employed to assess the quality of the reconstructed cortical surface. The Euler
number is a mathematical concept that summarizes the topological complexity of a surface and,
can be calculated as 2-2n, where n represents the number of defects such as holes or handles. A
high Euler number represents a surface with fewer defects, indicating high-quality cortical
surface reconstruction. The Euler number is a reliable and quantitative measure that can be used
to identify images unsuitable for analysis [18, 101, 103]. Similarly, the images with an Euler number
magnitude less than 1.5 times the IQR in the adverse direction from the study-specific
distribution (Q1–1.5*IQR) were identified as outliers and excluded. For functional images, scans
with large head motion (mean FD > 0.5 mm, or frames with FD over 0.5 mm > 20%) were
excluded, along with scans with fewer than 100 final time points or a ratio of final time points to
original time points < 90%. In total, 2,117 structural images and 3,573 functional images were
excluded.

Step 4: Visual double-check. During the initial three QC steps using automated assessment
approaches, 5,865 scans with structural imaging problems and 7,505 scans with functional
imaging problems were excluded. To further ensure the quality of the remaining scans, we
performed a detailed and comprehensive visual check QC. (1) A visual QC team was assembled,
comprising of four anatomically trained experts: Q.W., Q.Y., C.P., and L.S.. For each participant
who had passed the automated QC steps, three 2D pictures were generated (one for structural
MRI images and two for functional MRI images). (2) Based on these images, L.S. conducted the
initial round of visual QC on both structural and functional data for all participants, recording the
IDs of those with quality errors. (3) The pictures were then distributed evenly among Q.W., Q.Y.,
and C.P. for a secondary evaluation. The IDs of the participants exhibiting quality defects were
documented. The final list of participants who were excluded was determined based on the
combination of these records. Throughout the process, the QC team engaged in in-depth
discussions to ensure that the exclusion criteria were consistently applied across members. The
exclusion criteria were as follows: The T1-weighted structural images were primarily evaluated
for artifacts and quality of cortical segmentation, reconstruction, and registration. For
participants with T2-weighted images, those with abnormal myelination distribution (as

measured by the T1/T2 ratio) were also excluded. Functional images were assessed for brain
coverage, functional-to-structural and functional-to-standard space registration quality, and
volume-to-surface mapping quality. Participants were excluded if any of these issues were
present. A comprehensive tutorial on visual QC procedures is available at
https://github.com/sunlianglong/BrainChart-FC-Lifespan/blob/main/QC/README.md. In this
step, 651 structural images and 1,153 functional images were excluded. Finally, only scans that
successfully passed QC for both functional and structural images were retained.

The Application of the rigorous criteria outlined above resulted in the exclusion of 10,231 scans
in 9,564 participants. The final sample included 33,250 healthy participants (33,250 cross-
sectional scans and 1,481 longitudinal scans) and 1,216 patients (1,216 cross-sectional scans;
414 patients with ASD, 622 patients with MDD, and 180 patients with AD) with high-quality
functional and structural images.

Data processing pipeline

(i) Structural data preprocessing. Despite our efforts to employ a unified structural
preprocessing pipeline across all datasets to mitigate the impact of disparate methodologies, the
substantial variations in the structure and function of the human brain across the lifespan present
a significant challenge. This was particularly evident in the perinatal and infant periods, where
the anatomical characteristics differ markedly from those of adults. For example, in six-month-
old infants, the contrast between gray and white matter is extremely subtle, and at approximately
six months of age, there is a contrast inversion between gray and white matter. These factors
greatly complicate the segmentation of brain tissue during this period [104, 105]. In the absence of a
preprocessing pipeline suitable for all stages of life, it is necessary to find appropriate methods
for early developmental datasets while ensuring the uniformity of the pipelines in other datasets.

For individuals aged two years and older, we utilized the publicly available, containerized HCP
structural preprocessing pipelines (v4.4.0-rc-MOD-e7a6af9) [106], which have been standardized
through the QuNex platform (v0.93.2) [107]. Briefly, this pipeline consists of three stages: (1) The
PreFreeSurfer stage focused on the normalization of anatomical MRI data and involved a
sequence of preprocessing steps that included brain extraction, denoising, and bias field
correction on anatomical T1 weighted (T1w) and T2 weighted (T2w) MRI data (if T2w data
were available). (2) The FreeSurfer stage aimed to create cortical surfaces from the normalized
anatomical data, including anatomical segmentation; the construction of pial, white, and mid-
thickness surfaces; and surface registration to the standard atlas. (3) The PostFreeSurfer stage
converted the outputs from the previous steps into the HCP format (CIFTI). The volumes were
transformed to the standard MNI space using nonlinear registration, while the surfaces were
mapped to the standard fs_LR_32k space using spherical registration and surface downsampling.
To mitigate the computational burden of processing the large ABCD dataset, we chose to use the
community-shared, preprocessed data released through the ABCD-BIDS Community Collection
[108] (ABCD collection 3165; https://github.com/ABCD-STUDY/nda-abcd-collection-3165). The
multimodal neuroimaging data were preprocessed using the ABCD-HCP pipeline, a variant of
the HCP pipeline adapted to better suit the ABCD dataset. Modifications to the ABCD-HCP
structural pipeline include volume registration algorithms and bias field correction methods.
Further details of these modifications can be found in the online document
(https://collection3165.readthedocs.io/en/stable/pipelines/).

For participants in the postmenstrual age range of 32 to 44 weeks from the dHCP study, we
applied the officially recommended dHCP structural pipelines [109], which have been specifically
designed to account for the substantial differences between neonatal and adult MRI data. This
HCP-style pipeline included the following steps: (1) bias correction and brain extraction, which
were performed on the motion-corrected, reconstructed T2w images; (2) tissue segmentation; (3)
cortical reconstruction of the white matter surface; (4) surface topology correction; (5)
generation of pial and mid-thickness surfaces; (6) generation of inflated surfaces derived from
the white matter surface through an expansion-based smoothing process; and (7) projection of
the inflated surface onto a sphere for surface registration. Furthermore, we used the officially
recommended iBEAT V2.0 pipelines [110] for participants aged from 0-2 years (all from the BCP
study). This pipeline, which is optimized for preprocessing early-age neuroimaging data based
on advanced algorithms, has shown superior performance in tissue segmentation and cortical
reconstruction for BCP datasets compared to alternative approaches [110]. The main steps of this
pipeline included (1) inhomogeneity correction of T1w/T2w images; (2) skull stripping and
cerebellum removal (for participants with incomplete cerebellum removal, frame-by-frame
manual corrections were performed); (3) tissue segmentation; (4) cortical surface reconstruction;
(5) topological correction of the white matter surface; and (6) final reconstruction of the inner
and outer cortical surfaces. To ensure consistency in data preprocessing, we employed the iBEAT
pipeline for structural image preprocessing of participants aged 2-6 years (53 scans, representing
13% of the total BCP cohort) from the BCP site.

The individual cortical surface obtained from the dHCP and iBEAT V2.0 structural pipelines
were aligned with the adult fs_LR_32k standard space using a three-step registration method
(Supplementary Fig. 3). For participants aged 32 to 44 postmenstrual weeks, the following steps
were implemented. (1) Individual surfaces were registered to their respective postmenstrual week
templates [111]. (2) Templates for 32-39 postmenstrual weeks and 41-44 postmenstrual weeks were
registered to the 40-week template. (3) The 40-week template was then registered to the
fs_LR_32k surface template. For participants aged 1-24 months, the following steps were
undertaken. (1) Individual surfaces were registered to their corresponding monthly age templates
[112]. (2) All monthly templates were registered to the 12-month template. (3) The 12-month
template was then registered to the fs_LR_32k surface template. A supplementary analysis was
conducted to validate the normative growth pattern of the global functional connectome, which
involved avoiding cross-age surface registration (Supplementary Result 7 and Supplementary Fig.
30).

(ii) Functional data preprocessing in volumetric space. For individuals aged two years and
older, the HCP functional preprocessing pipelines were employed [106]. The fMRIVolume stage
consisted of the following steps. (1) Slice timing correction: This step was applied to single-band
acquisitions, as multi-band acquisitions did not require slice timing correction. (2) Motion
correction: EPI images were aligned to the single-band reference image using 6 DOF FLIRT
registration. In cases where the single-band imaging data were not available, the first frame of
the fMRI data was used as the reference. The motion parameters, including translations, rotations,
and their derivatives were recorded. Additionally, the demeaned and linearly detrended
parameter was provided for nuisance regression analysis. (3) EPI distortion correction:
Geometric distortion correction was conducted using either the opposite-phase encoded
spin□echo images (when LR-RL or AP-PA encoded acquisitions were available) or the regular
(gradient-echo) fieldmap images (when fieldmap acquisitions were available). If neither image

was available, this step was skipped. (4) Anatomical registration: The fMRI images were
registered to the T1w image using 6 DOF FLIRT with boundary-based registration (BBR). (5)
Intensity normalization: The fMRI data, masked by the final brain mask generated by the
PostFreeSurfer structural pipeline, were normalized to a 4D whole-brain average of 10,000.

For participants in the postmenstrual age range of 32 to 44 weeks from the dHCP study, we
applied the dHCP functional pipelines [113]. Building on the foundation of the HCP pipeline and
the FSL FEAT pipeline, this pipeline was tailored to address the unique challenges associated
with neonatal fMRI data. The key components of the pipeline included the following steps. (1)
Fieldmap preprocessing, which included estimation of the susceptibility distortion field based on
the opposite-phase encoded spin☐echo images and subsequent alignment of this field to the
functional data. (2) Registration, which included BBR of the fieldmap magnitude to the T2w
image, BBR of the single-band reference image to the T2w image with incorporation of field
map-based distortion correction, and 6 DOF FLIRT registration of the first volume of the
functional multiband EPI to the single-band reference image. (3) Susceptibility and motion
correction, which included slice-to-volume motion correction, motion-by-susceptibility
distortion correction, and estimation of motion nuisance regressors. These steps resulted in
distortion-corrected and motion-corrected 4D multiband EPI images in the T2w native
volumetric space. For participants from the BCP cohort, we implemented several steps to obtain
preprocessed volumetric fMRI data. (1) Motion correction: functional images were aligned to the
single-band reference image using 6 DOF FLIRT registration. In the absence of a single-band
reference, the mean functional images (with all frames aligned to the first frame) were employed
as the reference. (2) Distortion correction: we performed distortion correction based on the
opposite-phase encoding (AP-PA) spin☐echo images. This step was only performed for
participants with available images. (3) EPI to anatomical registration: the reference image was
aligned to the anatomical image (T1w or T2w) using 6 DOF FLIRT registration.

(iii) Functional data preprocessing in surface space. In the fMRISurface stage of the HCP
functional pipeline, the goal was to project the volume time series onto the standard CIFTI
grayordinates space. For the data from the dHCP and BCP cohorts, we followed the same steps
of the HCP preprocessing pipeline to achieve an accurate representation of cortical BOLD
signals on the surface. Specifically, the fMRI volumetric data in the cortical cortex were
separated into left and right hemispheres and mapped onto each participant's mid-thickness
surfaces using a partial-volume weighted, ribbon-constrained volume-to-surface mapping
algorithm [106]. Subsequently, the time courses were then transferred from the individual's native
space to the fs_LR_32k standard space using each participant's surface registration
transformations from the structural preprocessing stage.

(iv) Functional data postprocessing. For the ABCD dataset, the ABCD-HCP functional pipeline
used DCANBOLDProcessing software
(https://collection3165.readthedocs.io/en/stable/pipelines/) to reduce spurious variance that is
unlikely to reflect neural activity. For other datasets, the preprocessed fMRI data were post-
processed using SPM12 (v6470) and GRETNA (v2.0.0) with a uniform pipeline. Specifically, the
following steps were initially conducted on the time series for each vertex in fs_LR_32k space
(59,412 vertices in total): linear trend removal, regression of nuisance signals (24 head motion
parameters, white matter signal, cerebrospinal fluid signal, and global signal), and temporal
bandpass filtering (0.01–0.08 Hz). To mitigate the effects of head motion, the motion censoring

was further implemented. This process involved discarding volumes with an FD greater than 0.5
mm and adjacent volumes (one before and two after). To maintain the temporal continuity of the
fMRI time series, we subsequently filled these censored frames using a linear interpolation.
Participants with more than 20% of frames exceeding the 0.5 mm FD threshold were excluded
from our study. Surface-based smoothing was then applied using a 6-mm full-width at half-
maximum (FWHM) kernel. Finally, the data were resampled to a mesh of 2,562 vertices
(corresponding to the fsaverage4 standard space) for each hemisphere using the HCP Workbench
metric-resample command. The removal of the medial wall resulted in a combined total of 4,609
vertices exhibiting BOLD signals on both the left and right hemisphere surfaces.

Construction of the age-specific and individualized functional atlases across the lifespan

(i) Construction of population-level age-specific atlases. To improve the precise mapping of
individual-specific functional networks across the lifespan, we first developed a Gaussian-
weighted iterative age-specific group atlas (GIAGA) generation approach (Supplementary Fig.
6a) to create a set of age-specific population-level functional atlases (Fig. 2a, Supplementary
Figs 7 and 8). Given the dramatic functional changes that occur during early development [57], we
prioritized the generation of finer age-specific atlases for these stages compared to the later life
stages. To this end, we divided all individual scans into 26 different age subgroups, ranging from
32 postmenstrual weeks to 80 years of age. Each age group consisted of cross-sectional data only.
Then, we constructed an age-specific functional atlas for each subgroup. A total of 9 atlases were
constructed for the perinatal to early infant period, including 4 for perinatal development (34-
week, 36-week, 38-week, and 40-week (0-year) atlases) and 5 for the first year of life (1-month,
3-month, 6-month, 9-month, and 12-month atlases). 2 atlases were developed for toddlers (18-
month and 24-month atlases), while 9 atlases were created for childhood and adolescence (4-year,
6-year, 8-year, 10-year, 12-year, 14-year, 16-year, 18-year, and 20-year atlases). Finally, 6 atlases
were generated for adults and the elderly (30-year, 40-year, 50-year, 60-year, 70-year, and 80-
year atlases). A total of 300 participants were randomly selected for each age subgroup. In the
event that the available sample size was less than 300, all participants who passed the imaging
quality control were included. Further details on the age range, number of participants, and sex
ratio for each atlas can be found in Supplementary Table 13.

In recent studies of brain functional organization, Yeo's 7- and 17-network atlases [58] have been
widely used to map cortical functional systems [114]. By including hand sensorimotor areas based
on activations in a hand motor task [115], Wang and colleagues extended this classical functional
parcellation, resulting in an 18-network atlas [60]. In line with previous studies [47, 61, 62], we utilized
this updated classic 18-network map as the initial atlas for the construction of age-specific group
atlases. The detailed construction process for a given age subgroup (e.g., 17-19 years) was as
follows. First, to enrich the dataset for this age subgroup, we included the latter half of the
participants from the previous subgroup (15-17 years) and the earlier half of the participants
from the subsequent subgroup (19-21 years). We then used the individualized parcellation
iteration algorithm proposed by Wang and colleagues [60] to map the 18-network atlas to each
participant, generating the initial individualized functional parcellations (step 1 in Supplementary
Fig. 6a). We then proposed the GIAGA approach. Around the core age (i.e., 18 years) of this
given group, we generated a Gaussian probability distribution $N(\mu, \sigma^2)$ with mean $\mu = 0$ and
standard deviation $\sigma = 1$ and assigned weights to each participant based on their age position in
this Gaussian distribution. The weight quantified the participant's contribution to the population-

level atlas construction, with closer to the core age resulting in a greater contribution. For each
vertex, we calculated the cross-participant cumulative probability of belonging to each network
and assigned vertex labels to the network with the highest cumulative probability, resulting in an
initial age-specific population-level atlas (step 2 in Supplementary Fig. 6a). Finally, steps 1 and 2
were iteratively repeated until the overlap between the current and previous atlases exceeded 95%
or the total number of iterations exceeded 10, indicating convergence (step 3 in Supplementary
Fig. 6a).

(ii) Individualized atlas construction. For a given participant, we used the same iterative
parcellation method described above to generate an individualized functional parcellation based
on the corresponding population-level atlas specific to the participant's subgroup (Supplementary
Fig. 6b, adapted from [60]). Briefly, the influence of the population-level atlas on the individual
brain varied across participants and across brain regions; therefore, this method made flexible
modifications during the construction of the individualized atlas based on the distribution of
intersubject variability in the functional connectome and the tSNR of the functional BOLD
signals [60]. Over the iterations, the weight of population-based information was progressively
reduced, allowing the final individualized map to be completely driven by the individual-level
BOLD data. More information on this iterative functional parcellation approach can be found in
the study by Wang and colleagues [60].

Notably, given the potential variance of different interindividual variability patterns and tSNR
distributions across different age subgroups, we generated an interindividual variability map and
a tSNR map for each age subgroup. This was done to improve the accuracy of both the
individual and population-level atlases. We divided the time series data of each participant
within each age subgroup into two halves. For each half, we computed a vertex-by-vertex
functional connectome matrix. This allowed us to obtain the intersubject variability and the
intrasubject variability within the subgroup. By regressing the intrasubject variability from the
intersubject variability, we obtained a "purified" measure of intersubject variability in the
functional connectome [116, 117].

(iii) Construction of the reference atlas used for comparison. To mitigate the potential bias
introduced by specifying a reference atlas for 'mature age', we adopted a data-driven approach to
construct the reference atlas. Atlas similarity was assessed using the overlap index, which
quantifies the proportion of vertices with matching labels between two atlases. For instance, if
two atlases have 4,000 vertices with identical labels out of a total of 4,609 vertices, the overlap
index would be 4,000/4,609 = 86.8%. We computed the overlap index between each pair of the
26 atlases, resulting in a 26×26 similarity matrix. Hierarchical clustering was applied to this
matrix, as shown in Supplementary Fig. 10a. We selected a highly congruent cluster of atlases,
including the 18-, 20-, 30-, 40-, 50-, 60-, and 70-year atlases. For each vertex, we assigned the
label as the system that had the highest probability of occurrence across these selected atlases,
thereby generating the final reference atlas (Supplementary Fig. 10b).

(iv) Homogeneity of both the age-specific and personalized functional atlases. We evaluated
the functional homogeneity of three parcellation atlases at specific age intervals: the adult-based
group atlas established by Yeo et al. [58], the age-specific group atlas, and the individual-specific
atlas (Supplementary Fig. 9). For each age interval, we performed one-way repeated measures
analysis of variance (RANOVA) followed by post hoc multiple comparisons tests to determine

whether the homogeneity of the individualized atlas was significantly greater than that of the
age-specific group atlas and whether the homogeneity of the age-specific group atlas was
significantly greater than that of the adult-based group atlas.

The homogeneity of a system was assessed by calculating the average similarity between every
pair of vertices assigned to it. The commonly used metric is within-system homogeneity, which
is calculated as the average of Pearson's correlation coefficients between the time series of all
vertex pairs within each system, serving as a measure of internal consistency [48, 49]. To summarize
within-system homogeneity for comparisons across atlases, we averaged the homogeneity values
across systems [49]. For validation, we employed another commonly used metric, the functional
profile homogeneity, which defines system similarity as Pearson's correlation coefficient
between the "connectivity profiles" of vertices within a system [118, 119]. The connectivity profile of
a vertex is represented by the connections between this vertex with all other cortical vertices. The
global average functional profile homogeneity value was derived by averaging the homogeneity
values across all systems [119]. The RANOVA revealed significant differences in the global average
of functional homogeneity across different atlases for any given age interval (all $F > 267$, $p < 10^{-25}$,
Supplementary Fig. 9). Post hoc analysis revealed significant differences in functional
homogeneity between every pair of atlases (all $p < 10^{-8}$, individual-specific atlas > age-specific
group atlas > adult-based group atlas, Supplementary Fig. 9), regardless of the age groups
considered.

Individualized metrics of the functional connectome at global, system, and regional levels

For each pair of vertices among the 4,609 vertices in the fsaverage4 space, we computed the
Pearson's correlation coefficient to characterize the vertex-by-vertex functional connectivity,
resulting in a 4,609×4,609 functional connectome matrix for each participant. All negative
functional connectivity strengths were set to zero. For each participant, the global mean of
functional connectome was defined as the mean of all 4,609×4,609 connections (edges), and the
global variance of functional connectome was defined as the standard deviation of all
4,609×4,609 connections. For validation, we also calculated the global mean of the functional
connectome by averaging only the positive-weight edges, which yielded similar lifespan growth
patterns (Supplementary Result 8 and Supplementary Fig. 31). At a regional level, the FCS of a
given vertex was quantified as the average of the connections with all other vertices.

For a given brain system, an individual's within-system functional connectivity FC_w was defined
as the average connection strength among all vertices within that personalized system.
Conversely, the individual's between-system connectivity FC_b was represented by the average
strength of connections between this system and all other systems. System segregation [64] was
determined by calculating the difference between FC_w and FC_b, normalized by FC_w, as
described in the following formula:

932
$$\text{System segregation} = \frac{FC_w - FC_b}{FC_w}$$

Similarly, global system segregation was defined as the difference between global mean within-
system connectivity and global mean between-system connectivity, normalized by global mean
within-system connectivity.

The degree of global similarity between an individualized atlas and the reference atlas was
quantified by the overlap index. This was defined as the number of vertices with the same label
in the two atlases divided by the total number of vertices in both atlases. If there were 4,609
vertices with the same label in two atlases, the overlap index was 4,609/4,609 = 1.0. The degree
of similarity between an individualized system and its corresponding system in the reference
atlas was quantified using the Dice coefficient.

Modeling normative growth curves across the lifespan

To estimate the normative growth patterns for various metrics of the functional connectome in
healthy individuals combined across cohorts, we applied the GAMLSS [41, 42] to the cross-
sectional data using the *gamlss* package (version 5.4-3) in R 4.2.0. The GAMLSS procedure
were established with two steps: identification of the optimal data distribution, followed by
determination of the best-fitting parameters for each functional connectome metric. Using these
metric-specific GAMLSS models, we obtained nonlinear normative growth curves and their first
derivatives. Furthermore, the sex-stratified growth patterns were revealed. The goodness of fit of
the model was confirmed by out-of-sample metrics and visualized by traditional Q☐Q
(quantile☐quantile) plots and detrended transformed Owen's plots. The robustness of the
lifespan growth curves was assessed through bootstrap resampling analysis, leave-one-study-out
analysis, balanced resampling analysis, and split-half replication analysis.

(i) Model data distributions. While the World Health Organization provides guidelines for
modeling anthropometric growth charts (such as head circumference, height, and weight) using
the Box☐Cox t-distribution as a starting point [41], we recognized that the growth curves of brain
neuroimaging metrics do not necessarily follow the same underlying distributions. For instance,
Bethlehem et al. reported that the generalized gamma distribution provided the best fit for brain
tissue volumes [18]. Therefore, we evaluated all continuous distribution families (n=51) for model
fitting. To identify the optimal distribution, we fitted GAMLSS with different distributions to
four representative global functional metrics (global mean of the connectome, global variance of
the connectome, global atlas similarity, and global system segregation) and assessed model
convergence. The Bayesian information criterion (BIC) was used to evaluate the fits of the
converged models. A lower BIC value indicates a superior fit. As shown in Supplementary Fig.
32, the Johnson's Su (JSU) distribution consistently demonstrated the optimal fit performance
across all the evaluated models.

(ii) GAMLSS framework. We performed the GAMLSS procedure with the functional
connectome metric as the dependent variable, age as a smoothing term (using the B-spline basis
function), sex and in-scanner head motion (HM) as other fixed effects, and scanner sites as
random effects. The JSU distribution, which has four parameters: median (μ), coefficient of
variation (σ), skewness (ν), and kurtosis (τ), was chosen to fit the data distribution. Each
functional connectome metric, denoted by y, was modeled as:

$$y = JSU(\mu, \sigma, \nu, \tau),$$

$$\mu = f_\mu(age) + \beta_\mu^1(sex) + \beta_\mu^2(HM) + z_\mu(site),$$

$$\sigma = f_\sigma(age) + \beta_\sigma(sex),$$

$$\nu = \beta_\nu,$$

$$\tau = \beta_\tau.$$

Given the growth complexity across the lifespan, we sought to capture the underlying age-related
trends by exploring a range of model specifications. We fitted three GAMLSS models with
different degrees of freedom (df = 3-5) for the B-spline basis functions in the location (μ)
parameters, and set default degrees of freedom (df = 3) for the B-spline basis functions in the
scale (σ) parameters. Following the practice of previous studies [18, 80], only an intercept term was
included for the ν or τ parameter. For model estimation, we used the default convergence
criterion of log-likelihood = 0.001 between model iterations and set the maximum number of
iteration cycles as 200. Finally, the optimal model of a given functional metric was selected
based on the lowest BIC value among all converging models. In our study, we did not observe
instances of nonconvergence in the GAMLSS models for any metric, including those used in
sensitivity analyses.

(iii) Goodness of fit of the normative model. To assess the quality of the model fits, we
employed a training-test split strategy, which enabled us to recognize the importance of out-of-
sample metrics. The dataset was randomly divided into two halves, with one half being used for
training (N = 16,663) and the other for testing (N = 16,587). The stratification by site was
applied to both halves. Subsequently, the GAMLSS model was refitted using the training set and
the model's goodness of fit was evaluated using the testing set. This procedure was repeated by
interchanging the roles of the training and testing sets.

The model's goodness of fit for the central tendency was assessed using R-squared (R^2). The
calibration of the centiles was evaluated using quantile randomized residuals (also known as
randomized z-scores) [120]. If the modeled distribution closely aligns with the observed distribution,
the randomized z-scores should follow a normal distribution, regardless of the shape of the
modeled distribution [121, 122]. We used the Shapiro–Wilk test to determine the normality of the
distribution of the randomized z-scores, where a *W* value close to 1 indicated good normality.
Additionally, we examined the higher-order moments (skewness and kurtosis) of the randomized
residuals to gain deeper insights into the goodness of fit of the normative model [121]. Skewness
values close to 0 indicate symmetrically distributed residuals, showing no left or right bias, and
kurtosis values close to 0 indicate a desirable light-tailed distribution. The results demonstrated
that nearly all models had skewness and kurtosis values close to 0, with the Shapiro–Wilk *W*
values consistently above 0.99 (Supplementary Figs 33 and 34, Supplementary Table 14). The R^2
values for the global connectome mean, global connectome variance, global atlas similarity, and
global system segregation were 0.49, 0.48, 0.56, and 0.36, respectively. The R^2 values for the
system segregation of each network ranged from 0.14 to 0.32.

Furthermore, the normalized quantile residuals of the normative model were visually assessed
using two diagnostic methods. First, we inspected the plots related to the residuals. As shown in
Supplementary Fig. 35, the residuals against the fitted values of μ and the index were uniformly
distributed around the horizontal line at 0. In addition, the kernel density estimation of the
residuals showed an approximately normal distribution, and the normal quantile–quantile (Q–Q)
plots showed an approximately linear trend with an intercept of 0 and a slope of 1. Second, we
used the detrended transformed Owen's plots of the fitted normalized quantile residuals to
evaluate the performance of the models. This function uses Owen's method to construct a
nonparametric confidence interval for the true distribution. As shown in the resulting plots
(Supplementary Fig. 36), the zero horizontal line fell within the confidence interval, suggesting

that the residuals followed a normal distribution.

(iv) Sex differences across the lifespan. In the GAMLSS model, sex was included as a fixed
effect to evaluate its impact on the lifespan curves of the functional connectome. We obtained the
μ and σ coefficients, as well as their standard errors, *T*-values, and *P*-values, for the sex variable
using the *summary* function in R as detailed in Supplementary Tables 3 and 4. The estimated μ
and σ coefficients represent the adjusted mean and variance effect of sex on the functional
phenotype, considering control variables such as age, head motion (mean FD), and the random
effects of scanner site. The *T*-value, calculated as the coefficient divided by its standard error,
serves as a statistic to test the null hypothesis that the coefficient is equal to zero (no effect).

Sensitivity analysis of the connectome-based normative models

The lifespan normative growth patterns were validated at the global, system, and regional levels
using various analysis strategies. These analyses addressed key methodological concerns
including head motion, the impact of uneven sample and site distributions across ages,
replication using independent samples, model stability, and potential effects of specific site. At
the global and system level, we quantitatively assessed the similarity between these validated
growth patterns and the main results by sampling 80 points at one-year intervals for each growth
curve and growth rate and calculating Pearson's correlation coefficient between the
corresponding curves. The sampling was also conducted at six-month intervals (160 points) and
monthly intervals (1,000 points). At the regional level, we calculated the spatial association
between the lifespan growth axis in the sensitivity analyses and that shown in the main results.

(i) Analysis with stricter head motion threshold (mean FD threshold < 0.2 mm). Previous
research has indicated that head motion can significantly impact the quality of brain imaging data
[123-125]. To ensure that our findings were not influenced by the potential effects of head motion, we
implemented a stricter quality control threshold, excluding participants with a mean FD
exceeding 0.2 mm, and replicated all normative model analyses. Specifically, after excluding
8,756 participants from the initial cohort of 33,250 participants with a 0.5 mm mean FD
threshold, we used data from 24,494 participants to validate the lifespan growth curves of the
functional brain connectome at the global, system, and regional levels (Supplementary Fig. 22).

(ii) Balanced resampling analysis. To address potential biases arising from uneven sample and
site distributions across age groups, a balanced sampling strategy was performed (Supplementary
Fig. 23). This approach ensured equitable participant and site counts across various age groups
through random sampling. Specifically, we divided the entire age range across the lifespan into
sixteen age groups (each spanning five years) and then calculated the number of participants and
sites for each age group. Besides the age groups under 5 years of age or over 70, the (35, 40] age
group had the fewest participants at 464 and the (40, 45] age group contained the fewest sites at
23 (Supplementary Fig. 23-Ia). Thus, we selected all participants from the 23 most populated
sites within the (35,40] age group, comprising 457 participants. For other age groups, a random
sampling strategy was implemented to include 457 participants from the 23 most populated sites.
The resulting distribution of participants and sites across age groups after resampling is shown in
Supplementary Fig. 23-Ib.

For global and system metrics, sampling was repeated 1,000 times using the above procedure on
a pool of 33,250 participants. For each sampling, we randomly selected 6,770 participants and

re-performed the GAMLSS models, resulting in 1,000 sets of growth curves for each metric. We
then calculated the 95% CI for these curves, the 95% CI for the peak of the median (50th) centile,
and the correlations between the 1,000 median centile lines and the median centile line derived
from the original cohort of 33,250 participants. For regional metrics (i.e., FCS), we selected a
random resample and recalculated all results, including the normative growth curves and growth
rate of the regional FCS, the lifespan growth axis, and the association between the lifespan
growth axis and the S-A axis.

(iii) Split-half replication analysis. To assess model replicability in independent datasets, a split-
half strategy was conducted (Supplementary Fig. 24). Participants were randomly divided into
two subgroups, each comprising 50% of the participants ($N_{Subgroup1} = 16,663$, $N_{Subgroup2} = 16,587$),
with stratification by site. The lifespan normative growth patterns were independently evaluated
using Subgroup 1 and Subgroup 2.

(iv) Bootstrap resampling analysis. To assess the robustness of the lifespan growth curves and
obtain their confidence interval, a bootstrap resampling analysis was performed (Supplementary
Fig. 25). This involved the execution of 1,000 bootstrap repetitions using replacement sampling.
To ensure that the bootstrap replicates preserved the age and sex proportionality of the original
studies, the lifespan (from 32 weeks to 80 years) was segmented into 10 equal intervals and
stratified sampling was conducted based on both age and sex. For each functional metric, 1,000
growth curves were fitted and 95% CIs were computed for both the median (50th) centile curve
and the inflection points. The 95% CI were calculated based on the mean and standard deviation
of the growth curves and growth rates across all repetitions.

(v) Leave-one-study-out (LOSO) analysis. To ascertain whether the lifespan growth curves were
influenced by specific sites, the LOSO analyses were implemented (Supplementary Fig. 26). In
each instance, the samples were removed from one site at a time, the GAMLSS models were
refitted and the parameters and growth curves were estimated. We initially compared the curves
obtained after excluding the largest site (Site 1 from the UK Biobank dataset, 12,877 participants)
with those fitted using the entire dataset (N = 33,250). This reveals that both the growth curves
and growth rates were almost identical. The mean and standard deviation across all repetitions
were used to calculate the LOSO 95% CIs for both the normative growth curves and growth rates.
The narrow CI indicated that our models were robust when data from any single site were
removed.

Clinical relevance of connectome-based normative models in brain disorders

To ascertain the clinical relevance of the established lifespan connectome models, the present
study included quality-controlled structural and functional MRI data from three brain disorders.
All procedures of quality control, image preprocessing, and network analysis were identical to
those used for connectome-based normative modeling. The final analyses comprised data from
591 HCs and 414 patients with ASD from the ABIDE dataset (13 sites), 535 HCs and 622
patients with MDD from the DIDA-MDD dataset (5 sites), and 187 HCs and 180 patients with
AD from the MCADI dataset (5 sites).

(i) Individual deviation z scores. The standard protocol for normative model [82] emphasizes the
importance of incorporating some control samples from the same imaging sites as the patients to
the testing set. This is done to verify that the observed case☐control differences are not due to

the analysis with controls in the training set and cases in the testing set [78, 82]. This approach also
allows for the estimation of site effects within the case‒control datasets. To establish the
normative models for all three disorders using the same set of healthy participants, all the HCs of
the three case‒control datasets were randomly divided in half (N_{train} = 654; N_{test} = 659). This
was done in a stratified manner by age, sex, and site. Lifespan connectome-based normative
models were reconstructed by using the training set (N = 32,591), which consisted of half of the
HCs (N_{train} = 654) and all samples of other datasets (N = 31,937). The testing set, comprising
another half of HCs (N_{test} = 659) and the patient cases, was used as a completely independent set
to determine their deviation scores. Specifically, the individual quantile scores were first
estimated relative to the normative curves. Subsequently, the deviation z scores were derived
using quantile randomized residuals [120], an approach that transforms quantiles of the fitted JSU
distribution into standard Gaussian distribution z scores. This process was repeated 100 times,
generating 100 new models and 100 sets of deviation scores for both the patients and the testing
set of the healthy controls. The normality of the distribution of the deviation z scores was
assessed and confirmed using a two-tailed Kolmogorov‒Smirnov test. *P-values* < 0.05 were
observed for all functional metrics in all repetitions. Our subsequent analysis was based on these
independently derived deviation scores in the HCs (HC_{test}) and disease cases.

(ii) Stability of deviation scores across 100 repetitions. To quantitatively assess the similarity
between the estimated growth curves in 100 distinct normative models and the curves in the
main results, we sampled 80 points at one-year intervals for each growth curve and calculated
Pearson's correlation coefficients between the corresponding curves (Supplementary Fig. 15,
Supplementary Tables 5 and 6). The curves of all metrics demonstrated a high degree of
similarity to the main results (mean r > 0.95, mean MSE < 0.1). To evaluate the stability of
individual deviation, we computed the pairwise Pearson's correlation coefficients and MSEs of
the deviation scores among 100 distinct models. The results indicated a high degree of stability
in the estimates of individual deviations for patients within specific disease cohorts (mean r >
0.95, mean MSE < 0.2 for all metrics). For case‒control group comparison analysis and disease
classifications analysis, we replicated the analysis 100 times.

(iii) Individual heterogeneity of deviations. Extreme deviations were defined as z > |2.6|
(corresponding to a p < 0.005), consistent with the criteria used in previous studies [75, 76, 78]. The
extreme positive and negative deviation scores of each functional metric were calculated for each
patient. The percentage map of extreme deviations indicated substantial individual heterogeneity
within each disease group (Supplementary Fig. 16).

(iv) Disease subtypes identification based on individual functional deviations. Given the
substantial individual heterogeneity, we sought to identify subtype differences within each
disease cohort (ASD, MDD, and AD) by employing the data-driven k-means clustering
algorithm. The deviation features of each patient included the global mean of the connectome,
global variance of the connectome, global system segregation, system segregation of each
network, and regional level FCS, encompassing a total of 4,619 features. Dimensionality
reduction was performed on the normalized features using principal component analysis (PCA).
We identified the number of principal components that cumulatively explained more than 95% of
the variance, and these components were then used as the features for clustering analysis. The
similarity matrix of features across patients was calculated using the Euclidean distance. The
optimal number of clusters was determined to be between 2 and 8. A total of 30 different indices

were employed from the NbClust package [126] to determine the optimal number of clusters. The
most frequently identified optimal cluster number was selected as the final cluster count.

(v) Case☐control difference between patients of each subtype and their matched HCs. The
individual deviation scores of patients in each subtype were compared to the median of their
matched HCs. For each metric, the significance of the median differences between the patients
and HCs was assessed using the Mann☐Whitney U test. *P*-values were adjusted for multiple
comparisons using the Benjamini☐Hochberg false discovery rate (FDR) correction across all
possible pairwise tests ($p < 0.05$). For each metric, the case☐control difference analysis was
repeated 100 times. The proportion of tests that passed the significance threshold in 100
comparisons was reported.

(vi) Disease classification based on connectome-based deviations. We performed support vector
machines (SVM) analysis to evaluate the ability of connectome-based deviations in
discriminating patients from controls. For each disease group, two types of classification models
were conducted: classification between all patients and HCs and classification between each
subtype of patients and HCs. Each classification model was repeated 100 times. For each time, a
2-fold cross-validation framework was implemented, with each fold alternately serving as the
training and test sets. To mitigate the impact of features with greater numeric ranges, we
normalized each feature in the training set and applied the resulting parameters to the testing set.
We then plotted receiver operating characteristic (ROC) curves and calculated the areas under the
curve (AUC) to estimate the classification performance. The statistical significance of the AUC
was evaluated using the nonparametric permutation test (1,000 times). During each permutation,
the labels of the patients and controls were randomly shuffled before implementing SVM and
cross-validation. This process yielded a null distribution of the AUC value, and the *P*-value was
computed. Finally, the mean ROC curve was obtained by averaging 100 ROC curves, and the
mean AUC value was obtained by averaging 100 AUC values. The codes for the classification
analysis were modified from Cui et al. [127] (https://github.com/ZaixuCui/Pattern_Classification)
and the libsvm software (www.csie.ntu.edu.tw/~cjlin/libsvm/).

(vii) Predictions of clinical scores based on connectome-based deviations. Using support vector
regression (SVR) with a linear kernel, we sought to assess the ability of the connectome-based
functional deviations to predict the clinical scores of patients. A 2-fold cross-validation
framework was implemented to estimate the prediction accuracy. For a given disease cohort, the
patients were ordered by their target scores and subsequently distributed into alternate folds for
training and testing (e.g., 1[st], 3[rd], …, to the first fold; 2[nd], 4[th], …, to the second fold). Each fold
alternately served as the training and test sets. To mitigate the impact of features with greater
numeric ranges, we normalized each feature in the training set and applied the resulting
parameters to the testing set. The final predictive performance was quantified using Pearson's
correlation coefficients between the predicted and observed clinical scores. The statistical
significance of the prediction accuracy was evaluated using the nonparametric permutation test
(1,000 times). During each permutation, the observed scores of the patients were randomly
shuffled before implementing SVR and cross-validation. This process yielded a null distribution
of the correlation coefficients, and the *P*-value was computed. The codes for the prediction
analysis were modified from Cui and Gong [128]
(https://github.com/ZaixuCui/Pattern_Regression_Matlab) and the libsvm software
(www.csie.ntu.edu.tw/~cjlin/libsvm/).

Data availability

The MRI dataset listed in Supplementary Table 1 are partly available at the Adolescent Brain Cognitive Development Study (https://nda.nih.gov/), the Autism Brain Imaging Data Exchange Initiative (https://fcon_1000.projects.nitrc.org/indi/abide/), the Alzheimer's Disease Neuroimaging Initiative (https://adni.loni.usc.edu/), the Age_ility Project (https://www.nitrc.org/projects/age-ility), the Baby Connectome Project (https://nda.nih.gov/), the Brain Genomics Superstruct Project (https://doi.org/10.7910/DVN/25833), the Calgary Preschool MRI Dataset (https://osf.io/axz5r/), the Cambridge Centre for Ageing and Neuroscience Dataset (https://www.cam-can.org/index.php?content=dataset), the Developing Human Connectome Project (http://www.developingconnectome.org/data-release/second-data-release/), the Human Connectome Project (https://www.humanconnectome.org), the Lifespan Human Connectome Project (https://nda.nih.gov/), the Nathan Kline Institute-Rockland Sample Dataset (https://fcon_1000.projects.nitrc.org/indi/pro/nki.html), the Neuroscience in Psychiatry Network Dataset (https://nspn.org.uk/), the Pediatric Imaging, Neurocognition, and Genetics (PING) Data Repository (http://pingstudy.ucsd.edu/), the Pixar Dataset (https://openfmri.org/dataset/ds000228/), the Strategic Research Program for Brain Sciences (SRPBS) MRI Dataset (https://bicr-resource.atr.jp/srpbsopen/), the Southwest University Adult Lifespan Dataset (http://fcon_1000.projects.nitrc.org/indi/retro/sald.html), the Southwest University Longitudinal Imaging Multimodal Brain Data Repository (http://fcon_1000.projects.nitrc.org/indi/retro/southwestuni_qiu_index.html), and the UK Biobank Brain Imaging Dataset (https://www.ukbiobank.ac.uk/). The dhcpSym surface atlases in aged from 32 to 44 postmenstrual weeks is available at https://brain-development.org/brain-atlases/atlases-from-the-dhcp-project/cortical-surface-template/. The UNC 4D infant cortical surface atlases are available at https://bbm.web.unc.edu/tools/. The fs_LR_32k surface atlas is available at https://balsa.wustl.edu/. The subcortical atlases are available at https://github.com/yetianmed/subcortex. The brain charts and lifespan developmental atlases are shared online via GitHub (https://github.com/sunlianglong/BrainChart-FC-Lifespan).

Code availability

The codes for this manuscript are available on GitHub (https://github.com/sunlianglong/BrainChart-FC-Lifespan). Software packages used in this manuscript include MRIQC v0.15.0 (https://github.com/nipreps/mriqc), QuNex v0.93.2 (https://gitlab.qunex.yale.edu/), HCP pipeline v4.4.0-rc-MOD-e7a6af9 (https://github.com/Washington-University/HCPpipelines/releases), ABCD-HCP pipeline v1 (https://github.com/DCAN-Labs/abcd-hcp-pipeline), dHCP structural pipeline v1 (https://github.com/BioMedIA/dhcp-structural-pipeline), dHCP functional pipeline v1 (https://git.fmrib.ox.ac.uk/seanf/dhcp-neonatal-fmri-pipeline), iBEAT pipeline v1.0.0 (https://github.com/iBEAT-V2/iBEAT-V2.0-Docker), MSM v3.0 (https://github.com/ecr05/MSM_HOCR), FreeSurfer v6.0.0 (https://surfer.nmr.mgh.harvard.edu/), FSL v6.0.5 (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki), Connectome Workbench v1.5.0 (https://www.humanconnectome.org/software/connectome-workbench), MATLAB R2018b (https://www.mathworks.com/products/matlab.html), SPM12 toolbox v6470 (https://www.fil.ion.ucl.ac.uk/spm/software/spm12), GRETNA toolbox v2.0.0 (https://www.nitrc.org/projects/gretna), BrainNet Viewer toolbox v 20191031 (https://www.nitrc.org/projects/bnv), cifti-matlab toolbox v2 (https://github.com/Washington-

University/cifti-matlab), HFR_ai toolbox v1.0-beta-20181108
(https://github.com/MeilingAva/Homologous-Functional-Regions), System segregation code
(https://github.com/mychan24/system-segregation-and-graph-tools), Python v3.8.3
(https://www.python.org), neuroharmonize package v2.1.0
(https://github.com/rpomponio/neuroHarmonize), scikit-learn package v1.1.3 (https://scikit-
learn.org). R v4.2.0 (https://www.r-project.org), GAMLSS package v5.4-3
(https://www.gamlss.com/),and ggplot2 package v3.4.2 (https://ggplot2.tidyverse.org/).

1239

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China (82021004, 82327807, and 31830034 to Y.HE.), the scientific and technological innovation 2030 - the major project of the Brain Science and Brain-Inspired Intelligence Technology (2021ZD0200500 to Q.D. and 2022ZD0211500 to M.R.X.), the Changjiang Scholar Professorship Award (T2015027 to Y.HE.), the Beijing Natural Science Foundation (JQ23033 to M.R.X.), the Beijing United Imaging Research Institute of Intelligent Imaging Foundation (CRIBJZD202102 to M.R.X.), the National Natural Science Foundation of China (31521063 and 31221003 to Q.D., 82071998 to M.R.X., T2325006 to G.L.G., 82202245 to Q.L.L., 81971690 to X.H.L., 32130045 to S.Z.Q., 81571062 and 82172018 to Y.L., 81471120 to X.Z., 61633018 to Y.HAN., 81901101 to P.W., 81400890 to D.W.W., 81920108019, 82330058, 91649117, 81771344, and 81471251 to S.J.Q.), the Beijing Brain Initiative of Beijing Municipal Science & Technology Commission (Z181100001518003 to S.T.), the Fund of Shenzhen Institute for Neuroscience Research (S.T.), the Science and Technology Plan Project of Guangzhou (2018-1002-SF-0442 to S.J.Q.), the Guangzhou Key Laboratory (09002344 to S.J.Q.), and the Key R&D Program of Sichuan Province (2023YFS0076 to T.L.C.). We are grateful to the Adolescent Brain Cognitive Development (ABCD) Study, the Autism Brain Imaging Data Exchange (ABIDE) Initiative, the Alzheimer's Disease Neuroimaging Initiative (ADNI), the Age_ility Project, the Baby Connectome Project (BCP), the Brain Genomics Superstruct Project (BGSP), the Calgary Preschool MRI Dataset, the Cambridge Centre for Ageing and Neuroscience (Cam-CAN) Dataset, the developing Human Connectome Project (dHCP), the Human Connectome Project (HCP), the Lifespan Human Connectome Project (HCPA & HCPD), the Nathan Kline Institute-Rockland Sample (NKI-RS) Dataset, the Neuroscience in Psychiatry Network (NSPN) Dataset, the Pixar Dataset, the Southwest University Adult Lifespan Dataset (SALD), the Southwest University Longitudinal Imaging Multimodal (SLIM) Brain Data Repository, the UK Biobank (UKB) Brain Imaging Dataset, the Disease Imaging Data Archiving: major depressive disorder (DIDA-MDD) Working Group (PI: Yong He, Lingjiang Li, Jingliang Cheng, Qiyong Gong, Ching-Po Lin, Jiang Qiu, Shijun Qiu, Tianmei Si, Yanqing Tang, Fei Wang, Peng Xie, Xiufeng Xu, and Mingrui Xia), the Multi-center Alzheimer Disease Imaging (MCADI) Consortium (PI: Yong Liu, Xi Zhang, Yuying Zhou, Ying Han, and Qing Wang). We thank the National Center for Protein Sciences at Peking University in Beijing, China, for assistance with MRI data acquisition.

ABCD: data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not

necessarily participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. The ABCD data repository grows and changes over time. The ABCD data used in this report came from https://nda.nih.gov/edit_collection.html?id=3165, shared by the DCAN Labs ABCD-BIDS Community Collection (ABCC) (Collection Investigators: Damien Fair).

ABIDE I: primary support for the work by Adriana Di Martino was provided by the (NIMH K23MH087770) and the Leon Levy Foundation. Primary support for the work by Michael P. Milham and the INDI team was provided by gifts from Joseph P. Healy and the Stavros Niarchos Foundation to the Child Mind Institute, as well as by an NIMH award to MPM (NIMH R03MH096321).

ABIDE II: primary support for the work by Adriana Di Martino and her team was provided by the National Institute of Mental Health (NIMH 5R21MH107045). Primary support for the work by Michael P. Milham and his team provided by the National Institute of Mental Health (NIMH 5R21MH107045); Nathan S. Kline Institute of Psychiatric Research). Additional Support was provided by gifts from Joseph P. Healey, Phyllis Green and Randolph Cowen to the Child Mind Institute.

ADNI: data used in preparing this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, many investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators may be found at http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf. Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.;Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

BCP: data used herein is supported by NIH grant (1U01MH110274) and the efforts of the UNC/UMN Baby Connectome Project Consortium.

dHCP: data were provided by the developing Human Connectome Project, KCL-Imperial-
Oxford Consortium funded by the European Research Council under the European Union
Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement no. [319456]. We are
grateful to the families who generously supported this trial.

HCP: data were provided by the Human Connectome Project, WU-Minn Consortium (Principal
Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH
Institutes and Centers which support the NIH Blueprint for Neuroscience Research; and by the
Mc-Donnell Center for Systems Neuroscience at Washington University.

HCP Lifespan: data used in this publication was supported by the National Institute of Mental
Health of the National Institutes of Health under Award Number U01MH109589 and by funds
provided by the McDonnell Center for Systems Neuroscience at Washington University in St.
Louis. The content is solely the responsibility of the authors and does not necessarily represent
the official views of the National Institutes of Health.

NKI-RS: Funding for key personnel provided in part by the New York State Office of Mental
Health and Research Foundation for Mental Hygiene. Additional project support provided by the
NKI Center for Advanced Brain Imaging (CABI), the Brain Research Foundation (Chicago, IL),
the Stavros Niarchos Foundation, and NIH grant P50 MH086385-S1.

NSPN: The NSPN study was funded by a Wellcome Trust award to the University of Cambridge
and the University College London.

UK Biobank: this research has been conducted using data from UK Biobank
(www.ukbiobank.ac.uk). UK Biobank is generously supported by its founding funders the
Wellcome Trust and UK Medical Research Council, as well as the Department of Health,
Scottish Government, the Northwest Regional Development Agency, British Heart Foundation,
and Cancer Research UK.

Competing interests

The authors declare that they have no competing interests.

Author contributions

L.L.S. and Y.H. conceptualized the study. Y.H. supervised the project. L.L.S., T.D.Z., X.Y.L.,
M.R.X., and Y.H. designed the methodology. L.L.S. developed visualizations. Q.L.L., X.H.L.,
D.N.D., Z.L.Z., Z.L.X., and Z.X.C. provided guidance on data analysis and result interpretation.
L.L.S., X.Y.L., Q.W., C.X.P., Q.Y., Q.L.L., Y.H.X. and M.R.X. performed data quality control;
G.L.G., Y.C.B., P.D.C., R.C., Y.C., T.L.C., J.L.C., Y.Q.C., Z.J.D., Y.D., Y.Y.D., Q.D., J.H.G.,
Q.Y.G., Y.H., Z.Z.H., C.C.H., R.W.H., L.J.L., C.P.L., Q.X.L., B.S.L., C.L., N.Y.L., Y.L., J.L.,
L.L.M., W.W.M., S.Z.Q., J.Q., T.M.S., S.P.T., Y.Q.T., S.T., D.W.W., F.W., J.L.W., P.W., X.Q.W.,
Y.P.W., D.T.W., Y.K.W., P.X., X.F.X., L.Y.Y., H.B.Z., X.Z., G.Z., Y.T.Z., S.Y.Z collected a subset
of the data for this study. L.L.S. and Y.H. wrote the manuscript. All authors reviewed the final
manuscript.

1365

Reference

1. Sporns, O., Tononi, G. & Kotter, R. The human connectome: A structural description of the human brain. *PLoS Comput Biol* **1**, e42 (2005).
2. Smith, S.M.*, et al.* Functional connectomics from resting-state fMRI. *Trends Cogn Sci* **17**, 666-682 (2013).
3. Park, H.J. & Friston, K. Structural and functional brain networks: from connections to cognition. *Science* **342**, 1238411 (2013).
4. Bassett, D.S. & Sporns, O. Network neuroscience. *Nat Neurosci* **20**, 353-364 (2017).
5. Biswal, B., Yetkin, F.Z., Haughton, V.M. & Hyde, J.S. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn Reson Med* **34**, 537-541 (1995).
6. Zuo, X.N.*, et al.* Human Connectomics across the Life Span. *Trends Cogn Sci* **21**, 32-45 (2017).
7. Gao, W., Lin, W., Grewen, K. & Gilmore, J.H. Functional Connectivity of the Infant Human Brain: Plastic and Modifiable. *Neuroscientist* **23**, 169-184 (2017).
8. Cao, M., Huang, H. & He, Y. Developmental Connectomics from Infancy through Early Childhood. *Trends Neurosci* **40**, 494-506 (2017).
9. Keunen, K., Counsell, S.J. & Benders, M. The emergence of functional architecture during early brain development. *Neuroimage* **160**, 2-14 (2017).
10. Di Martino, A.*, et al.* Unraveling the miswired connectome: a developmental perspective. *Neuron* **83**, 1335-1353 (2014).
11. Ecker, C., Bookheimer, S.Y. & Murphy, D.G.M. Neuroimaging in autism spectrum disorder: brain structure and function across the lifespan. *The Lancet Neurology* **14**, 1121-1134 (2015).
12. Di Martino, A.*, et al.* Shared and distinct intrinsic functional network centrality in autism and attention-deficit/hyperactivity disorder. *Biol Psychiatry* **74**, 623-632 (2013).
13. Perovnik, M., Rus, T., Schindlbeck, K.A. & Eidelberg, D. Functional brain networks in the evaluation of patients with neurodegenerative disorders. *Nat Rev Neurol* **19**, 73-90 (2023).
14. Chai, Y.*, et al.* Functional connectomics in depression: insights into therapies. *Trends Cogn Sci* (2023).
15. Fornito, A., Zalesky, A. & Breakspear, M. The connectomics of brain disorders. *Nat Rev Neurosci* **16**, 159-172 (2015).
16. Gong, Q. & He, Y. Depression, neuroimaging and connectomics: a selective overview. *Biol Psychiatry* **77**, 223-235 (2015).
17. Cole, T.J. The development of growth references and growth charts. *Ann Hum Biol* **39**, 382-394 (2012).
18. Bethlehem, R.A.I.*, et al.* Brain charts for the human lifespan. *Nature* **604**, 525-533 (2022).
19. Rutherford, S. Charting brain growth and aging at high spatial precision. *eLife* **11**, e72904 (2021).
20. Chen, L.Z., Holmes, A.J., Zuo, X.N. & Dong, Q. Neuroimaging brain growth charts: A road to mental health. *Psychoradiology* **1**, 272-286 (2021).
21. Deery, H.A., Di Paolo, R., Moran, C., Egan, G.F. & Jamadar, S.D. The older adult brain is less modular, more integrated, and less efficient at rest: A systematic review of large-scale resting-state functional brain networks in aging. *Psychophysiology* **60**, e14159 (2023).
22. Edde, M., Leroux, G., Altena, E. & Chanraud, S. Functional brain connectivity changes across the human life span: From fetal development to old age. *J Neurosci Res* **99**, 236-262 (2021).
23. Biswal, B.B.*, et al.* Toward discovery science of human brain function. *Proc Natl Acad Sci U S A* **107**, 4734-4739 (2010).
24. Betzel, R.F.*, et al.* Changes in structural and functional connectivity among resting-state networks across the human lifespan. *Neuroimage* **102 Pt 2**, 345-357 (2014).
25. Cao, M.*, et al.* Topological organization of the human brain functional connectome across the lifespan. *Dev Cogn Neurosci* **7**, 76-93 (2014).
26. Chen, Y.*, et al.* Age-related early/late variations of functional connectivity across the human lifespan. *Neuroradiology* **60**, 403-412 (2018).
27. Petrican, R., Taylor, M.J. & Grady, C.L. Trajectories of brain system maturation from childhood to older adulthood: Implications for lifespan cognitive functioning. *Neuroimage* **163**, 125-149 (2017).
28. Vij, S.G., Nomi, J.S., Dajani, D.R. & Uddin, L.Q. Evolution of spatial and temporal features of functional brain networks across the lifespan. *Neuroimage* **173**, 498-508 (2018).
29. Wang, L., Su, L., Shen, H. & Hu, D. Decoding lifespan changes of the human brain using resting-state functional connectivity MRI. *PLoS One* **7**, e44530 (2012).
30. Rutherford, S.*, et al.* Evidence for embracing normative modeling. *eLife* **12**, e85082 (2023).

31. Fjell, A.M., *et al.* Relationship between structural and functional connectivity change across the adult lifespan: A longitudinal investigation. *Hum Brain Mapp* **38**, 561-573 (2017).

32. Luo, A.C., *et al.* Functional connectivity development along the sensorimotor-association axis enhances the cortical hierarchy. *Nature Communications* **15** (2024).

33. Wu, K., *et al.* Topological organization of functional brain networks in healthy children: differences in relation to age, sex, and intelligence. *PLoS One* **8**, e55347 (2013).

34. Gu, S., *et al.* Emergence of system roles in normative neurodevelopment. *Proc Natl Acad Sci U S A* **112**, 13681-13686 (2015).

35. Sanders, A.F.P., *et al.* Age-related differences in resting-state functional connectivity from childhood to adolescence. *Cereb Cortex* (2023).

36. Sherman, L.E., *et al.* Development of the Default Mode and Central Executive Networks across early adolescence: A longitudinal study. *Developmental Cognitive Neuroscience* **10**, 148-159 (2014).

37. Marek, S., Hwang, K., Foran, W., Hallquist, M.N. & Luna, B. The Contribution of Network Organization and Integration to the Development of Cognitive Control. *PLoS Biol* **13**, e1002328 (2015).

38. Mulders, P.C., van Eijndhoven, P.F., Schene, A.H., Beckmann, C.F. & Tendolkar, I. Resting-state functional connectivity in major depressive disorder: A review. *Neurosci Biobehav Rev* **56**, 330-344 (2015).

39. Hull, J.V., *et al.* Resting-State Functional Connectivity in Autism Spectrum Disorders: A Review. *Front Psychiatry* **7**, 205 (2016).

40. Yu, M., Sporns, O. & Saykin, A.J. The human connectome in Alzheimer disease - relationship to biomarkers and genetics. *Nat Rev Neurol* **17**, 545-563 (2021).

41. Borghi, E., *et al.* Construction of the World Health Organization child growth standards: selection of methods for attained growth curves. *Stat Med* **25**, 247-265 (2006).

42. D. Mikis Stasinopoulos, R.A.R. Generalized Additive Models for Location Scale and Shape (GAMLSS) in R. *Journal of Statistical Software* **23** (2007).

43. Peterson, M.R., *et al.* Normal childhood brain growth and a universal sex and anthropomorphic relationship to cerebrospinal fluid. *J Neurosurg Pediatr* **28**, 458-468 (2021).

44. Zhang, H., *et al.* Growth charts of brain morphometry for preschool children. *Neuroimage* **255**, 119178 (2022).

45. Dong, H.-M., *et al.* Charting brain growth in tandem with brain templates at school age. *Science Bulletin* **65**, 1924-1934 (2020).

46. Gordon, E.M., *et al.* Precision Functional Mapping of Individual Human Brains. *Neuron* **95**, 791-807 e797 (2017).

47. Li, M., *et al.* Performing group-level functional image analyses based on homologous functional regions mapped in individuals. *PLoS Biol* **17**, e2007032 (2019).

48. Kong, R., *et al.* Spatial Topography of Individual-Specific Cortical Networks Predicts Human Cognition, Personality, and Emotion. *Cereb Cortex* **29**, 2533-2551 (2019).

49. Cui, Z., *et al.* Individual Variation in Functional Topography of Association Networks in Youth. *Neuron* (2020).

50. Bijsterbosch, J.D., *et al.* The relationship between spatial configuration and functional connectivity of brain regions. *Elife* **7** (2018).

51. Molloy, M.F. & Saygin, Z.M. Individual variability in functional organization of the neonatal brain. *Neuroimage* **253**, 119101 (2022).

52. Eyre, M., *et al.* The Developing Human Connectome Project: typical and disrupted perinatal functional connectivity. *Brain* **144**, 2199-2213 (2021).

53. Thomason, M.E., *et al.* Age-related increases in long-range connectivity in fetal functional neural connectivity networks in utero. *Dev Cogn Neurosci* **11**, 96-104 (2015).

54. Gao, W., *et al.* Functional Network Development During the First Year: Relative Sequence and Socioeconomic Correlations. *Cereb Cortex* **25**, 2919-2928 (2015).

55. Gao, W., *et al.* The synchronization within and interaction between the default and dorsal attention networks in early infancy. *Cereb Cortex* **23**, 594-603 (2013).

56. Sylvester, C.M., *et al.* Network-specific selectivity of functional connections in the neonatal brain. *Cereb Cortex* **33**, 2200-2214 (2023).

57. Grayson, D.S. & Fair, D.A. Development of large-scale functional networks from birth to adulthood: A guide to the neuroimaging literature. *Neuroimage* **160**, 15-31 (2017).

58. Yeo, B.T., *et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165 (2011).

59. Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human brain connectomics. *PLoS One* **8**, e68910 (2013).

60. Wang, D., *et al.* Parcellating cortical functional networks in individuals. *Nature Neuroscience* **18**, 1853-1860
(2015).
61. Brennan, B.P., *et al.* Use of an Individual-Level Approach to Identify Cortical Connectivity Biomarkers in
Obsessive-Compulsive Disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging* **4**, 27-38 (2019).
62. Cui, W., *et al.* Personalized fMRI Delineates Functional Regions Preserved within Brain Tumors. *Ann Neurol*
91, 353-366 (2022).
63. Li, M., *et al.* Individualized functional connectome identified generalizable biomarkers for psychiatric
symptoms in transdiagnostic patients. *Neuropsychopharmacology* **48**, 633-641 (2023).
64. Chan, M.Y., Park, D.C., Savalia, N.K., Petersen, S.E. & Wig, G.S. Decreased segregation of brain systems
across the healthy adult lifespan. *Proc Natl Acad Sci U S A* **111**, E4997-5006 (2014).
65. Huntenburg, J.M., Bazin, P.L. & Margulies, D.S. Large-Scale Gradients in Human Cortical Organization.
Trends Cogn Sci **22**, 21-31 (2018).
66. Sydnor, V.J., *et al.* Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for
psychopathology. *Neuron* (2021).
67. Xia, Y., *et al.* Development of functional connectome gradients during childhood and adolescence. *Science
Bulletin* (2022).
68. Pines, A.R., *et al.* Dissociable multi-scale patterns of development in personalized brain networks. *Nat
Commun* **13**, 2647 (2022).
69. Alexander-Bloch, A.F., *et al.* On testing for spatial correspondence between maps of human brain structure and
function. *Neuroimage* **178**, 540-551 (2018).
70. Kaczkurkin, A.N., Raznahan, A. & Satterthwaite, T.D. Sex differences in the developing brain: insights from
multimodal neuroimaging. *Neuropsychopharmacology* **44**, 71-85 (2019).
71. Cahill, L. Why sex matters for neuroscience. *Nat Rev Neurosci* **7**, 477-484 (2006).
72. Zhang, C., *et al.* Sex and Age Effects of Functional Connectivity in Early Adulthood. *Brain Connect* **6**, 700-
713 (2016).
73. Dorfschmidt, L., *et al.* Sexually divergent development of depression-related brain networks during healthy
human adolescence. *Science Advances* **8**, eabm7825 %@ 2375-2548 (2022).
74. Zhang, R., Rolls, E.T., Cheng, W. & Feng, J. Different cortical connectivities in human females and males
relate to differences in strength and body composition, reward and emotional systems, and memory. *Brain Struct
Funct* **229**, 47-61 (2024).
75. Sun, X., *et al.* Mapping Neurophysiological Subtypes of Major Depressive Disorder Using Normative Models
of the Functional Connectome. *Biol Psychiatry* (2023).
76. Wolfers, T., *et al.* Mapping the Heterogeneous Phenotype of Schizophrenia and Bipolar Disorder Using
Normative Models. *JAMA Psychiatry* **75**, 1146-1155 (2018).
77. Rutherford, S. Evidence for embracing normative modeling. *Elife* **12**, e85082 (2023).
78. Segal, A., *et al.* Regional, circuit and network heterogeneity of brain abnormalities in psychiatric disorders. *Nat
Neurosci* (2023).
79. Marquand, A.F., Rezek, I., Buitelaar, J. & Beckmann, C.F. Understanding Heterogeneity in Clinical Cohorts
Using Normative Models: Beyond Case-Control Studies. *Biol Psychiatry* **80**, 552-561 (2016).
80. Tian, Y.E., *et al.* Evaluation of Brain-Body Health in Individuals With Common Neuropsychiatric Disorders.
JAMA Psychiatry **80**, 567-576 (2023).
81. Tian, Y.E., *et al.* Heterogeneous aging across multiple organ systems and prediction of chronic disease and
mortality. *Nat Med* **29**, 1221-1231 (2023).
82. Rutherford, S., *et al.* The normative modeling framework for computational psychiatry. *Nat Protoc* **17**, 1711-
1734 (2022).
83. van den Heuvel, M.I. & Thomason, M.E. Functional Connectivity of the Human Brain in Utero. *Trends Cogn
Sci* **20**, 931-939 (2016).
84. Raz, G. & Saxe, R. Learning in Infancy Is Active, Endogenously Motivated, and Depends on the Prefrontal
Cortices. *Annual Review of Developmental Psychology* **2**, 247-268 (2020).
85. Gao, W. Evidence on the emergence of the brain's default network from 2-week-old to 2-year-old healthy
pediatric subjects. *PNAS* (2009).
86. Fjell, A.M., *et al.* What is normal in normal aging? Effects of aging, amyloid and Alzheimer's disease on the
cerebral cortex and the hippocampus. *Prog Neurobiol* **117**, 20-40 (2014).
87. Pascoal, T.A., *et al.* Abeta-induced vulnerability propagates via the brain's default mode network. *Nat Commun*
10, 2353 (2019).
88. Wales, R.M. & Leung, H.-C.J.B.C. The effects of amyloid and tau on functional network connectivity in older

populations. **11**, 599-612 (2021).

89.   Jagust, W. Imaging the evolution and pathophysiology of Alzheimer disease. *Nat Rev Neurosci* **19**, 687-700 (2018).

90.   Bedford, S.A., Seidlitz, J. & Bethlehem, R.A.I. Translational potential of human brain charts. *Clin Transl Med* **12**, e960 (2022).

91.   Zhou, Z.X., Chen, L.Z., Milham, M.P., Zuo, X.N. & Lifespan Brain Chart, C. Six cornerstones for translational brain charts. *Sci Bull (Beijing)* (2023).

92.   Cash, R.F.H., Cocchi, L., Lv, J., Fitzgerald, P.B. & Zalesky, A. Functional magnetic resonance imaging–guided personalization of transcranial magnetic stimulation treatment for depression. *JAMA psychiatry* **78**, 337-339 %@ 2168-2622X (2021).

93.   Lotter, L.D., *et al.* Human cortex development is shaped by molecular and cellular brain systems. *bioRxiv* (2023).

94.   Brouwer, R.M., *et al.* Genetic variants associated with longitudinal changes in brain structure across the lifespan. *Nat Neurosci* **25**, 421-432 (2022).

95.   Kopal, J., Uddin, L.Q. & Bzdok, D. The end game: respecting major sources of population diversity. *Nat Methods* (2023).

96.   Stenger, V.A. Technical considerations for BOLD fMRI of the orbitofrontal cortex. in *The Orbitofrontal Cortex* (ed. D. Zald & S. Rauch) 0 (Oxford University Press, 2006).

97.   Deichmann, R., Gottfried, J.A., Hutton, C. & Turner, R. Optimized EPI for fMRI studies of the orbitofrontal cortex. *Neuroimage* **19**, 430-441 (2003).

98.   Kia, S.M., *et al.* Closing the life-cycle of normative modeling using federated hierarchical Bayesian regression. *PLoS One* **17**, e0278776 (2022).

99.   Bayer, J.M.M., *et al.* Accommodating site variation in neuroimaging data using normative and hierarchical Bayesian models. *Neuroimage* **264**, 119699 (2022).

100. Di Biase, M.A., *et al.* Mapping human brain charts cross-sectionally and longitudinally. *Proc Natl Acad Sci U S A* **120**, e2216798120 (2023).

101. Elyounssi, S., *et al.* Uncovering and mitigating bias in large, automated MRI analyses of brain development. *bioRxiv* (2023).

102. Esteban, O., *et al.* MRIQC: Advancing the automatic prediction of image quality in MRI from unseen sites. *PLoS One* **12**, e0184661 (2017).

103. Rosen, A.F.G., *et al.* Quantitative assessment of structural image quality. *Neuroimage* **169**, 407-418 (2018).

104. Wang, L., *et al.* Benchmark on Automatic 6-month-old Infant Brain Segmentation Algorithms: The iSeg-2017 Challenge. *IEEE Trans Med Imaging* (2019).

105. Zeng, Z., *et al.* 3D☐MASNet: 3D mixed☐scale asymmetric convolutional segmentation network for 6☐month☐old infant brain MR images. *Human Brain Mapping* (2022).

106. Glasser, M.F., *et al.* The minimal preprocessing pipelines for the Human Connectome Project. *Neuroimage* **80**, 105-124 (2013).

107. Ji, J.L., *et al.* QuNex—An integrative platform for reproducible neuroimaging analytics. *Frontiers in Neuroinformatics* **17** (2023).

108. Feczko, E., *et al.* Adolescent Brain Cognitive Development (ABCD) Community MRI Collection and Utilities. (2021).

109. Makropoulos, A., *et al.* The developing human connectome project: A minimal processing pipeline for neonatal cortical surface reconstruction. *Neuroimage* **173**, 88-112 (2018).

110. Wang, L., *et al.* iBEAT V2.0: a multisite-applicable, deep learning-based pipeline for infant cerebral cortical surface reconstruction. *Nat Protoc* (2023).

111. Williams, L.Z.J., *et al.* Structural and functional asymmetry of the neonatal cerebral cortex. *Nat Hum Behav* (2023).

112. Wu, Z., *et al.* Construction of 4D infant cortical surface atlases with sharp folding patterns via spherical patch-based group-wise sparse representation. *Hum Brain Mapp* **40**, 3860-3880 (2019).

113. Fitzgibbon, S.P., *et al.* The developing Human Connectome Project (dHCP) automated resting-state functional processing framework for newborn infants. *Neuroimage* **223**, 117303 (2020).

114. Eickhoff, S.B., Yeo, B.T.T. & Genon, S. Imaging-based parcellations of the human brain. *Nat Rev Neurosci* **19**, 672-686 (2018).

115. Buckner, R.L., Krienen, F.M., Castellanos, A., Diaz, J.C. & Yeo, B.T. The organization of the human cerebellum estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 2322-2345 (2011).

116. Mueller, S., *et al.* Individual variability in functional connectivity architecture of the human brain. *Neuron* **77**,

586-595 (2013).

117. Sun, L., *et al.* Structural insight into the individual variability architecture of the functional brain connectome. *Neuroimage*, 119387 (2022).

118. Power, J.D., *et al.* Functional network organization of the human brain. *Neuron* **72**, 665-678 (2011).

119. Craddock, R.C., James, G.A., Holtzheimer, P.E., 3rd, Hu, X.P. & Mayberg, H.S. A whole brain fMRI atlas generated via spatially constrained spectral clustering. *Hum Brain Mapp* **33**, 1914-1928 (2012).

120. Dunn, P.K. & Smyth, G.K. Randomized Quantile Residuals. *Journal of Computational and Graphical Statistics* **5**, 236-244 (1996).

121. Dinga, R., *et al.* Normative modeling of neuroimaging data using generalized additive models of location scale and shape. *bioRxiv* (2021).

122. Flexible Regression and Smoothing Using GAMLSS in R.

123. Satterthwaite, T.D., *et al.* Heterogeneous impact of motion on fundamental patterns of developmental changes in functional connectivity during youth. *Neuroimage* **83**, 45-57 (2013).

124. Ciric, R., *et al.* Mitigating head motion artifact in functional connectivity MRI. *Nat Protoc* **13**, 2801-2826 (2018).

125. Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L. & Petersen, S.E. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* **59**, 2142-2154 (2012).

126. Charrad, M., Ghazzali, N., Boiteau, V. & Niknafs, A.J.J.o.s.s. NbClust: an R package for determining the relevant number of clusters in a data set. **61**, 1-36 (2014).

127. Cui, Z., Xia, Z., Su, M., Shu, H. & Gong, G. Disrupted white matter connectivity underlying developmental dyslexia: A machine learning approach. *Hum Brain Mapp* **37**, 1443-1458 (2016).

128. Cui, Z. & Gong, G. The effect of machine learning regression algorithms and sample size on individualized behavioral prediction with functional connectivity features. *Neuroimage* **178**, 622-637 (2018).