Systemic 4-1BB stimulation augments extrafollicular memory B cell formation and recall responses during *Plasmodium* infection.

Carolina Caloba[1,2,3], Alexandria J. Sturtz[4], Taylor A. Lyons[3], Lijo John[1,5], Akshaya Ramachandran[1,2], Allen M. Minns[6], Anthony M. Cannon[7], Justin P. Whalley[1,3], Tania H. Watts[8], Mark H. Kaplan[7], Scott E. Lindner[5], Rahul Vijay[1,3]

[1]Discipline of Microbiology and Immunology, [2]School of Graduate and Postdoctoral Studies, [3] Center for Cancer Biology, Immunology and Infection, Rosalind Franklin University of Medicine and Science, North Chicago, IL, USA. [4]Division of Biology and Biomedical Sciences, Washington University, St. Louis, Missouri, USA. [5]Deparment of Veterinary Biochemistry, Kerala Veterinary and Animal Sciences University, Kerala, India. [6]Department of Biochemistry and Molecular Biology, Huck Center for Malaria Research, Pennsylvania State University, University Park, Pennsylvania, USA. [7]Department of Microbiology and Immunology, Indiana University School of Medicine, Indianapolis, IN, USA. [8]Department of Immunology, Temerty Faculty of Medicine, University of Toronto, Toronto, ON, Canada.

Running Title: 4-1BB stimulation and memory B cells in malaria

Summary:
T-dependent germinal center (GC) output, comprising plasma cells (PC) and memory B cells (MBC), is crucial for the clearance of *Plasmodium* infection and protection against reinfection. In this study, we examined the effect of an agonistic antibody targeting 4-1BB (CD137), a member of the Tumor Necrosis Factor Receptor Superfamily (TNFRSF), during experimental malaria. Here we show that exogenous 4-1BB stimulation, despite delaying the effector GC response, surprisingly enhanced humoral memory recall and protection from reinfection. Single cell RNA and ATAC sequencing of MBCs from mice that received 4-1BB stimulation revealed distinct populations with transcriptional and epigenetic signatures indicative of superior recall and proliferative potential. Importantly, our results indicate that the effects of 4-1BB stimulation are dependent on IL-9R signaling in B cells but independent of parasite load during primary infection. Our study proposes an immunomodulatory approach to enhance the quality of the MBC pool, providing superior protection during infection and vaccination, particularly in the context of malaria.

Introduction

Despite decades of research, malaria remains a major public health threat, with an estimated 249 million cases and 608,000 deaths in 2022[1]. It predominantly affects poorer regions, with almost 95% of cases and 98% of deaths documented in Africa. The severity of the disease is especially prominent in individuals with limited or no prior exposure to the parasite, making children under the age of 5 the most impacted[1,2]. The newly approved anti-malarial vaccines RTS,S/AS01 and R21 provide waning efficacy even after a multi-dose regimen[3]. Clearance of the malarial parasite relies on CD4 T cell-dependent B cell activation, culminating in the production of parasite-specific antibodies during the blood stage of infection[4,5]. Prior to or concomitant with the clearance of infection, a fraction of activated B cells differentiates into memory B cells (MBCs) which quickly become antibody-secreting plasma cells (PCs) upon reinfection[6-11,12]. However, increasing evidence suggests that even after multiple exposures, naturally acquired anti-*Plasmodium* humoral immune memory affords just clinical immunity and not sterilizing immunity, resulting in repeat infections[13].

Leveraging numerous studies in tumor models, the expression and engagement of coinhibitory receptors such as PD-1, LAG-3, and CTLA-4 on CD4 T cells have been identified as contributing to the inefficient anti-*Plasmodium* immune responses [14-18]. In one such study, exogenous

activation of OX40, a member of the tumor necrosis factor receptor superfamily (TNFRSF)
expressed on T cells improved *Plasmodium* clearance following primary infection and also
rendered protection from rechallenge in a model of experimental malaria [16,17]. These studies
highlight substantial room for improvement in the natural immune response against *Plasmodium*
and underscore the potential of exogenous costimulatory agonists to fully harness the full breadth
of the immune system.
4-1BB (CD137), another TNFRSF member predominantly expressed on T cells, has been
shown to have non-redundant roles with OX40 in enhancing CD4 T cell function[19,20]. Given that
CD4 T cells play a key role in anti-*Plasmodium* immunity by providing help to B cells, we
investigated the role of exogenous 4-1BB stimulation in a rodent malaria model. Here, we show
that exogenous ligation of 4-1BB using a monoclonal antibody significantly delayed the effector
humoral response while paradoxically enhancing humoral immune memory. Transcriptomic and
epigenetic analysis of MBCs from treated mice suggests they have a distinct genetic signature
that facilitate superior recall and strongly points to an extrafollicular origin. Additionally, this
enhanced protection following 4-1BB activation depended on IFNγ-induced expression of T-BET
in B cells during the effector phase and, more importantly, IL-9:IL-9R signaling on MBCs.
Collectively, our study establishes a previously unidentified pathway by which MBCs can be
programmed to attain superior recall potential and identify key molecules that may be targeted to
enhance vaccine efficacy especially in *Plasmodium* infections.
69
70 **Materials and Methods**
71
72 **Animals, infection, treatments and parasitemia**
73
C57BL/6 mice and *Tbx21*[-/-] mice were purchased from The Jackson Laboratory. 4-1BB[-/-] mice
were originally obtained from Byoug S. Kwon, Korean National University[21] and transferred to
THW and RV labs for these studies. *Plasmodium yoelli* (*Py*) clone 17XNL and *P. berghei* clone
ANKA (*PbA*) were obtained from the Malaria Research and Reference Reagent Resource Center
(MR4; American Type Culture Collection), stocks generated and mice infected as previously
described[22]. Briefly, frozen parasite stocks (200μL aliquots) were resuspended in 4mL of normal
saline or PBS (1:20 dilution) and animals were infected with 1×10^6 parasitized RBCs
intravenously. Treatments with all biologics were done intraperitoneally in a 200μL volume (in
PBS) as follows: 50 μg of anti-4-1BB antibody (3H3; BioXCell), 200μg of IFNγ blocking antibody
(XMG1.2; Bioxcell), 100μg IL-9 blocking antibody (9C1; Bioxcell) or rat IgG (rIgG) or mouse IgG
(mIgG) on indicated timepoints. Mice were treated with artemether (Biotechne) at 20mg/kg,
dissolved in 50μL of mineral oil at indicated timepoints. Parasitemia was measured by flow
cytometry as previously described[23,24]. All experiments and procedures were approved by the
Rosalind Franklin University of Medicine and Science Institutional Animal Care and Use
Committee.
89
90 ***Competitive mixed bone marrow chimeras***
91
C57BL/6 mice (CD45.1/.1) were lethally irradiated (950 rads) and transferred with 10×10^6 cells
containing a mixture of μMT cells (CD45.1/.2) along with *Tbx21*[-/-] (CD45.2/.2) or *Il9r*[-/-] (CD45.2/.2)
or WT (CD45.2/.2) cells at an 8:2 ratio, respectively. For CD4-specific competitive mixed bone
marrow chimeras, TCRα[-/-] were used as recipients and transferred with 10×10^6 cells of CD4[-/-] cells
(CD45.2/CD45.2) along with *4*-1BB[-/-] (CD45.1/.1) or WT (CD45.1/.1) cells at an 8:2 ratio,
respectively. At 6 weeks post transfer mice were checked for reconstitution (>90% donor; 80%

μMT or TCRα cells and 10% KO cells) by flow cytometry and at 8 weeks post transfer,
reconstituted mice were used for the experiments listed elsewhere.
100
101   **Confocal imaging**
102
103   Spleens were processed as previously described[22]. Antibodies used for staining are B220-AF488
104   (clone RA3-6B2; eBioscience), CD4-AF594 (clone GK1.5; BioLegend) and GL7-AF647 (BD
105   Pharmingen). Imaging was done using a Zeiss LSM710 confocal microscope and images were
106   processed using IMARIS x64 software (version 9.2.1).
107
108   **ELISpot and ELISA**
109
For ELISPOT, Nunc MaxiSorp white polystyrene plates were coated with $0.5\,\mu g\,ml^{-1}$ of
recombinant $MSP1_{19}$ and blocked for at least 2 h with supplemented Roswell Park Memorial
Institute (RPMI) 1640 media. Bone marrow cells from *PbA*-infected mice were serially diluted in
supplemented RPMI 1640 media and plated for 20 h at 37 °C with 5% CO_2. Plates were washed
with PBS + 0.05% Tween 20 and incubated overnight with horseradish peroxidase (HRP)-
conjugated IgM at 4 °C. Spots were developed with 3-amino-9-ethylcarbazole*.*
For ELISA, Nunc MaxiSorp plates were coated with $0.5\,\mu g\,ml^{-1}$ $MSP1_{19}$ and blocked with 2.5%
w/v BSA + 5% v/v fetal bovine serum. Serum samples from naive and *Py*-infected mice were
serially diluted and incubated for 18 h at 4 °C. $MSP1_{19}$-specific antibodies were detected with
HRP-conjugated IgM, IgG2b or IgG2c at 1:1000 dilution. Plates were developed using a SureBlue
reserve TMB Kit (KPL), according to the manufacturer's protocol, and absorbance was measured
at an optical density of 450 nm using a Accuris ELISA plate reader. End-point titers were
extrapolated from a sigmoidal 4PL (where x is the log concentration) standard curve for each
sample. The threshold for end-point titers was the mean plus 0.5–8 multiplied by the standard
deviation recorded for naive mouse sera.
125
126   *Ex vivo B cell activation*
127
128   Splenic cells were enriched for B cells using the MojoSort™ Mouse Pan B Cell Isolation Kit
129   (Biolegend) according to the manufacturer's protocol. 7.5 x10^5 B cells were plated in triplicates
130   and were activated with anti-IgM (10 μg/mL) and anti-CD40 (10 μg/mL). Some wells received rIFNγ
131   (20ng/mL).
132
133   *Intracellular cytokine stimulation*
134
135   Single cell suspension of splenocytes were prepared and 1x 10^6 were seeded in U-bottom wells
136   in 96-well plate. The cells were stimulated with cell activation cocktail (BioLegend) containing
137   phorbol myristate acetate (PMA) and Ionomycin in the presence of brefeldin A (BioLegend) for 4
138   hours at 37°C and 5% CO_2. After incubation, cells were stained as described elsewhere in the
139   manuscript.
140
141   *SpyCage Reagent for detecting and enriching MSP1$_{-19}$ specific B cells*
142
SpyCage reagents bearing 60 copies of either mScarlet ("RedCage") or mNeonGreen
("GreenCage") were covalently loaded with 60 copies of the immunodominant *P. yoelii* antigen
MSP-1 or the liver stage antigen *Py*UIS4 as a decoy control, respectively. Antibodies specific to
SpyCage were biotinylated using amine-reactive biotinylated crosslinkers and were used to enrich

SpyCage-bound cells[25]. Briefly, single cell suspensions were stained with 1.25mg of *Py*UIS4 for
10 minutes at room temperature followed by *Py*MSP-1 (1.25mg) staining for 30 minutes at 4°C.
Cells were washed and incubated with 1mg of anti-aldolase biotynilated antibody for 30 minutes
at 4°C. Stained cells were then washed and enriched with MojoSort™ Streptavidin nanobeads
(Biolegend), according to the manufacturer's protocol. Enriched cells were used for surface and
intranuclear staining as detailed below.
153
154   **Flow cytometry**
155
Spleens were harvested and homogenized through wire meshes to obtain single cell
suspensions. After RBCs lysis, cells were filtered and counted. Samples were blocked using Fc
block (clone 2.4G2) in FACS buffer (PBS + 0.002% w/v sodium azide + 2% v/v fetal bovine serum)
and cells were stained with fluorescently labeled antibodies. Transcription factors were stained
using True-Nuclear™ Transcription Factor Buffer Set (Biolegend), according to manufacturer's
protocol. For intracellular cytokine staining Cyto-Fast Fix/Perm (Biolegend) kit was used
according to manufacturer's protocol. Samples were acquired on BD-LSR-II or Cytek Aurora and
analyzed using FlowJo software (TreeStar Inc).
164
165   *MBC transfer*
166
167   Memory B cells (B220$^+$IgD$^-$CD38$^+$GL7$^-$) were sort-purified from *Py*-infected mice treated with rIgG
168   or 3H3. 2x10^5 cells were transferred intravenously to naive μMT mice, which were infected with
169   1 × 10^6 *Py* -parasitized RBCs, the following day.
170
171   *Single-cell RNA sequencing analysis*
172
Single cell suspensions were stained with Live/Dead dye (Biolegend) and then divided in a ratio
of 1:4 in which 1 part was used for sorting of polyclonal cells and 4 parts were used for MSP-1
specific cells. Before staining for surface markers, MSP-1 specific cells were stained and enriched
with SpyCage reagents as previously described. Both polyclonal and MSP-1 cells were incubated
with fluorescently labeled antibodies and Total-seq C antibodies (Biolegend) for 30 minutes and
washed before sorting for resting B cells (B220$^+$IgD$^+$), polyclonal (B220$^+$IgD$^-$GL7$^-$CD38$^+$) and
MSP-1 specific (B220$^+$IgD$^-$GL7$^-$CD38$^+$Decoy$^-$MSP1$^+$) MBCs. Post-sorted cells were run on the
10X Chromium (10X Genomics) and library preparation was performed according to the
manufacturer's protocol for Chromium Next GEM Single Cell 5' Reagent Kit v2 with Feature
Barcode technology and V(D)J amplification for mouse B cells (10X Genomics). Libraries were
pooled and sequenced in the Genomics and Microbiome Core Facility at Rush University using
NovaSeqX Plus (Illumina). 10X Cell Ranger Multi (v.7.1.0, 10X Genomics) was used to process
gene expression, CITE-seq and V(D)J libraries in each sample individually.
Downstream analysis was performed using the Seurat package[26] (version 5.1.0) with R
(version 4.4.1). Samples were merged (rIgG and 3H3 groups in Polyclonal or MSP-1 specific pool)
and the SCTransform[27] function using the glmGamPoi[28] package was utilized for normalization,
selection of highly variable features and scaling of data. Principal component analysis (PCA) was
performed and 30 dimensions were used for dimensionality reduction with Uniform Manifold
Approximation and Projection (UMAP). Due to observed batch effects, the SCT datasets were
integrated using the IntegrateLayers() function. Unsupervised clustering was performed using
FindNeighbors() and FindClusters() function and RunUMAP() was used with reduction set to
"integrated.dr". For CITE-seq, NormalizeData() and ScaleData() were used for normalization.

Differential expression analysis was performed by PrepSCTFindMarkers() followed by
FindMarkers() function, which were used to identify the markers of each cluster and differentially
expressed genes (DEGs) in the 3H3 compared to rIgG group. Of the identified cluster markers,
only genes with adjusted p-value < 0.05 and average fold change > 0.3 were used for the
heatmaps, which show the aggregated expression of all cells within a specific cluster. Of the
DEGs between 3H3 versus rIgG group, only the ones with adjusted p-value < 0.05 were kept for
gene ontology (GO) analysis. Enrichment for GO analysis was performed with the clusterProfiler
package (v4.12.6)[29] with q- and p-value cutoffs of 0.05 and ontology set to "Biological Processes".
To remove redundancy of enriched terms, the function simplify() was used. To generate a gene
signature score for each cell in the SCT dataset, the function AddModuleScore() was used with
gene lists of GC-derived, EF-derived or atypical MBCs (**Supp. Table 1**) . Single-cell trajectories
(pseudotime analysis) were constructed using the Monocle3 package[30-34]. Cells were ordered by
choosing the root cluster based on expression of genes associated with stemness and the earliest
principal node was defined programmatically according to Monocle3's vignette. RColorBrewer[35],
ggplot2[36], scCustomize[37] and viridis[38] packages were used for visualization of the data.
210
211   *V(D)J analysis*
212
BCR repertoire analysis was performed using Immcantation packages[39-43]. First, V, D and J genes
were assigned using IgBLAST followed by removal of non-productive sequences and cells with
multiple or without heavy chains. Then, cell IDs were matched with the gene expression data, cell
type annotations were transferred and cells without gene expression data were removed. Clonal
analysis was performed followed by generation of germline sequences. The V gene somatic
hypermutation was calculated and used to compute the median mutation frequency of a clone.
For visualization of data, the package ggplot2[44] was used.
220
221   **Single-cell ATAC sequencing analysis**
222
Polyclonal and MSP-1 specific MBCs were flow-sorted as described for scRNA-seq and nuclei
was isolated. Transposition, barcoding on the 10X Chromium (10X Genomics) and library
construction were performed according to the manufacturer's protocol for Chromium Next GEM
Single Cell ATAC Reagents Kits v2 (10X Genomics). Libraries were pooled and sequenced in the
Genomics and Microbiome Core Facility at Rush University using NovaSeqX Plus (Illumina). 10X
Cell Ranger ATAC (v.2.1.0, 10X Genomics) was used to process the data.
Downstream analysis were performed using Seurat[26] (version 5.1.0) and Signac[45] (version
1.14.0) packages. Cells with count lower than 500 were excluded and common peak sets (for
polyclonal and MSP-1 specific MBCs) were created and filtered out based on length (10000 >
peak width > 20). Peaks were quantified in each sample individually and the common set of peaks
was passed as the feature argument. Samples were merged (rIgG and 3H3 groups in Polyclonal
or MSP-1 specific pool) as Seurat objects and features that corresponded to sequences other
than the standard chromosomes were removed. After gene annotation and based on the quality
control metrics, cells were filtered based on the quantified peaks (100000 > quantified peaks >
100), frequency of reads in peaks (20%), ratio of reads mapping in blacklist regions compared to
peaks (0.08), nucleosome signal (signal < 4) and transcription start site enrichment (TSS > 2).
Normalization, selection of top features and dimensional reduction were performed by
RunTFIDF(), FindTopFeatures() and RunSVD() functions, respectively. The first LSI component
was removed since it strongly correlated to sequencing depth. Clustering was performed using
the FindNeighbors() function with reduction set to "lsi" followed by FindClusters() function.
GeneActivity() was used to create a gene activity matrix, which was normalized and scaled by

NormalizeData() and ScaleData(), respectively. FindTransferAnchors() and TransferData()
functions were used to integrate scATAC-seq data with gene expression data from the scRNA-
seq. Differentially accessible regions (DARs) were calculated using the FindMarkers() function
using a logistic regression (LR) model. For the total DARs, it was used a threshold of adjusted p-
value < 0.05 and average log2foldchange > 0.3 and for the top DARs to be used for motif analysis
it was used a threshold of adjusted p-value < 0.005 and of frequency of cells where the DAR is
detected > 0.2. The list of total DARs was used to calculate the closest genes to the identified
peaks, while the top DARs were used to find overrepresented motifs in the accessible regions.
The list of motif position frequency matrices was obtained from JASPAR2020[46] database. was
used RColorBrewer[35], scCustomize[37] and ggseqlogo[47] packages were used for visualization of
the data.
255
256 ***Quantitative real-time PCR***
257
MBCs were flow sorted from *Py*-infected mice that were treated with 3H3 or rIgG and then with
artemether on time points indicated elsewhere. RNA was extracted using Trizol reagent according
to the manufacturer's protocol. 2μg of RNA was used for cDNA synthesis using MMLV reagent.
cDNA was added to the 2X PCR PowerUP SYBR Green Master mix at a 1:20 ratio, along with
0.2μM of forward and reverse primers. Primer sequences used in the experiments are as follows:
Gapdh Fwd: 5'-GAG AAC TTT GGC ATT GTG G-3'; *Gapdh* Rev: 5'-ATG CAG GGA TGA TGT
TCT G-3'; *Il9r* Fwd: 5'-GGA CAG TTG GCA GTA AGT CAC C-3'; *Il9r* Rev: 5'-CCA CTC TCT CCA
AGG TCC AA-3'. *Ct* value were normalized to *Gapdh* using the following equation: $\Delta Ct = Ct\ (Il9r)$
$-Ct(Gapdh).$ Results are represented as a ratio of *Gapdh* $2^{-(\Delta Ct)}$
267
268
269 ***Statistical analysis***
270
271 Statistical analyses were performed using Prism 10 software (GraphPad). Specifics of each tests
272 are detailed in the figure legends.
273
274 **Results**
275
276 **Systemic 4-1BB stimulation delays effector GC response and parasite control.**
277
In our preliminary experiments with a non-lethal rodent *Plasmodium* strain, *Plasmodium yoelii*
(*Py*), we observed that 4-1BB was preferentially upregulated on splenic CD4 T cells during the
course of infection (**Supp Fig. 1a**). Given the upregulation of 4-1BB, we infected (WT) C57BL/6
mice or 4-1BB[-/-] mice with *Py* and tracked the parasite load. As shown in **Supp Fig. 1b** 4-1BB[-/-]
mice had an elevated parasite load compared to WT mice, suggesting a role for this receptor in
anti-*Plasmodium* immune response. We next sought to investigate the effect of exogenous
ligation of 4-1BB during infection. To this end, we infected WT mice with *Py* and treated them with
an agonistic monoclonal antibody to 4-1BB (3H3), or the isotype control, on 5 and 7 days post-
infection (dpi) (**Fig. 1a**). We then monitored the kinetics of parasite load by measuring the
frequency of parasitized RBCs during the course of infection using flow cytometry as previously
described[23]. As a TNFRSF member with a co-stimulatory role like OX40[48], we anticipated that 4-
1BB stimulation would enhance the anti-*Plasmodium* humoral immune response. Contrary to this
expectation, mice treated with 3H3 exhibited a prolonged infection course with an 8-fold higher
total parasite load compared to controls, eventually resolving by 39-40 dpi (**Fig. 1b**). Additionally,

we observed a delayed onset of germinal center (GC) responses—both GC B cells (**Supp Fig. 1c and Fig. 1c**) and GC Tfh cells (**Supp Fig. 1d and Fig. 1d**)—in 3H3-treated mice, peaking only around 39 dpi, suggesting that parasite clearance may be GC-dependent. Not surprisingly, serum parasite-specific antibody levels were also slow to accumulate in 3H3-treated mice (**Fig. 1e**). Confocal imaging of spleens from either group of mice revealed a pronounced disruption of lymphoid architecture in 3H3-treated mice, with no apparent follicular organization until 21 dpi, after which the follicles grouped into GCs as evidenced by the appearance of GL7 staining within the follicles (**Fig. 1f**). Importantly, 3H3-driven loss of parasite control was reversed in mice that lacked 4-1BB in CD4 T cells (CD4$^{\text{4-1BBL-/-}}$) (**Supp Fig. 1e**). These data collectively demonstrate that exogenous ligation of 4-1BB on CD4 T cells in *Plasmodium*-infected mice results in derailed GC responses and heightened parasite load.

Enhanced protection following rechallenge in mice exposed to systemic 4-1BB stimulation.

Infection of mice with *P. berghei ANKA* (*PbA*) is invariably lethal even in mice that recovered from *Py* infection. This challenge protocol is commonly used and is considered a stringent test for MBC recall. Given that exogenous 4-1BB ligation delayed GC responses and increased the parasite burden during the primary *Py* infection, we hypothesized that mice treated with 3H3 would exhibit impaired memory responses and, consequently, reduced protection upon challenge. To test this, we monitored survival in *Py*-infected mice treated with either 3H3 or rIgG after challenge with *PbA* on day 90 post *Py* infection, when both groups were in convalescence (**Fig. 2a**). Surprisingly, 3H3-treated mice exhibited significantly longer survival compared to the rIgG group (**Fig. 2b**). To directly measure MBC recall, we assessed the fold change in antibody titers between day 0 and day 5 post *PbA* challenge. Consistent with the enhanced survival of 3H3-treated mice, we observed a greater fold increase in anti-merozoite surface protein 1 (MSP-1$_{-19}$)-specific antibody (IgM) titers between day 0 and day 5 post *PbA* challenge (**Fig. 2c**), indicating an enhanced MBC recall response. An increase in parasite-specific antibody titers during this short window would indicate that the antibodies are derived from PCs differentiating from an existing MBC pool, rather than from a *de novo* GC reaction. The higher parasite load observed during the primary *Py* infection following 3H3 treatment (**Fig. 1b**) raises the possibility that the enhanced protection following *PbA* challenge may result from elevated inflammation and/or residual antigen load. To investigate whether these factors influenced the outcome, we truncated the primary infection and the initial parasite load by treating mice with the antimalarial drug artemether at indicated time points (**Fig. 2d**). As expected, artemether treatment resulted in complete parasite clearance in both groups, with 3H3 group showing no evidence of a productive GC response even by 21 dpi, while rIgG group retained the GC B cell and GC Tfh cells (**Supp Fig. 2a-c**) as was seen when mice were not treated with artemether (**Fig. 1**). Upon challenge with *PbA* on day 38 post-*Py* infection, we observed significantly enhanced protection in 3H3-treated mice that received chemoprophylaxis (**Fig. 2e**). Correspondingly, these mice also showed a higher fold change in anti-MSP1 antibody titers post *PbA* challenge (**Fig. 2f**). To further test whether 3H3-mediated protection is relevant to attenuated *Plasmodium* parasite vaccination, we immunized mice with irradiated blood-stage *Py* parasites and administered either 3H3 or rIgG on days 5 and 7 post-infection. On days 30-35 post-vaccination, the mice were challenged with *PbA* and monitored for survival (**Fig. 2g**). As with live parasite infections, vaccination with irradiated parasites also conferred superior protection when mice were treated with 3H3 (**Fig. 2h**), which correlated with an enhanced MBC recall response, as evidenced by higher fold changes in serum antibody titers (**Fig. 2i**).

In summary, these data demonstrate that systemic 4-1BB ligation, while delaying the effector GC response, significantly enhances humoral immune memory. This enhancement in memory function appears to be independent of early antigen load and holds true even when attenuated parasites are used.

Enhanced humoral immune memory stems not from larger number but higher recall potential of MBCs.

To verify whether the higher fold change in parasite-specific antibodies on day 5 post *PbA* challenge was due to larger number of antibody secreting cells (ASC), we enumerated PCs in the bone marrow of *Py*-recovered 3H3- and rIgG-treated mice challenged with *PbA.* As expected, we observed significantly larger numbers of MSP-1 specific IgM$^+$ PCs in 3H3-treated mice (**Fig. 3a**). As MBCs are the precursors for PCs upon antigen re-exposure[10], we determined the frequency and number of MBCs (IgD$^-$CD38$^+$GL7$^-$B220$^+$) prior to *PbA* challenge on 90 dpi. Contrary to our expectation, we observed significantly lower numbers of both polyclonal and antigen-specific (MSP-1 specific) (**Supp Fig. 3a, b**) MBCs in 3H3-treated mice compared to rIgG-treated ones. Upon further characterization of the MBC pool using previously described markers such as CD73, CD80 and PD-L2[49,50] to identify functionally distinct subsets, we observed significantly lower number of polyclonal CD73$^+$CD80$^+$ (**Fig. 3b**), CD73$^+$PD-L2$^+$ and CD80$^+$PD-L2$^+$ (**Supp Fig. 3c, d**) MBCs. Evaluation of MSP-1 specific MBCs also showed reduced numbers of CD73$^+$PD-L2$^+$, CD80$^+$PD-L2$^+$ (**Supp Fig. 3e, f**) pools. MSP-1 specific MBCs within the CD73$^+$CD80$^+$ fraction was also lower in 3H3-treated mice (**Fig. 3c**). The double positive MBC fraction has been shown to readily differentiate into ASCs while the double negative pool has been shown to reseed a secondary GC response upon antigen reencounter[12, 50]. Given that we see a rapid increase in antibody titers in just 5 days of *PbA* challenge (**Fig. 2a**), it is likely that the PCs may have readily differentiated from the existing double positive MBCs.

Given that chemoprophylaxis using artemether preserved the protective effect of 3H3 (**Fig. 2e and f**), we asked whether the PC and MBC compartments showed similar trends to what was observed without artemether. To this end, mice were infected and treated as shown in **Fig. 2d** and splenic MBCs before *PbA* challenge as well as PCs in the bone marrow 5 days post challenge were characterized. We observed lower number of polyclonal (**Fig. 3d**) as well as the MSP-1 specific (**Fig. 3e**) CD73$^+$CD80$^+$ MBCs in 3H3-treated mice. Despite the reduced number of MBCs and in agreement with the survival data following *PbA* challenge (**Fig. 2e**), we observed a larger frequency of PCs that stained positive for Ki67 and TACI, indicative of a high proliferative burst (**Fig. 3f**). Taken together, these data suggest that, MBCs from 3H3-treated mice are qualitatively different and are capable to differentiate and proliferate into larger number of PCs upon antigen encounter. Since artemether treatment preserves the survival advantage and cellular phenotypes, we decided to adopt this regimen as a standard approach in future experiments. This will not only allow the mice to reach convalescence early (35-40 dpi with artemether compared to 85-90 dpi without), but will also simulate the chemoprophylactic approach often used in malaria-endemic regions.

Confirming our prediction that MBCs from 3H3-treated mice may be functionally superior, they showed a higher binding affinity to MSP-1 as evidenced by a higher geometric mean fluorescence intensity of MSP-1 staining (**Fig. 3g**). To formally compare the functional capacity of these MBCs upon antigen re-exposure, we flow-sorted MBCs from 3H3- and rIgG-treated mice and adoptively transferred them into separate groups of B cell deficient (μMT) recipients that were then infected with *Py* the following day. *Py*-infected μMT mice that received MBCs from 3H3-

389 treated donors exhibited lower parasite load on 12 dpi compared to those receiving MBCs from
390 rIgG-treated donors (**Fig. 3h**) and this correlated with higher serum antibody titers (**Fig. 3i**) on 4
391 dpi. Collectively, our data so far show that MBCs derived from 3H3-treated mice exhibit
392 intrinsically superior recall potential and capacity to differentiate into antibody secreting PCs.
393
394 **Exogenous 4-1BB ligation drives extrafollicular MBCs.**
395
To understand the genetic basis for the functional differences between MBCs from 3H3- and rIgG-
treated mice, we performed scRNA-, CITE- and V(D)J- sequencing on flow-sorted polyclonal
(B220$^+$IgD$^-$GL7$^-$CD38$^+$) and MSP-1 specific (B220$^+$IgD$^-$GL7$^-$CD38$^+$Decoy$^-$MSP1$^+$) MBCs (**Supp**
Fig. 4a) on day 38 post *Py* infection from both groups as detailed in **Fig. 4a.** As internal controls
to verify our workflow, resting B cells (B220$^+$IgD$^+$) from either group were also spiked in to
represent ~ 25% of all cells sequenced in their respective polyclonal fraction. As shown in the
UMAP (**Supp Fig. 4b**), the polyclonal MBCs grouped into 10 different clusters with the spiked-in
resting B cells (IgD$^+$) grouping separately in cluster 1 and representing roughly 25-27% of all cells
sequenced (**Supp Fig. 4c**). To better highlight the potential differences among different MBC
populations, polyclonal cells were re-clustered without the IgD$^+$ population (**Fig. 4b**) into 9
clusters. Notably, cluster 0 was significantly more represented in the 3H3 group, comprising 55%
of all MBCs compared to 28% in the rIgG group (**Fig. 4c**). This cluster exhibited increased
expression of molecules involved with B cell activation (*Cd79a*, *Cd79b* and *Cd40*)[51-53] and survival
(*Tnfrsf13b* and *Tnfrsf13c*)[54-57] (**Fig. 4d**). Cluster 2 was equally represented in both groups (19-
20%) (**Fig 4c**) and expressed genes associated with robust activation and recall potential such
as *Fcrl5*[58] and *Il9r*[59], respectively (**Fig. 4d**). In contrast, clusters 1 and 3 were reduced in the 3H3
group relative to the rIgG group (**Fig. 4c**) and were characterized by markers associated with
stemness (*Sox4*, *Myb*, *Il7r* and *Cd93*)[60-63] and genes associated with GC-derived MBCs (*Plxnb2*,
Basp1, *Scimp* and *Tox*)[64], respectively (**Fig. 4d**). Intriguingly, a closer look revealed that clusters
0 and 2, comprising around 75% of cells derived from 3H3-treated mice (**Fig. 4c**), exhibited a
gene signature (**Supp table. 1**) indicative of an extrafollicular (EF) origin[64] (**Fig. 4e and f**). This is
consistent with our data showing a higher recall potential of MBCs from 3H3-treated mice, as EF-
derived MBCs have been shown to be more proliferative and result in larger number of PCs[65].
Finally, clusters 4-8 that showed lower EF-derived signature comprised only 6% of MBCs in the
3H3 group compared to nearly 22% (**Fig. 4c**) in the rIgG group and contained mostly atypical B
cells (**Fig. 4d, e and f**).
To further investigate if 3H3 treatment programs antigen-specific MBCs in a similar
fashion, we analyzed sequences from MSP-1 specific cells. As shown in the UMAP (**Fig. 4g**), we
identified 5 clusters, with cluster 0 being overrepresented in the rIgG group (67%), compared to
only 28% in the 3H3 group (**Fig. 4h**) and marked by high expression of genes characteristic of
atypical MBCs, such as *Itgax*, *Itgam*, *Itgb2* and *Fgr*[66,67] (**Fig. 4i**). Conversely, cluster 1 comprised
71% of MSP-1 specific MBCs in the 3H3 group, but only 20% in the rIgG group (**Fig. 4h**) and
characterized by a combination of markers present in clusters 0 and 2 of the polyclonal pool (**Fig.**
4d and Fig. 4i). Cluster 2 was less represented in the 3H3 group (1%) compared to the rIgG
group (8%) and expressed markers associated with stemness, such as *Il7r*[68,69] and *Lef1*[70] (**Fig 4h**
and Fig 4i). Clusters 3 and 4 made up less than 6% in the rIgG group and less than 0.5% in the
3H3 group (**Fig 4h**). Much like clusters 0 and 2 in the polyclonal pool (**Fig. 4e**), cluster 1 of the
MSP-1 specific pool was enriched for an EF-derived signature (**Fig 4j**), which was more
pronounced in 3H3-group (**Fig. 4k**)**,** while cluster 0 (MSP-1 specific) from the rIgG group showed
the highest atypical signature (**Fig. 4j and k**). These data collectively demonstrate that MBCs
generated in mice following systemic 4-1BB stimulation is predominantly of EF origin, whereas
those from control mice resemble atypical MBCs.

3H3 treatment generates MBCs with a distinct transcriptional profile supporting enhanced recall.

Characterization of the major clusters revealed differences in the functional composition of the polyclonal and MSP-1 specific MBC pools from rIgG- and 3H3-treated mice. Importantly, cluster 0 in the polyclonal pool expressed the highest level of IgM (*Ighm*) (**Fig. 5a and Supp Fig. 5a**)**,** a hallmark of a highly plastic and rapidly responding MBCs following *Plasmodium* rechallenge[65]; while cluster 2 co-expressed CD73 (*Nt5e*)*,* CD80 (*Cd80*) (**Fig. 5b and Supp Fig. 5b**) and *Pdcd1lg2* (**Supp Fig. 5c**), markers indicative of superior recall[50]. Utilizing pseudotime inference using cluster 1 as the origin, the IgM+ population (cluster 0) appeared to be in the early stage of its differentiation process, while clusters 2, 4 and 5 were closer to being terminally differentiated (**Fig. 5c**). In contrast to MBCs from rIgG-treated mice (blue arrows, **Fig. 5c**), polyclonal MBCs from 3H3-treated mice favored the early and plastic stage[65] (cluster 0, IgM+) (thick red arrow, **Fig. 5c**) and had their differentiation skewed away from the terminal atypical trajectories (thin red arrows, **Fig. 5c**). This will result in the preferential accumulation of IgM+ EF MBCs and reduction of atypical MBCs in the 3H3 group (**Fig. 4c**). Characterization of MSP-1 specific pool showed that cluster 1 from 3H3-treated mice co-expressed high levels of IgM (*Ighm*) (**Fig. 5d and Supp Fig. 5d**), CD73 (*Nt5e*), CD80 (*Cd80*) (**Fig. 5e and Supp Fig. 5e**) and *Pdcd1lg2* (**Supp Fig. 5f**). Pseudotime analysis using cluster 2 as the origin showed that MBCs from 3H3-treated mice shifted their differentiation towards the EF trajectory (thick red arrow, **Fig. 5f**) and away from the atypical trajectory (thin red arrow, **Fig. 5f**). While MBCs from the rIgG group showed a higher propensity to take the atypical trajectory, away from the EF route (blue arrow, **Fig. 5f**). The data so far showing that MBCs from 3H3-treated mice are more likely to have an EF origin is further corroborated by our V(D)J sequencing data showing they have significantly lower mutational frequency in their BCRs (**Supp Fig. 5g and h**).

To explore functional differences within the EF clusters, we analyzed differentially expressed genes (DEGs) between rIgG and 3H3 in clusters 0 (IgM+) and 2 (CD73+CD80+) from the polyclonal pool as well as cluster 1 from the MSP-1 specific pool **(Supp. Table 2)**. Notably, 122 DEGs were upregulated across all 3 clusters (**Supp**. **Fig 5i**). Approximately 88% of polyclonal cluster 2 DEGs were shared with polyclonal cluster 0 and/or MSP1+ cluster 1, with only 21 unique genes. In addition, more than half of the DEGs in polyclonal cluster 0 and MSP-1 specific cluster 1 were shared, underscoring the similarities between these two clusters (**Supp**. **Fig 5i**). Given that most DEGs in polyclonal cluster 2 overlapped with the other 2 clusters (**Supp**. **Fig 5i**), we further investigated the biological processes regulated by the combined DEGs in polyclonal clusters 0 and 2 (**Fig. 5g**) and MSP-1 specific cluster 1 (**Fig. 5h**). Notably, both polyclonal and MSP-1 specific MBCs from 3H3 group showed increased expression of genes associated with heightened capacity for activation, proliferation and antigen presentation as well as certain metabolic pathways, possibly to meet the energetic and biomass demands to support these processes (**Fig 5g and Fig 5h**). In addition, MBCs from 3H3-treated mice also showed enhanced cellular response to IL-4, a cytokine shown to downregulate GC related genes and enhance MBC generation and activation[71,72], which may enforce the MBCs to take the extrafollicular route. Aligning with these data, both the polyclonal (clusters 0 and 2, **Supp Fig. 5j**) and MSP-1 specific MBCs (cluster 1, **Supp Fig. 5k**) also downregulated genes involved in pathways related to increased apoptosis, reduced transcription factor binding, gene silencing and metabolism among others. Collectively, these data confirm that exogenous ligation of 4-1BB results in the generation of MBCs with an EF origin that are, although numerically fewer, functionally superior.

3H3-treatment induced protection is dependent on IFNγ and B cell intrinsic T-BET:IL-9R axis.

Previous studies have highlighted the role of B cell intrinsic IFNγ signaling and T-BET expression [73-75] for enhanced MBC recall and PC differentiation. Importantly, exogenous stimulation of 4-1BB was shown to result in higher levels of IFNγ secretion by CD4 T cells[76]. When splenocytes were stimulated in the presence of PMA/Ionomycin and brefeldin A, we observed that CD4 T cells derived from 3H3-treated mice were indeed capable of secreting higher levels of IFNγ (**Supp Fig 6a**). Additionally, B cells cultured *ex vivo* with IFNγ showed a significant increase in T-BET expression within 18 hours (**Supp Fig. 6b**). In line with this, we also observed significantly higher expression of T-BET in activated (IgD⁻) B cells derived from 3H3-treated mice around 15 dpi (**Supp Fig. 6c**). T-BET expression in B cells during the effector phase was shown to enhance protection from rechallenge by mediating class-switching to IgG2c antibody isotype[73]. Consistent with this finding, we observed higher increases in IgG2c levels following *PbA* challenge in 3H3-treated mice (**Supp. Fig. 6d**). Given these data, we asked whether 3H3-mediated superior protection is IFNγ dependent. To test this, mice were infected with *Py* and treated with 3H3 or rIgG and IFNγ was blocked using IFNγ blocking antibody (XMG1.2) on indicated timepoints. Mice were then challenged with *PbA* (around 90 dpi) and survival was monitored (**Supp Fig. 6e**). As expected, 3H3-treated mice lost protection following *PbA* challenge when IFNγ was blocked (**Supp Fig. 6f**). Given the link between IFNγ and T-BET, we then asked whether B cell intrinsic T-BET expression is necessary for 3H3-mediated protection. To this end, we generated mixed bone marrow chimeras in which T-BET deficiency was restricted to the B cell compartment (B^Tbx21-/-) (**Supp Fig. 6g**) and infected them with *Py,* followed by 3H3 treatment. Similar to our results from XMG1.2 treatment, *PbA* challenge of convalescent B^Tbx21-/- mice showed significantly reduced survival compared to B^WT mice (**Supp Fig. 6h**). Taken together, these highlight the importance of IFNγ:T-BET axis during *Plasmodium* infection and show that the enhanced protection observed in mice subjected to exogenous 4-1BB ligation is dependent on this axis.

IL-9 receptor (IL-9R) signaling on MBCs has also been shown to result in superior recall[59]. In line with this, MSP-1 specific MBCs from 3H3-treated mice subjected to IFNγ blockade and B^Tbx21-/- mice showed significantly lower levels of IL-9R expression (**Supp Fig. 6i and j**) suggesting that B cell intrinsic IFNγ:T-BET axis is key to this phenomenon. To investigate the contribution of IL-9R to higher MBC recall, we searched for distinct *Il9r*-expressing cluster(s) in our scRNA-seq data and identified polyclonal (cluster 2) (**Fig. 6a**) and MSP-1 specific (cluster 1) (**Fig. 6b**) which also co-expressed other signature MBC markers such *as Nt5e, Cd80, Pdcd1lg2* among others (**Fig. 4d**). Quantitative real-time PCR analysis on flow-sorted MBCs from 3H3-treated mice also showed significantly higher levels of *Il9r* expression compared to those from rIgG-treated mice (**Supp. Fig 7a**). In an attempt to draw parallels among populations from our transcriptomic and flow cytometry datasets, we clustered our multi-parameter flow cytometry data in a UMAP space (polyclonal, **Fig. 6c** and MSP-1 specific, **Fig. 6f**). As shown in **Fig. 6c** and **d,** the polyclonal pool clustered into 9 populations (*A-I*), of which populations *D* and *I* denoted the MSP-1 specific pool. While populations *E, F* and *I* expressed IL-9R, population E failed to express key functional markers such as CD73 and CD80 making it less likely to be involved in an immediate recall following antigen re-exposure (**Fig. 6d** and **Supp Fig. 7b**). Importantly and in agreement with this, we observed that the CD73⁺CD80⁺ MBC population from 3H3-treated mice expressed significantly higher levels of IL-9R compared to controls (**Fig. 6e**). These data also suggested that population *I* may be functionally similar to population *F* and together they may represent antigen-specific MBC pools that are swiftly recalled following antigen-reencounter. Phenotypic dissimilarity of populations *D* and *I* also highlights the functional heterogeneity within a given

antigen-specific pool. This was confirmed by further clustering the MSP-1 specific pool which yielded 5 different populations, mostly differing in their expression of key functional MBC markers (**Fig. 6f and g**). As with the polyclonal cells (**Fig.6e**), MSP-1 specific MBCs from 3H3-treated mice also expressed higher levels of IL-9R (**Fig. 6h**) suggestive of their superior recall potential. Given the parallels between the gene and surface maker expression data, polyclonal cluster 2 may be identical to population *F* and they may represent MBCs equipped to readily engage in a recall response. Given the dependency on IFNγ and B cell intrinsic T-BET expression for heightened survival of 3H3-treated mice, we hypothesized that IL-9R expression may also be regulated by this axis. To test this, B cells from naïve mice were activated *ex vivo* in the presence or absence of IFNγ and IL-9R expression was evaluated after 3 days in culture. Although activation of B cells alone resulted in increase in IL-9R expression, it was further increased when IFNγ was added to the culture (**Supp Fig. 7c**).

To investigate whether the enhanced protection following *PbA* challenge in 3H3-treated mice is dependent on IL-9:IL-9R signaling, we blocked IL-9 in convalescent mice prior to *PbA* challenge and monitored survival (**Supp Fig. 7d)**. As shown in **Fig**. **6i**, the enhanced protection observed in 3H3-treated mice was reversed in mice treated with IL-9 blocking antibody compared to controls; rIgG-treated mice did not show any difference in survival. Supporting our observation, 3H3-treated mice subjected to IL-9 neutralization exhibited a blunted antibody recall following *PbA* challenge (**Fig. 6j**), indicating that IL-9 signaling is important for 3H3-mediated protection. Next, we interrogated whether B cell-intrinsic IL-9R expression is important for the enhanced protection following 3H3 treatment. To accomplish this, we generated competitive mixed bone marrow chimeras in which IL-9R deficiency is restricted to B cells (B$^{Il9r-/-}$) (**Supp Fig. 7e**). Since we did not see any difference in the rIgG treated mice following IL-9 blockade (**Fig. 6i**), we confined our experiment to 3H3-treatment in B$^{Il9r-/-}$ and B^{WT} mice. On 38 dpi, mice were challenged with *PbA* and survival was monitored. Notably, we observed that B$^{Il9r-/-}$ mice showed significantly shorter survival than B^{WT} mice following *PbA* challenge (**Fig. 6k**) and a dampened MBC recall response as evident from the lower fold change in serum antibody titers (**Fig. 6l**). These data show that the MBC recall following *PbA* challenge in 3H3-treated mice is dependent on the B cell-intrinsic IL-9R expression. Collectively, our data highlight a previously underappreciated IFNγ:T-BET:IL-9R axis present in MBCs following *Plasmodium* infection, which could potentially be modulated to enhance humoral immune memory.

Exogenous 4-1BB ligation poises MBCs for swift recall by altering their epigenome.

Given that both polyclonal and MSP-1 specific MBCs from 3H3-treated mice downregulated genes associated with transcriptional silencing and regulation of transcription factor activity (**Supp. Fig 5j and k**), we wanted to determine whether these MBCs are imprinted with an epigenetic signature supporting superior recall. To this end, we flow-sorted polyclonal and MSP-1 specific MBCs from 3H3- or rIgG-treated mice (**Fig. 7a**) as before and performed scATAC-seq. Initial clustering based on transposase activity populated 4 different clusters in the MSP-1 specific cells (**Supp. Fig 8a**) and 6 clusters in the polyclonal pool (**Supp. Fig 8b**). Given the importance of B cell-intrinsic IL-9R expression in mice treated with 3H3, we wanted to identify the IL-9R expressing population in our scATAC-seq dataset and examine the accessibility of genes that can potentiate PC differentiation. Thus, we integrated our scRNA-seq (from **Fig. 4 and 5**) with the scATAC-seq datasets and used the transferred labels to annotate the ATAC clusters from MSP1-specific (**Fig. 7b**) and polyclonal MBCs (**Supp. Fig. 8c**). As shown in **Fig. 7b**, following integration, most cells fell in either cluster 0 or 1. Since cluster 1 in MSP-1 specific MBCs expressed the highest level of *Il9r* (**Fig. 6b**), we examined whether the *Il9r* gene locus in this cluster is still

accessible. As shown in **Fig. 7c**, we observed significantly higher accessibility in the *Il9r* locus in
cluster 1 compared to cluster 0, showing that MBCs in cluster 1 are imprinted with an epigenome
that favors higher *Il9r* expression. In line with our prediction that IL-9R expressing MBCs are
primed to quickly differentiate into PCs, cluster 1 also showed enhanced accessibility for *Prdm1*
locus (**Fig. 7d**) as well as for key transcription factors (TF) binding motifs (XBP1, ATF3, MYC,
ATF6, KLF6 and BHLHE40) (**Fig. 7e**) associated with enhanced PC differentiation. Importantly
and similar to cluster 1 in the MSP-1 specific pool, cluster 2 in the polyclonal pool also showed
significantly enhanced *Il9r* locus accessibility (**Supp Fig. 8d**) as well as markers of higher
activation potential (*Fclr5*, **Supp Fig. 8e**), enhanced PC differentiation (*Atf6*, **Supp. Fig 8f**) and
antiapoptotic signaling (*Bcl2*, **Supp. Fig. 8g**). In addition, TF binding motifs important in PC
differentiation (KLF4, ATF3, XBP1 and IRF4) as well as cell proliferation (MYC) and IFNγ signaling
(IRF1) were also accessible in cluster 2 from the polyclonal pool (**Supp Fig.8h**). Given that *Il9r*
expression correlates with enhanced protection in 3H3-treated mice, we compared the chromatin
accessibility of *Il9r*-expressing MBCs between 3H3- and rIgG-treated groups. In strong agreement
with our survival and phenotypic data, MBCs from 3H3-treated mice in cluster 1 not only showed
continued accessibility for *Il9r* locus (**Fig. 7f**), but were also enriched for the key transcription
factor binding motifs associated with enhanced PC differentiation and proliferation (**Fig. 7g**).
Notably, we also observed that T-BET and IRF1 binding motifs also showed higher accessibility
further reinforcing our finding that the superior recall we observe in 3H3-treated mice is dependent
on IFNγ:T-BET:IL-9R axis.
602
603 **Discussion**
604
In this study, we investigated the effects of exogenous 4-1BB ligation and observed an
unexpected outcome in MBC generation in an experimental mouse model of malaria. Despite the
disruption of the effector GC response and a total reduction in MBC numbers, the MBC recall
response was significantly enhanced, providing protection upon reinfection. While our data draw
some parallels with a previous study using a viral infection model [77] showing a derailed GC
response, our findings on humoral immune memory response are unique. Single-cell
transcriptomic analysis of the MBCs showed a predominant extrafollicular rather than a GC-
derived signature, with IL-9R emerging as a key determinant for the enhanced recall. Using a
combination of genetic, chimeric, and biochemical approaches, we identify a previously
unappreciated axis comprising IFNγ and B cell intrinsic T-BET expression during the effector
phase, potentially driving the expression of IL-9R. Sequencing for transposase-accessible
chromatin regions in flow-sorted MBCs from *Plasmodium*-infected mice treated with exogenous
4-1BB agonist also revealed an epigenomic landscape that primed them for rapid differentiation
into PCs.
Similar to our study, *Ehrlichia muris (E. muris)* infection alone without the need of any
agonistic antibody to 4-1BB, resulted in a derailed GC response[78]. Notably, the MBCs that
appeared in *E. muris* infection were CD11c[+] and T-dependent but seldom expressed GC-related
markers[79]. Thus, the antigen-specific IgM[+] MBCs (cluster 1) in our study are largely similar to
MBCs mentioned in the study above[79], but express FCLR5 and not CD11c. Co-expression of
CD11c, CD11b and FCLR5 are considered markers for atypical MBCs, with predominantly
inhibitory functions [80-82]. Based on the lack of CD11b and CD11c expression combined with the
expression of various activation markers (**Fig. 4 and 6**) including FCLR5[58], we contend that the
IgM[+] MBCs from 3H3-treated mice (cluster 1) are different from the inhibitory atypical MBCs.
IFNγ-driven T-BET expression was shown to be pivotal to increased protection following
Plasmodium rechallenge[73]. Reliance on IFNγ:T-BET axis in 3H3-treated mice for enhanced recall,
along with a pronounced EF signature of these MBCs, is in agreement with previous findings

showing that B cell-intrinsic IFNγ signaling reinforces EF B cell responses[83]. Previous studies
have also independently shown the role of IL-9R expression for enhanced recall[59]. 3H3-treated
but not rIgG-treated mice, seem to depend on IL-9R expression specifically on B cells for
protection, which may be explained by the far lower IL-9R expression in MBCs derived from rIgG-
treated mice. Our scATAC-seq data shows that IRF1 binding motifs are readily accessible in *Il9r*
expressing MBCs from both groups showing responsiveness to IFNγ[84]. More importantly, MBCs
from 3H3-treated mice have a more pronounced accessibility to this region along with T-BET
binding motifs. This further strengthens the possibility that IFNγ-driven T-BET may influence the
expression of IL-9R in MBCs from 3H3-treated mice. We also observed lower IL-9R levels in B
cells lacking T-BET expression or when IFNγ was blocked, which is complemented by our finding
that IL-9R expression can be enhanced in B cells when subjected to IFNγ *ex vivo*. Together, these
data suggest a potential role of IFNγ:T-BET axis in IL-9R expression and raises the possibility that
T-BET levels may have to reach a specific threshold to enforce IL-9R expression. While these
observations are B cell-centric, the source of IL-9 during the memory phase to signal via IL-9R
remains to be investigated. As such, stimulation of various TNFRSF members have been shown
to promote IL-9 secretion from CD4 T cells in multiple experimental settings[85,86-88]. Although our
IL-9 neutralization experiments suggest the presence of an IL-9 source prevalent during the
memory phase, a direct determination of the identity of the cellular source is yet to be done. IL-9
neutralization during the memory phase compromised survival and MBC recall only in 3H3-treated
mice, signifying the role of IL-9:IL-9R signaling following 4-1BB stimulation. To our knowledge,
our study remains the first of its kind to link the IFNγ:T-BET axis with IL-9R in MBC recall. Despite
mining numerous publicly available human malaria datasets studying MBCs, we were unable to
find a high IL-9R expressing population that could be correlated with enhanced protection upon
pathogen re-exposure. This is due to the fact that none of these studies involved the use of
immunomodulatory approaches that potentially drive high T-BET in B cells, which may be key to
driving IL-9R. Indeed, our own *ex vivo* data with murine B cells stimulated with IFNγ resulted in
the upregulation of T-BET first and IL-9R later (T-BET expression preceding IL-9R). A fate
mapping approach capable of tracking T-BET expression would be the ideal strategy to directly
ask this question, but it remains beyond the scope of this manuscript.

Although many studies have focused on the effect of 4-1BB on CD8 T cells[89,90], systemic
ligation of 4-1BB has been shown to reprogram CD4 T cells in a virus infection setting as well[76].
It is unlikely that 4-1BB stimulation on CD8 T cells may play a significant role in our studies, as
CD8 T cell depletion in 3H3-treated mice still disrupted the effector germinal center (GC) response
and resulted in substantial loss of parasite control, similar to observed in CD8 T cell replete WT
mice (*Caloba and Vijay unpublished data*). B cells are also known to express 4-1BB[91], but we did
not explore the role of 4-1BB stimulation on B cells as CD4-specific 4-1BB$^{-/-}$ bone marrow
chimeras failed to respond to 3H3 treatment as it did not derail the parasite control. These findings
indicate that in rodent malaria models, the effects of exogenous 4-1BB stimulation are dependent
on CD4 T cell intrinsic 4-1BB expression.

Here we provide data showing that exogenous 4-1BB stimulation is capable of programing
a predominantly EF B cell response that generates antigen-specific MBCs with superior recall
potential. Importantly, although the MBCs in mice that received 4-1BB stimulation is at a numerical
disadvantage, a large fraction of those cells possess a distinct epigenetic landscape and
transcriptional signature poising them for swift and efficient recall. Single-cell trajectory analysis
also reveals that MBCs from 3H3-treated mice have a tendency to stay more plastic within the
extrafollicular space. Our data also brings up the argument that GC response during effector
phase cannot always be a predictor for the ensuing memory response. We show that 4-1BB is a
compelling target for host-directed immunotherapy in infections that are reliant on MBC-driven
antibody response for protection. While the widescale use of a monoclonal antibody as an

immunotherapy may be cost prohibitive, the prospect of synthesizing small molecule activators of
4-1BB for achieving potent humoral memory immune activation and thus durable and long-lasting
protection may be practical.
683
684

Acknowledgements

The authors thank all past and present members of the Vijay Lab for the helpful dicussions and
technical assistance. We thank Drs. Fabio Re (RFUMS) and Danielle Stanisic (Griffith University)
for the critical review of the manuscript. We thank the RFUMS flow cytometry facility for the
instrumentation as well as the Biological Research Facility for animal husbandry and
maintenance. We thank Dr. Stefan Green, Kevin Kunstman, Edith Perez and Ping Li at the
Genomics and Microbiome Core Facility at Rush University for help with sequencing. This work
was supported by the RFUMS startup funds and Schweppe Scholar Award to R.V and the AAI
2024 Careers in Immunology Fellowship to C.C. and R.V.

Author Contributions
696

R.V. conceived the study. C.C. and R.V. designed, executed, analyzed and wrote the manuscript.
C.C. also performed the bioinformatic data analyses. A.J.S. executed experiments. T.A.L., L.J.,
and A.R., provided technical assistance. A.M.M and S.E.L. generated and provided reagents,
T.H.W. provided reagents as well as provided intellectual input, A.M.C. and M.H.K. generated
and provided reagents, M.H.K. provided intellectual input, J.P.W. provided technical assistance
for bioinformatic analyses.

References

1.   WHO World malaria report 2023. Geneva: World Health Organization;2023.
2.   Cowman, A.F., Healer, J., Marapana, D., and Marsh, K. (2016). Malaria: Biology and Disease. Cell *167*, 610-624. 10.1016/j.cell.2016.07.055.
3.   Laurens, M.B. (2020). RTS,S/AS01 vaccine (Mosquirix): an overview. Hum Vaccin Immunother *16*, 480-489. 10.1080/21645515.2019.1669415.
4.   MacLennan, I.C. (1994). Germinal centers. Annual review of immunology *12*, 117-139. 10.1146/annurev.iy.12.040194.001001.
5.   Cohen, S., Mc, G.I., and Carrington, S. (1961). Gamma-globulin and acquired immunity to human malaria. Nature *192*, 733-737. 10.1038/192733a0.
6.   Liu, Y.J., Malisan, F., de Bouteiller, O., Guret, C., Lebecque, S., Banchereau, J., Mills, F.C., Max, E.E., and Martinez-Valdez, H. (1996). Within germinal centers, isotype switching of immunoglobulin genes occurs after the onset of somatic mutation. Immunity *4*, 241-250. 10.1016/s1074-7613(00)80432-x.
7.   Jacob, J., Kelsoe, G., Rajewsky, K., and Weiss, U. (1991). Intraclonal generation of antibody mutants in germinal centres. Nature *354*, 389-392. 10.1038/354389a0.
8.   Berek, C., Berger, A., and Apel, M. (1991). Maturation of the immune response in germinal centers. Cell *67*, 1121-1129. 10.1016/0092-8674(91)90289-b.
9.   Nutt, S.L., Hodgkin, P.D., Tarlinton, D.M., and Corcoran, L.M. (2015). The generation of antibody-secreting plasma cells. Nat Rev Immunol *15*, 160-171. 10.1038/nri3795.
10.   Inoue, T., and Kurosaki, T. (2024). Memory B cells. Nat Rev Immunol *24*, 5-17. 10.1038/s41577-023-00897-3.
11.   Good-Jacobson, K.L., and Shlomchik, M.J. (2010). Plasticity and heterogeneity in the generation of memory B cells and long-lived plasma cells: the influence of germinal center interactions and dynamics. Journal of immunology *185*, 3117-3125. 10.4049/jimmunol.1001155.
12.   Kurosaki, T., Kometani, K., and Ise, W. (2015). Memory B cells. Nat Rev Immunol *15*, 149-159. 10.1038/nri3802.
13.   Tran, T.M., Li, S., Doumbo, S., Doumtabe, D., Huang, C.Y., Dia, S., Bathily, A., Sangala, J., Kone, Y., Traore, A., et al. (2013). An intensive longitudinal cohort study of Malian children and adults reveals no evidence of acquired immunity to Plasmodium falciparum infection. Clin Infect Dis *57*, 40-47. 10.1093/cid/cit174.
14.   Butler, N.S., Moebius, J., Pewe, L.L., Traore, B., Doumbo, O.K., Tygrett, L.T., Waldschmidt, T.J., Crompton, P.D., and Harty, J.T. (2011). Therapeutic blockade of PD-L1 and LAG-3 rapidly clears established blood-stage Plasmodium infection. Nat Immunol *13*, 188-195. 10.1038/ni.2180.
15.   Kurup, S.P., Obeng-Adjei, N., Anthony, S.M., Traore, B., Doumbo, O.K., Butler, N.S., Crompton, P.D., and Harty, J.T. (2017). Regulatory T cells impede acute and long-term immunity to blood-stage malaria through CTLA-4. Nat Med *23*, 1220-1225. 10.1038/nm.4395.
16.   Zander, R.A., Obeng-Adjei, N., Guthmiller, J.J., Kulu, D.I., Li, J., Ongoiba, A., Traore, B., Crompton, P.D., and Butler, N.S. (2015). PD-1 Co-inhibitory and OX40 Co-stimulatory Crosstalk Regulates Helper T Cell Differentiation and Anti-Plasmodium Humoral Immunity. Cell Host Microbe *17*, 628-641. 10.1016/j.chom.2015.03.007.
17.   Zander, R.A., Vijay, R., Pack, A.D., Guthmiller, J.J., Graham, A.C., Lindner, S.E., Vaughan, A.M., Kappe, S.H.I., and Butler, N.S. (2017). Th1-like Plasmodium-Specific Memory CD4(+) T Cells Support Humoral Immunity. Cell Rep *21*, 1839-1852. 10.1016/j.celrep.2017.10.077.
18.   Illingworth, J., Butler, N.S., Roetynck, S., Mwacharo, J., Pierce, S.K., Bejon, P., Crompton, P.D., Marsh, K., and Ndungu, F.M. (2013). Chronic exposure to Plasmodium

falciparum is associated with phenotypic evidence of B and T cell exhaustion. Journal of
immunology *190*, 1038-1047. 10.4049/jimmunol.1202438.
19.   Dawicki, W., Bertram, E.M., Sharpe, A.H., and Watts, T.H. (2004). 4-1BB and OX40 act
independently to facilitate robust CD8 and CD4 recall responses. Journal of immunology *173*,
5944-5951. 10.4049/jimmunol.173.10.5944.
20.   Dawicki, W., and Watts, T.H. (2004). Expression and function of 4-1BB during CD4
versus CD8 T cell responses in vivo. Eur J Immunol *34*, 743-751. 10.1002/eji.200324278.
21.   Kwon, B.S., Hurtado, J.C., Lee, Z.H., Kwack, K.B., Seo, S.K., Choi, B.K., Koller, B.H.,
Wolisi, G., Broxmeyer, H.E., and Vinay, D.S. (2002). Immune responses in 4-1BB (CD137)-
deficient mice. Journal of immunology *168*, 5483-5490. 10.4049/jimmunol.168.11.5483.
22.   Vijay, R., Guthmiller, J.J., Sturtz, A.J., Surette, F.A., Rogers, K.J., Sompallae, R.R., Li,
F., Pope, R.L., Chan, J.A., de Labastida Rivera, F., et al. (2020). Infection-induced plasmablasts
are a nutrient sink that impairs humoral immunity to malaria. Nat Immunol *21*, 790-801.
10.1038/s41590-020-0678-5.
23.   Malleret, B., Claser, C., Ong, A.S., Suwanarusk, R., Sriprawat, K., Howland, S.W.,
Russell, B., Nosten, F., and Renia, L. (2011). A rapid and robust tri-color flow cytometry assay
for monitoring malaria parasite development. Sci Rep *1*, 118. 10.1038/srep00118.
24.   Villarino, N.F., LeCleir, G.R., Denny, J.E., Dearth, S.P., Harding, C.L., Sloan, S.S.,
Gribble, J.L., Campagna, S.R., Wilhelm, S.W., and Schmidt, N.W. (2016). Composition of the
gut microbiota modulates the severity of malaria. Proc Natl Acad Sci U S A *113*, 2235-2240.
10.1073/pnas.1504887113.
25.   Johnson, J.T., Surette, F.A., Ausdal, G.R., Shah, M., Minns, A.M., Lindner, S.E., Zander,
R.A., and Butler, N.S. (2024). CD4 T Cell-Derived IL-21 Is Critical for Sustaining Plasmodium
Infection-Induced Germinal Center Responses and Promoting the Selection of Memory B Cells
with Recall Potential. Journal of immunology *212*, 1467-1478. 10.4049/jimmunol.2300683.
26.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and species. Nat
Biotechnol *36*, 411-420. 10.1038/nbt.4096.
27.   Choudhary, S., and Satija, R. (2022). Comparison and evaluation of statistical error
models for scRNA-seq. Genome Biol *23*, 27. 10.1186/s13059-021-02584-9.
28.   Ahlmann-Eltze, C., and Huber, W. (2021). glmGamPoi: fitting Gamma-Poisson
generalized linear models on single cell count data. Bioinformatics *36*, 5701-5702.
10.1093/bioinformatics/btaa1009.
29.   Xu, S., Hu, E., Cai, Y., Xie, Z., Luo, X., Zhan, L., Tang, W., Wang, Q., Liu, B., Wang, R.,
et al. (2024). Using clusterProfiler to characterize multiomics data. Nature protocols *19*, 3292-
3320. 10.1038/s41596-024-01020-z.
30.   Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J.,
Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate
decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol *32*, 381-386.
10.1038/nbt.2859.
31.   Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).
Reversed graph embedding resolves complex single-cell trajectories. Nat Methods *14*, 979-982.
10.1038/nmeth.4402.
32.   Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos,
S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional landscape of
mammalian organogenesis. Nature *566*, 496-502. 10.1038/s41586-019-0969-x.
33.   Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden:
guaranteeing well-connected communities. Sci Rep *9*, 5233. 10.1038/s41598-019-41695-z.

34.   Levine, J.H., Simonds, E.F., Bendall, S.C., Davis, K.L., Amir el, A.D., Tadmor, M.D.,
Litvin, O., Fienberg, H.G., Jager, A., Zunder, E.R., et al. (2015). Data-Driven Phenotypic
Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. Cell *162*, 184-
197. 10.1016/j.cell.2015.05.047.
35.   Neuwirth, E. (2022). RColorBrewer: ColorBrewer Palettes.
36.   Planche, T., Dzeing, A., Emmerson, A.C., Onanga, M., Kremsner, P.G., Engel, K.,
Kombila, M., Ngou-Milama, E., and Krishna, S. (2002). Plasma glutamine and glutamate
concentrations in Gabonese children with Plasmodium falciparum infection. QJM *95*, 89-97.
10.1093/qjmed/95.2.89.
37.   Marsh, S. (2021). scCustomize: Custom Visualizations & Functions for Streamlined
Analyses of Single Cell Sequencing.
38.   Garnier, S., Ross, Noam, Rudis, Robert, Camargo, Pedro A, Sciaini, Marco, Scherer,
Cédric (2024). Colorblind-Friendly Color Maps for R. 10.5281/zenodo.4679423.
39.   Gupta, N.T., Vander Heiden, J.A., Uduman, M., Gadala-Maria, D., Yaari, G., and
Kleinstein, S.H. (2015). Change-O: a toolkit for analyzing large-scale B cell immunoglobulin
repertoire sequencing data. Bioinformatics *31*, 3356-3358. 10.1093/bioinformatics/btv359.
40.   Nouri, N., and Kleinstein, S.H. (2018). A spectral clustering-based method for identifying
clones from high-throughput B cell repertoire sequencing data. Bioinformatics *34*, i341-i349.
10.1093/bioinformatics/bty235.
41.   Nouri, N., and Kleinstein, S.H. (2020). Somatic hypermutation analysis for improved
identification of B cell clonal families from next-generation sequencing data. PLoS Comput Biol
16, e1007977. 10.1371/journal.pcbi.1007977.
42.   Gupta, N.T., Adams, K.D., Briggs, A.W., Timberlake, S.C., Vigneault, F., and Kleinstein,
S.H. (2017). Hierarchical Clustering Can Identify B Cell Clones with High Confidence in Ig
Repertoire Sequencing Data. Journal of immunology *198*, 2489-2499.
10.4049/jimmunol.1601850.
43.   Hoehn, K.B., Pybus, O.G., and Kleinstein, S.H. (2022). Phylogenetic analysis of
migration, differentiation, and class switching in B cells. PLoS Comput Biol *18*, e1009885.
10.1371/journal.pcbi.1009885.
44.   Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York.
45.   Stuart, T., Srivastava, A., Madad, S., Lareau, C.A., and Satija, R. (2021). Single-cell
chromatin state analysis with Signac. Nat Methods *18*, 1333-1341. 10.1038/s41592-021-01282-
5.
46.   Fornes, O., Castro-Mondragon, J.A., Khan, A., van der Lee, R., Zhang, X., Richmond,
P.A., Modi, B.P., Correard, S., Gheorghe, M., Baranasic, D., et al. (2020). JASPAR 2020:
update of the open-access database of transcription factor binding profiles. Nucleic Acids Res
48, D87-D92. 10.1093/nar/gkz1001.
47.   Wagih, O. (2024). ggseqlogo: A 'ggplot2' Extension for Drawing Publication-Ready
Sequence Logos. https://CRAN.R-project.org/package=ggseqlogo.
48.   Arch, R.H., and Thompson, C.B. (1998). 4-1BB and Ox40 are members of a tumor
necrosis factor (TNF)-nerve growth factor receptor subfamily that bind TNF receptor-associated
factors and activate nuclear factor kappaB. Mol Cell Biol *18*, 558-565. 10.1128/MCB.18.1.558.
49.   Brown, S.L., Bauer, J.J., Lee, J., Ntirandekura, E., and Stumhofer, J.S. (2022). IgM(+)
and IgM(-) memory B cells represent heterogeneous populations capable of producing class-
switched antibodies and germinal center B cells upon rechallenge with P. yoelii. Journal of
leukocyte biology *112*, 1115-1135. 10.1002/JLB.4A0921-523R.
50.   Zuccarino-Catania, G.V., Sadanand, S., Weisel, F.J., Tomayko, M.M., Meng, H.,
Kleinstein, S.H., Good-Jacobson, K.L., and Shlomchik, M.J. (2014). CD80 and PD-L2 define

functionally distinct memory B cell subsets that are independent of antibody isotype. Nat
Immunol *15*, 631-637. 10.1038/ni.2914.
51.   Huse, K., Bai, B., Hilden, V.I., Bollum, L.K., Vatsveen, T.K., Munthe, L.A., Smeland,
E.B., Irish, J.M., Walchli, S., and Myklebust, J.H. (2022). Mechanism of CD79A and CD79B
Support for IgM+ B Cell Fitness through B Cell Receptor Surface Expression. Journal of
immunology *209*, 2042-2053. 10.4049/jimmunol.2200144.
52.   Tkachenko, A., Kupcova, K., and Havranek, O. (2023). B-Cell Receptor Signaling and
Beyond: The Role of Igalpha (CD79a)/Igbeta (CD79b) in Normal and Malignant B Cells. Int J
Mol Sci *25*. 10.3390/ijms25010010.
53.   Carpenter, E.L., Mick, R., Ruter, J., and Vonderheide, R.H. (2009). Activation of human
B cells by the agonist CD40 antibody CP-870,893 and augmentation with simultaneous toll-like
receptor 9 stimulation. J Transl Med *7*, 93. 10.1186/1479-5876-7-93.
54.   Luo, W., Weisel, F., and Shlomchik, M.J. (2018). B Cell Receptor and CD40 Signaling
Are Rewired for Synergistic Induction of the c-Myc Transcription Factor in Germinal Center B
Cells. Immunity *48*, 313-326 e315. 10.1016/j.immuni.2018.01.008.
55.   He, B., Santamaria, R., Xu, W., Cols, M., Chen, K., Puga, I., Shan, M., Xiong, H.,
Bussel, J.B., Chiu, A., et al. (2010). The transmembrane activator TACI triggers immunoglobulin
class switching by activating B cells through the adaptor MyD88. Nat Immunol *11*, 836-845.
10.1038/ni.1914.
56.   Eslami, M., Schuepbach-Mallepell, S., Diana, D., Willen, L., Kowalczyk-Quintas, C.,
Desponds, C., Peter, B., Vigolo, M., Renevey, F., Donze, O., et al. (2024). Unique and
redundant roles of mouse BCMA, TACI, BAFF, APRIL, and IL-6 in supporting antibody-
producing cells in different tissues. Proc Natl Acad Sci U S A *121*, e2404309121.
10.1073/pnas.2404309121.
57.   Lau, A.W.Y., Turner, V.M., Bourne, K., Hermes, J.R., Chan, T.D., and Brink, R. (2021).
BAFFR controls early memory B cell responses but is dispensable for germinal center function.
The Journal of experimental medicine *218*. 10.1084/jem.20191167.
58.   Kim, C.C., Baccarella, A.M., Bayat, A., Pepper, M., and Fontana, M.F. (2019). FCRL5(+)
Memory B Cells Exhibit Robust Recall Responses. Cell Rep *27*, 1446-1460 e1444.
10.1016/j.celrep.2019.04.019.
59.   Takatsuka, S., Yamada, H., Haniuda, K., Saruwatari, H., Ichihashi, M., Renauld, J.C.,
and Kitamura, D. (2018). IL-9 receptor signaling in memory B cells regulates humoral recall
responses. Nat Immunol *19*, 1025-1034. 10.1038/s41590-018-0177-0.
60.   Rumfelt, L.L., Zhou, Y., Rowley, B.M., Shinton, S.A., and Hardy, R.R. (2006). Lineage
specification and plasticity in CD19- early B cell precursors. J Exp Med *203*, 675-687.
10.1084/jem.20052444.
61.   Mallampati, S., Sun, B., Lu, Y., Ma, H., Gong, Y., Wang, D., Lee, J.S., Lin, K., and Sun,
X. (2014). Integrated genetic approaches identify the molecular mechanisms of Sox4 in early B-
cell development: intricate roles for RAG1/2 and CK1epsilon. Blood *123*, 4064-4076.
10.1182/blood-2013-12-543801.
62.   Greig, K.T., de Graaf, C.A., Murphy, J.M., Carpinelli, M.R., Pang, S.H., Frampton, J.,
Kile, B.T., Hilton, D.J., and Nutt, S.L. (2010). Critical roles for c-Myb in lymphoid priming and
early B-cell development. Blood *115*, 2796-2805. 10.1182/blood-2009-08-239210.
63.   Clark, M.R., Mandal, M., Ochiai, K., and Singh, H. (2014). Orchestrating B cell
lymphopoiesis through interplay of IL-7 receptor and pre-B cell receptor signalling. Nat Rev
Immunol *14*, 69-80. 10.1038/nri3570.
64.   Callahan, D., Smita, S., Joachim, S., Hoehn, K., Kleinstein, S., Weisel, F., Chikina, M.,
and Shlomchik, M. (2024). Memory B cell subsets have divergent developmental origins that are

coupled to distinct imprinted epigenetic states. Nat Immunol *25*, 562-575. 10.1038/s41590-023-
01721-9.

65.   Krishnamurty, A.T., Thouvenel, C.D., Portugal, S., Keitany, G.J., Kim, K.S., Holder, A.,
Crompton, P.D., Rawlings, D.J., and Pepper, M. (2016). Somatically Hypermutated
Plasmodium-Specific IgM(+) Memory B Cells Are Rapid, Plastic, Early Responders upon
Malaria Rechallenge. Immunity *45*, 402-414. 10.1016/j.immuni.2016.06.014.

66.   Sutton, H.J., Aye, R., Idris, A.H., Vistein, R., Nduati, E., Kai, O., Mwacharo, J., Li, X.,
Gao, X., Andrews, T.D., et al. (2021). Atypical B cells are part of an alternative lineage of B cells
that participates in responses to vaccination and infection in humans. Cell Rep *34*, 108684.
10.1016/j.celrep.2020.108684.

67.   Perez-Mazliah, D., Gardner, P.J., Schweighoffer, E., McLaughlin, S., Hosking, C.,
Tumwine, I., Davis, R.S., Potocnik, A.J., Tybulewicz, V.L., and Langhorne, J. (2018).
Plasmodium-specific atypical memory B cells are short-lived activated B cells. Elife *7*.
10.7554/eLife.39800.

68.   Kikuchi, K., Kasai, H., Watanabe, A., Lai, A.Y., and Kondo, M. (2008). IL-7 specifies B
cell fate at the common lymphoid progenitor to pre-proB transition stage by maintaining early B
cell factor expression. Journal of immunology *181*, 383-392. 10.4049/jimmunol.181.1.383.

69.   Corfe, S.A., and Paige, C.J. (2012). The many roles of IL-7 in B cell development;
mediator of survival, proliferation and differentiation. Semin Immunol *24*, 198-208.
10.1016/j.smim.2012.02.001.

70.   Reya, T., O'Riordan, M., Okamura, R., Devaney, E., Willert, K., Nusse, R., and
Grosschedl, R. (2000). Wnt signaling regulates B lymphocyte proliferation through a LEF-1
dependent mechanism. Immunity *13*, 15-24. 10.1016/s1074-7613(00)00004-2.

71.   Shehata, L., Thouvenel, C.D., Hondowicz, B.D., Pew, L.A., Pritchard, G.H., Rawlings,
D.J., Choi, J., and Pepper, M. (2024). Interleukin-4 downregulates transcription factor BCL6 to
promote memory B cell selection in germinal centers. Immunity *57*, 843-858 e845.
10.1016/j.immuni.2024.02.018.

72.   Duan, L., Liu, D., Chen, H., Mintz, M.A., Chou, M.Y., Kotov, D.I., Xu, Y., An, J., Laidlaw,
B.J., and Cyster, J.G. (2021). Follicular dendritic cells restrict interleukin-4 availability in
germinal centers and foster memory B cell generation. Immunity *54*, 2256-2272 e2256.
10.1016/j.immuni.2021.08.028.

73.   Ly, A., Liao, Y., Pietrzak, H., Ioannidis, L.J., Sidwell, T., Gloury, R., Doerflinger, M.,
Triglia, T., Qin, R.Z., Groom, J.R., et al. (2019). Transcription Factor T-bet in B Cells Modulates
Germinal Center Polarization and Antibody Affinity Maturation in Response to Malaria. Cell Rep
29, 2257-2269 e2256. 10.1016/j.celrep.2019.10.087.

74.   Stone, S.L., Peel, J.N., Scharer, C.D., Risley, C.A., Chisolm, D.A., Schultz, M.D., Yu, B.,
Ballesteros-Tato, A., Wojciechowski, W., Mousseau, B., et al. (2019). T-bet Transcription Factor
Promotes Antibody-Secreting Cell Differentiation by Limiting the Inflammatory Effects of IFN-
gamma on B Cells. Immunity *50*, 1172-1187 e1177. 10.1016/j.immuni.2019.04.004.

75.   Nellore, A., Zumaquero, E., Scharer, C.D., Fucile, C.F., Tipton, C.M., King, R.G., Mi, T.,
Mousseau, B., Bradley, J.E., Zhou, F., et al. (2023). A transcriptionally distinct subset of
influenza-specific effector memory B cells predicts long-lived antibody responses to vaccination
in humans. Immunity *56*, 847-863 e848. 10.1016/j.immuni.2023.03.001.

76.   Curran, M.A., Geiger, T.L., Montalvo, W., Kim, M., Reiner, S.L., Al-Shamkhani, A., Sun,
J.C., and Allison, J.P. (2013). Systemic 4-1BB activation induces a novel T cell phenotype
driven by high expression of Eomesodermin. The Journal of experimental medicine *210*, 743-
755. 10.1084/jem.20121190.

77.   Hong, J.P., Reynoso, G.V., Andhey, P.S., Swain, A., Turner, J.S., Boon, A.C.M.,
Krammer, F., Ellebedy, A.H., Zanini, F., Artyomov, M., et al. (2020). An Agonistic Anti-CD137

Antibody Disrupts Lymphoid Follicle Structure and T-Cell-Dependent Antibody Responses. Cell
Rep Med *1*. 10.1016/j.xcrm.2020.100035.
78.   Racine, R., Jones, D.D., Chatterjee, M., McLaughlin, M., Macnamara, K.C., and
Winslow, G.M. (2010). Impaired germinal center responses and suppression of local IgG
production during intracellular bacterial infection. Journal of immunology *184*, 5085-5093.
10.4049/jimmunol.0902710.
79.   Yates, J.L., Racine, R., McBride, K.M., and Winslow, G.M. (2013). T cell-dependent IgM
memory B cells generated during bacterial infection are required for IgG responses to antigen
challenge. Journal of immunology *191*, 1240-1249. 10.4049/jimmunol.1300062.
80.   Moir, S., Ho, J., Malaspina, A., Wang, W., DiPoto, A.C., O'Shea, M.A., Roby, G., Kottilil,
S., Arthos, J., Proschan, M.A., et al. (2008). Evidence for HIV-associated B cell exhaustion in a
dysfunctional memory B cell compartment in HIV-infected viremic individuals. The Journal of
experimental medicine *205*, 1797-1805. 10.1084/jem.20072683.
81.   Portugal, S., Tipton, C.M., Sohn, H., Kone, Y., Wang, J., Li, S., Skinner, J., Virtaneva, K.,
Sturdevant, D.E., Porcella, S.F., et al. (2015). Malaria-associated atypical memory B cells
exhibit markedly reduced B cell receptor signaling and effector function. Elife *4*.
10.7554/eLife.07218.
82.   Sullivan, R.T., Kim, C.C., Fontana, M.F., Feeney, M.E., Jagannathan, P., Boyle, M.J.,
Drakeley, C.J., Ssewanyana, I., Nankya, F., Mayanja-Kizza, H., et al. (2015). FCRL5 Delineates
Functionally Impaired Memory B Cells Associated with Plasmodium falciparum Exposure. PLoS
Pathog *11*, e1004894. 10.1371/journal.ppat.1004894.
83.   Elsner, R.A., Smita, S., and Shlomchik, M.J. (2024). IL-12 induces a B cell-intrinsic IL-
12/IFNgamma feed-forward loop promoting extrafollicular B cell responses. Nat Immunol *25*,
1283-1295. 10.1038/s41590-024-01858-1.
84.   Kimura, T., Nakayama, K., Penninger, J., Kitagawa, M., Harada, H., Matsuyama, T.,
Tanaka, N., Kamijo, R., Vilcek, J., Mak, T.W., and et al. (1994). Involvement of the IRF-1
transcription factor in antiviral responses to interferons. Science *264*, 1921-1924.
10.1126/science.8009222.
85.   Kim, I.K., Kim, B.S., Koh, C.H., Seok, J.W., Park, J.S., Shin, K.S., Bae, E.A., Lee, G.E.,
Jeon, H., Cho, J., et al. (2015). Glucocorticoid-induced tumor necrosis factor receptor-related
protein co-stimulation facilitates tumor regression by inducing IL-9-producing helper T cells. Nat
Med *21*, 1010-1017. 10.1038/nm.3922.
86.   Niese, M.L., Pajulas, A.L., Rostron, C.R., Cheung, C.C.L., Krishnan, M.S., Zhang, J.,
Cannon, A.M., and Kaplan, M.H. (2024). TL1A priming induces a multi-cytokine Th9 cell
phenotype that promotes robust allergic inflammation in murine models of asthma. Mucosal
Immunol *17*, 537-553. 10.1016/j.mucimm.2024.03.006.
87.   Ming, S., Zhang, M., Liang, Z., Li, C., He, J., Chen, P., Zhang, S., Niu, X., Deng, S.,
Geng, L., et al. (2021). OX40L/OX40 Signal Promotes IL-9 Production by Mucosal MAIT Cells
During Helicobacter pylori Infection. Front Immunol *12*, 626017. 10.3389/fimmu.2021.626017.
88.   Xiao, X., Balasubramanian, S., Liu, W., Chu, X., Wang, H., Taparowsky, E.J., Fu, Y.X.,
Choi, Y., Walsh, M.C., and Li, X.C. (2012). OX40 signaling favors the induction of T(H)9 cells
and airway inflammation. Nat Immunol *13*, 981-990. 10.1038/ni.2390.
89.   Shuford, W.W., Klussman, K., Tritchler, D.D., Loo, D.T., Chalupny, J., Siadak, A.W.,
Brown, T.J., Emswiler, J., Raecho, H., Larsen, C.P., et al. (1997). 4-1BB costimulatory signals
preferentially induce CD8+ T cell proliferation and lead to the amplification in vivo of cytotoxic T
cell responses. The Journal of experimental medicine *186*, 47-55. 10.1084/jem.186.1.47.
90.   Tan, J.T., Whitmire, J.K., Murali-Krishna, K., Ahmed, R., Altman, J.D., Mittler, R.S.,
Sette, A., Pearson, T.C., and Larsen, C.P. (2000). 4-1BB costimulation is required for protective

anti-viral immunity after peptide vaccination. Journal of immunology *164*, 2320-2325.
10.4049/jimmunol.164.5.2320.
91.   Wong, H.Y., Prasad, A., Gan, S.U., Chua, J.J.E., and Schwarz, H. (2020). Identification
of CD137-Expressing B Cells in Multiple Sclerosis Which Secrete IL-6 Upon Engagement by
CD137 Ligand. Front Immunol *11*, 571964. 10.3389/fimmu.2020.571964.
1000
1001

 FIGURE LEGENDS

1003

Fig. 1: Exogenous 4-1BB stimulation leads to loss of parasite control due to derailed GC
response during *Plasmodium* infection. a, Experimental design. C57BL/6 mice were infected
with *Plasmodium yoelii*-parasitized red blood cells (*Py* pRBCs) and treated with 50μg of 4-1BB
agonist antibody (3H3) or isotype control (rIgG) on days 5 and 7 post infection. **b,** Kinetics of
parasite burden (% infected RBCs). Data are mean $\pm$ s.e.m and are pooled from *n=2* independent
experiments using *n=7* mice for either group. **c,d,** Representative flow plots (left) and kinetics
(right) of germinal center B cells (**c**) and germinal center Tfh cells (**d**) in the spleens of IgG- and
3H3-treated mice. **e,** Summary graph of anti-MSP-1$_{-19}$ IgG2b serum antibody end-point titers
(EPT) on 21 and 39 dpi. For **c-e,** data are mean $\pm$ s.d and are representative of *n=2* biologically
independent with similar results using *n=3* mice for either group. **f,** Confocal micrographs of rIgG-
and 3H3-treated spleens on days 21 and 39 post infection showing total B cells (green), CD4 T
cells (purple) and GC B cells (red). Data are representative of at least 2 biologically independent
experiments. For **f**, scale bar: 200 μm. For **b, c** and **d,** two-way ANOVA and for **e,** two-tailed
Students' *t* test was used for statistical analysis.

1018

Fig. 2: 4-1BB stimulation enhances MBC recall and protection against rechallenge
independently of antigen load. a, Experimental design. C57BL/6 mice were infected with *Py*
and treated with 3H3 or isotype control as previously described. During the convalescent phase,
mice were challenged with *Plasmodium berghei* ANKA-parasitized (*PbA*) RBCs. **b,** Survival graph
of mice following *PbA* challenge. Data are pooled from two independent experiments. **c,** Fold
change in anti-MSP-1$_{-19}$ IgM serum antibody EPT between 0 and 5 dpi. Data are pairwise
comparison of the two time points in each mouse and are pooled from *n=2* biologically
independent experiments using *n=6* (rIgG) and *n=5* (3H3). **d,** Experimental design. C57BL/6 mice
were infected with *Py* and treated with 3H3 or isotype control as previously described. Mice were
treated with the Artemether (Artem) to control the parasite load and challenged with *PbA*. **e,**
Survival of mice following *PbA* challenge. Data are pooled from two independent experiments. **f,**
Fold change in anti-MSP-1$_{-19}$ IgM serum antibody EPT between 0 and 5 dpi. Data represents
pairwise comparison of the two time points in each mouse and are pooled from *n=2* biologically
independent experiments with similar data using *n=7* (rIgG) and *n=7* (3H3). **g,** Experimental
design. C57BL/6 mice were infected with irradiated parasites and treated with 3H3 or isotype
control as previously described. During convalescence mice were rechallenged with *PbA*. **h,**
Survival of mice following *PbA* infection. Data are pooled from two independent experiments. **i,**
Fold change in anti-MSP-1$_{-19}$ IgM serum antibody EPT between 0 and 5 dpi. Data are pairwise
comparison of the two time points in each mouse and are pooled from *n=2* biologically
independent experiments using *n=5* (rIgG) and *n=5* (3H3). For **b**, **e** and **h**, data were analyzed
by Mantel-Cox test. For **c, f** and **i**, two-tailed Mann-Whitney U tests were used for statistical
analysis.

1041

Fig. 3: Enhanced recall potential and not abundance of MBCs drives protection following
4-1BB stimulation. a, Anti-MSP-1$_{-19}$ IgM ELISPOT of bone marrow plasma cells (BMPCs) on 5
dpi post *PbA* challenge. Data are representative of *n=2* biologically independent experiments
using *n=4* mice/group. **b-e**, *Py*-infected C57BL/6 mice were treated with 3H3 or isotype control as
previously described. **d-f,** Mice were additionally treated with the antimalarial drug artemether at
regular intervals as detailed elsewhere. Representative flow plots (left) and total numbers (right)
of polyclonal (**b,d**) and MSP-1 specific (**c,e**) MBCs in the spleen. Data are either representative
(**b,c**) or are pooled from *n=2* biologically independent experiments using *n=3-4* mice/group. **f,**

Representative flow plots (left) and frequency (right) of proliferating BMPCs as detected by Ki67
and TACI staining. **g,** Geometric mean fluorescence intensity (gMFI) of MSP-1 staining in MSP-
1 specific MBCs. Data are representative of *n=2* biologically independent experiments using *n=3*
mice/group. **h,** Total MBCs from rIgG- and 3H3-treated mice were flow-sorted and transferred to
μMT recipients that were infected with *Py* the following day. **h**, Summary graphs showing parasite
burden on 12 dpi and **i,** anti-MSP-1$_{-19}$ IgM serum antibody titers on 4 dpi. Data are representative
of *n=2* biologically independent experiments using *n=4* mice/group. Data were analyzed by two-
tailed Mann-Whitney U tests (**a-e** and **g-l**), and Student's *t* test (**f**).

Fig. 4: Exogenous 4-1BB ligation induces MBCs with an extrafollicular gene signature. a,
Experimental design. C57BL/6 mice were infected and treated as indicated. Polyclonal and MSP-
1 specific MBCs were flow-sorted and scRNA-seq, V(D)J sequencing and CITE-seq were
performed. **b,** UMAP showing the identified clusters of polyclonal MBCs. **c,** Pie diagram depicting
the frequency of respective clusters in the polyclonal MBC pool derived from rIgG and 3H3-treated
mice. **d,** Heatmap depicting markers from each cluster with selected genes highlighted on the left.
e,f, Cells were scored based on gene signatures of MBCs derived from a GC response, EF
response or from atypical B cells and were visualized per cluster (**e**) and the prevalence of each
of these signatures are compared between treatments as shown (**f**). **g,** UMAP showing clusters
in MSP-1 specific MBCs. **h,** Frequency of clusters in MSP-1 specific MBCs from rIgG or 3H3-
treated mice cells are shown in the pie chart. **i,** Heatmap of markers from each cluster with
selected genes highlighted on the left. **j,k,** Cells were scored based on gene signatures of MBCs
derived from a GC response, EF response or atypical B cells and visualized per cluster (**j**) and
between treatments (**k**). Data from **f** and **k** were analyzed by Wilcoxon test. Cells were pooled
from 3 mice per group prior to sequencing.

Fig. 5: 4-1BB stimulation skews MBC differentiation towards a highly functional
population. C57BL/6 mice were infected and treated as indicated in **Fig. 4a**. Distribution of IgM
(**a**), CD73 and CD80 (**b**) expression among the polyclonal MBCs clusters. **c,** Pseudo time
trajectories (left) and their representative contributions towards each cluster in polyclonal MBCs
in rIgG- and 3H3-treated mice (right) based on their frequencies in **Fig. 4c**. Distribution of IgM (**d**),
CD73 and CD80 (**e**) expression among the MSP-1 specific MBCs clusters. **f,** Pseudo time
trajectories (left) and their representative contributions in MSP-1 specific MBCs in rIgG- and 3H3-
treated mice (right) based on their frequencies in **Fig. 4h**. Gene ontology (GO) analysis of
upregulated genes in MBCs from 3H3-treated mice compared to MBCs from rIgG-treated mice in
clusters 0 and 2 of polyclonal MBCs combined (**g**) and cluster 1 of MSP-1 specific MBCs (**h**).

Fig. 6: 3H3-driven protection is dependent on IL-9:IL-9R signaling. C57BL/6 mice were
infected and treated as indicated in **Fig. 4a**. UMAP clustering of scRNA-seq data showing *Il9r*
expression among polyclonal (**a**) and MSP-1 specific MBC clusters (**b**). **c,** UMAP clustering of
polyclonal MBCs on 38 dpi generated by FlowSOM. **d,** Expression levels of various proteins using
FlowSOM clustering of polyclonal MBCs. **e,** Geometric mean fluorescence intensity (gMFI) of IL-
9R in polyclonal MBCs on 38 dpi. Data is pooled from *n=2* biologically independent experiments
using n=4 mice/group/experiment. **f,** UMAP clustering of MSP-1 specific MBCs on 38 dpi
generated by FlowSOM. **g,** Expression levels of various proteins using FlowSOM clustering of
MSP-1 specific MBCs. **h,** gMFI of IL-9R in MSP-1 specific MBCs on 38 dpi. Data is pooled from
n=2 biologically independent experiments using n=4 mice/group/experiment. **i,j,** WT mice were
infected with *Py* and treated with rIgG or 3H3 and then with the antimalarial drug artemether at
regular intervals. One day prior and after *PbA* rechallenge, mice were treated with αIL-9
neutralizing antibody or isotype control (mIgG) (as detailed in **Supp Fig. 7d**). Survival was

monitored (**i**) and fold change in anti-MSP-1$_{-19}$ IgM serum antibody EPT between 0 and 5 dpi was
assessed (**j**). Data is pooled from *n=2* biologically independent experiments using *n=6*
mice/group. **k**,**l**, B cell-specific WT (B^{WT}) and IL-9R KO (B$^{IL-9R\ KO}$) chimeras generated and were
infected with *Py* and treated with 3H3 on days 5 and 7 post infection and then with the antimalarial
drug artemether at regular intervals (**Supp Fig.7e**). On 38 dpi, during the convalescent phase,
mice were rechallenged with *PbA*. Survival was monitored (**k**) and fold change in anti-MSP-1$_{-19}$
IgM serum antibody EPT between 0 and 5 dpi was assessed (**l**). Data is pooled from *n=2*
biologically independent experiments using *n=11* mice/group (**k**) and are representative of *n=2*
biologically independent experiments using *n=5-6* mice/group (**l**). For **e**, **h**, **j** and **l** two-tailed Mann-
Whitney U tests were used for statistical analysis. For **i** and **k**, data were analyzed by Mantel-
Cox.
1110
Fig. 7: Altered chromatin landscape in MBCs following 4-1BB stimulation poises them to
readily become PCs upon recall. a, Experimental design. C57BL/6 mice were infected and
treated as indicated. Polyclonal and MSP-1 specific MBCs were flow-sorted and scATAC-seq was
performed. **b,** UMAP clustering of MSP-1 specific MBCs using scATAC-seq data with transferred
labels from scRNA-seq post data integration. Comparison of *Il9r* (**c**) and *Prdm1* (**d**) accessibility
between clusters 0 and 1 of MSP-1 specific MBCs. **e,** Motif analysis of differentially accessible
regions (DARs) in cluster 1 of MSP-1 specific MBCs compared to the other clusters. **f,**
Comparison of *Il9r* accessibility in cluster 1 between MSP-1 specific MBCs obtained from rIgG
and 3H3-treated mice. **g,** Motif analysis of differentially accessible regions (DARs) in 3H3
compared to rIgG in cluster 1 of MSP-1 specific MBCs. Sequenced nuclei were pooled from 4
mice per group.
1122

a

i.v
10^6 *Py* pRBCs

i.p
rIgG/3H3

Kinetics of:
-Parasite load
-GC responses
-Serum antibody
-Confocal imaging

b

c Germinal Center B cells

d Germinal Center Tfh cells

e

f

B220 CD4 GL7

a Day 5 post *PbA* challenge, ELISPOT

b Polyclonal MBCs

c MSP-1 specific

d Polyclonal MBCs

e MSP-1 specific

f

g

h

i

a

Py rIgG/3H3 Artemether Sort MBCs

Polyclonal MBCs
(B220⁺ IgD⁻ GL7⁻ CD38⁺)

MSP-1 specific MBCs
(B220⁺ IgD⁻ GL7⁻ CD38⁺ Decoy⁻ MSP1⁺)

scATAC-seq

b MSP-1 specific MBCs
Post scRNA-seq integration

c MSP-1 cluster 1 coverage

d

e MSP-1 cluster 1 motifs

f 3H3 vs. rIgG (cluster 1) coverage

g 3H3 vs. rIgG (cluster 1) motifs