Exogenous 4-1BB co-stimulation enhances memory B cell response during malaria.

Carolina Caloba[1,2], Alexandria J. Sturtz[3,6], Akshaya Ramachandran[1,7], Lijo John[1,2], Kateryna Malkina[1,4], Allen M. Minns[5], Scott E. Lindner[5], Rahul Vijay[1,2]

[1]Center for Cancer Cell Biology, Immunology and Infection, [2]Department of Microbiology and Immunology, Rosalind Franklin University of Medicine and Science, North Chicago, IL, USA. [3]Department of Microbiology and Immunology, University of Iowa, Iowa City, Iowa, USA, [4]Lake Forest College, Lake Forest, Illinois, USA.[5]Department of Biochemistry and Molecular Biology, Hucks Center for Malaria Research, Pennsylvania State University, University Park, Pennsylvania, USA. [6]Current Address: Division of Biology and Biomedical Sciences, Washington University, St.Louis, Missouri, USA. [7]Current Address: Harvard Medical School, Boston, Massachusetts, USA

Running Title: 4-1BB stimulation and memory B cells

Abstract:
Clearance of the malarial parasite and protection from reinfection is dependent on T helper cell-dependent germinal center (GC) -derived plasma cells (PC) and memory B cells (MBC). Enhancing or blocking T cell help during experimental malaria has been shown to either improve or impede GC output, respectively. Using a mouse model of malaria, we show that an agonistic antibody against the co-stimulatory molecule, 4-1BB (CD137), hinders an optimal GC response, but enhances humoral immune memory. We demonstrate that this apparent paradox is at least partially driven by CD4 T cell-intrinsic *Eomesodermin* expression. By using chemoprophylaxis or irradiated parasites we show that the beneficial effects of 4-1BB ligation on humoral immune memory are neither influenced by the inflammatory milieu, nor due to an increased number of MBCs, but rather due to their heightened recall capacity and their ability to readily differentiate into antibody secreting PCs.

Introduction

Despite decades of research, malaria still remains a major public health threat with an estimated 247 million cases and 619000 related deaths in 2021(1). The burden of the disease is further exacerbated by *Plasmodium*-immune evasion strategies as well as the continuing emergence of drug-resistant strains. Humoral immune memory following *Plasmodium* exposure is suboptimal, resulting in repeat infections. While the approval of the anti-malarial vaccine RTS,S/AS01, to be used in children is a major breakthrough, it provides limited efficacy (~30%) even after the regimented 4 doses (2).

Clearance of the malarial parasite relies on spatially and temporally layered CD4 T cell-dependent B cell activation events culminating in the production of parasite-specific antibodies. Concomitant with the clearance of infection, a fraction of activated B cells differentiates into MBCs, that quickly become antibody-secreting PC upon reinfection. However, increasing evidence suggest that, while humoral immune memory against *Plasmodium* provides clinical immunity, sterilizing immunity is rarely achieved (3). Expression and engagement of coinhibitory receptors such as PD1, LAG3, and CTLA-4 on CD4 T cells have been attributed to this suboptimal anti-*Plasmodium* immune responses *(4-8)*. These studies used combinatorial approaches to simultaneously block the inhibitory receptors while also activating costimulatory molecule (such as OX-40) to improve both effector and humoral immune memory (4, 6, 7). Previous studies have reported that agonizing OX40, a member of the tumor necrosis factor receptor superfamily (TNFRSF) expressed on T cells would improve both effector and humoral immune memory following *Plasmodium* infection (6, 7). 4-1BB or CD137 is another TNFRSF member expressed on T cells whose precise role in modulating anti-*Plasmodium* immunity have not been explored.

In this study, we investigated the role of exogenous 4-1BB stimulation using a monoclonal antibody, 3H3, in a rodent malaria model and show an enhancement in memory B cell response (MBC). We show that while 3H3 treatment derails effector humoral immune response and early parasite control, it

51 significantly improves the humoral immune memory-mediated protection from an otherwise acutely lethal
52 *Plasmodium* challenge.
53
54 **Materials and Methods**
55
56 *Animals, infection, treatments and parasitemia*
57
C57BL/6 mice wild-type were purchased from The Jackson Laboratory. $CD4^{Ert2\text{-}Cre}$ and $Eomes^{fl/fl}$ mice were
obtained from Dr. Noah Butler (University of Iowa), and 4-1BB KO mice from Dr.Tania Watts (University
of Toronto) and are bred in-house. *Plasmodium yoelli* (*Py*) clone 17XNL and *P. berghei* clone ANKA
(*Pb.ANKA*) were obtained from the Malaria Research and Reference Reagent Resource Center (MR4;
American Type Culture Collection), stocks generated and mice infected as previously described (9). *Py* -
infected mice were treated intraperitoneally with either 50 μg of anti-4-1BB antibody (3H3; BioXCell) or
Rat IgG (rIgG) on days 5 and 7 post infection. For $CD4^{Ert2\text{-}Cre}Eomes^{fl/fl}$ experiments, 4 mg of tamoxifen
(TAM) (Sigma, MO) were administered intraperitoneally on days 2, 3 and 4 post infection. In some
experiments, mice were treated with either 100 μL of Artemether (10mg/mL) or Mineral Oil. Parasitemia
was measured by flow cytometry as previously described.(10, 11). All experiments and procedures were
approved by the Rosalind Franklin University of Medicine and Science Institutional Animal Care and Use
Committee.
70
71 *Confocal imaging*
72
73 Spleens were processed as previously describedn(9). Antibodies used for staining are B220-AF488 (clone
74 RA3-6B2; eBioscience), CD4-AF594 (clone GK1.5; BioLegend) and GL7-AF647 (BD Pharmingen).
75 Imaging was done using a Zeiss LSM710 confocal microscope and images were processed using IMARIS
76 x64 software (version 9.2.1).
77
78 *ELISpot and ELISA*
79
Nunc MaxiSorp white polystyrene plates were coated with 0.5 μg ml^{-1} of recombinant MSP1$_{19}$ and blocked
for at least 2 h with PBS + 2.5% w/v BSA + 5% v/v goat serum. Bone marrow cells from *P. berghei* ANKA-
infected mice were serially diluted in supplemented Iscoves modified Dulbecco's media and plated for 20
h at 37 °C with 5% CO_2. Plates were washed with PBS + 0.05% Tween 20 and incubated overnight with
horseradish peroxidase (HRP)-conjugated IgM at 4 °C. Spots were developed with 3-amino-9-
ethylcarbazole.
⁠   Nunc MaxiSorp plates were coated with 0.5 μg ml^{-1} MSP1$_{19}$ and blocked with 2.5% w/v BSA +
5% v/v normal goat serum. Serum samples from naive and *Py*-infected mice were serially diluted and
incubated for 18 h at 4 °C. MSP1$_{19}$-specific antibodies were detected with HRP-conjugated IgG2b or IgG2c.
Plates were developed using a SureBlue reserve TMB Kit (KPL), according to the manufacturer's protocol,
and absorbance was measured at an optical density of 450 nm using a SpectraMax 340 (Molecular Devices).
End-point titers were extrapolated from a sigmoidal 4PL (where x is the log concentration) standard curve
for each sample. The threshold for end-point titers was the mean plus 0.5–8 multiplied by the standard
deviation recorded for naive mouse sera.
94
95 *Flow cytometry*
96
97 After generation of single-cell suspension of splenocytes, RBCs were lysed and the cells were counted.
98 Samples were blocked using Fc block (clone 2.4G2) in FACS buffer (PBS + 0.09% w/v sodium azide + 2%
99 v/v fetal calf serum) and cells were stained with fluorescently labeled antibodies. Transcription factors were
100 stained using Foxp3 Transcription factor staining buffer (Tonbo), according to manufacturer's protocol.
101 Samples were acquired on BD-LSR-II or Cytek Aurora and analyzed using FlowJo software (TreeStar Inc).

SpyCage Reagent for detecting MSP1-19 specific B cells

SpyCage reagents bearing 60 copies of either mScarlet ("RedCage") or mNeonGreen ("GreenCage") were covalently loaded with 60 copies of the immunodominant *P. yoelii* antigen MSP1 or the liver stage antigen PyUIS4 as a decoy control, respectively. Antibodies specific to SpyCage were biotinylated using amine-reactive biotinylated crosslinkers, and were used to enrich SpyCage-bound cells.

MBC transfer

Memory B cells (B220$^+$IgD$^-$CD38$^+$GL7$^-$) were sort-purified from *Py*-infected mice treated with rIgG or 3H3. 2x10^5 cells were transferred intravenously to naive µMT mice, which were infected with 1×10^6 *Py* - parasitized RBCs.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 10 software (GraphPad). Specific tests are detailed in the figure legends.

Results and Discussion

Exogenous 4-1BB co-stimulation delays GC response

Our initial studies revealed ~2-3-fold higher numbers of 4-1BB$^+$ CD4 T cells compared to their CD8 T cell counterparts both in naïve as well as *Plasmodium yoelii* (*Py*) infected mice (up to 7 dpi) (**Supplemental Fig. 1***A*) and that 4-1BB KO mice infected with *Py* showed heightened parasite load compared to WT mice (**Fig. 1***A*). Enhanced susceptibility of KO mice was also associated with a significantly reduced GC response (data not shown). Notably, and contrary to the role of OX40 ligation(6), ligation of 4-1BB using an agonistic monoclonal antibody (Ab) (clone 3H3) showed a prolonged infection course with an 8-fold higher total parasite load, which eventually resolved by day 39-40 p.i. (**Fig. 1***B*). We next investigated whether delayed parasite control in 3H3-treated mice was associated with perturbations of GC-associated effector populations (**Supplemental Fig 1***B,C*). We found reduced frequencies and numbers of GC B cells (**Fig. 1***C*) and GC Tfh cells (**Fig. 1***D*) and serum-anti-*Plasmodium* antibody titers (**Fig.1** *E*) until at least day 39 post-infection (p.i.). Concomitant with the delay in GC response, we also observed a pronounced disruption of lymphoid architecture following 3H3 treatment until 21 d.p.i (**Fig. 1***F*). However, by 39 d.p.i the follicles regrouped into GCs coinciding with the resolution of infection in 3H3-treated, *Plasmodium*-infected mice (**Fig. 1***F)*. These data show that 4-1BB is necessary for optimal *Plasmodium* parasite control and that agonistic ligation of 4-1BB distorts lymphoid architecture and GC orchestration during experimental malaria.

Deletion of CD4-intrinsic Eomes rescues GC response following exogenous 4-1BB co-stimulation

3H3-mediated engagement of 4-1BB upregulates eomesodermin (EOMES) in CD4 T cells(12) which we also observed in *Plasmodium* infection-induced effector CD4 T cells by day 9 p.i (**Supplemental Fig. 2***A*). However, the number of EOMES$^+$ cells waned by day 21 p.i, which coincided with the appearance of BCL6$^+$ CD4 T cells in 3H3 treated mice (**Supplemental Fig. 2***B,C*). This observation led us to speculate that EOMES and BCL6 in CD4 T cells may counter-repress one another during *Py* infection. To test whether the lack of CD4 T cell- intrinsic *BCL6* facilitates EOMES upregulation, we infected *CD4.Cre.Bcl6$^{fl/fl}$* mice with *Py* and treated them with 3H3 or rIgG. 3H3 treatment resulted in EOMES upregulation independent of BCL6 expression, showing that BCL6 does not regulate EOMES expression (**Supplemental Fig. 2***D)*. To test the converse notion of whether EOMES regulates BCL6 expression, we infected

CD4.Ert2.Cre.Eomes$^{fl/fl}$ mice with *Py*, followed by Tamoxifen (TAM) treatment to trigger deletion of
EOMES in CD4 T cells (**Supplemental Fig. 2E, left panel**). Deletion of EOMES in CD4 T cells neither
affected the expression of BCL6 (**Supplemental Fig. 2F**) nor altered parasite control in *Py-*
CD4.Ert2.Cre.Eomes$^{fl/fl}$ infected mice (**Supplemental Fig. 2G**). However, when EOMES in CD4 T cells
was deleted in *Py*-infected mice (Cre$^+$) treated with 3H3, we observed a marked enhancement of parasite
control compared to Cre$^-$ mice (**Fig. 2A**). This enhanced parasite control was associated with elevated
numbers of GC B cells (**Fig. 2B**) and GC Tfh cells (**Fig. 2C**). Thus, no apparent counterregulatory loop
exists between EOMES and BCL6, yet the detrimental effects of 3H3-mediated ligation of 4-1BB on GC
responses requires CD4 T cell-intrinsic expression of EOMES.

Exogenous 4-1BB costimulation enhances survival following lethal Plasmodium challenge

A delayed effector GC response following 3H3 treatment (**Fig. 1**) is predicted to also impair protective
humoral immune memory responses against a subsequent parasite challenge. To test this, we monitored
survival and humoral immune memory responses in convalescent *Py*-infected-3H3/rIgG treated WT mice
that were challenged (day 93 p.i) with a virulent and acutely lethal *Plasmodium* strain, *P. berghei ANKA*
(Pb.ANKA). We observed that 3H3-treated, convalescent mice showed significantly higher survival rates
compared to rIgG-treated mice following *Pb.ANKA* challenge (**Fig. 3A**). Enhanced survival of 3H3-treated
mice correlated with greater fold increases in anti-merozoite surface protein (MSP1$_{-19}$)-specific antibody
titers between day 0 and day 5 post-*Pb.ANKA* challenge (**Fig. 3B**), a measure of enhanced memory B cell
(MBC) recall response. Because derailed effector GC responses in 3H3-treated mice requires CD4 T cell-
intrinsic *EOMES* expression (**Fig. 2**), we investigated whether EOMES expression also governs immune
memory-mediated protection against heterologous *Pb.ANKA* challenge. *Py*-infected
CD4.Ert2.Cre.Eomes$^{fl/fl}$ (Cre$^+$) mice and their genetic littermate control cohorts (Cre-) were treated with
TAM on days 2-4 p.i followed by 3H3 on 5 and 7 dpi (as in **Fig. 2**) and subsequently challenged with
Pb.ANKA on day 93 post-*Py* infection. We found that CD4 T cell-intrinsic expression of EOMES is also
necessary for immune memory-mediated resistance to *Pb.ANKA* challenge (**Fig. 3C**). The loss of protection
in *Cre$^+$* mice strongly correlated with a significantly subdued MBC recall as evidenced by a lower fold
increase in anti-MSP1$_{-19}$-specific antibody levels, compared to their *Cre$^-$* littermates (**Fig. 3D**). These data
show that CD4 T cell-intrinsic EOMES expression is essential for enhanced MBC recall following 3H3
treatment.
To investigate whether the enhanced MBC recall and subsequent improvement in survival
following 3H3 treatment (**Fig. 1A**) is independent of the heightened inflammation that may ensue higher
and prolonged parasite load (**Fig. 1A**), we truncated infection in *Py*-infected, 3H3- or rIgG-treated WT mice
by administering the antimalarial drug artemether on 12-14 dpi (**Supplemental Fig. 3A**). Notably,
compared to vehicle (mineral oil)-treated mice, artemether treatment neither compromised protection (**Fig.**
3E) nor impaired the enhanced MBC recall response (**Fig. 3F**) mediated by 3H3 treatment following
Pb.ANKA challenge. Notably, artemether treatment did not alter EOMES expression in CD4 T cells
(**Supplemental Fig. 3B**), and did not rescue the effector (day 21 p.i.) GC responses in *Py* infected 3H3-
treated mice (**Supplemental Fig. 3B,C**), which matched infection outcomes in mice treated with 3H3 alone
*(***Supplemental Fig. 2A**). To test whether 3H3-mediated enhanced immune memory protection is relevant
to attenuated *Plasmodium* parasite vaccination, we immunized mice with irradiated blood stage *Py* parasites
and subsequently administered either 3H3 or rIgG on days 5 and 7 p.i. On day 30 p.i, vaccinated mice were
challenged with *Pb.ANKA* and we observed superior protection and MBC recall in 3H3-treated, radiation
attenuated parasite vaccinated mice (**Fig. 3G,H**), which aligns with our results in mice infected with live
Py parasites. Collectively, our data reveal a critical pathway that enhances both infection- and vaccination-
induced MBC recall responses following exogenous 4-1BB stimulation.

Enhanced immune memory responses following 4-1BB ligation are MBC-intrinsic

Pathogenic infections or vaccinations typically establish GC-driven protective immune memory responses
that are mediated by the presence and function of MBCs and plasma cells (PC)(13). The elevated class-
switched, parasite-specific antibody titers measured between d0 and d5 p.i following *Pb.ANKA* challenge
(**Fig. 3**) in 3H3-treated mice is suggestive of an enhanced memory response that may result from either an
elevated number of MBCs or an enhanced differentiation capability of MBCs, both of which will result in
larger number of antibody secreting PCs. Enumeration of splenic MBCs showed a significantly reduced
number of both polyclonal (**Fig. 4***A, B)* and antigen (MSP1$_{-19}$) specific fractions (**Fig. 4***C, D)* in *Py*-infected
3H3- treated mice, compared to controls on 90 dpi. Further characterization of these MBCs based on CD73,
CD80 and PDL2(14, 15) also revealed that 3H3-treated mice harbored significantly lower numbers of each
of these subsets compared to control mice (**Supplemental Fig. 4***A-F*). Despite the lower steady state
numbers of MBC in 3H3-treated mice, we observed that there was a significantly larger number of antibody
secreting PCs in the bone marrow following *Pb.ANKA* challenge (**Fig. 4***E),* suggesting that MBCs from
3H3-treated mice have a superior per-cell recall potential compared to MBC in rIgG-treated mice. To
directly test the per-cell protective and functional capacity of both MBC populations, we flow-sorted MBCs
from 3H3 or rIgG-treated mice and transferred an equal number into separate groups of B cell deficient
(μMT) mice that were infected with *Py* the following day. We observed that *Py*-infected μMT mice that
received MBCs from 3H3-treated donor mice exhibited lower parasite load compared to those receiving
MBCs from rIgG-treated donor mice (**Fig. 4***F*), which further correlated with higher serum antibody titers
(**Fig. 4***G*). These data collectively show that MBCs derived from 3H3-treated mice exhibit intrinsically
superior function and capacity to differentiate into PCs that can mediate protection against *Plasmodium*
challenge.
In this study we report that exogenous 4-1BB ligation during primary *Plasmodium* infection while
derailing early GC responses, enhances humoral immune memory during experimental malaria. These data
are in contrast to the paradigm that the quality of the effector GC response predicts the subsequent memory
response. We show that that the beneficial effects of 4-1BB ligation (enhancement of memory response)
are independent of the early parasite burden and inflammatory milieu that follows. Our experiments
suggests that 4-1BB expression on CD4 T cells is required for this enhanced protection (*unpublished*
observations, Caloba and Vijay) and that it is at least partially contributed by CD4 T cell-intrinsic EOMES
expression (**Fig. 3***B, C*). Our observation that 3H3-treated mice begin with fewer MBCs of all subtype s
(**Supplemental Fig. 4***A-F*), but do differentiate into greater number of antibody-secreting PCs upon antigen
reexposure, strengthens our prediction that exogenous 4-1BB stimulation imprints a superior MBC program
on B cells. In summary we show that despite a delayed and rather inefficient effector immune response,
exogenous 4-1BB stimulation has the potential to enhance humoral immune memory in a CD4 T cell
dependent manner. Our findings reported here are especially relevant in the context of the anti-malarial
vaccine, RTS,S which currently offers only limited efficacy (~30%)(2). The possibility of using such an
immunomodulatory approach following vaccination may enhance immunogenicity and overall efficacy of
the vaccine.
239
240

References

241
242
1.   WHO. World malaria report 2022. Geneva: World Health Organization;2022.
2.   Laurens, M. B. 2020. RTS,S/AS01 vaccine (Mosquirix): an overview. *Hum Vaccin Immunother*
16: 480-489.
3.   Tran, T. M., S. Li, S. Doumbo, D. Doumtabe, C. Y. Huang, S. Dia, A. Bathily, J. Sangala, Y.
Kone, A. Traore, M. Niangaly, C. Dara, K. Kayentao, A. Ongoiba, O. K. Doumbo, B. Traore, and P. D.
Crompton. 2013. An intensive longitudinal cohort study of Malian children and adults reveals no
evidence of acquired immunity to Plasmodium falciparum infection. *Clin Infect Dis* 57: 40-47.
4.   Butler, N. S., J. Moebius, L. L. Pewe, B. Traore, O. K. Doumbo, L. T. Tygrett, T. J. Waldschmidt,
P. D. Crompton, and J. T. Harty. 2011. Therapeutic blockade of PD-L1 and LAG-3 rapidly clears
established blood-stage Plasmodium infection. *Nat Immunol* 13: 188-195.
5.   Kurup, S. P., N. Obeng-Adjei, S. M. Anthony, B. Traore, O. K. Doumbo, N. S. Butler, P. D.
Crompton, and J. T. Harty. 2017. Regulatory T cells impede acute and long-term immunity to blood-stage
malaria through CTLA-4. *Nat Med* 23: 1220-1225.
6.   Zander, R. A., N. Obeng-Adjei, J. J. Guthmiller, D. I. Kulu, J. Li, A. Ongoiba, B. Traore, P. D.
Crompton, and N. S. Butler. 2015. PD-1 Co-inhibitory and OX40 Co-stimulatory Crosstalk Regulates
Helper T Cell Differentiation and Anti-Plasmodium Humoral Immunity. *Cell Host Microbe* 17: 628-641.
7.   Zander, R. A., R. Vijay, A. D. Pack, J. J. Guthmiller, A. C. Graham, S. E. Lindner, A. M. Vaughan,
S. H. I. Kappe, and N. S. Butler. 2017. Th1-like Plasmodium-Specific Memory CD4(+) T Cells Support
Humoral Immunity. *Cell Rep* 21: 1839-1852.
8.   Illingworth, J., N. S. Butler, S. Roetynck, J. Mwacharo, S. K. Pierce, P. Bejon, P. D. Crompton, K.
Marsh, and F. M. Ndungu. 2013. Chronic exposure to Plasmodium falciparum is associated with
phenotypic evidence of B and T cell exhaustion. *Journal of immunology* 190: 1038-1047.
9.   Vijay, R., J. J. Guthmiller, A. J. Sturtz, F. A. Surette, K. J. Rogers, R. R. Sompallae, F. Li, R. L.
Pope, J. A. Chan, F. de Labastida Rivera, D. Andrew, L. Webb, W. J. Maury, H. H. Xue, C. R. Engwerda,
J. S. McCarthy, M. J. Boyle, and N. S. Butler. 2020. Infection-induced plasmablasts are a nutrient sink
that impairs humoral immunity to malaria. *Nat Immunol* 21: 790-801.
10.   Malleret, B., C. Claser, A. S. Ong, R. Suwanarusk, K. Sriprawat, S. W. Howland, B. Russell, F.
Nosten, and L. Renia. 2011. A rapid and robust tri-color flow cytometry assay for monitoring malaria
parasite development. *Sci Rep* 1: 118.
11.   Villarino, N. F., G. R. LeCleir, J. E. Denny, S. P. Dearth, C. L. Harding, S. S. Sloan, J. L. Gribble,
S. R. Campagna, S. W. Wilhelm, and N. W. Schmidt. 2016. Composition of the gut microbiota modulates
the severity of malaria. *Proc Natl Acad Sci U S A* 113: 2235-2240.
12.   Curran, M. A., T. L. Geiger, W. Montalvo, M. Kim, S. L. Reiner, A. Al-Shamkhani, J. C. Sun, and
J. P. Allison. 2013. Systemic 4-1BB activation induces a novel T cell phenotype driven by high expression
of Eomesodermin. *The Journal of experimental medicine* 210: 743-755.
13.   Dhenni, R., and T. G. Phan. 2020. The geography of memory B cell reactivation in vaccine-
induced immunity and in autoimmune disease relapses. *Immunol Rev* 296: 62-86.
14.   Brown, S. L., J. J. Bauer, J. Lee, E. Ntirandekura, and J. S. Stumhofer. 2022. IgM(+) and IgM(-)
memory B cells represent heterogeneous populations capable of producing class-switched antibodies and
germinal center B cells upon rechallenge with P. yoelii. *Journal of leukocyte biology* 112: 1115-1135.
15.   Zuccarino-Catania, G. V., S. Sadanand, F. J. Weisel, M. M. Tomayko, H. Meng, S. H. Kleinstein,
K. L. Good-Jacobson, and M. J. Shlomchik. 2014. CD80 and PD-L2 define functionally distinct memory
B cell subsets that are independent of antibody isotype. *Nat Immunol* 15: 631-637.
286
287

FIGURE LEGENDS

FIGURE 1. Exogenous 4-1BB co-stimulation delays GC response. *A*, WT or 4-1BB KO mice were infected with $1x10^6$ *Py*-parasitized RBCs. Parasitemia was measured by flow cytometry on indicated timepoints (n=5/group). *B-F*, WT mice were infected $1x10^6$ *Py*-parasitized RBCs and treated with 50µg of 3H3 or rIgG on days 5 and 7 p.i. Parasitemia (*B, n=6*/group), splenic GC B cells (*C, n=3*/group), GC-Tfh cells (*D, n=3*/group) was measured on indicated time points using flow cytometry and anti-MSP1$_{-19}$ IgG2b antibody titers were measured by ELISA (*E*) and splenic GCs were visualized using confocal microscopy on indicated timepoints (*F*). *A-D* – 2-way ANOVA and *E*, Student's *t* test used. All experiments are representative of at least two independent experiments.

FIGURE 2. Deletion of CD4-intrinsic *Eomes* rescues GC response following exogenous 4-1BB co-stimulation. *A-C*, *CD4.Ert2.Cre.Eomes$^{fl/fl}$* mice were infected with $1x10^6$ *Py*-parasitized RBCs and were treated with 400µg of Tamoxifen orally on days 2-4 dpi and with 3H3 on 5 and 7 dpi. Parasite load (*A*), splenic GC B cell (*B*) and GC Tfh cells (*C*) were evaluated on indicated time points using flow cytometry. *A-C*, n=3 data pooled from two independent experiments, *Mann Whitney* test used.

FIGURE 3. Exogenous 4-1BB stimulation enhances survival following lethal challenge. WT (*A-B, E-F, G-H*) or *CD4.Ert2.Cre.Eomes$^{fl/fl}$* (*C-D*) mice were infected with $1x10^6$ *Py*-parasitized RBCs (*A-F*) or irradiated $1x10^6$ *Py*-parasitized RBCs (*G-H*) were treated with 3H3 or rIgG as indicated on day 5 and 7 p.i.). Mice in *C, D* were treated with 400µg of Tamoxifen orally on days 2-4 dpi and mice in *E, F* were treated with 100 µL of Artemether (10mg/mL) on 12-14 dpi. Mice were challenged with *Pb.ANKA* on indicated timepoints and survival was monitored. Serum was collected on the day (d0) of *Pb.ANKA* challenge and 5 days after (d5) and ELISA was performed to measure change in antibody titers (*B, C, D, F* and *H*) as a measure of MBC recall. Data pooled from (A-F and H) or representative of 2 independent experiments. Kaplan Meyer survival (*A, C, E* and *G*) or paired *t* test (*B, D, F* and *H*) used.

FIGURE 4. Superior MBCs recall following exogenous 4-1BB stimulation. *A-E*, WT mice infected with $1x10^6$ *Py*-parasitized RBCs were treated with 50µg of 3H3 or rIgG on 5 and 7 dpi. On 90 dpi, splenic polyclonal (*A, B*) and antigen specific (*C, D*) MBCs were evaluated. *E*, different cohort of 3H3 or rIgG-treated mice were challenged with *Pb.ANKA* and bone marrow plasma cell numbers were evaluated using ELIspot assay. *F, G*, MBCs from 3H3 or rIgG-treated mice were flow sorted and transferred into mMT mice, which were infected with *Py* the next day, and parasitemia (*F*) on 12 dpi and serum antibody titers were measured using ELISA (*G*). Data are representative (*A-E*) or are pooled (*n=6*/group) from two independent experiments. Mann Whitney test used.

A

Parasitemia (%)

○ WT
■ 41BB KO

P<0.0001

Py → Days p.i

B

Parasitemia (%)

◇ rIgG
■ 3H3

P<0.0001

3H3/rIgG

Py → Days p.i

C Germinal Center B cells

rIgG   Day 21   3H3

CD95   16.0   1.7

Day 39

CD95   4.0   16.0

GL7

○ rIgG
■ 3H3   P<0.0001

GC B cells (%)

P<0.0001

GC B cells (x 10⁶)

Days p.i

D Germinal Center Tfh cells

rIgG   Day 21   3H3

PD1   9.0   0.4

Day 39

PD1   7.0   9.0

CXCR5

P<0.0001

GC Tfh (%)

GC Tfh (#)

○ rIgG
■ 3H3

P=0.0022

Days p.i

E

IgG2b EPT

n.s   P=0.044

>200X

○ rIgG
■ 3H3

Days p.i

F

rIgG   3H3

Day 21

Day 39

B220 CD4 GL7

A

B GC B cells, 23 dpi
CD4-ERT2-Cre()EOMES fl/fl

C GC Tfh cells, 23 dpi
CD4-ERT2-Cre()EOMES fl/fl

A rIgG 3H3

CD38

10.7   10.2

IgD

B

IgD^-CD38^+ B cells (×10⁶)

P = 0.0159

rIgG   3H3

C rIgG 3H3

MSP1

0.91   1.27

UIS4

D

IgD^-CD38^+ MSP-1⁺ (×10³)

P = 0.0159

rIgG   3H3

E rIgG 3H3

Number of spot per 10⁶ cells

P=0.0079

rIgG 3H3

F

Parasitemia (%)

P=0.0286

P=0.0286

3H3 rIgG PBS

G

Anti MSP1₋₁₉ IgM (OD ₄₅₀)

P=0.0286

P=0.0065

P=0.0022

● 3H3
○ rIgG

1:20 1:40 1:80
dilutions

SUPPLEMENTAL FIGURE 1. A, Relative frequency of CD4 and CD8 T cell that express 4-1BB. Gating strategy for GC B cells (B) and GC-Tfh (C) shown.

SUPPLEMENTAL FIGURE 2. Representative histograms and kinetics of EOMES (*A*) and BCL6 (*B*) expression in CD4 T cell from *Py*-infected 3H3 or rIgG-treated mice shown. *C*, Relative expression of EOMES and BCL6 in CD4 T cells in *Py*-infected 3H3 mice during the course of infection shown. *D*, Representative histogram of EOMES expression in CD4 T cells in *Py*-infected 3H3 or rIgG-treated *CD4.Cre.Bcl6*$^{fl/fl}$ mice. *E*, Representative histogram showing deletion of EOMES in CD4 T cells in *CD4.Ert2.Cre.Eomes*$^{fl/fl}$ mice treated as in Fig.2. *F*, Representative histogram of BCL6 expression in CD4 T cells in *Py*-infected *CD4.Ert2.Cre.Eomes*$^{fl/fl}$ mice and *G*, kinetics of parasite load in Tam treated *Py*-infected *CD4.Ert2.Cre.Eomes*$^{fl/fl}$ mice.

SUPPLEMENTAL FIGURE 3. WT mice were infected with 1x10^6 *Py*-parasitized RBCs and treated with 50 μg of 3H3 or rIgG on days 5 and 7 p.i. and with 100 μL of Artemether (10 mg/mL) on 12-14 dpi and parasite load (*A*), EOMES expression in CD4 T cells (*B*), GC B cells (*C*) and GC-Tfh cells (*D*) measured using flow cytometry.

SUPPLEMENTAL FIGURE 4. WT mice were infected with 1x10^6 *Py*-parasitized RBCs and treated with 50 μg of 3H3 or rIgG on days 5 and 7 p.i. were evaluated for different subsets of polyclonal (*A, B, E*) or antigen-specific MBCs (*C, D, F*).