Title:

Simple, reference-independent analyses help optimize hybrid assembly of microbial community metagenomes

Authors:

Garrett J. Smith[1,2,#], Theo van Alen[1], Maartje van Kessel[1], Sebastian Lücker[1,#]

Affiliations:

[1] Department of Microbiology, Radboud Institute for Biological and Environmental Sciences, Radboud University, Nijmegen, The Netherlands

[2] Department of Microbiology, The Ohio State University, Ohio, USA

[#] Corresponding author

Corresponding authors and contact information:

Smith.10284@osu.edu

s.luecker@science.ru.nl

Abstract

Hybrid metagenomic assembly, leveraging both long- and short-read sequencing technologies, of microbial communities is becoming an increasingly accessible approach, yet its widespread application faces several challenges. High-quality references may not be available for assembly accuracy comparisons common for benchmarking, and certain aspects of hybrid assembly may require dataset-dependent, empirically-guided optimization rather than application of a uniform approach. In this study, several simple, reference-free characteristics – gene lengths and read recruitment – were analyzed as reliable proxies of assembly quality to guide hybrid assembly optimization. These characteristics were further explored in relation to reference-dependent genome- and gene-centric analyses that are common for microbial community metagenomic studies. Here, two laboratory-scale bioreactors were sequenced with short and long read platforms, and assembled with commonly used software packages. Following long read assembly, long read correction and short read polishing were iterated to resolve errors. Each iteration in this process was shown so have a substantial effect on gene- and genome-centric community composition. Simple, reference-free assembly characteristics, specifically changes in gene fragmentation and short read recruitment, explored throughout this process replicated patterns of more advanced analyses seen in published comparative studies, and therefore are suitable proxies for hybrid metagenome assembly accuracy to save computational resources. Hybrid metagenomic sequencing approaches will likely remain relevant due to the low costs of short read sequencing, therefore it is imperative that users are equipped to estimate assembly accuracy prior to downstream gene- and genome-centric analyses.

Background/Introduction

Though the increasing variety of high-throughput short- and long-read (meta)genomic sequencing technologies are only within their first decades of existence, both the sequencing technologies and software development have flourished and has already been implemented to study microbial ecosystems[1–15]. Integrating both short- and long-read platforms for single microorganisms or microbial communities is gaining popularity because they compensate for the

other's weaknesses – shorter reads achieve higher accuracy while longer reads

offer better contiguity[7,11,16–20]. Despite the technical costs decreasing and

technologies improving, hybrid strategies currently are more cost-effective for

recovering high-quality Metagenome-Assembled Genomes (MAGs) from

microbial communities than either platform alone[12]. However, many challenges,

some biological and some technical, remain that may deter users from seeking

relatively complicated approaches like hybrid (meta)genomic sequencing of

microorganisms in complex ecosystems.

Hybrid (meta)genomic assembly implements multiple and/or iterative

processes to help overcome the limitations of the sequencing technologies.

One main strategy is to assemble short reads and then bridge gaps with long

reads, leading to assemblies that are accurate but less contiguous[1,5,8,11,14,18,19].

The other main strategy is to assemble long reads and then iteratively use short

reads to resolve the errors, leading to a more contiguous assembly that retains

errors[1,5,8,11,14,17–26]. In the recent past and near future, as long read accuracy

increases tremendously, this second strategy may become overall more

favorable for the field because, intuitively, better contiguity enables better gene-

and genome-resolved analyses. Most benchmarks for the second strategy

have provided a general set of guidelines for implementing the iterative

processes to improve upon long read assemblies: using several tools is

preferred, long read correction prior to short read polishing is advantageous,

and iterative processes have diminishing returns[4,12,13,17,22,26–31]. These

concepts are incorporated into pipelines for automating the optimization of

hybrid microbial (meta)genomic assemblies[19,29], suggesting they are core

aspects of hybrid metagenomic assembly even for microorganisms.

Intriguingly, despite the existence in literature essentially since the

development of long read sequencing, the iterative processes for fixing errors

in long read assemblies has been less thoroughly investigated. In particular,

one challenge in performing hybrid assembly for a microbial community

metagenome is identifying the number of iterations to maximize the quality of

the data. Generally, there is little consistency of tool choice and the number of

iterations among published studies of microbial communities or other datasets

reconstructed using LR assembly[1–3,5–11,13,14,26,30], and arguably even less

consistency among benchmarking studies[3,11,14,17,18,20–24,26,27,31–43]. While most

studies have determined the approach *a priori*, some evidence suggests that

an unsupervised but empirically-guided approach combining various tools

optimizes the hybrid microbial assemblies because certain tools are better able

to fix certain errors, and some may even degrade the quality by re-introducing

errors[4,17,21,26,31]. This suggests that is very likely that ideal hybrid assemblies for

microbial communities may not be achieved using a universal or standard

methodology, but rather likely varies depending not only on the biology of the

system but also on tool choices.

An additional challenge of applying hybrid assembly to a complex

microbial community is determining the quality of the assembly itself. Typically

during benchmarking, assembly qualities are assessed by comparing to high-

quality references, *e.g.*, count of differences (mis-assemblies), genomic

alignments, or specific marker gene presences[5,12,18,22,23,30,32,33,37,44], which can

lead to poor interpretation of multiple or divergent genomes, as well as high

computational costs. In the absence of high-quality reference genomes, the

characteristics of an ideal hybrid assembly of a complex microbial communities

are less clear. Common, reference-independent statistics for comparing

assemblies, for example contig counts, total base pairs assembled, and L/N50

and similar metrics, might not significantly change at a metagenome assembly

scale during iterative processes that fix relatively small-scale errors[17,26,28,30,31].

Therefore, typical assembly quality assessments are not suitable for many

complex microbial community metagenomic datasets.

Benchmarking of individual bacterial genomes suggested that read

recruitment, and to a lesser degree gene counts and lengths, are useful

reference-free proxy indicators of assembly quality[17]. While there are advanced

statistics, both of gene length distributions and read recruitment profiles are

relatively simple to calculate, and often precede downstream analyses. In

several cases, gene fragmentation and read recruitment profiles, often in

comparison to references, have been used to asses recovered MAG quality or

the accuracy of assemblies themselves[4,8,11–13,15,17,18,22,45], but in many cases

there was no apparent evaluation or optimization of hybrid assembly of

microbial community metagenomes[1,2,5,9–11,14,45]. Given the challenges of

applying hybrid assembly approaches for microbial community metagenomes,

and the likelihood that no universal approach works best for all datasets,

characteristics that are reference-independent and relative to the dataset may be the most suitable to estimate quality and empirically guide optimization.

Here, we examined multiple reference-independent and -dependent assembly characteristics in order to determine which would be effective for empirically optimizing hybrid metagenomic assembly of a uncharacterized, complex microbial communities. Two long-term laboratory-scale nitrifying bioreactors were sequenced using both Illumina MiSeq (short-read, SR) and Oxford Nanopore Technology (long-read, LR) platforms and assembled with multiple programs to allow for initial biological and computational variation. LR assemblies were then corrected with LRs and polished with SRs using prevalent tools in the field with low computation demands. Rather than assess the efficacy of tool, we sought to (1) observe the impact of iterative processes on community reconstruction, and (2) identify reference-independent assembly characteristics to help estimate assembly accuracy to optimize the approach. We first established that there is significant variation in community composition during these iterative processes as a result of the errors being fixed, but these patterns largely match published studies. We then demonstrated that simple, reference-independent assembly characteristics, for example coding gene counts and bps or SR recruitment, not only follow these same patterns but also were robustly correlated with more advanced reference-agnostic metrics, thus serving as practical proxies for assembly accuracy and community reconstruction. Finally, we offer practical guidelines to aid users for empirically optimizing hybrid assembly of microbial community metagenomes.

Methods

The tools, their purpose, and the rationale for or advantages of their use in this study are described in Text S1, and summarized in Table S1.

Long and short read sequencing

Biomass was collected in November 2020 and March 2021 from long-term, autotrophic nitrifying enrichment cultures maintained in either oxygen- (OLR) or nitrogen-limited (NLR) conditions in a tandem laboratory-scale bioreactor system inoculated in 2015 with wastewater treatment plant activated sludge from the Bavaria Brewery in Lieshout, The Netherlands (51.518666 N,

5.613009 E). The cultivation of these bioreactors and the reconstruction of their microbial communities is an on-going project and will be published separately in a following study. Genomic DNA was extracted from the biomass using a conventional N-cetyl-N,N,N,-trimethyl ammonium bromide (CTAB) protocol in 2020, and the Powersoil DNA Isolation kit (Qiagen) in 2021 with minimal modifications to the manufacturer's directions to reduce DNA shearing such as inverting rather than vortexing or pipetting to mix. wo DNA isolation protocols were used because it is standard in our group due to strong evidence that it introduces sufficient bias to aid differential coverage binning[46–48].

In total, 1 ng of DNA for both the OLR and NLR reactors was used to prepare a library using the Nextera XT kit (Illumina, San Diego, California U.S.A.) according to manufacturer's instructions. After quality and quantity check of the libraries, they were paired-end sequenced (2x 300bp) using the Illumina MiSeq sequencing machine and the MiSeq Reagent Kit v3 (San Diego, California USA) according to manufacturer's protocols. Oxford Nanopore Technologies (ONT) sequencing was done with 840 and 1670 ng DNA for the OLR and NLR reactors, respectively, after library preparation using the Ligation Sequencing Kit 1D (SQK-LSK108) and the Native Barcoding Expansion Kit (EXP-NBD104), according to the manufacturer's protocols (Oxford Nanopore Technologies, Oxford, UK). The libraries were loaded on a Flow Cell (R9.4.1) and sequenced on a MinION device (Oxford Nanopore Technologies, Oxford, UK), according to the manufacturer's instructions. Guppy (version 4.0.11)[49] was used to basecall fast5 files using the dna_r9.4.1_450bps_hac.cfg model, both provided by Oxford Nanopore Technologies.

Raw or basecalled sequencing reads for both bioreactors and technologies are available at NCBI via BioProject PRJNA1005948 as the following BioSamples: SAMN37004618, raw MiSeq reads for the OLR; SAMN37004620, raw Oxford Nanopore Technologies reads for the OLR; SAMN37004619, raw MiSeq reads for the NLR; SAMN37004621, raw Oxford Nanopore Technologies reads for the NLR.

Long-read, short-read, and hybrid microbial community metagenomic assembly

169 An overview of the experimental design is shown in Fig. 1. All computational

170 programs were employed with default settings unless explicitly stated. Generic

171 example code is available in File S1.

172   Sequencing yielded 3.1-4.6 Gbp per library (Table S2) following read

173 trimming and length and quality control using BBduk (BBtools version 37.76)[50]

174 with a minimum phred score 18 and length of 200 bps for the MiSeq reads, and

175 porechop (version 0.2.3_seqan2.1.1)[51] with minimum split length of 3000 bps

176 for the ONT reads.

177   Three common SPAdes-dependent programs were used to generate

178 SR-only, LR-only, and/or SR-first hybrid, *i.e.*, short read assembly followed by

179 connecting contigs using long read alignments, assemblies: hybridSPAdes

180 (version 3.15.4)[40,52] known for high-quality metagenomic assemblies, Unicycler

181 (version 0.4.9b)[19] with utilities for optimizing SPAdes to enable recovery of

182 circular and high-quality single genomes, and OPERA-MS (version 0.9.0)[3]

183 capable of automated refinement of high-quality individual genomes within a

184 multi-species metagenome (hybrid only). Note that Unicycler, when given only

185 long reads, does not use SPAdes and instead uses miniasm[38] and Racon[27].

186 The k-mer list generated by automated selection during Unicycler hybrid

187 assembly was used for hybridSPAdes assemblies. While the focus here was

188 optimizing a hybrid approach that leveraged the continuity of the long-read

189 assemblies, these SPAdes-dependent assemblies served as a baseline for

190 comparing SR and LR assembly of these datasets.

191   Two programs were used for assembly of LR data: Canu (version 1.8)[34]

192 that performs read correction prior to assembly and is generally thought to

193 result in more accurate assemblies, and Flye (version 2.9-b1768)[33] that

194 performs correction of the assembly using the input reads. Both assemblers

195 were used in three different ways: (1) default settings ("def") developed for

196 single-genomes, (2) metagenomic settings ("meta") to improve assembly of

197 sequences with uneven depths[2,33], and (3) metagenomic settings for

198 approximately even bp sub-samples ("sub") of the reads to artificially reduce

199 the sequencing depth and possibly uncover biological variation. Long reads

200 were sub-sampled into 12 read pools with approximately the same quantity of

201 bps using the "subsample" utility of Trycycler (version 0.4.1)[22]. All assemblies

were filtered to a 4 kbp minimum contig length using BBtools utilities prior to
any downstream analyses.

Long read correction and short read polishing

Consensus correction of individual LR "meta" assemblies was performed
using Racon (version 1.3.1)[27] up to 10 iterations, the greatest number of
iterations in a hybrid assembly approach that we identified among literature[37].
For each iteration of Racon correction, minimap2 (version 2.17-r941)[53] was
used with default settings to recruit LRs to the original assembly or previous
iteration's corrected assembly to generate the overlap information. Contigs that
were not corrected were retained with the optional flag "-u". SR pile-up polishing
of LR "meta" assemblies was performed using Pilon (version 1.23)[54] to 10
iterations after 0 (original assembly), 2, 5, and 10 rounds of LR correction.
These stages were chosen to include common (0 and 2) LR correction
endpoints found in literature and online guidance, and extensive (5[22] and
10[37,55]) LR correction endpoints that would help demonstrate the maximum
possible value of this process. Note that many benchmarks focus on individual
microbial genomes, and particularly studies exploring the quality during iterative
processes, and therefore it seemed plausible that additional iterations may aid
in improving assembly quality. We did not test SR polishing after 1 iteration of
LR correction because it was rarely observed in published literature or online
guidelines, and essentially never when Racon was used for LR correction. For
each iteration of SR polishing, BBmap (BBtools version 37.76) was used to
recruit SRs to the original assembly, Racon-corrected assemblies, or previous
iteration's polished assembly, using a 97.5% identity filter and retaining all
ambiguous alignments. Because we found no examples of LR correction after
SR polishing and it is expected to re-introduce errors, this was not performed.

Assembly characteristics determination

Programs for estimating quality were used with default settings except when
explicitly stated otherwise. Assemblies were compared using metaQuast
(version 5.0.2)[56] without references to examine the distribution of contig counts
and lengths. Recruitment of SRs to the assemblies to quantify aligned reads
and ambiguity was performed using BBmap (BBtools version 37.76) with

retaining only perfect mappings of paired reads and ambiguous mappings
randomly assigned. Open reading frames (ORFs) were predicted using
Prodigal (version 2.6.3)[57] using the "meta" procedure for quantification of gene
counts and lengths, and as input for downstream analyses of assembled genes.
The "meta" procedure of Prodigal was implemented prior to downstream
analytical pipelines because it led to increased marker gene recovery
estimates, though following similar trends (data not shown), compared to the
outputs of the analytical pipelines that assume inputs are single genomes and
therefore run prodigal in "single" mode by default. Phylogenetic marker gene
recovery, fragmentation, and redundancy were estimated by Benchmarking
Universal Single-Copy Orthologs (BUSCO, version 5.1.2)[58] using "genome"
mode and only the "bacteria_odb10" lineage, as well as CheckM (version
1.1.3)[59] using the taxonomy workflow for the domain "Bacteria". Gene
fragmentation of entire assemblies was estimated by comparing to a DIAMOND
(version 0.9.31)[60] database of the uniref50 (downloaded 2023-03-01)[61] dataset
using IDEEL (downloaded 2023-05-13)[13]. To more concretely associate
potential complete genome contigs with the domain bacteria, circular and long
(>1 Mbp) contigs were also analyzed individually using the Genome Taxonomy
DataBase toolkit (GTDB-tk, version 1.6.0 with reference database version
r202)[62]. Based on the GTDB-tk output for circular contigs alone, there were
members of at least 6 and 5 distinct Bacterial phyla present in the OLR and
NLR, respectively (data not shown). Microbiomes with abundant Archaea or
Eukaryotes may need to adjust these taxonomically constrained pipelines to
better suit their ecosystem.
260
Automated binning and beta diversity
Programs for automated binning and estimation of genome quality, as well as
identification and read depth calculation of marker genes, were used with
default settings except where noted otherwise.
Both LRs and SRs from both bioreactor datasets were used for read
depth calculation of contigs using the mmlong (version 0.1.2)[63] utility
readcoverage2. Automated binning using composition (*i.e.*, tetranucleotide
frequency) and coverage (*i.e.*, read depth) was then performed using Metabat2
(version 2.12.1)[64]. The quality of automated bins was estimated using CheckM

(version 1.1.3)[59], with cutoffs of >50% completeness and <10% contamination
scores as "medium quality" (MQ). These oversimplified, and somewhat low,
thresholds are no longer *en vogue*, but represent computationally simple, rapid,
and bulk estimates compared to contemporary, thorough requirements that
include other information like rRNA presence and tRNA counts[65].

The RNA polymerase subunit B (*rpoB*) is a protein-coding gene typically
found only once in a genome and is universally conserved among Bacteria,
Archaea, and Eukarya, and therefore is tractable to serve as a marker gene for
individual species to complement the often poorly assembling 16S rRNA gene
for gene-centric phylogenetic analyses of metagenomes. *RpoB* genes were
identified in assembled contigs using hmmer (version 3.1b2)[66] with the
available model and thresholds for the Protein FAMily (pfam) identifier
PF04563.15 (downloaded 2020-06-09). To observe beta-diversity, *rpoB*-
containing contigs throughout the iterative correction and polishing processes
were compiled for the each bioreactor and assembler separately, and then the
LRs and SRs from both bioreactor datasets were used for read depth estimation
for *rpoB*-containing contigs using the mmlong (version 0.1.2)[63] utility
readcoverage2. The Bray-Curtis abundance-rank dissimilarity between each
read set was calculated for subsequent two-axis Non-Metric multi-Dimensional
Scaling (NMDS) performed using vegan (version 2.5-7)[67] in R (version 4.1.2)[68],
and the values were then manually mean-centered and scaled for better
comparability. These NMDS analyses were unable to converge after 50 tries
due to low stress, but here the species scores were used to view the clustering
and trajectories of the *rpoB*-containing contigs' "species scores" for each
assembly rather than the four reads' "site scores".

Further data analysis and visualization
Logs and outputs were mined for data using various bash commands. Data
were ultimately imported into R (versions 4.1.2)[68] for analysis and visualization
relying primarily on tidyverse (version 1.3.1)[69]. Most calculations were
additionally normalized to contig length in Mbps to make bioreactors and
assemblers more comparable. No specific code was developed for this project
due to the focus on applying end-user tools.

Results

Short read, long read, and hybrid assembly baselines

SR sequencing yielded 7.56 million paired-end reads totaling 4.06 Gbp, and 8.73 million paired-end reads totaling 4.60 Gbp of data after low quality read removal and base trimming for the OLR and NLR, respectively. LR sequencing yielded 318 thousand reads totaling 3.13 Gbp, and 416 thousand reads totaling 4.01 Gbp of data after low quality read removal and base trimming data for the OLR and NLR, respectively (Table S2). Reads were either assembled alone (SR-alone or LR-alone) or used as inputs for automated SR-first hybrid assembly. There were substantial differences between assemblies of the bioreactors of approximately 63 total assembled Mbp on average (Fig. S1), which we infer was due to the greater sequencing depths for the NLR compared to the OLR rather than any biological differences.

In addition to biological and/or sequencing depth effects on the assemblies, the assembly approach implemented by the different programs had a substantial effect on the assembly outcome. This was apparent across many aspects of these assemblies: LR assemblies were larger in terms of total bps (on average 1.35-fold), they were less fractured such that a greater fraction of bps were on larger contigs (1.69-fold), and they yielded more circularized contigs (2.34-fold for count and 5.5-fold for total bps) (Text S1, Figs. S1, S2A, S3). However, probable complete microbial genomes, *i.e.*, a single contig greater than 0.5 Mbp and ideally circular, reconstructed with SR and SR-first hybrid assemblies tended to have higher completeness scores on average (1.73-fold for long contigs and 1.38-fold for circular, Text S1, Fig. S2B, Fig. S3). Thus, baseline assemblies recapitulated most benchmarking studies showing that LR assemblies lead to better contiguity, but SR assemblies tend to be more accurate[1,5,8,11,17,21]. Still, the contiguity of the LR assemblies offers advantages for microbial community reconstruction with sufficient LR correction and/or SR polishing to resolve mis-assemblies.

From here, we explored these metagenome-optimized LR assemblies over the iterative LR correction and SR polishing iterations to ultimately determine the ideal assembly for these datasets with empirical information. We assessed the microbial composition and its variability using standard gene- and genome-centric analyses like beta-diversity and automated bin recovery. We

then further analyzed characteristics that should noticeably change, in contrast

to for example the size and contiguity of the assembly, and that ideally would

also be reference-free: reduction in gene fragmentation because errors in LR

and their assemblies lead to frameshifts that fracture genes, and increase in

SR recruitment because these more high-accuracy reads should recruit to more

accurate assemblies. Additionally, we followed automated bin recovery and

single phylogenetic marker gene beta diversity to see how well these simple,

reference-free characteristics may serve as indicators of community

reconstruction.

Marker gene beta diversity

Complementing genome-centric approaches, many studies of microbial

communities applying hybrid metagenomic assembly performed gene-centric

analyses[2,5–10,13,14], often as a means of inferring metabolic capabilities and

therefore ecosystem roles, but can also help access less well assembled

community members to better assess extant diversity. While genome recovery

has been examined as proxies for microbial metagenomic assembly accuracy,

there is substantially less effort put into examining the remainder of the

community members and how this might impact downstream analyses of the

communities. The RNA Polymerase subunit B (*rpoB*) gene is a phylogenetic

marker for all domains of life that was used here for beta-diversity clustering

analysis by using their SR recruitment profiles via Non-Metric multi-

Dimensional Scaling (NMDS).

As expected given the issues with LR assembly, error-fixing substantially

improved marker gene recovery but surprisingly decreased read depth. Initial

LR assemblies and assemblies with LR correction alone contained fewer (2

total genes) and more fractured (1/3 the mean length) *rpoB* genes compared to

assemblies after the first few iterations of SR polishing (Text S1, Fig. S4). While

read depths stabilized due to SR polishing, they unexpectedly decreased on

average (Text S1, Fig. S5). We speculate that this phenomenon was caused

by the identification of additional *rpoB* genes on low abundance contigs once

SR polishing made these genes sufficiently intact to be identified.

Both increased *rpoB* recovery and changes to read depth profiles had

clear effects on the trajectories of the communities during error-fixing

processes, thus influencing apparent beta-diversity. Assemblies with only LR

correction were noisy and remained distant from one another, compared to the

convergence observed due to the first few SR polishing iterations (Fig. 2). In

most cases, assemblies without LR correction were distant from assemblies

with both LR correction and SR polishing, highlighting the beneficial impact of

LR correction that may only be apparent after SR polishing (Fig. 2).

Unfortunately, also in most cases, communities that converged during SR

polishing were still distinct from those with different preceding LR iterations,

indicating that this impacts the community's even with SR polishing (Fig. 2).

These results show that iterative LR correction and SR polishing processes

ultimately impact downstream community reconstruction analyses.

Automated bin recovery

A primary goal for many microbial community metagenomic sequencing

approaches is to reconstruct high-quality, ideally essentially complete genomes

of as many of the members as possible[1–10,12,13]. Hybrid assembly of microbial

metagenomes has already been employed to recover microbial genomes from

a variety of ecosystems from waste sludge to human guts, and more[1–10,12,13].

To maximize the value of the genomes, the assembly itself needs to be

accurate because both binning of contigs into genomes and their resulting

quality, estimated via marker gene identification, depends on the accuracy of

the assembly. In some cases, genome recovery has already been used to

demonstrate or compare the accuracy of LR and/or hybrid assemblies of

microbial (meta)genomes[1,7,10,15,18,33]. Here, assemblies at each stage of LR

correction and SR polishing were automatically binned, and both the

assemblies and bins were assessed with CheckM to estimate the completion

of microbial genomes and estimate the redundancy by counting the 104

bacterial marker genes and reporting the copy number of each gene and the

estimated completion and contamination percentages.

Error-fixing of LR assemblies was required for genome recovery, and in

particular SR polishing greatly improved automated bin recovery. Initial

assemblies led to the recovery of few medium-quality or better (MQ) automated

bins, only 9.5 on average (Text S1). By following contigs across bins, we noted

that binning prior to LR correction and SR polishing led to worse automated

binning outcomes compared to binning after iterative error-fixing processes (data not shown). LR correction increased the number of MQ bins by three to 12.3 on average, but SR polishing alone almost doubled the number of MQ bins, and in combination with LR correction more than doubled the initial MQ bin count on average (Fig. 3, Text S1, Fig. S6). Additionally, the mean completeness scores for MQ bins remained in the low 70% range after only LR correction but increased to over 80% as a result of SR polishing (Fig. 3, Text S1, Fig. S6), indicating an increase in the quality of recovered genomic contents by combining LR correction and SR polishing. Furthermore, the substantial increases in MQ bin count and quality due to SR polishing were matched by large increases in the quantity of both number of contigs (7.17-fold) and bps (1.92-fold) contained in MQ bins (Fig. 3, Text S1, Fig. S6), indicating that SR polishing improved the ability for contigs to be binned. However, automated bin recovery metrics remained somewhat noisy and often never reached a clear plateau (Fig. 3), suggesting that genome recovery may not be a robust assembly quality indicator. In summary, automated bin analyses suggest that the iterative LR correction and SR polishing processes strongly influence genomic recovery.

Bacterial marker gene redundancy estimated for the full assemblies followed similar patterns as automated bin recovery, and therefore may serve as a proxy to save computational resources. In contrast to bin recovery, marker gene recovery showed more clearly curves that saturated throughout the iterative error-fixing processes (Fig. S6). The estimated copy number of marker genes, as well as redundancy, *i.e.*, the sum of CheckM completion and contamination scores, reached their maximum values after two iterations of SR polishing, with greatest redundancy after both LR correction and SR polishing (Fig S6, Text S1). In fact, the redundancy of the assemblies was strongly correlated to the number of MQ bins recovered (adjusted $R^2 \gtrsim 0.85$, p $<<0.05$, Fig. S7, Text S1). Intriguingly, slope of these regressions was approximately 0.01 (Fig. S7), or 1 genome per CheckM score of 100%, implying a near equal relationship between marker gene redundancy and complete genome recovery. However, the distance from an idealized line (Fig. S7) suggests that this hybrid assembly approach was still unable to reconstruct the genomes of all members of the microbial community. Yet, given the correlation with MQ bin recovery but

an obvious plateau in redundancy, assembly-level marker gene recovery could

serve as a preliminary indicator of genomic yield during the iterative processes

without the need for automated binning.

Gene fragmentation

Small scale errors (*i.e.,* insertions, deletions, and, to a lesser extent,

substitutions) in LRs are known issues for using these platforms that cause

gene fragmentation in the resulting assemblies[7,12,13,17,20,23,26,30,31,45,54,70].

Theoretically, as errors are fixed during the iterative LR correction or SR

polishing processes then gene fragmentation should decrease, *e.g.*, fewer

genes and more bps in them, and ultimately stabilize. In benchmarking studies,

this is often indirectly measured by comparing differences in gene counts or

marker gene inventories (*i.e.*, completion scores), or have already

demonstrated that directly assessing gene lengths works as a reasonable proxy

for the accuracy of hybrid assemblies of microbial (meta)genomes[4,7,11–

13,17,18,22,37,45]. As a precursor to many downstream analyses, coding gene

identification with programs like Prodigal allow direct estimates of coding gene

lengths, while other programs like IDEEL and BUSCO can estimate gene

fragmentation by comparison to reference databases and are commonly used

by benchmarking or other comparative studies[4,12,13,22,30,37,71].

Gene fragmentation substantially improved and plateaued during

iterative error-fixing processes. Initial LR assemblies varied due to both

bioreactor and assembler, consistent with the baseline information (Text S1).

On average, gene fragmentation did not markedly improve, either in reducing

the number of coding genes or increasing the quantity of bps in coding genes,

due to LR correction alone (Text S1, Fig. S8). However, it did substantially

improve after just two iterations of SR polishing (0.81-fold for gene count, 1.12-

fold for bps in genes) after which these assembly statistics appeared saturated

(Text S1, Fig. S8, Fig. S9). While small, there was a non-negligible advantage

to LR correction preceding SR polishing (Text S1, Fig. S8, Fig. S9). The same

patterns were noticeable not only for the entire assembly, but also for several

relevant assembly fractions principally ready for downstream analyses –

circular and long contigs that may represent essentially complete microbial

sequences, and medium quality or better automated bins (Text S1, Figure S8,

Fig. S9). As expected based on both the theory behind the iterative error fixing
processes and most published studies assessing this information[4,11–
13,17,18,22,37,45], gene fragmentation was improved primarily due to the first few
iterations of SR polishing.

Comparison of coding genes agnostically to reference databases
showed similar patterns. After just two iterations of SR polishing, assessing the
entire assemblies with BUSCO indicated that bacterial marker genes were no
longer found only fragmented (Text S1, Fig. S9), and IDEEL indicated that the
number of coding genes within 5% of their most similar reference sequence's
length considerably increased (2.7-fold, Text S1, Fig. S10). All gene length
analyses followed similar patterns, and linear regression demonstrated that the
number of coding genes within 5% of their most similar reference gene length
was strongly correlated to the total number of base pairs in coding genes (R^2
$\geqq 0.94$, p-adj <<0.05, Fig. 4). We acknowledge that the resulting data were
unevenly distributed across both dependent and independent variables in this
analysis because of the large differences due to, and the inflated sample sizes
of, SR polishing but the correlations hold when looking at only the first three SR
polishing iterations as well, albeit weaker (adjusted R^2 0.6-0.75, p-value <<0.05,
data not shown). Additionally, similar reference-independent assembly
characteristics including total number of coding genes, as well as median count
of coding genes and bps in coding genes normalized to contig length, were also
correlated (adjusted R^2 >0.7, p-value <<0.05, data not shown). Therefore a
less-resource demanding analysis such as gene length summarization alone is
a practical proxy for more demanding analyses like gene fragmentation.

Short read recruitment
Multiple iterations of LR correction and SR polishing are often performed in
benchmarks and published studies because it is expected that newly fixed
errors allow different read pools to align. Then theoretically, SR recruitment
would reach a maximum, stable pool as the assembly approaches high
accuracy because fewer errors are fixed each iteration leading to fewer
changes in read alignments. Several studies have already shown a relationship
between microbial (meta)genome quality and read recruitment[4,11,12,15,17,22,45].
Additionally, in a community where multiple strains co-exist and/or species

encode duplicated genes, the quantity ambiguously recruited reads (*i.e.,*
aligning to two or more positions) should decrease as the minor sequence
differences are resolved. Here, information-rich outputs from BBmap were
analyzed for the total number of aligned SR bps, assembled contigs to which
no SRs aligned, and ambiguously aligned SRs to observe the magnitudes in
which these features change during LR correction and SR polishing.

Consistent with expectations and gene fragmentation results, increases
to SR recruitment increased greatly due to SR polishing and then saturated
after the first few iterations. The bioreactor had little apparent effect on SR
recruitment to the initial assemblies, in contrast to the assembler program and
(Text S1). These data, along with the larger proportion of SRs that aligned
ambiguously (2.02-fold, Text S1), suggests that the Flye assemblies may be of
higher quality or possibly capable of reconstructing similar yet distinct strains.
On average, LR correction led to approximately 25-35% increases in total SR
bps recruited, SR bps recruited normalized to contig length, and SR alignment
ambiguity, but only about a 10% reduction in contigs lacking SR recruitment
(Text S1). In contrast, SR polishing greatly improved SR recruitment across all
measured features, including increased total SR bps recruited (3-fold) and
median SR bps recruited normalized to contig length (10-fold), fewer contigs
lacking any SR recruitment (30% reduction), and a smaller fraction of
ambiguously aligned SRs (26% reduction, Fig. 5, Text S1). There was a clear
plateau in SR recruitment by 3 iterations of SR polishing, and a noticeable
improvement due to LR correction but only after some SR polishing. These
observed trends match common implementation of at least 2 SR polishing
iterations[2,9–13,15,17,22,29,33,45,54]. The saturation patterns of SR recruitment also
appeared consistent with the decreases in changes made during SR polishing
(Fig. S11), strengthening the link between using SR to improve assembly
accuracy and SR recruitment profiles themselves. In summary, SR recruitment
followed similar trends as all other assembly characteristics, and as a quick and
common predecessor to many downstream analyses, offers a practical
estimation of assembly accuracy during these iterative processes to help
empirically determine their endpoints.

Discussion

Correction and polishing affect gene- and genome-centric community reconstruction

Hybrid assembly approaches, leveraging complementary beneficial attributes of both LR and SR platforms to overcome their limitations, are already being used to study microbial communities in various evniromenments[1–15]. Assembly of LRs offers a major advantage over SR data alone, due to the ability to achieve greater contiguity, however, it comes at a significant cost due to lower accuracy[1,5,8,11,14,17,21–24,32]. Each study, from benchmarking to community reconstructions, applied different methods in their hybrid assembly approaches, and it's probable that the biological and technical complexity of microbial community metagenomes limits the implementation of uniform methods across different studies. One of the least consistently applied, and likely dataset- and tool-dependent, aspects of applying a hybrid assembly approach for a microbial community metagenome is the iterative processes of LR correction and SR polishing to improve the accuracy of LR assemblies.

Few studies have explored the impacts of these iterative processes, despite being implemented by most of them. Generally, the consensus we understood was that while SR polishing offers the greatest improvements, some LR correction is advantageous, and both offer diminishing returns[13,17,21,26,31,37]. We first showed that microbial community composition changes during these iterative processes, which was, as far as we know, the first demonstration of this phenomenon. We further showed that genome recovery was also greatly impacted by these processes, which combined with marker gene beta diversity, demonstrates that the exact number of iterations for these processes has a significant impact on reconstruction, and thus also interpretation, of the microbial communities. Fortunately, results here followed similar patterns with each other, and also in many aspects mirrored published studies comparing different aspects of hybrid assembly across multiple tools and over several iterations[1,2,4,7,10,12–15,17,18,22,26–31,37]. However, there is not only substantial methodological variation between published approaches and the performed here, but also biological variation between each dataset which probably precludes the development and application of a universal or standardized approach for applying hybrid metagenomics to microbial

communities. Therefore, it is likely that each dataset needs to be empirically
evaluated and its hybrid assembly approach optimized.

Simple, reference-independent assembly characteristics as proxies for the
quality of hybrid assemblies of microbial community metagenomes
Further complicating hybrid assembly for microbial community metagenomes
is the lack of a clear assessment of quality. While mis-assembly and contiguity
are the most common metrics, these assembly characteristics either require
high-quality reference genomes that may not exist for poorly characterized
ecosystems, or would not substantially change while optimizing LR correction
or SR polishing[5,12,17,18,22,23,26–28,30–33,37,44]. Therefore, assembly characteristics
appropriate for a microbial community should be dataset-dependent and
expected to change as assembly accuracy also changes.

Here, several assembly characteristics were tracked during the iterative
processes to determine which were suitable proxies for assembly quality, and
to help optimize hybrid assembly of uncharacterized microbial community
metagenomes. Specifically, gene fragmentation and SR recruitment were
focused on during LR correction and SR polishing because these features have
already been proposed as suitable proxies for hybrid assembly of microbial
genomes[4,11–13,17,22,45]. These simple assembly characteristics followed patterns
consistent many benchmarking and comparative studies[4,12,13,17,22,26–31]: (1) the
greatest improvements occurred within the first three iterations of SR polishing;
(2) LR correction improved the accuracy of the assemblies but was not always
apparent until after SR polishing; (3) beyond five iterations LR correction or SR
polishing offered severely diminished returns. Furthermore, simpler and
reference-independent assembly characteristics were strongly correlated with
more advanced, reference-dependent characteristics, *i.e.*, gene counts,
lengths, and bps in genes were all correlated with the gene fragmentation
profiles, as well as automated bin recovery and marker gene beta diversity
trajectories. Thus, these simple, reference-independent assembly
characteristics are reliable indicators of assembly accuracy and can be used to
empirically optimize hybrid assembly of uncharacterized microbial community
metagenomes.

 Unexplored factors and limitations of this study

The approach in this study was not without issue, and therefore here we

describe expected deviations from our approach that may be necessary for

certain users or datasets.

First, recent technological advances for both LR sequencing and

application may ease the approach for integrating hybrid metagenomic

assembly of microbial communities, but this study was performed using older

sequencing chemistry and basecalling algorithms. Some differences remain,

but both major LR platforms, Pacific Biosciences (PacBio) and Oxford

Nanopore Technologies (ONT) have achieved large improvements in chemistry

and basecalling that have increased both depth and accuracy of LR

sequencing[4,11,12,24,45,72,73]. Therefore, not only will LR-alone assemblies be

more accurate as a result of these advances, they already may be sufficient for

retrieving HQ genomes without SR polishing[12,74]. However, resources may

restrict users to older and more accessible technologies, or projects may

already have older datasets that still need analysis that integrate short and long

read technologies. Additionally, for the foreseeable future, SR complements to

LR datasets will likely continue to increase the sample count for differential

coverage binning. Furthermore, SRs still appear to aid assembly accuracy for

high-quality microbial genomes reconstructed from LR[75]. Therefore, combining

both sequencing technologies will be attractive as a cost-effective approach

yielding the highest quality data compared to any single platform[12].

Second, the myriad of programs and pipelines to process LRs from

basecalling, to read correction, to assembly, to assembly correction, SR

recruitment and polishing, and more, continuously improve and

expand[3,4,14,17,18,21–24,26,27,29,31,33–38,40–43,45,49,52,54,55,70,71,76–89]. To reasonably

perform this study and analyze the results, only two assembly, one LR

correction, and one SR polishing programs were tested, thus introducing

technical variation by assembler but limiting it during LR correction and SR

polishing. These programs were chosen largely based on their observed

prevalence in literature and online resources, but we acknowledge that, in

particular, Racon and Pilon may be outcompeted by others that yield better

results[17,21,26,31,35,37,41,43,76–80]. The programs we used are also dataset agnostic,

and have relatively minimal computational requirements in contrast to tools that

are dataset-dependent or demand advanced/specific computational
capabilities , making them a relatively universal option. For several available LR
or hybrid assembly programs, Racon or Pilon have are already
implemented[19,26,29], indicating their value to the field despite the fact that others
or various combinations may yield more accurate assemblies. We further note
that each of these computational tools also come with a suite of settings to fine-
tune performance that were not explored. One setting of particular interest that
was not investigated here, nor could we find literature on, was read recruitment
stringency. Theoretically, this could greatly impact optimization of the iterative
processes, and thus accurate reconstructions, of microbial communities with
multiple similar strains, genomes with multicopy genes or repeat regions, or
other factors that have plagued metagenomic studies since their inception.

Third, contig pools were not separated for some analyses in order to
simplify the workflow. Specifically, in this workflow circular contigs that likely
constituted nearly complete genomes were not removed from the assemblies
prior LR correction or SR polishing. We speculate that this simplification could
result in mis-assemblies derived from reads, particularly SR, that should align
to different strains ambiguously aligning to conserved regions. However, SR
ambiguity decreased during iterative error fixing processes, possibly due to the
availability of multiple reference sequences to differentially recruit ambiguous
reads during SR polishing.

Fourth and lastly, we acknowledge that the biomass collected for DNA
isolation and sequencing with the different platforms were separated by several
months. Though largely stable after enrichment for over five years, the
communities still shift slowly, thought largely due to strains competing for the
same niches that will be examined in follow-up studies. These differences could
impact ideal integration the LR and SR data if strains substantially shifted in
abundance, leading to improper error-fixing events and could produce
inaccurate consensus genomes rather than a realistic strain representation.
672

Conclusions
Integrating LR and SR sequencing platforms for hybrid assembly of a microbial
community metagenome is challenging due to both biological and technical
complexities. While improvements to the quality of LR themselves may be able

to preclude the need for SRs, we expect that SRs will still be beneficial as
relatively inexpensive additional data for differential coverage binning, and
users may still desire to explore SR polishing to maximize their assembly
quality. Benchmarks combining these technologies have led to some
consensus in the methods for hybrid assembly, but direct replication of the
methods may not serve all applications. Here, we've demonstrated the that
simple, reference-free characteristics, specifically gene lengths and SR
recruitment, serve as dataset-dependent indicators of microbial community
metagenomic assembly quality, and even approximate advanced reference-
dependent characteristics. Additionally, the trajectories of reference-
independent assembly characteristics were consistent with benchmarks and
published workflows, but did also show that the number of iterations can have
substantial impact on downstream analyses. In summary, assembly
characteristics that are simple, reference-independent, and precursors to
downstream analyses, *e.g.*, gene counts and bps, or recruited short read
profiles, can help optimize hybrid assembly approach for microbial community
metagenomes. Rather than inform other users of the best approach for their
own hybrid metagenomic microbial community projects, we provide others with
assembly-level characteristics that are proxies quality to help optimize their own
approach. Finally, we offer practical recommendations to enable other users to
integrate LR and SR sequencing platforms for microbial community
metagenomes with fewer obstacles and guided by empirical data from their own
projects.
700
Additional practical recommendations
• Use the same pool or replicate extractions of DNA for both SR and LR,
which ideally minimizes community differences that might lead to mis-
assembly.
• Calculate read depth using bps rather than read counts when mixing LR
and SR profiles, otherwise LR depth will be vastly and unevenly
underestimated.
• If possible, use multiple assembly programs prior to LR correction and
SR polishing, and then consolidate the genome outputs (circular, MQ

automated bins) because different algorithms may have a substantial
impact on which community members assemble well[22].

- 712 LR correction should precede SR polishing to avoid the (re-)introduction
  713 of errors.

- 714 A mixture of different SR polishing tools may yield the best overall
  715 quality[17], likely due to differential abilities to polish specific errors.

- 716 These analyses are all relative, *i.e.*, dataset-dependent, and therefore
  717 should not be used to compare different systems.

- 718 While the ultimate goal might be genome recovery, we suggest
  719 monitoring gene fragmentation, read recruitment, or data collected
  720 during LR correction or SR polishing for empirically guiding optimization.
  - 721 Logged data from some programs can be information-rich and
    722 should be mined when possible. For example, Pilon records the
    723 changes it makes, which also followed the same patterns as the
    724 other analyses, *i.e.*, the most and largest changes occur in the
    725 first few iterations but plateaued on a log-scale by ~4 iterations
    726 (Fig. S11).

References

(1) Overholt, W. A.; Hölzer, M.; Geesink, P.; Diezel, C.; Marz, M.; Küsel, K. Inclusion of Oxford Nanopore Long Reads Improves All Microbial and Viral Metagenome-Assembled Genomes from a Complex Aquifer System. *Environmental Microbiology* **2020**, *22* (9), 4000–4013. https://doi.org/10.1111/1462-2920.15186.

(2) Singleton, C. M.; Petriglieri, F.; Kristensen, J. M.; Kirkegaard, R. H.; Michaelsen, T. Y.; Andersen, M. H.; Kondrotaite, Z.; Karst, S. M.; Dueholm, M. S.; Nielsen, P. H.; Albertsen, M. Connecting Structure to Function with the Recovery of over 1000 High-Quality Metagenome-Assembled Genomes from Activated Sludge Using Long-Read Sequencing. *Nat Commun* **2021**, *12* (1), 2009. https://doi.org/10.1038/s41467-021-22203-2.

(3) Bertrand, D.; Shaw, J.; Kalathiyappan, M.; Ng, A. H. Q.; Kumar, M. S.; Li, C.; Dvornicic, M.; Soldo, J. P.; Koh, J. Y.; Tong, C.; Ng, O. T.; Barkham, T.; Young, B.; Marimuthu, K.; Chng, K. R.; Sikic, M.; Nagarajan, N. Hybrid Metagenomic Assembly Enables High-Resolution Analysis of Resistance Determinants and Mobile Elements in Human Microbiomes. *Nat Biotechnol* **2019**, *37* (8), 937–944. https://doi.org/10.1038/s41587-019-0191-2.

(4) Liu, L.; Yang, Y.; Deng, Y.; Zhang, T. Nanopore Long-Read-Only Metagenomics Enables Complete and High-Quality Genome Reconstruction from Mock and Complex Metagenomes. *Microbiome* **2022**, *10* (1), 209. https://doi.org/10.1186/s40168-022-01415-8.

(5) Ye, L.; Dong, N.; Xiong, W.; Li, J.; Li, R.; Heng, H.; Chan, E. W. C.; Chen, S. High-Resolution Metagenomics of Human Gut Microbiota Generated by Nanopore and Illumina Hybrid Metagenome Assembly. *Frontiers in Microbiology* **2022**, *13*.

(6) Ravi, A.; Troncoso-Rey, P.; Ahn-Jarvis, J.; Corbin, K. R.; Harris, S.; Harris, H.; Aydin, A.; Kay, G. L.; Le Viet, T.; Gilroy, R.; Pallen, M. J.; Page, A. J.; O'Grady, J.; Warren, F. J. Hybrid Metagenome Assemblies Link Carbohydrate Structure with Function in the Human Gut Microbiome. *Commun Biol* **2022**, *5* (1), 1–13. https://doi.org/10.1038/s42003-022-03865-0.

(7) Gounot, J.-S.; Chia, M.; Bertrand, D.; Saw, W.-Y.; Ravikrishnan, A.; Low, A.; Ding, Y.; Ng, A. H. Q.; Tan, L. W. L.; Teo, Y.-Y.; Seedorf, H.; Nagarajan, N. Genome-Centric Analysis of Short and Long Read Metagenomes Reveals Uncharacterized Microbiome Diversity in Southeast Asians. *Nat Commun* **2022**, *13* (1), 6044. https://doi.org/10.1038/s41467-022-33782-z.

(8) Van Goethem, M. W.; Osborn, A. R.; Bowen, B. P.; Andeer, P. F.; Swenson, T. L.; Clum, A.; Riley, R.; He, G.; Koriabine, M.; Sandor, L.; Yan, M.; Daum, C. G.; Yoshinaga, Y.; Makhalanyane, T. P.; Garcia-Pichel, F.; Visel, A.; Pennacchio, L. A.; O'Malley, R. C.; Northen, T. R. Long-Read Metagenomics of Soil Communities Reveals Phylum-Specific Secondary Metabolite Dynamics. *Commun Biol* **2021**, *4* (1), 1–10. https://doi.org/10.1038/s42003-021-02809-4.

(9) Jin, H.; You, L.; Zhao, F.; Li, S.; Ma, T.; Kwok, L.-Y.; Xu, H.; Sun, Z. Hybrid, Ultra-Deep Metagenomic Sequencing Enables Genomic and Functional Characterization of Low-Abundance Species in the Human Gut Microbiome. *Gut Microbes* **2022**, *14* (1), 2021790. https://doi.org/10.1080/19490976.2021.2021790.

(10) Zhang, Z.-F.; Liu, L.-R.; Pan, Y.-P.; Pan, J.; Li, M. Long-Read Assembled Metagenomic Approaches Improve Our Understanding on Metabolic Potentials

of Microbial Community in Mangrove Sediments. *Microbiome* **2023**, *11* (1),
188. https://doi.org/10.1186/s40168-023-01630-x.

(11) Meslier, V.; Quinquis, B.; Da Silva, K.; Plaza Oñate, F.; Pons, N.; Roume, H.;
Podar, M.; Almeida, M. Benchmarking Second and Third-Generation
Sequencing Platforms for Microbial Metagenomics. *Sci Data* **2022**, *9* (1), 694.
https://doi.org/10.1038/s41597-022-01762-z.

(12) Sereika, M.; Kirkegaard, R. H.; Karst, S. M.; Michaelsen, T. Y.; Sørensen, E. A.;
Wollenberg, R. D.; Albertsen, M. Oxford Nanopore R10.4 Long-Read
Sequencing Enables the Generation of near-Finished Bacterial Genomes from
Pure Cultures and Metagenomes without Short-Read or Reference Polishing.
Nature Methods **2022**, *19* (7), 823–826. https://doi.org/10.1038/s41592-022-
01539-7.

(13) Stewart, R. D.; Auffret, M. D.; Warr, A.; Walker, A. W.; Roehe, R.; Watson, M.
Compendium of 4,941 Rumen Metagenome-Assembled Genomes for Rumen
Microbiome Biology and Enzyme Discovery. *Nat Biotechnol* **2019**, *37* (8), 953–
961. https://doi.org/10.1038/s41587-019-0202-3.

(14) Brown, C. L.; Keenum, I. M.; Dai, D.; Zhang, L.; Vikesland, P. J.; Pruden, A.
Critical Evaluation of Short, Long, and Hybrid Assembly for Contextual
Analysis of Antibiotic Resistance Genes in Complex Environmental
Metagenomes. *Scientific Reports* **2021**, *11* (1), 3753.
https://doi.org/10.1038/s41598-021-83081-8.

(15) Tao, Y.; Xun, F.; Zhao, C.; Mao, Z.; Li, B.; Xing, P.; Wu, Q. L. Improved
Assembly of Metagenome-Assembled Genomes and Viruses in Tibetan Saline
Lake Sediment by HiFi Metagenomic Sequencing. *Microbiology Spectrum* **2022**,
11 (1), e03328-22. https://doi.org/10.1128/spectrum.03328-22.

(16) Chen, L.-X.; Anantharaman, K.; Shaiber, A.; Eren, A. M.; Banfield, J. F.
Accurate and Complete Genomes from Metagenomes. *Genome Res.* **2020**, *30*
(3), 315–333. https://doi.org/10.1101/gr.258640.119.

(17) Wick, R. R.; Holt, K. E. Polypolish: Short-Read Polishing of Long-Read
Bacterial Genome Assemblies. *PLOS Computational Biology* **2022**, *18* (1),
e1009802. https://doi.org/10.1371/journal.pcbi.1009802.

(18) Zhang, Z.; Yang, C.; Veldsman, W. P.; Fang, X.; Zhang, L. Benchmarking
Genome Assembly Methods on Metagenomic Sequencing Data. *Briefings in
Bioinformatics* **2023**, *24* (2), bbad087. https://doi.org/10.1093/bib/bbad087.

(19) Wick, R. R.; Judd, L. M.; Gorrie, C. L.; Holt, K. E. Unicycler: Resolving
Bacterial Genome Assemblies from Short and Long Sequencing Reads. *PLOS
Computational Biology* **2017**, *13* (6), e1005595.
https://doi.org/10.1371/journal.pcbi.1005595.

(20) Weirather, J. L.; Cesare, M. de; Wang, Y.; Piazza, P.; Sebastiano, V.; Wang, X.-
J.; Buck, D.; Au, K. F. Comprehensive Comparison of Pacific Biosciences and
Oxford Nanopore Technologies and Their Applications to Transcriptome
Analysis. F1000Research June 19, 2017.
https://doi.org/10.12688/f1000research.10571.2.

(21) Zimin, A. V.; Salzberg, S. L. The Genome Polishing Tool POLCA Makes Fast
and Accurate Corrections in Genome Assemblies. *PLOS Computational Biology*
2020, *16* (6), e1007981. https://doi.org/10.1371/journal.pcbi.1007981.

(22) Wick, R. R.; Judd, L. M.; Cerdeira, L. T.; Hawkey, J.; Méric, G.; Vezina, B.;
Wyres, K. L.; Holt, K. E. Trycycler: Consensus Long-Read Assemblies for
Bacterial Genomes. *Genome Biology* **2021**, *22* (1), 266.
https://doi.org/10.1186/s13059-021-02483-z.

(23) Wick, R. R.; Holt, K. E. Benchmarking of Long-Read Assemblers for
828   Prokaryote Whole Genome Sequencing. F1000Research February 1, 2021.
829   https://doi.org/10.12688/f1000research.21782.4.
830   (24) Wick, R. R.; Judd, L. M.; Holt, K. E. Performance of Neural Network
831   Basecalling Tools for Oxford Nanopore Sequencing. *Genome Biology* **2019**, *20*
832   (1), 129. https://doi.org/10.1186/s13059-019-1727-y.
833   (25) Watson, M.; Warr, A. Errors in Long-Read Assemblies Can Critically Affect
834   Protein Prediction. *Nat Biotechnol* **2019**, *37* (2), 124–126.
835   https://doi.org/10.1038/s41587-018-0004-z.
836   (26) Huang, N.; Nie, F.; Ni, P.; Luo, F.; Gao, X.; Wang, J. NeuralPolish: A Novel
837   Nanopore Polishing Method Based on Alignment Matrix Construction and
838   Orthogonal Bi-GRU Networks. *Bioinformatics* **2021**, *37* (19), 3120–3127.
839   https://doi.org/10.1093/bioinformatics/btab354.
840   (27) Vaser, R.; Sovic, I.; Nagarajan, N.; Sikic, M. Fast and Accurate de Novo
841   Genome Assembly from Long Uncorrected Reads. *Genome Res.* **2017**,
842   gr.214270.116. https://doi.org/10.1101/gr.214270.116.
843   (28) Chen, Z.; Erickson, D. L.; Meng, J. Polishing the Oxford Nanopore Long-Read
844   Assemblies of Bacterial Pathogens with Illumina Short Reads to Improve
845   Genomic Analyses. *Genomics* **2021**, *113* (3), 1366–1377.
846   https://doi.org/10.1016/j.ygeno.2021.03.018.
847   (29) Damme, R. V.; Hölzer, M.; Viehweger, A.; Müller, B.; Bongcam-Rudloff, E.;
848   Brandt, C. Metagenomics Workflow for Hybrid Assembly, Differential
849   Coverage Binning, Metatranscriptomics and Pathway Analysis (MUFFIN).
850   *PLOS Computational Biology* **2021**, *17* (2), e1008716.
851   https://doi.org/10.1371/journal.pcbi.1008716.
852   (30) Belser, C.; Istace, B.; Denis, E.; Dubarry, M.; Baurens, F.-C.; Falentin, C.;
853   Genete, M.; Berrabah, W.; Chèvre, A.-M.; Delourme, R.; Deniot, G.; Denoeud,
854   F.; Duffé, P.; Engelen, S.; Lemainque, A.; Manzanares-Dauleux, M.; Martin, G.;
855   Morice, J.; Noel, B.; Vekemans, X.; D'Hont, A.; Rousseau-Gueutin, M.; Barbe,
856   V.; Cruaud, C.; Wincker, P.; Aury, J.-M. Chromosome-Scale Assemblies of
857   Plant Genomes Using Nanopore Long Reads and Optical Maps. *Nature Plants*
858   **2018**, *4* (11), 879–887. https://doi.org/10.1038/s41477-018-0289-4.
859   (31) Hu, J.; Fan, J.; Sun, Z.; Liu, S. NextPolish: A Fast and Efficient Genome
860   Polishing Tool for Long-Read Assembly. *Bioinformatics* **2020**, *36* (7), 2253–
861   2255. https://doi.org/10.1093/bioinformatics/btz891.
862   (32) Chen, Z.; Erickson, D. L.; Meng, J. Benchmarking Hybrid Assembly
863   Approaches for Genomic Analyses of Bacterial Pathogens Using Illumina and
864   Oxford Nanopore Sequencing. *BMC Genomics* **2020**, *21* (1), 631.
865   https://doi.org/10.1186/s12864-020-07041-8.
866   (33) Kolmogorov, M.; Bickhart, D. M.; Behsaz, B.; Gurevich, A.; Rayko, M.; Shin,
867   S. B.; Kuhn, K.; Yuan, J.; Polevikov, E.; Smith, T. P. L.; Pevzner, P. A.
868   metaFlye: Scalable Long-Read Metagenome Assembly Using Repeat Graphs.
869   *Nat Methods* **2020**, *17* (11), 1103–1110. https://doi.org/10.1038/s41592-020-
870   00971-x.
871   (34) Koren, S.; Walenz, B. P.; Berlin, K.; Miller, J. R.; Bergman, N. H.; Phillippy, A.
872   M. Canu: Scalable and Accurate Long-Read Assembly via Adaptive k-Mer
873   Weighting and Repeat Separation. *Genome Res.* **2017**, *27* (5), 722–736.
874   https://doi.org/10.1101/gr.215087.116.
875   (35) Firtina, C.; Kim, J. S.; Alser, M.; Senol Cali, D.; Cicek, A. E.; Alkan, C.; Mutlu,
876   O. Apollo: A Sequencing-Technology-Independent, Scalable and Accurate

Assembly Polishing Algorithm. *Bioinformatics* **2020**, *36* (12), 3669–3679.
https://doi.org/10.1093/bioinformatics/btaa179.

(36) Zeng, J.; Cai, H.; Peng, H.; Wang, H.; Zhang, Y.; Akutsu, T. Causalcall:
Nanopore Basecalling Using a Temporal Convolutional Network. *Frontiers in*
Genetics **2020**, *10*.

(37) Lee, J. Y.; Kong, M.; Oh, J.; Lim, J.; Chung, S. H.; Kim, J.-M.; Kim, J.-S.; Kim,
K.-H.; Yoo, J.-C.; Kwak, W. Comparative Evaluation of Nanopore Polishing
Tools for Microbial Genome Assembly and Polishing Strategies for Downstream
Analysis. *Sci Rep* **2021**, *11* (1), 20740. https://doi.org/10.1038/s41598-021-
00178-w.

(38) Li, H. Minimap and Miniasm: Fast Mapping and de Novo Assembly for Noisy
Long Sequences. *Bioinformatics* **2016**, *32* (14), 2103–2110.
https://doi.org/10.1093/bioinformatics/btw152.

(39) De Maio, N.; Shaw, L. P.; Hubbard, A.; George, S.; Sanderson, N. D.; Swann, J.;
Wick, R.; AbuOun, M.; Stubberfield, E.; Hoosdally, S. J.; Crook, D. W.; Peto, T.
E. A.; Sheppard, A. E.; Bailey, M. J.; Read, D. S.; Anjum, M. F.; Walker, A. S.;
Stoesser, N.; consortium, on behalf of the R. Comparison of Long-Read
Sequencing Technologies in the Hybrid Assembly of Complex Bacterial
Genomes. *Microbial Genomics,* 2019, *5*.
https://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.00
0294.

(40) Antipov, D.; Korobeynikov, A.; McLean, J. S.; Pevzner, P. A. hybridSPAdes:
An Algorithm for Hybrid Assembly of Short and Long Reads. *Bioinformatics*
2016, *32* (7), 1009–1015. https://doi.org/10.1093/bioinformatics/btv688.

(41) Hu, K.; Huang, N.; Zou, Y.; Liao, X.; Wang, J. MultiNanopolish: Refined
Grouping Method for Reducing Redundant Calculations in Nanopolish.
Bioinformatics **2021**, *37* (17), 2757–2760.
https://doi.org/10.1093/bioinformatics/btab078.

(42) Huang, N.; Nie, F.; Ni, P.; Luo, F.; Wang, J. SACall: A Neural Network
Basecaller for Oxford Nanopore Sequencing Data Based on Self-Attention
Mechanism. *IEEE/ACM Transactions on Computational Biology and*
Bioinformatics **2022**, *19* (1), 614–623.
https://doi.org/10.1109/TCBB.2020.3039244.

(43) Warren, R. L.; Coombe, L.; Mohamadi, H.; Zhang, J.; Jaquish, B.; Isabel, N.;
Jones, S. J. M.; Bousquet, J.; Bohlmann, J.; Birol, I. ntEdit: Scalable Genome
Sequence Polishing. *Bioinformatics* **2019**, *35* (21), 4430–4432.
https://doi.org/10.1093/bioinformatics/btz400.

(44) Dida, F.; Yi, G. Empirical Evaluation of Methods for de Novo Genome
Assembly. *PeerJ Comput. Sci.* **2021**, *7*, e636. https://doi.org/10.7717/peerj-
cs.636.

(45) Dohm, J. C.; Peters, P.; Stralis-Pavese, N.; Himmelbauer, H. Benchmarking of
Long-Read Correction Methods. *NAR Genomics and Bioinformatics* **2020**, *2* (2),
lqaa037. https://doi.org/10.1093/nargab/lqaa037.

(46) Albertsen, M.; Karst, S. M.; Ziegler, A. S.; Kirkegaard, R. H.; Nielsen, P. H.
Back to Basics – The Influence of DNA Extraction and Primer Choice on
Phylogenetic Analysis of Activated Sludge Communities. *PLOS ONE* **2015**, *10*
(7), e0132783. https://doi.org/10.1371/journal.pone.0132783.

(47) Weber, L.; DeForce, E.; Apprill, A. Optimization of DNA Extraction for
Advancing Coral Microbiota Investigations. *Microbiome* **2017**, *5* (1), 18.
https://doi.org/10.1186/s40168-017-0229-y.

(48) Martin-Laurent, F.; Philippot, L.; Hallet, S.; Chaussod, R.; Germon, J. C.;
Soulas, G.; Catroux, G. DNA Extraction from Soils: Old Bias for New Microbial
Diversity Analysis Methods. *Applied and Environmental Microbiology* **2001**, *67*
(5), 2354–2359. https://doi.org/10.1128/AEM.67.5.2354-2359.2001.
(49) *Nanopore Community*. Oxford Nanopore Technologies.
https://nanoporetech.com/community (accessed 2023-04-05).
(50) *BBMap*. SourceForge. https://sourceforge.net/projects/bbmap/ (accessed 2023-
02-14).
(51) Wick, R. Porechop, 2023. https://github.com/rrwick/Porechop (accessed 2023-
02-14).
(52) Nurk, S.; Meleshko, D.; Korobeynikov, A.; Pevzner, P. A. metaSPAdes: A New
Versatile Metagenomic Assembler. *Genome Res* **2017**, *27* (5), 824–834.
https://doi.org/10.1101/gr.213959.116.
(53) Li, H. Minimap2: Pairwise Alignment for Nucleotide Sequences. *Bioinformatics*
2018, *34* (18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191.
(54) Walker, B. J.; Abeel, T.; Shea, T.; Priest, M.; Abouelliel, A.; Sakthikumar, S.;
Cuomo, C. A.; Zeng, Q.; Wortman, J.; Young, S. K.; Earl, A. M. Pilon: An
Integrated Tool for Comprehensive Microbial Variant Detection and Genome
Assembly Improvement. *PLOS ONE* **2014**, *9* (11), e112963.
https://doi.org/10.1371/journal.pone.0112963.
(55) Zhang, Y.; Akdemir, A.; Tremmel, G.; Imoto, S.; Miyano, S.; Shibuya, T.;
Yamaguchi, R. Nanopore Basecalling from a Perspective of Instance
Segmentation. *BMC Bioinformatics* **2020**, *21* (3), 136.
https://doi.org/10.1186/s12859-020-3459-0.
(56) Mikheenko, A.; Saveliev, V.; Gurevich, A. MetaQUAST: Evaluation of
Metagenome Assemblies. *Bioinformatics* **2016**, *32* (7), 1088–1090.
https://doi.org/10.1093/bioinformatics/btv697.
(57) Hyatt, D.; Chen, G.-L.; LoCascio, P. F.; Land, M. L.; Larimer, F. W.; Hauser, L.
J. Prodigal: Prokaryotic Gene Recognition and Translation Initiation Site
Identification. *BMC Bioinformatics* **2010**, *11* (1), 119.
https://doi.org/10.1186/1471-2105-11-119.
(58) Manni, M.; Berkeley, M. R.; Seppey, M.; Simão, F. A.; Zdobnov, E. M. BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral
Genomes. *Molecular Biology and Evolution* **2021**, *38* (10), 4647–4654.
https://doi.org/10.1093/molbev/msab199.
(59) Parks, D. H.; Imelfort, M.; Skennerton, C. T.; Hugenholtz, P.; Tyson, G. W.
CheckM: Assessing the Quality of Microbial Genomes Recovered from Isolates,
Single Cells, and Metagenomes. *Genome Res.* **2015**, *25* (7), 1043–1055.
https://doi.org/10.1101/gr.186072.114.
(60) Buchfink, B.; Xie, C.; Huson, D. H. Fast and Sensitive Protein Alignment Using
DIAMOND. *Nat Methods* **2015**, *12* (1), 59–60.
https://doi.org/10.1038/nmeth.3176.
(61) Suzek, B. E.; Wang, Y.; Huang, H.; McGarvey, P. B.; Wu, C. H.; the UniProt
Consortium. UniRef Clusters: A Comprehensive and Scalable Alternative for
Improving Sequence Similarity Searches. *Bioinformatics* **2015**, *31* (6), 926–932.
https://doi.org/10.1093/bioinformatics/btu739.
(62) Chaumeil, P.-A.; Mussig, A. J.; Hugenholtz, P.; Parks, D. H. GTDB-Tk: A
Toolkit to Classify Genomes with the Genome Taxonomy Database.

Bioinformatics **2020**, *36* (6), 1925–1927.
https://doi.org/10.1093/bioinformatics/btz848.

(63) SorenKarst. Mmlong, 2022. https://github.com/SorenKarst/mmlong (accessed
2023-02-14).

(64) Kang, D. D.; Li, F.; Kirton, E.; Thomas, A.; Egan, R.; An, H.; Wang, Z.
MetaBAT 2: An Adaptive Binning Algorithm for Robust and Efficient Genome
Reconstruction from Metagenome Assemblies. *PeerJ* **2019**, *7*, e7359.
https://doi.org/10.7717/peerj.7359.

(65) Shaffer, M.; Borton, M. A.; McGivern, B. B.; Zayed, A. A.; La Rosa, S. L.;
Solden, L. M.; Liu, P.; Narrowe, A. B.; Rodríguez-Ramos, J.; Bolduc, B.;
Gazitúa, M. C.; Daly, R. A.; Smith, G. J.; Vik, D. R.; Pope, P. B.; Sullivan, M.
B.; Roux, S.; Wrighton, K. C. DRAM for Distilling Microbial Metabolism to
Automate the Curation of Microbiome Function. *Nucleic Acids Research* **2020**,
48 (16), 8883–8900. https://doi.org/10.1093/nar/gkaa621.

(66) Eddy, S. R. Accelerated Profile HMM Searches. *PLOS Computational Biology*
2011, *7* (10), e1002195. https://doi.org/10.1371/journal.pcbi.1002195.

(67) Oksanen, J.; Simpson, G. L.; Blanchet, F. G.; Kindt, R.; Legendre, P.; Minchin,
P. R.; O'Hara, R. B.; Solymos, P.; Stevens, M. H. H.; Szoecs, E.; Wagner, H.;
Barbour, M.; Bedward, M.; Bolker, B.; Borcard, D.; Carvalho, G.; Chirico, M.;
Caceres, M. D.; Durand, S.; Evangelista, H. B. A.; FitzJohn, R.; Friendly, M.;
Furneaux, B.; Hannigan, G.; Hill, M. O.; Lahti, L.; McGlinn, D.; Ouellette, M.-
H.; Cunha, E. R.; Smith, T.; Stier, A.; Braak, C. J. F. T.; Weedon, J. *Vegan:*
Community Ecology Package; 2022.

(68) R Core Team. *R: A Language and Environment for Statistical Computing*; R
Foundation for Statistical Computing: Vienna, Austria, 2021.

(69) Wickham, H.; Averick, M.; Bryan, J.; Chang, W.; McGowan, L. D.; François,
R.; Grolemund, G.; Hayes, A.; Henry, L.; Hester, J.; Kuhn, M.; Pedersen, T. L.;
Miller, E.; Bache, S. M.; Müller, K.; Ooms, J.; Robinson, D.; Seidel, D. P.;
Spinu, V.; Takahashi, K.; Vaughan, D.; Wilke, C.; Woo, K.; Yutani, H.
Welcome to the Tidyverse. *Journal of Open Source Software* **2019**, *4* (43), 1686.
https://doi.org/10.21105/joss.01686.

(70) Amarasinghe, S. L.; Su, S.; Dong, X.; Zappia, L.; Ritchie, M. E.; Gouil, Q.
Opportunities and Challenges in Long-Read Sequencing Data Analysis. *Genome*
Biology **2020**, *21* (1), 30. https://doi.org/10.1186/s13059-020-1935-5.

(71) Krakau, S.; Straub, D.; Gourlé, H.; Gabernet, G.; Nahnsen, S. Nf-Core/Mag: A
Best-Practice Pipeline for Metagenome Hybrid Assembly and Binning. *NAR*
Genomics and Bioinformatics **2022**, *4* (1), lqac007.
https://doi.org/10.1093/nargab/lqac007.

(72) *PacBio Revio | Long-read sequencing at scale*. PacBio.
https://www.pacb.com/revio/ (accessed 2023-04-04).

(73) *Continuous development and improvement*. Oxford Nanopore Technologies.
https://nanoporetech.com/about-us/continuous-development-and-improvement
(accessed 2023-04-04).

(74) Zhao, W.; Zeng, W.; Pang, B.; Luo, M.; Peng, Y.; Xu, J.; Kan, B.; Li, Z.; Lu, X.
Oxford Nanopore Long-Read Sequencing Enables the Generation of Complete
Bacterial and Plasmid Genomes without Short-Read Sequencing. *Frontiers in*
Microbiology **2023**, *14*.

(75) Zhang, H.; Jain, C.; Aluru, S. A Comprehensive Evaluation of Long Read Error
Correction Methods. *BMC Genomics* **2020**, *21* (6), 889.
https://doi.org/10.1186/s12864-020-07227-0.

(76) Kundu, R.; Casey, J.; Sung, W.-K. HyPo: Super Fast & Accurate Polisher for Long Read Genome Assemblies. bioRxiv December 20, 2019, p 2019.12.19.882506. https://doi.org/10.1101/2019.12.19.882506.

(77) Shafin, K.; Pesout, T.; Chang, P.-C.; Nattestad, M.; Kolesnikov, A.; Goel, S.; Baid, G.; Kolmogorov, M.; Eizenga, J. M.; Miga, K. H.; Carnevali, P.; Jain, M.; Carroll, A.; Paten, B. Haplotype-Aware Variant Calling with PEPPER-Margin-DeepVariant Enables High Accuracy in Nanopore Long-Reads. *Nat Methods* **2021**, *18* (11), 1322–1332. https://doi.org/10.1038/s41592-021-01299-w.

(78) Medaka, 2023. https://github.com/nanoporetech/medaka (accessed 2023-11-21).

(79) Huang, Y.-T.; Liu, P.-Y.; Shih, P.-W. Homopolish: A Method for the Removal of Systematic Errors in Nanopore Sequencing by Homologous Polishing. *Genome Biology* **2021**, *22* (1), 95. https://doi.org/10.1186/s13059-021-02282-6.

(80) Ruan, J.; Li, H. Fast and Accurate Long-Read Assembly with Wtdbg2. *Nat Methods* **2020**, *17* (2), 155–158. https://doi.org/10.1038/s41592-019-0669-3.

(81) Shafin, K.; Pesout, T.; Lorig-Roach, R.; Haukness, M.; Olsen, H. E.; Bosworth, C.; Armstrong, J.; Tigyi, K.; Maurer, N.; Koren, S.; Sedlazeck, F. J.; Marschall, T.; Mayes, S.; Costa, V.; Zook, J. M.; Liu, K. J.; Kilburn, D.; Sorensen, M.; Munson, K. M.; Vollger, M. R.; Monlong, J.; Garrison, E.; Eichler, E. E.; Salama, S.; Haussler, D.; Green, R. E.; Akeson, M.; Phillippy, A.; Miga, K. H.; Carnevali, P.; Jain, M.; Paten, B. Nanopore Sequencing and the Shasta Toolkit Enable Efficient de Novo Assembly of Eleven Human Genomes. *Nat Biotechnol* **2020**, *38* (9), 1044–1053. https://doi.org/10.1038/s41587-020-0503-6.

(82) Vaser, R.; Šikić, M. Time- and Memory-Efficient Genome Assembly with Raven. *Nat Comput Sci* **2021**, *1* (5), 332–336. https://doi.org/10.1038/s43588-021-00073-4.

(83) Chen, Y.; Nie, F.; Xie, S.-Q.; Zheng, Y.-F.; Dai, Q.; Bray, T.; Wang, Y.-X.; Xing, J.-F.; Huang, Z.-J.; Wang, D.-P.; He, L.-J.; Luo, F.; Wang, J.-X.; Liu, Y.-Z.; Xiao, C.-L. Efficient Assembly of Nanopore Reads via Highly Accurate and Intact Error Correction. *Nat Commun* **2021**, *12* (1), 60. https://doi.org/10.1038/s41467-020-20236-7.

(84) Hu, J.; Wang, Z.; Sun, Z.; Hu, B.; Ayoola, A. O.; Liang, F.; Li, J.; Sandoval, J. R.; Cooper, D. N.; Ye, K.; Ruan, J.; Xiao, C.-L.; Wang, D.-P.; Wu, D.-D.; Wang, S. An Efficient Error Correction and Accurate Assembly Tool for Noisy Long Reads. bioRxiv March 12, 2023, p 2023.03.09.531669. https://doi.org/10.1101/2023.03.09.531669.

(85) Pagès-Gallego, M.; de Ridder, J. Comprehensive Benchmark and Architectural Analysis of Deep Learning Models for Nanopore Sequencing Basecalling. *Genome Biology* **2023**, *24* (1), 71. https://doi.org/10.1186/s13059-023-02903-2.

(86) Konishi, H.; Yamaguchi, R.; Yamaguchi, K.; Furukawa, Y.; Imoto, S. Halcyon: An Accurate Basecaller Exploiting an Encoder–Decoder Model with Monotonic Attention. *Bioinformatics* **2021**, *37* (9), 1211–1217. https://doi.org/10.1093/bioinformatics/btaa953.

(87) Xu, Z.; Mai, Y.; Liu, D.; He, W.; Lin, X.; Xu, C.; Zhang, L.; Meng, X.; Mafofo, J.; Zaher, W. A.; Koshy, A.; Li, Y.; Qiao, N. Fast-Bonito: A Faster Deep Learning Based Basecaller for Nanopore Sequencing. *Artificial Intelligence in the Life Sciences* **2021**, *1*, 100011. https://doi.org/10.1016/j.ailsci.2021.100011.

(88) Lv, X.; Chen, Z.; Lu, Y.; Yang, Y. An End-to-End Oxford Nanopore Basecaller Using Convolution-Augmented Transformer. bioRxiv November 10, 2020, p 2020.11.09.374165. https://doi.org/10.1101/2020.11.09.374165.

(89) Miculinić, N.; Ratković, M.; Šikić, M. MinCall - MinION End2end
Convolutional Deep Learning Basecaller. arXiv April 22, 2019.
https://doi.org/10.48550/arXiv.1904.10337.
(90) Schulz, F.; Lagkouvardos, I.; Wascher, F.; Aistleitner, K.; Kostanjšek, R.; Horn,
M. Life in an Unusual Intracellular Niche: A Bacterial Symbiont Infecting the
Nucleus of Amoebae. *ISME J* **2014**, *8* (8), 1634–1644.
https://doi.org/10.1038/ismej.2014.5.
(91) Birtles, R. J.; Rowbotham, T. J.; Michel, R.; Pitcher, D. G.; Lascola, B.;
Alexiou-Daniel, S.; Raoult, D. "Candidatus Odyssella Thessalonicensis" Gen.
Nov., Sp. Nov., an Obligate Intracellular Parasite of Acanthamoeba Species.
International Journal of Systematic and Evolutionary Microbiology **2000**, *50*
(1), 63–72. https://doi.org/10.1099/00207713-50-1-63.
(92) Horn, M.; Wagner, M.; Müller, K.-D.; Schmid, E. N.; Fritsche, T. R.; Schleifer,
K.-H.; Michel, R. Neochlamydia Hartmannellae Gen. Nov., Sp. Nov.
(Parachlamydiaceae), an Endoparasite of the Amoeba Hartmannella
Vermiformis. *Microbiology (Reading)* **2000**, *146 (Pt 5)*, 1231–1239.
https://doi.org/10.1099/00221287-146-5-1231.
(93) Zablocki, O.; Michelsen, M.; Burris, M.; Solonenko, N.; Warwick-Dugdale, J.;
Ghosh, R.; Pett-Ridge, J.; Sullivan, M. B.; Temperton, B. VirION2: A Short-
and Long-Read Sequencing and Informatics Workflow to Study the Genomic
Diversity of Viruses in Nature. *PeerJ* **2021**, *9*, e11088.
https://doi.org/10.7717/peerj.11088.
(94) Sieber, C. M. K.; Paul, B. G.; Castelle, C. J.; Hu, P.; Tringe, S. G.; Valentine, D.
L.; Andersen, G. L.; Banfield, J. F. Unusual Metabolism and Hypervariation in
the Genome of a Gracilibacterium (BD1-5) from an Oil-Degrading Community.
mBio **2019**, *10* (6), e02128-19. https://doi.org/10.1128/mBio.02128-19.
1101

Figure 1. Schematic overview of assembly, correction, and polishing.
(A) Assembly, correction, and polishing experimental design. Broken lines represent read recruitment steps for either correction or polishing. Acronyms: SR, short read; LR, long read; Def., default settings; Met., metagenome-optimized settings; Sub., sub-sampled LRs with metagenome-optimized settings. (B) Simplified diagram of analyses performed and the tools used to perform them for each assembly and after each iteration of correction or polishing.

Figure 2. Beta diversity estimated from RNA Polymerase subunit B (*rpoB*) gene profiles for each bioreactor and Long Read (LR) assembly throughout the iterative LR correction and Short Read (SR) polishing processes.

The two bioreactors are separated over vertical panels, the two LR assemblers are separated over the horizontal panels. Points are the mean-centered species scores for all rpoB genes per assembly, with gray lines showing the standard errors of the mean. Each point is colored by the SR polishing iteration, with colored lines connecting points to the preceding LR correction and subsequent SR polishing iteration. White text over points indicates the LR correction iteration prior to any SR polishing. Inset shows the overlapping points of the second LR correction stage as the communities converge due to SR polishing for the Nitrogen-Limited Bioreactor's Flye assembly.

Figure 3. Stair plots of automated medium-quality or better (MQ) bin recovery for each bioreactor and Long Read (LR) assembly throughout the iterative LR correction and Short Read (SR) polishing processes.

The two bioreactors are separated over vertical panels, the two LR assemblers are separated over the horizontal panels. The LR correction and SR polishing iterations are spread across the x-axis so that the ten SR polishing steps are immediately to the right of the respective preceding LR correction step. Each colored line represents the proportion of the maximum observed value per combination of bioreactor and assembler of one of the following characteristics MQ automated bins: the number of MQ bins recovered ("count"), the mean completeness score of the bins estimated by CheckM ("mean completeness"), the number of contigs in MQ bins ("contigs"), and the quantity of bps in MQ bins ("bps"). Matching color diamonds indicate the LR correction and/or SR polishing stage with the maximum observed value for each characteristic.

Figure 4. Linear correlation between calculated bps in genes and the number of these genes estimated to be similar length to reference database entries for each bioreactor and Long Read (LR) assembly throughout the iterative LR correction and Short Read (SR) polishing processes.

The two bioreactors are separated over vertical panels, and the two LR assemblers are separated over the horizontal panels. Each point is colored by the SR polishing iteration, with lines connecting the points from the same LR correction iteration, all of which are partially transparent. The light blue solid lines and shaded regions are the linear regressions for the displayed data and their 95% confidence intervals. Slopes, correlation coefficients (adjusted R^2) and p-values are displayed in upper-left corner of each panel.

Figure 5. Polar coordinate plot of Short Read (SR) recruitment for each bioreactor and Long Read (LR) assembly throughout the iterative LR correction and SR polishing processes.

The two bioreactors are separated over vertical panels, and the two LR assemblers are separated over the horizontal panels. Each panel shows a polar coordinate plot of the LR correction or SR polishing iteration in a circle with the bps in SR recruited per contig per Mbp sequence length as the distance from the center. The innermost ten points (due to lower values) indicate the amount of bps in SR recruited after up to ten LR correction iterations without SR polishing. The colored lines connecting points of the same color link assemblies with the same number of LR correction steps preceding SR polishing. The size of the points indicates the fraction of SR that were ambiguously recruited, i.e., SR aligning to 2 or more positions.