The αC-β4 loop controls the allosteric cooperativity between nucleotide and substrate in the catalytic subunit of protein kinase A

Cristina Olivieri[1,†,‡], Yingjie Wang[1,2,†,§], Caitlin Walker[1], Manu V. Subrahmanian[1], Kim N. Ha[3],

David A. Bernlohr[1], Jiali Gao[2], Carlo Camilloni[4,◊], Michele Vendruscolo[4], Susan S. Taylor[5,6], and

Gianluigi Veglia[1,2]*

[1]Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, MN 55455, USA

[2]Department of Chemistry and Supercomputing Institute, University of Minnesota, MN 55455, USA

[3]Departmenf of Chemistry and Biochemistry, St. Catherine University, MN 55105, USA

[4]Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, UK.

[5]Department of Pharmacology, University of California at San Diego, CA 92093, USA.

[6]Department of Chemistry and Biochemistry, University of California at San Diego, CA 92093, USA.

[†]These authors contributed equally to this work.

Present address:

[‡] Department of Chemistry, University of Milan, 20133, Milan, Italy

[§] Institute of Systems and Physical Biology, Shenzhen Bay Laboratory, Shenzhen, 518055, China

[◊] Department of Bioscience, University of Milano, 20133, Milano, Italy

*Corresponding Author

Gianluigi Veglia
Department of Chemistry and Department of Biochemistry, Molecular Biology, and Biophysics,
321 Church Street SE
Minneapolis, MN 55455
Telephone: (612) 625-0758
Fax: (612) 625-5780
e-mail: vegli001@umn.edu

ABSTRACT

Allosteric cooperativity between ATP and substrates is a prominent characteristic of the cAMP-dependent catalytic subunit of protein kinase A (PKA). Not only this long-range synergistic action is involved in substrate recognition and fidelity, but it is also likely to regulate PKA association with regulatory subunits and other binding partners. To date, a complete understanding of the molecular determinants for this intramolecular mechanism is still lacking.

Here, we integrated NMR-restrained molecular dynamics simulations and a Markov State Model to characterize the free energy landscape and conformational transitions of the catalytic subunit of protein kinase A (PKA-C). We found that the apoenzyme populates a broad free energy basin featuring a conformational ensemble of the active state of PKA-C (ground state) and other basins with lower populations (excited states). The first excited state corresponds to a previously characterized inactive state of PKA-C with the αC helix swinging outward. The second excited state displays a disrupted hydrophobic packing around the regulatory (R) spine, with a flipped configuration of the F100 and F102 residues at the αC-β4 loop. To experimentally validate the second excited state, we mutated F100 into alanine (F100A) and used NMR spectroscopy to characterize the structural response of the kinase to ATP and substrate binding. While the catalytic efficiency of PKA-C^{F100A} with a canonical peptide substrate remains unaltered, this mutation rearranges the αC-β4 loop conformation, interrupting the structural coupling of the two lobes and abolishing the allosteric binding cooperativity of the enzyme. The highly conserved αC-β4 loop emerges as a pivotal element able to control the synergistic binding between nucleotide and substrate. These results may explain how mutations or insertions near or within this motif affect the function and drug sensitivity in other homologous kinases.

Keywords:
Protein Kinases, cAMP-dependent protein kinase A, αC-β4 loop, allosteric mutations, binding co-operativity.

INTRODUCTION

Eukaryotic protein kinases (EPKs) are plastic enzymes of paramount importance in signaling processes, catalyzing phosphoryl transfer reactions, or acting as scaffolds for other enzymes and/or binding partners [1,2]. Of all kinases, the catalytic subunit of PKA (PKA-C) was the first to be structurally characterized by X-ray crystallography.[3,4] In its inhibited state, PKA-C assembles into a heterotetrametric holoenzyme comprising two catalytic (C) and two regulatory (R) subunits.[5] The canonical activation mechanism of PKA involves the binding of two cAMP molecules that disassemble the holoenzyme, unleashing active PKA-C monomers, which target signaling part-ners.[6] In 2017, however, Scott and coworkers proposed an alternative activation mechanism in which the holoenzyme does not disassemble under physiological conditions; rather, it is clustered in *signaling islands*, localized near the substrates by A-kinase anchoring proteins (AKAPs), facil-itating targeted signaling interactions.[7] To date, the activation mechanism of PKA is still under active investigation since PKA-C appears to be fully liberated under pathological conditions.[8-10]

X-ray crystallography studies revealed that PKA-C is a bilobal enzyme, with a dynamic N-terminal lobe (N-lobe) comprising four β-sheets and an αC helix, while the C-terminal lobe (C-lobe) is more rigid and mostly composed of α-helices (**Figure 1A**).[3,4,11] The N-lobe harbors the nucleotide-binding site, whereas the substrate binding cleft lays at the interface between the N- and C-lobe. The three-dimensional structure of PKA-C features a highly conserved hydrophobic core decorated with catalytically important motifs, *i.e.*, the Gly-rich loop, DFG-loop, activation loop, and magnesium and peptide positioning loops.[12] In the catalytically active state, these motifs are all poised for phosphoryl transfer,[13-15] a condition necessary but not sufficient to define an active kinase. More recent studies revealed a critical role of the enzyme's hydrophobic core, which is crossed by a catalytic (C) spine and a regulatory (R) spine surrounded by shell residues (**Figure 1B**).[16,17] The C spine comprises an array of hydrophobic residues and is assembled upon binding

ATP, whereas the R spine is engaged when the activation loop is phosphorylated at Thr197,

which positions the αC-helix in an active configuration.[18]

A distinct property of PKA-C is the binding cooperativity between ATP and substrate.[19] In the

catalytic cycle, the kinase binds ATP and unphosphorylated substrates with positive binding co-

operativity, whereas a negative binding cooperativity between ADP and phosphorylated substrate

characterizes the exit complex.[20] The biological importance of these phenomena has been em-

phasized by our recent studies on disease-driven mutations,[21-23] which all feature disrupted co-

operativity between nucleotides and protein kinase inhibitor (PKI) or canonical substrates.[21-23]

Since the recognition sequence of substrates is highly homologous to that of the R subunits,[5] a

loss of cooperativity may affect not only substrate binding fidelity but also regulatory processes.

Thus far, the molecular determinants for the binding cooperativity between nucleotide and sub-

strate and its role in PKA signalosome remain elusive.

Figure 1. Structural and catalytic motifs of PKA-C. (**A**) Backbone representation of the ter-
nary complex of PKA-C bound to ATP and the PKI$_{5-24}$ peptide (not depicted), PDB Code 4WB5.
Highlighted are key motifs including the αA, αC, and αE helices, C-terminal tail, activation loop,
peptide-positioning loop, Gly-rich loop, and the DFG and APE motifs. (**B**) The hydrophobic core
of PKA-C features the regulatory spine (R-spine, gold), catalytic spine (C-spine, blue), shell resi-
dues (cyan), and the αC-β4 loop (hot pink), which locks the αE helix and couples the two lobes
of PKA-C.

Aiming to define the allosteric mechanism for PKA-C cooperativity, we combined NMR-restrained

replica-averaged metadynamics (RAM)[24,25] and Markov State Model (MSM)[26] and charted the

conformational landscape and dynamic transitions of the main forms of PKA-C. We found that the

apo kinase occupies three distinct basins: (*i*) a most populated ground state (GS) with constitu-

tively active conformations competent for catalysis, (*ii*) a first high free energy basin representative

of typical inactive states with a dislodged configuration of the αC helix (ES1), and (*iii*) a second

high free energy basin with a disrupted hydrophobic array of residues at the core of the enzyme

(ES2). Notably, the equilibrium between the most populated ground state and the other low-pop-

ulated states agrees with previous NMR Carr-Purcell-Meiboom-Gill (CPMG) relaxation dispersion

and chemical exchange saturation transfer (CEST) measurements.[27] To compound the existence

of the second inactive basin, we mutated F100 at the αC-β4 loop into Ala (PKA-C^{F100A}) and char-

acterized its ligand binding thermodynamics and structural response by isothermal titration calo-

rimetry (ITC) and NMR spectroscopy, respectively. We found that PKA-C^{F100A} phosphorylates a

canonical peptide with a catalytic efficiency similar to the wild-type kinase (PKA-C^{WT}); however,

this mutation abolishes the binding cooperativity between nucleotide and substrate. The NMR

experiments reveal that F100A perturbs the hydrophobic packing around the αC-β4 loop and

interrupts the allosteric communication between the two lobes of the enzyme. Overall, these re-

sults emphasize the pivotal role of the αC-β4 loop in kinase function and may explain why single-

site mutations or insertion mutations stabilizing this motif in homologous kinases result in onco-

genes and confer differential drug sensitivity.[28-30]

RESULTS

The free energy landscape of PKA-C charted by NMR-restrained replica-averaged metadynamics (RAM) simulations. To characterize the experimentally accessible conformational landscape of PKA-C in the μs-to-ms timescale, we performed NMR-restrained metadynamics simulations within the RAM framework.[24,25] Using four replicas restrained with backbone chemical shifts (CS), we simulated the apo, binary (PKA-C/ATP), and ternary (PKA-C/ATP/PKI$_{5-24}$) complexes of PKA-C. Incorporating the CS restrains in the simulations ensures a close agreement with the conformational space explored by the kinase under our NMR experimental conditions,[31,32] whereas the enhanced sampling through metadynamics simulations boosts the conformational plasticity of the enzyme along different degrees of freedom (*i.e.*, collective variables, CVs. See Methods) (**Figure 2 - figure supplement 1**).[33] In our case, back-calculations of the CS values with Sparta+ [34] show that the restrained simulations improved the agreement between theoretical and experimental CS values of ~0.2 ppm for the amide N atoms and ~0.1 ppm for the remainder backbone atoms (**Figure 2 - figure supplement 2**). Also, the bias-exchange metadynamics allow each replica to span a significantly broader conformational space relative to classical MD simulations (**Figure 2 - figure supplement 3**). The metadynamics simulations were equilibrated and converged in 300 ns (**Figure 2A, Figure 2 - figure supplement 4**). The full free energy landscape was then reconstructed by sampling an extra 100 ns of production phase with reduced bias along each CV. We found that the population of the PKA-C conformers is modulated by ATP and pseudosubstrate binding within the NMR detection limit of sparsely populated states (~ 0.5% or ΔG < 3.2 kcal/mol).

Figure 2. Free energy landscape of PKA-C obtained from replica-averaged metadynamics (RAM) simulations. (**A**) Convergence of the bias deposition along the first three collective variables (CVs). The free energy (expressed in kcal/mol) of the different CVs was averaged over the last 100 ns of RAM simulations. The standard deviations are reported as red error bars. (**B-D**) Free energy landscape along the first two principal components (PC1 and PC2) of PKA-C in the apo, ATP-bound, and ATP/PKI bound forms. PC1 and PC2 are projected from the first three CVs. The vertices represent conformational states. In the apo form, multiple states have comparable free energy with ΔG < 5 kcal/mol, whereas in the binary form, fewer states have ΔG < 5 kcal/mol. For the ternary form only a major ground state is populated.

Specifically, the apo PKA-C populates preferentially a ground state and five readily accessible low-populated excited states (**Figure 2B**, **Figure 2 – supplementary table 1**). The binary complex populates the same ground state and a broad higher energy basin (**Figure 2C**, **Figure 2 – supplementary table 1**). Finally, the ternary complex occupies a narrower dominant ground state (**Figure 2D**, **Figure 2 – supplementary table 1**). Overall, these calculations revealed that the apo kinase explores multiple minima along each CV. The binding of ATP shifts the population of the ensemble into an additional minimum, featuring structures that are committed to substrate binding. Finally, the conformational heterogeneity of the ensemble is significantly reduced upon formation of the ternary complex, which represents a catalytically committed state[13] (**Figure 2A**).

The free energy landscape obtained from these RAM simulations is consistent with the qualitative

picture previously inferred from our NMR spin relaxation experiments and hydrogen/deuterium

exchange studies,[35-37] while providing a detailed structural characterization of the ground and

sparsely populated conformationally excited states.

MSM reveals the conformational transitions of PKA-C from ground to high free energy

(excited) states. To refine the free energy landscape and delineate the kinetics of conformational

transitions of the kinase, we performed additional unbiased sampling to build Markov State Mod-

els (MSM).[38,39] MSMs are commonly used to describe the dynamic transitions of various meta-

stable states for macromolecules in terms of their populations and transition rates. MSMs are

typically created by combining thousands of short unbiased simulations.[38,39] Following this strat-

egy, we performed several short simulations (10 – 20 ns) using thousands of the low free energy

conformations (ΔG < 3.2 kcal/mol) chosen randomly from the three forms of PKA-C as starting

structures. The conformational ensembles were clustered into microstates and seeded to start a

second round of adaptive sampling (see Methods). This iterative process was repeated three

times to assure convergence and yielded 100 μs trajectories for both the apo and binary forms,

whereas 60 μs trajectories were collected for the less dynamic ternary complex. Once we reached

a sufficient sampling, we built an MSM including L95, V104, L106, M118, M120, Y164, and F185

to investigate the dynamic transitions of the hydrophobic R spine and shell residues (**Figure 3 –**

figure supplement 1). These residues are ideal reporters of the dynamic processes governing

the activation/deactivation of the kinase.[40] To compare the free energy landscape of different

complexes, we projected the conformational ensembles of three forms and existing crystal struc-

tures of PKA-C onto the first two time-lagged independent components (tICs) of the apo form,

which were obtained by a time-lagged independent component analysis (tICA). These tICs rep-

resent the directions of the slowest motion of the kinase and visualize the conformational transi-

tions of V104, L95, and F185 (**Figure 3 – figure supplement 1**). The MSM shows that the hydro-

phobic core of the apo PKA-C accesses three major basins. The broadest basin represents the

ground state (GS) and corresponds to the conformations of PKA-C captured in several crystal

structures (**Figure 3A**). The other two states (conformationally excited states) are less populated.

The first excited state (ES1) features a disrupted hydrophobic packing of L95, V104, and F185.

This conformation matches an inactive state with an outward orientation of the αC-helix typical of

the inhibited states of PKA-C (**Figure 3 – Figure supplement 2**). The second state (ES2) displays

a flipped configuration of the V104 side chain and disrupted hydrophobic interactions responsible

for anchoring the αC-β4 loop to the C-lobe, with F100 and F102 adopting a gauche+ configuration

and forming a stable π-π stacking. (**Figure 3A**). In contrast, the active GS ensemble features a

trans configuration for the F100 and F102 side chains that stabilize the π-π stacking with Y156 at

the αE-helix and cation- π interactions with R308 at the C-tail (**Figure 3A**). To our knowledge, the

ES2 state was never observed in the available crystal structures. Upon ATP binding, the confor-

mational space spanned by the kinase becomes narrower, and the conformers populate mostly

the GS, with a small fraction in the ES1 state (**Figure 3B**). This is consistent with the role of the

nucleotide as an allosteric effector that enhances the affinity of the enzyme for the substrate.[41] In

the ternary form (ATP and PKI-bound), PKA-C populates only the GS consistent with the compe-

tent conformation observed in the ternary structure (*i.e.*, 1ATP). In this case, the αC-β4 loop of

the enzyme is locked in a well-defined, active configuration (**Figure 3A**).

Figure 3. Free energy surfaces and dynamic transitions determined by a Markov State Model (MSM) for apo, ATP-bound, and ATP/PKI-bound PKA-C. (**A**) Free energy landscape projected along the first two time-lagged independent components (tICs) of apo PKA-C, featuring three basins, GS, ES1, and ES2. The transition from GS to ES1 (arrow) highlights the changes around the αB-αC loop, with the disruption of the K72-E91 salt bridge and the PIF pocket (V80-I85-F347) hydrophobic interactions. The GS to ES2 transition (arrow) displays the rearrangement of the hydrophobic packing around the αC-β4 loop. (**B**, **C**) Free energy surfaces projected along the first two tICs for the ATP-bound and ATP/PKI-bound PKA-C, respectively. Known crystal structures for the three forms are indicated by small white triangles.

Moreover, in the apo form, the αC-β4 loop is quite dynamic due to transient hydrophobic inter-

actions involving F100 and V104, as well as W222 and the APE motif (A206 and P207) (**Figure**

4A). The binding of both nucleotide and PKI rigidifies the residues near F100 and V104, *e.g.*,

V103 and F185 (**Figure 4A**) and makes more persistent several electrostatic interactions essen-

tial for catalysis (D166-N171, K168-T201, and Y204-E230), which are transient in the apo PKA-

C. Finally, we used kinetic Monte Carlo sampling to characterize the slow transition between dif-

ferent states.[42] In the GS state of the apo PKA-C, F100 and F102 adopt *trans* configurations that

stabilize the interactions with the αE and αJ helices, and, together with the nucleotide, they lock

the αC-β4 loop in an active state (**Figure 3A**). The GS to ES1 transition features the disruption of

the K72-E91 salt bridge, typical of the inactive kinase. In contrast, the GS to ES2 transition in-

volves a 120° flip of the F100 aromatic group that interacts with V104, a conformation found only

in the catalytically uncommitted apoenzyme (**Figure 3A**). Interestingly, the GS to ES2 transition

involves a concerted disruption of the D166-N171 and K168-T201 electrostatic interactions and

the destabilization of the packing between W222 and the APE motif (A206 and P207) required for

substrate recognition (**Figure 3C**). Overall, these calculations show that conformational transi-

tions from the active GS to ES1 and ES2 represent two independent pathways toward inactive

states of the kinase.

Figure 4. Conformational transitions of apo and ATP-bound PKA-C from GS to ES1 and ES2 states as shown by the kinetic Monte Carlo trajectories. (**A**, **B**) Time course of the structural transitions from GS to ES1 GS to ES2 for apo and ATP-bound PKA-C, respectively. The GS to ES1 transition is characterized by the disruption of the K72-E91 salt bridge and occurs frequently for both PKA-C forms. In contrast, the structural transition from GS to ES2 occurs only for the apo PKA-C, and it features the interactions between F100 and V104 that cause allosteric changes between D166-N171, K168-T201, and W222-A206-P207. The dark color traces indicate the moving averages calculated every 10 frames. (**C**) Structural snapshot of the GS conformation showing that the key catalytic motifs are poised for phosphoryl transfer. (**D**) Structural snapshot of the ES2 conformation with a disrupted configuration of key catalytic motifs typical of inactive kinase.

Direct correspondence between the conformationally excited states identified by MD simulations and NMR data. NMR CPMG relaxation dispersion and CEST experiments performed on the apo PKA-C revealed the presence of conformationally excited states for several residues embedded into the hydrophobic core of the enzyme.[27] The chemical shifts of one of these states follow the previously observed open-to-closed transitions of the enzyme,[41] whereas another excited state observable in the CEST experiments does not follow the same trend. We surmised that this new state may represent an alternate inactive state.[27] Since ES2 features a disrupted hydrophobic packing of the αC-β4 loop and several of the methyl groups displaying the excited state in the CEST experiments are near the αC-β4 loop, we compared the CS differences of the methyl group from CEST experiments[27] ($\Delta\omega^{Exp}$) with those from the MD simulations ($\Delta\omega^{Pred}$). We analyzed the methyl groups of L103, V104, I150, L172, and I180, sampling more than 500 snapshots as representative conformations of the kinase in the ES2 and GS states (**Figure 3 – figure supplement 3**). We then computed the distribution of the methyl ^{13}C CS using ShiftX2.[43] For Val104 and Ile150, these calculations yielded $\Delta\omega^{Pred}$ values of 0.87 ± 0.02 and 0.85 ± 0.02 ppm between the ES2 and GS conformations, respectively, which are in good agreement with $\Delta\omega^{Exp}$ (1.10 ± 0.06 and 0.94 ± 0.09 ppm) obtained from the CEST experiments or fitting the CPMG dispersion curves (**Figure 5A-B**)**.** The $\Delta\omega^{Pred}$ values and the directions of the CS changes obtained for the remainder three sites are also in good agreement with the CEST profiles (**Figure 5 – figure supplement 1**). **Figure 4C** shows a linear fitting of $\Delta\omega^{Pred}$ and $\Delta\omega^{Exp}$ with a slope of 0.86 and R^2 of 0.82. The latter supports the hypothesis that the excited state observed by NMR may correspond to the simulated structural ensemble of the ES2 basin. Finally, MSM estimates a population of ES2 of 6 ± 2%, which is consistent with the 5 ± 1% population found by fitting the NMR data.[40] Altogether, the combination of NMR and MD simulations support the presence of an alter-

nate inactive state that features disrupted hydrophobic interactions near the αC-β4 loop that per-

turb the anchoring with the αE helix, and in turn, the structural couplings between the two lobes

of the kinase.

Figure 5. Comparison of experimental versus calculated ^{13}C chemical shifts of methyl
groups of PKA-C. (**A**) Experimental and calculated CS for Val104-Cγ1 of apo PKA-C. The GS is
in blue and the ES in magenta. (**B**) Corresponding CS profiles for Ile150-Cδ1. The experimental
CS is shown in dotted lines for GS (black) and ES (red). (**C**) Correlation between predicted | Δω^{Pred}|
and experimental | Δω^{Exp}| CS differences for methyl groups near the αC-β4 loop. The fitted linear
correlation has a slope of 0.86 and R^2 of 0.82.

The F100A mutation disrupts the allosteric network of the kinase. The above analysis

suggests that the destabilization of the αC-β4 loop may promote the GS to ES2 transition of the

kinase. To validate this hypothesis, we generated the F100A mutant seeking to increase the flex-

ibility of the αC-β4 loop. Starting from the coordinates of the X-ray structure of the ternary complex

(PDB ID: 4WB5), we simulated the F100A dynamics in an explicit water environment. First, we

performed a short equilibration using classical MD simulations. During 1 μs of MD simulations,

the αC-β4 loop of the F100A mutant undergoes a significant motion as manifested by increased

values of the backbone root mean square deviation (rmsd) and a conformational change (flipping)

of the F102 side chain (**Figure 6A**). This region in PKA-C^WT adopts a stable β-turn, with a persis-

tent hydrogen bond between the backbone oxygen of F100 and the amide hydrogen of L103. In

contrast, the hydrogen bond in F100A is formed more frequently between A100 and F102, result-

ing in an average γ-turn conformation (**Figure 6B**). Such local rearrangement not only disrupts

the hydrogen bond between N99 and Y156, altering the interactions between the αC-β4 loop and

the αE helix but also destabilizes the cation-π interaction between F102 and R308 at the C-tail

(**Figure 6B – supplementary figure 1**)

Figure 6. F100A mutation increases the dynamics of the αC-β4 loop, perturbing the local hydrophobic packing and its anchoring to the αE helix. (**A**) Time series of the αC-β4 loop dynamics, H-bond occurrence for the $\beta-$ and γ-turns, F102 χ_1 angle, and N99 and Y156 for WT (black) and F100A (red) in the ATP-bound state. (**B**) Representative structural snapshots showing the β-turn conformation for PKA-C^{WT} (green) and γ-turn for PKA-C^{F100A} (magenta).

The structural changes of the αC-β4 loop caused by the F100A mutation propagate through the protein core across the R spine, C spines, and shell residues (**Figure 7A**), and alter the response of the kinase to nucleotide binding. In the PKA-C^{WT}, ATP binding shifts the conformational ensemble of the kinase toward an intermediate state competent to substrate binding, as represented by the changes in the population densities of the hydrophobic core residues as a function of the rmsd (**Figure 7B**). For F100A, the population of the C spine follows the same trend of the WT kinase, with the shell residues already populating the intermediate state. In contrast, the R spine residues fail to adopt a competent state (**Figure 7B**). The latter is due to the perturbation of hydrophobic packing of L95 and L106 in the R spine, and V104 in the shell near the mutation site. Using the lowest principal components, we also analyzed the global dynamic response of the kinase to ATP binding. Not only does F100A change the breathing mode of the two lobes (PC1), but it also alters the shearing motion (PC2) of the binary complex, emphasizing the importance of this allosteric site for the inter-lobe communication (**Figure 7C-E**).

Figure 7. Structural responses to ATP binding of PKA-C^{F100A} mutant. (**A**) Superposition of the hydrophobic cores (C spine, R spine, and shell residues) for PKA-C^{WT} (lime) and PKA-C^{F100A} (hot pink), highlighting the structural perturbations of the R spine and shell residues. (**B**) Structural perturbation upon ATP binding for the hydrophobic core residues of PKA-C^{WT} and PKA-C^{F100A} shown as changes in the population densities vs. rmsd (**C, D**) First (PC1) and second (PC2) principal components describing breathing and shearing motions of the two lobes. (**E**) 2D projections and distributions of PC1 and PC2 for PKA-C^{WT} and PKA-C^{F100A}.

To analyze the internal correlated motions of the kinase, we further analyzed the MD trajectories of both wild-type and mutant enzyme using mutual information (**Figure 8**).[44] This method identifies correlated changes of backbone and side chain rotamers in proteins and detects clusters of residues that are responsible for intramolecular communication between active sites.[44] For PKA-C^{WT}, we observed numerous strong correlations that are clustered within each lobe and across the enzyme, connecting key motifs, such as the Gly-rich loop, αC-β4 loop, activation loop, and C-terminal tail (**Figure 8A**). These correlation patterns are reminiscent of the chemical shift correlations experimentally found for PKA-C^{WT},[20] which constitute the hallmark of the dynamically-committed state.[35,36] In contrast, the mutual information matrix for the F100A mutant displays

weaker and interspersed correlations across the enzyme. New correlations also are present

across the helical C-lobe suggesting a rewiring of the internal allosteric communication (**Figure**

8B). A possible explanation is that the increased motion caused by the elimination of the F100

aromatic side chain at the αC-β4 loop propagates throughout the entire hydrophobic core. This

structural reorganization causes the F100A kinase to adopt a dynamically uncommitted state.[35]

Taken together, the simulations of F100A suggest that perturbations of the αC-β4 loop increase

the local flexibility and break the structural connection between the two lobes, disrupting the cor-

related breathing/shearing motions highlighted in previous MD simulation studies[45,46]. Although

the single F100A mutation does not drive the kinase into a completely inactive state (ES2), it is

sufficient to abolish the structural couplings between the two lobes.

Figure 8. Mutual information analysis of backbone and side chain rotamers of ATP-bound
PKA-C^WT and PKA-C^F100A. (**A**) Mutual information matrix for PKA-C^WT showing well-organized
clusters of interactions within each lobe and distinct inter-lobe communication typical of an active
kinase. (**B**) Mutual information matrix for PKA-C^F100A revealing an overall reorganization of the
allosteric network caused by the disruption of the hydrophobic core.

Effects of the F100A mutation on the catalytic efficiency and binding thermodynamics

of PKA-C. Based on the MD simulations of the F100A mutant, we hypothesized that the disruption

of the structural coupling between the N-lobe and C-lobe would affect the binding cooperativity

between nucleotide (ATPγN) and pseudosubstrate inhibitor (PKI$_{5-24}$). To test this hypothesis, we

expressed, purified, and evaluated the catalytic efficiency for the WT and F100A kinases by car-

rying out steady-state coupled enzyme assays[47] using the standard substrate Kemptide.[48] F100A

showed only a slight increase in K_M and V_{max} compared to PKA-C^{WT}, resulting in a small reduction

of the catalytic efficiency (k_{cat}/K_M = 0.50 ± 0.04 and 0.41 ± 0.08 for PKA-C^{WT} and PKA-C^{F100A},

respectively; **Supplementary table 2**). We then performed ITC[49] to obtain ΔG, ΔH, $-T\Delta S$, and K_d,

and determine the cooperativity coefficient (σ) for ATPγN and PKI$_{5-24}$ binding. We first analyzed

the binding of ATPγN to the apo PKA-C^{F100A} and, subsequently, the binding of PKI$_{5-24}$ to the

ATPγN-saturated PKA-C^{F100A} (**Supplementary Table 3**). We found that PKA-C^{WT} and PKAC^{F100A}

have similar binding affinities for ATPγN (K_d = 83 ± 8 μM and 73 ± 2 μM, respectively). However,

in the apo form, F100A showed a 3-fold higher binding affinity for the pseudosubstrate relative to

PKA-C^{WT} (K_d = 5 ± 1 μM and 17 ± 2 μM, respectively - **Supplementary Table 3**). Upon saturation

with ATPγN, PKA-C^{F100A} displayed a 12-fold reduction in binding affinity for PKI$_{5-24}$, resulting in a

σ of ~3, a value significantly lower than the WT enzyme (σ greater than 100). These data support

the predictions of MD simulations and mutual information that ATP binding to F100A does not

promote a conformational state fully competent with substrate binding.

NMR mapping of nucleotide/pseudosubstrate binding response. To elucidate the atomic

details of the disrupted structural coupling and binding cooperativity for PKA-C^{F100A}, we used so-

lution NMR spectroscopy mapping its response to the nucleotide (ATPγN) and pseudosubstrate

(PKI$_{5-24}$) binding. Specifically, we monitored the ^{1}H and ^{15}N chemical shift perturbations (CSPs,

$\Delta\delta$) of the amide fingerprints for the binary (PKA-C^{F100A}/ATPγN) and ternary (PKA-

C^{F100A}/ATPγN/PKI$_{5-24}$) complexes and compared them with the corresponding complexes of PKA-

C^{WT} (**Figure 9** and **Figure 9 – figure supplement 1**). The interaction with ATPγN causes a dra-

matic broadening of several amide resonances throughout the structure of PKA-C^{F100A}, suggest-

ing the presence of an intermediate conformational exchange as previously observed.[50] For the

remainder of the amide peaks, both PKA-C^{WT} and PKA-C^{F100A} exhibit similar CSP profiles upon

binding ATPγN (**Figure 9A-B**), although the extent of the CS changes is significantly attenuated

for F100A (**Figure 9 – figure supplement 1B**). This reduction is apparent for residues in the N-

lobe (β2- β3 region), around the mutation site (β4), and at the C-terminal tail. A similar pattern

emerges upon binding PKI$_{5-24}$ to the ATPγN-saturated PKA-C^{F100A}, where a substantial decrease

in CSP is observed for residues of the C-lobe localized near the motifs critical for substrate-bind-

ing (*i.e.*, αE, αF, and αG helices, **Figure 9C-D**).

To determine the global response to ligand binding for WT and F100A, we analyzed the chem-

ical shifts of the amide fingerprints of the two proteins using the COordiNated ChemIcal Shifts

bEhavior (CONCISE).[51] CONCISE describes the overall changes of the protein fingerprint reso-

nances by providing the probability density (population) of each state along the conformational

equilibrium for binding phenomena that follow linear CS trajectories.[51] For PKA-C^{WT}, both nucle-

otide and pseudosubstrate binding shift the overall population of the amides from an open to an

intermediate and a fully closed state (**Figure 9 – figure supplement 2**). A similar trend is ob-

served for PKA-C^{F100A}, though the probability densities are broader, indicating that the amide res-

onances follow a less coordinated response.[51] Also, the maximum of the probability density for

the ternary complex is shifted toward the left, indicating that the mutant adopts a more open struc-

ture than the corresponding wild-type enzyme. Overall, the shapes and the positions of the prob-

ability distributions of the resonances suggests that several residues do not respond in a coordi-

nated manner to the nucleotide binding, and PKI$_{5-24}$ shifts the conformation of the kinase toward

a partially closed state, which explains the loss in binding cooperativity as previously observed.[21-

23]

Figure 9. NMR map of the structural response of PKA-C^{F100A} to nucleotide and PKI binding.
(**A**) Comparison of the chemical shift perturbation (CSP) of the amide resonances for PKA-C^{F100A} (black) and PKA-C^{WT} (cyan) upon ATPγN binding. The dashed line indicates one standard deviation from the average CSP. (**B**) CSPs of PKA-C^{F100A}/ATPγN amide resonances mapped onto the crystal structure (PDB: 4WB5). (**C**) Comparison of the CSPs of the amide resonances for PKA-C^{F100A} and PKA-C^{WT} upon binding ATPγN and PKI$_{5-24}$ (black). (**D**) CSPs for the F100A/ATPγN/PKI complex mapped onto the crystal structure (PDB: 4WB5).

To define the allosteric network of the kinase upon binding nucleotides and substrate, we examined the CS using CHEmical Shift Covariance Analysis (CHESCA),[22,23,27] a statistical method that identifies correlated responses of residue pairs to a specific perturbation (*i.e.*, ligand binding, mutations, *etc.*). CHESCA works under the assumption that pairwise correlated CS changes of residues identify possible intramolecular allosteric networks.[52,53] For PKA-C, we found that coordinated structural rearrangements, as identified by CHESCA, are directly related to the extent of binding cooperativity.[22,23,27] Therefore, we compared the CHESCA maps for PKA-C^{WT} and PKAC^{F100A} for four different states: *apo*, ATPγN-, ADP-, and ATPγN/PKI$_{5-24}$-bound. For PKA-C^{F100A},

the CHESCA matrix exhibits sparser and more attenuated correlations (*i.e.*, lower correlation co-

efficient value) relative to PKA-C^{WT} (**Figure 10A**). Although many inter-lobe correlations are still

present for F100A, several other correlations in specific structural domains such as the αG-, αH-,

and αI-helices are absent or attenuated. For instance, the F100A mutation does not display cor-

relations between the αA-helix and the C-terminal tail that constitute a critical "complement to the

kinase core".[54] We also utilized CHESCA to assess the allosteric communication among the PKA-

C communities as defined by McClendon *et al.*[55] The CHESCA community map for PKA-C^{WT}

shows strong correlations across the enzyme, especially for structurally adjacent communities

and at the interface between the two lobes (see for instance the correlations among *ComA*, *ComB*,

ComC, *ComE*, and *ComH*) (**Figure 10B-C**). For F100A, the CHESCA community map shows that

the cross-talk between the nucleotide-binding (*ComA*) and positioning of αC-helix (*ComB*) com-

munities, as well as the R-spine assembly (*ComC*) and the activation loop (*ComF*) communities

are preserved (**Figure 10B-C**). However, the correlations between *ComE*, responsible for stabi-

lizing the C spine, and *ComC*, involved in the assembly of the R spine, are absent. Similarly, the

long-range correlations between the C and R spines (*i.e.*, *ComD* with *ComC*) are missing. Finally,

several correlations between *ComF1*, *ComG*, and *ComH* are no longer present. These commu-

nities orchestrate substrate recognition and R subunits binding. Overall, the CHESCA analysis

for PKA-C^{F100A} suggests that the reduced degree of cooperativity we determined thermodynami-

cally corresponds to a decrease in coordinated structural changes upon ligand binding. The latter

is apparent from the loss of correlated structural changes among the structural communities, in-

cluding the hydrophobic spines, substrate binding cleft, and the docking surface for PKA interac-

tions with other binding partners.

Figure 10 – Correlated chemical shift changes reveal the uncoupling of the intramolecular allosteric network in PKA-C[F100A]. (**A**) Comparison of the CHESCA matrices obtained from the analysis of the amide CS of PKA-C[WT] (blue correlations) and PKA-C[F100A] (black correlations). The correlations coefficients (R_{ij}) were calculated using the apo, ADP-bound, ATPγN-bound, and ATPγN/PKI[5-24]-bound states. For clarity, only correlation with $R_{ij} > 0.98$ are displayed. For the enlarged CHESCA map of F100A see **Figure 10 – figure supplement 1.** The data for the PKA-C[WT] matrix were taken from Walker *et al.*[22]. (**B**) Community CHESCA analysis of PKA-C[WT] (blue correlations) and PKA-C[F100A] (black correlations). Only correlations with $R_{A,B} > 0.98$ are shown. (**C**) Spider plot showing the extent of intramolecular correlations identified by the community CHESCA analysis for PKA-C[F100A] mapped onto the crystal structure (PDB: 4WB5). The thickness of each line in the spider plot indicates the extent of coupling between the communities.

DISCUSSION

Our structural and dynamic studies suggest that the binding cooperativity of PKA-C originates from the allosteric coupling between the nucleotide-binding pocket and the interfacial region between the two lobes, which harbors the substrate binding cleft.[21-23] The latter has been emphasized for other kinases such as Src and ERK2.[56,57] The biological significance of this intramolecular communication has emerged from our recent studies on pathological mutations situated in the activation loop of PKA-C and linked to Cushing's syndrome.[21-23] These mutations drastically reduce substrate binding affinity, and more importantly, disrupt the communication between the two ligand binding pockets of the kinase.[21-23] Interestingly, the E31V mutation distal from the active site and also related to the Cushing's syndrome, displays disfunction similar to the other orthosteric mutations, suggesting that it is possible to modulate substrate recognition allosterically.[23]

463 Indeed, these mutations are connected to allosteric nodes that, once perturbed, radiate their ef-

464 fects towards the periphery of the enzyme interrupting the coordinated dynamic coupling between

465 the two lobes.[23] Interestingly, these mutations do not prevent Kemptide phosphorylation; rather,

466 they cause a loss of substrate fidelity with consequent aberrant phosphorylation of downstream

467 substrates.[10,58-62] Additionally, recent thermodynamic and NMR analyses of PKA-C binding to dif-

468 ferent nucleotides and inhibitors demonstrated that it is possible to control substrate binding af-

469 finity by changing the chemistry of the ligand at the ATP binding pocket.[22] Altogether, these stud-

470 ies put forward the idea that substrate phosphorylation and cooperativity between ATP and sub-

471 strates may be controlled independently.

472 By integrating our NMR data with RAM simulations and MSM, we comprehensively mapped

473 the free energy landscape of PKA-C in various forms. We found that the active kinase unleashed

474 from the regulatory subunits occupies a broad energy basin (GS) that corresponds to the confor-

475 mation of the ternary structure of PKA-C with ATP and pseudosubstrate (PKI$_{5-24}$), exemplifying a

476 catalytic competent state poised for phosphoryl transfer.[15] We also identified two orthogonal con-

477 formationally excited states, ES1 and ES2. While ES1 corresponds to canonical inactive kinase

478 conformations, the ES2 state was never observed in the crystallized structures. Our previous

479 CEST NMR measurements suggested the presence of an additional sparsely populated state that,

480 at that time, we were unable to characterize structurally.[27] The chemical shifts of this state did not

481 follow the opening and closing of the kinase active cleft and were assigned to a possible alternate

482 inactivation pathway.[27] These new simulations and MSM show that this sparsely populated state

483 may be attributed to the transition from GS to ES2, which features a disruption of hydrophobic

484 packing, with a conformational rearrangement for the αC-β4 loop that causes a partial disruption

485 of the hydrophobic R spine. These structural changes rewire the allosteric coupling between the

486 two lobes, as shown by mutual information analysis. A single mutation (F100A), suggested by our

simulations, promotes the flip of the αC-β4 loop and reproduces the hypothesized structural un-

coupling between the two lobes of PKA-C. We experimentally tested the effects of the F100A

mutation and found that it prevents the enzyme from adopting a conformation competent for sub-

strate binding, resulting in a drastic reduction of the cooperativity between ATP and substrate.

These NMR data further support our working model, showing that if the inter-lobe communication

is interrupted, the binding response of the kinase is attenuated. Indeed, our investigation further

emphasizes the integral role of the kinase hydrophobic core and demonstrates that alterations in

the spines and shell residues may lead to a dysfunctional kinase by either preventing phosphor-

ylation (see V104G and I150A mutations)[40] or by disrupting the binding cooperativity as for F100A.

The αC-β4 loop is a regulatory element present in all EPKs and its importance has been

stressed in bioinformatics studies[63] and supported by computational work.[55] F100 and F102 in the

αC-β4 loop are at intersections of various structural communities and constitute a critical hydro-

phobic node that anchors the N- to the C-lobe. Also, studies on EGFR and ErbB2 kinases led to

the hypothesis that the αC-β4 loop may act as a molecular brake [64,65] or an autoinhibitory switch.[66]

Therefore, it is not surprising that activating mutations and in-frame insertions in the αC-β4 loop

are frequently found in kinase-related cancers and somatic oncogenic mutations such as P101S,

P101L, and L103F.[67] Also, an elegant study by Kannan and coworkers emphasized the role of

the αC-β4 loop in dimerization and aberrant activation of EGFR by insertion mutations.[28] Moreo-

ver, in a recent paper Zhang and co-workers found a similar behavior for the HER2 exon 20

insertions.[30] These latter two studies propose that in-frame insertions alter the conformational

landscape of the kinase rigidifying the structure around the αC-β4 loop and restricting the kinase

conformational ensemble in a constitutively active state. Remarkably, in the case of EGRF inser-

tion mutations, the activation of the kinase occurs in a length-dependent manner. [28] Additionally,

these human mutations display a gradual response to drugs, which can be exploited for designing

selective inhibitors against EGFR pathological mutants.[68]

The data presented here show that it is possible to abolish the binding cooperativity of a kinase

by turning the dial in the opposite direction, *i.e.*, increasing the flexibility of the αC-β4 loop and

disconnecting the allosteric network between the N- and C-lobes. The identification of this new,

partially inactivating pathway provides further understanding of how to control the dynamics and

function of kinases.

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Gene (*Homo sapiens)*	PKA-Cα		PKA-C	Uniprot ID: P17612
strain, strain background (*Escherichia coli*)	BL21(DE3) pLyss	Agilent	Cat. #200132	Chemically competent cells
Sequence-based reagent	PKA-C^{F100A}	This study	PCR primer (Forward)	tattctgcaagcggtg aacgccccgtttctg gttaagctg
Sequence-based reagent	PKIα 5-24	Synthetic peptide	PKI$_{5-24}$	Chemically synthesized
Sequence-based reagent	Kemptide	Synthetic peptide	Kemptide	LRRASLG
Commercial assay or kit	QuikChange Lightning Multi Mutagenesis Kit	Agilent genomics	Cat #210519	Commercial mutagenesis kit
Recombinant DNA reagent	PKA-^{CF100A}	This study	PKA-C^{F100A}	Single Ala mutant of PKA-C
Chemical compound, drug	AMP-PNP or ATPγN	Roche Applied Science	CAS 25612-73-1	ATP analogous
Chemical compound, drug	ADP	Sigma-Aldrich	CAS 20398-34-9	nucleotide
Software, algorithm	TopSpin 4.1	Bruker Inc.	https://www.bruker .com/	
Software, algorithm	NMRFAM-Sparky	NMRFAM	https://nmr-fam.wisc.edu/nmr-fam-sparky-distri-bution/	

Software, algorithm	NMRPipe	Delaglio F., NIH	https://www.ibbr.umd.edu/nmrpipe/install.html	
Software, algorithm	POKY	Lee W.,	https://sites.google.com/view/pokynmr	
Software, algorithm	COordiNated ChemIcal Shift bEhavior (CONCISE)	Veglia G.	https://conservancy.umn.edu/handle/11299/217206 https://conservancy.umn.edu/handle/11299/227294	Matlab script
Software, algorithm	CHEmical Shift Covariance Analysis (CHESCA)	Melacini G.	https://academic.oup.com/bioinformatics/article/37/8/1176/5905475?login=true	NMRFAM-Sparky&POKY tools
Software, algorithm	PyMol	Schrödinger, LLC	https://pymol.org	
Software, algorithm	MatLab2022b	MathWorks	https://www.math-works.com/products/matlab.html	
Software, algorithm	GraphPad Prism 9	GraphPad Software Inc.	https://www.graphpad.com/	
Software, algorithm	GROMACS 4.6	Hess B et al.	http://www.gromacs.org/	
Software, algorithm	CHARMM36a1	Best RB et al. (2012)	https://pubs.acs.org/doi/10.1021/ct300400x	
Software, algorithm	PLUMED 2.1.1	Bonomi M, et al. (2009)	https://www.plumed.org/doc-v2.5/user-doc/html/_c_h_a_n_g_e_s-2-1.html	
Software, algorithm	ALMOST 2.1	Kohlhoff et al. (2009).	svn://svn.code.sf.net/p/al-most/code/almost-code	
Software, algorithm	METAGUI	Biarnés X et al. (2012)	https://www.sciencedirect.com/scienc	

			e/article/pii/S0010 465511003079	
Software, algo-rithm	MDTraj	McGibbon RT et al. (2015)	https://www.scienc edirect.com/scienc e/article/pii/S0006 349515008267	
Software, algo-rithm	SPARTA+	Yang Shen, Ad Bax (2010)	https://spin.niddk.b ov/bax/software/S PARTA+/	
Software, algo-rithm	MSMbuilder	Beauchamp KA et al. (2011)	https://msmbuilder .org/	
Software, algo-rithm	Mutinf	McClendon C et al. (2012)	https://simtk.org/pr ojects/mutinf/	

MATERIAL AND METHODS

Replica-averaged metadynamics (RAM) simulations

I. System setup. As a starting structure, we used the coordinates of the crystal structure of PKA-C^{WT} (PDB ID: 1ATP) and added the missing residues 1-14 at the N terminus. The protonation state of histidine residues was set as previously reported.[69] The kinase was solvated in a rhombic dodecahedron solvent box using the three-point charge TIP3P model[70] for water, which extended approximately 10 Å away from the surface of the proteins. Counter ions (K+ and Cl-) were added to ensure electrostatic neutrality to a final ionic concentration of ~ 150 mM. All protein covalent bonds were constrained using the LINCS algorithm[71] and long-range electrostatic interactions were simulated using the particle-mesh Ewald method with a real-space cut-off of 10 Å.[72] The simulations of apo, binary (one Mg^{2+} ion and one ATP), and ternary (two Mg^{2+} ions, one ATP, and PKI_{5-24}) forms were performed simultaneously using GROMACS 4.6[73] with CHARMM36a1 force field.[74] For F100A, the corresponding residues were mutated through the mutagenesis wizard of Pymol.[75]

II. Standard MD simulations. Each system was minimized using the steepest descent algorithm to remove the geometric distortions, and then gradually heated to 300 K at a constant volume

over 1 ns using harmonic restraints with a force constant of 1000 kJ/(mol *$Å^2$) on heavy atoms for

both the kinase and nucleotide. The restraints were gradually released over the following 12 ns

of simulations at constant pressure (1 atm) and temperature (300 K). The systems were equili-

brated for an additional 20 ns without positional restraints. A Parrinello-Rahman barostat[76] was

used to keep the pressure constant, while a V-rescale thermostat [77] with a time step of 2 fs was

used to keep the temperature constant. Each system was simulated for 1.05 μs, with snapshots

recorded every 20 ps.

III. Replica exchange (REX) simulations. Following standard MD simulations, parallel replica ex-

change (REX) simulations were set up on the apo, binary, and ternary forms of PKA-C. Four

replicas were used for each REX simulation, and the initial structures were randomly chosen from

the μs-scale unbiased simulations. Chemical shifts of PKA-C for N, CA, CO, CB, and HN from

NMR experiments were imposed as restraints based on the following penalty function:

547
$$E^{cs} = \alpha \sum_{k=1}^{N} \sum_{l=1}^{5} \left(\delta_{kl}^{exp} - \frac{1}{M} \sum_{m=1}^{M} \delta_{kl}^{calc} \right)^2$$

where α is the force constant, k runs over all residues of the protein, l denotes the different back-

bone atoms, and *m* runs over the four replicas. δ_{kl}^{calc} was computed using CamShift, a module of

ALMOST-2.1.[78] The force constant was gradually increased from 0 (unbiased) to 20 (maximum

restraints for production) over 50 ns. All other settings were identical to the unbiased simulations.

REX simulations were carried out with GROMACS 4.6,[73] with the replica exchange controlled by

the PLUMED 2.1.1 module.[79] Approximately, 100 ns of REX simulations were further carried out

for each replica in the three PKA-C forms.

IV. Replica-averaged metadynamics (RAM) simulations. RAM simulations were started from the

final structures of REX simulations. The CS restraints were imposed in the same manner as for

the REX simulations. Four CVs are chosen to increase the conformational plasticity around the

catalytic cores (detailed in **Figure S1**): (CVI) the ψ angles of the backbone of all the loops that

are not in contact with ATP (Back-far), (CVII) the ϕ angles of the backbone of all the loops that

are in contact with ATP (Back-close), (CVIII) the χ_1 angles of side-chains of all the loops that are

in contact with ATP (Side-close), (CVIV) the radius of gyration calculated over the rigid part (*i.e.*,

residues 50-300) of the protein (rgss). Gaussian deposition rate was performed with an initial rate

of 0.125 kJ/mol/ps, where the σ values were set to 0.5, 0.2, 0.2, and 0.01 nm for the four CVs,

respectively. The RAM simulations were also carried out with GROMACS 4.6 in conjunction with

PLUMED 2.1 and ALMOST 2.1, and continued for ~400 ns for each replica with exchange trails

every 1 ps.

V. Reconstruction of free energy surface (FES) from the RAM simulations. After about 300 ns,

the sampling along the first 3 CVs reached convergence with fluctuations of bias within 1 kcal/mol,

allowing a reliable reconstruction of the corresponding FES. The production run was continued

for another 100 ns to sample enough conformations. These conformations were first clustered

into microstates using the regular spatial method (cut-off radius of 0.13), and then the free energy

of each state was reweighted according to the deposited potential along each CV, which can be

obtained from the analysis module of METAGUI.[80] To visualize the distribution of these mi-

crostates and their relative energy differences, we further performed principal component analysis

to project the microstates represented in the 3-dimensional CV space into 2-dimensional space

spanned by PC1 and PC2. Then we plotted these microstates in a 3-dimensional space spanned

by PC1, PC2, and free energy differences ΔG.

VI. Independent validation of chemical shifts with SPARTA+. During the REX and RAM simula-

tions, the chemical shifts were computed via CamShift in ALMOST 2.1.[78] As an independent

validation for the efficacy of the bias, we further calibrated the chemical shifts of 2000 snapshots

with MDTraj[81] and SPARTA+.[34]

VII. Adaptive sampling. The first round of adaptive sampling started from the snapshots of low-

energy microstates obtained from the previous step, *i.e.*, 1200 structures for the apo form, 400

structures for the binary form, and 200 structures for the ternary form. The initial velocities were

randomly generated to satisfy the Maxwell distribution at 300K. For the apo form, a 10 ns simula-

tion was performed for each run, whereas for the binary, each simulation lasted 30 ns, resulting

in a total of 12 µs trajectories for both the apo and binary forms. Three rounds of adaptive sam-

pling were started from the 400 microstates, derived via K-mean clustering of all snapshots of

previous ensembles, to obtain a converged free energy landscape. Therefore, a total of 100 µs

trajectories and 100,0000 snapshots (100 ps per frame) were collected for both the apo and binary

form after three rounds of adaptive sampling, and a total of 60 µs trajectories were collected for

the ternary form.

VIII. Markov state model (MSM) and time-lagged independent component analysis (tICA). The

Cartesian coordinates of key hydrophobic residues, including R-spine residues, L95, L106, Y164,

and F185, and the shell residues, V104, M118, and M120, were chosen as the metrics to char-

acterize the conformational transition of the hydrophobic core of PKA-C. Specifically, each snap-

shot was first aligned to the same reference structure by superimposing the αE (residues 140-

160) and αF helices (residues 217-233) and represented by the deviation of the Cartesian coor-

dinates of key residues. The representation in this metric space was further reduced to 10-dimen-

sion vectors using time-lagged independent component analysis (tICA)[82] at a lag time of 1 ns. All

snapshots were clustered into 400 microstates with K-mean clustering. An MSM was built upon

the transition counts between these microstates.

IX Kinetic Monte Carlo trajectory of PKA-C in different forms. Long trajectories were generated

using a kinetic Monte Carlo method based on the MSM transition probability matrix of the three

forms of PKA-C. Specifically, the discrete jumps between the 100 microstates were sampled for

60 µs. Random conformations were chosen for each state from all the available snapshots. Sub-

sequently, the time series of various order parameters were analyzed and plotted.

Protein expression and purification

The recombinant human Cα subunit of cAMP-dependent protein kinase with the Phe to Ala mutation in position 100 (PKA-C^{F100A}) was generated from the human PKA-Cα wild-type using Quik-Change Lightning mutagenesis kit (Agilent genomics). The key resource table lists the PCR primers used to modify the pET-28a expression vector encoding for the wild-type human PKA-Cα gene (*PRKACA* – uniprot P17612) [21-23]. The unlabeled and uniformly ^{15}N-labeled PKA-C^{F100A} mutant was expressed and purified following the same protocols used for the wild-type protein.[41] Briefly, transformed *E. coli* BL21 (DE3) *pLyss* cells (Agilent) were cultured overnight at 30 °C in Luria-Bertani (LB) medium. The next morning, the cells were transferred to fresh LB medium for the overexpression of the unlabeled protein or to M9 minimal medium supplied with ^{15}NH$_4$Cl (Cambridge Isotope Laboratories Inc.) as the only nitrogen source for the labeled protein overexpression. In both cases, protein overexpression was induced with 0.4 mM of β-D-thiogalactopyranoside (IPTG) and carried out for 16 hours at 20 °C. The cells were harvested by centrifugation and resuspended in 50 mM Tris-HCl, 30 mM KH$_2$PO$_4$, 100 mM NaCl, 5 mM 2-mercaptoethanol, 0.15 mg/mL lysozyme, 200 µM ATP, DNaseI, 1 tablet of cOmplete ULTRA EDTA-free protease inhibitors (Roche Applied Science) (pH 8.0) and lysed using French press at 1,000 psi. The cell lysate was cleared by centrifugation (60,000 × g, 4 °C, 45 min), and the supernatant was batch-bound with Ni^{2+}-NTA agarose affinity resin (Thermo Scientific). The his-tagged PKA-C^{F100A} was eluted with 50 mM Tris-HCl, 30 mM KH$_2$PO$_4$, 100 mM NaCl, 5 mM 2-mercaptoethanol, 0.5 mM phenylmethylsulfonyl fluoride (PMSF) (pH 8.0) supplied with 200 mM of imidazole. The tail of poly-His was cleaved using a stoichiometric amount of recombinant tobacco etch virus (TEV) protease in 20 mM KH$_2$PO$_4$, 25 mM KCl, 5 mM 2-mercaptoethanol, 0.1 mM PMSF (pH 6.5), overnight at 4 °C. The different phosphorylation states of PKA-C^{F100A} were separated using a cation exchange column (HiTrap Q-SP, GE Healthcare Life Sciences) using a linear gradient of KCl in 20 mM KH$_2$PO$_4$ at pH6.5.[83] The purified protein isoforms were then stored in phosphate buffer

containing 10 mM dithiothreitol (DTT), 10 mM MgCl$_2$, and 1 mM NaN$_3$ at 4 °C. The protein purity

was assessed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS–PAGE).

Peptide synthesis

The Kemptide (LRRASLG) and PKI$_{5–24}$ (TTYADFIASGRTGRRNAIHD) peptides were synthesized

using a CEM Liberty Blue microwave synthesizer using standard Fmoc chemistry. All peptides

were cleaved with Reagent K (82.5% TFA, 5% phenol, 5% thioanisole, 2.5% ethanedithiol, and

5% water) for 3 h and purified using a semipreparative Supelco C18 reverse-phase HPLC column

at 3 mL/min. The purified peptides were concentrated, lyophilized, and stored at −20 °C. Molecu-

lar weight and quantity were verified by MALDI-TOF and/or amino-acid analysis (Texas A&M Uni-

versity).

Isotherm titration calorimetry (ITC) measurements

PKA-C^{F100A} was dialyzed into 20 mM MOPS, 90 mM KCl, 10 mM DTT, 10 mM MgCl$_2$, and 1 mM

NaN$_3$ (pH 6.5) and concentrated using conical spin concentrator (10 kDa membrane cut-off, Mil-

lipore) to a solution at 80-100 μM, as confirmed by A280 = 55,475 M^{-1} cm^{-1}. Approximately 300 μL

of protein was used for each experiment, with 50 μL of 2 mM ATPγN and/or 1 mM PKI$_{5–24}$ in the

titrant syringe. All measurements were performed at 300 K in triplicates with a low-volume

NanoITC (TA Instruments). The binding was assumed to be 1:1, and curves were analyzed with

the NanoAnalyze software (TA Instruments) using the Wiseman isotherm [49]

652
$$\frac{d[MX]}{d[X_{tot}]} = \Delta H^\circ V_0 \left[\frac{1}{2} + \frac{1 - \dfrac{1 - r}{2} - R_m/2}{(R_m^2 - 2R_m(1 - r) + (1 + r)^2)^{1/2}} \right]$$

653   where $d[MX]$ is the change in total complex relative to the change in total protein concentration,

654     $d[X_{tot}]$ is dependent on r (the ratio of K_d relative to the total protein concentration), and R_m (the

655   ratio between total ligand and total protein concentration). The heat of dilution of the ligand into

656   the buffer was considered for all experiments and subtracted.

657   The free energy of binding was determined from:

$$\Delta G = RTlnK_d$$

659 where R is the universal gas constant and T is the temperature at measurement (300 K). The

660 entropic contribution to binding was calculated using:

$$T\Delta S = \Delta H - \Delta G$$

662 The degree of cooperativity (σ) was calculated as:

$$\sigma = \frac{K_{d\,apo}}{K_{d\,nucleotide}}$$

664 where $K_{d\,apo}$ is the dissociation constant of PKI$_{5-24}$ binding to the apo-enzyme, and $K_{d\,Nucleotide}$ is the

665 corresponding dissociation constant for PKI$_{5-24}$ binding to the nucleotide-bound kinase.

Enzyme assays

667 Steady-state coupled enzyme activity assays using Kemptide as substrate were performed under

668 saturating ATP concentrations and spectrophotometrically at 298 K, as reported by Cook *et al.*[47]

669 The values of V_{max} and K_M were obtained from a nonlinear fit of the initial velocities to the Mich-

670 aelis-Menten equation.

NMR spectroscopy

NMR measurements were performed on a Bruker Avance NEO spectrometer operating at a ^{1}H

Larmor frequency of 600 MHz equipped with a cryogenic probe or on a Bruker Avance III 850

MHz spectrometer equipped with a TCI cryoprobe. The NMR experiments were recorded at 300K

in 20 mM KH$_2$PO$_4$ (pH 6.5), 90 mM KCl, 10 mM MgCl$_2$, 10 mM DTT, 1 mM NaN$_3$, 5% D$_2$O, and

0.1% 4-benzene sulfonyl fluoride hydrochloride (AEBSF, Pefabl–c - Sigma-Aldrich). Concentra-

tions for samples were 0.15 mM of uniformly ^{15}N-labeled PKA-C^{F100A}, as determined by A$_{280}$ meas-

urements, 12 mM ATPγN or ADP was added for the nucleotide-bound form, and 0.3 mM PKI$_{5-24}$

for the ternary complex. [^{1}H, ^{15}N]-WADE-TROSY-HSQC pulse sequence[84] was used to record the

amide fingerprint spectra of PKA-C^{F100A} in the apo, nucleotide-bound (ADP- or ATPγN-bound –

binary form), and ternary complex (PKAC^{F100A}/ATPγN/PKI$_{5-24}$). All [^{1}H, ^{15}N]-WADE-TROSY-HSQC

experiments were acquired with 2048 (proton) and128 (nitrogen) complex points, processed us-

ing NMRPipe[85] and visualized using NMRFAM-SPARKY [86] and POKY.[87] Combined chemical shift

perturbations (CSP) were calculated using ^{1}H and ^{15}N chemical shifts according to:

$$\Delta\delta = \sqrt{(\Delta\delta H)^2 + (0.154 \times \Delta\delta N)^2}$$

in which $\Delta\delta$ is the CSP; $\Delta\delta_H$ and $\Delta\delta_N$ are the differences of ^{1}H and ^{15}N chemical shifts, respectively,

between the first and last point of the titration; and 0.154 is the scaling factor for nitrogen.[88]

COordiNated ChemIcal Shift bEhavior (CONCISE). The normal distributions reported in the

CONCISE plot were calculated using principal component analysis for residues whose chemical

shifts responded linearly to ligand binding.[51] In this work, we use the ^{1}H and ^{15}N chemical shifts

derived from the [^{1}H, ^{15}N]-WADE-TROSY-HSQC experiments for the apo, ADP, ATPγN, and

ATPγN/PKI$_{5-24}$ bound forms of PKA-C.

CHEmical Shift Covariance Analysis (CHESCA). This analysis was used to identify and char-

acterize allosteric networks of residues showing concerted responses to nucleotide and pseudo-

substrate binding. To identify inter-residue correlations, four states were used: apo, ATPγN-bound,

ADP-bound, and ATPγN/PKI$_{5-24}$. The identification of inter-residue correlations by CHESCA relies

on agglomerative clustering (AC) and singular value decomposition (SVD).[53] Pairwise correlations

of chemical shift changes were calculated to identify networks. When plotted on a correlation

matrix, the 2D correlations allow the identification of local and long-range correlated residues. For

each residue, the maximum change in the chemical shift was calculated in both the ^{1}H (*x*) and

^{15}N (*y*) dimensions ($\Delta\delta_{x,y}$). The residues included in the CHESCA analysis were those satisfying

$\Delta\delta_{x,y} > \frac{1}{2}\,\Delta\nu_{xA,yA} + \frac{1}{2}\,\Delta\nu_{xB,yB}$, where A and B correspond to two different forms analyzed, and $\Delta\nu$

denotes the linewidth. Correlation scores were used to quantify the CHESCA correlation of a

single residue or a group of residues with another group. Correlation scores were evaluated for

single residues or for the entire protein using the following expression:

706
$$Corr\ Score = \frac{number\ of\ (R_{ij} > R_{cutoff})}{total\ number\ of\ R_{ij}}$$

where R_{ij} is the correlation coefficient and i and j denote (a) a single residue and the remainder

residues of the protein, respectively, or (b) both represent all the assigned residues in the entire

protein. For all the analyses a R_{cutoff} of 0.98 was used. Community CHESCA analysis utilizes a

similar approach to map correlations between functional communities within the kinase. Each

community represents a group of residues associated with a function or regulatory mechanism.[55]

To represent community-based CHESCA analysis, we utilized R_{cutoff} > 0.8. For instance, to repre-

sent a chemical shift correlation between two community (X and Y) with n_A and n_B number of

assigned residues, respectively, the correlation score between A and B is defined as

$$R_{A,B} = number\ of\ (R_{ij} > R_{cutoff})/(n_A * n_B)$$

where R_{ij} is the correlation coefficient between residue i (belonging to community A) and residue

j (belonging to community B). R_{cutoff} is the correlation value cutoff. $R_{A,B}$ assumes values from 0 (no

correlation between residues in A and B) to 1 (all residues in A have correlations > cut-off with all

residues in B).

Acknowledgments

721   This work was supported by the National Institute of Health GM 100310 and HL 144130 to GV.

722   The authors would like to acknowledge the Minnesota Supercomputing Institute for MD calcula-

723   tions. YW would like to thank the Guangdong Pearl River Talent Program (2021QN02Y618) and

724   the National Natural Science Foundation of China (22007069, 92269102) for part of the MD anal-

725   ysis carried out at the Shenzhen Bay Laboratory Supercomputing Centre.

Competing interest

727   The authors declare no financial and non-financial competing interests.

Author contributions

Y.W., C.O., and G.V. designed research. Y.W. performed the MD simulations and analysis. C.O. prepared the kinase samples, executed all NMR and ITC experiments, and analyzed the results. C.W. performed the kinase activity assay and prepared the peptides for the assay. C.O., Y.W. contributed to the manuscript, preparing all the text and figures. M.V.S., K.N.H., C.C., J.G., and M.V. contributed to the critical analysis of the data and writing of the manuscript. D.A.B. contributed to the critical analysis of the data and writing the manuscript. S.S.T. contributed to the writing of the manuscript. G.V. designed and directed the experiments and contributed to the writing of the manuscript. All authors have approved the final version of the manuscript.

To whom correspondence should be addressed. Email: vegli001@umn.edu.

Data Availability

All the data generated or analyzed in this study are included in the manuscript and supporting files. The NMR chemical shifts and MD trajectories are deposited in the Data Repository for the University of Minnesota, https://hdl.handle.net/11299/261043.

References

1. Manning, G., Whyte, D.B., Martinez, R., Hunter, T. & Sudarsanam, S. The Protein Kinase Complement of the Human Genome. *Science* **298**, 1912-1934 (2002).
2. Huse, M. & Kuriyan, J. The Conformational Plasticity of Protein Kinases. *Cell* **109**, 275-282 (2002).
3. Knighton, D.R. et al. Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 407-14 (1991).
4. Knighton, D.R. et al. Structure of a peptide inhibitor bound to the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 414-20 (1991).
5. Taylor, S.S., Ilouz, R., Zhang, P. & Kornev, A.P. Assembly of allosteric macromolecular switches: lessons from PKA. *Nat Rev Mol Cell Biol* **13**, 646-58 (2012).
6. Walsh, D.A., Perkins, J.P. & Krebs, E.G. An adenosine 3',5'-monophosphate-dependant protein kinase from rabbit skeletal muscle. *J Biol Chem* **243**, 3763-5 (1968).
7. Smith, F.D. et al. Local protein kinase A action proceeds through intact holoenzymes. *Science* **356**, 1288-1293 (2017).
8. Tillo, S.E. et al. Liberated PKA Catalytic Subunits Associate with the Membrane via Myristoylation to Preferentially Phosphorylate Membrane Substrates. *Cell Rep* **19**, 617-629 (2017).
9. Xiong, W.H., Qin, M. & Zhong, H. PKA regulation of neuronal function requires the dissociation of catalytic subunits from regulatory subunits. *bioRxiv* (2023).
10. Omar, M.H. et al. Mislocalization of protein kinase A drives pathology in Cushing's syndrome. *Cell Rep* **40**, 111073 (2022).
11. Knighton, D.R., Xuong, N.H., Taylor, S.S. & Sowadski, J.M. Crystallization studies of cAMP-dependent protein kinase. Cocrystals of the catalytic subunit with a 20 amino acid residue peptide inhibitor and MgATP diffract to 3.0 A resolution. *J Mol Biol* **220**, 217-20 (1991).
12. Johnson, D.A., Akamine, P., Radzio-Andzelm, E., Madhusudan, M. & Taylor, S.S. Dynamics of cAMP-dependent protein kinase. *Chem Rev* **101**, 2243-70 (2001).
13. Bastidas, A.C. et al. Phosphoryl Transfer by Protein Kinase A Is Captured in a Crystal Lattice. *Journal of the American Chemical Society* **135**, 4788-4798 (2013).
14. Gerlits, O. et al. Phosphoryl Transfer Reaction Snapshots in Crystals: INSIGHTS INTO THE MECHANISM OF PROTEIN KINASE A CATALYTIC SUBUNIT. *J Biol Chem* **290**, 15538-15548 (2015).
15. Gerlits, O. et al. Zooming in on protons: Neutron structure of protein kinase A trapped in a product complex. *Sci Adv* **5**, eaav0482 (2019).
16. Kornev, A.P., Haste, N.M., Taylor, S.S. & Eyck, L.F. Surface comparison of active and inactive protein kinases identifies a conserved activation mechanism. *Proc Natl Acad Sci U S A* **103**, 17783-8 (2006).
17. Ten Eyck, L.F., Taylor, S.S. & Kornev, A.P. Conserved spatial patterns across the protein kinase family. *Biochim Biophys Acta* **1784**, 238-43 (2008).
18. Hu, J. et al. Kinase regulation by hydrophobic spine assembly in cancer. *Mol Cell Biol* **35**, 264-76 (2015).
19. Lew, J., Coruh, N., Tsigelny, I., Garrod, S. & Taylor, S.S. Synergistic binding of nucleotides and inhibitors to cAMP-dependent protein kinase examined by acrylodan fluorescence spectroscopy. *J Biol Chem* **272**, 1507-13 (1997).
20. Wang, Y. et al. Globally correlated conformational entropy underlies positive and negative cooperativity in a kinase's enzymatic cycle. *Nat Commun* **10**, 799 (2019).

21.   Olivieri, C. et al. Defective internal allosteric network imparts dysfunctional
ATP/substrate-binding cooperativity in oncogenic chimera of protein kinase A. *Commun*
Biol **4**, 321 (2021).

22.   Walker, C. et al. Cushing's syndrome driver mutation disrupts protein kinase A allosteric
network, altering both regulation and substrate specificity. *Sci Adv* **5**, eaaw9298 (2019).

23.   Walker, C. et al. Is Disrupted Nucleotide-Substrate Cooperativity a Common Trait for
Cushing's Syndrome Driving Mutations of Protein Kinase A? *J Mol Biol* **433**, 167123
(2021).

24.   Camilloni, C., Cavalli, A. & Vendruscolo, M. Assessment of the use of NMR chemical
shifts as replica-averaged structural restraints in molecular dynamics simulations to
characterize the dynamics of proteins. *J Phys Chem B* **117**, 1838-43 (2013).

25.   Camilloni, C. & Vendruscolo, M. Statistical mechanics of the denatured state of a protein
using replica-averaged metadynamics. *J Am Chem Soc* **136**, 8982-91 (2014).

26.   Husic, B.E. & Pande, V.S. Markov State Models: From an Art to a Science. *J Am Chem*
Soc **140**, 2386-2396 (2018).

27.   Olivieri, C. et al. ATP-competitive inhibitors modulate the substrate binding cooperativity
of a kinase by altering its conformational entropy. *Sci Adv* **8**, eabo0696 (2022).

28.   Ruan, Z. & Kannan, N. Altered conformational landscape and dimerization dependency
underpins the activation of EGFR by alpha C-beta 4 loop insertion mutations.
Proceedings of the National Academy of Sciences of the United States of America **115**,
Eb162-Eb171 (2018).

29.   Kannan, N., Neuwald, A.F. & Taylor, S.S. Analogous regulatory sites within the αC-β4
loop regions of ZAP-70 tyrosine kinase and AGC kinases. *Biochimica Et Biophysica*
Acta-Proteins and Proteomics **1784**, 27-32 (2008).

30.   Zhao, S. et al. Conformational Landscapes of HER2 Exon 20 Insertions Explain Their
Sensitivity to Kinase Inhibitors in Lung Adenocarcinoma. *J Thorac Oncol* **15**, 962-972
(2020).

31.   Robustelli, P., Kohlhoff, K., Cavalli, A. & Vendruscolo, M. Using NMR chemical shifts as
structural restraints in molecular dynamics simulations of proteins. *Structure* **18**, 923-33
(2010).

32.   Qi, G. et al. Enhancing Biomolecular Simulations with Hybrid Potentials Incorporating
NMR Data. *J Chem Theory Comput* **18**, 7733-7750 (2022).

33.   Piana, S. & Laio, A. A bias-exchange approach to protein folding. *J Phys Chem B* **111**,
4553-9 (2007).

34.   Shen, Y. & Bax, A. SPARTA+: a modest improvement in empirical NMR chemical shift
prediction by means of an artificial neural network. *J Biomol NMR* **48**, 13-22 (2010).

35.   Masterson, L.R. et al. Dynamically committed, uncommitted, and quenched states
encoded in protein kinase A revealed by NMR spectroscopy. *Proc Natl Acad Sci U S A*
108, 6969-74 (2011).

36.   Masterson, L.R. et al. Dynamics connect substrate recognition to catalysis in protein
kinase A. *Nat Chem Biol* **6**, 821-8 (2010).

37.   Li, G.C., Srivastava, A.K., Kim, J., Taylor, S.S. & Veglia, G. Mapping the Hydrogen Bond
Networks in the Catalytic Subunit of Protein Kinase A Using H/D Fractionation Factors.
Biochemistry **54**, 4042-4049 (2015).

38.   Kohlhoff, K.J. et al. Cloud-based simulations on Google Exacycle reveal ligand
modulation of GPCR activation pathways. *Nat Chem* **6**, 15-21 (2014).

39.   Plattner, N., Doerr, S., De Fabritiis, G. & Noe, F. Complete protein-protein association
kinetics in atomic detail revealed by molecular dynamics simulations and Markov
modelling. *Nature Chemistry* **9**, 1005-1011 (2017).

40.   Kim, J. et al. A dynamic hydrophobic core orchestrates allostery in protein kinases. *Sci*
Adv **3**, e1600663 (2017).

41. Masterson, L.R., Mascioni, A., Traaseth, N.J., Taylor, S.S. & Veglia, G. Allosteric
cooperativity in protein kinase A. *Proc Natl Acad Sci U S A* **105**, 506-11 (2008).
42. Shukla, D., Meng, Y., Roux, B. & Pande, V.S. Activation pathway of Src kinase reveals
intermediate states as targets for drug design. *Nat Commun* **5**, 3397 (2014).
43. Han, B., Liu, Y.F., Ginzinger, S.W. & Wishart, D.S. SHIFTX2: significantly improved
protein chemical shift prediction. *Journal of Biomolecular Nmr* **50**, 43-57 (2011).
44. McClendon, C.L., Hua, L., Barreiro, A. & Jacobson, M.P. Comparing Conformational
Ensembles Using the Kullback-Leibler Divergence Expansion. *J Chem Theory Comput*
8, 2115-2126 (2012).
45. Tsigelny, I. et al. 600 ps molecular dynamics reveals stable substructures and flexible
hinge points in cAMP dependent protein kinase. *Biopolymers* **50**, 513-24 (1999).
46. Lu, B., Wong, C.F. & McCammon, J.A. Release of ADP from the catalytic subunit of
protein kinase A: a molecular dynamics simulation study. *Protein Sci* **14**, 159-68 (2005).
47. Cook, P.F., Neville, M.E., Jr., Vrana, K.E., Hartl, F.T. & Roskoski, R., Jr. Adenosine
cyclic 3',5'-monophosphate dependent protein kinase: kinetic mechanism for the bovine
skeletal muscle catalytic subunit. *Biochemistry* **21**, 5794-9 (1982).
48. Kemp, B.E., Graves, D.J., Benjamini, E. & Krebs, E.G. Role of Multiple Basic Residues
in Determining Substrate-Specificity of Cyclic Amp-Dependent Protein-Kinase. *Journal of*
Biological Chemistry **252**, 4888-4894 (1977).
49. Wiseman, T., Williston, S., Brandts, J.F. & Lin, L.-N. Rapid measurement of binding
constants and heats of binding using a new titration calorimeter. *Analytical Biochemistry*
179, 131-137 (1989).
50. Srivastava, A.K. et al. Synchronous Opening and Closing Motions Are Essential for
cAMP-Dependent Protein Kinase A Signaling. *Structure* **22**, 1735-43 (2014).
51. Cembran, A., Kim, J., Gao, J. & Veglia, G. NMR mapping of protein conformational
landscapes using coordinated behavior of chemical shifts upon ligand binding. *Phys*
Chem Chem Phys **16**, 6508-18 (2014).
52. Akimoto, M. et al. Signaling through dynamic linkers as revealed by PKA. *Proc Natl Acad*
Sci U S A **110**, 14231-6 (2013).
53. Selvaratnam, R., Chowdhury, S., VanSchouwen, B. & Melacini, G. Mapping allostery
through the covariance analysis of NMR chemical shifts. *Proc Natl Acad Sci U S A* **108**,
6133-8 (2011).
54. Veron, M., Radzio-Andzelm, E., Tsigelny, I., Ten Eyck, L.F. & Taylor, S.S. A conserved
helix motif complements the protein kinase core. *Proc Natl Acad Sci U S A* **90**, 10618-22
(1993).
55. McClendon, C.L., Kornev, A.P., Gilson, M.K. & Taylor, S.S. Dynamic architecture of a
protein kinase. *Proc Natl Acad Sci U S A* **111**, E4623-31 (2014).
56. Foda, Z.H., Shan, Y., Kim, E.T., Shaw, D.E. & Seeliger, M.A. A dynamically coupled
allosteric network underlies binding cooperativity in Src kinase. *Nat Commun* **6**, 5939
(2015).
57. Iverson, D.B., Xiao, Y., Jones, D.N., Eisenmesser, E.Z. & Ahn, N.G. Activation Loop
Dynamics Are Coupled to Core Motions in Extracellular Signal-Regulated Kinase-2.
Biochemistry **59**, 2698-2706 (2020).
58. Beuschlein, F. et al. Constitutive activation of PKA catalytic subunit in adrenal Cushing's
syndrome. *N Engl J Med* **370**, 1019-28 (2014).
59. Calebiro, D., Bathon, K. & Weigand, I. Mechanisms of Aberrant PKA Activation by Cα
Subunit Mutations. *Horm Metab Res* **49**, 307-314 (2017).
60. Calebiro, D. et al. PKA catalytic subunit mutations in adrenocortical Cushing's adenoma
impair association with the regulatory subunit. *Nature Communications* **5**, 5680 (2014).

61.   Di Dalmazi, G. et al. Novel somatic mutations in the catalytic subunit of the protein
kinase A as a cause of adrenal Cushing's syndrome: a European multicentric study. *J*
Clin Endocrinol Metab **99**, E2093-100 (2014).
62.   Lubner, J.M. et al. Cushing's syndrome mutant PKA(L)(205R) exhibits altered substrate
specificity. *FEBS Lett* **591**, 459-467 (2017).
63.   Kannan, N. & Neuwald, A.F. Did protein kinase regulatory mechanisms evolve through
elaboration of a simple structural component? *Journal of Molecular Biology* **351**, 956-972
(2005).
64.   Chen, H. et al. A molecular brake in the kinase hinge region regulates the activity of
receptor tyrosine kinases. *Mol Cell* **27**, 717-30 (2007).
65.   Klein, T. et al. Structural and dynamic insights into the energetics of activation loop
rearrangement in FGFR1 kinase. *Nature Communications* **6**(2015).
66.   Fan, Y.X. et al. Mutational activation of ErbB2 reveals a new protein kinase
autoinhibition mechanism. *J Biol Chem* **283**, 1588-1596 (2008).
67.   McSkimming, D.I. et al. ProKinO: a unified resource for mining the cancer kinome. *Hum*
Mutat **36**, 175-86 (2015).
68.   Yeung, W., Ruan, Z. & Kannan, N. Emerging roles of the alphaC-beta4 loop in protein
kinase structure, function, evolution, and disease. *IUBMB Life* **72**, 1189-1202 (2020).
69.   Cembran, A. et al. Conformational equilibrium of N-myristoylated cAMP-dependent
protein kinase A by molecular dynamics simulations. *Biochemistry* **51**, 10186-96 (2012).
70.   Jorgensen, W.L., Chandrasekhar, J., Madura, J.D., Impey, R.W. & Klein, M.L.
Comparison of Simple Potential Functions for Simulating Liquid Water. *Journal of*
Chemical Physics **79**, 926-935 (1983).
71.   Hess, B., Bekker, H., Berendsen, H.J.C. & Fraaije, J.G.E.M. LINCS: A linear constraint
solver for molecular simulations. *Journal of Computational Chemistry* **18**, 1463-1472
(1997).
72.   Darden, T., York, D. & Pedersen, L. Particle Mesh Ewald - an N.Log(N) Method for
Ewald Sums in Large Systems. *Journal of Chemical Physics* **98**, 10089-10092 (1993).
73.   Hess, B., Kutzner, C., van der Spoel, D. & Lindahl, E. GROMACS 4: Algorithms for
Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J Chem Theory*
Comput **4**, 435-47 (2008).
74.   Best, R.B. et al. Optimization of the additive CHARMM all-atom protein force field
targeting improved sampling of the backbone phi, psi and side-chain chi(1) and chi(2)
dihedral angles. *J Chem Theory Comput* **8**, 3257-3273 (2012).
75.   DeLano, W. The PyMOL molecular graphics system, version 0.99 rc6. *DeLano*
Scientific: San Carlos, CA (2002).
76.   Parrinello, M. & Rahman, A. Crystal-Structure and Pair Potentials - a Molecular-
Dynamics Study. *Physical Review Letters* **45**, 1196-1199 (1980).
77.   Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling. *J*
Chem Phys **126**, 014101 (2007).
78.   Kohlhoff, K.J., Robustelli, P., Cavalli, A., Salvatella, X. & Vendruscolo, M. Fast and
accurate predictions of protein NMR chemical shifts from interatomic distances. *J Am*
Chem Soc **131**, 13894-5 (2009).
79.   Bonomi, M. et al. PLUMED: A portable plugin for free-energy calculations with molecular
dynamics. *Computer Physics Communications* **180**, 1961-1972 (2009).
80.   Biarnes, X., Pietrucci, F., Marinelli, F. & Laio, A. METAGUI. A VMD interface for
analyzing metadynamics and molecular dynamics simulations. *Computer Physics*
Communications **183**, 203-211 (2012).
81.   McGibbon, R.T. et al. MDTraj: A Modern Open Library for the Analysis of Molecular
Dynamics Trajectories. *Biophys J* **109**, 1528-32 (2015).

82. Schwantes, C.R. & Pande, V.S. Improvements in Markov State Model Construction
Reveal Many Non-Native Interactions in the Folding of NTL9. *J Chem Theory Comput* **9**,
2000-2009 (2013).
83. Yonemoto, W., Garrod, S.M., Bell, S.M. & Taylor, S.S. Identification of phosphorylation
sites in the recombinant catalytic subunit of cAMP-dependent protein kinase. *J Biol*
Chem **268**, 18626-32 (1993).
84. Manu, V.S., Olivieri, C., Pavuluri, K. & Veglia, G. Design and applications of water
irradiation devoid RF pulses for ultra-high field biomolecular NMR spectroscopy.
Physical Chemistry Chemical Physics **24**, 18477-18481 (2022).
85. Delaglio, F. et al. NMRPipe: a multidimensional spectral processing system based on
UNIX pipes. *J Biomol NMR* **6**, 277-93 (1995).
86. Lee, W., Tonelli, M. & Markley, J.L. NMRFAM-SPARKY: enhanced software for
biomolecular NMR spectroscopy. *Bioinformatics* **31**, 1325-1327 (2015).
87. Manthey, I. et al. POKY software tools encapsulating assignment strategies for solution
and solid-state protein NMR data. *J Struct Biol X* **6**, 100073 (2022).
88. Williamson, M.P. Using chemical shift perturbation to characterise ligand binding. *Prog*
Nucl Magn Reson Spectrosc **73**, 1-16 (2013).

A	**B**	**C**	**D**
Back-far	Back-close	Side-close	Rgss

Figure 2 – figure supplement 1. **Illustration of the collective variables (CVs) used in the RAM simulations.** (**A**) Backbone ψ angles of loops not in contact with ATP (Back-far). The Cα atoms are depicted as blue spheres. (**B**) Backbone ψ angles of loops in contact with ATP (Back-close). The Cα atoms are colored in magenta. (**C**) Side chains χ_1 angles of loops in contact with ATP (Side-close). The side chains are represented in sticks colored in magenta. (**D**) The radius of gyration is calculated over the rigid part of the protein (Rgss), where the residues involved are colored in cyan.

Figure 2 – figure supplement 2. **Distribution of the Root-Mean-Square-Error (RMSE) of the chemical shifts for the different simulation schemes**. (**A**) RMSE of CS for apo PKA-C calculated from standard MD (left), REX (middle), and RAM (right). (**B**) RMSE of CS for PKA-C/ATP calculated from standard MD (left), REX (middle), and RAM (right). (**C**) RMSE of CS for PKA-C/ATP/PKI$_{5-24}$ from standard MD (left), REX (middle), and RAM (right). Color codes for different backbone atoms (C, Cα, CO, H, and N) are indicated in the left figures.

Figure 2 – figure supplement 3. **Replica-averaged metadynamics (RAM) simulations ex-plore a larger conformational space than standard MD and replica exchange (REX) simu-lations**. (**A**) Comparison of conformational space (CV1 vs. CV2) sampled by RAM Replica 1, standard MD, and REX Replica 1 for PKA-C. (**B**) Comparison of conformational space (CV3 vs. CV2) sampled by RAM Replica 1, standard MD, and REX Replica 1 for apo PKA-C.

Figure 2 – figure supplement 4. Accumulative deposition of history-dependent biases along the first three CVs for the RAM simulations of the apo PKA-C. The accumulative biases converged after ~300 ns in the three CVs.

Figure 3 – figure supplement 1. **R spine and shell residues selected for two time-lagged independent components (tICA) and Markov State Model (MSM) analysis.** (**A**) Atom motions of key residues that define tIC1 of apo PKA-C colored according to the superposition deviations. Backbone atoms of Val104 show the largest change in tIC1. (**B**) Atom motions of key residues that define tIC2 of apo PKA-C colored according to the superposition deviations. Backbone atoms of Phe185 and Val104 show the largest change in tIC2.

Figure 3 – figure supplement 2. Structural features of αC-out transition (ES1) in various inactive kinases. (**A**) Crystal structures of PKA-C in active and inactive states highlight the αB-αC loop alterations. Crystal structures of PKA-C in the active (1ATP) and inactive conformations (3AG9, 1SZM, 4DFY), highlighting the electrostatic interactions between K72 in β3 and E91 in the αC. (**B**) Disruption of the K72 - E91salt bridge in the inactive structures of Abl (1OPJ), Src (1FMK), and CDK2 (4EK3). (**C**) The αC helix orientation of active PKA-C compared to the orientation in inactive kinases. (**D**) Structural transition from the GS state to ES1 state characterized by the outward movement of the αC helix (i.e., kinase inactivation).

1006

1007

Figure 3 – figure supplement 3. **Distinct hydrophobic packing for residues around the αC-**
β4 loop in the GS and ES states of apo PKA-C. (**A**) Projections of randomly selected confor-
mations for the GS (blue) and ES (magenta) onto the conformational landscape of apo PKA-C.
Snapshots with tIC1 < 1.2 were clustered to separate the ES and GS, whereas those conformers
with tIC1 > 0.2 were clustered as GS. (**B,C**) Close up of the hydrophobic packing in the ES and
GS (**C**) states, highlighting Leu103, Val104, Ile150, Leu172, and Ile180 that show slow exchange
in the CPMG experiments.

1015

Figure 5 - figure supplement 1. **Distribution of predicted and experimental ^{13}C CS of se-**
lected methyl groups. **(A-C)** ES (magenta) and GS (blue) of apo PKA-C for Leu103-Cδ2 (**A**),
Leu172-Cδ1 (**B**), and Ile180-Cδ1 (**C**). The experimental CSs are shown as dotted lines for GS
(black) and ES (red).

1023 **Figure 6 - figure supplement 1. Time series of the distance between F102 and R308 for**
1024 **PKA-C^{WT} (black) and the PKA-C^{F100A} mutant (magenta)**. The π-cation interactions between the
1025 aromatic ring of F102 and the guanidine group of R308 are more persistent in the WT enzyme
1026 than in the F100A mutant.

1027

1028

1029

1030

1031   **Figure 9 - figure supplement 1. NMR fingerprints of PKA-C^{F100A}.** (**A**) [^{1}H,^{15}N]-WADE-TROSY
1032   spectra of apo, ADP-bound, ATPγN-bound, and ATPγN/PKI$_{5-24}$ bound PKA-C^{F100A}. (**B**) Changes
1033   in the chemical shift perturbation (CSP) between PKA-C^{WT} and PKA-C^{F100A} bound to ATPγN. (**C**)
1034   Changes in CSP ($\Delta\delta_{WT}$ - $\Delta\delta_{F100A}$) upon binding ATPγN and PKI$_{5-24}$.

1035

1036

1037

1038 **Figure 9 - figure supplement 2. CONCISE plot showing the shifts of the probability distri-**
1039 **bution of the amide resonances as a function of nucleotides and substrate binding.** The
1040 per-residue chemical shift information is averaged into the average principal component (PC)
1041 score indicative of the position of each conformational state of the kinase along the open-to-
1042 closed equilibrium.

1043

1044

1045

Figure S10 - figure supplement 2. Intermolecular allosteric network of F100A mapped using CHESCA and community CHESCA. (**A**) CHESCA matrix obtained from the amide chemical shift trajectories of PKA-C^{F100A} in the apo, ADP-bound, ATPγN-bound, and ATPγN/PKI$_{5-24}$-bound states. Only correlations with $R_{ij} > 0.98$ are displayed. (**B**) Plot of the correlation scores vs. residue calculated for PKA-C^{WT} (blue) and PKA-C^{F100A} (black). (**C**) Community CHESCA analysis of PKA-C^{F100A}. Only correlations with $R_{A,B} > 0.98$ are shown. (**D**) Spider plots indicating the correlated structural communities of PKA-C^{F100A} and PKA-C^{WT} plotted on their corresponding structures. The size of each node is independent of the number of residues it encompasses, and the weight of each line indicates the strength of coupling between the individual communities.

1055

1056

1057

1058 **Figure 2 – supplementary Table 1**. ΔG (kcal/mol) and relative population of the ground state
1059 and the first 6 excited states in different forms of PKA-C obtained from the RAM simulations.

	GS	ES1	ES2	ES3	ES4	ES5	ES6
Apo	0 (58.0%)	0.38 (30.8%)	1.22 (7.6%)	2.10 (1.8.%)	2.28 (1.3%)	2.85 (0.5%)	8.16* (<1e-4)
Binary	0 (99.4%)	3.11 (0.6%)	5.80 (<1e-4)	6.66 (<1e-4)	7.25 (<1e-4)	8.27 (<1e-4)	8.47 (<1e-4)
Ternary	0 (100.0%)	4.85 (<1e-4)	5.68 (<1e-4)	6.92 (<1e-4)	7.04 (<1e-4)	7.58 (<1e-4)	7.87 (<1e-4)

1060 * Numbers in red refer to populations of excited states below 0.5%.

1061

1062 **Supplementary Table 2. Kinetic parameters of Kemptide phosphorylation by PKA-C^{WT} and**
1063 **PKA-C^{F100A} obtained from coupled assays.** The K_M and V_{max} values were obtained from a non-
1064 linear least squares analysis of the concentration-dependent initial phosphorylation rates. Errors
1065 in the k_{cat}/K_M ratios were propagated from the individual errors in K_M and k_{cat}.

	PKA-C^{WT}	PKA-C^{F100A}
V_{max}	0.322 ± 0.005	0.379 ± 0.009
K_M	30 ± 1	42 ±3
k_{cat}	15 ± 1	17 ± 1
k_{cat}/K_M	0.50 ± 0.04	0.41 ± 0.08

1066

1067

1068 **Supplementary Table 3. Changes in enthalpy, entropy, free energy, and dissociation con-**
1069 **stants for nucleotide binding to PKA-C^{WT} and PKA-C^{F100A}.** All errors were calculated from trip-
1070 licate measurements. Values for PKA-C^{WT} are taken from Walker *et al.* [22]

	K_d (μM)	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)	σ
PKA-C^{WT}	83 ± 8	-5.61 ± 0.06	-3.6 ± 0.1	-2.0 ± 0.1	N/A
PKA-C^{F100A}	73 ± 2	-5.7 ± 0.2	-21 ± 5	7 ± 3	N/A

1071

1072

1073 **Supplementary Table 4. Changes in enthalpy, entropy, free energy, and dissociation con-**
1074 **stants for PKI$_{5-24}$ binding to apo and ATPγN - saturated PKA-C^{WT} and PKA-C^{F100A}.** All errors
1075 were derived from triplicate measurements. The error for the cooperativity coefficient (σ) was
1076 propagated from the errors in K_d. Values for PKA-C^{WT} were originally published in Walker *et al.* [22].

PKI$_{5-24}$ to apo kinase					
	K_d (μM)	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)	σ
PKA-C^{WT}	17 ± 2	-6.57 ± 0.08	-10.8 ± 0.5	4.2 ± 0.5	N/A
PKA-C^{F100A}	5 ± 1	-7.7 ± 0.1	-17 ± 5	7 ± 3	N/A

1077

PKI$_{5-24}$ to ATPγN-saturated kinase					
	K_d (μM)	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)	σ
PKA-C^{WT}	0.16 ± 0.02	-9.33 ± 0.07	-13.9 ± 0.5	4.6 ± 0.4	106 ± 18
PKA-C^{F100A}	2 ± 1	-7.9 ± 0.3	-17 ± 1	9 ± 1	3 ± 1

1078

1079