The αC-β4 loop controls the allosteric cooperativity between nucleotide and substrate in the catalytic subunit of protein kinase A

Cristina Olivieri[1,†,‡], Yingjie Wang[1,2,†,§], Caitlin Walker[1], Manu V. Subrahmanian[1], Kim N. Ha[3],

David A. Bernlohr[1], Jiali Gao[2], Carlo Camilloni[4,◊], Michele Vendruscolo[4], Susan S. Taylor[5,6], and

Gianluigi Veglia[1,2]*

[1]Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, MN 55455, USA

[2]Department of Chemistry and Supercomputing Institute, University of Minnesota, MN 55455, USA

[3]Departmenf of Chemistry and Biochemistry, St. Catherine University, MN 55105, USA

[4]Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, UK.

[5]Department of Pharmacology, University of California at San Diego, CA 92093, USA.

[6]Department of Chemistry and Biochemistry, University of California at San Diego, CA 92093, USA.

[†]These authors contributed equally to this work.

Present address:

[‡] Department of Chemistry, University of Milan, 20133, Milan, Italy

[§] Institute of Systems and Physical Biology, Shenzhen Bay Laboratory, Shenzhen, 518055, China

[◊] Department of Bioscience, University of Milano, 20133, Milano, Italy

*Corresponding Author

Gianluigi Veglia
Department of Chemistry and Department of Biochemistry, Molecular Biology, and Biophysics,
321 Church Street SE
Minneapolis, MN 55455
Telephone: (612) 625-0758
Fax: (612) 625-5780
e-mail: vegli001@umn.edu

ABSTRACT

Allosteric cooperativity between ATP and substrates is a prominent characteristic of the cAMP-dependent catalytic (C) subunit of protein kinase A (PKA). Not only this long-range synergistic action is involved in substrate recognition and fidelity, but it is likely to regulate PKA association with regulatory subunits and other binding partners. To date, a complete understanding of the molecular determinants for this intramolecular mechanism is still lacking.

Here, we used an integrated NMR-restrained molecular dynamics simulations and a Markov Model to characterize the free energy landscape and conformational transitions of the catalytic subunit of protein kinase A (PKA-C). We found that the apo-enzyme populates a broad free energy basin featuring a conformational ensemble of the active state of PKA-C (ground state) and other basins with lower populations (excited states). The first excited state corresponds to a previously characterized inactive state of PKA-C with the αC helix swinging outward. The second excited state displays a disrupted hydrophobic packing around the regulatory (R) spine, with a flipped configuration of the F100 and F102 residues at the tip of the αC-β4 loop. To experimentally validate the second excited state, we mutated F100 into alanine and used NMR spectroscopy to characterize the binding thermodynamics and structural response of ATP and a prototypical peptide substrate. While the activity of PKA-C^{F100A} toward a prototypical peptide substrate is unaltered and the enzyme retains its affinity for ATP and substrate, this mutation rearranges the αC-β4 loop conformation interrupting the allosteric coupling between nucleotide and substrate. The highly conserved αC-β4 loop emerges as a pivotal element able to modulate the synergistic binding between nucleotide and substrate and may affect PKA signalosome. These results may explain how insertion mutations within this motif affect drug sensitivity in other homologous kinases.

Keywords:
Protein Kinases, cAMP-dependent protein kinase A, allosteric mutations, binding cooperativity.

INTRODUCTION

Eukaryotic protein kinases (EPKs) are plastic enzymes of paramount importance to signaling

processes, catalyzing phosphoryl transfer reactions, or acting as scaffolds for other enzymes

and/or binding partners [1]. Of all kinases, the catalytic subunit of PKA (PKA-C) was the first to be

structurally characterized by X-ray crystallography.[2,3] In its inhibited state, PKA-C forms a heter-

otetrametric complex comprising two catalytic (C) and two regulatory (R) subunits.[4] The canonical

activation mechanism of PKA involves binding two cAMP molecules and disassembling the holo-

enzyme, which unleashes active PKA-C monomers that target signaling partners.[5] In 1997, how-

ever, Scott and coworkers suggested that the holoenzyme does not disassemble under physio-

logical conditions; rather, the intact holo-enzyme forms *signaling islands* localized by A-kinase

anchoring proteins (AKAPs) in the proximity of substrates.[6]

X-ray crystallography studies revealed that PKA-C is a bilobal enzyme, with a dynamic N-

terminal lobe comprising β-sheets and the αC helix and a more rigid C-terminal lobe, with mostly

α-helices (**Figure 1**).[2,3,7] The N-lobe harbors the nucleotide-binding site, whereas the substrate

binding cleft lays at the interface between the N-lobe and C-lobe. The three-dimensional structure

of the enzyme features a highly conserved hydrophobic core decorated by catalytically important

motifs, *i.e.*, the Gly-rich loop, DFG-loop, activation loop, positioning loop, and magnesium and

peptide positioning loops.[8] In the catalytically active state, these motifs are all poised for phos-

phoryl transfer. However, the conformation of these motifs is necessary but not sufficient to define

an active kinase. More recent studies revealed a critical role of the hydrophobic core, which is

crossed by a catalytic (C) spine, a regulatory (R) spine, and shell residues (**Figure 1**).[9,10] The C

spine comprises an array of hydrophobic residues and is assembled upon binding ATP, whereas

the R spine is engaged when the activation loop is phosphorylated, which contributes to the po-

sitioning of the αC-helix.[11]

A distinct property of PKA-C is the binding cooperativity between ATP and substrate.[12] During

the catalytic cycle, the kinase recognizes and binds substrates with positive binding cooperativity

between ATP and unphosphorylated substrates, whereas a negative binding cooperativity be-

tween ADP and phosphorylated substrate characterizes the exit complex.[13] The biological im-

portance of binding cooperativity has been emphasized by our recent studies on disease-driven

mutations of PKA-C,[14-16] which all feature disrupted cooperativity between nucleotides and protein

kinase inhibitor (PKI) or typical substrates.[14-16] Since the recognition sequence of the substrate is

highly homologous to the regulatory subunits,[4] a loss of cooperativity of binding may affect not

only substrate binding fidelity but also the regulation by the R subunit. However, the molecular

determinants for the binding cooperativity between ATP and substrate and its role in PKA signal-

osome remain elusive to date.

Figure 1. Structural and catalytic motifs of PKA-C. (**A**) Surface representation of the X-ray
structure of PKA-C bound to the endogenous inhibitor, PKI (PDB: 4WB5). (**B**) Hydrophobic or-
ganization of the PKA-C core, with the R-spine (gold), C-spine (blue), shell residues (cyan), and
the αC-β4 loop (hot pink) that locks into αE helix.

Here, we combined NMR-restrained replica-averaged metadynamics (RAM)[17,18] and Markov

State Model (MSM)[19] to define the conformational landscape and the corresponding dynamics of

PKA-C. We found that both the apo kinase and the nucleotide-bound kinase occupy three distinct

basins: (*i*) a most populated ground state with constitutively active conformations competent for

catalysis, (*ii*) a first high free energy basin representative of typical inactive states with a dislodged

configuration of the αC helix, and (*iii*) a second high free energy basin with a disrupted hydropho-

bic array of residues at the core of the enzyme. Notably, the equilibrium between the most popu-

lated ground state and the other low-populated states agrees with previous NMR relaxation dis-

persion and CEST measurements.[20] To compound the existence of the second inactive basin, we

mutated F100 at the tip of the αC-β4 loop into Ala (PKA-C^{F100A}) and characterized its binding

thermodynamics and ligand binding response by isothermal titration calorimetry and NMR spec-

troscopy. We found that PKA-C^{F100A} phosphorylates canonical peptides and preserves the bind-

ing affinity for both nucleotide and substrate. However, PKA-C^{F100A} lacks binding cooperativity

between nucleotide and substrate. NMR revealed that perturbing the hydrophobic packing around

the αC-β4 loop interrupts the allosteric communication between the two lobes of the enzyme.

These results further support the pivotal role of the αC-β4 loop in kinases function and may ex-

plain why single-site mutations or insertion mutations in the homologous kinase that stabilize this

motif result in oncogenes and confer different drug sensitivity.[21]

RESULTS

The free energy landscape of PKA-C charted by NMR-restrained replica-averaged

metadynamics (RAM) simulations. To characterize the experimentally accessible conforma-

tional landscape of PKA-C in the µs-to-ms range, we performed NMR-restrained metadynamics

simulations within the RAM framework.[17,18] We simulated the apo, binary (PKA-C/ATP), and ter-

nary (PKA-C/ATP/PKI$_{5-24}$) forms of PKA-C, using four replicas restrained using backbone chemi-

cal shifts (CS) to ensure a close agreement with conformational space explored by the kinase

under the conditions used in this work. The enhanced sampling was achieved through bias-ex-

change metadynamics along different collective variables (CVs) to boost the conformational plas-

ticity of the enzyme (**Figure 2 - figure supplement 1**).[22] Back-calculation of the CS using the

Sparta+ software[23] shows that the CS restraints improved the agreement between back-calcu-

lated and experimental CS. Specifically, we obtained an overall improvement of ~0.2 ppm on the

amide N atom and ~0.1 ppm on the remainder backbone atoms (**Figure 2 - figure supplement**

2). The bias-exchange metadynamics allowed for each replica to span a broader conformational

space relative to classical simulations (**Figure 2 - figure supplement 3**). The deposition of

Gaussian biases required by the metadynamics approach converged after 300 ns, where the

fluctuations along the first 3 CVs were less than 1 kcal/mol (**Figure 2A, Figure 2 - figure supple-**

ment 4). The converged bias revealed that the *apo* kinase accesses multiple minima along each

CV, and the conformational heterogeneity is significantly reduced upon ligand binding, especially

in the ternary form (**Figure 2A**). The full free energy landscape was then reconstructed by sam-

pling an extra 100 ns production phase with reduced biases along each CV. The free energy

landscape shows how the population of the conformers is modulated by ligands within the NMR

detection limit of sparsely populated states (~ 0.5% or $\Delta G < 3.2$ kcal/mol).

Figure 2. Free energy landscape (FEL) of PKA-C in various ligated forms obtained from replica-averaged metadynamics (RAM) simulations. (**A**) Convergence of the bias deposition along the first three collective variables (CVs). The free energy (expressed in kcal/mol) of the different CVs were averaged over the last 100 ns of RAM simulations. The standard deviations are reported as red error bars. (**B-D**) FEL along the first two principal components (PC1 and PC2) of PKA-C in the apo, ATP-bound, and ATP and the model substrate PKI bound forms. PC1 and PC2 are projected from the first three CVs. The vertices represent conformational states. In the apo form, multiple states have comparable free energy with ΔG < 5 kcal/mol, whereas in the binary form, fewer states have ΔG < 5 kcal/mol, whereas for the ternary form only a major ground state is populated.

According to these simulations, apo PKA-C populates preferentially a ground state and five readily accessible low-populated excited states (**Figure 2B**, **Figure 2 – supplementary table 1**). The nucleotide-bound PKA-C (binary form) features a similar ground state and a broad higher energy basin (**Figure 2C**, **Figure 2 – supplementary table 1**). Finally, the ternary complex occupies a narrow dominant ground state (**Figure 2D**, **Figure 2 – supplementary table 1**). This free energy landscape obtained from the RAM simulations is consistent with the qualitative picture

previously inferred from our NMR spin relaxation experiments,[24] while providing a detailed struc-

tural characterization of the excited states.

MSM reveals the conformational transitions of PKA-C from ground to high free energy

(excited) states. To explore the conformational transitions of the kinase upon ligand binding, we

performed additional unbiased sampling to build a Markov State Model (MSM). MSMs are com-

monly used to describe the dynamic transitions of macromolecules in terms of (1) the probabilities

of occupation of a specific set of states and (2) the transition probabilities of moving between

these states. In practice, a MSM is typically created by combining thousands of short unbiased

simulations. [25,26] Following this strategy, we performed several short simulations (10 – 20 ns)

using thousands of the low free energy conformations ($\Delta G < 3.2$ kcal/mol) chosen randomly from

the three forms of PKA-C as starting structures. The conformational ensembles were clustered

into microstates and seeded to start a second round of adaptive sampling (see Methods). The

iterative process was repeated three times to assure convergence and yielded a total of 100 μs

trajectories for both the apo and binary forms, whereas for the less dynamic ternary complex, we

collected trajectories of 60 μs. Once we reached a sufficient sampling, we built a MSM including

L95, V104, L106, M118, M120, Y164, and F185 to investigate the dynamic transitions of the hy-

drophobic R spine and shell residues (**Figure 3 – figure supplement 1**). These residues are ideal

reporters of the dynamic processes governing the activation/deactivation of the kinase.[27] To com-

pare the free energy landscape of different complexes, we first projected the conformational en-

sembles of three forms and existing crystal structures along the first two time-lagged independent

components (tICs) of the *apo* form, which were obtained by a time-lagged independent compo-

nent analysis (tICA) analysis (see Methods). These tICs represent the directions of the slowest

motion of the kinase and visualize the conformational transitions of V104, L95, and F185 (**Figure**

3 – figure supplement 1). Apo PKA-C accesses three major basins. The broadest basin repre-

sents the ground state (GS) and represents the conformations of the kinase captured by essen-

tially all crystal structures (**Figure 3A**). Additionally, there are two distinct excited states: the first

excited state (ES1) features a disrupted hydrophobic packing of L95, V104, and F185. This con-

formation features an inactive state with an orientation of the αC-helix typical of the inhibited

states as found for the PKA-C bound to regulatory subunits RIα and RIIβ. The second excited

state (ES2), to our knowledge, was never captured in crystal structures. The ES2 state displays

a flipped configuration of the V104 side chain and a rearrangement of the αC-β4 loop, with the

dihedral angles of F100 and F102 adopting a gauche$^+$ configuration. This orientation of the F100

and F102 aromatic rings breaks the hydrophobic packing of the αC-β4 loop with the C-lobe, caus-

ing steric contacts between F100 and V104 (**Figure 3A**). In contrast, the active GS ensemble

features a *trans* configuration of the F100 and F102 side chains that stabilizes the hydrophobic

interactions with αE- and αJ-helices in the C-lobe (**Figure 3A**). Upon binding ATP, the confor-

mational space span by the kinase becomes narrower, and the conformers populate mostly the

GS, with a small fraction in the ES1 state (**Figure 3B)**. This is consistent with the role of the

nucleotide as an allosteric effector, enhancing the affinity of the enzyme for the substrate. In the

ternary form (ATP and PKI-bound), PKA-C populates only the GS consistent with the competent

conformation observed in the first ternary complex structure. In this case, the αC-β4 loop of the

enzyme is locked in a well-defined configuration, as shown in **Figure 3A**.

Figure 3. The apo, ATP-bound, and ATP/PKI-bound PKA-C reveal distinct free energy sur-face (FES) and dynamics, as determined by a Markov State Model (MSM). (**A**) Free energy landscape projected along the first two time-lagged independent components (tICs) of the apo PKA-C, the projections of known crystal structures, and characteristic features of GS, ES1, and ES2. The transition from GS to ES1 highlights the changes around the αB-αC loop, where the salt bridges between K72-E91 and H87-T197 and the PIF pocket (V80-I85-F347) are all disrupted. The transition from GS to ES2 highlights the rearrangement around the αC-β4 loop, with distinct local hydrophobic packing. (**B** and **C**) FES projected along the first two tICs for the ATP-bound PKA-C (**B**), ATP/PKI bound PKA-C (**C**), and the projections of known crystal structures.

In the apo form, the αC-β4 loop is quite dynamic due to transient hydrophobic interactions between F100 and V104 as well as W222 and the APE motif (A206 and P207) (**Figure 4A**). The binding of both nucleotide and PKI increases the rigidity of residues near F100 and V104 such as V103 and F185 (**Figure 4A**). In addition, several electrostatic interactions essential for catalysis

(D166-N171, K168-T201, and Y204-E230), which are transient in the apo PKA-C, become more

persistent (**Figure 4B**).

The MSM makes it possible to use kinetic Monte Carlo sampling to characterize the slow tran-

sition between different states.[28] In the apo form, the GS features F100 and F102 in *trans*, a

configuration that stabilizes the interactions with the αE and αJ helices and, together with the

nucleotide, locks the αC-β4 loop to elicit an active kinase conformation (**Figure 3A**). The GS to

ES1 transition features the disruption of the K72-E91 salt bridge, whereas the GS to ES2 transi-

tion involves a 120º flip of the F100 aromatic group that interacts with V104, a conformation found

only in the uncommitted apo enzyme (**Figure 3A**). The GS to ES2 transition involves a concerted

disruption of the D166-N171 and K168-T201 electrostatic interactions, essential for catalysis. Also,

this event destabilizes the packing between W222 and the APE motif (A206 and P207) required

for substrate recognition (**Figure 3C**). All these conformational transitions suggest that ES1 and

ES2 represent destabilized states of the kinase.

Figure 4. Conformational transition between GS and ES1, ES2 along the kinetic Monte
Carlo trajectory in apo (A) and ATP-bound (B) forms. (A, B) The transition from GS to ES1,
revealed as breaking of the K72-E91 salt bridge, is frequently found in both forms, whereas the
transition to ES2, revealed as the contact between F100 and V104, only occurs in the apo form
and in concert with allosteric changes between D166-N171, K168-T201, and W222-A206-P207.
The darker colors in (A) and (B) highlight moving averages over every 10 frames. (C) GS confor-
mation reveals the assembly of key catalytic features across the core region. **(D)** ES2 confor-
mation revealed disruption of key structure motifs across the core region, indicative of inactivation.

Direct correspondence between the conformationally excited states identified by MD simulations and NMR. NMR relaxation dispersion and CEST experiments performed on the apo PKA-C revealed the presence of conformationally excited states for several residues embedded into the hydrophobic core of the enzyme.[20] Based on the free energy landscape and the conformational transitions in the core of the kinase, we reasoned that the inactive states identified by the simulations would correspond to a disruption of the hydrophobic packing near the αC-β4 loop. To test this hypothesis, we calculated the values of the change in CS ($\Delta\omega$) for the methyl groups of L103, V104, I150, L172, and I180 from the MD simulations and compared them with those obtained from the fitting of the CPMG dispersion curves and CEST profiles.[20] We first sampled more than 500 snapshots as representative structures of the ES2 and GS states (**Figure 3 – figure supplement 2**), and computed the distribution of the methyl ^{13}C chemical shifts on these sites using ShiftX2.[29] These calculations yield $\Delta\omega$ values of 0.87 ± 0.02 and 0.85 ± 0.02 ppm between ES2 and GS for Val104 and Ile150, respectively. These values are in good agreement with the experimental $\Delta\omega$ values (1.10 ± 0.06 and 0.94 ± 0.09 ppm) obtained from the fitting of the CPMG dispersion curves (**Figure 5A-B**)**.** The CS difference obtained for the other three sites also yielded a good agreement, with the same direction of the chemical shift changes found experimentally (**Figure 5 – figure supplement 1**). Remarkably, the fitting of the calculated vs. experimental $\Delta\omega$ values gives a slope of 0.86 with R^2 of 0.82 (**Figure 4C)** supporting that the excited state observed experimentally may correspond to the calculated ensemble. In addition, MSM estimates a population of ES2 of 6 ± 2%, consistent with the 5 ± 1% reported in NMR experiments.[27] These results support that the alternate conformations identified by NMR represent inactive states and may correspond to the disruption of the hydrophobic anchor of the αC-β4 loop via F100 and F102.

Figure 5. Transition from GS to ES2 shown in the apo PKA-C recapitulated the structural changes near the αC- β4 loop probed by NMR experiments. (**A** and **B**) Distribution of predicted ^{13}C CS of selected methyl groups in ES (magenta) and GS (blue) of the apo PKA-C for Val104-Cγ1 (**A**) and Ile150-Cδ1 (**B**). The experimental CS is shown in dotted lines for GS (black) and ES (red). (**C**). Correlation of the predicted chemical shift differences $|\Delta\omega^{Pred}|$ and the experimental result $|\Delta\omega^{Exp}|$ for a set of hydrophobic residues near the αC-β4 loop. The fitted linear correlation has a slope of 0.86 and R^2 of 0.82.

The F100A mutation disrupts the allosteric network of the kinase. The above analysis

suggests that F100 located at the αC-β4 loop is critical for the GS to ES1 transition path of the

kinase. Therefore, we modeled the F100A mutant and analyzed its dynamic trajectories. First, we

performed a short equilibration using classical MD simulations starting from the coordinates of the

X-ray structure of the ternary complex (PDB ID: 4WB5). During 1 µs of MD simulations, the αC-

β4 loop of the F100A mutant undergoes a significant motion as manifested by the increased val-

ues of the backbone rmsd and the conformational transition (flipping) of the F102 side chain (**Fig-**

ure 6A). This region in the WT PKA-C adopts a stable β-turn in the WT, with a persistent H-bond

between the backbone oxygen of F100 and the amide hydrogen of L103. In the simulation of

F100A, H-bond is formed more frequently between A100 and F102, and this region adopts a γ-

turn conformation. Such local rearrangement disrupts the hydrogen bond between N99 and Y156

of αE, altering the anchoring of the αC-β4 to the αE helix (**Figure 6B**).

Figure 6. Increased dynamics at the αC-β4 loop upon F100A mutation, perturbing the local
hydrophobic packing and its anchoring to the αE helix. (**A**) Time series of the αC-β4 loop,
H-bond occurrence for the β− and γ-turns, F102 χ_1 angle, and N99 and Y156 for WT (black) and
F100A (red) in the ATP-bound state. (**B**) Representative structural snapshots showing the for-
mation of the β-turn for the PKA-C[WT] (green) and γ-turn for the PKA-C[F100A] mutant.

In the F100A mutant, the local changes caused by the αC-β4 loop structural transitions propagate

and alter the response of the nucleotide binding as shown on the rmsd changes for key hydro-

phobic motifs such as the R- and C- spines and shell residues (**Figure 7A**). While the nucleotide

binding decreases the average RMSD for these hydrophobic motifs for WT PKA-C, it stabilizes

only the C-spine and fails to drive the R-spine and shell residues to an intermediate state compe-

tent for substrate binding (**Figure 7B**). The latter can be attributed to the perturbation of hydro-

phobic packing of L95, L106 of R-spine, and V104 of the shell residues close to the mutation site.

Using the lowest principal components, we also analyzed the global dynamic response to ATP

binding. Not only does F100A change the breathing mode of the two lobes (PC1), but it also alters

the shearing motion (PC2) of the kinase, emphasizing the importance of this allosteric mutation

on the internal communication across the hydrophobic core (**Figure 7C-E**).

Figure 7. Distinct global structural response to ATP binding in the F100A mutant. (A) Struc-
ture superposition of the C Spine, R Spine, and Shell residues between WT (lime) and F100A
(hot pink), highlighting the differences between Shell and R Spine. **(B)** Change of RMSD upon
ATP binding at C Spine, Shell, and R spine, for WT and F100A, respectively. **(C)** Structural illus-
tration of the first principal component (PC1), i.e., the breathing motion of the two lobes. **(D)** Struc-
tural illustration of the second principal component (PC2), i.e., the shearing motion of the two
lobes. **(E)** Comparison of the 2D projection and distribution along PC1 and PC2 for WT and F100A,
highlighting their dramatic differences along both axes.

In fact, these altered motions have global repercussions on the allosteric networks as detected

by mutual information for rotameric states (**Figure 8**). For PKA-C^{WT}, there are numerous correla-

tions observable within both lobes of the enzyme, involving the Gly-rich loop, αC-β4 loop, activa-

tion loop, and the C-terminal tail These dynamic correlations between the two lobes are the hall-

mark for the dynamically-committed state. In contrast, the mutual information for F100A mutants

shows significantly less correlations in the N-lobe and the correlations within the C-lobe are re-

distributed.

Figure 8. Mutual information of dihedral angles for the (A) WT and (B) F100A upon binding
ATP. This analysis reveals the prominent loss of allosteric communication of F100A, especially
at multiple key motifs as is highlighted by purple strips.
324

Taken together, these simulations on F100A suggest that the local perturbation of the αC-β4

loop diminishes the structural connection between the two lobes, disrupting the correlated shear-

ing motion highlighted in other simulation studies. Although the single F100A mutation does not

drive the kinase into a completely dysfunctional state (ES2), it is sufficient to abolish the binding

cooperativity of the enzyme.

The F100A mutation disrupts nucleotide-substrate binding cooperativity in PKA-C.

Based on the results from the MD simulations, we hypothesized that the disruption of the coupling

between the N-lobe and C lobe would affect the binding cooperativity between nucleotide (ATPγN)

and pseudosubstrate inhibitor (PKI$_{5-24}$) binding. We first evaluated the catalytic efficiency for the

wild-type and F100A mutant by carrying out steady-state coupled enzyme assays[30] using the

standard substrate, Kemptide.[31] F100A showed a slight increase in K_M and V_{max} compared to PKA-

C^{WT}, resulting in a reduction of the catalytic efficiency (k_{cat}/K_M = 0.50 ± 0.04 and 0.41 ± 0.08 for

PKA-C^{WT} and PKA-C^{F100A}, respectively; **Supplementary table 2**). We then performed isothermal

titration calorimetry (ITC)[32] to obtain ΔG, ΔH, $-T\Delta S$, K_d, and determine the cooperativity coefficient

(σ) for ATPγN and PKI$_{5-24}$ binding. We first analyzed the binding of ATPγN to the apo PKA-C^{F100A}

and, subsequently, the binding of PKI$_{5-24}$ to the ATPγN-saturated PKA-C^{F100A} (**Supplementary**

Table 3). We found that PKA-C^{WT} and PKAC^{F100A} have similar binding affinities for ATPγN (K_d =

83 ± 8 μM and 73 ± 2 μM, respectively). However, in the apo form F100A showed a 3-fold higher

binding affinity for the pseudosubstrate relative to PKA-C^{WT} (K_d = 5 ± 1 μM and 17 ± 2 μM, re-

spectively - **Supplementary table 3**). Upon saturation with ATPγN, PKA-C^{F100A} displayed a 12-

fold reduction in binding affinity for PKI$_{5-24}$, resulting in a σ of ~3, a value significantly lower than

the wild-type (σ greater than 100).

NMR mapping of nucleotide/pseudosubstrate binding response. To understand the

atomic details of the disrupted binding cooperativity for PKA-C^{F100A}, we used solution NMR spec-

troscopy and mapped the amide fingerprints of PKA-C^{F100A} using [^{1}H, ^{15}N]-WADE-TROSY-HSQC

experiments [20] upon adding nucleotide (ATPγN) and pseudosubstrate (PKI$_{5-24}$). Specifically, we

monitored the ^{1}H and ^{15}N chemical shift perturbations (CSPs, Δδ) of the amide fingerprints for the

PKA-C^{F100A} saturated with ATPγN and in complex with ATPγN/PKI$_{5-24}$ and we compared them with

PKA-C^{WT} (**Figure 9 – figure supplement 1**). ATPγN binding causes the dramatic broadening of

several amide resonances throughout the structure of PKA-C^{F100A}, suggesting the presence of an

intermediate conformational exchange [33]. The fingerprints of both PKAC^{WT} and PKA-C^{F100A} show

similar CSP profiles upon binding ATPγN (**Figure 9 – figure supplement 1B**), although the extent

of the changes is significantly attenuated for PKA-C^{F100A}. The reduction of CSP is apparent for

residues in the N-lobe (β2- β3 region), around the mutation (β4), and at the C-terminal tail. A

similar pattern is observed upon binding PKI$_{5-24}$ to the ATPγN-saturated PKA-C^{F100A} (**Figure 9**). A

substantial decrease in CSP is observed for the C-lobe near the domains critical for substrate-

binding (*i.e.*, αE, αF, and αG).

To determine the global response to ligand binding for WT and F100A, we analyzed the chem-

ical shifts of the amide fingerprints of the two proteins using the COordiNated ChemIcal Shifts

bEhavior (CONCISE).[34] CONCISE describes the global response of the protein fingerprint reso-

nances by providing the probability density (population) of each state along the conformational

equilibrium for binding phenomena that follow linear chemical shift trajectories. For PKA-C^{WT}, both

nucleotide and pseudosubstrate binding shift the overall population of the residues from an open

to an intermediate and a fully closed state (**Figure 10 – figure supplement 1**). A similar trend is

observed for PKA-C^{F100A}, though the probability densities are broader, indicating that the amide

resonances follow a less coordinated response.[34] Also, the maximum probability density for the

closed state shows that the ternary complex of the mutant is slightly more open than the corre-

sponding wild-type (**Figure 10 – figure supplement 1**). Overall, the shape of the probability dis-

tributions of the amide chemical shifts suggests that several residues do not respond in a coordi-

nated manner and that PKI$_{5-24}$ binding shifts the conformation of the kinase toward a partially

closed state, which may explain the loss in binding cooperativity as previously observed.[14-16]

**Figure 9. Structural response of PKA-C^{F100A} binding to nucleotide and protein kinase inhib-
itor.** (**A**) Histogram shows the chemical shift perturbation (CSP) of the amide fingerprint for PKA-
C^{F100A} (black) in response to ATPγN binding compared to the CSP obtained for the wild-type pro-
tein (cyan). The dashed line on the histogram indicates one standard deviation from the average
CSP. (**B**) CSPs of PKA-C^{F100A}/ATPγN amide resonances mapped onto the structure (PDB: 4WB5).
(**C**) CSP of amide fingerprint for PKA-C^{F100A} bound to ATPγN and PKI$_{5-24}$ (black), compared to the
CSP of the wild-type protein obtained in the same conditions. (**D**) CSP for the F100A/ATPγN/PKI
complex mapped onto the crystal structure (PDB: 4WB5).

To analyze these coordinated chemical shift changes in detail and define the allosteric network

of the kinase upon binding ligands and substrate, we utilized the CHEmical Shift Covariance Anal-

ysis (CHESCA),[15,16,20] a statistical method that identifies correlated responses of residues to a

specific perturbation. CHESCA works under the assumption that pairwise correlated chemical

shift changes between residues identify possible allosteric networks.[35,36] We found that coordi-

nated structural rearrangements, as identified by CHESCA, are a strong indication of binding co-

operativity in PKA-C.[15,16,20] Therefore, we compared the CHESCA maps for PKA-C^{WT} and

PKAC^{F100A} for four different states: *apo*, ATPγN-, ADP-, and ATPγN/PKI$_{5-24}$-bound. The CHESCA

matrix of PKA-C^{F100A} exhibits sparser and more attenuated (i.e., lower value of correlation coeffi-

cient) correlations relative to PKA-C^{WT} (**Figure 10**). Although many inter-lobe correlations are still

present for the mutant, several correlations in specific structural domains such as the αG-, αH-,

and αI- helices are absent or attenuated. For instance, the F100A mutation does not display cor-

relations between the αA-helix and the C-terminal tail. We also utilized the CHESCA maps to

assess the allosteric communication among the communities as defined by McClendon *et al.*[37]

Using the community definition, the CHESCA map for PKA-C^{WT} shows a strong correlation across

the enzyme, especially for structurally adjacent communities and at the interface between the two

lobes (see for instance the correlations among ComA, ComB, ComC, ComE, and ComH). The

community CHESCA for F100A shows that the cross-talk between the nucleotide-binding (ComA)

and positioning of αC-helix (ComB) is preserved, as well as the coupling between the R-spine

assembly (ComC), and the activation loop (ComF). Except for ComD, the correlations among

these communities in the N-lobe and ComF1, ComG, and Com H are completely abolished. Note

that ComF1, ComG, and ComH are involved in the substrate binding and are instrumental in

docking the R-subunits. Overall, the CHESCA analysis shows that the reduced degree of coop-

erativity we determined thermodynamically for PKA-C^{F100A} corresponds to a decrease in coordi-

nated structural changes upon ligand binding, and these changes affect the C-lobe communities

involved in the substrate binding and protein/protein interactions.

**Figure 10 – Changes of the intramolecular allosteric network in F100A as mapped by cor-
related chemical shift changes. (A)** Comparison of the CHESCA matrices obtained from the
analysis of the amide chemical shifts of PKA-C$_{WT}$ (top diagonal, blue) and PKA-C$_{F100A}$ (bottom
diagonal, black) in the apo, ADP-bound, ATPγN-bound, and ATPγN/PKI$_{5-24}$-bound states. Only
correlations with R$_{ij}$ > 0.98 are reported. The enlarged CHESCA map of F100A is available in
Figure 10 – figure supplement 1 while the data for the PKA-C^{WT} matrix are taken from Walker
et al.[15]. **(B)** Community CHESCA analysis of PKA-C^{WT} (top diagonal, blue) and PKA-C^{F100A} (bot-
tom diagonal, black). Only correlations with R$_{A,B}$ > 0.98 are shown. **(C)** Community CHESCA
matrix plotted on its corresponding structures. The size of each node is independent of the num-
ber of residues it encompasses while the weight of each line indicates the strength of coupling
between the individual communities.

DISCUSSION

A distinct feature of PKA-C is the binding cooperativity between nucleotide and substrate that

originate from the allosteric coupling between the nucleotide-binding pocket and the interfacial

region between the two lobes that harbors the substrate binding cleft. Structural and dynamic

NMR data suggested that a well-tuned coupling between the two lobes of the kinase is required

for efficient substrate binding. Also, additional NMR and functional studies showed that mutations

in the activation loop linked to Cushing's syndrome reduce drastically substrate binding affinity,

and more importantly, reduce the communication between the ligand binding pockets.[14-16] Inter-

estingly, a mutation (E31V) distal from the active site and linked to the progression of Cushing's

syndrome has a similar effect, suggesting a possible allosteric modulation of the kinase substrate

recognition.[16] NMR chemical shift perturbation data suggested that these mutations are con-

nected to allosteric nodes that, once perturbed, radiate their effects in the periphery of the enzyme

and prevent an efficient dynamic coupling between the two lobes of the enzyme. Interestingly,

these mutations do not prevent Kemptide phosphorylation. Rather, they cause a loss of substrate

fidelity with consequent aberrant phosphorylation of downstream substrates.[38-43] Additionally,

thermodynamics and recent NMR studies using different nucleotides and inhibitors demonstrated

that it is possible to control substrate binding affinity by changing the chemistry of the ligand at

the ATP binding pocket. Altogether, these studies suggest that phosphorylation reaction and bind-

ing synergy between ATP and substrates may be controlled independently.

Our NMR data combined with RAM simulations and MSM enabled us to comprehensively map

the free energy landscape of PKA-C in various forms. We found that the active kinase unleashed

from the regulatory subunits occupies a broad energy basin (GS) that corresponds to the confor-

mation of the ternary structure of PKA-C with ATP and pseudosubstrate (PKI$_{5-24}$) that exemplifies

a catalytic competent state, poised for phosphoryl transfer.[44] We also found two orthogonal con-

formationally excited states ES1 and ES2. While the ES1 state corresponds to the inactive kinase

conformations, the ES2 state was never observed in the crystallized structures. Our previous

CEST NMR measurements suggested the presence of a sparsely populated state that, at that

time, we were unable to characterize structurally. These new simulations and MSM show that the

transition from GS to ES2 is due to a disruption of hydrophobic packing, featuring a conformational

rearrangement for the αC-β4 loop, which causes a partial disruption of the hydrophobic R-spine.

These structural changes interrupt the allosteric coupling between the two lobes, as shown by

mutual information analysis. A single mutation (F100A), suggested by our simulations, promotes

the flip of the αC-β4 loop and reproduces the hypothesized structural uncoupling between the two

lobes of PKA-C. We experimentally tested the effects of the F100A mutations and found that it

prevents the enzyme from adopting a conformation competent for substrate binding, resulting in

a drastic reduction of the cooperativity between ATP and nucleotide. These NMR data further

support our working model, showing that the inter-lobe communication is interrupted and the bind-

ing response of the F100A kinase is attenuated based on chemical shift perturbation. Furthermore,

our study further emphasizes the active role of the hydrophobic interior of the kinase and shows

that small alterations in the spines and shell sequences may lead to a dysfunctional kinase by

either preventing phosphorylation (see V104G and I150A mutations)[27] or by disrupting the binding

cooperativity as for F100A.

The αC-β4 loop is a regulatory element present in all EPKs and its importance has been

stressed in bioinformatics studies[45] and supported by computational work, showing that F100 and

F102 are at intersections of various communities and constitute a critical hydrophobic node, an-

choring the N- to the C-lobe.[37] More importantly, studies on EGFR and ErbB2 kinases lead to the

hypothesis that the αC-β4 loop may act as a molecular brake [46,47] or an autoinhibitory switch.[48]

Therefore, it is not surprising that activating mutations and in-frame insertions in the αC-β4 loop

are frequently found in kinase-related cancers, such as somatic oncogenic mutations such as

P101S, P101L, and L103F.[49] Finally, an elegant study by Kannan and coworkers emphasized the

role of the αC-β4 loop in dimerization of EGFR.[21] These researchers found that in-frame insertions

rigidify and activate the kinase in a length dependent manner. More importantly, these human

mutations display a gradual response to drugs, a factor that may exploited for designing mutant-

selective inhibitors of EGFR.[50] The data presented here show that it is possible to abolish the

binding cooperativity of a kinase by turning the dial in the opposite direction, *i.e.*, increasing the

dynamics of the αC-β4 loop and disconnecting the allosteric network between the N- and C-lobes.

The identification of this new, partially inactivating pathway provides further understanding of how

to control the dynamics and function of kinases.

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Gene (*Homo sapiens)*	PKA-Cα		PKA-C	Uniprot ID: P17612
strain, strain background (*Escherichia coli*)	BL21(DE3) pLyss	Agilent	Cat. #200132	Chemically competent cells
Sequence-based reagent	PKA-C^{F100A}	This study	PCR primer (Forward)	tattctgcaagcggtg aacgccccgtttctg gttaagctg
Sequence-based reagent	PKIa 5-24	Synthetic peptide	PKI$_{5-24}$	Chemically synthesized
Sequence-based reagent	Kemptide ()	Synthetic peptide	Kemptited	LRRASLG
Commercial assay or kit	QuikChange Lightning Multi Mutagenesis Kit	Agilent genomics	Cat #210519	Commercial mutagenesis kit
Recombinant DNA reagent	PKA-^{CF100A}	This study	PKA-C^{F100A}	Single Ala mutant of PKA-C
Chemical compound, drug	AMP-PNP or ATPγN	Roche Applied Science	CAS 25612-73-1	ATP analogous
Chemical compound, drug	ADP	Sigma-Aldrich	CAS 20398-34-9	nucleotide
Software, algorithm	TopSpin 4.1	Bruker Inc.	https://www.bruker.com/	
Software, algorithm	NMRFAM-Sparky	NMRFam	https://nmr-fam.wisc.edu/nmr-fam-sparky-distribution/	

Software, algorithm	NMRPipe	Delaglio F, NIH	https://www.ibbr.umd.edu/nmrpipe/install.html	
Software, algorithm	POKY	Lee W,	https://sites.google.com/view/pokynmr	
Software, algorithm	COordiNated ChemIcal Shift bEhavior (CONCISE)	Veglia G	https://conservancy.umn.edu/handle/11299/217206 https://conservancy.umn.edu/handle/11299/227294	Matlab script
Software, algorithm	CHEmical Shift Covariance Analysis (CHESCA)	Melacini G	https://academic.oup.com/bioinformatics/article/37/8/1176/5905475?login=true	NMRFAM-Sparky&POKY tools
Software, algorithm	PyMol	Schrödinger, LLC	https://pymol.org	
Software, algorithm	MatLab2022b	MathWorks	https://www.mathworks.com/products/matlab.html	
Software, algorithm	GraphPad Prism 9	GraphPad Software Inc.	https://www.graphpad.com/	
Software, algorithm	GROMACS 4.6	Hess B et al	http://www.gromacs.org/	
Software, algorithm	CHARMM36a1	Best RB et al (2012)	https://pubs.acs.org/doi/10.1021/ct300400x	
Software, algorithm	PLUMED 2.1.1	Bonomi M, et al. (2009)	https://www.plumed.org/doc-v2.5/user-doc/html/_c_h_a_n_g_e_s-2-1.html	
Software, algorithm	ALMOST 2.1	Kohlhoff et al (2009).	svn://svn.code.sf.net/p/almost/code/almost-code	
Software, algorithm	METAGUI	Biarnés X et al. (2012)	https://www.sciencedirect.com/scienc	

			e/article/pii/S0010 465511003079	
Software, algo-rithm	MDTraj	McGibbon RT et al (2015)	https://www.scienc edirect.com/scienc e/article/pii/S0006 349515008267	
Software, algo-rithm	SPARTA+	Yang Shen, Ad Bax (2010)	https://spin.niddk.b ov/bax/software/S PARTA+/	
Software, algo-rithm	MSMbuilder	Beauchamp KA et al (2011)	https://msmbuilder .org/	
Software, algo-rithm	Mutinf	Mcclendon C et al (2012)	https://simtk.org/pr ojects/mutinf/	

MATERIAL AND METHODS

Replica-averaged metadynamics (RAM) simulations

I. System setup. We used the crystal structure of the wild type PKA-C (PDB ID: 1ATP) as the

template and added the missing residues 1-14 at the N terminus. The protonation state of histidine

residues followed our previous settings.[51] The protein was solvated in a rhombic dodecahedron

solvent box with the tree-point charge TIP3P model[52] for water extended approximately 10 Å away

from the surface of the proteins. Counter ions (K^+ and Cl^-) were added to ensure electrostatic

neutrality corresponding to an ionic concentration of ~ 150 mM. All protein covalent bonds were

constrained with the LINCS algorithm.[53] and long-range electrostatic interactions are treated with

the particle-mesh Ewald method with a real-space cut-off of 10 Å.[54] Parallel simulations on the

apo form, the binary form with one Mg^{2+} ion and one ATP, and the ternary form with two Mg^{2+} ions,

one ATP and one PKI_{5-24} are performed simultaneously using GROMACS 4.6[55] with

CHARMM36a1 force field.[56] For the two mutants, F100A and V104G, the corresponding residues

were mutated through the mutagenesis wizard of PYMOL.

II. Standard MD simulations. Each system was minimized using the steepest decent algorithm to

remove the bad contacts, and then gradually heated to 300 K at a constant volume over 1 ns,

using harmonic restraints with a force constant 1000 kJ/(mol *Å^2) on heavy atoms of both proteins

and nucleotides. Over the following 12 ns of simulations at constant pressure (1 atm) and tem-

perature (300 K), the restraints were gradually released. The systems were equilibrated for an

additional 20 ns without positional restraints. A Parrinello-Rahman barostat[57] was used to keep

the pressure constant, while a V-rescale thermostat [58] with a time step of 2 fs was used to keep

the temperature constant. Each system was simulated for 1.05 μs, with snapshots recorded every

20 ps.

III. Replica exchange (REX) simulations. Following the standard MD simulations, parallel replica

exchange (REX) simulations on the apo, binary and ternary form of PKA-C were set up. For each

REX simulations, four replicas were used, and the initial structures were randomly chosen from

the μs-scale unbiased simulations. Chemical shifts of PKA-C for N, CA, CO, CB, HN from NMR

experiments were imposed as restraints based on the penalty function

513
$$E^{cs} = \alpha \sum_{k=1}^{N} \sum_{l=1}^{5} \left(\delta_{kl}^{exp} - \frac{1}{M} \sum_{m=1}^{M} \delta_{kl}^{calc} \right)^2$$

where α is the force constant, k runs over all residues of the protein, l denotes the different back-

bone atoms, and m runs over the four replicas. δ_{kl}^{calc} is computed using CamShift that is a module

of ALMOST-2.1.[59] The force constant was gradually increased from 0 (unbiased) to 20 (maximum

restraints for production) over 50 ns. All other settings are the same as the previous unbiased

simulations. REX simulations were carried out with GROMACS 4.6,[55] with the replica exchange

controlled by the module PLUMED 2.1.1.[60] These simulations were further conducted ~100 ns for

each replica of the three forms.

IV. Replica-averaged metadynamics (RAM) simulations. The subsequent RAM simulations were

started from the final structures of REX simulations. The CS restraints were imposed in the same

way as the REX simulations. Four CVs are chosen to increase the conformational plasticity

around the catalytic cores (detailed in Figure S1): (CVI) the ψ angles of backbone of all the loops

that are not in contact with ATP (Back-far), (CVII) the ϕ angles of backbone of all the loops that

are in contact with ATP (Back-close), (CVIII) the χ_1 angles of side-chains of all the loops that are

in contact with ATP (Side-close), (CVIV) the radius of gyration calculated over the rigid part (*i.e.,*

residues 50-300) of the protein (rgss). Gaussian deposition rate was performed with an initial rate

of 0.125 kJ/mol/ps, where the σ values were set to 0.5, 0.2, 0.2, and 0.01 nm for the four CVs,

respectively. The RAM simulations were also carried out with GROMACS 4.6 in conjunction with

PLUMED 2.1 and ALMOST 2.1, and continued for ~400 ns for each replica, with exchange trails

every 1 ps.

V. Reconstruction of free energy surface (FES) from the RAM simulations. After about 300 ns,

the sampling along the first 3 CVs reached convergence, allowing a reliable reconstruction of the

corresponding FES. The production run was continued for another 100 ns to sample enough

conformations. These conformations were first clustered into microstates using the regular spatial

method (cut-off radius of 0.13), and then the free energy of each state is reweighted according to

the deposited potential along each CV, which can be obtained from the analysis module of

METAGUI.[61] To visualize the distribution of these microstates and their relative energies differ-

ences, we further performed principal component analysis to project the microstates represented

in the 3-dimennsional CV space into 2-dimensional space spanned by PC1 and PC2. Then we

can plot these microstates in a 3-dimensional space spanned by PC1, PC2 and free energy dif-

ferences *ΔG*.

VI. Independent validation of chemical shifts with SPARTA+. During the REX and RAM simula-

tions, chemical shifts were computed via CamShift in ALMOST 2.1. As an independent validation

for the efficacy of the bias, we further calibrated chemical shifts of 2000 snapshots with MDTraj[62]

and SPARTA+.[23]

VII. Adaptive sampling. The first round of adaptive sampling started from the snapshots of low-

energy microstates obtained from the previous step, *i.e.*, 1200 structures for the apo form, 400

structures for the binary form and 200 structures for the ternary form. The initial velocities were

randomly generated to satisfy the Maxwell distribution at 300K. For the apo form, a 10 ns simula-

tion was performed for each run, whereas for the binary, each simulation lasted 30 ns, resulting

in a total of 12 μs trajectories for both the apo and binary forms. To obtain converged free energy

landscape, a total of three rounds of adaptive sampling was started from the 400 microstates that

was obtained by K-mean clustering of all snapshots of previous ensembles. Therefore, a total of

100 μs trajectories and 100,0000 snapshots (100 ps per frame) were collected for both the apo

and binary form after three rounds of adaptive sampling, and a total of 60 μs trajectories were

collected for the ternary form.

VIII. Markov state model (MSM) and time-lagged independent component analysis (tICA). The

Cartesian coordinates of key hydrophobic residues, include R-spine residues, L95, L106, Y164

and F185, and the shell residues, V104, M118 and M120, were chosen as the metrics to charac-

terize the conformational transition of the hydrophobic core of PKA-C. Specifically, each snapshot

was first aligned to the same reference structure by superimposition of αE (residues 140-160)

and αF helices (residues 217-233), and represented by the deviation of Cartesian coordinates of

the key residues. The representation in this metric space was further reduced to 10-dimension

vectors using time-lagged independent component analysis (tICA)[63] at a lag time of 1 ns. All the

snapshots were clustered into 400 microstates with K-mean clustering. A MSM was built upon

the transition counts between these microstates.

IX Kinetic Monte Carlo trajectory of PKA-C in different forms. Long trajectories were generated

using a kinetic Monte Carlo method based on the MSM transition probability matrix of the three

forms of PKA-C. Specifically, the discrete jumps between the 100 microstates were sampled for

60 us. And then random conformations were chosen for each state from all the available snap-

shots. Time series of various order parameters were analyzed subsequently.

Protein expression and purification

The recombinant human Cα subunit of cAMP-dependent protein kinase with the Phe to Ala mu-

tation in position 100 (PKA-C^{F100A}) was generated from the human PKA-Cα wild-type using Quik-

Change Lightning mutagenesis kit (Agilent genomics). The key resource table lists the PCR pri-

mers used to modify the pET-28a expression vector encoding the wild-type human PKA-Cα gene

(*PRKACA* – uniprot P17612) [14-16]. The unlabeled and uniformly ^{15}N-labeled PKA-C^{F100A} mutant

was expressed and purified following the same protocols used for the wild-type protein.[64] Briefly,

transformed *E. coli* BL21 (DE3) *pLyss* cells (Agilent) were cultured overnight at 30 °C in Luria-

Bertani (LB) medium. The next morning, the cells were transferred to fresh LB medium for the

overexpression of the unlabeled protein or to M9 minimal medium supplied with ^{15}NH$_4$Cl (Cam-

bridge Isotope Laboratories Inc.) as the only nitrogen source for the labeled protein overexpres-

sion. In both cases, protein overexpression was induced with 0.4 mM of β-D-thiogalactopyra-

noside (IPTG) and carried out for 16 hours at 20 °C. The cells were harvested by centrifugation

and resuspended in 50 mM Tris-HCl, 30 mM KH$_2$PO$_4$, 100 mM NaCl, 5 mM 2-mercaptoethanol,

0.15 mg/mL lysozyme, 200 μM ATP, DNaseI, 1 tablet of cOmplete ULTRA EDTA-free protease

inhibitors (Roche Applied Science) (pH 8.0) and lysed using French press at 1,000 psi. The cell

lysate was cleared by centrifugation (60,000 × g, 4 °C, 45 min), and the supernatant was batch-

bound with Ni^{2+}-NTA agarose affinity resin (Thermo Scientific). The his-tagged PKA-C^{F100A} was

eluted with 50 mM Tris-HCl, 30 mM KH$_2$PO$_4$, 100 mM NaCl, 5 mM 2-mercaptoethanol, 0.5 mM

phenylmethylsulfonyl fluoride (PMSF) (pH 8.0) supplied with 200 mM of imidazole. The tail of

poly-His was cleaved using a stoichiometric amount of recombinant tobacco etch virus (TEV)

protease in 20 mM KH$_2$PO$_4$, 25 mM KCl, 5 mM 2-mercaptoethanol, 0.1 mM PMSF (pH 6.5), over-

night at 4 °C. The different phosphorylation states of PKA-C^{F100A} were separated using a cation

exchange column (HiTrap Q-SP, GE Healthcare Life Sciences) using a linear gradient of KCl in

20 mM KH$_2$PO$_4$ at pH6.5.[65] The purified protein isoforms were then stored in phosphate buffer

containing 10 mM dithiothreitol (DTT), 10 mM MgCl$_2$, and 1 mM NaN$_3$ at 4 °C. The protein purity

was assessed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS–PAGE).

Peptide synthesis

The Kemptide (LRRASLG) and PKI$_{5-24}$ (TTYADFIASGRTGRRNAIHD) peptides were synthesized

using a CEM Liberty Blue microwave synthesizer using standard Fmoc chemistry. All peptides

were cleaved with Reagent K (82.5% TFA, 5% phenol, 5% thioanisole, 2.5% ethanedithiol, and

5% water) for 3 h and purified using a semipreparative Supelco C18 reverse-phase HPLC column

at 3 mL/min.

The purified peptides were concentrated, lyophilized, and stored at −20 °C. Molecular weight and

quantity were verified by MALDI-TOF and/or amino-acid analysis (Texas A&M University).

Isotherm titration calorimetry (ITC) measurements

PKA-C^{F100A} was dialyzed into 20 mM MOPS, 90 mM KCl, 10 mM DTT, 10 mM MgCl$_2$, and 1 mM

NaN$_3$ (pH 6.5) and concentrated using conical spin concentrator (10 KDa membrane cut-off, Mil-

lipore) to a solution at 80-100 µM, as confirmed by A280 = 55,475 M^{-1} cm^{-1}. Approximately 300 µL

of protein was used for each experiment, with 50 µL of 2 mM ATPγN and/or 1 mM PKI$_{5-24}$ in the

titrant syringe. All measurements were performed at 300 K in triplicates with a low-volume

NanoITC (TA Instruments). The binding was assumed to be 1:1, and curves were analyzed with

the NanoAnalyze software (TA Instruments) using the Wiseman isotherm [32]

617
$$\frac{d[MX]}{d[X_{tot}]} = \Delta H^\circ V_0 \left[\frac{1}{2} + \frac{1 - \frac{1-r}{2} - R_m/2}{(R_m^2 - 2R_m(1-r) + (1+r)^2)^{1/2}} \right]$$

618   where $d[MX]$ is the change in total complex relative to the change in total protein concentration,

619     $d[X_{tot}]$ is dependent on r (the ratio of K_d relative to the total protein concentration), and R_m (the

620   ratio between total ligand and total protein concentration). The heat of dilution of the ligand into

621   the buffer was considered for all experiments and subtracted.

622   The free energy of binding was determined from:

623
$$\Delta G = RTlnK_d$$

624 where R is the universal gas constant and T is the temperature at measurement (300 K). The

625 entropic contribution to binding was calculated using:

626
$$T\Delta S = \Delta H - \Delta G$$

627 The degree of cooperativity (σ) was calculated as:

628
$$\sigma = \frac{K_{d\,apo}}{K_{d\,nucleotide}}$$

where $K_{d\,apo}$ is the dissociation constant of PKI_{5-24} binding to the apo-enzyme, and $K_{d\,Nucleotide}$ is the

corresponding dissociation constant for PKI_{5-24} binding to the nucleotide-bound the kinase.

Enzyme assays

Steady-state coupled enzyme activity assays using Kemptide as substrate were performed under

saturating ATP concentrations and spectrophotometrically at 298 K, as reported by Cook *et al.*[30]

The values of V_{max} and K_M were obtained from a nonlinear fit of the initial velocities to the Mich-

aelis-Menten equation.

NMR spectroscopy

NMR measurements were performed on a Bruker Avance NEO spectrometer operating at a ^{1}H

Larmor frequency of 600 MHz equipped with a cryogenic probe or on a Bruker Avance III 850

MHz spectrometer equipped with a TCI cryoprobe. The NMR experiments were recorded at 300K

in 20 mM KH_2PO_4 (pH 6.5), 90 mM KCl, 10 mM $MgCl_2$, 10 mM DTT, 1 mM NaN_3, 5% D_2O, and

0.1% 4-benzene sulfonyl fluoride hydrochloride (AEBSF, Pefabl–c - Sigma-Aldrich). Concentra-

tions for samples were 0.15 mM of uniformly ^{15}N-labeled PKA-C^{F100A}, as determined by A_{280} meas-

urements, 12 mM ATPγN or ADP was added for the nucleotide-bound form, and 0.3 mM PKI_{5-24}

for the ternary complex. [^{1}H, ^{15}N]-WADE-TROSY-HSQC pulse sequence [66] was used to record

the amide fingerprint spectra of PKA-C^{F100A} in the apo, nucleotide-bound (ADP- or ATPγN-bound

– binary form), and ternary complex (PKAC^{F100A}/ATPγN/PKI_{5-24}). All [^{1}H, ^{15}N]-WADE-TROSY-

HSQC experiments were acquired with 2048 (proton) and128 (nitrogen) complex points, pro-

cessed using NMRPipe[67] and visualized using NMRFAM-SPARKY [68] and POKY.[69] Combined

chemical shift perturbations (CSP) were calculated using ^{1}H and ^{15}N chemical shifts according to:

$$\Delta\delta = \sqrt{(\Delta\delta H)^2 + (0.154 \times \Delta\delta N)^2}$$

in which $\Delta\delta$ is the CSP; $\Delta\delta_H$ and $\Delta\delta_N$ are the differences of ^{1}H and ^{15}N chemical shifts, respectively,

between the first and last point of the titration; and 0.154 is the scaling factor for nitrogen.[70]

COordiNated ChemIcal Shift bEhavior (CONCISE). The normal distributions reported in the

CONCISE plot were calculated using principal component analysis for residues whose chemical

shifts responded linearly to ligand binding.[34] In this work, we use the ^{1}H and ^{15}N chemical shifts

derived from the [^{1}H, ^{15}N]-WADE-TROSY-HSQC experiments for the apo, ADP, ATPγN, and

ATPγN/PKI$_{5-24}$ bound forms of PKA-C.

CHEmical Shift Covariance Analysis (CHESCA). This analysis was used to identify and func-

tionally characterize allosteric networks of residues eliciting concerted responses to, in this case,

nucleotide and pseudosubstrate. To identify inter-residue correlations, four states were used: apo,

ATPγN-bound, ADP-bound, and ATPγN/PKI$_{5-24}$. The identification of inter-residue correlations by

CHESCA relies on agglomerative clustering (AC) and singular value decomposition (SVD). Pair-

wise correlations between chemical shift variations experienced by different residues were calcu-

lated to identify networks. When plotted on a correlation matrix, this allows for the identification of

regions that are correlated to one another. For each residue, the maximum change in chemical

shift was calculated in both the ^{1}H (*x*) and ^{15}N (*y*) dimensions ($\Delta\delta_{x,y}$). The residues included in the

CHESCA analysis were the ones that satisfied the following: $\Delta\delta_{x,y} > \frac{1}{2}\,\Delta\nu_{xA,yA} + \frac{1}{2}\,\Delta\nu_{xB,yB}$, where A

and B correspond to two different forms analyzed (note that there is no dependence on which two

forms satisfied this statement), and $\Delta\nu$ denotes the linewidth. Correlation scores were used to

quantify the CHESCA correlation of a single residue or a group of residues with another group.

Correlation scores were evaluated for both (a) a single residue and (b) the full protein. The gen-

eralized expression for evaluating either case is:

673
$$Corr\ Score = \frac{number\ of\ (R_{ij} > R_{cutoff})}{total\ number\ of\ R_{ij}}$$

where R_{ij} is the CHESCA correlation matrix and i and j denote (a) a single residue and all other

assigned residues in the protein, respectively, or (b) both represent all the assigned residues in

the protein. For all the analysis a R_{cutoff} of 0.98 was used.

Community CHESCA analysis is a chemical shift-based correlation map between functional com-

munities within the kinase. Each community is a group of residues [37] associated with a function

or regulatory mechanism. To represent community-based CHESCA analysis, we lowered the cor-

relation cut-off such that R_{cutoff} > 0.8.

Suppose communities X and Y have n_A and n_B number of assigned residues respectively, the

correlation score between A and B is defined as,

683
$$R_{A,B} = number\ of\ (R_{ij} > R_{cutoff})/(n_A * n_B)$$

684   where R_{ij} is the CHESCA correlation coefficient between residue i (belongs to community A) and

685   residue j (belongs to community B). R_{cutoff} is the correlation value cutoff. $R_{A,B}$ can take values from

686   0 (no correlation between residues in A and B) to 1 (all residues in A has correlation > cutoff with

687   all residues in B).

Acknowledgements

689   This work was supported by the National Institute of Health GM 100310 and HL 144130 to GV.

690   The authors would like to acknowledge the Minnesota Supercomputing Institute for MD calcula-

691   tions. YW would like to thank Guangdong Pearl River Talent Program (2021QN02Y618) and Na-

692   tional Natural Science Foundation of China (22007069, 92269102) for part of the MD analysis

693   carried out at the Shenzhen Bay Laboratory Supercomputing Centre.

Competing interest

The authors declare no financial and non-financial competing interests.

Author contributions

Y.W., C.O., and G.V. designed research. Y.W. performed the MD simulations and analysis. C.O. prepared the kinase samples, executed all NMR and ITC experiments, and analyzed the results. C.W. performed the kinase activity assay and prepared the peptides for the assay. C.O., Y.W. contributed to the manuscript, preparing all the text and figures. M.V.S., K.N.H., C.C., J.G., and M.V. contributed to the critical analysis of the data and writing of the manuscript. D.A.B. contributed to the critical analysis of the data and writing the manuscript. S.S.T. contributed to the writing of the manuscript. G.V. designed and directed the experiments and contributed to the writing of the manuscript. All authors have given approval to the final version of the manuscript.

To whom correspondence should be addressed. Email: vegli001@umn.edu.

References

1. Manning, G., Whyte, D.B., Martinez, R., Hunter, T. & Sudarsanam, S. The Protein Kinase Complement of the Human Genome. *Science* **298**, 1912-1934 (2002).
2. Knighton, D.R. et al. Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 407-14 (1991).
3. Knighton, D.R. et al. Structure of a peptide inhibitor bound to the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 414-20 (1991).
4. Taylor, S.S., Ilouz, R., Zhang, P. & Kornev, A.P. Assembly of allosteric macromolecular switches: lessons from PKA. *Nat Rev Mol Cell Biol* **13**, 646-58 (2012).
5. Walsh, D.A., Perkins, J.P. & Krebs, E.G. An adenosine 3',5'-monophosphate-dependant protein kinase from rabbit skeletal muscle. *J Biol Chem* **243**, 3763-5 (1968).
6. Smith, F.D. et al. Local protein kinase A action proceeds through intact holoenzymes. *Science* **356**, 1288-1293 (2017).
7. Knighton, D.R., Xuong, N.H., Taylor, S.S. & Sowadski, J.M. Crystallization studies of cAMP-dependent protein kinase. Cocrystals of the catalytic subunit with a 20 amino acid residue peptide inhibitor and MgATP diffract to 3.0 A resolution. *J Mol Biol* **220**, 217-20 (1991).
8. Johnson, D.A., Akamine, P., Radzio-Andzelm, E., Madhusudan, M. & Taylor, S.S. Dynamics of cAMP-dependent protein kinase. *Chem Rev* **101**, 2243-70 (2001).
9. Kornev, A.P., Haste, N.M., Taylor, S.S. & Eyck, L.F. Surface comparison of active and inactive protein kinases identifies a conserved activation mechanism. *Proc Natl Acad Sci U S A* **103**, 17783-8 (2006).
10. Ten Eyck, L.F., Taylor, S.S. & Kornev, A.P. Conserved spatial patterns across the protein kinase family. *Biochim Biophys Acta* **1784**, 238-43 (2008).
11. Hu, J. et al. Kinase regulation by hydrophobic spine assembly in cancer. *Mol Cell Biol* **35**, 264-76 (2015).
12. Lew, J., Coruh, N., Tsigelny, I., Garrod, S. & Taylor, S.S. Synergistic binding of nucleotides and inhibitors to cAMP-dependent protein kinase examined by acrylodan fluorescence spectroscopy. *J Biol Chem* **272**, 1507-13 (1997).
13. Wang, Y. et al. Globally correlated conformational entropy underlies positive and negative cooperativity in a kinase's enzymatic cycle. *Nat Commun* **10**, 799 (2019).
14. Olivieri, C. et al. Defective internal allosteric network imparts dysfunctional ATP/substrate-binding cooperativity in oncogenic chimera of protein kinase A. *Commun Biol* **4**, 321 (2021).
15. Walker, C. et al. Cushing's syndrome driver mutation disrupts protein kinase A allosteric network, altering both regulation and substrate specificity. *Sci Adv* **5**, eaaw9298 (2019).
16. Walker, C. et al. Is Disrupted Nucleotide-Substrate Cooperativity a Common Trait for Cushing's Syndrome Driving Mutations of Protein Kinase A? *J Mol Biol* **433**, 167123 (2021).
17. Camilloni, C., Cavalli, A. & Vendruscolo, M. Assessment of the use of NMR chemical shifts as replica-averaged structural restraints in molecular dynamics simulations to characterize the dynamics of proteins. *J Phys Chem B* **117**, 1838-43 (2013).
18. Camilloni, C. & Vendruscolo, M. Statistical mechanics of the denatured state of a protein using replica-averaged metadynamics. *J Am Chem Soc* **136**, 8982-91 (2014).
19. Husic, B.E. & Pande, V.S. Markov State Models: From an Art to a Science. *J Am Chem Soc* **140**, 2386-2396 (2018).
20. Olivieri, C. et al. ATP-competitive inhibitors modulate the substrate binding cooperativity of a kinase by altering its conformational entropy. *Sci Adv* **8**, eabo0696 (2022).

21. Ruan, Z. & Kannan, N. Altered conformational landscape and dimerization dependency
underpins the activation of EGFR by alpha C-beta 4 loop insertion mutations.
Proceedings of the National Academy of Sciences of the United States of America **115**,
Eb162-Eb171 (2018).
22. Piana, S. & Laio, A. A bias-exchange approach to protein folding. *J Phys Chem B* **111**,
4553-9 (2007).
23. Shen, Y. & Bax, A. SPARTA+: a modest improvement in empirical NMR chemical shift
prediction by means of an artificial neural network. *J Biomol NMR* **48**, 13-22 (2010).
24. Masterson, L.R. et al. Dynamically committed, uncommitted, and quenched states
encoded in protein kinase A revealed by NMR spectroscopy. *Proc Natl Acad Sci U S A*
108, 6969-74 (2011).
25. Kohlhoff, K.J. et al. Cloud-based simulations on Google Exacycle reveal ligand
modulation of GPCR activation pathways. *Nat Chem* **6**, 15-21 (2014).
26. Plattner, N., Doerr, S., De Fabritiis, G. & Noe, F. Complete protein-protein association
kinetics in atomic detail revealed by molecular dynamics simulations and Markov
modelling. *Nature Chemistry* **9**, 1005-1011 (2017).
27. Kim, J. et al. A dynamic hydrophobic core orchestrates allostery in protein kinases. *Sci
Adv* **3**, e1600663 (2017).
28. Shukla, D., Meng, Y., Roux, B. & Pande, V.S. Activation pathway of Src kinase reveals
intermediate states as targets for drug design. *Nat Commun* **5**, 3397 (2014).
29. Han, B., Liu, Y.F., Ginzinger, S.W. & Wishart, D.S. SHIFTX2: significantly improved
protein chemical shift prediction. *Journal of Biomolecular Nmr* **50**, 43-57 (2011).
30. Cook, P.F., Neville, M.E., Jr., Vrana, K.E., Hartl, F.T. & Roskoski, R., Jr. Adenosine
cyclic 3',5'-monophosphate dependent protein kinase: kinetic mechanism for the bovine
skeletal muscle catalytic subunit. *Biochemistry* **21**, 5794-9 (1982).
31. Kemp, B.E., Graves, D.J., Benjamini, E. & Krebs, E.G. Role of Multiple Basic Residues
in Determining Substrate-Specificity of Cyclic Amp-Dependent Protein-Kinase. *Journal of
Biological Chemistry* **252**, 4888-4894 (1977).
32. Wiseman, T., Williston, S., Brandts, J.F. & Lin, L.-N. Rapid measurement of binding
constants and heats of binding using a new titration calorimeter. *Analytical Biochemistry*
179, 131-137 (1989).
33. Srivastava, A.K. et al. Synchronous Opening and Closing Motions Are Essential for
cAMP-Dependent Protein Kinase A Signaling. *Structure* **22**, 1735-43 (2014).
34. Cembran, A., Kim, J., Gao, J. & Veglia, G. NMR mapping of protein conformational
landscapes using coordinated behavior of chemical shifts upon ligand binding. *Phys
Chem Chem Phys* **16**, 6508-18 (2014).
35. Akimoto, M. et al. Signaling through dynamic linkers as revealed by PKA. *Proc Natl Acad
Sci U S A* **110**, 14231-6 (2013).
36. Selvaratnam, R., Chowdhury, S., VanSchouwen, B. & Melacini, G. Mapping allostery
through the covariance analysis of NMR chemical shifts. *Proc Natl Acad Sci U S A* **108**,
6133-8 (2011).
37. McClendon, C.L., Kornev, A.P., Gilson, M.K. & Taylor, S.S. Dynamic architecture of a
protein kinase. *Proc Natl Acad Sci U S A* **111**, E4623-31 (2014).
38. Beuschlein, F. et al. Constitutive activation of PKA catalytic subunit in adrenal Cushing's
syndrome. *N Engl J Med* **370**, 1019-28 (2014).
39. Calebiro, D., Bathon, K. & Weigand, I. Mechanisms of Aberrant PKA Activation by Cα
Subunit Mutations. *Horm Metab Res* **49**, 307-314 (2017).
40. Calebiro, D. et al. PKA catalytic subunit mutations in adrenocortical Cushing's adenoma
impair association with the regulatory subunit. *Nature Communications* **5**, 5680 (2014).

41.   Di Dalmazi, G. et al. Novel somatic mutations in the catalytic subunit of the protein
kinase A as a cause of adrenal Cushing's syndrome: a European multicentric study. *J*
Clin Endocrinol Metab **99**, E2093-100 (2014).
42.   Omar, M.H. et al. Mislocalization of protein kinase A drives pathology in Cushing's
syndrome. *Cell Rep* **40**, 111073 (2022).
43.   Lubner, J.M. et al. Cushing's syndrome mutant PKA(L)(205R) exhibits altered substrate
specificity. *FEBS Lett* **591**, 459-467 (2017).
44.   Gerlits, O. et al. Zooming in on protons: Neutron structure of protein kinase A trapped in
a product complex. *Sci Adv* **5**, eaav0482 (2019).
45.   Kannan, N. & Neuwald, A.F. Did protein kinase regulatory mechanisms evolve through
elaboration of a simple structural component? *Journal of Molecular Biology* **351**, 956-972
(2005).
46.   Chen, H. et al. A molecular brake in the kinase hinge region regulates the activity of
receptor tyrosine kinases. *Mol Cell* **27**, 717-30 (2007).
47.   Klein, T. et al. Structural and dynamic insights into the energetics of activation loop
rearrangement in FGFR1 kinase. *Nature Communications* **6**(2015).
48.   Fan, Y.X. et al. Mutational activation of ErbB2 reveals a new protein kinase
autoinhibition mechanism. *J Biol Chem* **283**, 1588-1596 (2008).
49.   McSkimming, D.I. et al. ProKinO: a unified resource for mining the cancer kinome. *Hum*
Mutat **36**, 175-86 (2015).
50.   Yeung, W., Ruan, Z. & Kannan, N. Emerging roles of the alphaC-beta4 loop in protein
kinase structure, function, evolution, and disease. *IUBMB Life* **72**, 1189-1202 (2020).
51.   Cembran, A. et al. Conformational equilibrium of N-myristoylated cAMP-dependent
protein kinase A by molecular dynamics simulations. *Biochemistry* **51**, 10186-96 (2012).
52.   Jorgensen, W.L., Chandrasekhar, J., Madura, J.D., Impey, R.W. & Klein, M.L.
Comparison of Simple Potential Functions for Simulating Liquid Water. *Journal of*
Chemical Physics **79**, 926-935 (1983).
53.   Hess, B., Bekker, H., Berendsen, H.J.C. & Fraaije, J.G.E.M. LINCS: A linear constraint
solver for molecular simulations. *Journal of Computational Chemistry* **18**, 1463-1472
(1997).
54.   Darden, T., York, D. & Pedersen, L. Particle Mesh Ewald - an N.Log(N) Method for
Ewald Sums in Large Systems. *Journal of Chemical Physics* **98**, 10089-10092 (1993).
55.   Hess, B., Kutzner, C., van der Spoel, D. & Lindahl, E. GROMACS 4: Algorithms for
Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J Chem Theory*
Comput **4**, 435-47 (2008).
56.   Best, R.B. et al. Optimization of the additive CHARMM all-atom protein force field
targeting improved sampling of the backbone phi, psi and side-chain chi(1) and chi(2)
dihedral angles. *J Chem Theory Comput* **8**, 3257-3273 (2012).
57.   Parrinello, M. & Rahman, A. Crystal-Structure and Pair Potentials - a Molecular-
Dynamics Study. *Physical Review Letters* **45**, 1196-1199 (1980).
58.   Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling. *J*
Chem Phys **126**, 014101 (2007).
59.   Kohlhoff, K.J., Robustelli, P., Cavalli, A., Salvatella, X. & Vendruscolo, M. Fast and
accurate predictions of protein NMR chemical shifts from interatomic distances. *J Am*
Chem Soc **131**, 13894-5 (2009).
60.   Bonomi, M. et al. PLUMED: A portable plugin for free-energy calculations with molecular
dynamics. *Computer Physics Communications* **180**, 1961-1972 (2009).
61.   Biarnes, X., Pietrucci, F., Marinelli, F. & Laio, A. METAGUI. A VMD interface for
analyzing metadynamics and molecular dynamics simulations. *Computer Physics*
Communications **183**, 203-211 (2012).

62.   McGibbon, R.T. et al. MDTraj: A Modern Open Library for the Analysis of Molecular
Dynamics Trajectories. *Biophys J* **109**, 1528-32 (2015).
63.   Schwantes, C.R. & Pande, V.S. Improvements in Markov State Model Construction
Reveal Many Non-Native Interactions in the Folding of NTL9. *J Chem Theory Comput* **9**,
2000-2009 (2013).
64.   Masterson, L.R., Mascioni, A., Traaseth, N.J., Taylor, S.S. & Veglia, G. Allosteric
cooperativity in protein kinase A. *Proc Natl Acad Sci U S A* **105**, 506-11 (2008).
65.   Yonemoto, W., Garrod, S.M., Bell, S.M. & Taylor, S.S. Identification of phosphorylation
sites in the recombinant catalytic subunit of cAMP-dependent protein kinase. *J Biol*
Chem **268**, 18626-32 (1993).
66.   Manu, V.S., Olivieri, C., Pavuluri, K. & Veglia, G. Design and applications of water
irradiation devoid RF pulses for ultra-high field biomolecular NMR spectroscopy.
Physical Chemistry Chemical Physics **24**, 18477-18481 (2022).
67.   Delaglio, F. et al. NMRPipe: a multidimensional spectral processing system based on
UNIX pipes. *J Biomol NMR* **6**, 277-93 (1995).
68.   Lee, W., Tonelli, M. & Markley, J.L. NMRFAM-SPARKY: enhanced software for
biomolecular NMR spectroscopy. *Bioinformatics* **31**, 1325-1327 (2015).
69.   Manthey, I. et al. POKY software tools encapsulating assignment strategies for solution
and solid-state protein NMR data. *J Struct Biol X* **6**, 100073 (2022).
70.   Williamson, M.P. Using chemical shift perturbation to characterise ligand binding. *Prog*
Nucl Magn Reson Spectrosc **73**, 1-16 (2013).

A Back-far   **B** Back-close   **C** Side-close   **D** Rgss

Figure 2 – figure supplement 1. **Illustration of the collective variables (CVs) used in the**
RAM simulations. (**A**) The ψ angles of the backbone of all the loops not in contact with ATP
(Back-far), where the Cα atoms of the residues involved are highlighted in the blue sphere. (**B**)
The ψ angles of the backbone of all the loops in contact with ATP (Back-close), where the Cα
atoms of the residues involved are highlighted in magenta sphere. (**C**) The $\chi 1$ angles of side
chains of all the loops that are in contact with ATP (Side-close), where the side chains of the
residues involved are highlighted in magenta stick. (**D**) The radius of gyration is calculated over
the rigid part of the protein (rgss), where the residues involved are colored in cyan.

Figure 2 – figure supplement 2. **Distribution of the Root-Mean-Square-Error (RMSE) of the chemical shifts in different simulation schemes**. (**A**) RMSE of CS for the apo PKA-C from standard MD (left), REX (middle), and RAM (right). (**B**) RMSE of CS for PKA-C/ATP from standard MD (left), REX (middle), and RAM (right). (**C**) RMSE of CS for PKA-C/ATP/PKI$_{5-24}$ from standard MD (left), REX (middle), and RAM (right). Color codes for different backbone atoms (C, Cα, CO, H and N) are shown in the left figures.

Figure 2 – figure supplement 3. **Replica-averaged metadynamics (RAM) simulations explore a larger conformational space than standard MD and replica exchange (REX) simulations**. (**A**) Comparison of conformational space sampled by RAM Replica 1, standard MD, and REX Replica 1 of the apo PKA-C, along the CV1 and CV2. (**B**) Comparison of conformational space sampled by RAM Replica 1, standard MD, and REX Replica 1 of the apo PKA-C, along the CV3 and CV2.

Figure 2 – figure supplement 4. Accumulative deposition of history-dependent biases along the first three CVs for the RAM simulation of the apo PKA-C. The accumulative biased converged after around 300 ns along all three CVs.

Figure 3 – figure supplement 1. **Residues of the regulatory spine and shell are chosen as the metrics for two time-lagged independent components (tICA) and Markov State Model (MSM) analysis.** (**A**) Atom motions of key residues that define tIC1 of the apo PKA-C, colored by the superposition deviations. Backbone atoms of Val104 show the largest change in tIC1. (**B**) Atom motions of key residues that define tIC2 of the apo PKA-C, colored by the superposition deviations. Backbone atoms of Phe185 and Val104 show largest change in tIC2.

Figure 3 – figure supplement 2. **ES and GS in the apo PKA-C show distinct hydrophobic**
packing for residues around the αC-β4 loop. (**A**) Projections of randomly selected confor-
mations for ES (magenta) and GS (blue) onto the conformational landscape of the apo PKA-C.
To best separate ES from GS, snapshots with tIC1 < 1.2 were clustered as ES, whereas those
with tIC1 > 0.2 were clustered as GS. **(B,C)** Representative structure of ES (**B**) reveals different
hydrophobic packing from that of GS (**C**), highlighted by the distinction at Leu103, Val104, Ile150,
Leu172, and Ile180, where all show slow chemical exchanges in CPMG experiment of the apo
PKA-C.

Figure S7. **Distribution of predicted ^{13}C CS of selected methyl groups**. **(A-C)** ES (magenta)
and GS (blue) of the apo PKA-C for Leu103-Cδ2 (**A**), Leu172-Cδ1 (**B**) and Ile180-Cδ1 (**C**). The
experimental CS are shown in dotted line for GS (black) and ES (red).

Figure S8 - figure supplement 1. NMR fingerprints of PKA-C^{F100A}. (**A**) [^{1}H,^{15}N]-WADE-TROSY spectrum of apo PKA-C^{F100A} and bound to bound to ADP, ATPγN, and e ATPγN/PKI$_{5\text{-}24}$. (**B**) Change in chemical shift perturbation (CSP) between PKA-C^{WT} and PKA-C^{F100A} upon binding ATPγN. (**C**) Change in CSP ($\Delta\delta_{WT}$ - $\Delta\delta_{F100A}$) upon binding ATPγN and PKI$_{5\text{-}24}$.

937 **Figure S10 - figure supplement 1: CONCISE plot showing the probability distribution of**
938 **the amide resonances as a function of ligand binding.** The per-residue information is aver-
939 aged into the average principal component (PC) score indicative of the position of each confor-
940 mational state of the kinase along the equilibrium.

Figure S10 - figure supplement 2. Changes of the intermolecular allosteric network in F100A as mapped by correlated chemical shift changes. (A) CHESCA matrix obtained from the amide chemical shifts of PKA-CF100A in the apo, ADP-bound, ATPγN-bound, and ATPγN/PKI$_{5-24}$-bound states. Only correlations with $R_{ij} > 0.98$ are reported. **(B)** Plot of the correlation score vs. residue calculated for PKA-C^{WT} (blue) and PKA-C^{F100A} (black). **(C)** Community CHESCA analysis of and PKA-C^{F100A} (bottom diagonal, black). Only correlations with $R_{A,B} > 0.98$ are shown. (**D**) Community CHESCA matrices of PKA-C^{F100A} and PKA-C^{WT} plotted on their corresponding structures. The size of each node is independent of the number of residues it encompasses, meanwhile the weight of each line indicates the strength of coupling between the individual communities.

957 **Figure 2 – supplementary table 1**. ∆G (kcal/mol) and relative population of ground state and
958 the first 6 excited states in different forms of PKA-C by the RAM simulations.

	GS	ES1	ES2	ES3	ES4	ES5	ES6
Apo	0 (58.0%)	0.38 (30.8%)	1.22 (7.6%)	2.10 (1.8.%)	2.28 (1.3%)	2.85 (0.5%)	8.16* (<1e-4)
Binary	0 (99.4%)	3.11 (0.6%)	5.80 (<1e-4)	6.66 (<1e-4)	7.25 (<1e-4)	8.27 (<1e-4)	8.47 (<1e-4)
Ternary	0 (100.0%)	4.85 (<1e-4)	5.68 (<1e-4)	6.92 (<1e-4)	7.04 (<1e-4)	7.58 (<1e-4)	7.87 (<1e-4)

959 * Numbers in red refer to populations of excited states below 0.5%.

961 <mark>**Supplementary Table 2.**</mark> **Kinetic parameters of Kemptide phosphorylation by PKA-C^{WT} and**
962 **PKA-C^{F100A}.** The K_M and V_{max} values were obtained from a nonlinear least squares analysis of the
963 concentration-dependent initial phosphorylation rates using a standard coupled enzyme activity.
964 Error in k_{cat}/K_M was propagated from the error in K_M and k_{cat}.

	PKA-C^{WT}	PKA-C^{F100A}
V_{max}	0.322 ± 0.005	0.379 ± 0.009
K_M	30 ± 1	42 ±3
k_{cat}	15 ± 1	17 ± 1
k_{cat}/K_M	0.50 ± 0.04	0.41 ± 0.08

967 <mark>**Supplementary Table 3.**</mark> **Changes in enthalpy, entropy, free energy, and dissociation con-**
968 **stant for the binding of nucleotide to PKA-C^{WT} and PKA-C^{F100A}.** All errors were calculated
969 using triplicate measurements. Values for PKA-C^{WT} are re-printed for clarity but were originally
970 published in Walker *et al.* [15]

	K_d (μM)	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)	σ
PKA-C^{WT}	83 ± 8	-5.61 ± 0.06	-3.6 ± 0.1	-2.0 ± 0.1	N/A
PKA-C^{F100A}	73 ± 2	-5.7 ± 0.2	-21 ± 5	7 ± 3	N/A

973 **Supplementary Table 4. Changes in enthalpy, entropy, free energy, and dissociation con-**
974 **stant for the binding of PKI$_{5-24}$ to apo and nucleotide-saturated PKA-C^{WT} and PKA-C^{F100A}.**
975 All errors were calculated using triplicate measurements. The error in σ was propagated from the
976 error in K_d. Values for PKA-C^{WT} are re-printed for clarity but were originally published in Walker *et*
977 *al.* [15].

PKI$_{5-24}$ to apo kinase					
	K_d (μM)	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)	σ
PKA-C^{WT}	17 ± 2	-6.57 ± 0.08	-10.8 ± 0.5	4.2 ± 0.5	N/A
PKA-C^{F100A}	5 ± 1	-7.7 ± 0.1	-17 ± 5	7 ± 3	N/A

PKI$_{5-24}$ to ATPγN-saturated kinase					
	K_d (μM)	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)	σ
PKA-C^{WT}	0.16 ± 0.02	-9.33 ± 0.07	-13.9 ± 0.5	4.6 ± 0.4	106 ± 18
PKA-C^{F100A}	2 ± 1	-7.9 ± 0.3	-17 ± 1	9 ± 1	3 ± 1