Ancient origin of the rod bipolar cell pathway in the vertebrate retina

Ayana M Hellevik[1,*], Philip Mardoum[1,*], Joshua Hahn[2,*], Yvonne Kölsch[3,4], Florence D D'Orazi[1], Sachihiro C. Suzuki[1], Leanne Godinho[5], Owen Lawrence[1], Fred Rieke[6,7], Karthik Shekhar[2,8], Joshua R Sanes[3], Herwig Baier[4], Tom Baden[9,10], Rachel O Wong[1], Takeshi Yoshimatsu[11,12,#]

1, Department of Biological Structure, University of Washington, Seattle, WA 98195, USA; 2, Department of Chemical and Biomolecular Engineering; Helen Wills Neuroscience Institute; Vision Sciences Graduate Program; California Institute of Quantitative Biosciences (QB3), University of California Berkley, Berkeley, CA 94720, USA; 3, Department of Molecular & Cellular Biology and Center for Brain Science, Harvard University, Cambridge, MA 02138, USA; 4, Max Planck Institute for Biological Intelligence, Department Genes – Circuits – Behavior, 82152 Martinsried, Germany; 5, Institute of Neuronal Cell Biology, Technische Universität München, 80802 Munich, Germany; 6, Department of Physiology and Biophysics, University of Washington, Seattle, WA 98195, USA; 7, Vision Science Center, University of Washington, Seattle, WA 98195, USA; 8, Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA; 9, School of Life Sciences, University of Sussex, Brighton, BN1 9QG, UK; 10, Institute of Ophthalmic Research, University of Tübingen, Tübingen, 72076, Germany; 11, Department of Ophthalmology & Visual Sciences, Washington University in St Louis School of Medicine, St Louis, MO 63110, USA; 12, BioRTC, Yobe State University, Damatsuru, Yobe 620101, Nigeria

*These authors contributed equally to this study

#Lead contact: Takeshi Yoshimatsu takeshi@wustl.edu

Keywords

Rod, Rod bipolar cells, Retina, Evolution

Highlights

- Zebrafish have two rod bipolar cell types (RBC1/2).

- Synaptic connectivity of RBC1 resembles that of the mammalian RBCs.

- The primary rod pathway therefore probably evolved more than 400 million years ago.

- The second zebrafish RBC type, RBC2, forms a separate pathway from RBC1.

ABSTRACT

Vertebrates rely on rod photoreceptors for vision in low-light conditions[1]. Mammals have a specialized downstream circuit for rod signaling called the primary rod pathway, which comprises specific cell types and wiring patterns that are thought to be unique to this lineage[2–6]. Thus, it has been long assumed that the primary rod pathway evolved in mammals[3,5–7]. Here, we challenge this view by demonstrating that the mammalian primary rod pathway is conserved in zebrafish, which diverged from extant mammals ~400 million years ago. Using single-cell RNA-sequencing, we identified two bipolar cell (BC) types in zebrafish that are related to mammalian rod BCs (RBCs) of the primary rod pathway. By combining electrophysiology, histology, and ultrastructural reconstruction of the zebrafish RBCs, we found that, like mammalian RBCs[8], both zebrafish RBC types connect with all rods and red-cones in their dendritic territory, and provide output largely onto amacrine cells. The wiring pattern of the amacrine cells post-synaptic to one RBC type is strikingly similar to that of mammalian RBCs, suggesting that the cell types and circuit design of the primary rod pathway have emerged before the divergence of teleost fish and amniotes (mammals, birds, reptiles). The second RBC type in zebrafish, which forms separate pathways from the first RBC type, is either lost in mammals or emerged in fish to serve yet unknown roles.

INTRODUCTION

Rod photoreceptors of the vertebrate retina are capable of detecting very dim light, down to individual photons[9–14]. The mammalian cell types and circuitry that convey rod-driven signals, called the primary rod pathway, were identified and defined in the cat retina about 50 years ago[15]. Since its initial characterization, this rod pathway has been examined extensively in many mammalian species including in humans, where it consistently uses homologous cell types and connectivity patterns: Rod bipolar cells (RBCs), as well as A2 and A17 amacrine cells[3,5,6,16,17]. The RBC is a molecularly, structurally and functionally distinct retinal bipolar cell type that receives input from all rods within its dendritic field and is predominantly driven by rods[8,18–21]. All other bipolar cell types receive most of their photoreceptor input from cones, which operate in daylight conditions. In contrast to mammals, the bipolar cells that have been characterized in non-mammals lack clear distinctions regarding the ratio of rod and cone inputs[14,22]. Thus, it is not surprising that the RBC is thought to be unique to mammals, a notion that has led to the prevailing view that the rod pathway evolved separately in this class[3,5–7]. However, the molecular, structural and functional signatures that together define bipolar cell types in mammals are largely unknown in non-mammals, so it remains unclear whether the signatures characteristic of the mammalian RBC and its downstream pathway are present in non-mammals. To address this issue, we focused on zebrafish, a species that allows transgenic labeling of neuronal populations, to analyze single-cell transcriptomics, histology, physiology and circuit reconstructions of genetically-defined retinal bipolar cells.

There are more than a dozen BC types in mammals that are diverse in morphology, connectivity and molecular profiles[23–27], but can be classified into two main groups: ON BCs that depolarize and OFF BCs that hyperpolarize in response to increases in luminance[28]. RBCs are one type of ON BCs that are distinct in many ways from all other BCs, which mainly connect with cones and are called cone BCs (CBCs) here. Transcriptionally, mammalian RBCs can be distinguished from

CBCs by the expression of protein kinase C-alpha (PKCα)[29]. Morphologically, the axon terminals

of RBCs are generally larger than those of CBCs and end in the innermost layer of the inner

plexiform layer (IPL). The synaptic arrangement of the RBC axons differs from the common

synaptic arrangement of most cone bipolar cells. Whereas CBC axons directly synapse onto the

retinal output neurons, retinal ganglion cells (RGCs), along with a plethora of amacrine cells (ACs),

RBCs predominately form a 'dyadic' synapse with two types of inhibitory amacrine cells, small

field A2 (or A-II) and large-field A17 ACs[15,18,19,30]. The A17 AC almost exclusively makes reciprocal

feedback synapses onto RBC axon terminals[31]. In contrast, A2 ACs receive numerous synapses

from RBCs (~40 synapses per RBC in mice), but do not provide feedback onto the RBCs[18,32].

These RBC to A2 AC synapses are the critical sites for the amplification and gain control of rod

signals[33–35]. Rod signals are eventually relayed to RGCs by connections from A2 ACs on CBCs,

which split rod signals into ON and OFF channels via sign-conserving gap junctions with ON

CBCs and inhibitory synapses with OFF CBCs[15,36,37].

Here, by analyzing single-cell transcriptomic profiles of zebrafish BCs, we discovered two BC

types, RBC1 and RBC2, with molecular signatures similar to those of mammalian RBCs. Using

transgenic zebrafish lines that express a fluorescent protein in RBC1 or RBC2 cells, we identified

the inputs and outputs of RBC1 and RBC2. We found that both zebrafish RBC types connect with

all rods and red cones (or longwave-length sensitive, LWS, cones) inside their dendritic fields.

We further reconstructed the downstream circuits of both BC types using serial block-face

electron microscopy and found that RBC1 predominantly synapses onto three morphological

types of ACs. The circuit diagrams and synaptic arrangements of two of the ACs closely resemble

those of the mammalian A2 and A17 ACs. In contrast, RBC2 mainly connects to a different set of

ACs, which does not include A2-like ACs. These results suggest that (i) zebrafish possess two

separate pathways for processing rod signals, and that (ii) one of these is similar to the rod –>

RBC –> A2 AC –> CBC –> RGC pathway found in mammals. We conclude that the primary rod

pathway emerged >400 million years ago, before the divergence of teleosts and mammals in the

Devonian.

RESULTS

Two zebrafish BC types are transcriptionally analogous to the mammalian RBCs

We first determined the transcriptional similarity between each zebrafish BC type and the

mammalian RBCs by using single-cell RNA-sequencing (scRNA-seq) in adult zebrafish. BCs

were isolated using a fluorescent marker in the *Tg(vsx1:GFP)^nns5* transgenic line, in which all BCs

express GFP[38] (Fig. 1a). Clustering analysis of 19492 high-quality single cell transcriptomes

identified 23 molecularly distinct BC clusters (Fig. 1b,c). To identify the clusters most similar to

mammalian RBCs, we performed a hierarchical clustering analysis based on average

transcriptomic profiles (Fig. 1d) and combined this with the expression patterns of marker genes

identified in mice to tentatively annotate each cluster as ON CBC, OFF CBC or RBC (Fig. 1e; [27]).

In mice, RBCs are clearly separated from CBCs at the first dendrogram bifurcation[27]. Similarly,

the first dendrogram bifurcation separates two BCs from the other BCs in zebrafish. In contrast to

mice, however, the zebrafish RBC clade contained two molecularly distinct clusters 14 (c14) and

19 (c19) (Fig. 1d). We observed that *prkca* (the gene encoding PKCα), a common marker of

mammalian RBCs, is only highly expressed in c14. However, both c14 and c19 specifically

express *gramd1b*, which is an RBC-specific marker in mice. In addition to these genetic

signatures similar to mammalian RBCs, both c14 and c19 clusters express neurotransmitter

receptor, *grm6a* and *grm6b,* and it's downstream signaling molecules, *trpm1a, trpm1b*, *nyx* and

rgs11, which are essential for mediating rod inputs in mammals[39] (Fig. S1). Therefore, we

hypothesized that zebrafish, unlike mice, may possess two RBC types, which we call RBC1 (c14)

and RBC2 (c19).

Figure 1

Figure 1. Comparison of single-cell gene expressions identified two possible rod bipolar cells in zebrafish

a, Schematic representation of retinal circuits (left) and an image of a retinal slice from Tg(vsx1:GFP)^nns5 transgenic adult zebrafish (right). GFP expression in all bipolar cells (BCs). Nuclei was stained by DAPI. PR: photoreceptor, HC: horizontal cell, AC: amacrine cell, RGC: retinal ganglion cell. *b*, 2D visualization of single-cell clusters using Uniform Manifold Approximation (UMAP)[40]. Individual points correspond to single cells colored according to cluster identity. *c*, Marker genes for each cluster. *d*, Agglomerative hierarchical clustering of average gene signatures of clusters using the correlation metric and complete linkage. BC subclasses (colors) were assigned based on the known marker expressions shown in *e*. *e*, Gene expression patterns of known BC subclass markers in BC clusters. The size of each circle depicts the percentage of cells in the cluster in which the marker was detected (≥1 UMI), and its contrast depicts the scaled average expression level of cells within the cluster in *c,e*. Data for mouse is from Shekhar K, et al., 2016, Cell.

Figure S1

Figure S1. Expression patterns of the identified marker genes in BC clusters

a,b, The size of each circle depicts the percentage of cells in the cluster in which the marker was detected (≥1 UMI), and its grey scale depicts the scaled average expression level of cells within the cluster

RBC1 and RBC2 morphologies resemble mammalian RBCs

We next determined the morphological similarities of RBC1 and RBC2 with mammalian RBCs. In

mammals, RBC axons arborize in the innermost layer of the IPL[41]. By screening our zebrafish

transgenic lines, we identified two lines, *Tg(vsx1:memCerulean)^q19* (*vsx1:memCer*) and

Tg(vsx2:memCerulean)^wst01 (*vsx2:memCer*), that each label BCs with axon terminals in the

innermost layer of the IPL (Fig. 2). Fluorescent *in situ* hybridization for the identified gene markers,

s100a10b and *uts1,* which are selectively expressed by RBC1 and RBC2 (Fig. 1c), revealed that

vsx1:memCer and *vsx2:memCer* label RBC1 and RBC2, respectively (Fig. 2a,b). We also

observed that dendritic arbors of both RBC1 and RBC2 cover the retina in a non-overlapping

manner, an arrangement called 'tiling' that is considered a hallmark of a BC type (Fig. 2e,f)[42].

Therefore, both RBC1 and RBC2 represent single bipolar types that transcriptionally and

morphologically resemble mammalian RBCs.

We observed slight variations in morphology and molecular expression between RBC1 and RBC2.

The axon terminal of RBC1 is relatively spherical, similar to mammalian RBCs, in contrast to the

'flat-footed' axonal ending of RBC2 (Fig. 2c,d). RBC1 were immunoreactive for PKCα (Fig. 2c),

whereas RBC2 were not (Fig. 2d), consistent with the difference in their *prkca* expression (Fig.

1d). In addition, their abundance differed: RBC1s were more densely packed than RBC2s

(*p=0.0052*, Mann-Whitney two-tailed U test) (Fig. 2g,i). This difference in the densities is unlikely

due to regional variations as both RBCs are present in the dorsal and ventral-temporal retina at

similar densities (Fig. 2g,i and Fig. S2). The dendritic field sizes of the two RBC types were

inversely related to their cell density, consistent with their tiling arrangement (Fig. 2g-j).

Figure 2

Figure 2. Transgenic labeling of cluster 14 and 19 revealed morphological features of these BCs

a,b, En face view of retinal flat mount at the inner nucleus layer level. Cerulean fluorescent expression (colored yellow) transgenic lines, Tg(vsx1:memCerulean)^{q19} (vsx1:memCer) in *a* and

Tg(vsx2:memCerulean)[wst01] *(vsx2:memCer) in* **b**. *vsx1:memCer and vsx2:memCer BCs are*

positive for cluster specific genes, s100a10b and uts1, respectively, which are detected using in

situ hybridization chain reaction [43]. **c,d**, *Side views of the labeled cells and the distribution patterns*

*of their axon terminals in en face views of retinal flat mounts for RBC1 (**c**) and RBC2 (**d**) BCs.*

Immunolabeling for PKCα is in magenta. IPL: inner plexiform layer. Note that not all PKC-

immunoreactive cells are apparent in this image of the vsx1:memCer line, due to the incomplete

labeling of this line. **e,f**, *Dendritic tiling of RBC1 (**e**) and RBC2 (**f**) in en face view of retinal flat*

mounts at the outer plexiform layer level. Dendritic territories are marked by the red boundaries.

g-j, *Mean cell densities of RBC1 (**g**, n=3 and 4 for D and VT, respectively) and RBC2 (**I**, n=3 for*

both D and VT) BCs in different regions of the retina. Box and violin plots of dendritic field sizes

of RBC1 (h, n=37 and 33 or D and VT, respectively) and RBC2 (j, n=40 and 41 or D and VT,

respectively) BCs. White filled circles are medians. Grey circles indicate individual cells. D: dorsal,

VT: ventrotemporal.

Figure S2

a

vsx1:memCer

Dorsal   Ventro-temporal

10 µm

b

vsx2:memCer

Dorsal   Ventro-temporal

10 µm

Figure S2. Dendritic tiling of RBC1 and RBC2 BCs across the retina

Confocal images of retinal flat mount at outer plexiform layer level from Tg(vsx1:memCerulean)[q19] *(vsx1:memCer) and Tg(vsx2:memCerulean)*[wst01] *(vsx2:memCer). Note that the vsx1:memCer line occasionally labels OFF BCs. These BCs were distinguished by tracing the cells to the axon terminals in the confocal image volumes.*

Both RBC1 and RBC2 connect with all rods and red-cones in their dendritic territory

If RBC1 and RBC2 are authentic RBCs, they should synapse preferentially with rods. Using 4C12 antibodies to label rods, we found that the majority of the dendritic tips of both RBC types (RBC1: 84±3.9%, RBC2: 78±2.9%) contacted with rod spherules (Fig. 3a-d). We also found that some dendritic tips were not associated with rods (Fig. 3a-d). Using transgenic lines to label specific cone types, we identified that dendritic tips of both BC types contacted red-cones (or long-wavelength sensitive cones), labeled in the *Tg(trb2:tdtomato)^{q22}* line (Fig. 3a-e,g). Furthermore, both types connected with nearly all rods and red cones within their dendritic fields (Fig. 3f,h). We did not observe any dendritic tips that were not associated with either rods or red cones (Fig. 3e,g), indicating that they receive few if any inputs from the other cone types, which include green, blue, and violet cones[44,45]. Therefore, both RBC1 and 2 receive predominant rod input and share specificity for red cones among cones (Fig. 3e,g).

Interestingly, the dendritic tips of RBC1 and RBC2 terminating in rod spherules differed in structure (Fig. 3c,d). Specifically, the dendrites of RBC1 invaginating rod spherules appeared to form a horseshoe or 'doughnut' ending, whereas those of RBC2 ended in a simpler arrangement (Fig. 3c,d). The larger surface area of RBC1's dendritic tips at rod terminals may increase the sensitivity to rod inputs in this BC type compared to RBC2. We also observed differences in the distal axonal boutons of RBC1 and RBC2 (Fig. 3i,j). While axons of both types terminate close to the ganglion cells in the IPL, RBC2 axons have a bouton in the OFF layer of the IPL, next to the boundary with the ON layer (Fig. 3j). These distal boutons are likely pre-synaptic sites as they contain the pre-synaptic protein, Ribeye (Fig. 3j). These differences in the dendritic tip and axon bouton shapes between RBC1 and RBC2 suggest that, while both BCs receive input from the same combination of photoreceptor types, they may serve distinct visual functions.

Figure 3

Figure 3. RBC1 and RBC2 connect to rods and red cones but differ in dendritic and axonal

synaptic arrangements

***a,b**, Dendritic tips invaginating the rod and red cone axon terminals, visualized in retinal slices*

from   (a)   Tg(vsx1:memCerulean:trb2:tdtomato)^{q19,q22} and   (b)

Tg(vsx2:memCerulean:trb2:tdtomato)^{wst01,q22} adult zebrafish. Rods were immunolabeled using

*4C12 antibody. **c,d**, Doughnut and simple dendritic tip structures at rod terminals (arrow heads)*

*in RBC1 and RBC2, respectively. **e-h**, Box and violin plots of RBC1 (**e,f**, n=7) and RBC2 (**g,h**,*

*n=8) connectivity with photoreceptors. **i,j**, Distribution of ribbon synapses in the RBC1 (**i**) and 2*

*(**j**) axons. Ribbons were immunolabeled by anti-ribeye antibody. Ribeye signals outside the axons*

were digitally masked out in the right two images. RBC2 axon harbors a ribbon containing distal

bouton in the OFF layer (arrow head).

RBC1 receives rod inputs via mGluR6 receptors

We next asked whether zebrafish RBCs receive functional rod input via the metabotropic

glutamate receptor mGluR6 as seen in mammalian RBCs. We first investigated the expression of

mGluR6 in RBC1 and RBC2 dendritic tips at rod spherules. Super-resolution imaging of mGluR6

immunolabeling in *vsx1:memCerulean and vsx2:memCerulean* retinas showed that the dendritic

tips of RBC1, but not RBC2, robustly overlapped with mGluR6 immunoreactivity at contacts with

rod spherules (Fig. 4a,b). These findings are consistent with the transcriptional profiles, which

showed that RBC1 expresses higher mRNA levels of *grm6a* and *grm6b*, which encodes mGluR6,

than RBC2 (Fig. S1).

We then used electrophysiological recordings to ask whether mGluR6 mediates rod input to the

zebrafish RBCs. We prepared retinal wholemounts that preserve synaptic connections in the

outer retina, and performed whole-cell patch-clamp recordings on the axon terminals of RBC1

and RBC2 (Fig. 4c). Both RBC1 (n = 10) and RBC2 cells (n = 3) exhibited ON responses to a

cone-activating flash (red LED), confirming the successful patch-clamp recordings of light

responses in these BCs and demonstrating that both cell types are ON cells (Fig. 4d), consistent

with the position of their axonal arbors (Fig. 2c,d).

Although measuring rod-mediated responses from RBC2 was infeasible for technical reasons

(see Methods), we were successful in recording rod responses from RBC1. We were therefore

able to ask whether these responses were mediated by mGluR6. We presented rod-isolating dim

blue flashes (10 ms) before and after introducing the mGluR6 receptor agonist 6-(2-

aminopropyl)benzofuran (APB) to the perfusion solution. To isolate excitatory inputs to the cell,

all recordings were performed near the reversal potential for chloride-mediated conductances (~-

60 mV) and in the presence of inhibitory receptor blockers, gabazine, strychnine, and TPMPA

((1,2,5,6-Tetrahydropyridin-4-yl)methylphosphinic acid). Our results showed that nearly all rod

inputs were blocked in the presence of APB, indicating that, like mammalian RBCs, mGluR6

mediates rod input to RBC1 (Fig. 4e).

Both RBCs primarily synapse onto amacrine cells

To determine the synaptic targets of RBC1 and RBC2, we reconstructed their connectomes using

serial block face scanning electron microscopy (SBFSEM). In the reconstructions, we observed

an array of large BC axon terminals in the innermost layer of the IPL, which are characteristics of

RBC1 and RBC2 axons (Fig. S3a,b). To confirm that these large axon terminals belong to RBC1

and RBC2, we reconstructed dendrites of some of these BCs (Fig. 5a). Consistent with our

observations in light microscopic experiments (Fig. 3a-d), the large axon BCs predominantly

connect with rods.

Figure 4

Figure 4. Rod input to RBC1 is mediated by mGluR6 receptors.

*a,b, Confocal microscopy images of mGluR6 immunolabeling at rod terminals. **c,d**, Colocalization of mGluR6 and RBC1 (**c**) and RBC2 (**d**) dendritic tips at rod terminals visualized by structured illumination microcopy. **e**, Whole-cell patch clamping of a RBC1 axon terminal, visualized by dye-filling Alexa Fluor 594). **f**, Voltage responses of RBC1 and RBC2 after a cone activating light flash (arrow heads). **g**, Population data of RBC1 responses to rod activating light flashes with and without the group III metabotropic glutamate receptor agonist, APB (6-(2-aminopropyl)benzofuran). Filled circles: mean; error bars, S.D.; open circles, individual cells. Traces on the right are an example of the cell's light evoked response before and during APB bath application. Inhibitory neurotransmitter receptors were blocked (inh lock) by a bath application of gabazine, strychnine, and TPMPA ((1,2,5,6-Tetrahydropyridin-4-yl)methylphosphinic acid).*

We then reconstructed all the large axons in the SBFSEM image volume. To distinguish RBC2

from RBC1, we used the ribbon containing axonal distal bouton in the OFF layer as a proxy for

RBC2 (Fig. 3i,j and Fig. 5b). These reconstructions revealed the regular mosaic arrangements of

both presumed RBC1 and RBC2 (Fig. 5c), indicating that we identified most, if not all, RBC1 and

RBC2 in the EM volume. Using this criterion, we also verified that dendritic tips of RBC1 are

doughnut shaped whereas those of RBC2 ended in a simple tip within the rod spherule, consistent

with our light microscopy data (Fig. 3c,d, 5a).

Figure 5

Figure 5. Identification of RBC1 and RBC2 in a SBFSEM volume.

a, Reconstructions of a RBC1 and a RBC2, and zoomed-in images of their dendritic tips at rod

and cone terminals. Ribbons in the rod and cones are painted red. b, Ribbon synapse distributions

in a RBC1 and a RBC2. The locations of ribbon synapses are marked in red. Arrow heads indicate

the locations of example ribbon synapses (arrows) shown in the insets. c, Reconstruction of all

RBC1s and RBC2s in the EM volume. Postsynaptic neurons of a centrally located RBC1 (open

arrow head) and RBC2 (closed arrow head) were reconstructed in Fig. 6, 7, and S4-8.

We then focused on one RBC1 and one RBC2 in the central area of the volume and traced all of

their post-synaptic neuronal processes (Fig. S3c,d). Amacrine cells (ACs), unlike reginal ganglion

cells (RGCs), make output synapses within the retina. Hence, we identified the neuronal class

(e.g. AC or RGC) for the majority (28/32) of the RBC1 postsynaptic processes and over the half

(18/31) of the RBC2 postsynaptic processes based on the presence or absence of presynaptic

structures. We found that both RBC1 and RBC2 predominantly synapse onto ACs (Fig. 6a,f). The

majority of the postsynaptic processes received 4 or fewer ribbon synapses from one RBC1 or

RBC2, with an exception of one process, which received 14 inputs from one RBC1 (Fig. 6b,g).

Figure S3

Figure S3. Identification of RBC1 and RBC2 postsynaptic neurons in SBFSEM volume

a, A partial image of an example SEM image of an adult zebrafish retina. OPL (outer plexiform layer), INL (inner nuclear layer), IPL (inner plexiform layer), GCL (ganglion cell

*layer). **b**, Magnified image of the region within the black box in **a** at the bottom layer of the IPL.*

*Characteristic large bipolar cell axons are painted in light yellow and green. **c,d,** Traces of*

neuronal processes and the location of somas of cells that are post-synaptic to RBC1 and RBC2

cells. Individual cells were color coded. IPL: inner plexiform layer.

We further morphologically classified the postsynaptic ACs that we traced, comprising 14 cells for

RBC1 (Fig. S4-6) and 10 cells for RBC2 (Fig. S7,8), respectively. Most ACs extended their

dendrites within a single sublamina in the IPL (Fig. 6c,h). Among these mono-stratifying ACs for

RBC1, we identified two groups based on their dendritic stratification depth within the IPL (Fig.

6d,e). These two groups of ACs differed in their synaptic arrangement. ACs stratifying in the lower

layer formed reciprocal synapses (RS) – a synaptic arrangement that includes both input and

output synapses with a BC axon - with RBC1, whereas ACs stratifying in the upper layer did not

(Fig. 6d,e, Fig. S4,5). For RBC2, all but one AC (1/8) formed local reciprocal synapses (Fig. 6i,

Fig. S7,8). In addition, RBC1 formed an exceptionally high number of ribbon connections with one

bi-stratifying AC (marked in red in Fig. 6b,c, morphology in Fig. 7d,e and Fig. S6). In contrast,

RBC2 does not have a post-synaptic partner with extensive synapses.

Taken together, these results demonstrated that RBC1 allocated the majority (91%) of synaptic

outputs to 3 types of AC: 14% to non-RS ACs, 37% to RS ACs, and 40% to one bi-stratifying ACs.

In contrast, the RBC2 we reconstructed synapsed primarily onto a mono-stratifying AC type (68%).

Among 24 ACs that we traced throughout the volume, only 5 were shared between RBC1 and

RBC2 (Fig. S4-8). Therefore, the downstream circuits of these BC types are largely separate at

least at the AC level.

Figure 6

Figure 6. Identification RBC1 and RBC2 post-synaptic neuron types

a-d, Quantification of morphological parameters of neurons postsynaptic to one of the RBC1s in

the EM volume (marked by an open arrow head in Fig. 5c). One postsynaptic neuron contained

*an exceptionally higher number of synapses (14) with the RBC1 (marked in red in **b** and **c**).*

Dendritic stratification is normalized to 0 and 1 at the lower and upper ends of the RBC1 axon

*terminals, respectively in **d**. **e,** Mono-stratifying ACs with (red) or without (blue) reciprocal*

synapses with RBC1s in the volume. The axon of the presynaptic RBC1 is also shown in the side

*view. Individual cells were color coded. **f-i**, Quantification of morphological parameters for*

neurons postsynaptic to one of the RBC2s in the EM volume (marked by a closed arrow head in

Fig. 5c). AC: amacrine cells, RGC: retinal ganglion cells.

The wiring diagram of RBC1 resembles that of the mammalian primary RBC pathway

Finally, we compared the targets of RBC1 and RBC2 with those known to comprise the primary

rod pathway in the mammalian retina. Mammalian RBCs synapse onto a mono-stratifed AC type

called A17 and a bi-stratified AC type called A2[3,5–7]. A17 ACs are wide field ACs that synapse

exclusively with RBCs and form reciprocal synapses with them[31,46], whereas A2 ACs are narrow

field ACs that receive numerous (~40) synapses from RBCs but do not form reciprocal synapses

with them[18,30]. Instead, A2 ACs form gap-junctions with ON CBCs through dendrites in the ON

layer and output synapses onto OFF CBCs through the bouton structures in the OFF layer called

lobular appendages[32].

We found that zebrafish RBC1s synapse onto both wide field ACs with reciprocal synapses (the

RS ACs)(Fig. 6,7a), resembling mammalian A17s (Fig. 7b), and a narrow field bi-stratifying ACs

with extensive synaptic connections, resembling mammalian A2s (Fig. 6,7c-f). By marking

synaptic sites of RS ACs throughout their dendrites, we found that RS ACs are dedicated to the

RBC1 pathway; synapsing predominantly (both input and output) with RBC1 and to a lesser extent

with RBC2 (Fig. S4), with no synapses with other BC types (n=7 cells). This synaptic specificity

and the reciprocal synapse arrangement in RS ACs mirror those of mammalian A17 ACs[31].

Figure 7

a Zebrafish RS AC

En face view   *Side view*

RBC1

10 μm

b Mammalian A17 AC

Side view

20 μm

c

RBC1
Input from RBC1

5 μm

d

En face view

10 μm

e Zebrafish bi-stratifying AC

Side view

IPL

Distribution

Output to all BCs
Input from all BCs
Non-synaptic contacts

10 μm

f Mammalian A2 AC

Side view

Distribution

Output to all cells
Input from rod BCs
Gap-junction

10 μm

g Zebrafish

Cone
Rod
non-RS
RS
A2
R1
CB CB
R2
RS

h Mammals

RB
A2
A17
CB CB

Figure 7. Circuit diagram of RBC1 is similar to mammalian RBC pathway

a, *An example monostratified amacrine cell (RS AC) classified in Fig. 6e that formed of reciprocal*

synapse with RBC1. ***b,*** *A mouse A17 amacrine cells (taken from [47]) (**b**).* ***c,d,*** *A2-like ACs that are*

postsynaptic to two neighboring RBC1s. ***e,d,*** *Locations and distributions of synaptic sites and*

*non-synaptic contacts with BCs in zebrafish bi-stratifying AC (**e**) and in rabbit A2 ACs (taken from*

*[32]) (**f**). Note that synapses or non-synaptic contacts with AC and RGC are not included in (**e**), and*

*inputs from CBC are not included in (**f**).* ***g,h,*** *Schematic diagrams of zebrafish (**g**) and mammalian*

*(**h**) RBC pathways. Mouse A17 and rabbit A2 images are from [47] and [32], respectively, used with*

permissions. Data for the distributions of synaptic sites within mammalian A2 ACs across the

inner plexiform layer (IPL) are taken from[48].

Next, we examined the synaptic arrangements of RBC1 with the A2-like ACs. First, we confirmed

that this type of AC is common to other RBC1s. By tracing the postsynaptic processes of

neighboring RBC1, we found another A2-like AC, which received a high number of ribbon inputs

from the neighboring RBC1 (Fig. 7c,d). We marked the locations of synapses with all BCs for

those two ACs (Fig. 7e). Gap-junctions are too small to be resolved in our SBFSEM images, but

as a proxy, we marked non-synaptic contacts (Fig. 7e). These revealed a striking similarity in the

distribution patterns of synapses and (potential) gap-junction sites across the IPL layers between

this AC type in zebrafish and mammalian A2 ACs (Fig. 7e,f), including the bouton structures in

the OFF layer that contain large presynaptic sites[48–50]. Taken together, we conclude that the circuit

diagram among mammalian RBC, A2, and A17 ACs are conserved in the zebrafish RBC1

pathway (Fig. 7g,h).

In contrast to the targets of RBC1, RBC2 formed synapses exclusively with wide-field ACs (Fig.

S7,8) and lack a synaptic partner with extensive synapses. Thus, RBC2 participates in a circuit

that differs from that of mammalian RBCs.

DISCUSSION

By combining scRNA-seq, electrophysiology, and light and electron microscopy circuit

reconstructions, we demonstrated that RBC1 shares many features with the mammalian primary

rod pathway (Fig. 7g,h), implying that the conserved rod pathway is evolutionarily ancient.

The number of BC types

In this study, we found 23 molecular types in adult zebrafish BCs. However, a previous

morphological characterization of zebrafish BCs, based on their dendritic connectivity with

photoreceptors and axon stratifications, identified 32 anatomical types[51]. The discrepancy in the

number of BC types between morphological and transcriptional characterizations may arise from

the regional specializations that have been documented in the larval zebrafish retina[52–55]. In any

event, it is clear that the adult zebrafish retina contains at least 23 BC types.

This number of molecular types of bipolar cells (BCs) in zebrafish, as identified in this study, is

higher than that found in mammals investigated to date (14-15 across mammals)[26,27,29,56], but

similar to that found in chick retina (22 molecular and 15 morphological BC types)[57,58]. The higher

number of BC types in zebrafish and chicken is not surprising, given that these species have

higher numbers of photoreceptor types: 5 in zebrafish and 7 in chicks, compared to >=3 in

mammals[59,60]. We demonstrate here one source of the increase: a single type of BC carries most

of the input from rods in mammals, whereas zebrafish has two RBC types.

The number of RBC types

Previous morphological characterization of zebrafish BCs found only one BC type that connects

rods and red cones. Axons of these BCs terminate in the innermost layer of the IPL[51]. We

speculate that this type actually includes both RBC1 and RBC2, which were combined owing to

their striking morphological similarity (Fig. 2,3). Consistent with this hypothesis, studies in goldfish

have reported two morphologically distinct "mixed" BC types that receive dominant inputs from

rods[22]. They have large axon terminals at the bottom of the IPL, but the axon of one mixed BC

type contains a smaller axonal distal bouton in the OFF layer, similar to RBC2[61]. Immunostaining

for PKC only labels mixed BCs without an axonal distal bouton, similar to RBC1[62]. The presence

of these features in goldfish suggests that RBC1 and RBC2 are conserved among teleost fish.

Much less is known about RBCs in other non-mammalian vertebrate species. In salamander, one

type of mixed ON BCs exhibit sensitivity close to that of rods[14,63]. They terminate their axons at

the bottom of the IPL, similar to teleosts and mammals, but it is unknown whether they express

the RBC marker PKC. Furthermore, the anatomical connections of BCs with rods are not yet

comprehensively studied in salamander. In birds, PKC labels some ON layer stratifying BC types

strongly[64,65]. Single-cell RNA-sequencing of chick BCs revealed a BC type that is

transcriptomically similar to the mammalian RBCs[58]. However, the physiological properties and

connections of these BCs are unknown. Moreover, a connectomic survey of chick BC types failed

to identify BC cells that connect with all rods in their dendritic field[57]. Unlike the species mentioned

above, the presumed rod bipolar cells, which have light sensitivity close to that of rods, in sea

lamprey are OFF type[66]. However, their connectivity with rods is unknown. Thus, it remains

unclear whether RBC2 orthologs are present in species other than zebrafish.

In mammals, morphological, molecular, and functional studies have identified only a single RBC

type[23,26,27,67]. Therefore, we speculate that either RBC2 evolved after the divergence between

teleost fish and mammals, or mammals lost this pathway.

Roles of cone inputs in RBCs

Cone inputs onto rod-dominant mixed BCs have been proposed to broaden the dynamic range of

light intensities to which they can respond[68]. Consistent with this idea, we found that both RBC1

and RBC2 are selective for red cones, which, with their broad spectral sensitivity, are suited for

encoding achromatic luminance information[69]. Because rods evolved from cones[1], we speculate

that RBCs may have emerged from red-cone specific CBCs. The red cone selectivity is also

conserved in at least one of the mixed rod dominant BC types in goldfish[70]. Although cone

selectivity is unknown in Salamander rod-driven mixed BCs, their spectral sensitivity curve is

broader at longer wavelengths than that of rods, indicating that they may connect to red cones[14].

Electrophysiological recording from rod-driven BCs in *Giant Danio*, a teleost fish species, showed

that rod and cone inputs onto rod-dominant BCs are mediated by different mechanisms: rod inputs

through mGluR6, whereas cone inputs through both mGluR6 and EAAT (excitatory amino acid

transporter)[68]. In this BC type, mGluR6 and EAAT-mediated inputs suppress each other, likely to

allow this cell to respond to both rod and cone dynamic ranges[68]. Electrophysiological recordings

in zebrafish found that some of ON BCs responded to glutamate via both mGluR6 and a

glutamate-gated chloride conductance increase mechanism, which is likely through EAATs[71].

However, the nature of EAAT contributions for cone responses in RBC1 and RBC2 is unknown.

While the study in *Giant Danio* suggest that mixed inputs expand the dynamic range of rod-

dominant BCs, electrophysiological recordings in goldfish and salamander have found that the

dynamic range of rod dominant BCs is similar to that of rods and that cone contributions to the

light response are small[14,72]. Therefore, the roles of red-cone inputs to RBCs remain to be

determined.

Unifying mixed BCs and RBCs

In mammals, it was initially thought that RBCs exclusively synapse with rods[73]. However, several

recent studies have demonstrated convincingly that RBCs also receive synapses from cones, at

least in mice, rabbits, and primates[8,73–75]. Indeed, mouse RBCs contact the majority of M-cones

(~80%), which are analog of zebrafish red-cone, in their dendritic territories[8]. RBCs were likely

thought to be exclusive to rods because of the high ratio of rods to cones in the outer nuclear

layer in mice and rabbits[76,77]. As a consequence, only a few cones, generally three or fewer,

synapse on a mouse RBC, compared to inputs from ~35 rods[8,75].

The rod-driven BCs in non-mammals are classically called "mixed" BCs because they connect

with both rods and cones. However, as argued above, this mixed connectivity is conserved in the

mammalian RBCs. Moreover, the dendritic specificity and connectivity of RBCs are conserved in

mice and zebrafish. In both species, RBCs connect with all rods and the majority of red-cones (or

M-cones) in their dendritic fields (Fig. 3f,h). Therefore, converging rod and red-cone inputs is likely

a conserved feature of RBCs in all vertebrates. This leads us to propose that the non-mammalian

mixed BCs and the mammalian RBCs represent a single class of neurons, RBCs. Finally, taken

together with the striking similarity in the downstream circuitry of RBCs between zebrafish and

mammals, we conclude that zebrafish RBC1 is transcriptomically, anatomically, and functionally

equivalent of mammalian RBC and that they share the same evolutionary origin.

ACKNOWLEDGEMENTS

We thank the Vision Core at the University of Washington for processing zebrafish retina samples

and acquiring serial images for SBF-SEM. We thank Rachael N. Swanstrom for helping with cell

tracing of the EM volume. Funding was provided by the MCSA fellowship ("ColourFish" 748716)

from the EU Horizon 2020 to TY, the NIH EY14358 to ROW, U01MH105960 and R01 EY022073

to JRS, EY01730 (Vision Core grant, P.I. M. Neitz) Developmental Biology Training Grant

HD07183 to FDD, The Wellcome Trust 220277/Z20/Z to TB, European Research Council ERC-

StG 677687 to TB, UKRI BBSRC, BB/R014817/1 and BB/W013509/1 to TB, Leverhulme Trust

PLP-2017-005; RPG-2021-026 to TB, NIH grant R00EY028625 to KS, Hellmann Foundation

Fellowship to KS, travel grant from Graduate School in Systemic Neuroscience to YK, and Max

Planck Society to YK and HB.

AUTHOR CONTRIBUTIONS

TY, ROW, AMH, PM, JH, and YK designed the study, with input from KS, JRS, HB, and TB. YK

performed single-cell RNA-seq, under the supervision of HB and JRS with guidance from KS. JH

processed and analyzed the data with guidance from KS; TY and SCS generated new plasmids;

TY and FDD generated novel lines; TY performed experiments, collected and analyzed the data

for light microscopy with help from OL; PM performed whole-cell patch recordings; FDD prepared

the sample for SBF-SEM; AMH, OL, and TY traced the EM images; TY analyzed the EM data;

TY wrote the manuscript with inputs from all authors.

DECLARATION OF INTERESTS

506   The authors declare no competing interests.

MATERIALS AND METHODS

Animals

All procedures were performed in accordance with the University of Washington Institutional Animal Care and Use Committee guidelines, the Harvard University/Faculty of Arts & Sciences Standing Committee on the Use of Animals in Research and Teaching (IACUC), and the UK Animals (Scientific Procedures) act 1986 and approved by the animal welfare committee of the University of Sussex. For all experiments, we used adult zebrafish (age 6-18 months) of either sex that were kept at 28°C in a room with a normal 14/10 light cycles.

The following previously published transgenic lines were used: *Tg(vsx1:GFP)^nns5* [78], *Tg(vsx1:memCerulean)^q19* [79], Tg(*trb2:tdtomato*)^q22 [80]. In the larva *Tg(vsx1:memCerulean)^q19* labels a subpopulation of OFF layer stratifying BCs[79]. In adults, while OFF stratifying BCs are still weakly labeled, Cerulean is now strongly expressed in RBC1 (Fig. 2a,c). In addition, *Tg(vsx2:memCerulean)^wst01* line was generated by injecting pBH-vsx2-memCerulean-pA plasmid into single-cell stage eggs. Plasmid was diluted in 1x Danieau's solution to a concentration of 50 ng/ml. Plasmid solution was loaded into a pulled-glass micropipette, mounted to a micromanipulator (Narishige), and pressure-injected via attachment to a Picospritzer II (Parker). Injections were made at 10 psi for durations from 100 to 200 ms. Injected fish were raised and out-crossed with wild-type fish to screen for founders. Positive progenies were raised to establish transgenic lines.

BC single cell RNA sequencing

BC purification and sequencing

Adult zebrafish carrying the *Tg(vsx1:GFP)nns5* transgene were used to isolate BCs for single-cell RNA sequencing. Retinas were dissected and digested in papain solution containing 20U/ml papain, 80U/ml DNaseI, and 1.5mM L-cysteine in oxygenated (ox) Ames solution at 28°C for 45

minutes. The digestion was stopped by replacing the papain solution with a papain inhibitor

solution containing 15mg/ml ovomucoid and 15mg/ml BSA. The tissue was gently dissociated by

trituration using a flamed glass pipette and washed with ox. Ames containing 0.4% BSA. The

resulting cell suspension was filtered through a 30μm strainer and fluorescence-activated cell

sorting (FACS) was performed. Non-transgenic wild-type retinas were used to determine

background fluorescence levels and adjust sorting gates. Live bipolar cells were distinguished

using Calcein blue. Cells were washed, resuspended in PBS 0.04% BSA, and loaded onto the

microfluidic device within ~45 minutes after FACS enrichment. Droplet RNA sequencing

experiments were conducted on the 10X chromium platform according to the manufacturer's

instructions with no modifications. Up to sixteen retinas from up to eight fish per batch were

dissected and dissociated. Eight cDNA libraries were generated across four experiments with two

replicates each. The cDNA libraries were sequenced on an Illumina HiSeq 2500 to a depth of

~30,000 reads per cell.

Single cell transcriptomics data analysis

We performed the initial preprocessing using the cellranger software suite (version 2.1.0, 10X

Genomics), following steps described previously in our study of Zebrafish RGCs[81]. The

sequencing reads were demultiplexed using "cellranger mkfastq" to obtain a separate set of

fastq.gz files for each of 8 samples, which were distributed across Y biological replicates. Reads

for each sample were aligned to the zebrafish reference transcriptome (ENSEMBL zv10, release

82) using "cellranger count" with default parameters to obtain a binary alignment file and a filtered

gene expression matrix (GEM) for each sample. To account for intronic reads, the binary

alignment files were processed using velocyto with default parameters[82], producing a loom file

containing a GEM for exonic reads and a separate matrix for intronic reads. The matrices were

combined for each sample, resulting in a total gene expression matrix (GEM; genes x cells)

summarizing transcript counts. We used the Seurat R package (Stuart et al., 2019) to combine

the GEMs from different channels and analyzed them for each of the biological replicates, unless

otherwise stated, with default parameter values. Additionally, we evaluated the robustness of our

clustering results to variations in select parameters. The full details of our analyses are

documented   in   markdown   scripts,   which   are   available   at

https://github.com/shekharlab/ZebrafishBC.

Preprocessing and batch integration

The combined GEM was filtered to remove genes expressed in fewer than 25 cells, and cells

expressing fewer than 50 genes resulting in 25,233 genes and 19,492 cells. Briefly, each cell was

normalized to a total library size of 10,000 and the normalized counts were log-transformed using

the function Seurat::NormalizeData. We used Seurat::FindVariableFetures with option

selection.method = "vst" to identify the top 2000 highly variable genes (HVGs)[83] in each batch.

Next, we performed scRNA-seq integration. We used Seurat::FindIntegrationAnchors and

Seurat::IntegrateData, both with options "dims=1:40" to perform Canonical Correlation Analysis

(CCA)-based batch correction on the reduced expression matrix consisting of the HVGs. The

"integrated" expression values were combined across replicates, and used for dimensionality

reduction and clustering.

Dimensionality Reduction, Clustering and Visualization

To remove scale disparities between genes arising from differences in average expression levels,

the integrated expression values for each HVG were z-scored across the cells using

Seurat::ScaleData. Next, we performed Principal Component Analysis (PCA) on the scaled matrix,

and used Seurat::ElbowPlot to select principal components (PCs). Using the top 20 PCs, we built

a k-nearest neighbor graph using Seurat::FindNeighbors and identified transcriptionally distinct

clusters using Seurat::FindClusters, using a resolution parameter of 0.5.

Using the top 20 PCs, we also embedded the cells onto a 2D embedding using Uniform Manifold

Approximation (Becht et al., 2019) using the Seurat function RunUMAP.

Identification of BCs and filtering contaminant classes

BC clusters were identified based on expression of the pan-BC markers *vsx1*, and other cell

classes were filtered based on well known gene markers Examples of such genes include *rlbp1a*

and *apoeb* for Muller glia[84], *rbpms2b* for retinal ganglion cells[85], *gad1* and *gad2* for amacrine

cells[86], *pde6* for photoreceptors[87], and *cldn19* for endothelial cells[88]. A total of 155 cells

corresponding to these cell classes were removed.

Hierarchical clustering

To identify transcriptional relationships between BC clusters, we used

Seurat::FindVariableFeatures to recalculate the top 2000 most variable genes. The average

expression values of genes in each cluster were used as input for hierarchical clustering,

performed using pvclust with parameters method.hclust = "complete" and method.dist =

"correlation". The resulting output was visualized as a dendrogram.

Plasmid construction

Plasmid pBH-vsx2-memCerulean-pA was made using the Gateway system (ThermoFisher,

12538120) with combinations of entry and destination plasmids as follows: pTol2CG2[89], p5E-vsx2,

pME-membrane-Cerulean, p3E-pA[89]. Plasmid p5E-vsx2 was generated by inserting a

polymerase chain reaction (PCR)-amplified vsx2 promoter genomic fragment into p5E plasmid

using BP clonase (ThermoFisher, 11789013). PCR reaction was performed using primers: 5'-

GGGGACAACTTTGTATAGAAAAGTTGATGCTAAACAACTTCAAACGACCAA-3'   and   5'-

GGGGACTGCTTTTTTGTACAAACTTGGCCTCTGAGACTATTCCCTTCTTTG-3'.

Immunostaining and light microscopy imaging

Adult zebrafish were humanely euthanized in ice-chilled fish water. After decapitation, retinal

tissues were dissected from the enucleated whole eyes by removing cornea, lens, and epithelial

layer in 1x in phosphate-buffered saline (PBS). The tissue were immediately fixed in 4%

paraformaldehyde (Agar Scientific, AGR1026) in PBS for 20 min at room temperature (RT)

followed by three washes in PBS. For retinal slice preparation, the tissues were mounted in 2%

agarose in PBS and sliced at 100 or 200 µm thickness using vibratome (TPI 1000). For rod

staining, the tissues were sliced horizontally, parallel to the outer plexiform layer (OPL), to

facilitate antibody penetration in the tissue while preserving bipolar cell dendrites in the OPL.

Sliced or the whole retinal samples were treated with PBS containing 0.2% Triton X-100 (Sigma-

Aldrich, X100) for at least 10 min and up to 1 day, followed by the addition of primary antibodies.

After 3 to 5 days of incubation at 4°C, samples were washed three times with PBS and 0.2%

Triton X-100 solution and treated with secondary antibodies. After 1 day of incubation, samples

were mounted in 1% agar in PBS on a coverslip, and subsequently, PBS was replaced with

mounting media (VECTASHIELD, H-1000) for imaging.

Primary antibodies were 4C12 antibody (mouse, 1:50; kindly provided by Jim Fadool[90] and anti-

mGluR6b antidoby (rabbit, 1:500; kindly provided by Stephan CF Neuhauss[91]). Secondary

antibodies were AlexaFluor594 anti-rabbit (donkey, 1:500; Jackson ImmunoResearch

Laboratories 711-586-152) and DyLight647 anti-mouse (donkey, 1:500; Jackson

ImmunoResearch Laboratories 715-606-150). Confocal image stacks were taken on a TCS SP8

(Leica) with a 63× oil immersion objective (HC PL APO CS2, Leica). Typical voxel size was 90

nm and 0.5 µm in xy and z, respectively. Super-resolution images were taken on an OMX (General

Electric). Contrast, brightness, and pseudo-color were adjusted for display in Fiji [National

Institutes of Health (NIH)].

Image stacks were median filtered in Fiji. For some images, maximum-intensity projections were

generated in Amira (FEI). 3D image reconstructions were digitally sliced using the Amira slice

functions. All measurements were made in Fiji.

Image Analysis

Quantitation of cell density

We obtained the cell density of RBC1 and RBC2 by counting the axon terminals of these cells

within regions of interest from confocal image stacks of the dorsal and ventro-temporal retina.

RBC2 was counted from images of the *Tg(vsx2:memCerulean)[wst01]* line. Because not all RBC1

labeled in the *Tg(vsx1:memCerulean)[q19]* line express mCerulean, we quantified the density of

PKC labeled cells with axon terminals in the bottom layer of the IPL. Counts were obtained from

3-4 retinas from 3 animals of each line. For RBC1, axons were quantified within an area between

37,000 and 85,000 μm^2, and for RBC2, the areas were between 11,000 -22,000 μm^2.

Dendritic field

The dendritic field was defined by tracing the extent of a given cell's dendrites with the polygonal

select tool, and removing any concavity using FIJI (see Figure S2). The dendritic arbor area was

then obtained by calculating the area enclosed by the polygon. Because the dendritic tips of some

neighboring cells of the same type overlapped and could not be distinguished readily, one

investigator repeatedly traced (3 to 4 measurements for a single cell) the dendrite boundary, and

obtained the respective area for a given cell until at least three measurements were within ± 2.5%

of the average of all previous measurements for that cell. Confocal images from three fish were

used, with images of RBC1 and RBC2 cells acquired from the dorsal and ventral regions of the

retina: 10 to 17 cells per fish were measured for each location and cell type, resulting in a total of

33 to 41 cells measured for each location and cell type.

Photoreceptor connectivity

Dendritic contacts with photoreceptors were defined by the co-localization of dendritic tips

extending towards outer nucleus layer and photoreceptor terminals by scrolling through confocal

image stack in Fiji. The percentage contacted was computed by dividing the number of

photoreceptors contacted by a given BC by the number of photoreceptors within the dendritic field

of the BC.

Electrophysiology

Fish (3-6 months old) used in physiology experiments were dark adapted for at least 2 hours, and

the retinas were isolated under infrared light following procedures approved by the Administrative

Panel on Laboratory Animal Care at the University of Washington. Retinas were continuously

superfused (~8 mL/min) with oxygenated (95% O2, 5% CO2) bicarbonate-buffered Ames solution

(Sigma) maintained at 25ºC–28ºC.

Recordings were conducted in a flat-mount preparation with photoreceptors facing down. Bipolar

cells were patched at the axon terminals. To access bipolar cell terminals for recording, small

groups of ganglion cells were suctioned off the top of the retina to expose the inner plexiform

layer. Terminals of RBC1 bipolar cells could be targeted for recording using only infrared

illumination, whereas RBC2 terminals, which were not easily visible without fluorescence imaging,

were targeted using a custom-built two-photon microscope. As a result, measuring rod-mediated

responses in RBC2 was unfeasible due to the compromise of rod responses by two-photon

imaging.

Whole-cell voltage-clamp recordings were obtained using patch pipettes filled with a Cs+-based

internal solution. This internal solution also included Alexa Fluor 594, which was used for two-

photon imaging of each cell after recording to confirm its type by morphology. To isolate excitatory

postsynaptic currents, we voltage-clamped cells near the reversal potential for chloride-mediated

conductances (~-60 mV). In addition, to block inhibitory synaptic transmission, we added the

GABAA receptor antagonist gabazine (20 µM), the GABAC receptor antagonist TPMPA (50 µM),

and the glycine receptor antagonist strychnine (3 µM) to the superfusion solution. In experiments

in which mGluR6-mediated input was blocked, the mGluR6 receptor agonist APB (10 µM) was

also added to the superfusion.

Light from blue or red light-emitting diodes (LEDs, peak output = 470 nm and 640 nm respectively)

was delivered to the recording chamber via fiber optic cable positioned beneath the microscope's

condenser lens. The light uniformly illuminated a circular area through an aperture 0.5 mm in

diameter centered on the recorded cell. Protocols for light stimulation were designed to either

activate rods only (using the blue LED) or both rods and cones (using the red LED).

EM data acquisition, reconstruction, and annotation

Dissected retinal tissues from wild type adult zebrafish were immediately transferred into a 1.5-

ml tube with the fixative (4% glutaraldehyde in 0.1M cacodylate buffer [pH7.4]) and incubated

overnight on a shaker at RT. Subsequently, the tissue was washed three times in 0.1 M

cacodylate buffer (pH7.4) and incubated in a solution containing 1.5% potassium ferrocyanide

and 2% osmium tetroxide ($OsO4$) in 0.1 M cacodylate buffer [0.66% lead in 0.03 M aspartic acid

(pH 5.5)] for 1 hour. After washing, the tissue was placed in a freshly made thiocarbohydrazide

(TCH) solution (0.1 g of TCH in 10 ml of double-distilled H_2O heated to 600°C for 1 hour) for 20

min at RT. After another rinse, at RT, the tissue was incubated in 2% $OsO4$ for 30 min at RT. The

samples were rinsed again and stained en bloc in 1% uranyl acetate overnight at 40°C, washed,

and stained with Walton's lead aspartate for 30 min. After a final wash, the retinal pieces were

dehydrated in a graded ice-cold alcohol series and placed in propylene oxide at RT for 10 min.

Last, the sample was embedded in Durcupan resin. Semithin sections (0.5 to 1 µm thick) were

cut and stained with toluidine blue, until the fiducial marks (box) in the ganglion cell layer (GCL)

appeared. The block was then trimmed and mounted in a serial block-face scanning electron

microscope (GATAN/Zeiss, 3View). Serial sections were cut at a thickness of 70 nm and imaged

at an xy resolution of 7 nm. Six tiles, each about 40 μm by 40 μm with an overlap of about 10%,

covering from the outer nucleus layer to the ganglion cell layer in a side view was obtained. Retinal

location was not recorded. The image stacks were concatenated and aligned using TrakEM2

(NIH). Neurons were traced or painted using the tracing and painting tools in TrakEM2.

Statistics

Mann-Whitney U test was used to determine the p-value for comparing dendritic field sizes.

DATA AVAILABILITY

Computational scripts detailing scRNA-seq analysis reported in this paper are available at

https://github.com/shekharlab/

ZebrafishBC. We have also provided R markdown (Rmd) files that show step-by-step

reproduction of the key results at https://github.com/shekharlab/ZebrafishBC. The raw and

processed scRNA-seq data reported in this paper was obtained from the Gene Expression

Omnibus (GEO) entry GSE237215 (subseries GSE237214).

Figure S4

Monostratifying-AC with reciprocal synapses
(A17-like)

Figure S4. Gallery of mono-stratifying AC making reciprocal synapses with RBC1

En face and side views of individual cells or processes. The numbers of input (open bar) and output (closed bar) synapses with each RBC1 (blue) and RBC2 (red) terminal are indicated as the height of the bars.

Figure S5

Monostratifying-AC without reciprocal synapses

10 μm

Figure S5. Gallery of mono-stratifying AC without reciprocal synapses with RBC1

En face and side views of individual cells. The numbers of input (open bar) and output (closed bar) synapses with each RBC1 (blue) and RBC2 (red) terminal are indicated as the height of the bars.

Figure S6

Bi-stratifying-AC
(A2-like)

RGC

10 μm

RBC1   RBC2

15 #Synapse
0

Input Output   Input Output

Figure S6. Gallery of bi-stratifying AC and RGC contacted to RBC1

En face and side views of individual cells. The numbers of input (open bar) and output (closed

bar) synapses with each RBC1 (blue) and RBC2 (red) terminal are indicated as the height of the

bars.

Figure S7

Monostratifying-AC with reciprocal synapses

Monostratifying-AC without reciprocal synapses

10 μm

RBC1   RBC2

Input Output   Input Output

15
0
#Synapse

748 *Figure S7. Gallery of mono-stratifying AC connected to RBC2*

749 *En face and side views of individual cells. The numbers of input (open bar) and output (closed*

750 *bar) synapses with each RBC1 (blue) and RBC2 (red) terminal are indicated as the height of the*

751 *bars.*

Figure S8

Bi-stratified-AC   RGC

754 *Figure S8. Gallery of bi-stratifying AC and RGC connected to RBC2*

755 *En face and side views of individual cells. The numbers of input (open bar) and output (closed*

756 *bar) synapses with each RBC1 (blue) and RBC2 (red) terminal are indicated as the height of the*

757 *bars.*

REFERENCES

1. Morshedian, A. & Fain, G. L. The evolution of rod photoreceptors. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, 20160074 (2017).

2. Bloomfield, S. A. & Dacheux, R. F. Rod Vision: Pathways and Processing in the Mammalian Retina. *Prog. Retin. Eye Res.* **20**, 351–384 (2001).

3. Field, G. D., Sampath, A. P. & Rieke, F. RETINAL PROCESSING NEAR ABSOLUTE THRESHOLD: From Behavior to Mechanism. *Annu. Rev. Physiol.* **67**, 491–514 (2005).

4. Grimes, W. N., Songco-Aguas, A. & Rieke, F. Parallel Processing of Rod and Cone Signals: Retinal Function and Human Perception. *Annu. Rev. Vis. Sci.* **4**, 123–141 (2018).

5. Strettoi, E. 1.11 - Mammalian Rod Pathways. in *The Senses: A Comprehensive Reference* (eds. Masland, R. H. et al.) 303–311 (Academic Press, 2008). doi:10.1016/B978-012370880-9.00266-8.

6. Wu, S. M. Synaptic Organization of the Vertebrate Retina: General Principles and Species-Specific Variations: The Friedenwald Lecture. *Invest. Ophthalmol. Vis. Sci.* **51**, 1264–1274 (2010).

7. Thoreson, W. B. & Dacey, D. M. Diverse Cell Types, Circuits, and Mechanisms for Color Vision in the Vertebrate Retina. *Physiol. Rev.* **99**, 1527–1573 (2019).

8. Behrens, C., Schubert, T., Haverkamp, S., Euler, T. & Berens, P. Connectivity map of bipolar cells and photoreceptors in the mouse retina. *eLife* **5**, e20041 (2016).

9. Baylor, D. A., Lamb, T. D. & Yau, K. W. Responses of retinal rods to single photons. *J. Physiol.* **288**, 613–634 (1979).

10. Baylor, D. A., Nunn, B. J. & Schnapf, J. L. The photocurrent, noise and spectral sensitivity of rods of the monkey Macaca fascicularis. *J. Physiol.* **357**, 575–607 (1984).

11. Field, G. D. & Rieke, F. Nonlinear Signal Transfer from Mouse Rods to Bipolar Cells and Implications for Visual Sensitivity. *Neuron* **34**, 773–785 (2002).

12. Field, G. D. & Rieke, F. Mechanisms regulating variability of the single photon responses of

mammalian rod photoreceptors. *Neuron* **35**, 733–747 (2002).

13. Ingram, N. T., Sampath, A. P. & Fain, G. L. Why are rods more sensitive than cones? *J. Physiol.* **594**,

5415–5426 (2016).

14. Yang, X.-L. & Wu, S. M. Response Sensitivity and Voltage Gain of the Rod- and Cone-Bipolar Cell

Synapses in Dark-Adapted Tiger Salamander Retina. *J. Neurophysiol.* **78**, 2662–2673 (1997).

15. Kolb, H. & Famiglietti, E. V. Rod and Cone Pathways in the Inner Plexiform Layer of Cat Retina.

Science **186**, 47–49 (1974).

16. Strettoi, E., Masri, R. A. & Grünert, U. AII amacrine cells in the primate fovea contribute to photopic

vision. *Sci. Rep.* **8**, 16429 (2018).

17. Lee, S. C. S., Martin, P. R. & Grünert, U. Topography of Neurons in the Rod Pathway of Human

Retina. *Invest. Ophthalmol. Vis. Sci.* **60**, 2848–2859 (2019).

18. Tsukamoto, Y., Morigiwa, K., Ueda, M. & Sterling, P. Microcircuits for Night Vision in Mouse Retina.

J. Neurosci. **21**, 8616–8623 (2001).

19. Kolb, H. & Nelson, R. Rod pathways in the retina of the cat. *Vision Res.* **23**, 301–312 (1983).

20. Dacheux, R. F. & Raviola, E. The rod pathway in the rabbit retina: a depolarizing bipolar and

amacrine cell. *J. Neurosci.* **6**, 331–345 (1986).

21. Freed, M. A., Smith, R. G. & Sterling, P. Rod bipolar array in the cat retina: pattern of input from rods

and GABA-accumulating amacrine cells. *J. Comp. Neurol.* **266**, 445–455 (1987).

22. Joselevitch, C. & Kamermans, M. Retinal parallel pathways: Seeing with our inner fish. *Vision Res.*

49, 943–959 (2009).

23. Franke, K. *et al.* Inhibition decorrelates visual feature representations in the inner retina. *Nature*

542, 439–444 (2017).

24. Ghosh, K. K., Bujan, S., Haverkamp, S., Feigenspan, A. & Wässle, H. Types of bipolar cells in the

mouse retina. *J. Comp. Neurol.* **469**, 70–82 (2004).

25. Grünert, U. & Martin, P. R. Cell types and cell circuits in human and non-human primate retina.

Prog. Retin. Eye Res. 100844 (2020) doi:10.1016/j.preteyeres.2020.100844.

26. Peng, Y.-R. *et al.* Molecular Classification and Comparative Taxonomics of Foveal and Peripheral

Cells in Primate Retina. *Cell* **176**, 1222-1237.e22 (2019).

27. Shekhar, K. *et al.* Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell

Transcriptomics. *Cell* **166**, 1308-1323.e30 (2016).

28. Masland, R. H. The neuronal organization of the retina. *Neuron* **76**, 266–80 (2012).

29. Hahn, J. *et al.* Evolution of neuronal cell classes and types in the vertebrate retina.

2023.04.07.536039 Preprint at https://doi.org/10.1101/2023.04.07.536039 (2023).

30. Strettoi, E., Dacheux, R. F. & Raviola, E. Synaptic connections of rod bipolar cells in the inner

plexiform layer of the rabbit retina. *J. Comp. Neurol.* **295**, 449–466 (1990).

31. Zhang, C., Hellevik, A., Takeuchi, S. & Wong, R. O. Hierarchical partner selection shapes rod-cone

pathway specificity in the inner retina. *iScience* **25**, 105032 (2022).

32. Strettoi, E., Raviola, E. & Dacheux, R. F. Synaptic connections of the narrow-field, bistratified rod

amacrine cell (AII) in the rabbit retina. *J. Comp. Neurol.* **325**, 152–168 (1992).

33. Dunn, F. A. & Rieke, F. Single-Photon Absorptions Evoke Synaptic Depression in the Retina to Extend

the Operational Range of Rod Vision. *Neuron* **57**, 894–904 (2008).

34. Grimes, W. N., Hoon, M., Briggman, K. L., Wong, R. O. & Rieke, F. Cross-synaptic synchrony and

transmission of signal and noise across the mouse retina. *eLife* **3**, e03892 (2014).

35. Pang, J.-J., Gao, F. & Wu, S. M. Light-evoked current responses in rod bipolar cells, cone depolarizing

bipolar cells and AII amacrine cells in dark-adapted mouse retina. *J. Physiol.* **558**, 897–912 (2004).

36. Demb, J. B. & Singer, J. H. Intrinsic properties and functional circuitry of the AII amacrine cell. *Vis.*

Neurosci. **29**, 51–60 (2012).

37. Strettoi, E., Dacheux, R. F. & Raviola, E. Cone bipolar cells as interneurons in the rod, pathway of the

rabbit retina. *J. Comp. Neurol.* **347**, 139–149 (1994).

38. Vitorino, M. *et al.* Vsx2 in the zebrafish retina: restricted lineages through derepression. *Neural*

Develop. **4**, 14 (2009).

39. Križaj, D., Cordeiro, S. & Strauß, O. Retinal TRP channels: Cell-type-specific regulators of retinal

homeostasis and multimodal integration. *Prog. Retin. Eye Res.* **92**, 101114 (2023).

40. Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat. Biotechnol.*

37, 38–44 (2019).

41. Wässle, H., Yamashita, M., Greferath, U., Grünert, U. & Müller, F. The rod bipolar cell of the

mammalian retina. *Vis. Neurosci.* **7**, 99–112 (1991).

42. Wässle, H., Puller, C., Müller, F. & Haverkamp, S. Cone Contacts, Mosaics, and Territories of Bipolar

Cells in the Mouse Retina. *J. Neurosci.* **29**, 106–117 (2009).

43. Choi, H. M. T. *et al.* Third-generation in situ hybridization chain reaction: multiplexed, quantitative,

sensitive, versatile, robust. *Development* **145**, dev165753 (2018).

44. Vihtelic, T. S., Doro, C. J. & Hyde, D. R. Cloning and characterization of six zebrafish photoreceptor

opsin cDNAs and immunolocalization of their corresponding proteins. *Vis. Neurosci.* **16**, 571–585

(1999).

45. Chinen, A., Hamaoka, T., Yamada, Y. & Kawamura, S. Gene Duplication and Spectral Diversification

of Cone Visual Pigments of Zebrafish. *Genetics* **163**, 663–675 (2003).

46. Sandell, J. H., Masland, R. H., Raviola, E. & Dacheux, R. F. Connections of indoleamine-accumulating

cells in the rabbit retina. *J. Comp. Neurol.* **283**, 303–313 (1989).

47. Grimes, W. N., Zhang, J., Graydon, C. W., Kachar, B. & Diamond, J. S. Retinal parallel processors:

more than 100 independent microcircuits operate within a single interneuron. *Neuron* **65**, 873–885

(2010).

48. Marc, R. E., Anderson, J. R., Jones, B. W., Sigulinsky, C. L. & Lauritzen, J. S. The AII amacrine cell

connectome: a dense network hub. *Front. Neural Circuits* **8**, (2014).

49. Gamlin, C. R., Zhang, C., Dyer, M. A. & Wong, R. O. L. Distinct Developmental Mechanisms Act

Independently to Shape Biased Synaptic Divergence from an Inhibitory Neuron. *Curr. Biol. CB* **30**,

1258-1268.e2 (2020).

50. Tsukamoto, Y. & Omi, N. Functional allocation of synaptic contacts in microcircuits from rods via rod

bipolar to AII amacrine cells in the mouse retina. *J. Comp. Neurol.* **521**, 3541–3555 (2013).

51. Li, Y. N., Tsujimura, T., Kawamura, S. & Dowling, J. E. Bipolar Cell-Photoreceptor Connectivity in the

Zebrafish (Danio rerio) Retina. *J. Comp. Neurol.* **520**, 3786–3802 (2012).

52. Zimmermann, M. J. Y. *et al.* Zebrafish Differentially Process Color across Visual Space to Match

Natural Scenes. *Curr. Biol. CB* **28**, 2018-2032.e5 (2018).

53. Zhou, M. *et al.* Zebrafish Retinal Ganglion Cells Asymmetrically Encode Spectral and Temporal

Information across Visual Space. *Curr. Biol.* **30**, 2927-2942.e7 (2020).

54. Yoshimatsu, T., Schröder, C., Nevala, N. E., Berens, P. & Baden, T. Fovea-like Photoreceptor

Specializations Underlie Single UV Cone Driven Prey-Capture Behavior in Zebrafish. *Neuron* **107**,

320-337.e6 (2020).

55. Schröder, C., Oesterle, J., Berens, P., Yoshimatsu, T. & Baden, T. Distinct synaptic transfer functions

in same-type photoreceptors. *eLife* **10**, e67851 (2021).

56. Yan, W. *et al.* Cell Atlas of The Human Fovea and Peripheral Retina. *Sci. Rep.* **10**, 9802 (2020).

57. Günther, A. *et al.* Double Cones and the Diverse Connectivity of Photoreceptors and Bipolar Cells in

an Avian Retina. *J. Neurosci.* **41**, 5015–5028 (2021).

58. Yamagata, M., Yan, W. & Sanes, J. R. A cell atlas of the chick retina based on single-cell

transcriptomics. *eLife* **10**, e63907 (2021).

59. Baden, T. & Osorio, D. The Retinal Basis of Vertebrate Color Vision. *Annu. Rev. Vis. Sci.* **5**, 177–200

(2019).

60. Lamb, T. D. Evolution of phototransduction, vertebrate photoreceptors and retina. *Prog. Retin. Eye*

Res. **36**, 52–119 (2013).

61. Sherry, D. M. & Yazulla, S. Goldfish bipolar cells and axon terminal patterns: A Golgi study. *J. Comp.*

Neurol. **329**, 188–200 (1993).

62. Behrens, U. D., Borde, J., Mack, A. F. & Wagner, H.-J. Distribution of phosphorylated protein kinase C

alpha in goldfish retinal bipolar synaptic terminals: control by state of adaptation and

pharmacological treatment. *Cell Tissue Res.* **327**, 209–220 (2007).

63. Pang, J.-J., Gao, F. & Wu, S. M. Stratum-by-stratum projection of light response attributes by retinal

bipolar cells of Ambystoma. *J. Physiol.* **558**, 249–262 (2004).

64. Balaji, V. *et al.* Immunohistochemical characterization of bipolar cells in four distantly related avian

species. *J. Comp. Neurol.* **531**, 561–581 (2023).

65. Caminos, E., Velasco, A., Jarrín, M., Aijón, J. & Lara, J. M. Protein kinase C-like immunoreactive cells

in embryo and adult chicken retinas. *Dev. Brain Res.* **118**, 227–230 (1999).

66. Frederiksen, R., Fain, G. L. & Sampath, A. P. A hyperpolarizing rod bipolar cell in the sea lamprey,

Petromyzon marinus. *J. Exp. Biol.* **225**, jeb243949 (2022).

67. Tsukamoto, Y. & Omi, N. Classification of Mouse Retinal Bipolar Cells: Type-Specific Connectivity

with Special Reference to Rod-Driven AII Amacrine Pathways. *Front. Neuroanat.* **11**, 92 (2017).

68. Wong, K. Y., Cohen, E. D. & Dowling, J. E. Retinal Bipolar Cell Input Mechanisms in Giant Danio. II.

Patch-Clamp Analysis of on Bipolar Cells. *J. Neurophysiol.* **93**, 94–107 (2005).

69. Yoshimatsu, T. *et al.* Ancestral circuits for vertebrate color vision emerge at the first retinal synapse.

Sci. Adv. **7**, eabj6815 (2021).

70. Ishida, A. T., Stell, W. K. & Lightfoot, D. O. Rod and cone inputs to bipolar cells in goldfish retina. *J.*

Comp. Neurol. **191**, 315–335 (1980).

71. Connaughton, V. P. & Nelson, R. Axonal stratification patterns and glutamate-gated conductance

mechanisms in zebrafish retinal bipolar cells. *J. Physiol.* **524 Pt 1**, 135–46 (2000).

72. Joselevitch, C. & Kamermans, M. Interaction between rod and cone inputs in mixed-input bipolar

cells in goldfish retina. *J. Neurosci. Res.* **85**, 1579–1591 (2007).

73. Pang, J.-J. *et al.* Direct rod input to cone BCs and direct cone input to rod BCs challenge the

traditional view of mammalian BC circuitry. *Proc. Natl. Acad. Sci.* **107**, 395–400 (2010).

74. Pang, J.-J., Yang, Z., Jacoby, R. A. & Wu, S. M. Cone synapses in mammalian retinal rod bipolar cells.

J. Comp. Neurol. **526**, 1896–1909 (2018).

75. Whitaker, C. M., Nobles, G., Ishibashi, M. & Massey, S. C. Rod and Cone Connections With Bipolar

Cells in the Rabbit Retina. *Front. Cell. Neurosci.* **15**, 662329 (2021).

76. Carter-Dawson, L. D. & LaVail, M. M. Rods and cones in the mouse retina. I. Structural analysis using

light and electron microscopy. *J. Comp. Neurol.* **188**, 245–262 (1979).

77. Famiglietti, E. V. & Sharpe, S. J. Regional topography of rod and immunocytochemically

characterized 'blue' and 'green' cone photoreceptors in rabbit retina. *Vis. Neurosci.* **12**, 1151–1175

(1995).

78. Kimura, Y., Satou, C. & Higashijima, S. V2a and V2b neurons are generated by the final divisions of

pair-producing progenitors in the zebrafish spinal cord. *Development* **135**, 3001–3005 (2008).

79. Randlett, O. *et al.* Cellular requirements for building a retinal neuropil. *Cell Rep.* **3**, 282–290 (2013).

80. Suzuki, S. C. *et al.* Cone photoreceptor types in zebrafish are generated by symmetric terminal

divisions of dedicated precursors. *Proc. Natl. Acad. Sci.* **110**, 15109–15114 (2013).

81. Kölsch, Y. *et al.* Molecular classification of zebrafish retinal ganglion cells links genes to cell types to

behavior. *Neuron* **109**, 645-662.e9 (2021).

82. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

83. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data

using regularized negative binomial regression. *Genome Biol.* **20**, 296 (2019).

84. Bernardos, R. L., Barthel, L. K., Meyers, J. R. & Raymond, P. A. Late-Stage Neuronal Progenitors in

the Retina Are Radial Müller Glia That Function as Retinal Stem Cells. *J. Neurosci.* **27**, 7028–7040

(2007).

85. Hörnberg, H. *et al.* RNA-binding protein Hermes/RBPMS inversely affects synapse density and axon

arbor formation in retinal ganglion cells in vivo. *J. Neurosci. Off. J. Soc. Neurosci.* **33**, 10384–10395

(2013).

86. Sandell, J. H., Martin, S. C. & Heinrich, G. The development of GABA immunoreactivity in the retina

of the zebrafish (Brachydanio rerio). *J. Comp. Neurol.* **345**, 596–601 (1994).

87. Abalo, X. M. *et al.* Circadian regulation of phosphodiesterase 6 genes in zebrafish differs between

cones and rods: Implications for photopic and scotopic vision. *Vision Res.* **166**, 43–51 (2020).

88. Kolosov, D., Bui, P., Chasiotis, H. & Kelly, S. P. Claudins in teleost fishes. *Tissue Barriers* **1**, e25391

(2013).

89. Kwan, K. M. *et al.* The Tol2kit: A multisite gateway-based construction kit forTol2 transposon

transgenesis constructs. *Dev. Dyn.* **236**, 3088–3099 (2007).

90. Morris, A. C., Scholz, T. L., Brockerhoff, S. E. & Fadool, J. M. Genetic Dissection Reveals Two Separate

Pathways for Rod and Cone Regeneration in the Teleost Retina. *Dev. Neurobiol.* **68**, 605–619 (2008).

91. Huang, Y.-Y., Haug, M. F., Gesemann, M. & Neuhauss, S. C. F. Novel Expression Patterns of

Metabotropic Glutamate Receptor 6 in the Zebrafish Nervous System. *PLOS ONE* **7**, e35256 (2012).