Inducer of CBF Expression 1 (ICE1) Promotes Cold-enhanced Immunity by Directly Activating Salicylic Acid Signaling

Shaoqin Li[a,b], Li He[c], Yongping Yang[a,b], Xiao Han[a], Yanru Hu[a], Yanjuan Jiang[a,c,*]

[a] CAS Key Laboratory of Tropical Plant Resources and Sustainable Use, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Kunming, Yunnan 650223, China.
[b] University of Chinese Academy of Sciences, Beijing 100049, China.
[c] State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, School of Life Sciences, Yunnan University, Kunming 650091, China

*Corresponding author
Email: yanjuanjiang@ynu.edu.cn

Short title: ICE1 activates SA signaling

ABSTRACT

Cold stress affects plant immune responses, and this process may involve the salicylic acid (SA) signaling pathway. However, the underlying mechanism by which low temperature signals coordinate with SA signaling to regulate plant immunity remains poorly characterized. Here, we found that low temperatures enhanced the disease resistance of Arabidopsis against *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000. This process required Inducer of CBF expression 1 (ICE1), the core transcription factor in cold-signal cascades. ICE1 physically interacted with Non-expresser of *PR* genes 1 (NPR1), the master regulator of the SA signaling pathway. Enrichment of ICE1 on the *PR1* promoter and its ability to transcriptionally activate *PR1* were enhanced by NPR1. Further analyses revealed that cold stress signals cooperate with SA signals to facilitate plant immunity against pathogen attack in an ICE1-dependent manner. Cold treatment promoted interactions of NPR1 and TGA3 with ICE1, and increased the ability of the ICE1–TGA3 complex to transcriptionally activate *PR1*. Together, our results characterize a previously unrecognized role of ICE1 as an indispensable regulatory node linking low temperature activated- and SA-regulated immunity. Discovery of a crucial role of ICE1 in coordinating multiple signals associated with immunity broadens our understanding of plant–pathogen interactions.

INTRODUCTION

In nature, plants can be attacked by a wide range of pathogens, including bacteria, fungi, oomycetes, viruses, and nematodes. In response, plants have developed a sophisticated innate immune system during co-evolution. The first layer of the defense system is governed by the plasma membrane (PM)-localized pattern recognition receptors (PRRs). The PRRs recognize the microorganism-derived pathogen-associated molecular patterns (PAMPs) and trigger the first layer of defense responses–this is known as PAMP-triggered immunity (PTI) (Jones and Dangl, 2006; Dodds and Rathjen, 2010). PTI is generally believed to be sufficient to prevent non-pathogenic microbes from proliferating *in planta.* However, some successful pathogens can deliver effector proteins to evade or suppress host PTI, allowing them to multiply aggressively (Kinya Nomur and He, 2005; Boller and Felix, 2009). To counter pathogen inhibition of PTI, plants have evolved intracellular nucleotide-binding leucine-rich repeat receptors (NLRs) that detect cellular effectors, directly or indirectly, and cause strong and robust immune responses. These responses, known as effector-triggered immunity (ETI), can stop virulent pathogens from causing severe disease symptoms (Jones and Dangl, 2006; Dodds and Rathjen, 2010; Deslandes and Rivas, 2012). Along with the two branches of immune system, the defense hormone salicylic acid (SA) plays a central role in these responses (Spoel and Dong, 2012). For example, the accumulation of SA at infection sites subsequently leads to systemic acquired resistance (SAR) against *Pseudomonas syringae* throughout the plant to limit pathogen propagation (Durrant and Dong, 2004; Vlot et al., 2009).

In the SA signaling transduction pathway, NONEXPRESSOR OF PATHOGENESIS-RELATED GENES 1 (NPR1) has been well documented as one of the SA receptors that perceives and transduces SA signals (Wang et al., 2020; Kumar et al., 2022). In the resting state, NPR1 exists in the cytoplasm as an oligomer. After induction, NPR1 monomers are released from oligomers and translocated to the nucleus, where they act as cofactors of transcription factors, such as the TGACG motif-binding transcription factors (TGAs), which regulate global transcriptional reprogramming and resistance to a broad spectrum of pathogens (Zhang et al., 1999; Després et al., 2000; Zhou et al., 2000; Johnson et al., 2003; Kumar et al., 2022). Recently, NPR1 has attracted

attention because of its central roles in plant growth and development, as well as in plant immune responses. For instance, it was reported that NPR1 interferes with the binding of ETHYLENE-INSENSITIVE 3 (EIN3), a key transcription factor involved in ethylene (ET) signaling, to the promoter regions of its target genes to mediate antagonism between SA and ET in apical hook formation (Huang et al., 2020). The transcriptional activation activity of EIN3 is enhanced by NPR1 in a process coordinated by SA and ET during the modulation of leaf senescence (Wang et al., 2021; Yu et al., 2021). Moreover, NPR1 promotes the polyubiquitination and degradation of GID1, the gibberellin receptor, to balance growth and defense (Yu et al., 2022). NPR1 is also involved in facilitating cell survival during ETI by ubiquitinating key proteins involved in immune responses, such as ENHANCED DISEASE SUSCEPTIBILITY 1 and WRKY transcription factors (Zavaliev et al., 2020). NPR1 inhibits the transcriptional activation activity of MYC2, the master regulator of jasmonic acid signaling pathway, to suppress pathogen virulence (Nomoto et al., 2021).

The TGA transcription factors are members of the basic leucine zipper (bZIP) family of proteins, which specifically bind to the TGACG-core activation sequence 1 (as-1) to regulate the transcriptional levels of their target genes (Després et al., 2000; Zhou et al., 2000; Johnson et al., 2003). Ten TGA transcription factors are encoded in the Arabidopsis genome, of which TGA1 to TGA7 have been characterized in terms of their ability to interact with NPR1 (Zhang et al., 1999; Després et al., 2000). Among them, TGA3 is mainly required for basal resistance and PR gene expression (Kesarwani et al., 2007; Kumar et al., 2022), and also integrates with other signals. For instance, TGA3 interacts with ARR2, the cytokinin-activated transcription factor, to promote plant immunity through NPR1/TGA3-dependent SA signaling (Choi et al., 2010). A protein complex of TGA3 and WRKY53 acts on the Cestrum yellow leaf curling virus promoter to achieve synergistic up-regulation of SA-associated gene expression (Sarkar et al., 2018). BRASSINOSTEROID INSENSITIVE2 (BIN2) balances SA-induced immunity and brassinosteroid-mediated plant growth through phosphorylating TGA3 to activate gene expression (Han et al., 2022).

Low temperature has significant effects on plant growth and development

(Orvar et al., 2000; Wang et al., 2009). The ICE1/C-REPEAT BINDING FACTOR/DRE-BINDING FACTOR 1 (ICE1/CBF/DREB1) pathway plays an essential role in plants' tolerance to low temperature stress, and ICE1 acts as an important transcriptional regulator of cold-responsive gene expression and cold tolerance (Stockinger et al., 1997; Kim et al., 2015). ICE1, a basic helix-loop-helix (bHLH) protein in the MYC class, directly binds to the MYC site (CANNTG) in the promoter of *CBF3* to induce its expression under cold stress (Gilmour et al., 1998; Chinnusamy et al., 2003). ICE1 also binds to the promoters of other genes encoding regulatory factors such as cold-responsive (COR), calmodulin binding transcription activator 2, mitogen-activated protein kinase 3 (MPK3), MPK4, and high expression of osmotically responsive genes 1 (HOS1), all of which are critical for the cold response (Tang et al., 2020). As an essential component of the cold signaling network, ICE1 functions as a key cross-talk node between low temperature signals and other signals. For instance, HOS1, SUMO E3 ligase (SIZ1), Open Stomata 1 (OST1), MPK3/6, and BIN2 are involved in regulating responses to freezing by maintaining homeostasis of the ICE1 protein under cold stress (Dong et al., 2006; Miura et al., 2007; Ding et al., 2015; Li et al., 2017; Zhao et al., 2017; Ye et al., 2019). Moreover, ICE1 integrates endogenous and environmental signals to mediate diverse physiological processes, such as seed germination, flowering time, and stomatal development (Lee et al., 2015, 2017; Hu et al., 2019; An et al., 2021; An et al., 2022).

Plant–pathogen interactions are affected by the ambient temperature, and low temperature can promote plant immune responses. For example, under cold temperatures, winter rye (*Secale cereale*) accumulates abundant pathogenesis-related (PR) proteins that help protect against pathogen attack (Griffith and Yaish, 2004). Singh (2014) found that 7 days of repeated cold stress activated the PTI immune response of Arabidopsis in a histone acetyltransferase 1 (HAT1)-dependent manner, thereby enhancing resistance to the virulent pathogen *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000. Wu (2019) reported that low temperatures induced a transient defense response to *Pst* DC3000 in plants. Specifically, low temperatures induced H_2O_2 accumulation, callose deposition, and defense gene expression (Wu et al., 2019). Notably, there is a growing body of evidence showing that SA-mediated

defense responses are activated at low temperature in the interaction between Arabidopsis and *Pseudomonas syringae*. In particular, SA biosynthetic genes, including *ISOCHORISMATE SYNTHASE1* (*ICS1*), *ISOCHORISMATE SYNTHASE2* (*ICS2*), *CALMODULIN BINDING PROTEIN 60-LIKE.g* (*CBP60g*), *SAR DEFICIENT1* (*SARD1*), and *PHENYLALANINE AMMONIA LYASE1* (*PAL1*), are up-regulated by cold stress, activating the SA-mediated immune pathway (Kim et al., 2013; Kim et al., 2017; Wu et al., 2019; Li et al., 2020). Cold temperatures induce expression of the SA signaling-responsive genes *PR1*, *PR2,* and *PR5,* whose encoded products increase plant immunity (Seo et al., 2010). These results suggest that there are broad signaling interactions between low temperature signals and SA responses. Nonetheless, the underlying molecular mechanisms and signaling pathways are unknown.

In this study, we found that the enhancement of plant immune responses by low temperatures largely depends on ICE1, the core transcription factor in the cold signaling pathway. Mechanistic analyses revealed that the ICE1 protein physically interacts with the SA receptor NPR1 and is involved in SA-mediated resistance against *Pst* DC3000. Further analyses demonstrated that ICE1 directly activates the transcription of *PR1* and NPR1 promotes its activities. We also found that ICE1 interacts with TGA transcription factors to cooperatively stimulate *PR1* expression under low temperatures. Taken together, our results demonstrate that the NPR1-TGA3/ICE1 regulatory module acts as a crucial node integrating cold signals and SA signaling in the synergistic regulation of plant immunity.

RESULTS

ICE1 Is Involved in Low Temperature-enhanced Plant Immunity

Low temperature has complicated effects on plant immunity (Yang and Hua, 2004; Huang et al., 2010; Yang et al., 2010; Wigge, 2013). To investigate the underlying relationship between low temperature and plant immunity, we assessed effect of low temperature on plant immune responses against the hemibiotropic pathogenic bacteria, *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000. For these analyses, we monitored disease symptoms and populations of *Pst* DC3000 in Arabidopsis Columbia-0 (Col-0) plants that were grown at normal temperature (22°C) for 4 weeks and then exposed to a

long-term (3 weeks) or short-term (10 hours) cold treatment (4°C) before infiltration with the pathogen. We observed that the pathogen-related water-soaking symptom, one of the earliest and most common symptoms of phyllosphere bacterial diseases, as well as bacterial proliferation in infected leaves, were largely reduced in Col-0 plants exposed to a short or long cold treatment, compared with those kept at 22°C (Fig. 1 A–D). These results confirmed that both long- and short-term low temperature treatments promote plant immunity against *Pst* DC3000, consistent with the results of previous studies (Seo et al., 2010; Kim et al., 2017; Wu et al., 2019).

ICE1 is a transcriptional inducer of genes that encode important components of chilling tolerance of Arabidopsis (Chinnusamy et al., 2003; Lee et al., 2005; Benedict et al., 2006; Chinnusamy et al., 2007). At the early stages of cold stress, ICE1 in its phosphorylated form activates the expression of *CBFs* to enhance plant freezing tolerance (Ding et al., 2015; Ye et al., 2019). Given the critical role of ICE1 at the early stage of the cold stress response, we hypothesized that ICE1 may be involved in short-term low temperature-enhanced immunity. To test this possibility, we investigated *Pst* proliferation in Col-0, the loss-of-function *ice1-2* mutant, and an *ICE1*-complemented mutant (*ice1-2/ICE1*) after short-term low temperature treatment. As shown in Fig. 1E, F, the leaves of Col-0 and *ice1-2/ICE1* exhibited clearly enhanced disease-resistance phenotypes, with less severe water-soaking symptoms and lower pathogen populations after cold treatment, compared with the control. However, in *ice1-2* mutants, the water-soaking symptoms were severe and the pathogen population was large (Fig. 1E, F). These results suggest that ICE1 may positively regulate cold stress-enhanced plant immunity. To confirm the role of ICE1, we used two stable *ICE1*-overexpressing transgenic lines, *35S::GFP-ICE1* (Chinnusamy et al., 2003); hereinafter *ICE1-OE 1*) and *35S::3Myc-ICE1* (hereinafter *ICE1-OE 2*) for bacterial propagation assays. As expected, compared with Col-0, the *ICE1*-overexpressing plants showed enhanced resistance, including less severe water soaking symptoms and lower pathogen proliferation after low temperature stress (Fig. 1G, H). Together, these results indicate that ICE1 acts as a positive regulator of cold stress-enhanced plant immunity.

ICE1 Physically Interacts with NPR1

ICE1 regulates diverse physiological processes through associating with other proteins (Ding et al., 2015; Lee et al., 2015; Hu et al., 2019; Ye et al., 2019; An et al., 2021; An et al., 2022). To explore the underlying molecular mechanism by which ICE1 regulates plant immune responses, we performed yeast two-hybrid screening analyses to identify its potential interacting protein partners. A potential candidate, NPR1, was identified and further characterized (Fig. 2A). To identify the functional region responsible for the interaction between ICE1 and NPR1, two truncated versions of the proteins were used in yeast two-hybrid assays. As shown in Fig. 2A, deletion of the N-terminal amino acid residues 1–260 of ICE1 (AD-ICE1$^{1-260}$) did not affect the ICE1–NPR1 interaction, whereas deletion of the C-terminal residues of ICE1 containing the bHLH domain (AD-ICE1$^{261-494}$) eliminated the interaction completely (Fig. 2A). Additionally, the C-terminal amino acid residues of NPR1 containing the BTB structural domain (BD-NPR1$^{1-194}$) strongly interacted with ICE1, while the N-terminal fragment (BD-NPR1$^{178-593}$) did not. These results demonstrate that the bHLH structural domain of ICE1 and the BTB structural domain of NPR1 are necessary for the ICE1–NPR1 interaction.

Bimolecular fluorescence complementation (BiFC) assays were conducted to further confirm the interaction between ICE1 and NPR1. For these assays, NPR1 fused to the C-terminal of YFP (NPR1-cYFP) and ICE1 fused to the N-terminal of YFP (ICE1-nYFP) were co-expressed in *Nicotiana benthamiana* leaves, and then fluorescence in the leaves was observed under a laser confocal microscope. Almost no YFP fluorescence was observed when NPR1-cYFP was co-expressed with ICE1-nYFP in *N. benthamiana* leaves at 22°C (Supplemental Fig. S1A), but the fluorescence signal became very strong after induction at 4°C for 4 h (Fig. 2B; Supplemental Fig. S2A), suggesting that the association between NPR1 and ICE1 may depend on low temperature *in vivo.* To further verify this interaction, we performed a split luciferase assay by fusing these proteins to the C-terminus (cLUC) or N-terminal (nLUC) of luciferase, with LUC fused with β-glucuronidase (GUS) as the negative control. Similarly, very faint LUC activity was detected when NPR1-cLUC was transiently co-expressed with ICE1-nLUC in *N. benthamiana*, but strong LUC signals were observed after low temperature induction (Fig. 2C; Supplemental

Fig. S1B).

Consistently, in a co-immunoprecipitation (Co-IP) assay, ICE1 was

immunoprecipitated by anti-GFP agarose beads from Arabidopsis protoplasts

co-expressing NPR1-YFP and ICE1-Myc, but not from those co-expressing

YFP and ICE1-Myc, and more precipitation products were detected after low

temperature induction (Fig. 2D). Taken together, these results indicate that

ICE1 interacts with NPR1 both *in vitro* and *in vivo*, and that low temperature

potentiates the association between ICE1 and NPR1 *in planta*.

ICE1 Is Required for SA-mediated Plant Immunity

Because ICE1 physically associates with the SA receptor NPR1 (Fig. 2), we

hypothesized that ICE1 may participate in SA signal transduction. To test this

hypothesis, we first detected changes in the transcript levels of *ICE1* in

response to benzothiadiazole (BTH), a synthetic analog of SA that effectively

induces SA responses and is extensively used as an alternative to SA

signaling (Huot et al., 2017; Kim et al., 2022). Interestingly, BTH treatment did

not affect the transcript level of *ICE1*, but led to the accumulation of the ICE1

protein to high levels (Supplemental Fig. S3). Next, we examined whether

ICE1 is involved in SA-regulated immunity by conducting a BTH protection

assay. For this assay, wild-type Col-0, *ice1-2,* and *ice1-2/ICE1* plants were

pre-treated with BTH for 24 h before pathogen infiltration, and then their

susceptibility to disease caused by *Pst* DC3000 was evaluated. Compared

with the mock pretreated plants, the BTH-pretreated Col-0 and *ice1-2/ICE1*

plants exhibited greatly reduced disease symptoms on the leaves after *Pst*

DC3000 infection (Fig. 3A and 3B). In contrast, the leaves of *ice1-2* displayed

severe disease symptoms, i.e., severe water soaking symptoms and strong

pathogen growth, after *Pst* DC3000 infection (Fig. 3A, B). The transcript levels

of *PR1* were also lower in BTH-pretreated *ice1-2* plants (Fig. 3C). These

results suggest that ICE1 may be required for effective SA-regulated immunity.

To further confirm the essential role of ICE1 in SA-regulated immunity, we

analyzed the disease symptoms of pretreated Col-0 and *ICE1*-overexpressing

plants (*ICE1-OE 1* and *ICE1-OE 2*). Compared with Col-0 plants,

BTH-pretreated *ICE1*-overexpressing plants showed less severe disease

symptoms after inoculation with *Pst* DC3000, accompanied by BTH-induced

high transcript levels of *PR1* (Fig. 3D–F).

PR1, whose expression is activated by the NPR1–TGA regulation module,

is a principal output gene of SA-associated immunity (Zhang et al., 1999;

Després et al., 2000; Zhou et al., 2000; Johnson et al., 2003). Given that ICE1

interacts with NPR1 and plays an essential role in the SA signaling pathway

(Figures 2 and 3), we wondered whether ICE1 directly regulates the

transcription of *PR1* in a similar manner as TGA transcription factors. ICE1 is a

MYC-like bHLH transcriptional activator that binds specifically to the MYC

recognition sequences (CANNTG, also known as the E-box) in the *CBF3*

promoter (Chinnusamy et al., 2003; Tang et al., 2020). We analyzed the

promoter of PR1 and found that some MYC recognition sequences exist in the

promoter of *PR1*, 1000 bp upstream of ATG (Supplemental Fig. S4). Hence,

we performed electrophoresis mobility shift assays (EMSAs) to determine

whether ICE1 could bind directly to the E-box *cis*-element in the promoter of

PR1. As shown in Fig. 3G, the ICE1 protein was able to bind to the E-box in the

PR1 promoter (Fig. 3G). To verify these findings, we conducted chromatin

immunoprecipitation (ChIP) assays with *ICE1-Myc* (*ICE1-OE 2*) plants treated

or not with BTH. We observed that ICE1 was enriched in the *PR1* promoter

region, and BTH promoted its enrichment (Fig. 3H). Next, a dual-luciferase

reporter assay was conducted to determine the effects of ICE1 on *PR1*

transcription. *LUC* expression driven by the *PR1* promoter was significantly

activated by ICE1 (Fig. 3I). Taken together, these results indicate that ICE1 is

required for effective SA-associated immunity *via* direct activation of *PR1*

transcription.

332   NPR1 Enhances the Ability of ICE1 to Transcriptionally Activate *PR1*

Having established that ICE1 interacts with NPR1 and positively regulates SA

responses through direct binding to the promoter region of *PR1*, we

investigated whether ICE1 functions in a NPR1-relevant manner. To test this

possibility, we generated *ice1-2 npr1-1* double mutant plants by crossing the

ice1-2 mutant with *npr1-1*, a loss-of-function mutant allele of *NPR1* (Col-0

background;(Cao et al., 1997)). Similar to the *ice1-2* and *npr1-1* single mutants,

the *ice1-2 npr1-1* double mutant displayed a highly susceptible phenotype to

Pst DC3000 (Fig. 4A, B), implying that ICE1 may act in the same pathway as

NPR1 to mediate plant immunity.

NPR1 has long been believed to associate with downstream transcription

factors, such as TGAs, to trigger their transcriptional activation of *PR* genes

(Zhang et al., 1999; Després et al., 2000; Zhou et al., 2000; Johnson et al.,

2003; Kumar et al., 2022). Thus, we wondered whether NPR1 could promote

the ability of ICE1 to activate the transcription of *PR1*. To test this possibility,

ChIP assays were conducted with *npr1-1 ICE1-Myc* and *NPR1-HA ICE1-Myc*

plants treated or not with BTH. As shown in Fig. 4C, D, ICE1 was enriched on

the *PR1* promoter in *ICE1-Myc*, and BTH promoted its enrichment. However,

the high enrichment of ICE1 on the promoter of *PR1* was impaired in the

background of *npr1-1*, but enhanced in the *NPR1*-overexpressing line (Fig. 4C,

D). These results suggest that the enrichment of ICE1 on the promoter of *PR1*

is promoted by NPR1.

Next, dual-luciferase reporter assays using Arabidopsis mesophyll

protoplasts were conducted to validate the effect of NPR1 on the

transcriptional activation activity of ICE1. The effectors consisted of GFP,

NPR1, and ICE1 under the control of *Pro35S*, and the reporter consisted of the

PR1 promoter fused to the *LUC* gene (Supplemental Fig. S5). Interestingly,

LUC expression driven by the *PR1* promoter was also induced when NPR1

and GFP were co-expressed, implying that basal TGA or other proteins were

involved. However, *LUC* expression was induced to a much higher level when

NPR1 and ICE1 were co-expressed than when GFP and ICE1 were

co-expressed (Fig. 4E). In contrast, *LUC* expression driven by the *PR1*

promoter was compromised by the loss-of-function of *NPR1* (Fig. 4F). These

results provide further evidence that the ability of ICE1 to activate the

transcription of *PR1* is stimulated by NPR1.

ICE1 Integrates Low Temperature-enhanced and SA-regulated Immunity

ICE1 is involved in low temperature-activated plant immunity (Fig. 1) and

plays a crucial role in the SA signaling pathway (Fig. 3), implying that it may

function as a central node linking these two signaling pathways. To test this

hypothesis, we assessed disease symptoms, including water-soaking

symptoms and pathogen proliferation, in Col-0, *ice1-2,* and *ice1-2/ICE1* plants

treated with BTH for 24 hours with or without a 10-hour low temperature

pretreatment. This allowed us to characterize the critical role of ICE1 in integrating low temperature signals and SA signals to induce pathogen resistance. As shown in Fig. 5A, B, in the leaves of Col-0 and *ice1-2/ICE1,* the water soaking symptoms and pathogen proliferation were alleviated by either BTH or 4°C pretreatment, and significantly diminished by BTH combined with cold treatment, compared with the mock treatment. In contrast, the leaves of *ice1-2* showed severe water soaking symptoms and high pathogen proliferation under all the different treatments. The transcript levels of *PR1* were also lower in *ice1-2* plants compared with Col-0 and *ice1-2/ICE1* plants under different conditions (Fig. 5C). We tested the enrichment of ICE1 on the promoter of *PR1* in the different treatments. As shown in Fig. 5D, accumulation of ICE1 on the promoter of *PR1* was enhanced by the combined low temperature and BTH treatment, compared with either of these treatments alone. Collectively, these results indicate that ICE1 functions as a crucial node by integrating low temperature signals and SA signals to promote plant disease immunity synergistically.

To further explore the underlying molecular mechanism by which ICE1 is synergistically regulated by these two important cues, we determined the amount of ICE1 protein that accumulated in response to a combined low temperature and BTH treatment. Consistent with previous results, ICE1 protein remained stable within 2 hours of 4°C treatment (Ding et al., 2015). Strikingly, ICE1 accumulated to a much higher level under 6 hours BTH treatment combined with 2 hours of 4°C treatment than in either the low temperature or BTH treatment alone (Fig. 5E-F).

ICE1 Physically Interacts with TGA Transcription Factors

TGA transcription factors were confirmed to be interacting proteins of NPR1 in yeast two-hybrid screening assays (Zhang et al., 1999; Zhou et al., 2000), and are believed to be the nuclear targets that bind to the *as-1* element required for SA-induced *PR1* gene expression (Zhang et al., 1999; Zhou et al., 2000; Johnson et al., 2003; Kesarwani et al., 2007; Ding et al., 2018). To explore the regulatory role of ICE1 in SA signaling comprehensively, we investigated whether ICE1 interacts with TGA transcription factors. The full-length coding sequences (CDSs) of *TGA1–TGA7* were fused to pGBK-T7 to construct

pGBK-TGAs vectors, and then each one was transformed into the yeast strain with pGAD-ICE1. Interestingly, ICE1 interacted with all tested TGA proteins in yeast (Fig. 6A). Among the seven members of the TGA family in Arabidopsis, TGA3 plays an important role to bind to the *as-1* element of *PR1* to induce its expression (Kesarwani et al., 2007; Choi et al., 2010; Kumar et al., 2022). Hence, we focused on TGA3 in subsequent experiments.

The results of the BiFC, split luciferase and Co-IP assays further confirmed that ICE1 interacted with TGA3 in plant cells. For the BiFC assays, full-length CDSs of *TGA1* (clade I), *TGA2* (clade II), *TGA3*, and *TGA7* (clade III) were fused to the C-terminal yellow fluorescent protein (cYFP) fragment to generate TGA1-cYFP TGA2-cYFP, TGA3-cYFP, and TGA7-cYFP, respectively, and the full-length CDS of *ICE1* was fused to the N-terminal fragment of YFP (nYFP) to produce ICE1-nYFP. Truncated TGA3$^{193-384}$-cYFP and ICE1$^{1-260}$-nYFP that could not interact with ICE1 or TGA3 were used as negative controls (Supplemental Fig. S6). When ICE1-nYFP was transiently co-expressed with TGAs-cYFP in leaf cells of *N. benthamiana*, recombinant YFP fluorescence was observed in the nucleus, whereas no fluorescence signals were detected when either TGA3$^{193-384}$-nYFP and ICE1-nYFP or ICE1$^{1-260}$-nYFP and TGA3-cYFP were co-expressed (Fig. 6B; Supplemental Fig. S2B).

For the split luciferase assay, TGA3 fused to the C-terminus (cLUC) and ICE1 fused to N-terminal (nLUC) of luciferase were co-expressed in *N. benthamiana* leaves. As shown in Fig. 6C, LUC activity could be detected when TGA3 interacted with ICE1. In the Co-IP assay, ICE1 was immunoprecipitated by anti-GFP agarose beads in Arabidopsis protoplasts co-expressing TGA3-YFP and ICE1-Myc, but not in those co-expressing YFP and ICE1-Myc. This result provided further evidence for the interaction between ICE1 and TGA3 *in vivo* (Fig. 6D). Taken together, these results demonstrate that ICE1 physically interacts with the TGA3 transcription factor *in planta*.

ICE1 and TGA3 Works Synergistically during Low Temperature-enhanced Immunity

Having demonstrated that ICE1 associates with TGA3, we wondered if these two proteins act cooperatively to modulate disease resistance in plants

exposed to cold treatment. Hence, we constructed the *TGA3*-knockout mutant *tga3^{CR}* (Supplemental Fig. S7), and then crossed *ice1-2* with *tga3^{CR}* to obtain *ice1-2 tga3^{CR}* double mutant plants. The disease phenotypes of these plants were then determined. As shown in Fig. 7A, B, compared with the *ice1-2* and *tga3^{CR}* single mutants, the *ice1-2 tga3^{CR}* double mutant showed a more susceptible phenotype, including severe water soaking symptoms and high pathogen proliferation after inoculation with *Pst* DC3000 both at normal and low temperatures, indicating a synergistic role of TGA3 and ICE1 in the regulation of plant immunity.

ICE1 acts as an essential node integrating SA-regulated and low temperature-enhanced immunity by direct enrichment on the promoter of *PR1* (Fig. 5). To further delineate the regulatory relationship between ICE1 and TGA3, we conducted ChIP assays to investigate if TGA3 promotes the enrichment of ICE1 on the promoter of *PR1 in vivo*. For this purpose, *tag3^{CR} ICE1-Myc* and *TGA3-HA ICE1-Myc* seedlings were treated at 4°C for 2 h. Seedlings grown at 22°C served as the control. As shown in Fig. 7C, the enrichment of ICE1 on the *PR1* promoter was decreased in the background of *tag3^{CR}* compared with Col-0. In contrast, the accumulation of ICE1 on the *PR1* promoter was increased in the background of *TGA3-HA-OE* plants compared with Col-0, and was further enhanced under low temperature (Fig. 7D). Dual-luciferase reporter assays using Arabidopsis mesophyll protoplasts were conducted to validate the effect of TGA3 on the transcriptional activation activity of ICE1. The effector consisted of GFP, TGA3, and ICE1 under the control of *Pro35S*, and the reporter consisted of the *PR1* promoter fused to the LUC gene (Supplemental Fig. S5). *LUC* expression driven by the *PR1* promoter was induced to higher level when TGA3 and ICE1 were co-expressed than when GFP and ICE1 were co-expressed (Fig. 7E). In contrast, ICE1 induced *LUC* expression was compromised in the *tag3^{CR}* mutant compared with Col-0 (Fig. 7F). These results indicate that the ability of ICE1 to activate the transcription of *PR1* is promoted by TGA3. Noticeably, the ability of TGA3 to increase the transcription of *PR1* was diminished in the *ice1-2* mutant, suggesting that ICE1 also stimulates the transcriptional activation activities of TGA3 (Fig. 7G). These results support the notion that ICE1 and TGA3 act cooperatively to mediate cold-enhanced plant immunity

through increased enrichment on the *PR1* promoter to activate its expression in association with SA.

The NPR1-TGA3 complex has been well documented that plays an indispensable role in *PR1* expression. Given ICE1 interacts with both NPR1 and TGA3, we questioned whether ICE1 could enhance the transcriptional activities of NPR1-TGA3 module on *PR1* expression. To clarify the potential role of ICE1 during the complex, we conducted the dual-luciferase reporter assay. Noticeably, compared with NPR1 and TGA3 were co-expressed, *LUC* expression driven by the *PR1* promoter was induced higher when NPR1, TGA3 and ICE1 were co-expressed (Fig. 7H), implying that ICE1 promotes the function of NPR1-TGA3 in *PR1* expression regulating.

DISCUSSION

In this study, we demonstrate a previously uncharacterized function of ICE1 as an essential node bridging cold stimulus and plant immunity. We found that ICE1 is involved in cold-activated immunity as well as SA-regulated immunity, as demonstrated by the fact that the *ice1-2* mutant exhibited enhanced susceptibility to pathogen infection, including severe pathogen-related water soaking symptoms and high pathogen proliferation, and decreased expression of *PR1* (Fig. 1 and Fig. 3). Most importantly, the coordinated function of cold- and SA-enhanced immunity was diminished in *ice1-2* mutant plants, suggesting that ICE1 is the crucial regulator linking these two cues (Fig. 5). The major underlying mechanism is that NPR1 associates with ICE1 to promote its ability to activate *PR1* transcription (Fig. 2 and Fig. 4). Additionally, TAG3 works synergistically with ICE1 under low temperature (Fig. 6 and Fig. 7). The relationship between cold stress and immunity has not been elucidated in plants (Wigge, 2013; Kim et al., 2017; Wu et al., 2019). The identification of the NPR1-TGA3-ICE1 regulatory module in this study represents a significant step in understanding SA signaling during cold-activated resistance of plants to pathogen attack.

NPR1, the master regulator of SA signaling, is required for basal and systemically acquired resistance in plants (Cao et al., 1997; Kesarwani et al., 2007). Because it lacks a typical DNA-binding domain, NPR1 was long believed to function as a transcriptional coactivator that interacts with other

proteins, including TGAs, EIN3, and Heat shock transcription factor 1 (HSFA1), to modulate physiological processes in plants (Zhang et al., 1999; Després et al., 2000; Olate et al., 2018; Huang et al., 2020; Wang et al., 2021; Yu et al., 2021). However, in terms of plant immune regulation, TGAs are so far the only group of transcription factors that are known to be direct targets of NPR1 and to activate *PR1* gene expression (Zhang et al., 1999; Després et al., 2000; Zhou et al., 2000; Johnson et al., 2003). Interestingly, in the present study, we demonstrate a novel function of ICE1 to activate SA signaling against pathogen infection. ICE1 interacts with NPR1 (Fig. 2) and functions in the same signal pathway with it (Fig. 4). As a MYC-like bHLH transcription factor, ICE1 can bind directly to the MYC recognition sequences of the *PR1* gene promoter to activate its expression (Supplemental Fig. S4; Fig. 3G-I), and its transcriptional activation activity is enhanced by NPR1 and TGA3 (Fig. 4C–F and Fig. 7C–F). In addition, ICE1 promotes the transcriptional activation of NPR1-TGA3 complex on expression of *PR1* (Fig. 7H). These findings not only provide evidence that ICE1 is an essential component of SA signal pathway at low temperature (Supplemental Fig. S8), but also increase our understanding of the role of the NPR1-ICE1-TGA3 regulatory module in integrating low temperature signal in activating plant immunity.

Recently, increasing attention has been paid to the interactions between low temperature stress and plant immunity. Previous studies have shown that the SA content increases in Arabidopsis plants exposed to cold stress (4°C) for more than 1 week, and that prolonged exposure (3 weeks) to low temperature promotes immunity against the bacterial plant pathogen, *Pst* DC3000 (Kim et al., 2013). Further study has clarified that low temperature can overcome the calmodulin-binding transcriptional activator 3 (CAMTA3) mediated genes inhibition of the SA pathway (Kim et al., 2017). Similarly, moderate low temperature (16°C) also enhances immunity accompanied by up-regulation of SA biosynthesis and signaling genes (Li et al., 2020). Wu et al. found that exposure to short-term cold stress (4°C for 10 hours) activates SA-dependent immunity as well as H_2O_2 and callose deposition (Wu et al., 2019). The results of those studies imply that SA may play a crucial role in low temperature-activated immunity, however, the underlying mechanism remained poorly characterized. In this study, we discovered that ICE1 acts an

indispensable regulatory node bridging low temperature cues and the SA signaling pathway. Our results show that low temperature treatment enhances disease resistance against the bacterial pathogen, *Pst* DC3000 (Fig. 1A–C), and that ICE1 is required for this modulation (Fig. 1E–H). Interestingly, we found that cold stress coordinates SA signaling to promote immunity, and this synergistic regulation occurs in an ICE1-dependent manner. This is highlighted by the fact that the enhanced immunity caused by low temperature combined with SA signaling was compromised in *ice1-2* mutant plants compared with Col-0 plants (Fig. 5A, B). Low temperature has multifaceted influences on the SA signaling pathway, which involves ICE1. For instance, in the Co-IP assays, low temperature facilitated interactions between ICE1 and NPR1 (Fig. 2D). Notably, in the BiFC and LUC assays, the NPR1 and ICE1 interaction only occurred after cold treatment (Fig. 2B, C; Supplemental Fig. S1), consistent with the results of previous studies showing that NPR1 accumulates in response to low temperature and enters the nucleus as a monomer to exert its function as a co-factor (Olate et al., 2018). The enrichment of ICE1–TGA3 on the *PR1* promoter and the ability of this complex to transcriptionally activate *PR1* were found to be significantly enhanced under low temperature (Fig. 7C–G). These results provide direct evidence for the co-operative regulation of low temperature cues and SA signaling *via* ICE1, illuminating the relationship between cold stress and SA-regulated immunity.

It has been proposed that NPR1 is present in both an inactive oligomer form and active monomer form and that SA treatment and low temperature promote the conformation change from oligomer to monomer (Mou et al., 2003; Tada et al., 2008; Yu et al., 2022). Although, in normal temperature and absence of SA, the majority of NPR1 proteins are in the oligomer form, some NPR1 proteins are in the monomer form, which may explain why NPR1 can interact with other proteins, including HSFA1, GID1 and EIN3 in normal temperature and absence of SA, however, the interactions can be enhanced by SA and low temperature (Olate et al., 2018; Huang et al., 2020; Yu et al., 2022). Consistent with these findings, we found that NPR1 could interact with ICE1 in normal temperature, and this process was enhanced by low temperature (Fig. 2B-D). Low temperature may promote NPR1-ICE1 interaction through two mechanisms. On one hand, the conformation of NPR1 may be changed by low temperature

and enhance its interaction with ICE1. On the other hand, low temperature may promote the accumulation of NPR1 monomer, leading to more active NPR1 proteins targeting ICE1.

Temperature is an essential environmental factor affecting plant growth and development, as well as resistance against pathogens. Recent studies have shown that elevated temperatures negatively affect pathogen resistance by suppressing SA production in Arabidopsis (Huot et al., 2017; Kim et al., 2022). Interestingly, the results of several studies, including ours, show that the SA pathway also plays indispensable role in the defense against bacterial pathogen attack under cold stress conditions (Fig. 5) (Seo et al., 2010; Kim et al., 2013; Kim et al., 2017; Wu et al., 2019; Li et al., 2020). Climate change will have a profound effect on plant immunity, so an in-depth understanding of the regulatory role of low temperature on plant immunity is instrumental for revealing the mechanisms of environmental adaptation in plants under cold stress. The results of studies on the temperature regulation of pathogen resistance may provide a theoretical basis for the development of new strategies to improve plants' resistance to pathogens in future agricultural production. In addition, because of their sessile lifestyle, plants are constantly exposed to a variety of abiotic stresses, and endogenous hormone signals often interact with multiple exogenous environmental cues to regulate their resistance to pathogens. Dissecting the regulatory mechanism of the integration of low temperature and SA signals in response to pathogen attack also provides new possibilities for research on the cross-talk between SA signaling and other environmental signals.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

The wild-type and mutant Arabidopsis plants used in this study were in the Columbia (Col-0) genetic background. All Arabidopsis plants were soil-grown for four weeks under an artificial growth chamber at 22°C and 60% relative humidity, and 10 h light/14 h dark photoperiod. The *ice1-2* (SALK_003155;(Kanaoka et al., 2008)) mutant and *npr1-1* (CS3726;(Cao et al., 1994)) mutant have been described. Seeds of *ice1-2/ICE1* were kindly provided by Prof. Shuhua Yang (China Agricultural University). The

35S::GFP-ICE1 (CS68099) line was obtained from the Arabidopsis Resource Center at Ohio State University (http://abrc.osu.edu). To generate the overexpression transgenic plants, the full-length cDNA of *ICE1, TGA3* were cloned into the pOCA30 vector in the sense orientation behind the CaMV 35S promoter. *tga3^{CR}* mutant was constructed by a CRISPR/ Cas9-mediated method as described previously (Liang et al., 2016).

Bacterial Disease Assays

Syringe-infiltration was performed in this study. Briefly, *Pst* DC3000 was cultured in modified Luria–Bertani medium (Xin et al., 2016) containing 100 mg l^{-1} rifampicin at 28 °C to an OD$_{600}$ of 0.8–1.0. Bacteria were collected by centrifugation and re-suspended in 0.25 mM MgCl$_2$. Cell density was adjusted to OD$_{600}$ = 0.2 (approximately 1 × 10^8 cfu ml^{-1}) and further diluted to cell densities of 1 × 10^5–1 × 10^6 cfu ml^{-1}. The infiltrated plants were first placed under ambient humidity for 1–2 h to allow water to evaporate, and then placed in high humidity (about 95%; by covering plants with a clear plastic dome) for disease development after the leaves returned to the state before infiltration.

For quantification of *Pst* DC3000 bacterial populations, leaf disks were taken using a disposable biopsy punch with 4 mm diameter (Integra LifeSciences) and ground in 0.25 mM MgCl$_2$. Colony-forming units were determined by serial dilutions and plating on Luria Marine plates containing 100 mg l^{-1} rifampicin. One biological replicate consists of 12 leaf disks (that is, from 3 leaves of one plant) and at least 3 biological replicates were included in each experiment. Experiments were repeated at least three times.

For BTH protection assay, four-week-old *Arabidopsis thaliana* were sprayed with benzo(1,2,3)thiadiazole-7-carbothioic acid-S-methyl ester (BTH; Chem Service, 0.1% DMSO, 0.05% Silwet L77) or Mock (0.1% DMSO, 0.05% Silwet L77) 24 h before bacterial syringe-infiltration.

RNA Extraction and qRT-PCR

Total RNA was extracted by using Trizol reagent (Invitrogen). The RT-qPCR analysis was conducted as described (He et al., 2023; Li et al., 2023). In brief, 1.0 μg DNase-treated RNA was reverse transcribed in a reaction volume of 20 μl containing oligo-(dT)19 primer and Moloney murine leukemia virus reverse

transcriptase (Thermo Fisher Scientific). RT-qPCR analysis was then

performed with 1.0 µl of 5-fold diluted cDNA using a SYBR premixed Ex Taq kit

(Takara), and a Light Cycler 480 Real-Time PCR System (Roche) for the

reaction. In each RT-qPCR experiment, at least three biological replicates

were used and three technical replicates were performed for each biological

replicate. The *ACTIN2* (*AT3G18780*) gene was used as an internal reference

gene.

Electrophoresis Mobility Shift Assay (EMSA)

To generate the ICE1 recombinant protein, the sequence encoding full-length

ICE1 was cloned into pET28a and transformed into *Escherichia coli* strain

BL21 (DE3). The protein was induced with 0.5 mM isopropyl

β-D-1-thiogalactopyranoside (IPTG) for 18 h at 16°C, and purified with Ni-NTA

resin. EMSA was performed using biotin-labeled probes and

Chemiluminescent EMSA Kit (Beyotime; GS009) referring to the instructions.

The primers used for vector construction are listed in Supplemental Table S1.

Yeast Two-Hybrid Screening and Confirmation

The full-length coding sequences (CDS) of *NPR1* and *TGA* were fused to the

bait vector pGBKT7 (Clontech, Palo Alto, CA, USA) to generate BD-NPR1 and

BD-TGA, while the full-length coding sequence of *ICE1* was ligated to the prey

vector pGADT7 to produce AD-ICE1. To identify structural domains essential

for protein interactions, several truncated NPR1 and TGA3 were cloned into

pGBKT7 and truncated ICE1 into pGADT7. The yeast two hybrid (Y2H) assays

were performed as described previously (Li et al., 2023). The vector pairs were

co-transformed into the yeast strain AH109, protein interactions were indicated

by the ability of cells to grow on dropout medium lacking Leu, Trp, His, and Ade

for 2 days after plating. The primers used for vector construction are listed in

Supplemental Table S1.

Split Luciferase Complementation Assay

678 The split luciferase complementation assays were performed as described

679 previously (Fujikawa and Kato, 2007; Chen et al., 2008). The full-length CDS

680 of *ICE1* was cloned into the pCAMBIA1300-nLUC vector under the control of

the 35S promoter. Full-length CDS of NPR1 and TGA3 were cloned into the pCAMBIA1300-cLUC vectors. After co-expression in *Nicotiana benthamiana* leaves for 48 h，the leaves were sprayed with 1 mM D-luciferin (0.01% Triton X-100) solution and then incubated for 5 min in the dark before detecting. Luciferase intensity was visualized on Tanon-5200 Chemiluminescent Imaging System (Tanon Science&Technology). The primers used for vector construction are listed in Supplemental Table S1.

Bimolecular Fluorescence Complementation Assays

The sequences encoding the C-terminal (64 amino acids) YFP (cYFP) fragment and the N-terminal 173 amino acid-enhanced YFP (nYFP) fragment were amplified by PCR and cloned into the labeled pFGC5941 plasmid to generate pFGC-nYFP and pFGC-cYFP, respectively (Kim et al., 2008). The full-length or truncated CDS sequences of *ICE1* were cloned into pFGC-nYFP to generate the N-terminal in-frame fusions with n-YFP (ICE1-nYFP, ICE1$^{1-260}$-nYFP and ICE1$^{261-494}$-nYFP), whereas full-length or truncated *NPR1* and *TGA* coding sequences were inserted into pFGC-cYFP to form a C-terminal intraframe fusion (NPR1-cYFP, NPR1$^{1-194}$- cYFP, NPR1$^{178-593}$- cYFP and TGA3- cYFP). The plasmids were transformed into *Agrobacterium tumefaciens* strain GV3101 and injected into tobacco (*N. benthamiana*) leaves. As described previously (He et al., 2023; Li et al., 2023), YFP and 4,6-diamidino-2-phenylindole (DAPI) fluorescence of infected leaves were observed under a confocal laser scanning microscope (Olympus) after 48 h of co-expression. Primers used for cloning are listed in Supplemental Table S1.

Co-immunoprecipitation Assays

To generate proteins fused with YFP or Myc tags, we amplified the full-length CDS of *NPR1* and *TGA3* by PCR and then cloned into the pEarleyGate 101 vector with YFP-tagged sequences under the control of *Pro35S* using the gateway method, whereas the full-length CDS of *ICE1* was inserted into the Myc-tagged pGWB517 vector. Arabidopsis protoplasts were transformed with different combinations of plasmids, including 35S::YFP and 35S:: ICE1-Myc, 35S::NPR1-YFP and 35S:ICE1-Myc, 35S::TGA3-YFP and 35S:ICE1-Myc at

22°C or 4°C for 16 h. Total protein was extracted with an IP buffer containing

50 mM MOPs, 5 mM EDTA, 0.2% v/v Triton X-100, 5 mM dithiothreitol, 1 mM

PMSF and 1 x complete protease inhibitor cocktail (Roche).

Immunoprecipitation experiments with GFP-trap beads were performed

according to the manufacturer's protocol. Briefly, cell lysates were incubated

with GFP-trap beads (ChromoTek) at 4°C overnight. After incubation, the

beads were washed three to five times with IP buffer, and then

co-immunoprecipitated proteins were detected by immunoblotting with an

anti-Myc antibody (Abmart; 1:10,000). Primers used for vector construction are

listed in Supplementary Table S1.

ChIP Analyses

ChIP Assays were performed substantially as described earlier

(Mukhopadhyay et al., 2008; Jiang et al., 2014). In brief, Col-0, *ICE1-Myc*,

npr1-1 ICE1-Myc, *NPR1-HA ICE1-Myc*, *TGA3-HA ICE1-Myc*, and *tga3^{CR}*

ICE1-Myc plants were treated with 1% formaldehyde (cross-linking treatment)

to isolate their chromatin. Proteins were immunoprecipitated using anti-Myc

antibody (1:1,000). The precipitated DNA was purified using a PCR purification

kit (Qiagen). To quantitatively determine the ICE1–DNA (target promoter)

binding, RT-qPCR analysis was performed with the *ACTIN2* 3' untranslated

region sequence as an endogenous control. Relative enrichment was

calculated based on DNA binding ratio.

Transient Transcriptional Activation Assays

The full-length CDSs of *ICE1*, *NPR1*, *TGA3*, and *GFP* were cloned into the

pGreenII 62-SK vector as effectors, respectively. The promoter sequence of

PR1 was amplified by PCR and inserted into the pGreenII 0800-LUC vector

(Biovector Science Lab) as the reporter (Hellens et al., 2005). Combinations of

plasmids were transformed into the Col-0 and/or mutant leaf mesophyll

protoplasts according to the Sheen laboratory protocol (Sheen, 2001).

Transfected cells were cultured for 16-18 h followed by analysis of relative

luciferase (LUC) activity using the Dual- Luciferase Reporter Assay System

(Promega), which detects the activities of firefly LUC and the internal control

Renilla reniformis (REN). Primers used for vector construction are listed in

Supplemental Table S1.

STATISTICAL ANALYSIS

Statistical analysis was performed by analysis of variance. The results of

statistical analyses are shown in Supplemental Data Set S1.

ACCESSION NUMBERS

Arabidopsis Genome Initiative numbers for the genes discussed in this article

are as follows: *ICE1*, AT3G26744; *NPR1*, AT1G64280; *TGA1*, AT5G65210;

TGA2, AT5G06950; *TGA3*, AT1G22070; *TGA4*, AT5G10030; *TGA5*,

AT5G06960; *TGA6*, AT3G12250; *TGA7*, AT1G77920; *PR1*, AT2G14610;

ACTIN2, AT3G18780.

ACKNOWLEDGMENTS

We thank Sheng Yang He for his thoughtful revision and constructive

comments on the manuscript. We would like to thank our lab members Diqiu

Yu, Houping Wang and Pengcheng Wang for their suggestions during this

work. We also thank Jennifer Smith, PhD, from Liwen Bianji (Edanz)

(www.liwenbianji.cn/) for editing the English text of a draft of this manuscript.

AUTHOR CONTRIBUTIONS

S.Q.L. and Y.J.J. designed and performed experiments; S.Q.L., L.H., Y.P.Y.

and X.H. and Y.R.H analyzed data; S.Q.L. and Y.J.J. wrote the manuscript. All

authors read and approved the final manuscript.

SUPPLEMENTAL DATA

The following materials are available in the online version of this article.

Supplemental Figure S1. No interacting signal of ICE1 and NPR1 was

detected at normal temperature.

Supplemental Figure S2. Percentage of transformed cells with YFP

fluorescence in the BiFC assays presented in Figures 2B and 6B.

Supplemental Figure S3. Transcriptional and protein levels of ICE1

responding to BTH treatment.

Supplemental Figure S4. The promoter sequence of *PR1*.

Supplemental Figure S5. Schematic of effectors and reporters used in transient transactivation assays.

Supplemental Figure S6. The bHLH motif and bZIP motif was required for interaction of ICE1 and TGA3.

Supplemental Figure S7. Generation of *tga3*^CR knockout by CRISPR/Cas9 technology.

Supplemental Figure S8. Working model of NPR1-TGA3/ICE1 regulating module against *Pst* DC3000 after cold stress.

Supplemental Figure S9. Uncropped images for protein gels.

Supplemental Table S1. Primers used for generating various clones.

Supplemental Data Set S1. The results of statistical analyses performed by ANOVA.

FUNDING

This study was supported by funding from the Natural Science Foundation of China (Grants 32360082 to Y.J.J.) and the Young and Middle Aged Academic and technical Leaders of Yunnan Province (202105AC160028 to Y.J.J.).

Conflict of interest statement. All authors state that they have no conflict of interest in relation to this research.

DATA AVAILABILITY

All data supporting the findings of this study are available within the article and its supplemental materials.

REFERENCES

An, J.P., Wang, X.F., Zhang, X.W., You, C.X., and Hao, Y.J. (2021). Apple B-box protein BBX37 regulates jasmonic acid mediated cold tolerance through the JAZ-BBX37-ICE1-CBF pathway and undergoes MIEL1-mediated ubiquitination and degradation. New Phytol 229, 2707-2729.

An, J.P., Xu, R.R., Liu, X., Su, L., Yang, K., Wang, X.F., Wang, G.L., and You, C.X. (2022). Abscisic acid insensitive 4 interacts with ICE1 and JAZ proteins

to regulate ABA signaling-mediated cold tolerance in apple. J Exp Bot 73,

980-997.

Benedict, C., Geisler, M., Trygg, J., Huner, N., and Hurry, V. (2006).

Consensus by democracy. Using meta-analyses of microarray and genomic

data to model the cold acclimation signaling pathway in Arabidopsis. Plant

Physiol 141, 1219-1232.

Boller, T., and Felix, G. (2009). A renaissance of elicitors: perception of

microbe-associated molecular patterns and danger signals by

pattern-recognition receptors. Annu Rev of Plant Biol 60, 379-406.

Cao, H., Bowling, S.A., Gordon, A.S., and Dong, X. (1994). Characterization

of an Arabidopsis mutant that is nonresponsive to inducers of systemic

acquired resistance. Plant Cell 6, 1583-1592.

Cao, H., Glazebrook, J., Clarke, J.D., Volko, S., and Dong, X. (1997). The

Arabidopsis *NPR1* gene that controls systemic acquired resistance encodes

a novel protein containing ankyrin repeats. Cell 88, 57-63.

Chen, H., Zou, Y., Shang, Y., Lin, H., Wang, Y., Cai, R., Tang, X., and Zhou,

J.M. (2008). Firefly luciferase complementation imaging assay for

protein-protein interactions in plants. Plant Physiol 146, 368-376.

Chinnusamy, V., Zhu, J., and Zhu, J.K. (2007). Cold stress regulation of gene

expression in plants. Trends Plant Sci 12, 444-451.

Chinnusamy, V., Ohta, M., Kanrar, S., Lee, B.H., Hong, X., Agarwal, M., and

Zhu, J.K. (2003). ICE1: a regulator of cold-induced transcriptome and

freezing tolerance in Arabidopsis. Genes Dev 17, 1043-1054.

Choi, J., Huh, S.U., Kojima, M., Sakakibara, H., Paek, K.H., and Hwang, I.

(2010). The cytokinin-activated transcription factor ARR2 promotes plant

immunity via TGA3/NPR1-dependent salicylic acid signaling in Arabidopsis.

Dev Cell 19, 284-295.

Deslandes, L., and Rivas, S. (2012). Catch me if you can: bacterial effectors

and plant targets. Trends Plant Sci 17, 644-655.

Després, C., DeLong, C., Glaze, S., Liu, E., and Fobert, P.R. (2000). The

Arabidopsis NPR1/NIM1 protein enhances the DNA binding activity of a

subgroup of the TGA family of bZIP transcription factors. Plant Cell 12,

279-290.

Ding, Y., Li, H., Zhang, X., Xie, Q., Gong, Z., and Yang, S. (2015). OST1

kinase modulates freezing tolerance by enhancing ICE1 stability in

Arabidopsis. Dev Cell 32, 278-289.

Ding, Y., Sun, T., Ao, K., Peng, Y., Zhang, Y., Li, X., and Zhang, Y. (2018).

Opposite roles of salicylic acid receptors NPR1 and NPR3/NPR4 in

transcriptional regulation of plant immunity. Cell 173, 1454-1467.e1415.

Dodds, P.N., and Rathjen, J.P. (2010). Plant immunity: towards an integrated

view of plant-pathogen interactions. Nature Rev Genet 11, 539-548.

Dong, C.H., Agarwal, M., Zhang, Y., Xie, Q., and Zhu, J.K. (2006). The

negative regulator of plant cold responses, HOS1, is a RING E3 ligase that

mediates the ubiquitination and degradation of ICE1. Proc Natl Acad Sci

USA 103, 8281-8286.

Durrant, W.E., and Dong, X. (2004). Systemic acquired resistance. Annu Rev

Phytopathol 42, 185-209.

Fujikawa, Y., and Kato, N. (2007). Split luciferase complementation assay to

study protein-protein interactions in Arabidopsis protoplasts. Plant J 52,

185-195.

Gilmour, S.J., Zarka, D.G., Stockinger, E.J., Salazar, M.P., Houghton, J.M.,

and Thomashow, M.F. (1998). Low temperature regulation of the

Arabidopsis CBF family of AP2 transcriptional activators as an early step in

cold-induced COR gene expression. Plant J 16, 433-442.

Griffith, M., and Yaish, M.W. (2004). Antifreeze proteins in overwintering

plants: a tale of two activities. Trends Plant Sci 9, 399-405.

Han, Q., Tan, W., Zhao, Y., Yang, F., Yao, X., Lin, H., and Zhang, D. (2022).

Salicylic acid-activated BIN2 phosphorylation of TGA3 promotes Arabidopsis

PR gene expression and disease resistance. EMBO J, e110682.

He, K., Du, J., Han, X., Li, H., Kui, M., Zhang, J., Huang, Z., Fu, Q., Jiang, Y.,

and Hu, Y. (2023). PHOSPHATE STARVATION RESPONSE1 (PHR1)

interacts with JASMONATE ZIM-DOMAIN (JAZ) and MYC2 to modulate

phosphate deficiency-induced jasmonate signaling in Arabidopsis. Plant Cell

35, 2132-2156.

Hellens, R.P., Allan, A.C., Friel, E.N., Bolitho, K., Grafton, K., Templeton,

M.D., Karunairetnam, S., Gleave, A.P., and Laing, W.A. (2005). Transient

expression vectors for functional genomics, quantification of promoter

activity and RNA silencing in plants. Plant Methods 1, 13.

Hu, Y., Han, X., Yang, M., Zhang, M., Pan, J., and Yu, D. (2019). The
transcription factor INDUCER OF CBF EXPRESSION1 interacts with
ABSCISIC ACID INSENSITIVE5 and DELLA proteins to fine-tune abscisic
acid signaling during seed germination in Arabidopsis. Plant Cell 31,
1520-1538.

Huang, P., Dong, Z., Guo, P., Zhang, X., Qiu, Y., Li, B., Wang, Y., and Guo,
H. (2020). Salicylic acid suppresses apical hook formation via
NPR1-mediated repression of EIN3 and EIL1 in Arabidopsis. Plant Cell 32,
612-629.

Huang, X., Li, J., Bao, F., Zhang, X., and Yang, S. (2010). A gain-of-function
mutation in the Arabidopsis disease resistance gene *RPP4* confers
sensitivity to low temperature. Plant Physiol 154, 796-809.

Huot, B., Castroverde, C.D.M., Velasquez, A.C., Hubbard, E., Pulman, J.A.,
Yao, J., Childs, K.L., Tsuda, K., Montgomery, B.L., and He, S.Y. (2017).
Dual impact of elevated temperature on plant defence and bacterial
virulence in Arabidopsis. Nat Commun 8, 1808.

Jiang, Y.J., Liang, G., Yang, S.Z., and Yu, D.Q. (2014). Arabidopsis WRKY57
functions as a node of convergence for jasmonic acid- and auxin-mediated
signaling in jasmonic acid-induced leaf senescence. Plant Cell 26, 230-245.

Johnson, C., Boden, E., and Arias, J. (2003). Salicylic acid and NPR1 induce
the recruitment of trans-activating TGA factors to a defense gene promoter
in Arabidopsis. Plant Cell 15, 1846-1858.

Jones, J.D., and Dangl, J.L. (2006). The plant immune system. Nature 444,
323-329.

Kanaoka, M.M., Pillitteri, L.J., Fujii, H., Yoshida, Y., Bogenschutz, N.L.,
Takabayashi, J., Zhu, J.K., and Torii, K.U. (2008). SCREAM/ICE1 and
SCREAM2 specify three cell-state transitional steps leading to Arabidopsis
stomatal differentiation. Plant Cell 20, 1775-1785.

Kesarwani, M., Yoo, J., and Dong, X. (2007). Genetic interactions of TGA
transcription factors in the regulation of pathogenesis-related genes and
disease resistance in Arabidopsis. Plant Physiol 144, 336-346.

Kim, J.H., Castroverde, C.D.M., Huang, S., Li, C., Hilleary, R., Seroka, A.,
Sohrabi, R., Medina-Yerena, D., Huot, B., Wang, J., Nomura, K., et al.
(2022). Increasing the resilience of plant immunity to a warming climate.

Nature 607, 339-344.

Kim, K.C., Lai, Z., Fan, B., and Chen, Z. (2008). Arabidopsis WRKY38 and
WRKY62 transcription factors interact with histone deacetylase 19 in basal
defense. Plant Cell 20, 2357-2371.

Kim, Y., Park, S., Gilmour, S.J., and Thomashow, M.F. (2013). Roles of
CAMTA transcription factors and salicylic acid in configuring the
low-temperature transcriptome and freezing tolerance of Arabidopsis. Plant
J 75, 364-376.

Kim, Y.S., Lee, M., Lee, J.H., Lee, H.J., and Park, C.M. (2015). The unified
ICE-CBF pathway provides a transcriptional feedback control of freezing
tolerance during cold acclimation in Arabidopsis. Plant Mol Biol 89, 187-201.

Kim, Y.S., An, C., Park, S., Gilmour, S.J., Wang, L., Renna, L., Brandizzi, F.,
Grumet, R., and Thomashow, M.F. (2017). CAMTA-mediated regulation of
salicylic acid immunity pathway genes in Arabidopsis exposed to low
temperature and pathogen infection. Plant Cell 29, 2465-2477.

Nomura, K., He, S.Y. (2005). Powerful screens for bacterial virulence proteins.
Proc Natl Acad Sci USA 102, 3527-3528.

Kumar, S., Zavaliev, R., Wu, Q., Zhou, Y., Cheng, J., Dillard, L., Powers, J.,
Withers, J., Zhao, J., Guan, Z., Borgnia, M.J., Bartesaghi, A., Dong, X.,
and Zhou, P. (2022). Structural basis of NPR1 in activating plant immunity.
Nature 605, 561-566.

Lee, B.H., Henderson, D.A., and Zhu, J.K. (2005). The Arabidopsis
cold-responsive transcriptome and its regulation by ICE1. Plant Cell 17,
3155-3175.

Lee, J.H., Jung, J.H., and Park, C.M. (2015). INDUCER OF CBF
EXPRESSION 1 integrates cold signals into FLOWERING LOCUS
C-mediated flowering pathways in Arabidopsis. Plant J 84, 29-40.

Lee, J.H., Jung, J.H., and Park, C.M. (2017). Light inhibits COP1-mediated
degradation of ICE transcription factors to induce stomatal development in
Arabidopsis. Plant Cell 29, 2817-2830.

Li, H., Ding, Y., Shi, Y., Zhang, X., Zhang, S., Gong, Z., and Yang, S. (2017).
MPK3- and MPK6-mediated ICE1 phosphorylation negatively regulates
ICE1 stability and freezing tolerance in Arabidopsis. Dev Cell 43,
630-642.e634.

Li, Z., Li, S., Jin, D., Yang, Y., Pu, Z., Han, X., Hu, Y., and Jiang, Y. (2023).
U-box E3 ubiquitin ligase PUB8 attenuates abscisic acid responses during
early seedling growth. Plant Physiol 191, 2519-2533.

Li, Z., Liu, H., Ding, Z., Yan, J., Yu, H., Pan, R., Hu, J., Guan, Y., and Hua, J.
(2020). Low temperature enhances plant immunity via salicylic acid pathway
genes that are repressed by ethylene. Plant Physiol 182, 626-639.

Liang, G., Zhang, H., Lou, D., and Yu, D. (2016). Selection of highly efficient
sgRNAs for CRISPR/Cas9-based plant genome editing. Sci Rep 6, 21451.

Miura, K., Jin, J.B., Lee, J., Yoo, C.Y., Stirm, V., Miura, T., Ashworth, E.N.,
Bressan, R.A., Yun, D.J., and Hasegawa, P.M. (2007). SIZ1-mediated
sumoylation of ICE1 controls *CBF3/DREB1A* expression and freezing
tolerance in Arabidopsis. Plant Cell 19, 1403-1414.

Mou, Z., Fan, W., and Dong, X. (2003). Inducers of plant systemic acquired
resistance regulate NPR1 function through redox changes. Cell 113,
935-944.

Mukhopadhyay, A., Deplancke, B., Walhout, A.J., and Tissenbaum, H.A.
(2008). Chromatin immunoprecipitation (ChIP) coupled to detection by
quantitative real-time PCR to study transcription factor binding to DNA in
Caenorhabditis elegans. Nat Protoc 3, 698-709.

Nomoto, M., Skelly, M.J., Itaya, T., Mori, T., Suzuki, T., Matsushita, T.,
Tokizawa, M., Kuwata, K., Mori, H., Yamamoto, Y.Y., et al. (2021).
Suppression of MYC transcription activators by the immune cofactor NPR1
fine-tunes plant immune responses. Cell Rep 37, 110125.

Olate, E., Jiménez-Gómez, J.M., Holuigue, L., and Salinas, J. (2018). NPR1
mediates a novel regulatory pathway in cold acclimation by interacting with
HSFA1 factors. Nat Plants 4, 811-823.

Orvar, B.L., Sangwan, V., Omann, F., and Dhindsa, R.S. (2000). Early steps
in cold sensing by plant cells: the role of actin cytoskeleton and membrane
fluidity. Plant J 23, 785-794.

Sarkar, S., Das, A., Khandagale, P., Maiti, I.B., Chattopadhyay, S., and Dey,
N. (2018). Interaction of Arabidopsis TGA3 and WRKY53 transcription
factors on *Cestrum yellow leaf curling virus* (CmYLCV) promoter mediates
salicylic acid-dependent gene expression in planta. Planta 247, 181-199.

Seo, P.J., Kim, M.J., Park, J.Y., Kim, S.Y., Jeon, J., Lee, Y.H., Kim, J., and

Park, C.M. (2010). Cold activation of a plasma membrane-tethered NAC
transcription factor induces a pathogen resistance response in Arabidopsis.
Plant J 61, 661-671.

Sheen, J. (2001). Signal transduction in maize and Arabidopsis mesophyll
protoplasts. Plant Physiol 127, 1466-1475.

Spoel, S.H., and Dong, X. (2012). How do plants achieve immunity? Defence
without specialized immune cells. Nat Rev Immunol 12, 89-100.

Stockinger, E.J., Gilmour, S.J., and Thomashow, M.F. (1997). *Arabidopsis*
thaliana CBF1 encodes an AP2 domain-containing transcriptional activator
that binds to the C-repeat/DRE, a *cis*-acting DNA regulatory element that
stimulates transcription in response to low temperature and water deficit.
Proc Natl Acad Sci USA 94, 1035-1040.

Tada, Y., Spoel, S.H., Pajerowska-Mukhtar, K., Mou, Z., Song, J., Wang, C.,
Zuo, J., and Dong, X. (2008). Plant immunity requires conformational
changes of NPR1 via S-nitrosylation and thioredoxins. Science 321,
952-956.

Tang, K., Zhao, L., Ren, Y., Yang, S., Zhu, J.K., and Zhao, C. (2020). The
transcription factor ICE1 functions in cold stress response by binding to the
promoters of *CBF* and *COR* genes. J Integr Plant Biol 62, 258-263.

Vlot, A.C., Dempsey, D.A., and Klessig, D.F. (2009). Salicylic Acid, a
multifaceted hormone to combat disease. Annu Rev Phytopathol 47,
177-206.

Wang, C., Dai, S., Zhang, Z.L., Lao, W., Wang, R., Meng, X., and Zhou, X.
(2021). Ethylene and salicylic acid synergistically accelerate leaf
senescence in Arabidopsis. J Integr Plant Biol 63, 828-833.

Wang, W., Withers, J., Li, H., Zwack, P.J., Rusnac, D.V., Shi, H., Liu, L., Yan,
S., Hinds, T.R., Guttman, M., Dong, X., and Zheng, N. (2020). Structural
basis of salicylic acid perception by Arabidopsis NPR proteins. Nature 586,
311-316.

Wang, Y., Bao, Z., Zhu, Y., and Hua, J. (2009). Analysis of temperature
modulation of plant defense against biotrophic microbes. Mol Plant Microbe
Interact 22, 498-506.

Wigge, P.A. (2013). Ambient temperature signalling in plants. Curr Opin Plant
Biol 16, 661-666.

Wu, Z., Han, S., Zhou, H., Tuang, Z.K., Wang, Y., Jin, Y., Shi, H., and Yang,
W. (2019). Cold stress activates disease resistance in *Arabidopsis thaliana*
through a salicylic acid dependent pathway. Plant Cell Environ 42,
2645-2663.

Xin, X.F., Nomura, K., Aung, K., Velasquez, A.C., Yao, J., Boutrot, F.,
Chang, J.H., Zipfel, C., and He, S.Y. (2016). Bacteria establish an aqueous
living space in plants crucial for virulence. Nature 539, 524-529.

Yang, H., Shi, Y., Liu, J., Guo, L., Zhang, X., and Yang, S. (2010). A mutant
CHS3 protein with TIR-NB-LRR-LIM domains modulates growth, cell death
and freezing tolerance in a temperature-dependent manner in Arabidopsis.
Plant J 63, 283-296.

Yang, S., and Hua, J. (2004). A haplotype-specific resistance gene regulated
by BONZAI1 mediates temperature-dependent growth control in Arabidopsis.
Plant Cell 16, 1060-1071.

Ye, K., Li, H., Ding, Y., Shi, Y., Song, C., Gong, Z., and Yang, S. (2019).
BRASSINOSTEROID-INSENSITIVE2 negatively regulates the stability of
transcription factor ICE1 in response to cold stress in Arabidopsis. Plant Cell
31, 2682-2696.

Yu, X., Xu, Y., and Yan, S. (2021). Salicylic acid and ethylene coordinately
promote leaf senescence. J Integr Plant Biol 63, 823-827.

Yu, X., Cui, X., Wu, C., Shi, S., and Yan, S. (2022). Salicylic acid inhibits
gibberellin signaling through receptor interactions. Mol Plant 15, 1759-1771.

Zavaliev, R., Mohan, R., Chen, T., and Dong, X. (2020). Formation of NPR1
condensates promotes cell survival during the plant immune response. Cell
182, 1093-1108.e1018.

Zhang, Y., Fan, W., Kinkema, M., Li, X., and Dong, X. (1999). Interaction of
NPR1 with basic leucine zipper protein transcription factors that bind
sequences required for salicylic acid induction of the *PR-1* gene. Proc Natl
Acad Sci USA 96, 6523.

Zhao, C., Wang, P., Si, T., Hsu, C.C., Wang, L., Zayed, O., Yu, Z., Zhu, Y.,
Dong, J., Tao, W.A., et al. (2017). MAP kinase cascades regulate the cold
response by modulating ICE1 protein stability. Dev Cell 43, 618-629.e615.

Zhou, J.M., Trifa, Y., Silva, H., Pontier, D., Lam, E., Shah, J., and Klessig,
D.F. (2000). NPR1 differentially interacts with members of the TGA/OBF

family of transcription factors that bind an element of the *PR-1* gene required

for induction by salicylic acid. Mol Plant Microbe Interact 13, 191-202.

Figure 1. ICE1 involves in low temperature-enhanced plant immunity. (A) Col-0 leaves infiltrated with *Pst* DC3000 (OD_{600} = 0.0001) after 3 weeks of 4°C treatments (down), 22°C as control (up). (B) Bacterial populations in leaves described in (A). Values are displayed as mean ± SEM (n ≥ 5 biological replicates). (C) Col-0 leaves infiltrated with *Pst* DC3000 (OD600 = 0.0001) after 10 hours of 4°C treatments (down), 22°C as control (up). (D) Bacterial populations on leaves described in (C). Values are displayed as mean ± SEM (n = 10 biological replicates). (E) Col-0, *ice1-2*, *ice1-2/ICE1* leaves infiltrated with *Pst* DC3000 (OD_{600} = 0.0001) after 10 hours of 4°C treatments (down), 22°C as control (up). (F) Bacterial populations on leaves described in (E). Values are displayed as mean ± SEM (n ≥ 5 biological replicates). (G) Col-0, *ICE1-OE 1* and *ICE1-OE 2* leaves infiltrated with *Pst* DC3000 (OD_{600} = 0.0001) after 10 hours of 4°C treatment (down), 22°C as control (up). (H) Bacterial populations of *Pst* DC3000 on plants described in (G). Values are displayed as mean ± SEM (n = 4 biological replicates). Photos were taken after 48 h pathogen inoculation. Different letters indicate statistically significant differences (two-way ANOVA, *P* < 0.05). Experiments were repeated at least three times with similar trends.

Figure 2. ICE1 interacts with NPR1. (A) Yeast two hybrid assays showing the interactions between ICE1, NPR1, and their truncated versions. Protein interactions were indicated by the growth of yeast cells after 2 days of incubation in dropout medium lacking Leu, Trp, His and Ade. The numbers indicate the positions of amino acids, pGBKT7 (BD) and pGADT7 (AD) were used as negative controls. (B) BiFC assay. Fluorescence was observed in the nucleus of transformed *N. benthamiana* cells co-expressing ICE1-nYFP (or ICE1$^{261-494}$-nYFP) with NPR1-cYFP or ICE1-nYFP with NPR1$^{1-191}$-cYFP at 4°C for 4 h. No signal was obtained in the negative controls which NPR1-cYFP (or NPR1$^{1-194}$-cYFP) with ICE1$^{1-260}$-nYFP or ICE1-nYFP (or ICE1$^{261-494}$-nYFP) with NPR1$^{370-593}$-cYFP were co-expressed. Nuclei are indicated by DAPI staining. (C) Interaction between ICE1 and NPR1 in split luciferase assay. ICE1-cLUC and NPR1-nLUC were co-expressed in *N. benthamiana* leaves, the luminescence intensity was detected at 4°C for 4 h after 48 h of incubation by an imaging system. GUS-nLUC and GUS-cLUC were set as a negative control. (D) Co-IP assay. Arabidopsis protoplasts expressing NPR1-YFP with ICE1-Myc, YFP with ICE1-Myc were incubated for 16 h. For low-temperature induction, protoplasts were processed at 4°C for 2 h before total protein was extracted. Then immunoprecipitated with GFP-trap agarose beads. Co-immunoprecipitated proteins were detected with anti-Myc antibody. Experiments were repeated at least three times with similar trends.

Figure 3. ICE1 involves in SA-regulated immunity. (A) Col-0, *ice1-2*, *ice1-2/ICE1* leaves infiltrated with *Pst* DC3000 (OD_{600} = 0.0001) after 24 hours pretreating with 100 μM BTH (down), mock as control (up). (B) Bacterial populations in leaves described in (A). Values are displayed as mean ± SEM (n ≥ 4 biological replicates). (C) qRT-PCR analysis of *PR1* expression in leaves described in (A). Values are displayed as mean ± SEM (n = 3 biological replicates). (D) Col-0, *ICE1-OE 1* and *ICE1-OE 2* leaves infiltrated with *Pst* DC3000 (OD_{600} = 0.0001) after 24 hours pretreating with 100 μM BTH (down), mock as control (up). (E) Bacterial populations in leaves described in (D). Values are displayed as mean ± SEM (n = 3 biological replicates). (F) qRT-PCR analysis of *PR1* expression in leaves described in (C). Values are displayed as mean ± SEM (n = 3 biological replicates). (G) EMSA assay examines ability of ICE1 binding to the E-box (CANNTG) of *PR1* promoter. Biotin-labeled probes were incubated with ICE1-His proteins, and the free and bound DNAs were separated on an acrylamide gel. As indicated, unlabeled probe (cold-PR1) was used as competitor and biotin labeled mutant probe (mutant-PR1) was used as negative control. Mutated probe in which the

CATTTG motif was replaced with AATTTC. (H) ChIP-qPCR analysis of the relative enrichment of ICE1 on the promoter of *PR1*. Col-0 and *ICE1*-overexpressing (*ICE1-Myc*) plants were treated with mock or 100 μM BTH for 6 h and pooled for ChIP assays using anti-Myc antibody, the *ACTIN2* untranslated region sequence (*pACTIN2*) as a negative control. Values are displayed as mean ± SEM (n ≥ 3 biological replicates). (I) Transient dual-LUC reporter assays showing that ICE1 activates the expression of *PR1*. Values are displayed as mean ± SEM (n ≥ 3 biological replicates). Each biological replicate was from different leaves of more than 60 Col-0 plants. Photos were taken after 48 h pathogen inoculation in A and D. Different letters indicate statistically significant differences (two-way ANOVA, *P* < 0.05). Experiments were repeated at least three times with similar trends.

Figure 4. NPR1 promotes the binding and transcriptional activation of ICE1 to *PR1*. (A) Col-0, *ice1-2*, *npr1-1* and *ice1-2 npr1-1* leaves infiltrated with *Pst* DC3000 (OD_{600} = 0.0001). Photos were taken after 48 h pathogen inoculation. (B) Bacterial populations in leaves described in (A). Values are displayed as mean ± SEM (n ≥ 6 biological replicates). (C and D) ChIP-qPCR analysis of the relative enrichment of ICE1 on the promoter regions of *PR1*. Col-0, *ICE1*-overexpressing (*ICE1-Myc*), *npr1-1 ICE1-Myc* and *NPR1-HA ICE1-Myc* plants were treated with mock or 100 µM BTH for 6 h and pooled for ChIP assays using anti-Myc antibody, the *ACTIN2* untranslated region sequence (*pACTIN2*) as a negative control. Values are displayed as mean ± SEM (n = 3 biological replicates). (E) Transient transcriptional activity assays showing that NPR1 enhances the transcriptional activation of ICE1 to *PR1*. Values are displayed as mean ± SEM (n = 3 biological replicates). (F) Transient transcriptional activity assays showing that activation of the *PR1* promoter by ICE1 is compromised in the *npr1-1* mutant. Values are displayed as mean ± SEM (n = 3 biological replicates). Each biological replicate was from different leaves of more than 60 plants in (E) and (F). Different letters indicate statistically significant differences (two-way ANOVA, *P* < 0.05). Experiments were repeated at least three times with similar results.

Figure 5. ICE1 links low temperature signals and SA pathway to promote plant immunity. (A) Col-0, *ice1-2* and *ice1-2/ICE1* leaves infiltrated with *Pst* DC3000 (OD$_{600}$ = 0.0001) after BTH with low temperature pretreating (briefly, 14 hours BTH treatment at 22°C and then move to 4°C growth chamber for 10 hours cold treatment). Photos were taken after 48 h pathogen inoculation. (B) Bacterial populations in leaves described in (A). Values are displayed as mean ± SEM (n = 5 biological replicates). (C) qRT-PCR analysis of *PR1* expression in leaves described in (A). Values are displayed as mean ± SEM (n = 3 biological replicates). (D) ChIP-qPCR analysis of the relative enrichment of ICE1 on the promoter regions of *PR1*. *ICE1*-overexpressing plants (*ICE1-Myc*) were treated with 100 μM BTH for 6 hours and 4°C for 2 hours and pooled for ChIP assays using anti-Myc antibody, the *ACTIN2* untranslated region sequence (*pACTIN2*) as a negative control. Values are displayed as mean ± SEM (n = 3 biological replicates). (E) Immunoblot analyzing the accumulation of ICE1 protein. Ten-day-old *ICE1*-overexpressing seedlings (*ICE1-Myc*) were treated with 100 μM BTH for 6 h and/or 4°C for 2 hours before protein extraction. The accumulation of ICE1 protein was detected with anti-Myc antibody. (F) Relative protein levels of ICE1-Myc in (E), which were quantified

by Image J. Values are displayed as mean ± SEM (n = 3 biological replicates). Different letters indicate statistically significant differences (two-way ANOVA, $P < 0.05$). Experiments were repeated at least three times with similar trends.

Figure 6. ICE1 interacts with TGA transcription factors. (A) Yeast two hybrid assays. Interactions of ICE1 with TGAs were indicated by the ability of yeast cells to grow on dropout medium lacking Leu, Trp, His, and Ade for 2 days after plating. BD and AD were used as negative controls. (B) BiFC assay. Fluorescence was observed in the nuclear compartment of transformed *N. benthamiana* cells, resulting from the complementation of ICE1-nYFP with TGA1/2/3/7-cYFP. No signal was obtained for the negative controls in which ICE1$^{1-260}$-nYFP with TGA3-cYFP or ICE1-nYFP with TGA3$^{193-384}$-cYFP were coexpressed. Nuclei are indicated by DAPI staining. (C) Split luciferase assay. nLUC-ICE1 and cLUC-TGA3 constructs were co-transformed into *N. benthamiana*. Luminescence intensity was measured after 48 h of incubation by an imaging system. GUS-nLUC and GUS-cLUC were set as a negative control. (D) Co-IP assay. Arabidopsis protoplasts expressing TGA3-YFP with ICE1-Myc, YFP with ICE1-Myc were incubated for 16 h, and total protein was extracted. Then immunoprecipitated with GFP-trap agarose beads. Co-immunoprecipitated proteins were detected with anti-Myc antibody. Experiments were repeated at least three times with similar trends.

Figure 7. ICE1 and TGA3 works synergistically to activate expression of *PR1* during low temperature stress. (A) Col-0, *ice1-2*, *tga3^CR* and *ice1-2 tga3^CR* leaves infiltrated with *Pst* DC3000 (OD$_{600}$ = 0.0001) after 10 hours of 4°C treatment (down), 22°C as control (up). Photos were taken after 48 h pathogen inoculation. (B) Bacterial populations in leaves described in (A). Values are displayed as mean ± SEM (n = 4 biological replicates). (C) ChIP-qPCR analysis of the *tga3^CR* mutation impaired the relative enrichment of ICE1 to the *PR1* promoter. Col-0, *ICE1*-overexpressing (*ICE1-Myc*) and *tga3^CR ICE1-Myc* plants were treated for 10 h of 4°C, 22°C as control. (D) ChIP-qPCR analyses of the relative enrichment of ICE1 to the *PR1* promoter enhanced by TGA3. Col-0, *ICE1*-overexpressing plants (*ICE1-Myc*) and *TGA3-HA ICE1-Myc* (hybridized F1 generation) were treated for 10 h of 4°C, 22°C as control. Treated leaves were pooled for ChIP assays using anti-Myc antibody, the *ACTIN2* untranslated region sequence (*pACTIN2*) as a negative control in (C) and (D). (E) Transient transcriptional activity assays showing that TGA3 enhances the transcriptional activation of ICE1 to *PR1*. (F) Transient transcriptional activity assays showing that activation of the *PR1* promoter by ICE1 is compromised in the *tga3^CR* mutant. (G) Transient transcriptional activity assays showing that activation of the *PR1* promoter by ICE1 is diminished in the *ice1-2* mutant. Each biological replicate was from different leaves of more than 60 plants in (E-G). (H) Transient transcriptional activity

assays showing that ICE1 enhances the transcriptional activation of NPR1-ICE1 complex to *PR1* expression. Values are displayed as mean ± SEM (n = 3 biological replicates) for (C-H). Different letters indicate statistically significant differences (two-way ANOVA, *P* < 0.05). Experiments were repeated at least three times with similar trends.

Parsed Citations

An, J.P., Wang, X.F., Zhang, X.W., You, C.X., and Hao, Y.J. (2021). Apple B-box protein BBX37 regulates jasmonic acid mediated cold tolerance through the JAZ-BBX37-ICE1-CBF pathway and undergoes MIEL1-mediated ubiquitination and degradation. New Phytol 229, 2707-2729.
  Google Scholar: Author Only Title Only Author and Title

An, J.P., Xu, R.R., Liu, X., Su, L., Yang, K., Wang, X.F., Wang, G.L., and You, C.X. (2022). Abscisic acid insensitive 4 interacts with ICE1 and JAZ proteins to regulate ABA signaling-mediated cold tolerance in apple. J Exp Bot 73, 980-997.
  Google Scholar: Author Only Title Only Author and Title

Benedict, C., Geisler, M., Trygg, J., Huner, N., and Hurry, V. (2006). Consensus by democracy. Using meta-analyses of microarray and genomic data to model the cold acclimation signaling pathway in Arabidopsis. Plant Physiol 141, 1219-1232.
  Google Scholar: Author Only Title Only Author and Title

Boller, T., and Felix, G. (2009). A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. Annu Rev of Plant Biol 60, 379-406.
  Google Scholar: Author Only Title Only Author and Title

Cao, H., Bowling, S.A., Gordon, A.S., and Dong, X. (1994). Characterization of an Arabidopsis mutant that is nonresponsive to inducers of systemic acquired resistance. Plant Cell 6, 1583-1592.
  Google Scholar: Author Only Title Only Author and Title

Cao, H., Glazebrook, J., Clarke, J.D., Volko, S., and Dong, X. (1997). The Arabidopsis NPR1 gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. Cell 88, 57-63.
  Google Scholar: Author Only Title Only Author and Title

Chen, H., Zou, Y., Shang, Y., Lin, H., Wang, Y., Cai, R., Tang, X., and Zhou, J.M. (2008). Firefly luciferase complementation imaging assay for protein-protein interactions in plants. Plant Physiol 146, 368-376.
  Google Scholar: Author Only Title Only Author and Title

Chinnusamy, V., Zhu, J., and Zhu, J.K. (2007). Cold stress regulation of gene expression in plants. Trends Plant Sci 12, 444-451.
  Google Scholar: Author Only Title Only Author and Title

Chinnusamy, V., Ohta, M., Kanrar, S., Lee, B.H., Hong, X., Agarwal, M., and Zhu, J.K. (2003). ICE1: a regulator of cold-induced transcriptome and freezing tolerance in Arabidopsis. Genes Dev 17, 1043-1054.
  Google Scholar: Author Only Title Only Author and Title

Choi, J., Huh, S.U., Kojima, M., Sakakibara, H., Paek, K.H., and Hwang, I. (2010). The cytokinin-activated transcription factor ARR2 promotes plant immunity via TGA3/NPR1-dependent salicylic acid signaling in Arabidopsis. Dev Cell 19, 284-295.
  Google Scholar: Author Only Title Only Author and Title

Deslandes, L., and Rivas, S. (2012). Catch me if you can: bacterial effectors and plant targets. Trends Plant Sci 17, 644-655.
  Google Scholar: Author Only Title Only Author and Title

Després, C., DeLong, C., Glaze, S., Liu, E., and Fobert, P.R. (2000). The Arabidopsis NPR1/NIM1 protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. Plant Cell 12, 279-290.
  Google Scholar: Author Only Title Only Author and Title

Ding, Y., Li, H., Zhang, X., Xie, Q., Gong, Z., and Yang, S. (2015). OST1 kinase modulates freezing tolerance by enhancing ICE1 stability in Arabidopsis. Dev Cell 32, 278-289.
  Google Scholar: Author Only Title Only Author and Title

Ding, Y., Sun, T., Ao, K., Peng, Y., Zhang, Y., Li, X., and Zhang, Y. (2018). Opposite roles of salicylic acid receptors NPR1 and NPR3/NPR4 in transcriptional regulation of plant immunity. Cell 173, 1454-1467.e1415.
  Google Scholar: Author Only Title Only Author and Title

Dodds, P.N., and Rathjen, J.P. (2010). Plant immunity: towards an integrated view of plant-pathogen interactions. Nature Rev Genet 11, 539-548.
  Google Scholar: Author Only Title Only Author and Title

Dong, C.H., Agarwal, M., Zhang, Y., Xie, Q., and Zhu, J.K. (2006). The negative regulator of plant cold responses, HOS1, is a RING E3 ligase that mediates the ubiquitination and degradation of ICE1. Proc Natl Acad Sci USA 103, 8281-8286.
  Google Scholar: Author Only Title Only Author and Title

Durrant, W.E., and Dong, X. (2004). Systemic acquired resistance. Annu Rev Phytopathol 42, 185-209.
  Google Scholar: Author Only Title Only Author and Title

Fujikawa, Y., and Kato, N. (2007). Split luciferase complementation assay to study protein-protein interactions in Arabidopsis protoplasts. Plant J 52, 185-195.

Google Scholar: Author Only Title Only Author and Title

Gilmour, S.J., Zarka, D.G., Stockinger, E.J., Salazar, M.P., Houghton, J.M., and Thomashow, M.F. (1998). Low temperature regulation of the Arabidopsis CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. Plant J 16, 433-442.
Google Scholar: Author Only Title Only Author and Title

Griffith, M., and Yaish, M.W. (2004). Antifreeze proteins in overwintering plants: a tale of two activities. Trends Plant Sci 9, 399-405.
Google Scholar: Author Only Title Only Author and Title

Han, Q., Tan, W., Zhao, Y., Yang, F., Yao, X., Lin, H., and Zhang, D. (2022). Salicylic acid-activated BIN2 phosphorylation of TGA3 promotes Arabidopsis PR gene expression and disease resistance. EMBO J, e110682.
Google Scholar: Author Only Title Only Author and Title

He, K., Du, J., Han, X., Li, H., Kui, M., Zhang, J., Huang, Z., Fu, Q., Jiang, Y., and Hu, Y. (2023). PHOSPHATE STARVATION RESPONSE1 (PHR1) interacts with JASMONATE ZIM-DOMAIN (JAZ) and MYC2 to modulate phosphate deficiency-induced jasmonate signaling in Arabidopsis. Plant Cell 35, 2132-2156.
Google Scholar: Author Only Title Only Author and Title

Hellens, R.P., Allan, A.C., Friel, E.N., Bolitho, K., Grafton, K., Templeton, M.D., Karunairetnam, S., Gleave, A.P., and Laing, W.A. (2005). Transient expression vectors for functional genomics, quantification of promoter activity and RNA silencing in plants. Plant Methods 1, 13.
Google Scholar: Author Only Title Only Author and Title

Hu, Y., Han, X., Yang, M., Zhang, M., Pan, J., and Yu, D. (2019). The transcription factor INDUCER OF CBF EXPRESSION1 interacts with ABSCISIC ACID INSENSITIVE5 and DELLA proteins to fine-tune abscisic acid signaling during seed germination in Arabidopsis. Plant Cell 31, 1520-1538.
Google Scholar: Author Only Title Only Author and Title

Huang, P., Dong, Z., Guo, P., Zhang, X., Qiu, Y., Li, B., Wang, Y., and Guo, H. (2020). Salicylic acid suppresses apical hook formation via NPR1-mediated repression of EIN3 and EIL1 in Arabidopsis. Plant Cell 32, 612-629.
Google Scholar: Author Only Title Only Author and Title

Huang, X., Li, J., Bao, F., Zhang, X., and Yang, S. (2010). A gain-of-function mutation in the Arabidopsis disease resistance gene RPP4 confers sensitivity to low temperature. Plant Physiol 154, 796-809.
Google Scholar: Author Only Title Only Author and Title

Huot, B., Castroverde, C.D.M., Velasquez, A.C., Hubbard, E., Pulman, J.A., Yao, J., Childs, K.L., Tsuda, K., Montgomery, B.L., and He, S.Y. (2017). Dual impact of elevated temperature on plant defence and bacterial virulence in Arabidopsis. Nat Commun 8, 1808.
Google Scholar: Author Only Title Only Author and Title

Jiang, Y.J., Liang, G., Yang, S.Z., and Yu, D.Q. (2014). Arabidopsis WRKY57 functions as a node of convergence for jasmonic acid- and auxin-mediated signaling in jasmonic acid-induced leaf senescence. Plant Cell 26, 230-245.
Google Scholar: Author Only Title Only Author and Title

Johnson, C., Boden, E., and Arias, J. (2003). Salicylic acid and NPR1 induce the recruitment of trans-activating TGA factors to a defense gene promoter in Arabidopsis. Plant Cell 15, 1846-1858.
Google Scholar: Author Only Title Only Author and Title

Jones, J.D., and Dangl, J.L. (2006). The plant immune system. Nature 444, 323-329.
Google Scholar: Author Only Title Only Author and Title

Kanaoka, M.M., Pillitteri, L.J., Fujii, H., Yoshida, Y., Bogenschutz, N.L., Takabayashi, J., Zhu, J.K., and Torii, K.U. (2008). SCREAM/ICE1 and SCREAM2 specify three cell-state transitional steps leading to Arabidopsis stomatal differentiation. Plant Cell 20, 1775-1785.
Google Scholar: Author Only Title Only Author and Title

Kesarwani, M., Yoo, J., and Dong, X. (2007). Genetic interactions of TGA transcription factors in the regulation of pathogenesis-related genes and disease resistance in Arabidopsis. Plant Physiol 144, 336-346.
Google Scholar: Author Only Title Only Author and Title

Kim, J.H., Castroverde, C.D.M., Huang, S., Li, C., Hilleary, R., Seroka, A, Sohrabi, R., Medina-Yerena, D., Huot, B., Wang, J., Nomura, K., et al. (2022). Increasing the resilience of plant immunity to a warming climate. Nature 607, 339-344.
Google Scholar: Author Only Title Only Author and Title

Kim, K.C., Lai, Z, Fan, B., and Chen, Z. (2008). Arabidopsis WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. Plant Cell 20, 2357-2371.
Google Scholar: Author Only Title Only Author and Title

Kim, Y., Park, S., Gilmour, S.J., and Thomashow, M.F. (2013). Roles of CAMTA transcription factors and salicylic acid in configuring the low-temperature transcriptome and freezing tolerance of Arabidopsis. Plant J 75, 364-376.
Google Scholar: Author Only Title Only Author and Title

Kim, Y.S., Lee, M., Lee, J.H., Lee, H.J., and Park, C.M. (2015). The unified ICE-CBF pathway provides a transcriptional feedback control of freezing tolerance during cold acclimation in Arabidopsis. Plant Mol Biol 89, 187-201.
Google Scholar: Author Only Title Only Author and Title

Kim, Y.S., An, C., Park, S., Gilmour, S.J., Wang, L., Renna, L., Brandizzi, F., Grumet, R., and Thomashow, M.F. (2017). CAMTA-mediated regulation of salicylic acid immunity pathway genes in Arabidopsis exposed to low temperature and pathogen infection. Plant Cell 29, 2465-2477.
Google Scholar: Author Only Title Only Author and Title

Nomura, K., He, S.Y. (2005). Powerful screens for bacterial virulence proteins. Proc Natl Acad Sci USA 102, 3527-3528.
Google Scholar: Author Only Title Only Author and Title

Kumar, S., Zavaliev, R., Wu, Q., Zhou, Y., Cheng, J., Dillard, L., Powers, J., Withers, J., Zhao, J., Guan, Z, Borgnia, M.J., Bartesaghi, A., Dong, X., and Zhou, P. (2022). Structural basis of NPR1 in activating plant immunity. Nature 605, 561-566.
Google Scholar: Author Only Title Only Author and Title

Lee, B.H., Henderson, D.A, and Zhu, J.K. (2005). The Arabidopsis cold-responsive transcriptome and its regulation by ICE1. Plant Cell 17, 3155-3175.
Google Scholar: Author Only Title Only Author and Title

Lee, J.H., Jung, J.H., and Park, C.M. (2015). INDUCER OF CBF EXPRESSION 1 integrates cold signals into FLOWERING LOCUS C-mediated flowering pathways in Arabidopsis. Plant J 84, 29-40.
Google Scholar: Author Only Title Only Author and Title

Lee, J.H., Jung, J.H., and Park, C.M. (2017). Light inhibits COP1-mediated degradation of ICE transcription factors to induce stomatal development in Arabidopsis. Plant Cell 29, 2817-2830.
Google Scholar: Author Only Title Only Author and Title

Li, H., Ding, Y., Shi, Y., Zhang, X., Zhang, S., Gong, Z, and Yang, S. (2017). MPK3- and MPK6-mediated ICE1 phosphorylation negatively regulates ICE1 stability and freezing tolerance in Arabidopsis. Dev Cell 43, 630-642.e634.
Google Scholar: Author Only Title Only Author and Title

Li, Z, Li, S., Jin, D., Yang, Y., Pu, Z., Han, X., Hu, Y., and Jiang, Y. (2023). U-box E3 ubiquitin ligase PUB8 attenuates abscisic acid responses during early seedling growth. Plant Physiol 191, 2519-2533.
Google Scholar: Author Only Title Only Author and Title

Li, Z, Liu, H., Ding, Z, Yan, J., Yu, H., Pan, R., Hu, J., Guan, Y., and Hua, J. (2020). Low temperature enhances plant immunity via salicylic acid pathway genes that are repressed by ethylene. Plant Physiol 182, 626-639.
Google Scholar: Author Only Title Only Author and Title

Liang, G., Zhang, H., Lou, D., and Yu, D. (2016). Selection of highly efficient sgRNAs for CRISPR/Cas9-based plant genome editing. Sci Rep 6, 21451.
Google Scholar: Author Only Title Only Author and Title

Miura, K., Jin, J.B., Lee, J., Yoo, C.Y., Stirm, V., Miura, T., Ashworth, E.N., Bressan, R.A, Yun, D.J., and Hasegawa, P.M. (2007). SIZ1-mediated sumoylation of ICE1 controls CBF3/DREB1A expression and freezing tolerance in Arabidopsis. Plant Cell 19, 1403-1414.
Google Scholar: Author Only Title Only Author and Title

Mou, Z., Fan, W., and Dong, X. (2003). Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. Cell 113, 935-944.
Google Scholar: Author Only Title Only Author and Title

Mukhopadhyay, A, Deplancke, B., Walhout, A.J., and Tissenbaum, H.A. (2008). Chromatin immunoprecipitation (ChIP) coupled to detection by quantitative real-time PCR to study transcription factor binding to DNA in Caenorhabditis elegans. Nat Protoc 3, 698-709.
Google Scholar: Author Only Title Only Author and Title

Nomoto, M., Skelly, M.J., Itaya, T., Mori, T., Suzuki, T., Matsushita, T., Tokizawa, M., Kuwata, K., Mori, H., Yamamoto, Y.Y., et al. (2021). Suppression of MYC transcription activators by the immune cofactor NPR1 fine-tunes plant immune responses. Cell Rep 37, 110125.
Google Scholar: Author Only Title Only Author and Title

Olate, E., Jiménez-Gómez, J.M., Holuigue, L., and Salinas, J. (2018). NPR1 mediates a novel regulatory pathway in cold acclimation by interacting with HSFA1 factors. Nat Plants 4, 811-823.
Google Scholar: Author Only Title Only Author and Title

Orvar, B.L., Sangwan, V., Omann, F., and Dhindsa, R.S. (2000). Early steps in cold sensing by plant cells: the role of actin cytoskeleton and membrane fluidity. Plant J 23, 785-794.
  Google Scholar: Author Only Title Only Author and Title

Sarkar, S., Das, A, Khandagale, P., Maiti, I.B., Chattopadhyay, S., and Dey, N. (2018). Interaction of Arabidopsis TGA3 and WRKY53 transcription factors on Cestrum yellow leaf curling virus (CmYLCV) promoter mediates salicylic acid-dependent gene expression in planta. Planta 247, 181-199.
  Google Scholar: Author Only Title Only Author and Title

Seo, P.J., Kim, M.J., Park, J.Y., Kim, S.Y., Jeon, J., Lee, Y.H., Kim, J., and Park, C.M. (2010). Cold activation of a plasma membrane-tethered NAC transcription factor induces a pathogen resistance response in Arabidopsis. Plant J 61, 661-671.
  Google Scholar: Author Only Title Only Author and Title

Sheen, J. (2001). Signal transduction in maize and Arabidopsis mesophyll protoplasts. Plant Physiol 127, 1466-1475.
  Google Scholar: Author Only Title Only Author and Title

Spoel, S.H., and Dong, X. (2012). How do plants achieve immunity? Defence without specialized immune cells. Nat Rev Immunol 12, 89-100.
  Google Scholar: Author Only Title Only Author and Title

Stockinger, E.J., Gilmour, S.J., and Thomashow, M.F. (1997). Arabidopsis thaliana CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. Proc Natl Acad Sci USA 94, 1035-1040.
  Google Scholar: Author Only Title Only Author and Title

Tada, Y., Spoel, S.H., Pajerowska-Mukhtar, K., Mou, Z, Song, J., Wang, C., Zuo, J., and Dong, X. (2008). Plant immunity requires conformational changes of NPR1 via S-nitrosylation and thioredoxins. Science 321, 952-956.
  Google Scholar: Author Only Title Only Author and Title

Tang, K., Zhao, L., Ren, Y., Yang, S., Zhu, J.K., and Zhao, C. (2020). The transcription factor ICE1 functions in cold stress response by binding to the promoters of CBF and COR genes. J Integr Plant Biol 62, 258-263.
  Google Scholar: Author Only Title Only Author and Title

Vlot, A.C., Dempsey, D.A., and Klessig, D.F. (2009). Salicylic Acid, a multifaceted hormone to combat disease. Annu Rev Phytopathol 47, 177-206.
  Google Scholar: Author Only Title Only Author and Title

Wang, C., Dai, S., Zhang, Z.L., Lao, W., Wang, R., Meng, X., and Zhou, X. (2021). Ethylene and salicylic acid synergistically accelerate leaf senescence in Arabidopsis. J Integr Plant Biol 63, 828-833.
  Google Scholar: Author Only Title Only Author and Title

Wang, W., Withers, J., Li, H., Zwack, P.J., Rusnac, D.V., Shi, H., Liu, L., Yan, S., Hinds, T.R., Guttman, M., Dong, X., and Zheng, N. (2020). Structural basis of salicylic acid perception by Arabidopsis NPR proteins. Nature 586, 311-316.
  Google Scholar: Author Only Title Only Author and Title

Wang, Y., Bao, Z, Zhu, Y., and Hua, J. (2009). Analysis of temperature modulation of plant defense against biotrophic microbes. Mol Plant Microbe Interact 22, 498-506.
  Google Scholar: Author Only Title Only Author and Title

Wigge, P.A. (2013). Ambient temperature signalling in plants. Curr Opin Plant Biol 16, 661-666.
  Google Scholar: Author Only Title Only Author and Title

Wu, Z, Han, S., Zhou, H., Tuang, Z.K., Wang, Y., Jin, Y., Shi, H., and Yang, W. (2019). Cold stress activates disease resistance in Arabidopsis thaliana through a salicylic acid dependent pathway. Plant Cell Environ 42, 2645-2663.
  Google Scholar: Author Only Title Only Author and Title

Xin, X.F., Nomura, K., Aung, K., Velasquez, A.C., Yao, J., Boutrot, F., Chang, J.H., Zipfel, C., and He, S.Y. (2016). Bacteria establish an aqueous living space in plants crucial for virulence. Nature 539, 524-529.
  Google Scholar: Author Only Title Only Author and Title

Yang, H., Shi, Y., Liu, J., Guo, L., Zhang, X., and Yang, S. (2010). A mutant CHS3 protein with TIR-NB-LRR-LIM domains modulates growth, cell death and freezing tolerance in a temperature-dependent manner in Arabidopsis. Plant J 63, 283-296.
  Google Scholar: Author Only Title Only Author and Title

Yang, S., and Hua, J. (2004). A haplotype-specific resistance gene regulated by BONZAI1 mediates temperature-dependent growth control in Arabidopsis. Plant Cell 16, 1060-1071.
  Google Scholar: Author Only Title Only Author and Title

Ye, K., Li, H., Ding, Y., Shi, Y., Song, C., Gong, Z, and Yang, S. (2019). BRASSINOSTEROID-INSENSITIVE2 negatively regulates the stability of transcription factor ICE1 in response to cold stress in Arabidopsis. Plant Cell 31, 2682-2696.

Google Scholar: Author Only Title Only Author and Title

Yu, X., Xu, Y., and Yan, S. (2021). Salicylic acid and ethylene coordinately promote leaf senescence. J Integr Plant Biol 63, 823-827.
Google Scholar: Author Only Title Only Author and Title

Yu, X., Cui, X., Wu, C., Shi, S., and Yan, S. (2022). Salicylic acid inhibits gibberellin signaling through receptor interactions. Mol Plant 15, 1759-1771.
Google Scholar: Author Only Title Only Author and Title

Zavaliev, R., Mohan, R., Chen, T., and Dong, X. (2020). Formation of NPR1 condensates promotes cell survival during the plant immune response. Cell 182, 1093-1108.e1018.
Google Scholar: Author Only Title Only Author and Title

Zhang, Y., Fan, W., Kinkema, M., Li, X., and Dong, X. (1999). Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the PR-1 gene. Proc Natl Acad Sci USA 96, 6523.
Google Scholar: Author Only Title Only Author and Title

Zhao, C., Wang, P., Si, T., Hsu, C.C., Wang, L., Zayed, O., Yu, Z., Zhu, Y., Dong, J., Tao, W.A., et al. (2017). MAP kinase cascades regulate the cold response by modulating ICE1 protein stability. Dev Cell 43, 618-629.e615.
Google Scholar: Author Only Title Only Author and Title

Zhou, J.M., Trifa, Y., Silva, H., Pontier, D., Lam, E., Shah, J., and Klessig, D.F. (2000). NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the PR-1 gene required for induction by salicylic acid. Mol Plant Microbe Interact 13, 191-202.
Google Scholar: Author Only Title Only Author and Title