Seasonal migration alters energetic trade-off optimization and shapes life history

Allison K. Pierce[1][*], Scott W. Yanco[2,3], Michael B. Wunder[1]

[1] University of Colorado Denver, Department of Integrative Biology, Denver, CO 80217, USA.
[2] Center for Biodiversity and Global Change, Yale University, New Haven, CT 06520, USA.
[3] Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06520, USA

[*]Corresponding author email: allison.pierce@ucdenver.edu phone: 720-300-1063

Short title: Migration alters trade-offs and shapes POL

Keywords: migration, pace of life, life history, trade-offs, energy budget, movement, evolution, seasonality, DEB

Authorship statement: AKP and SWY contributed equally to this work. AKP and SWY conceived the project, designed the model, performed analyses, and drafted the manuscript with support from MBW. Simulation and optimization model code written by AKP with input from SWY and MBW. Empirical data analysis and code lead by SWY with input from AKP and MBW. All authors contributed critical input to the manuscript.

Article type: Letter
Word count: Abstract 144; Main text 4996
Reference count: 75
Figure count: 6 (5 figures, 1 table)

Abstract

Trade-offs between current and future reproduction manifest as a set of co-varying life

history and metabolic traits, collectively referred to as "pace of life" (POL). Seasonal migration

modulates realized environmental dynamics and putatively affects realized POL, however, the

mechanisms by which migratory behavior shapes POL remain unclear. We explored how

migratory behavior interacts with environmental and metabolic dynamics to shape POL. Using

an individual based model of movement and metabolism we compared fitness-optimized trade-

offs among migration strategies. We found annual experienced seasonality and migration

distance primarily drove POL differentiation through developmental and migration phenology

trade-offs. Similarly, our analysis of empirically-estimated metabolic data from 265 bird species

suggested seasonal niche conservatism and migration distance interact to drive POL. We show

multiple viable life history strategies are conducive to a migratory lifestyle. Overall, our findings

suggest metabolism mediates complex interactions between behavior, environment, and life

history.

Introduction

Time and energy are limited resources used by organisms to survive and reproduce

(Levins 1962; Cody 1966). Organisms balance use of these resources by varying strategic

emphasis on current versus future reproduction (Williams 1966; van Noordwijk & de Jong 1986;

Stearns 1989, 1992). This fundamental trade-off manifests as a set of co-varying life history and

metabolic trade-offs, collectively referred to as 'pace of life' (POL; Williams 1966; Stearns

1989, 1992; Wikelski & Ricklefs 2001; Ricklefs & Wikelski 2002) driven by traits which can

include age at first reproduction (Harvey & Zammuto 1985; Sæther 1988), fecundity (Lack 1947;

Sæther 1988), growth rate (Gebhardt-Henrich & Richner 1998), investment in survival (Harvey

& Zammuto 1985), and metabolic rate (Wikelski *et al.* 2003a; Wiersma *et al.* 2007b).

At biogeographic scales, latitude covaries with both life history trade-offs (Moreau 1944;

Lack 1947; Martin 1996; Ricklefs 2000; Jetz *et al.* 2008; Yanco *et al.* 2022) and organismal

metabolism (Wikelski et al. 2003a; Wiersma et al. 2007a, b; Yanco et al. 2022) implying that

environment contributes to realized POL. However, despite the extensive literature on the topic,

the mechanistic drivers of covariation between environment and POL remain poorly understood,

largely because latitude is actually a proxy for several confounded variables (e.g., mean resource

availability, periodic amplitude of resource variance, ambient temperature, predation pressure,

intraspecific competition, etc.) all of which could plausibly drive variance in POL (Ghalambor &

Martin 2001). Yanco et al. (2022) showed that high amplitude seasonal variation in resource

availability mechanistically constrains organisms to relatively "fast" POL strategies (emphasis

on fecundity over survival, high metabolic rate). Conversely, low amplitude seasonal resource

fluctuation was associated with expanded viability of "slow" POL strategies (emphasis on

survival over fecundity, low metabolic rates). These findings imply that environmental

conditions, and the dynamics of those conditions over time, influence realized POL strategies.

POL also exhibits complicated covariance patterns with animal behavior. Behavior can

modify extrinsic mortality risk, which is expected to drive POL; High extrinsic mortality risk

favors a fast POL whereas low extrinsic mortality risk favors a slow POL (Ricklefs 2000; Réale

et al. 2010; Dammhahn *et al.* 2018). Behaviors such as movement may also mediate the

environmental conditions an organism experiences, making the optimal life history strategy the

outcome of an interaction between environment and behavior rather than environment alone

(Spiegel *et al.* 2015; Campos-Candela *et al.* 2019; Laskowski *et al.* 2021). For example,

populations of Corsican Blue Tits (Cyanistes caeruleus) in evergreen-dominated habitats

exhibited slow POL along with reduced handling aggression, exploration speed, and nest defense

whereas individuals in deciduous habitats showed the converse (Dubuc-Messier *et al.* 2017).

Seasonal migration is a behavior with putatively substantial effects on realized POL but

empirical findings about the direction and magnitude of those effects vary depending on the

taxonomic scale of comparison, environmental context, and characterization of the behavior

(Soriano-Redondo *et al.* 2020; Winger & Pegan 2021). Migration radically alters the

environmental conditions an organism experiences, typically leading to niche conservatism,

relatively invariant conditions across the annual cycle (Nakazawa *et al.* 2004; Gómez *et al.* 2016;

Merkle *et al.* 2016; Thorup *et al.* 2017; Zurell *et al.* 2018; Alerstam *et al.* 2019; Abrahms *et al.*

2021; Yanco *et al.* 2021), which might be expected to favor a slow POL. On the other hand,

migration is a potentially costly endeavor both in terms of increased mortality risk (Sillett &

Holmes 2002; Rushing *et al.* 2016, 2017; but see Zúñiga *et al.* 2017) and metabolic expenditures

(Wikelski *et al.* 2003b; Alves *et al.* 2013; Brown *et al.* 2023), suggesting that migration might be

associated with a fast POL. Intriguingly, some studies suggest that migrant birds exhibit

intermediate POL: faster than temperate counterparts, but slower than residents breeding at

similar latitudes. For example, avian clutch size is a classic case study of biogeographic variance

in life history across a gradient of seasonality (Moreau 1944; Lack 1947). Jetz et al. (2008)

showed that clutch sizes of migrants were smaller than would be expected from phylogenetically

equivalent residents breeding at similar latitudes. Similarly, Weirsma et al. (2007b) showed that

although tissue basal metabolic rate (BMR; assumed to be proportional to POL) was positively

related with latitude, migrants had lower BMR than their resident temperate counterparts and

higher BMR than tropical residents. Ultimately, how migration is mechanistically linked to

realized POL remains ambiguous because, to date, few studies have simultaneously accounted

for the complex feedbacks between the bioenergetics of migration and the emergent

environmental dynamics.

In this study we sought to explore how migratory behavioral mitigation of seasonality can

affect dynamics of metabolic and temporal trade-offs to shape POL and life history based on a

model of avian migration. For example, if migration solely balances the energetic cost of

movement against the benefit from mitigating seasonal resource variance, we would expect

similar life histories across migratory strategies. However, if migration substantially affects the

net energy available to balance metabolic tradeoffs, then we would expect covariation between

migratory strategy and life history. Furthermore, energetic trade-offs also interact with

phenological constraints imposed by migration. Time-consuming migrations might restrict

breeding season duration and ultimately favor fewer and/or faster developing young despite

increases in net energy from migration. Thus, certain migrations may be more time- than energy-

limited than others.

To model energetic and temporal trade-offs we combined an individual-based model

(IBM) of one-dimensional movement within a simulated environment of seasonally fluctuating

energy with a generalized Dynamic Energy Budget (Kooijman 2010) model of metabolism

(sensu Yanco *et al.* 2022). We compared life-history optimization trade-offs between migratory

and stationary strategies using a heuristic optimization algorithm to solve for fitness-optimized

IBM parameterizations across a gradient of seasonality and migration strategies. Finally, we

show empirically that seasonal niche conservatism and migration distance interact to drive

realized POL across 265 species of birds. We suggest that behavioral phenotype may play an

underappreciated role in driving life history strategies, especially in cases where the interaction

of behavior and the environment produce novel adaptive landscapes.

Methods

DEB model description

Dynamic Energy Budget (DEB) theory (Kooijman 2010) proposes a dynamic

mechanistic model of how individuals uptake, use, and dissipate energy over their lifespans and

across different life stages based on the principles of energy and mass conservation,

thermodynamics, and stoichiometric constraints. To simulate energy acquisition and allocation

dynamics of metabolic trade-offs we implemented a unitless specification of a standard DEB

model modified to incorporate metabolic costs of migratory movements (Kooijman 2010). We

scaled DEB parameters for energy and organism structural size by maximum reserve energy

density and maximum length respectively. Resultant system dynamics are therefore more

generalizable as they quantify proportional changes rather than absolute changes in units of

energy or body size.

In the standard model (Fig. 1), organisms acquire energy from the environment and

assimilate it into reserves ($\dot{p}_A$). Energy is then mobilized from the reserve at a fixed proportion κ

to cover energetic requirements of growth ($\dot{p}_G$) and somatic maintenance ($\dot{p}_S$), with the remainder

diverted to maturation maintenance ($\dot{p}_J$) and reproduction ($\dot{p}_R$). In the juvenile stage, energy

allocated to reproduction is applied to advancing maturity level rather than offspring production.

Locomotory costs are not explicitly defined but are assumed to come from somatic

maintenance costs when they are independent from foraging rate (Kooijman 2010). Therefore,

we modified the standard model to isolate migration movement costs from somatic maintenance

costs via the compound parameter m which we defined as the ratio between metabolic cost of

locomotion to travel one body length (l) during migration relative to somatic maintenance costs

unrelated to migration. Thus, total somatic maintenance rate ($\dot{p}_S$) is the sum of the locomotory

cost of migration ($\dot{p}_L$) which scales with structural surface area (l^2) and distance, and somatic

maintenance costs unrelated to migration ($\dot{p}_M$) which are proportional to biovolume (l^3).

In starvation conditions, energy use is prioritized by somatic maintenance first, growth

second, followed by maturation/reproduction. In our implementation, length or maturation level

cannot shrink to reduce energy costs and death occurs when energy drawn for somatic

maintenance exceeds available reserves. Detailed model parameterization and derivation is

available in supplement Appendix S1 and Kooijman (2010).

IBM simulation

We used an IBM simulation to evaluate daily energy acquisition and allocation for an

individual over 365 days from birth, through sexual maturation, to reproduction. At each time-

step a set of differential equations was numerically solved to update values of the DEB state

variables given individual trait parameters (Table 1; see Appendix S1 for detail). Environmental

energy available at each time-step depended on the individual's latitudinal location, where

available energy f varied across 365 discrete time-steps as one oscillation of a sine function, for

which the amplitude and phase varied spatially across the 180 continuous latitude units.

Maximum amplitude occurred at the lowest and highest latitudes and decreased towards the mid-

point. Phase is shifted by a half wavelength on either side of the mid-point to reflect the

alternating seasonality between hemispheres (sensu Yanco et al. 2022).

Individuals are initiated at the peak of seasonal energy availability for their initial

location and reproduction occurs such that offspring are born when energy availability peaks

again on day 365 ($t = 365 - a_b$). If individuals are migratory, at the end of each time-step they

move towards non-breeding locations once sexually mature or depart to breeding locations after

remaining at their non-breeding location for their specified duration (d_{NB}). For simplicity, we

assumed constant migration speed. Distance traveled in each time-step was calculated as

migration distance divided by specified duration of migration (d_M).

We calculated fecundity for viable solutions as the amount of cumulative energy in the

reproductive buffer divided by the cost of one egg, u_E^0 . Egg cost is based on the individual's

scaled reserve density at egg production with the assumption that length at birth and growth rate

of offspring are inherited traits.

At the final time-step, solutions (individual sets of parameters) were classified as

nonviable if energy reserves were depleted ($e < 0$), reserves at initialization were insufficient for

offspring to reach birth size ($e_b < l_b$), less than one offspring was produced, or migratory

individuals did not return to breeding locations in time to complete reproduction.

IBM parameter optimization

Heuristic optimization algorithms are useful for efficiently solving multi-dimensional

optimization problems that are analytically intractable and too computationally intensive to solve

numerically by brute force (Hamblin 2013), such as the problem we studied here. We maximized

for fecundity using a genetic algorithm, a heuristic optimization of an objective function

(represented here as an IBM) using iterative search techniques inspired by genetic mechanisms

of natural selection such as inheritance, mutation, and crossover. Despite the biological

inspiration for the algorithm, our implementation is not intended to be interpreted as simulating

evolutionary processes.

Optimization by movement strategy

To investigate life history optimization across a gradient of seasonality and migration

distance we optimized parameterizations for 9 different movement strategies: 6 migratory and 3

stationary. Stationary strategies occurred at latitudes we classified as high seasonality (80, -80),

mid seasonality (40, -40) and aseasonal (0). Migratory strategies were defined by breeding and

non-breeding end point locations and varied in distance such that migrants bred in high or mid-

seasonal latitudes and migrated to high or mid-seasonal locations in the opposite "hemisphere", a

mid-seasonal location in the same hemisphere, or the aseasonal location.

Migration strategies were further classified by migration distance, seasonality of

endpoints, and energy management strategy. Migrations between seasonally out of phase

endpoints (cross hemisphere) were defined as energy-buffering strategies because they buffer

seasonal oscillations of energy over time by tracking seasonal peaks in energy availability. We

classified migrations between seasonally in-phase locations (within hemisphere) as energy

mitigating strategies because they dampen amplitude of seasonal oscillations in energy over

time. Stationary strategies do not behaviorally manipulate seasonality and so we classified them

as coping strategies.

For each strategy, we initialized IBM parameters with randomly generated values for

5000 solutions and iteratively optimized values for 200 iterations (see Appendix S1 for detail).

We repeated this process 1000 times (runs) to ensure adequate exploration of the solution space.

Parameters were not simultaneously optimized across migratory strategies, so different strategies

might have explored the solution space to different extents (Hamblin 2013). To address this, we

used the same random number generation seeds across migration strategies for each run so that

each optimization process started with the same set of strategy invariant parameters for each

strategy but with different random sets for each run. We also performed an additional set of 90

runs per migratory strategy pre-initialized with parameters from five of the highest fecundity

viable solutions for each strategy from the initial 1000 runs (10 runs of 9 pre-initializations) to

equalize exploration of the solution space between strategies. Finally, we pooled together

solutions from all 1090 runs, removed duplicate solutions within strategies, and retained

solutions in the top 1% for fecundity from each for analysis.

We conducted all simulations, optimizations, and analyses in R (R Core Team 2022).

Pace of Life Trade-offs

We used a correlation-based principal components analysis (PCA) of optimized

individual parameter values from each movement strategy to analyze trait covariation patterns

and evaluate metabolic and temporal trade-off dynamics. We analyzed energy mobilization and

somatic allocation parameters $\dot{v}$ and κ, in addition to parameters governing downstream somatic

and reproductive allocation ($\dot{k}_M$, m, and g). Additionally, we analyzed parameters related to

timing of migration (a_p, d_{NB}, and d_M). We used bi-plots of parameter loadings to qualitatively

evaluate covariance relationships (especially including trade-offs) between parameter

combinations and between migration strategies.

In silico experiment

198   To investigate the effects of migration on relative life-history optimization across

199   migratory strategies, we evaluated the fecundity values of optimized parameterizations for each

200   migration strategy in simulation scenarios wherein individuals were forced to stay in breeding

201   locations (could not migrate) as compared to optimized parameterizations for stationary

202   strategies from the same respective breeding location.

Empirical Analysis

We leveraged publicly available datasets to explore the extent of empirical support for

the influence of migratory phenotype, biogeography, and realized niche dynamics on life history

in birds. We used the Add-my-Pet dataset which includes empirically estimated DEB parameters

for 4035 species (Marques *et al.* 2018) to calculate a general proxy for POL as a size-normalized

estimate of relative maximum somatic mobilization rate $\kappa \dot{p}_C$ where higher values represent

increased investment of reserves into somatic processes and thus slower pace of life. To account

for non-linear scaling of energy assimilation and mobilization as a function of body size, we

calculated this as $POL = \frac{\kappa \dot{v}}{z}$, where z is the ratio of species-specific maximum length to the

maximum length of 1 cm for a theoretical reference species, allowing for comparisons across

species. We used this POL metric as the response variable in the models described below.

To assess the effects of biogeography, migratory phenotype, and environmental dynamics

on these life history traits we considered three predictors: breeding latitude, migration distance,

and environmental niche dissimilarity (roughly analogous to energy management strategy). We

used the Birdlife International dataset (BirdLife International and Handbook of the Birds of the

World 2016) to calculate the latitude of the centroid of breeding season ranges and calculated

species-level migration distance as the great-circle distance between seasonal range centroids. To

characterize seasonal environmental dynamics for each species we used results from Cohen &

Jetz (2023) which estimates niche dissimilarity as the Mahalanobis distance in environmental

space between niche hypervolumes of breeding locations and overwintering locations.

Combining these sources resulted in a final dataset of 265 species. All predictor variables were

centered and scaled before model fitting.

We modeled macroecological patterns in our POL metric as a function of the three

predictor variables under a Bayesian framework using the `brms` package in R (Bürkner 2017).

We modeled POL as beta distributed (expected domain [0,1]) with niche dissimilarity as a

predictor on the beta distribution precision parameter (ϕ) based on previous reports (Yanco et al.

2022). To control for phylogenetic relatedness among the species in our sample we implemented

a phylogenetic generalized mixed model approach (Lynch 1991; Garamszegi 2014). Briefly, this

approach includes a random intercept by species and enforces a correlation structure among

species matching a known phylogeny. We downloaded 100 phylogenies for our target species

from vertlife.org (Jetz et al. 2012) and ran analyses using a single randomly selected tree. We re-

ran analyses with new randomly selected trees to assess sensitivity to the specific phylogeny

used and found results to be invariant across candidate trees. We fit the model using four MCMC

chains of 10,000 draws (with a burn in of 5,000) thinned by 5. We assessed model convergence

with a combination of $\hat{R} < 1.05$ and visual inspection of MCMC traceplots and posterior

predictive plots. We interpreted the relative significance of individual parameters using the

Bayesian probability of direction (PD), the proportion of the estimated posterior distribution on

the same side of zero as the estimate.

Results

Migration IBM

Movement paths of model optimized parameterizations for each migratory strategy

generally tracked maximal available energy over time but all strategies optimized for later

departure times from non-breeding locations than expected if perfectly tracking peak resources

(Fig 2A). Migratory strategies realized higher annual mean energy availability than stationary

strategies irrespective of breeding location with annual mean gain increasing with migratory

distance (Fig. 2B).

Pace of Life Trade-offs

Three axes of variation explained a total of 95.8% of the variation in optimized temporal

and energetic trade-off parameters across all migratory and stationary strategies with components

1, 2, and 3 each explaining 58.3%, 27.4%, and 10.1% respectively. Secondary and tertiary

somatic stream allocation parameters $\dot{k}_M$, g, and m were the primary contributors to the first

component in addition to energy mobilization ($\dot{v}$) and migration timing parameter d_{NB} (PC1, Fig.

S3). Somatic maintenance rate $\dot{k}_M$ exhibited opposing loadings to all other parameters ($\dot{k}_M$=

0.44, g =-0.38, m =-0.43, $\dot{v}$ = -0.38, and d_{NB}= -0.41) indicating trade-offs between somatic

maintenance in metabolic cost of growth and migration as well as migration timing. Higher

values of PC1 were associated with slower mobilization of reserves, prioritized investment into

somatic maintenance over growth, and less time spent in the non-breeding location. Upstream

energy allocation κ and parameters governing migratory timing a_p and d_M were the primary

contributors of the second component (PC2, Fig. S3), all of which exhibited positive loadings

(κ=0.58, a_p=0.56, and d_M =0.37) indicating higher values of PC2 were associated with higher

proportional allocation to somatic processes, later maturation (and thus migration), and longer

migration. Component 3 (PC3) was primarily associated with metabolic parameters κ, g, and $\dot{v}$ in

addition to a_p (Fig. S3). Loadings for g and a_p (g =-0.83, a_p =-0.38) opposed both κ and $\dot{v}$

(κ=0.41, $\dot{v}$=0.42) indicating higher values of PC3 represent increased somatic allocation and

mobilization of energy, lower investment into growth, and earlier maturation (and thus

migration).

Optimized parameterizations were differentiated by energy management strategy along

the PC1 and PC2 axes (Fig. 3A) indicating that trade-offs between metabolism and migratory

timing were associated with behavioral manipulation of experienced seasonality (Fig. 3B).

Unsurprisingly and by design, stationary strategies that cope with seasonality differentiated from

migratory strategies based on parameters for metabolic cost and timing of migration. More

interestingly, stationary coping strategies were associated with lower per-volume growth costs,

prioritized investment into somatic maintenance over growth, and slower mobilization. Buffering

strategies were generally associated with longer migrations, shorter non-breeding stationary

periods, and decreased migratory locomotion costs compared to mitigating strategies.

Migratory strategies were further differentiated by migration distance along the PC2 and

PC3 axes (Fig 3B) such that somatic allocation and migration duration generally increased with

distance and investment into growth and mobilization rate decreased. We did not observe

appreciable differentiation with seasonality of breeding or non-breeding locations (Fig. S5).

In silico experiment

Overall, energy-buffering long-distance migratory strategies realized the highest

fecundity followed by the stationary aseasonal strategy. All optimized parameterizations for

migratory movement strategies realized higher or similar relative fecundity than stationary

strategies for their respective breeding locations with fecundity gain increasing with migratory

distance. When model optimized parameterizations for migratory strategies were simulated to

remain stationary at breeding locations, all parameterizations realized lower relative fecundity

than optimized stationary parameterizations. Declines in relative fecundity generally increased

with migration distance (Fig. 4).

Empirical Analysis

Our macroecological analysis broadly agreed with the general patterns proposed in our

theoretical model - life history covaries with biogeography, behavior, and environmental

dynamics (Fig. 5). Both migration distance and environmental niche dissimilarity exhibited

positive effects on POL (migration distance: avg. marginal effect = 0.002, PD = 99.35%; niche

dissimilarity: avg. marginal effect = 0.037, pd = 84.42%) whereas breeding latitude had no

apparent influence on POL (avg. marginal effect < 0.001, PD = 58.83%). Variance

decomposition showed a strong influence of phylogeny on POL (variance ratio = 0.82).

Discussion

Migration integrates biogeography, behavior, and energetics

Our model suggests a complex relationship between behavior, environment, and life

history is fundamentally mediated by metabolism. We observed multiple distinct viable life

history strategies within any given migratory strategy (Fig 2A). Some of this diversity in strategy

clearly arises from the interaction of behavioral modes and biogeographic factors which are also

relevant to life history. For example, the three stationary strategies we considered showed

markedly different life histories despite shared behavior (Fig 3). Consistent with previous

reports, high energy availability and steep seasonal dynamics at high latitudes push organisms

towards fast POL strategies (Mueller & Diamond 2001; Wiersma et al. 2007b; Halali et al. 2020;

Yanco et al. 2022). Our results extend these findings to show that behavioral manipulation of the

realized seasonal dynamics (via movement) can further modify energetic strategies by interacting

with developmental constraints and affecting net energy availability.

Migratory phenology emerged as a fundamental axis of life history variation, primarily

trading off the timing and speed of migration with energy allocation strategies. Previous work in

boreal breeding birds found that migration distance (a proxy for breeding season duration)

negatively covaried with clutch size and annual fecundity such that species with the longest

breeding seasons were able to increase the number of young produced (Winger & Pegan 2021).

Similarly, we found greater investment in self in strategies that maximized time on migration and

non-breeding grounds. This result could relate to previous reports showing, counterintuitively,

longer migrations were associated with better body condition (Ortega 2023) and higher annual

survival (Zuniga et al 2017) in partially migratory systems. Finally, the apparent importance of

phenology suggests that incorporating variable migration speeds could be especially fruitful

extensions of the model.

We found that energy management strategy was related to but distinct from migration

distance alone. Together, these factors explained substantial variation in optimized life history

parameters (Fig 3A) and acted primarily via differential modulation of migration and

developmental timing, and energy allocation. Longer migrations tended to result in more

buffered energy strategies (by design) but did not perfectly covary, highlighting the utility of

using energetics as the common currency to parse behavior-life history relationships.

Our theoretical results suggest that migration mediates metabolic and temporal trade-offs

by altering net energy availability (Fig 2.) but differentially based on energy management

strategy and migration distance (Fig. 3). Energy-buffering strategies exhibited slower POL than

mitigating and coping strategies (Fig. 3A); Short-distance strategies exhibited faster POL than

other migratory strategies (Fig. 3B). Similarly, relative investment in self, at the expense of

reproduction, was driven by a combination of niche dissimilarity (roughly analogous to energy

management strategy) and migration distance in 265 species of migrant birds. Slower POL was

associated with greater migration distance but with greater rather than less niche dissimilarity.

However, the metric only measures differences in niche between endpoints and not across the

annual cycle, which might explain the converse result. Additionally, some of the environmental

variables used in the dissimilarity metric may be unrelated to energy availability. Despite this,

empirical results were congruent with the general theoretical prediction that variation in POL is

driven by a combination of migratory distance and differences in endpoint seasonality.

Implications for the evolution of avian migration

Our finding that multiple combinations of life history parameters are theoretically

conducive to a migratory lifestyle sheds new light on the apparent facility with which birds can

switch between migratory and sedentary lifestyles. Among birds, migration can emerge and

disappear over very short timescales (Berthold *et al.* 1992; Garcia-Perez *et al.* 2013). Indeed,

phylogenetic analyses suggest that such transitions between migratory states have occurred so

frequently over the evolutionary history of class Aves that the potential to migrate is widely

viewed as basal to the clade, requiring only minor phenotypic changes to switch states (Zink

2011; Zink & Gardner 2017). In addition to modifying behavioral, morphological, and

physiological traits related to migration, any population in the process of switching between

migratory states must also move toward a metabolic strategy that maximizes evolutionary

fitness. The expected distance in metabolic trait space between a resident life history strategy and

a migratory strategy would be lower if, as our results suggest, there are multiple potential

metabolic arrangements amenable to a migratory lifestyle. In other words, transitioning from

migrant to non-migrant (or vice versa) may not require a radical metabolic reorganization. Thus,

with respect to life history, it is perhaps unsurprising that migration has emerged multiple times

because there are so many putatively viable migratory life histories which are only partially

separable from entirely non-migratory life histories.

Relatedly, our *in silico* experiment showed the steepest declines in reproductive potential

when restricting high latitude migrants to a resident strategy. The net energy benefit of migration

increased with seasonality of the breeding location and migratory distance, implying that the

emergence of a long-distance migration from high latitude residents is associated with a large

increase in reproductive potential that far exceeds the best-performing stationary solution. Recent

work has suggested a high-latitude origin for most avian migrations as opposed to earlier theories

which posit a "southern home" (Bruderer & Salewski 2008; Winger *et al.* 2011, 2019; Yanco et

al. 2021). Our results provide theoretical justification for these observations in showing large

realized fitness benefits of migration as compared to otherwise optimized non-migratory

strategies.

Model assumptions and extensions

While our model accounts for several elements of dynamic environments and labile

behaviors, it is not a comprehensive account of all factors potentially influencing life history

optimization. The current instantiation of our model focuses on intrinsically controlled vital rates

in concert with environmental dynamism and behavioral plasticity. Future work could extend our

model to account for any number of additional factors, including especially other potential

extrinsic drivers or phenomena that transpire over shorter timescales. For example, in our model

individuals acquire energy at the same rate during migratory movements as while stationary.

Forcing dynamic activity budgeting at a sub-daily resolution (rather than assuming constant

foraging rates and efficiencies regardless of migration activity) could affect optimized values of

$\dot{v}$. We hypothesize that reduced foraging efficiency during migration might result in solutions

showing slower mobilization and/or larger reserve capacity for migratory strategies rather than

the faster mobilizations that we observed. Additionally, our simple movement model assumes

constant migration speed and symmetry in duration of migrations to and from breeding locations.

A more complex movement model could investigate effects of stepwise migration and finer scale

speed regulation on trade-off dynamics.

We also did not incorporate extrinsic sources of mortality which have the potential to

strongly influence fitness of otherwise viable and fecund life history strategies. Extrinsic

mortality during migration, while traditionally difficult to estimate (Rushing 2019), is potentially

substantial (Rushing *et al.* 2016; Ward *et al.* 2018; Buechley *et al.* 2021). Similarly, juvenile

survival and recruitment is a notoriously important phenomenon for understanding population

vital rates (Charnov & Schaffer 1973) for which our model does not account. Finally, differential

predation between winter and breeding areas is thought to be an important factor influencing the

fitness benefit of migration (Martin 2015). Model extensions that stochastically integrate

extrinsic mortality, especially environmentally mediated mortality, would shed light on how

animals may compensate for extrinsic mortality via energetic buffering (i.e., aiming for "excess"

intrinsic survival rates to counter unpredictable extrinsic mortality).

In addition to energetics in the form of food, ambient temperatures are thought to play an

important role in shaping migrations (Corkeron & Connor 1999; Wikelski *et al.* 2003b; Wiersma

et al. 2007a; Clerc & McGuire 2021). Our model assumes that seasonal temperature variation is

reflected in seasonal availability of energy and acquisition rates which scale with organismal

surface area. More complex thermoregulatory dynamics would allow exploration of how energy

acquisition from the environment relates to the costs of maintaining homeostasis in that

environment. For example, accounting for trade-offs between metabolic costs associated with

molt/maintenance of body insulation and movement performance could shed light on interactions

between molting phenology, migratory strategy, and life history.

Furthermore, even though we sought to maintain model generality, our IBM simulation

assumed oviparity, first year reproduction, and initiation of migration at maturation; all of which

are generally most consistent with avian life histories. Alternate IBM specifications of

reproductive timing, simulation length, movement constraints, or metabolism would allow for

more nuanced investigation into developmental trade-off dynamics in taxa that deviate from

these constraints. For example, the standard DEB model can be modified to account for fetal

development to examine gestational trade-off dynamics in mammals or to account for multiple

life stages to examine life-cycle dynamics in invertebrates.

Conclusion

Our work shows that rather than a singular response to environment, seasonal migrations

are actually a diverse set of adaptive strategies that integrate reciprocal relationships between

dynamic and spatially-varying environments, plastic behavioral adaptations, and apparently

labile metabolic organizations, all of which jointly produce fitness (Hutchinson 1948; Piersma &

van Gils 2019). Viewed as such, our mechanistic model provides novel theoretical insights into

the emergence of migration across taxa. Moreover, our results clearly demonstrate that behaviors

- which both cost energy and modify environmental context - are integral components of life

history strategy, providing mathematical support to the Extended POL Theory (Dammhahn *et al.*

2018). Finally, our work shows the promise of approaches that simultaneously consider these

reciprocally related aspects of organismal ecology (behavior, environment, and life history) and

the unifying power of individual energetics as common currency to relate the three.

Acknowledgements

427 We thank Bas Kooijman whose correspondence about our previous work helped to

428 inform aspects of this study. AKP was supported by the Fritz L. Knopf Fellowship in Avian

429 Conservation. SWY was partially supported by the Max Planck – Yale Center for Biodiversity

430 Movement and Global Change.

Literature Cited

Abrahms, B., Aikens, E.O., Armstrong, J.B., Deacy, W.W., Kauffman, M.J. & Merkle, J.A.
(2021). Emerging Perspectives on Resource Tracking and Animal Movement Ecology.
Trends Ecol. Evol., 36, 308–320.

Alerstam, T., Bäckman, J., Grönroos, J., Olofsson, P. & Strandberg, R. (2019). Hypotheses and
tracking results about the longest migration: The case of the arctic tern. *Ecol. Evol.*, 9,
9511–9531.

Alves, J.A., Gunnarsson, T.G., Hayhow, D.B., Appleton, G.F., Potts, P.M., Sutherland, W.J., *et
al.* (2013). Costs, benefits, and fitness consequences of different migratory strategies.
Ecology, 94, 11–17.

Augustine, S., Lika, K., & Kooijman, S. A. (2019). Altricial-precocial spectra in animal
kingdom. *Journal of Sea Research*, *143*, 27-34.

Berthold, P., Helbig, A.J., Mohr, G. & Querner, U. (1992). Rapid microevolution of migratory
behaviour in a wild bird species. *Nature*, 360, 360668a0.

BirdLife International and Handbook of the Birds of the World. (2016). Bird species distribution
maps of the world.
452
453 Brown, J.M., Bouten, W., Camphuysen, K.C.J., Nolet, B.A. & Shamoun-Baranes, J. (2023).
454   Energetic and behavioral consequences of migration: an empirical evaluation in the
455   context of the full annual cycle. *Sci. Rep.*, 13, 1210.
456
457 Bruderer, B. & Salewski, V. (2008). Evolution of bird migration in a biogeographical context. *J.*
458   *Biogeogr.*, 35, 1951–1959.
459
460 Buechley, E.R., Oppel, S., Efrat, R., Phipps, W.L., Carbonell Alanís, I., Álvarez, E., *et al.*
461   (2021). Differential survival throughout the full annual cycle of a migratory bird presents
462   a life-history trade-off. *J. Anim. Ecol.*
463
464 Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal*
465   *of Statistical Software*.
466
467 Campos-Candela, A., Palmer, M., Balle, S., Álvarez, A. & Alós, J. (2019). A mechanistic theory
468   of personality-dependent movement behaviour based on dynamic energy budgets. *Ecol.*
469   *Lett.*, 22, 213–232.
470
471 Charnov, E.L. & Schaffer, W.M. (1973). Life-History Consequences of Natural Selection: Cole's
472   Result Revisited. *Am. Nat.*, 107, 791–793.
473
474 Clerc, J. & McGuire, L.P. (2021). Considerations of varied thermoregulatory expressions in
475   migration theory. *Oikos*, 130, 1739–1749.
476
477 Cody, M.L. (1966). A GENERAL THEORY OF CLUTCH SIZE. *Evolution*, 20, 174–184.
478
479 Cohen, J. & Jetz, W. (2023). Diverse strategies for tracking seasonal environmental niches at
480   hemispheric scale. *Global Ecology and Biogeography* 32.9 (2023): 1549-1560.
481
482 Corkeron, P.J. & Connor, R.C. (1999). Why do baleen whales migrate?1. *Mar. Mamm. Sci.*, 15,
483   1228–1245.
484
485 Dammhahn, M., Dingemanse, N.J., Niemelä, P.T. & Réale, D. (2018). Pace-of-life syndromes: a
486   framework for the adaptive integration of behaviour, physiology and life history. *Behav.*
487   *Ecol. Sociobiol.*, 72, 62.
488
489 Dubuc-Messier, G., Réale, D., Perret, P. & Charmantier, A. (2017). Environmental heterogeneity
490   and population differences in blue tits personality traits. *Behav. Ecol.*, 28, 448–459.
491
492 Garamszegi, L.Z. (Ed.). (2014). *Modern Phylogenetic Comparative Methods and Their*
493   *Application in Evolutionary Biology: Concepts and Practice*. Springer, Berlin,
494   Heidelberg.
495

496 Garcia-Perez, B., Hobson, K.A., Powell, R.L., Still, C.J. & Huber, G.H. (2013). Switching
497   hemispheres: a new migration strategy for the disjunct Argentinean breeding population
498   of Barn Swallow (Hirundo rustica). *PLoS One*, 8, e55654.
499
500 Gebhardt-Henrich, S. & Richner, H. (1998). Causes of growth variation and its consequences for
501   fitness. *Oxford Ornithology Series*, 8, 324–339.
502
503 Ghalambor, C.K. & Martin, T.E. (2001). Fecundity-survival trade-offs and parental risk-taking in
504   birds. *Science*, 292, 494–497.
505
506 Gómez, C., Tenorio, E., Montoya, P. & Cadena, C. (2016). Niche-tracking migrants and niche-
507   switching residents: evolution of climatic niches in New World warblers (Parulidae).
508   *Proc R Soc B*, 283, 20152458.
509
510 Halali, S., van Bergen, E., Breuker, C.J., Brakefield, P.M. & Brattström, O. (2020). Seasonal
511   environments drive convergent evolution of a faster pace-of-life in tropical butterflies.
512   *Ecol. Lett.*
513
514 Hamblin, S. (2013). On the practical usage of genetic algorithms in ecology and
515   evolution. *Methods in Ecology and Evolution*, 4(2), 184-194.
516
517 Harvey, P.H. & Zammuto, R.M. (1985). Patterns of mortality and age at first reproduction in
518   natural populations of mammals. *Nature*, 315, 319–320.
519
520 Hutchinson, G.E. (1948). Circular causal systems in ecology. *Ann. N. Y. Acad. Sci.*, 50, 221–246.
521
522 Jetz, W., Sekercioglu, C.H. & Böhning-Gaese, K. (2008). The worldwide variation in avian
523   clutch size across species and space. *PLoS Biol.*, 6, 2650–2657.
524
525 Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. & Mooers, A.O. (2012). The global diversity of
526   birds in space and time. *Nature*, 491, 444–448.
527
528 Kooijman, S.A.L.M. (2010). *Dynamic Energy Budget Theory for Metabolic Organisation*.
529   Cambridge University Press.
530
531 Lack, D. (1947). The significance of clutch-size. *Ibis*, 89, 302–352.
532
533 Laskowski, K.L., Moiron, M. & Niemelä, P.T. (2021). Integrating Behavior in Life-History
534   Theory: Allocation versus Acquisition? *Trends Ecol. Evol.*, 36, 132–138.
535
536 Levins, R. (1962). Theory of Fitness in a Heterogeneous Environment. I. The Fitness Set and
537   Adaptive Function. *Am. Nat.*, 96, 361–373.
538
Marques, G.M., Augustine, S., Lika, K., Pecquerie, L., Domingos, T. & Kooijman, S.A.L.M.
(2018). The AmP project: Comparing species on the basis of dynamic energy budget
parameters. *PLoS Comput. Biol.*, 14, e1006100.

Martin, T.E. (1996). Life History Evolution in Tropical and South Temperate Birds: What Do
We Really Know? *J. Avian Biol.*, 27, 263–272.

Martin, T.E. (2015). Age-related mortality explains life history strategies of tropical and
temperate songbirds. *Science*, 349, 966–970.

Merkle, J.A., Monteith, K.L., Aikens, E.O., Hayes, M.M., Hersey, K.R., Middleton, A.D., *et al.*
(2016). Large herbivores surf waves of green-up during spring. *Proc. Biol. Sci.*, 283,
20160456.

Lynch. M. (1991). Methods for the Analysis of Comparative Data in Evolutionary Biology.
Evolution, 45, 1065–1080.

Moreau, R.E. (1944). Clutch-size: a comparative study, with special reference to African birds.
Ibis, 86, 286–347.

Mueller, P. & Diamond, J. (2001). Metabolic rate and environmental productivity: well-
provisioned animals evolved to run and idle fast. *Proc. Natl. Acad. Sci. U. S. A.*, 98,
12550–12554.

Muller, E.B. & Nisbet, R.M. (2000). Survival and production in variable resource environments.
Bull. Math. Biol., 62, 1163–1189.

Nakazawa, Y., Peterson, A.T., Martínez-Meyer, E. & Navarro-Sigüenza, A.G. (2004). Seasonal
Niches of Nearctic-Neotropical Migratory Birds: Implications for the Evolution of
Migration. *Auk*, 121, 610–618.

van Noordwijk, A.J. & de Jong, G. (1986). Acquisition and Allocation of Resources: Their
Influence on Variation in Life History Tactics. *Am. Nat.*, 128, 137–142.

Ortega, A. C. (2023). *Evaluating Resource Tracking and the Benefit of Migration for a
Temperate Ungulate* (Doctoral dissertation, University of Wyoming).

Piersma, T. & van Gils, J.A. (2019). *The Flexible Phenotype: A Body-Centred Integration of
Ecology, Physiology, and Behaviour*. Oxford University Press, New York, NY, USA.

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Réale, D., Dingemanse, N.J., Kazem, A.J.N. & Wright, J. (2010). Evolutionary and ecological
approaches to the study of personality. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 365,
3937–3946.

Ricklefs, R.E. (2000). Density Dependence, Evolutionary Optimization, and the Diversification
of Avian Life Histories. *Condor*, 102, 9–22.

588
589 Ricklefs, R.E. & Wikelski, M. (2002). The physiology/life-history nexus. *Trends Ecol. Evol.*, 17,
590   462–468.
591
592 Rushing, C.S. (2019). Estimability of migration survival rates from integrated breeding and
593   winter capture–recapture data. *Ecol. Evol.*
594
595 Rushing, C.S., Hostetler, J.A., Sillett, T.S., Marra, P.P., Rotenberg, J.A. & Ryder, T.B. (2017).
596   Spatial and temporal drivers of avian population dynamics across the annual cycle.
597   *Ecology*, 98, 2837–2850.
598
599 Rushing, C.S., Ryder, T.B. & Marra, P.P. (2016). Quantifying drivers of population dynamics for
600   a migratory bird throughout the annual cycle. *Proc. Biol. Sci.*, 283.
601
602 Sæther, B.-E. (1988). Pattern of covariation between life-history traits of European birds. *Nature*.
603
604 Sillett, S.T. & Holmes, R.T. (2002). Variation in survivorship of a migratory songbird
605   throughout its annual cycle. *J. Anim. Ecol.*, 71, 296–308.
606
607 Soriano-Redondo, A., Gutiérrez, J.S., Hodgson, D. & Bearhop, S. (2020). Migrant birds and
608   mammals live faster than residents. *Nat. Commun.*, 11, 5719.
609
610 Spiegel, O., Leu, S.T., Sih, A., Godfrey, S.S. & Bull, C.M. (2015). When the going gets tough:
611   behavioural type-dependent space use in the sleepy lizard changes as the season dries.
612   *Proc. Biol. Sci.*, 282.
613
614 Stearns, S.C. (1989). Trade-Offs in Life-History Evolution. *Funct. Ecol.*, 3, 259–268.
615
616 Stearns, S.C. (1992). *The Evolution of Life Histories*. OUP Oxford.
617
618 Thorup, K., Tøttrup, A.P., Willemoes, M., Klaassen, R.H.G., Strandberg, R., Vega, M., *et al.*
619   (2017). Resource tracking within and across continents in long-distance bird migrants.
620   *Sci Adv*, 3, e1601360.
621
622 Ward, M.P., Benson, T.J., Deppe, J., Zenzal, T.J., Diehl, R.H., Celis-Murillo, A., *et al.* (2018).
623   Estimating apparent survival of songbirds crossing the Gulf of Mexico during autumn
624   migration. *Proceedings of the Royal Society B: Biological Sciences*, 285, 20181747.
625
626 Wiersma, P., Chappell, M.A. & Williams, J.B. (2007a). Cold- and exercise-induced peak
627   metabolic rates in tropical birds. *Proc. Natl. Acad. Sci. U. S. A.*, 104, 20866–20871.
628
629 Wiersma, P., Muñoz-Garcia, A., Walker, A. & Williams, J.B. (2007b). Tropical birds have a
630   slow pace of life. *Proc. Natl. Acad. Sci. U. S. A.*, 104, 9340–9345.
631
632 Wikelski, M. & Ricklefs, R.E. (2001). The physiology of life histories. *Trends Ecol. Evol.*, 16,
633   479–481.

Wikelski, M., Spinney, L., Schelsky, W., Scheuerlein, A. & Gwinner, E. (2003a). Slow pace of life in tropical sedentary birds: a common-garden experiment on four stonechat populations from different latitudes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270, 2383–2388.

Wikelski, M., Tarlow, E.M., Raim, A., Diehl, R.H., Larkin, R.P. & Henk Visser, G. (2003b). Costs of migration in free-flying songbirds. *Nature*, 423, 704–704.

Williams, G.C. (1966). Natural Selection, the Costs of Reproduction, and a Refinement of Lack's Principle. *Am. Nat.*, 100, 687–690.

Winger, B.M., Auteri, G.G., Pegan, T.M. & Weeks, B.C. (2019). A long winter for the Red Queen: rethinking the evolution of seasonal migration. *Biol. Rev. Camb. Philos. Soc.*, 94, 737–752.

Winger, B.M., Lovette, I.J. & Winkler, D.W. (2011). Ancestry and evolution of seasonal migration in the Parulidae. *Proc Royal Soc B Biological Sci*, 279, 610–618.

Winger, B.M. & Pegan, T.M. (2021). Migration distance is a fundamental axis of the slow-fast continuum of life history in boreal birds. *Ornithology*.

Yanco, S.W., Linkhart, B.D., Marra, P.P., Mika, M., Ciaglo, M., Carver, A., *et al.* (2021). Niche dynamics suggest ecological factors influencing migration in an insectivorous owl. *Ecology*.

Yanco, S.W., Pierce, A.K. & Wunder, M.B. (2022). Life history diversity in terrestrial animals is associated with metabolic response to seasonally fluctuating resources. *Ecography* , 2022.

Zink, R.M. (2011). The evolution of avian migration. *Biol. J. Linn. Soc. Lond.*, 104, 237–250.

Zink, R.M. & Gardner, A.S. (2017). Glaciation as a migratory switch. *Sci Adv*, 3, e1603133.

Zúñiga, D., Gager, Y., Kokko, H., Fudickar, A.M., Schmidt, A., Naef-Daenzer, B., *et al.* (2017). Migration confers winter survival benefits in a partially migratory songbird. *Elife*, 6.

Zurell, D., Gallien, L., Graham, C.H. & Zimmermann, N.E. (2018). Do long-distance migratory birds track their niche through seasons? *J. Biogeogr.*, 45, 1459–1468.

674   **Table 1.** IBM state variables, individual parameters, and equations used to evaluate DEB and

675   movement model dynamics with potential value ranges for each parameter. During the model

676   optimization process we further restricted value ranges for some parameters to account for

677   covariation between parameters and to reduce generation of inviable solutions (see supplement

678   Appendix S1 for detail).

Symbol	Name	Detail	Domain
State variables			
f	Scaled energy in the environment available for assimilation	A Holling type II functional response of the density of food in the environment relative to uptake rate. Uptake rate held constant in simulations.	$(0,1)$
l	Scaled structural length	Proportion of current structural length relative to the maximum.	$(0,1)$
e	Scaled energy reserve density	Reserve energy available per unit of structural volume relative to maximum.	$(0,1)$
u_H	Scaled cumulative energy invested into maturation	Reflects current level of maturity.	$(0,u_H^p]$
u_R	Scaled cumulative energy invested into offspring	Accumulated buffer of energy to be converted into offspring.	$(0,\infty)$
Individual parameters			
κ	Proportion of mobilized energy invested into somatic processes	Complement of proportion invested into reproduction.	$(0,1)$
$\dot{v}$	Energy conductance rate	Ratio of maximum energy assimilation rate relative to maximum reserve capacity; determines speed of mobilization of energy from reserves. Lower values imply smaller reserve capacities or higher maximum assimilation rates.	$(0,1)$

m	Scaled migratory locomotion maintenance rate	Cost to move one volume of structure one unit of body length relative to somatic maintenance rate of one unit of volume.	(0,1)
g	Energy investment ratio	Cost to grow one volume of structure relative to maximum reserve energy density that can be committed to somatic processes.	$(0,\infty)$
u_H^b	Scaled maturity at birth	Cumulative scaled energy invested into maturation at birth.	$(0, u_H^p)$
u_H^p	Scaled maturity at puberty/sexual maturation	Cumulative scaled energy invested into maturation at puberty.	(u_H^b,∞)
d_M	Migration duration	uration in days for one-way migration between breeding and non-breeding location.	[0,365]
d_{NB}	Duration at non-breeding location	Stationary duration in days at non-breeding location.	[0,365]
Derived individual parameters			
$\dot{k}_M$	Somatic maintenance rate coefficient	Ratio between somatic maintenance rate per unit of volume and volume-specific growth costs of structure. Higher values imply higher maintenance costs or lower growth costs per unit of structural volume.	$(0,\infty)$
a_b	Age at birth	Analogous to incubation duration. Calculated based on DEB embryo development dynamics given l_b, $\dot{k}_M$, and e_b (see Appendix S1 for detail).	[0,365]
a_p	Age at puberty/sexual maturation	Age in days when sexual maturity threshold is met. $u_H^p = u_H$	[0,365]
u_E^0	Scaled initial energy reserve of embryo	Equal to scaled energy cost of one egg. Back calculated from l_b and e_b based on DEB embryo development dynamics (see Appendix S1 for detail).	$(0,\infty)$
Dynamic equations			

$\frac{de}{dt} = \frac{g\dot{k}_M}{l^3}\left(\dot{p}_A - \dot{p}_C - \frac{e}{g\kappa}\dot{p}_G\right)$	Reserve expenditure	Change in reserve energy density is the difference between assimilation and mobilization adjusted for change in structural volume due to growth.
$\frac{dl}{dt} = \dot{p}_G\left(\frac{\dot{k}_M}{3\kappa l^2}\right)$	Growth	Conversion of energy flux for growth into length.
$\frac{du_H}{dt} = \dot{k}_M * \dot{p}_R$, $if\ u_H < u_H^P$ $0, if\ u_H = u_H^P$	Maturation	Scaled energy allocated to reproduction is invested into maturation until the cumulative investment is equal to the level required to reach sexual maturity.
$\frac{du_R}{dt} = \dot{k}_M * \dot{p}_R$, $if\ u_H = u_H^P$ $0, if\ u_H < u_H^P$	Reproductive buffer investment	After maturity, scaled energy allocated to reproduction is invested into the reproductive buffer for later conversion to offspring.

Figure 1. Dynamic energetic budget model flow diagram illustrating energy uptake and

dissipation during the adult age stage where energy from the environment (*f*) is assimilated into

the organism's reserves (*e*) then mobilized for allocation to somatic and reproductive processes.

Figure 2. Model optimized migration movement paths (A) and available enroute energy (B) for

migration strategies from breeding locations with high seasonality (1-4) and mid-seasonality

(5,6) arranged by increasing migration distance top to bottom per location. In both panels, high

seasonality is represented in red, mid in green, and aseasonality in yellow and egg images depict

timing of birth and reproduction events. In panel A, solid black lines depict model optimized

movement paths between latitudes of migration endpoints colored by seasonality for each

strategy. In panel B, model optimized energy available enroute over time (black lines) overlay

energy available in breeding locations (thick opaque red and green lines) and non-breeding areas

(thick transparent lines colored by seasonality). Dashed lines depict mean annual energy

available in breeding locations (red and green) and realized enroute (black). Shaded areas

represent overwintering differences in energy availability between breeding and non-breeding

locations.

Figure 3. PCA of model optimized parameters depicting energetic trade-offs for stationary and

migratory strategies differentiated by energy management strategy (A) and migration distance

(B). Arrow length and direction indicate component loadings (correlations) of each parameter

onto PCA axes.

703
Figure 4. Fecundity of optimized parameterizations of migratory strategies in migratory

scenarios compared to simulations in which they do not migrate and remain in the breeding area.

Strategies breeding in high seasonality locales in the left panel and mid-seasonal locales on the

right with line darkness increasing with migration distance. Solid greyscale lines indicate

buffering energy management strategies and mitigation strategies are depicted as dashed lines.

Fecundity of stationary strategies depicted as colored lines in each panel for comparison. Values

for fecundity are scaled.

Figure 5. Average posterior marginal effects of niche dissimilarity (top row), migration distance

(middle row), and breeding latitude (bottom row; all scaled and centered) on POL based on

Bayesian phylogenetic mixed models. POL is measured as a size-normalized estimate of relative

maximum somatic mobilization; (see methods; small values indicate fast POL and vice versa).

The Bayesian probability of direction (PD) for each marginal effect is displayed at the bottom

right of each panel.