A first comprehensive analysis of Transcribed Ultra Conserved Regions uncovers
important regulatory functions of novel non-coding transcripts in gliomas.

Myron K Gibert Jr[1], Ying Zhang[†1], Shekhar Saha[†1], Pawel Marcinkiewicz[1], Collin Dube[1], Kadie
Hudson[1], Yunan Sun[1], Sylwia Bednarek[1], Bilhan Chagari[1], Aditya Sarkar[1], Christian Roig-Laboy[1],
Natalie Neace[1], Karim Saoud[1], Initha Setiady[1], Farina Hanif[1], David Schiff[2], Pankaj Kumar[4],
Benjamin Kefas, Markus Hafner[5], and Roger Abounader[1,2,3]

University of Virginia Departments of Microbiology, Immunology & Cancer Biology[1] Neurology[2],

Cancer Center[3], and Department of Public Health Sciences and Bioinformatics Core[4],
Charlottesville, VA, 22908, USA, and National Institutes of Health[5], Bethesda, MD, USA.

† These authors contributed equally to this work.

Corresponding author: Roger Abounader, University of Virginia, PO Box 800168, Charlottesville
VA 22908, USA, Phone: (434) 982-6634, Fax: 434-243-6843 E-mail: ra6u@virginia.edu.

<u>ABSTRACT</u>

Transcribed Ultra-Conserved Regions (TUCRs) represent a severely understudied class of
putative non-coding RNAs (ncRNAs) that are 100% conserved across multiple species. We
performed the first-ever analysis of TUCRs in glioblastoma (GBM) and low-grade gliomas (LGG).
We leveraged large human datasets to identify the genomic locations, chromatin accessibility,
transcription, differential expression, correlation with survival, and predicted functions of all 481
TUCRs, and identified TUCRs that are relevant to glioma biology. Of these, we investigated the
expression, function, and mechanism of action of the most highly upregulated intergenic TUCR,
uc.110, identifying it as a new oncogene. Uc.110 was highly overexpressed in GBM and LGG,
where it promoted malignancy and tumor growth. Uc.110 activated the WNT pathway by
upregulating the expression of membrane frizzled-related protein (MFRP), by sponging the tumor
suppressor microRNA miR-544. This pioneering study shows important roles for TUCRs in
gliomas and provides an extensive database and novel methods for future TUCR research.

<u>INTRODUCTION</u>

Transcribed Ultra-conserved Regions (TUCRs) represent 481 unique transcribed RNA molecules
that are "ultraconserved" across multiple species, including in the human, mouse (100%), rat
(100%), dog (98%), and chicken (95%) genomes. [1] TUCR expression has been found to be
highly deregulated in some cancers. Because of their ultra-conservation and their deregulation,
it is believed that TUCRs may have important regulatory roles in cancer. [2-11] About 90% of the
genome is transcribed, but only ~2 percent of the transcriptome is translated. The remainder of
the transcriptome is made up of non-coding elements that serve key regulatory roles. Of these
elements, long non-coding RNAs (lncRNAs) serve as important regulators of malignancy and
potential therapeutic targets in cancer. [2, 12-19] Due to their size and lack of known associated
protein products, it has been suggested that many TUCRs may function as lncRNAs.[2]   The
putative existence of "ultraconserved" lncRNAs is significant, as lncRNAs are typically poorly
conserved as a class of molecules.[2] Very little is known about TUCRs. [2] In particular, the
literature elucidating the expressions, functions, and mechanisms of action of TUCRs in
glioblastoma (GBM) and low-grade glioma (LGG) is nonexistent. GBM and LGG represent over
80% of primary malignant brain tumors in humans, of which GBM is the deadliest, with a median
survival of approximately 15 months. [20-28] Studying TUCRs in gliomas is therefore an
untouched avenue for understanding novel oncogenic mechanisms and discovering new
biomarkers and therapeutic targets.

In this study, we leveraged large human datasets to identify the genomic locations, chromatin
accessibility, transcription, differential expression, correlation with survival, and predicted
functions of all 481 TUCRs, and identified TUCRs that are relevant to glioma biology (Figure 1A).
Of these, we investigated the expression, function, and mechanism of action of the most highly
upregulated intergenic TUCR, uc.110, identifying it as a new oncogene. Uc.110 was highly
overexpressed in GBM and LGG, where it promoted malignancy parameters and tumor growth.
Uc.110 activated the WNT pathway by upregulating the expression of membrane frizzled-related
protein (MFRP), by sponging the tumor suppressor microRNA miR-544. This work shows
important roles for TUCRs in gliomas and provides an extensive database and novel methods for
future TUCR research in any disease context.

RESULTS

TUCRs are encoded throughout the genome, resistant to variation, and actively
transcribed.

We analyzed TUCR genomic locations published in Bejerano et al. [1] using hg38 genome
coordinates lifted over from the provided hg19 coordinates. We found that some TUCRs are
exonic and are contained within an exon of the "host" gene. Others are contained within an intron.
Some TUCRs straddle a region that spans exonic and intronic regions of the host gene
(exonic/intronic), and others are not contained within any known genetic element (intergenic)
(Figure 1B). We manually annotated each TUCR using a combination of UCSC Genome Browser
tracks, [29, 30] Quinlan Laboratory's bedtools, [31, 32] and TUCR genomic locations lifted over
to hg38 from hg19. [1] We identified 45 exonic, 231 intronic, 68 intronic/exonic, and 137 intergenic
TUCRs (Figure 1C). We found that TUCRs are located on all but one 21 numbered chromosomes
and the X chromosome. There were no annotated TUCRs on chromosome 21 (chr21), the Y
chromosome (chrY) or in the mitochondrial DNA (chrM) (Figure 1D). Detailed TUCR annotation
information for every single TUCR is provided in the supplementary materials (Supplementary
Master Table).

Since TUCRs are expected to be resistant to variation [2], we characterized the overlap of current
dbSNP (build 156) single nucleotide polymorphism (SNP) annotations to the lifted over hg38
TUCR genomic coordinates. We found that TUCRs overlap with fewer SNPs than protein coding
genes and non-coding RNAs, indicating that they are more resistant to variation (Figure 1E).

We also investigated TUCR transcription levels in comparison to transcription of known protein-
coding and non-coding genes. To accomplish this, we first analyzed their spatial associations
with markers for active chromatin (H3K4me3), active enhancers (H3K27ac), lncRNA transcription
(RNA Pol.II) and open chromatin (ATAC-Seq). We determined the significance of the spatial
relationships between these marks and TUCRs utilizing publicly available U87 CHIP- and ATAC-
Seq datasets. Then, we compared the data to TUCR intervals that were randomly shuffled to
create a negative control, other classes of non-coding RNAs, and TUCRs subset by genomic
annotation (Figure 1F and Supplementary Figure 1A). We found that TUCRs displayed a
significant enrichment for all transcriptional activity markers over expected and compared to
control. The above data show that TUCRs are distributed throughout the genome, resistant to
variation, and actively transcribed in GBM and LGG.

Figure 1

Figure 1. Annotation, localization, and expression of TUCRs in GBM and LGG. A) Experimental workflow for identifying
and studying TUCRs of interest. B) TCGA analysis shows that TUCRs can be exonic (red), intronic (blue), exonic/intronic
(purple) or intergenic (green). C) Circle graph showing the distribution of genomic annotations across all 481 TUCRs,
with colors matching 1B. D) Karyoplot showing that TUCRs exist on all chromosomes except for Chr21, the Y
chromosome and mitochondrial DNA, vertical lines show TUCRs with colors matching 1B. E) Bar chart demonstrating
that TUCRs are more resistant to single nucleotide variants (SNVs/SNPs) than other gene annotation categories. F)
Bar chart showing that TUCRs are enriched for markers for open and active chromatin in GBM cells, suggesting that
they represent transcriptionally active sites. Red bars represent chi-square expected overlaps, and teal bars represent
observed values. G) Heatmap representing TUCR absolute expression (RPKM) across multiple gene annotations. Blue
represents poorly expressed genes (<1 RPKM), White/Pink genes are moderately expressed (>=1 RPKM) and Red

104 *represents highly expressed genes (RPKM >=10). TUCRs demonstrate an expression profile that is most comparable*
105 *with protein coding genes. * = p < 0.05*

TUCRs are highly expressed in GBM and LGG tumors.

TUCR expression has not been characterized in GBM or LGG before. We performed the first
comprehensive analysis of TUCR expression in these cancers by comparing GBM (n = 166) and
LGG (n = 505) tumor samples from the Cancer Genome Atlas (TCGA) [33] to their normal brain
cortex counterparts in TCGA (n = 5) and the Genotype-Tissue Expression Database (GTEx, n =
255). [34] We first analyzed absolute TUCR expression, as measured by reads-per-kilobase
million (RPKM). The absolute expression, in GBM, of all TUCRs was compared to the expression
of lncRNAs, coding genes, antisense RNAs, and small noncoding RNAs (< 200 nt length), and
the expression of TUCRs separated by genomic annotation into exonic, intronic, exonic/intronic,
and intergenic. All gene annotations were obtained using the CHESS gene catalog, which
contains most Refseq and Ensembl genes, while also including a series of understudied novel
genes.[35] Highly expressed genes are visualized via heatmap (>=10 RPKM, red) along with
moderately (>=1 RPKM, white) and lowly expressed genes (<1 RPKM, blue). These analyses
were repeated in LGG (Supplementary Figure 1B). The data show that intragenic TUCRs are
expressed at magnitudes that are like those of protein coding genes in both GBM and LGG, while
intergenic TUCRs demonstrate expression levels that are closer to those of lncRNAs (Figure 1G).
122

TUCRs are deregulated in gliomas, and deregulation is associated with clinical outcomes.

We analyzed TCGA tumor data and GTEx normal brain cortex data and found that in addition to
being highly expressed in gliomas, TUCRs are highly deregulated in GBM and LGG as compared
to normal brain cortex. Of the 481 annotated TUCRs, we identified 87 that were upregulated and
67 that were downregulated in GBM (Figure 2A). We also identified 59 TUCRs that were
upregulated and 53 TUCRs that were downregulated in LGG. (Figure 2B). Of the 154 deregulated
TUCRs in GBM, 86 were also deregulated in LGG, a 56% overlap (Figure 2C). We then sought
to determine whether deregulation of TUCR expression correlates with patient outcomes in GBM
and LGG. For each of the 481 TUCRs, we generated a Kaplan-Meier plot tracking differences in
survival for high expressing (upper quartile) and low expressing (lower quartile) tumor groups. Of
the TUCRs that are expressed in GBM TCGA RNA-Seq data, only 4 were correlated with survival
in a statistically significant manner in both of our workflows (p <= 0.05, Supplementary Figure 2B).
We considered that this low prevalence of survival associated TUCRs in GBM was due to the
short overall survival of GBM patients (~15 months). We also studied survival differences in LGG
patients, as they have a longer median survival (~84 months). Of the TUCRs that are expressed
in LGG TCGA RNA-Seq data, 93 were correlated with survival in both of our workflows (Figure
2D). We have highlighted two TUCRs that represent a statistically significant correlation with good
(uc.338, Figure 2E) or poor (uc.75, Figure 2F) prognosis using both methods. When separated
by annotation category, intragenic TUCR deregulation has a greater association with patient
outcomes than intergenic TUCR deregulation (Supplementary Figure 2C).   Expression,

deregulation, and survival analyses were performed on all 481 TUCRs. Detailed results for
individual TUCRs can be found at www.abounaderlab.org/tucr-database/.
145

TUCRs are coregulated with genes that have specific functions.

We predicted TUCR functions by identifying coregulated genes with known functions via weighted
gene co-expression network analysis (WGCNA).[36] We aggregated the 42,644 genes in our
dataset into 60 colored modules based on clustered gene ontology (GO) terms. Each of these
modules contains genes with known functions, such as RNA binding, cell signaling, immune
response, metabolic response, etc. These modules can also be used to identify genes that
associate with clinical traits, such as the tumor tissue-type (Figure 2G). The data can also be
used to predict gene function for novel genes. To do this, we aggregated all 481 TUCRs into our
modules. We identified TUCRs that correlate with each of the 60 modules, with some having
positive correlations and others negative. For example, TUCRs that exhibit a positive correlation
with the #004C54 "midnight green" module (Supplementary Figure 6) could have a promoting
effect on nucleic acid binding and regulation, while those that are negatively correlated with the
#f4a460 module (Supplementary Figure 7) may have a negative effect on G-protein coupled
receptor and metabolic functions. Since many different TUCRs show associations with different
modules, and every module has at least one TUCR that is associated with it, these results suggest
that TUCRs may have a broad range of potential functions in GBM and LGG (Figure 2H). WGCNA
analyses were performed on all 481 TUCRs. Detailed results for individual TUCRs can be found
at www.abounaderlab.org/tucr-database/.

Figure 2

86 were also deregulated in LGG, a 56% overlap. Dark Red/Blue are TUCRs deregulated in GBM. Light Pink/Blue are
TUCRs deregulated in LGG. Intermediate Red/Blue represent TUCRs deregulated in both. D) Volcano plot showing
that several TUCRs are significantly associated with patient outcomes in LGG. Pink dots represent TUCRs significantly
associated with poor prognosis. Blue dots represent TUCRs significantly associated with good prognosis. E) Kaplan-
Meier showing that TUCR uc.338 is significantly associated with good prognosis. Red line represents the uc.338 high
expression group. Teal line represents the uc.338 low expression group. F) Kaplan-Meier showing that uc.75 is
significantly associated with poor prognosis (Line colors as described in E). G) Gene similarity dendrograms from
weighted gene correlation network analysis (WGCNA). 42,644 genes were aggregated into 3 "blocks" by gene similarity
and were then further aggregated into 51 linkage modules using TUCR expression as trait data. Modules are denoted
with distinct color hex codes. (e.g. #004C54 is the "midnight green" module). H) Heatmap showing that TUCRs
demonstrate association with all 60 modules, suggesting broad potential functions. Red and Blue represent positive and
negative correlations, respectively. I) Volcano plot showing that the uc.110 TUCR is the most upregulated TUCR in
*GBM and LGG (Line colors as described in E). * = p < 0.05*

183 **TUCR, uc.110, is highly upregulated in gliomas and is predicted to bind nucleic acids.**

The expression and deregulation of intergenic TUCRs is of particular interest as they may
represent novel lncRNAs due to their similar expression levels, genomic location, and lack of
coding potential.[2] These TUCRs are also easier to study experimentally; they are often
thousands of kilobases (kb) from the nearest protein-coding gene and likely function in a manner
that is independent of a "host gene". Because of this, we focused on intergenic TUCRs for our
experimental studies. Of the deregulated intergenic TUCRs in GBM and in LGG (Figure 2I), we
found that uc.110 is the most upregulated as compared to normal brain; 30-fold in GBM and 61.4-
fold in LGG (Figure 3A). It has near binary expression; it is rarely expressed at all in normal brain
but is very highly expressed in GBM and LGG (Figure 3B). Due to its high expression, we
hypothesized that this TUCR is functioning as an oncogene.

Since many TUCRs exist as a part of a larger transcript [2], we first determined the sequence of
the uc.110 full transcript. We utilized machine learning and *de novo* transcript reassembly using
TCGA and GTEx RNA-seq data to reconstruct RNA-Seq transcripts in the absence of a reference
genome (Supplementary Figure 3A). [35] We identified a 2,158 nucleotide (nt) long RNA molecule
that contains the 243 nucleotide (nt) uc.110 ultraconserved sequence (Figure 3C) as a novel
transcript. We confirmed the existence of this transcript experimentally using PCR amplifications
and sequencing (Supplementary Figure 3B).

After identifying the sequence for the full uc.110 transcript (Supplementary Figure 3C), we utilized
our WGCNA workflow to identify genes and modules (Figure 3D) that are significant to this
transcript. Of note, one of the top modules for uc.110 by module association is the #004C54
module, which represents genes that are involved in nucleic acid and protein binding
(Supplementary Figure 6). This is a published function for some TUCRs. [2] Genes that are
members of these modules are positively coregulated with uc.110 (Figure 3E). Based on these
findings, we hypothesized that uc.110 may be operating as an oncogenic RNA-binding molecule.
We also performed similar analyses for all 481 TUCRs to identify potential functional roles for
each TUCR in gliomas. Examples of an oncogenic TUCR (uc.2, Supplementary Figure 5) and a

tumor suppressor TUCR (uc.15, Supplementary Figure 5) are depicted in this manuscript, while
the analyses of the rest of the 481 TUCRs can be found at www.abounaderlab.org/tucr-database/.

Figure 3. The uc.110 TUCR is the most upregulated intergenic TUCR in gliomas and is predicted to bind nucleic acids.
A) Box- and dotplot showing that uc.110 is 30-fold upregulated in GBM and ~60-fold upregulated in LGG based on
TCGA and GTEx data analyses. Red boxes represent upregulated TUCRs. B) Box- and dotplot showing that uc.110 is
expressed in tumors but is poorly expressed in normal brain cortex based on TCGA and GTEx data analyses. C) Cartoon
showing that uc.110 is a 243 nt region in a 2,158 nt transcript. D) Heatmap depicting uc.110 gene module association.
Positive correlations are red, while negative correlations are blue, with weak correlations in white. Modules with no
linkage are gray. E) Scatter plots depicting uc.110 association with top 3 positive (top row) and negative (bottom row)

correlation modules. Caption lists the module name, number of genes in the module, and the significance of uc.110
association with the module.

uc.110 has oncogenic effects in GBM.

To determine the function of uc.110 in GBM, we first used qPCR to investigate the expression of
uc.110 in our banked tumor samples compared to normal brain cortex and cell lines compared to
normal human astrocytes. We independently confirmed the results from our TCGA analysis by
showing uc.110 is highly upregulated in GBM tumors (Figure 4A, 4B and Supplementary Figure
8). We then designed two siRNAs that target separate regions on the uc.110 RNA, one that
begins at nucleotide 96/243 (si-uc.110-1) and one that begins at nucleotide 195/243 (si-uc.110-
2), as well as a scrambled control (si-SCR) (Supplementary Figure 8A). We generated stable
A172 and U251 GBM cell lines that express uc.110 (LV-uc.110) or the empty expression vector
(LV-pCDH). We subjected these cell lines to siRNA transfection and assessed the effects on cell
counting, survival and invasion assays (Supplementary Figure 8B). We used qPCR to show that
uc.110 is generally, though not uniformly, upregulated in GBM cells (Supplementary Figure 8).
Based on these data, we prioritized the use of cell lines that overexpress uc.110 (A172, U251) for
knockdown experiments, and cells that express low levels of uc.110 (U87, GSC-28) for
overexpression experiments. We confirmed that siRNAs targeting of uc.110 lead to knockdown
of uc.110 expression in A172 and U251 cells. (Figure 4C) We also confirmed that LV-uc.110
overexpresses uc.110, and that this overexpression rescues uc.110 bioavailability in A172 and
U251 cells (Figure 4C).

Next, we performed cell counting assays [20, 37-39] to determine the effects of uc.110 knockdown
and rescue on cell accumulation. When we reduced uc.110 expression, we reduced cell
accumulation in A172 and U251 cells (Figure 4D). When we rescued uc.110 bioavailability by
restoring its expression, the cell accumulation phenotype was restored in A172 and U251 cells
(Figure 4E). We then used AlamarBlue [40, 41] to measure cell viability. When we reduced uc.110
expression, A172 and U251 cell viability was reduced. We were able to rescue this phenotype
by increasing uc.110 bioavailability (Figure 4F). We observed a similar phenotype in a glioma
stem cell line that overexpresses uc.110 (GSC-34, Figure 4G).

We then investigated the invasive potential of uc.110 using a transwell invasion assay. [42-44]
Knockdown of uc.110 reduced A172 and U251 cell invasion through a collagen IV matrix (Figure
4H). When uc.110 bioavailability was increased, a partial recovery of the phenotype was
observed (Supplementary Figure 8F). Lastly, we overexpressed uc.110 in U87 and GSC-28 cells
(Figure 4I) and determined that this leads to increased cell accumulation compared to the empty
vector after 7 days (Figure 4J) in U87 and GSC-28 cells. These data show that uc.110 has
oncogenic effects in GBM cells and stem cells.

Figure 4

Figure 4. The uc.110 TUCR operates as an oncogene. A) Bar graph depicting uc.110 upregulation in banked UVA GBM
tumors versus normal brain cortex. B) Boxplot representing uc.110 expression in pooled tumors versus normal brain.
Red boxes indicate an upregulated TUCR. C) Bar graph depicting qPCR validation of uc.110 siRNA knockdown and
rescue in A172 and U251 cell lines. Facets represent cell lines. si-SCR = scrambled control siRNA (red), si-uc.110-1 =
siRNA targeting uc.110 at nucleotide 96/243 (green), si-uc.110-2 = siRNA targeting uc.110 at nucleotide 195/243 (blue).
D) Line graph showing that knockdown of uc.110 reduces A172 and U251 cell accumulation over a 5–7 day period.
Facets represent cell lines. E) Bar graph depicting that the cell accumulation phenotype is rescued when uc.110 is
overexpressed in the presence of siRNA. Facets represent cell lines. Images are representative of the listed sample. F)
Bar graph showing that knockdown of uc.110 reduces A172 and U251 cell viability via Alamar Blue assay and can be
rescued with uc.110 overexpression. Facets represent cell lines. G) Bar graph showing that knockdown of uc.110
reduces GSC-34 glioma stem cell viability via Alamar Blue. H) Bar graph showing knockdown of uc.110 reduces A172
and U251 cell invasion and migration. Images are representative of the listed sample. I) Bar graph showing the qPCR
validation of overexpression in U87 cells. J) Bar graph showing that overexpression of uc.110 increases cell
*accumulation in U87 and GSC-28 cells. Facets represent cell lines. * = p < 0.05*

After determining that uc.110 displays an oncogenic phenotype *in vitro*, we sought to determine
whether this effect is recapitulated *in vivo*. U251 GBM cells were transfected with si-uc.110-1 or
si-uc.110-2. After 2 days, these cells were implanted into immunodeficient mice using intracranial
injection (Supplementary Figure 9). [37, 38, 45, 46] Tumor growth was monitored by MRI and
mouse survival was observed over a period of 70 days. Mice that were xenografted with U251
cells that were transfected with si-uc.110-1 and si-uc.110-2 expression developed smaller tumors,
as depicted, and quantified by MRI (Figure 5A, 5B). The mice that received si-uc.110 also
displayed better overall survival than mice that received scrambled control siRNA cells (Figure
5C).

Figure 5

Figure 5. The uc.110 TUCR promotes tumor growth in vivo. A) MRI images reveal a reduction in tumor size when uc.110
is knocked down via siRNAs. B) Bar graph showing that knockdown of uc.110 leads to a reduction in tumor volume. si-
SCR = scrambled control siRNA (red), si-uc.110-1 = siRNA targeting uc.110 at nucleotide 96/243 (green), si-uc.110-2
= siRNA targeting uc.110 at nucleotide 195/243 (blue). C) Kaplan-Meier plot showing that knockdown of uc.110 leads
to increased mouse survival.

uc.110 regulates the expression of the Wnt pathway member, Membrane Frizzled Related
Protein (MFRP).

LncRNAs can have various functions that depend on their subcellular localization. Nuclear
lncRNAs are usually involved in transcriptional regulation, while cytosolic lncRNAs are usually
involved in translational and spatial regulation. [2] We fractioned four GBM cell lines (A172, U251,
U87, U1242) into nuclear and cytosolic fractions. When compared to nuclear (U44, U48) and
cytosolic (GADPH, PPIA) controls, uc.110 appears to be localized to both the nucleus (mainly in
U87, U251, and U1242 cells), and the cytoplasm (mainly in A172 cells) (Supplementary Figure
8G). We then performed RNA-Seq on A172 cells that had been transfected with si-SCR, si-
uc.110-1, or si-uc.110-2 for 48 hrs. and found several genes that are deregulated when uc.110
expression is downregulated (Figure 6A). To identify genes that are particularly related to uc.110
function, we focused on genes that demonstrated coregulation with uc.110 in our WGCNA
analysis (Figure 3E). Of particular interest was the membrane frizzled related protein, also known
as MFRP. [47, 48] MFRP serves as a shuttle for the Wnt-ligand, and functions as an activator of
the Wnt-signaling pathway. This gene was the only gene in our analysis that correlated with
uc.110 expression, was upregulated in GBM tumors, and downregulated when uc.110 is knocked
down in A172 cells, suggesting MFRP coregulation with uc.110. (Figure 6B).

uc.110 sponges the tumor suppressor microRNA miR-544 to increase the bioavailability
of MFRP and WNT activity in GBM.

One common lncRNA mechanism of action is as a miRNA sponge, acting as a binding competitor
for various miRNAs and therefore increasing the bioavailability of those miRNAs' targets. [2, 49,
52-53] Based on the WGCNA data that we generated above, we hypothesized that uc.110 may
function by sponging miRNAs away from MFRP transcripts, as their expression relationship is
consistent with such an interaction. We hypothesized that a tumor suppressor miRNA can
successfully target and suppress MFRP in the normal brain (Supplementary Figure 10A). This
leads to downstream activation of Wnt target genes involved in biological processes such as cell
accumulation, invasion, and stem cell differentiation (Supplementary Figure 10B). We further
hypothesized that in glioma tumors, uc.110 is activated and acts as a binding competitor for this
miRNA (Supplementary Figure 10C), increasing the bioavailability of MFRP and increasing Wnt
pathway signaling (Supplementary Figure 10D). To identify candidate miRNAs that are consistent
with the afore mentioned hypothesis, we screened public databases and published literature for
GBM tumor suppressor miRNAs that are predicted to bind to both uc.110 and MFRP. The only
miRNA that fulfilled these criteria was miR-544. We first investigated the functional effects of miR-
544 in GBM cells. Transfection of miR-544 into U251, A172, and T98G GBM cell lines reduced
cell accumulation after 5 days (Figure 6C). Expression of both uc.110 and MFRP in GBM cells
was reduced when transfected with miR-544 or si-uc.110 (Figure 6D). To further test the
hypotheses, we asked if miR-544 targets both uc.110 and MFRP, and if this binding affects Wnt
signaling. To determine whether MFRP and uc.110 are direct targets of miR-544, we constructed
luciferase reporter vectors by inserting the uc.110 ultraconserved region and MFRP 3'UTR
downstream of hRluc followed by Synthetic Poly(A) using psiCHECK-2 backbone vector

(Promega) (Figure 7). We first measured target binding by transfecting the reporter constructs
followed by transfection with miR-544 or miR-SCR (scrambled control) in GBM cells. Ectopic
expression of miR-544 significantly decreased luciferase activity compared to miR-SCR (Figure
7D, left panel and figure 7E, left panel).   These binding sites for miR-544 were predicted via
computational algorithms and validated via sequencing. We then mutated the binding sites for
MFRP and uc.110 (Supplementary Figure 10, Figure 7C) and assessed signal strength again.
The data showed that luciferase activity was not significantly altered in mutant-reporter-vectors
transfected cells (Figure 7D, right panel and figure 7E, right panel), indicating that miR-544 binds
to both uc.110 and MFRP in GBM cells, and that this binding is lost when the miRNA binding sites
are mutated.

Figure 6

Figure 6. The uc.110 TUCR activates Wnt-signaling by sponging miR-544 from membrane frizzled related protein
(MFRP) 3'UTR. A) Volcano plot depicting transcriptome deregulation in RNA-Seq data on A172 GBM cells transfected
with si-uc.110. Purple dots represent genes that are significantly deregulated >= 2-fold. Blue dots represent genes that
are significantly deregulated. Pink dots represent genes that are deregulated >= 2-fold. Gray dots are neither
deregulated nor significant. B) Dot plot showing that, of the genes that are predicted miR-544 targets, MFRP is the only
gene that is upregulated in GBM Tumors and downregulated when uc.110 is downregulated. Purple dots represent
predicted miR-544 targets that are deregulated in A172 cells from 6A and TCGA RNA-Seq data. Pink dots represent
predicted miR-544 targets that are deregulated in A172s from 6A only. C) Bar graph showing that miR-544 transfection
reduces cell accumulation in A172, T98G, and U251 cells, confirming its tumor suppressor role. Facets represent cell

lines. si-SCR = scrambled control siRNA (red), si-uc.110-2 = siRNA targeting uc.110 at nucleotide 195/243 (blue). miR-
544 = miR-544 (yellow). D) Bar graph showing that transfection with miR-544 or si-uc.110-2 reduces uc.110 and MFRP
*expression. Facets represent genes and cell lines. * = p < 0.05*

Lastly, we asked if uc.110 expression alters Wnt pathway activity. To answer this question, we
studied one of the most established downstream targets of Wnt-signaling, the T cell
factor/lymphoid enhancer factor family (TCF/LEF). When Wnt-signaling is activated, TCF/LEF is
produced downstream and activates Wnt-signaling target genes. Therefore, TCF/LEF activity can
be used as a proxy for pathway activity and can be measured with a TCF/LEF luciferase reporter
assay. (Figure 7F). The activity of this reporter can be regulated by either directly reducing Wnt
bioavailability with miR-544 or indirectly by targeting uc.110 with siRNA. If upstream Wnt signaling
is reduced, the luciferase construct will bind fewer activators and exhibit decreased signal.
Likewise, we would expect that overexpression of uc.110 would rescue the bioavailability of MFRP
and consequently also downstream activation of the TCF/LEF construct. We found that
transfection of A172 and U251 cells with si-uc.110 and miR-544 reduced reporter activity in A172
(Figure 7G) and U251 (Figure 7H) cells, and that this effect can be rescued via uc.110
overexpression. These data taken in conjunction provide strong support for a miRNA sponge
model for the uc.110 oncogene. Altogether, the above data demonstrate an important role for
uc.110 in regulating the Wnt pathway in GBM by sponging the Wnt inhibitory miRNA miR-544
(model shown in Supplementary Figure 10).

Figure 7

Figure 7. The uc.110 TUCR activates Wnt-signaling by sponging miR-544 from membrane frizzled related protein
(MFRP) 3'UTR. A) Schematic depicting the uc.110 luciferase construct used to demonstrate binding to miR-544 and
si-uc.110-2. Binding of miR-544 to binding sites (orange) leads to a degradation of construct and a reduction in Renilla
signal (Green) B) Schematic depicting the MFRP luciferase construct used to demonstrate binding miR-544. C)
Schematic depicting miR-544 binding site mutations for uc.110 (two sites) and MFRP (one site). Top row represents
wild-type binding sites. Middle row is the miR-544 binding region. Bottom row are mutated sites. D) Bar graph showing
that transfection of miR-544 reduces uc.110 and MFRP luciferase expression signal in A172 and E) U251 glioma cells,
and that mutating miR-544 binding sites rescues luciferase signal. Facets represent cell lines and miR-544 binding site
mutation status. F) Schematic depiction of TCF/LEF luciferase reporter construct used to measure downstream Wnt-
signaling pathway activation. TCF/LEF binds to the reporter region (green) and activates luciferase (yellow). F) Bar
graph showing that transfection of si-uc.110-2 and miR-544 reduces TCF/LEF reporter signal in A172 and H) U251
*glioma cells. Signal is rescued when uc.110 is overexpressed in the presence of siRNA or miR-544. * = p< 0.05 the*
letters.

DISCUSSION

This study investigated Transcribed Ultraconserved Regions (TUCRs), a set that might contain
long noncoding RNA sequences that are fully conserved across human, mouse, and rat genomes.
These TUCRs are distinct due to their exceptional conservation, which often signifies functional
importance. Despite their potential significance, TUCRs have been minimally explored, especially
in relation to cancer. Of note, the findings of this study represent the first of their kind on TUCRs
in gliomas and the first comprehensive analysis of TUCR expressions and functions in any cancer.
They contribute critical new insights into an uncharted area of glioma biology, while also providing
a novel framework for studying TUCRs in other cancers and other diseases, where they are also
understudied.

We found that TUCRs are located across the genome, resistant to variation, and actively
transcribed. We manually annotated each as either exonic, intronic, exonic/intronic, or intergenic.
We identified distinct signatures for intergenic and intragenic (exonic, intronic, exonic/intronic)
RNAs. Intragenic TUCRs are expressed at a level that is most like coding genes, while intergenic
TUCRs more closely resemble lncRNAs. We then performed the first analysis of TUCR
expression in gliomas and found that the majority of TUCRs are deregulated >= 2-fold in GBM
and LGG, with a 56% overlap. This shows that TUCRs are not only expressed, but also frequently
dysregulated in gliomas compared to normal brain tissue. This is critical, as their high degree of
conservation and dysregulation suggests that they may serve critical biological functions. We then
extended our analysis to TUCR correlation with patient survival. In GBM, the extremely short
survival times (15 months) limit the detection of significant correlations. However, patients with
LGG live substantially longer (84 months), and therefore more TUCRs are associated with patient
outcomes in this disease, suggesting a potential impact on glioma patients' prognoses and
indicating possible novel biomarkers. Another facet of our research involved predicting the
functions and mechanisms of action of TUCRs in gliomas. We studied this for the first time in
gliomas WGCNA workflows to cluster TUCRs and provide functional predictions based on shared
functions between coregulated genes. This approach identifies a wide range of potential functions
for TUCRs, encompassing activities such as nucleic acid binding regulation, stem cell
differentiation, organ development, immune response, and cell signaling.

We found intergenic TUCRs to be of notable interest because they resemble lncRNAs but are
much more highly conserved and experience less sequence variation. Notably, these TUCRs do
not overlap with known genes, suggesting they might represent novel lncRNAs. Of these TUCRs,
uc.110 is the most upregulated in both GBM and LGG. Knocking down uc.110 reduces cancer
cell characteristics *in vitro* and *in vivo* and improves survival in mouse models. On the other hand,
increasing uc.110 expression increases malignancy in cells that do not express it, further
indicating its potential oncogenic role. We explored uc.110's function via WGCNA, revealing its
membership in modules associated with oncogenic nucleic acid binding. We integrated these data
with transcriptome deregulation data (RNA-Seq) post-uc.110 knockdown, revealing a close

relationship between uc.110 and the oncogenic membrane frizzled-related protein (MFRP). This
protein is involved in activating the Wnt-signaling pathway, impacting cell proliferation, invasion,
migration, and stem cell differentiation. From these data we hypothesized that uc.110 might
sponge tumor suppressor miRNAs from MFRP, enhancing Wnt signaling activation. Accordingly,
we demonstrated that one mechanism of action for the uc.110 oncogene is as a miRNA sponge
for miR-544, therefore increasing the bioavailability of MFRP and Wnt activation.

In conclusion, our results indicate that TUCRs are an important class of regulatory RNAs. They
are more highly conserved than typical genes and more resistant to variation, which suggests
biological importance. They are perturbed in gliomas, and this perturbation is associated with
clinical outcomes. Our predicted functions reveal that TUCRs are widely involved in cancer-
related biological processes. Some TUCRs previously thought to be intergenic may represent
previously undiscovered genes. Our findings also identify and characterize uc.110 as a new
oncogene in gliomas. Each of the experiments performed in our study represents the first of its
kind in gliomas. We have developed, adapted, and presented novel methods for studying TUCRs
that can be used in other cancers and other diseases, where TUCRs remain very understudied.
These methods and the data derived from them represent a "TUCR database" that will serve the
scientific community in future TUCR studies in gliomas and other diseases, where they remain
unstudied or understudied.

SUPPLEMENTARY DATA

Supplementary Figure 1

Supplementary Figure 1. Annotation, localization, and expression of TUCRs in GBM and LGG. A) Bar chart showing
that TUCRs are enriched for markers for open and active chromatin in GBMU87 cells, suggesting that they represent
transcriptionally active sites. Red bars represent chi-square expected overlaps, and teal bars represent observed
values. B) Heatmap representing TUCR absolute expression (RPKM) across multiple gene annotations. Blue
represents poorly expressed genes (<1 RPKM), White/Pink genes are moderately expressed (>=1 RPKM) and Red
represents highly expressed genes (RPKM >=10). TUCRs demonstrate an expression profile that is comparable with
*protein coding genes. * = p < 0.05*

Supplementary Figure 2

Supplementary Figure 2. TUCRs are deregulated and associated with patient outcomes in gliomas. *All experiments were performed using TCGA GBM and LGG RNA-Seq data. A) Volcano plots showing that TUCRs are deregulated in every TUCR annotation category in GBM and LGG. Red dots are upregulated. Blue are downregulated. B) Volcano plot showing that few TUCRs are significantly associated with patient outcomes in GBM. Red dots represent TUCRs significantly associated with poor prognosis. Blue dots represent TUCRs significantly associated with good prognosis. C) Volcano plot showing that TUCRs in every TUCR annotation category are associated with survival in gliomas. Red dots represent TUCRs significantly associated with poor prognosis. Blue dots represent TUCRs significantly associated with good prognosis.*

Supplementary Figure 3

uc.110

A

Chrom	Start	End	Length	ID
chr2	236,162,737	236,162,980	243	uc.110
chr2	236,161,295	236,164,636	3341	TUC110

C

TCGGCTTAGGCCCCCACTTCTGAACTGTTCAAATAGACAATCGGTTTTGATGATGTAATTTATAATTTATGCAGGTT
GTCTACATACATATGGTGTTGCTTATGTGCGTTGGGGTTTCAGGTGCTTTATTACTCTGCCGGTGTGATTTGCATTC
ATGTTTGGTTACCAAGATCATATTTCTCCTGTTTACACCCTAAGGGCCTAAAATGAAGATGAATGCCACCTCCCTTC
GTCTGGGCTGTCCCTGTAGAACCCCGGCCTTAGCTTCCCCCAAGACCCCAGAGCCCTGCTACCCTGGAACTGGGCC
CTGAGCACTGCGCTGAAGCTGCCTGGGAGGACCCAGCGGCCAGAAGGTCGGAGCAGGCCACCTGCTCTGGCTGT
GCATGCCTGGGCAGATGCCTGGGGTCTCTGGGGACAGAAATAGGTGTGTGCACTGGAGCCTCCTTCCCTGGGAA
GCCACGGCTGGGCATAAGCGAGCCACTCTGCCACACCGCCACTGAAGTCCAGCAGGGGAAAGAAGAGGGCCCCG
AGAGGGTCTCTGCTGCCCGCGCTGCTGAATGCGTGGGCTCCAGCCACTTGGGACTAGGAAGGCCCCGAGGTGAG
AACCAGAGTGTGATCCAAAGGCTCCTGAGGCCGGAAAGCTAGAAGGTCACTCCCGGCTTTCAGTGCACCTAGAAC
CACCCGCCATGGGGCATAACATGGGACTCTGTTCCAGACAGGTCTGGGGCTGGAGGGGAGCAGCCTTGGAAGCCC
TGTGCGAGCTTCACCAGATGCAGCTGAAATAATTAAGCCCATTAAGCTCAAACAAACAGAAACAAGAGGGAGAG
GGTGCTGACCCTGAGAGAAAGGGTGTCTTGGAATGCAGGTACCAGCGATGTGCCCAGGCTTGTACAAGCACCTA
GGCTGAAGGCCAGACGCACAGACAGCCATAGAGTCAGGATGTCCTCCGAGGCAAACCCGGGCTCAAAAAAGCCT
GGGGGGGGCCACTCCCTGCTCTCCATTTAGCTGACAAGGCATCTGTGAAAGAACCACCCAGGGGGCTGAAGACAG
GAAGCAAAGGCTGGATGCACAAGCATGCACTAAGTCTCGAAAATTCCAGCGCAGGTTGGGATTTGAGCAGCTTCC
AGTGCAGGTGGGTCCCGGCTGCTGGGAAGGTAGTCCTGGGCTCCTTCTGGAAGGGGAAGTTGCTTGAGGCCCAG
GACCAAACTAAAGAGGGTCCTGCCCCAAGAACAGGGAGATGGCTGCGGATTTTCAGCATACCTTGGGTTCTCCAA
TGGGGCAGGGGTGCCTGGCTTGCCCAGGGAGCAAAGGCAGTCACTGGGTGGAGTGACTGTTCTTGTGGCCCCCCT
CCGACCTCTGCCTGGGTCGGTGGCTGCGGGGTGCAGGCAGCCCTGAGGCGTCCCAGTGCACGGCTGGGGGGCGC
AGGTCCAAGGAGCTCCCAAGTGTAATTATTTGATctgcagatgtaagtggaatttattgtaacaacaaatatagtgatgtcaaaaccc
aaccttggattaaagccggcagccaaaggggaagtgtattaatttccaacagcatctcaggccagagcctattagaacaagctattcttcaggccccc
ccatcagaattaatcattgggagttaatttgaagctcagacaagttgctgttaatttagtgcagggaggacgggatggaataaaaagctgaggtgctg
gccgagcctcgtgggtctcattaatcagccagctgagacggcctggcttgattacaatttgcaaaaaaattcattagggcccggcgattgacaattttt
ctctctgcctgtgatgtgactcattaggcagccaaatgaaagccatgatgacggcccatgatgacagcggccatcaagcgtgcttgtttatcttcctccc
ctcggcctgtctcccgggccttggcctccctgccgggctctcggggcgccccggggctcaggggtttaggagggagggggtcagcgccccgaggag
caggaggggaggcggcatcggccagcttggggagctcgggccctctcctcctcgtgcctccatagggcgaggacctcgaggctgggagggcccgga
gctcggtcccaggagcgggcttcctgacggcggcctccggttcctagtgggcttcgcttt

Supplementary Figure 3. Elucidation of the uc.110 TUCR full transcript sequence. A) We used de novo transcript
reassembly of TCGA glioma RNA-Seq data and experimental PCR validation to identify the predicted sequence for the
novel full RNA transcript containing uc.110. Table depicts uc.110 ultraconserved and predicted full transcript genomic
locations and length in nucleotides (nt). B) PCR gel electrophoresis demonstrating validated uc.110 transcript variants.
C) The validated full sequence of the 2,158 nt uc.110 transcript is provided. The ultraconserved uc.110 region is colored
red and primer sequences are colored green. The additional variants (1-4 and 1-5) were also validated but de-
emphasized due to variant 1-3 having the strongest functional effect.

Supplementary Figure 4. An exploration of a putative oncogenic TUCR, uc.2 in gliomas. _A) Box- and dotplot showing_
uc.2 deregulation in GBM and LGG. Facets represent disease type. Red boxes represent upregulated TUCRs. Green
boxes represent downregulated TUCRs. Gray boxes represent TUCRs that are not deregulated. B) Box- and dotplot
showing uc.2 absolute expression in GBM and LGG. Facets represent disease type. C) Heatmap depicting uc.2 gene
module association. Positive correlations are red, while negative correlations are blue, with weak correlations in white.
Modules with no linkage are gray. D) Scatter plots depicting uc.2 association with top 3 positive (top row) and negative
(bottom row) correlation modules. E) Kaplan-Meier showing uc.2 association with GBM prognosis. Red line represents
the TUCR high expression group. Teal line represents the TUCR low expression group. F) Kaplan-Meier showing uc.2
association with LGG prognosis. Red line represents the TUCR high expression group. Teal line represents the TUCR
low expression group. (Similar analyses and figures for all 481 other TUCRs available at www.abounaderlab.org/tucr-
database/)

Supplementary Figure 5

uc.15

Supplementary Figure 5. An exploration of a putative oncogenic TUCR, uc.15 in gliomas. A) Box- and dotplot showing
uc.15 deregulation in GBM and LGG. Facets represent disease type. Red boxes represent upregulated TUCRs. Green
boxes represent downregulated TUCRs. Gray boxes represent TUCRs that are not deregulated. B) Box- and dotplot
showing uc.15 absolute expression in GBM and LGG. Facets represent disease type. C) Heatmap depicting uc.15 gene
module association. Positive correlations are red, while negative correlations are blue, with weak correlations in white.
Modules with no linkage are gray. D) Scatter plots depicting uc.15 association with top 3 positive (top row) and negative
(bottom row) correlation modules. E) Kaplan-Meier showing uc.15 association with GBM prognosis. Red line represents
the TUCR high expression group. Teal line represents the TUCR low expression group. F) Kaplan-Meier showing uc.15
association with LGG prognosis. Red line represents the TUCR high expression group. Teal line represents the TUCR
low expression group. (Similar analyses and figures for all other 481 TUCRs available at www.abounaderlab.org/tucr-
database/)

Supplementary Figure 6

A

B

Supplementary Figure 6. Top positively correlated TUCR modules in gliomas. A) The #0000FF (blue) module is the
most positively correlated with TUCRs. B) The #004C54 module (midnight green) is the second most positively
correlated module with TUCRs.

Supplementary Figure 7

Supplementary Figure 7. Top positively correlated TUCR modules in gliomas. A) The #f4a460 (sandybrown) module is the most positively correlated with TUCRs. B) The #FFA500 module (orange) is the second most positively correlated module with TUCRs.

Supplementary Figure 8

Supplementary Figure 8. The uc.110 TUCR operates as an oncogene (cont.) A) Cartoon depicting two siRNAs that
target different regions of the uc.110 TUCR. One starts at nt 96/243 (blue), and the other at nt 195/243 (green). B)
Cartoon schematic depicting transfection protocol. Cells were transfected with siRNAs using Lipofectamine 2000 at D0.
RNA was collected at D2, and functional assays were performed from D3-D7. C) Bar graph depicting uc.110
upregulation in banked GBM cell lines. D) Boxplot representing uc.110 expression in pooled glioma adherent cell lines
versus normal human astrocytes. Red boxes indicate an upregulated TUCR. D) Boxplot representing uc.110 expression
in pooled glioma adherent cell lines versus normal human astrocytes. Red boxes indicate an upregulated TUCR. E)
Boxplot representing uc.110 expression in pooled glioma stem cell lines versus normal human astrocytes. Green boxes
indicate a downregulated TUCR. F) Bar graph depicting that the cell invasion phenotype is rescued in A172 and U251
cells with uc.110 overexpression in the presence of siRNAs. Images are representative of the listed sample. si-SCR =
scrambled control siRNA (red), si-uc.110-1 = siRNA targeting uc.110 at nucleotide 96/243 (green), si-uc.110-2 = siRNA
targeting uc.110 at nucleotide 195/243 (blue). G) Cell fractionation bar graph depicting that uc.110 is a predominantly
nuclear RNA molecule, with cytosolic expression in A172s cells. Facets represent cytosolic (red) control genes (GAPDH,
PPIA), nuclear (teal) control genes (U44, U48), and the uc.110 TUCR. $* = p < 0.05$

Supplementary Figure 9. The uc.110 TUCR promotes tumor growth in vivo *A) Cartoon depiction of mouse experiment*
workflow. Cells were transfected with siRNAs using Lipofectamine 2000 at D0 and injected into mice at D2. Tumor
growth was assessed weekly, starting at D16, via MRI.

Supplementary Figure 10

526   _Supplementary Figure 10. PCR confirmation of mutation of miR-544 binding sites for MFRP and uc.110. A) PCR gel_
527   _showing expected products from uc.110 (first site) and MFRP mutations. B) PCR gel showing expected product from_
528   _the second miR-544 binding site in uc.110._

Supplementary Figure 11

Supplementary Figure 11. The uc.110 TUCR activates Wnt-signaling by sponging miR-544 from membrane frizzled
related protein (MFRP) 3'UTR. A) Schematic depicting model for miR-544 sponging by uc.110. B) Schematic depicting
simplified repressed Wnt-signaling pathway. In the normal brain, MFRP is downregulated by miR-544 as depicted in
6A. C) Activation of uc.110 in glioma tumors leads to decreased bioavailability of miR-544. This increases the
bioavailability of MFRP. D) Schematic depicting simplified activated Wnt-signaling pathway. When MFRP bioavailability
is increased by uc.110 activation, as depicted in 6C, Wnt-signaling is also increased.

MATERIALS AND METHODS

Data Availability Statement

RNA-Seq data for Figure 6A will be made available on the Gene Expression Omnibus (GEO) prior
to publication. Detailed TUCR results can be found at www.abounaderlab.org/tucr-database/.
Please refer to the corresponding author for any data access questions.

Detailed Computational Methodologies

Detailed methods, including access to information for all datasets used, can be found in a
repository at: **github.com/abounaderlab/tucr_project.**

TUCR Annotations [29, 30]

TUCR annotations were performed manually by overlaying consensus TUCR genomic annotation
tracks to the hg38 human genome in the UCSC Genome Browser. In parallel, bedtools closest
was used to identify genes that are intergenic or intragenic. These results were then cross
referenced to identify a consensus genomic annotation for each TUCR. Detailed methods can be
found at **github.com/abounaderlab/tucr_project**

TUCR Chromatin Landscaping

U87 H3K4me3, RNA Pol.II, and H3K27ac CHIP-Seq data and U87 ATAC-Seq data were acquired
from the Gene Expression Omnibus. Randomized control TUCRs were generated using Quinlan
Labs' bedtools [31, 32] and the shuffle command.[31, 32] Bedtools fisher and R/RStudio [53, 54]
were used to perform chi-square tests to compare predicted overlaps of peaks to expected peaks.
Detailed methods can be found at **github.com/abounaderlab/tucr_project**

TCGA AND GTEx RNA-Seq Data [33, 34]

GBM (n = 161) and LGG (n = 505) RNA-Seq data were acquired from the Cancer Genome Atlas
and were compared to normal brain cortex from the Genotype-Tissue Expression Database
(GTEx, n = 260) using a workflow including bedtools, bowtie, the SRA toolkit, and R/RStudio.
Detailed methods can be found at **github.com/abounaderlab/tucr_project**

TUCR Expression, Deregulation, and Survival Analyses [33, 34, 54, 55]

TUCR expression, deregulation, and survival analyses, were analyzed using processed TCGA
and GTEx RNA-Seq data and a workflow using R/RStudio. Detailed methods can be found at
github.com/abounaderlab/tucr_project

TUCR weighted gene correlation network analysis (WGCNA) [36]

TUCR WGCNA was performed using processed TCGA and GTEx RNA-Seq data using a modified
version of the R/RStudio workflow designed by Drs. Peter Langfelder and Steve Horvath at UC
Los Angeles. Detailed methods can be found at **github.com/abounaderlab/tucr_project**

De novo transcript reassembly and validation [35]

De novo transcript assembly was performed on TCGA GBM and LGG RNA-Seq data using
standard protocols and the *stringtie* bioinformatics package. Results were validated using PCR:
10 min at 95ºC, followed by 40 cycles of 10 seconds at 95ºC and 1 minute at 60ºC. Detailed
methods can be found at **github.com/abounaderlab/tucr_project.**

Patient Samples

GBM Tumor samples were acquired from the UVA Tumor Bank. Detailed patient information can
be found as a supplement (UVATumorBank_data.csv).

Cell Lines and stem cells

U87, U251, A172, and T98G glioblastoma cell lines were used in *in vitro* experiments and were
acquired from ATCC. U87 cells were cultured in 500 mL minimum essential media (MEM) Earles
(Gibco, #.11095-080) containing 5 mL penicillin/streptomycin (pen/strep, Gibco, Cat #.15140-
133), 5 mL MEM non-essential amino acids (NEAA, Gibco, #.11140-050), 5 mL sodium pyruvate
(Gibco, 100 nM, #.11360-070), 10 mL sodium bicarbonate (Gibco, 7.5%, #.25080-094), and 50
mL fetal bovine serum (FBS). T98G cells were cultured in 500 mL MEM Earles media containing
5 mL pen/strep, 5 mL NEAA, 5 mL sodium pyruvate, and 50 mL FBS. A172 cells were cultured
in 500 mL Dulbecco's modified eagle media (DMEM, Gibco, #.11965-092) containing 5 mL
pen/strep, and 50 mL FBS. U251 cells were cultured in 500 mL RPMI L-Glutamine media (Gibco,

#.11875093) containing 5 mL pen/strep and 25 mL FBS. GSC-34 and GSC-28 glioblastoma stem
cells were cultured in neurobasal (L-glutamine negative) media (Gibco, #.21103-049) containing
5 mL pen/strep, 5 mL B-27 (without Vit-A, Gibco, #.12587-010), 2.5 mL N-2 (Gibco, #.17502-048),
1 mL EGF, 1 mL FGF, and 1.25 mL L-Glutamine. All cell media contained in 5 µL Plasmocure
reagent to prevent mycoplasma contamination.

Primer and Oligo Design

Primers and siRNAs were designed using the Primer3 and Thermofisher design portals,
respectively. uc.110 forward primer sequence is 5'-CAGCCAAAGGGGAAGTGTAT-3', and the
reverse sequence is 5'-CCGTCCTCCCTGCACTAAAT-3'.

MFRP forward primer sequence is 5'- GCATCTATTCATGTGGCAGGC-3', and the reverse
sequence is 5'- TACTCCGGACCCTCCAGTTG-3'.

The miR-544 precursor was ordered from Invitrogen (#.AM17100). Negative control oligos were
ordered from Ambion (#.AM4635).

uc.110 stable overexpression

The full uc.110 transcript from "de novo transcript reassembly and validation" was cloned into the
pCDH-EF1-MCS-BGH-PGK-GFP-T2A-Puro vector (# CD550A-1) using stbl3 competent E. coli
cells and ampicillin selection. Amplified vector was extracted using the miniprep kit (Qiagen,
#.27106). 0.75 µg of this vector, 0.75 µg of psPAX2 lentiviral gag-pol packaging vector, and 0.5
µg of pMD.2G VSV-G enveloping protein was transfected in 6 µL X-tremeGENE transfection
reagent (#.06366236001) into 293T cells per manufacturer instructions to generate a lentivirus
that was transduced to U87, U251, and A172 cells in media without antibiotics. These cells were
subjected to antibody (puromycin) selection for uc.110-positive cells at D3.

uc.110 quantitative (q)PCR

Total RNA was isolated using the RNEasy+ kit (Qiagen, #.74134) according to manufacturer
instructions. RNA concentration and purity was measured via nanodrop. 800 ng of cDNA was
synthesized (BIORAD T100 Thermal Cycler) using the iScript (BIORAD, #. 1708890) synthesis
kit per manufacturer instructions. A 20 µL reaction mixture was then created for each condition
with the following concentrations: 1 µL of combined forward/reverse primers (5 µM), 10 µL of iQ
SYBR Green master mix (#1798880), 4 µL of nuclease free water, and 5 µL of synthesized cDNA.
These reactions were cycled (BIORAD CFX Real Time System) in 96-well plates: 10 min at 95°C,
followed by 40 cycles of 10 seconds at 95°C and 1 minute at 60°C.

Cell Counting (Accumulation) Assay [37-39, 44]

Cells were seeded in 6-well culture plates with full serum media at 30,000/well density at D-1. At
D0, each well was transfected via master mix 3 µL of siRNAs (20 µM) via 9 µL Lipofectamine
2000 (Invitrogen, #.11668-019) in 300 µL OPTI-MEM (Gibco, #.31985-070) and 700 µL antibiotic
and empty media for 6 hours. At 6 hours, media were replaced with fresh media containing
antibiotics and FBS. Cells were then counted via haemocytometer at Days 1, 3, 5, and 7 for each
cell line.

Transwell Cell Invasion/Migration Assay [42-44]

Cells were seeded in 6-well culture plates with full serum media at 300k/well density at D-1. At
D0, each well was transfected via master mix 3 µL of siRNAs (20 µM) via 9 µL Lipofectamine
2000 in 300 µL OPTI-MEM and 700 µL antibiotic and empty media for 6 hours. At 6 hours, the
media were replaced with fresh media containing antibiotics and FBS. The cells were then seeded
in empty media at 200k/chamber into Transwell Invasion Chambers coated with Collagen IV.
After 8 hours, non-invading cells were cleared and invading cells were stained with Crystal Violet.
646

AlamarBlue Cell Viability Assay [40-41]

Cells were seeded in 96-well culture plates with full serum media at 10k/well density at D-1.
Border wells were filled with media to account for edge effects. At D0, each well was transfected
via master mix 1 µL of siRNAs (20 µM) via 3 µL Lipofectamine 2000 in 30 µL OPTI-MEM and 70
µL antibiotic and empty media for 6 hours. At 6 hours, media were replaced with fresh media
containing antibiotics and FBS. Functional assays were performed using the AlamarBlue kit (Life
Technologies #. A50100) per manufacturer instructions. Reactions were allowed to proceed for
1 hour.

Ex vivo knockdown of uc.110

Cells were seeded in 6-well culture plates with full serum media at 300k/well density at D-1. At
D0, each well was transfected with 3 µL of siRNAs (20 µM) via 9 µL Lipofectamine 2000 in 300
µL OPTI-MEM and 700 µL antibiotic and empty media for 6 hours. At 6 hours, the media were
replaced with fresh media containing antibiotics and FBS. Mouse experiments were performed
using xenograft models and intracranial injections of U251 cells post transfection with siRNA

oligonucleotides. Cells were injected at D2 and were imaged at two-week intervals via MRI.
Survival was assessed sured daily and tumor volume was measured at the end of life.
664

RNA-seq post-uc.110 knockdown

666   Cells were seeded in 6-well culture plates with full serum media at 300k/well density at D1. At
667   D0, each well was transfected with 3 µL of siRNAs (20 µM) using 9 µL Lipofectamine 2000 in 300
668   µL OPTI-MEM and 700 µL antibiotic and empty media for 6 hours. At 6 hours, the media were
669   replaced with fresh media containing antibiotics and FBS. RNA Libraries were collected and
670   sequenced via RNA-Seq on Day 2 (post transfection).
671

Luciferase Reporter Vector Construction

The Luciferase reporter vector were constructed via insertion of uc.110 conserved region and
3'UTR of MFRP downstream of Renilla luciferase stop codon in psi-CHECK2 dual luciferase
vectors (Promega, Madison, WI, USA). The insertions were validated by sequencing. Uc.110
and MFRP primer pairs with XhoII and NotI sequence at 5' and 3' respectively, uc.110-FW: 5'-
ATATATctcgagCGAGGTGAGAACCAGAGTGT-3', uc.110-RW: 5'-
AATAATgcggccgcTTGGCTGCCTAATGAGTCACA-3', MFRP-FW: 5'-
ATATATctcgagAAATGGGGTCTGGTCCTTGG-3' and MFRP-RW: 5'-
AATAATgcggccgcTCGCCTTTCTCTCCCGGA-3' were used for PCR amplification.   Site-
directed mutagenesis of predicted miR-544 target sites for both uc.110 and MFRP were
performed to generate mutant vectors.
684

3'UTR Reporter Assays

To determine whether miR-544 directly binds to the MFRP 3'UTR and uc.110, cells were
transfected with miR-544 or miR-scr (control) for 24 hour. The cells were then transfected with
luciferase reporter control or 3'UTR-MFRP or uc.110 as well as corresponsive mutant vectors for
24 hours. Luciferase assays wered performed using the Luciferase System Kit (Promega) and
luminescence was measured. Renilla luciferase activity was double normalized by dividing each
well first by firefly activity and then by average luciferase/firefly value in a parallel set done with
constitutive luciferase plasmid.
694

TCF/LEF reporter Assays

Cells were seeded in 6-well culture plates with full serum media at 300k/well density at D-1. At
D0, each well was transfected with 3 µL of siRNA/miRNA (20 µM) using 9 µL Lipofectamine 2000
in 300 µL OPTI-MEM and 700 µL antibiotic and empty media for 6 hours. At 6 hours, the media

were replaced with fresh media containing antibiotics and FBS. MFRP and uc.110 sequences
were cloned into the PROMEGA pmirGLO Luciferase vector (E1330). BPS Dual reporter
luciferase assays were ordered for TCF/LEF (#.60500) and uc.110/MFRP (#.60683) experiments.

In Vivo Tumor Formation

Adult male and female Nude: Hsd:Athymic Nude-Foxn1 mice were purchased from Harlan. All the
animal work was conducted at the Animal Research Core Facility at the University of Virginia
School of Medicine in accordance with the institutional guidelines. Mice used for this study were
anesthetized with ketamine (17.4 mg/20g), xylazine (2.6 mg/20g) and placed on a sterotactic
frame. Tumor xenografts were generated by implantation of U251 cells transfected with si-uc.110-
1, si-cu.110-2 or si-Scr. U251 cells ($3x10^5$ cells; n=5) were stereotactically implanted into mice in
their right striata at the coordinates from the bregma 1mm anterior, 1.5 mm lateral and 2.5 mm
intraparenchymal. Three weeks after tumor implantation, the animals were subjected to brain MRI.
To measure the tumor size, 20 ul of gadopentetate dimeglumine (Magnevist, Bayer Healthcare)
was intraperitoneally injected 15 minutes before scanning. Tumor volumes were measured using
MicroDicom.

Statistical Analyses

Comparisons between means of samples were performed using Student's t-test and one-way
ANOVAs. Comparisons between categorical variables were performed using chi-squared and
Fisher's exact test. Comparisons were considered statistically significant if the p-value was less
than 0.05. Molecular experiment tests were performed in SigmaPlot 14.0, while computational
experiment tests were performed using bedtools and/or RStudio. Detailed methods can be found
at **github.com/abounaderlab/tucr_project**

AUTHOR CONTRIBUTIONS

Author contributions are defined using Elsevier's CRediT format:

Myron Gibert Jr: Conceptualization, Methodology, Software, Validation, Formal Analysis, Investigation, Writing, Visualization, Supervision, Project Administration, Funding Acquisition

Ying Zhang: Methodology, Investigation, Resources, Formal Analysis, Data Curation, Review and Editing

Shekhar Saha: Methodology, Investigation, Resources, Formal Analysis, Data Curation, Review and Editing

Pawel Marcinkiewicz: Investigation, Resources, Formal Analysis, Data Curation, Review and Editing

Sylwia Bednarek: Methodology, Validation, Formal Analysis, Investigation, Review and Editing, Visualization

Collin Dube: Methodology, Writing, Visualization, Review and Editing

Kadie Hudson: Methodology, Writing, Visualization, Review and Editing

Yunan Sun: Methodology, Writing, Visualization, Review and Editing

Bilhan Chagari: Formal Analysis, Investigation, Writing, Data Curation

Aditya Sarkar: Formal Analysis, Investigation, Writing, Data Curation

Christian Roig-Laboy: Formal Analysis, Investigation, Writing, Data Curation

Natalie Neace: Formal Analysis, Investigation, Writing, Data Curation

Karim Saoud: Formal Analysis, Investigation, Writing, Data Curation

Initha Setiady: Formal Analysis, Investigation, Writing, Data Curation

Farina Hanif: Methodology, Investigation, Review and Editing

David Schiff: Tumor tissue contribution, Review and Editing

Pankaj Kumar: Methodology, Software, Validation, Formal Analysis, Resources, Data Curation

Benjamin Kefas: Methodology, Investigation, Review and Editing

Markus Hafner: Conceptualization and editing

Roger Abounader: Conceptualization, Methodology, Resources, Review and Editing, Supervision, Project Administration, Funding Acquisition

752
ACKNOWLEDGMENTS

This article was supported by NIH UO1 CA220841, NINDS 1R21NS122136-01, NINDS RO1
NS122222, NCI Cancer Center Support Grant 5P30CA044579, and a Schiff Foundation grant (all
to R.A.). The work was also supported by the UVA Cancer Center Bioinformatics Core and the
Molecular Imaging Core. We thank the dbGAP and the TCGA data management teams for
providing access to raw RNAseq data. We would also like to express our gratitude to the patients
for their participation in TCGA. Special thanks goes to Drs. Stephen Turner and Alex Koeppel for
their work with the Bioinformatics Core and early contributions to some of the computational
studies.

REFERENCES

1.   Bejerano, G., et al., *Ultraconserved elements in the human genome.* Science, 2004. **304**(5675): p. 1321-5.

2.   Gibert, M.K., Jr., et al., *Transcribed Ultraconserved Regions in Cancer.* Cells, 2022. **11**(10).

3.   Calin, G.A. and C.M. Croce, *Chronic lymphocytic leukemia: interplay between noncoding RNAs and protein-coding genes.* Blood, 2009. **114**(23): p. 4761-70.

4.   Calin, G.A., et al., *Ultraconserved regions encoding ncRNAs are altered in human leukemias and carcinomas.* Cancer Cell, 2007. **12**(3): p. 215-29.

5.   Colwell, M., et al., *Evolutionary conservation of DNA methylation in CpG sites within ultraconserved noncoding elements.* Epigenetics, 2018. **13**(1): p. 49-60.

6.   Edwards, J.K., et al., *MicroRNAs and ultraconserved genes as diagnostic markers and therapeutic targets in cancer and cardiovascular diseases.* J Cardiovasc Transl Res, 2010. **3**(3): p. 271-9.

7.   Fabris, L. and G.A. Calin, *Understanding the Genomic Ultraconservations: T-UCRs and Cancer.* Int Rev Cell Mol Biol, 2017. **333**: p. 159-172.

8.   Lujambio, A., et al., *CpG island hypermethylation-associated silencing of non-coding RNAs transcribed from ultraconserved regions in human cancer.* Oncogene, 2010. **29**(48): p. 6390-401.

9.   McCole, R.B., et al., *Abnormal dosage of ultraconserved elements is highly disfavored in healthy cells but not cancer cells.* PLoS Genet, 2014. **10**(10): p. e1004646.

10. Scaruffi, P., *The transcribed-ultraconserved regions: a novel class of long noncoding RNAs involved in cancer susceptibility.* ScientificWorldJournal, 2011. **11**: p. 340-52.

11. Zambalde, E.P., et al., *Transcribed Ultraconserved Regions Are Associated with Clinicopathological Features in Breast Cancer.* Biomolecules, 2022. **12**(2).

12. Braicu, C., et al., *The Function of Non-Coding RNAs in Lung Cancer Tumorigenesis.* Cancers (Basel), 2019. **11**(5).

13. Catana, C.S., et al., *Non-coding RNAs, the Trojan horse in two-way communication between tumor and stroma in colorectal and hepatocellular carcinoma.* Oncotarget, 2017. **8**(17): p. 29519-29534.

14. Irimie, A.I., et al., *The Unforeseen Non-Coding RNAs in Head and Neck Cancer.* Genes (Basel), 2018. **9**(3).

15. Ling, H., et al., *Junk DNA and the long non-coding RNA twist in cancer genetics.* Oncogene, 2015. **34**(39): p. 5003-11.

16. Liz, J., et al., *Regulation of pri-miRNA processing by a long noncoding RNA transcribed from an ultraconserved region.* Mol Cell, 2014. **55**(1): p. 138-47.

17. Sakamoto, N., et al., *Non-coding RNAs are promising targets for stem cell-based cancer therapy.* Noncoding RNA Res, 2017. **2**(2): p. 83-87.

800   18.   Kiran, M., et al., *A Prognostic Signature for Lower Grade Gliomas Based on Expression*
801   *of Long Non-Coding RNAs.* Mol Neurobiol, 2019. **56**(7): p. 4786-4798.
802   19.   Dey, B.K., A.C. Mueller, and A. Dutta, *Long non-coding RNAs as emerging regulators of*
803   *differentiation, development, and disease.* Transcription, 2014. **5**(4): p. e944014.
804   20.   Cruickshanks, N., et al., *Role and Therapeutic Targeting of the HGF/MET Pathway in*
805   *Glioblastoma.* Cancers (Basel), 2017. **9**(7).
806   21.   Davis, M.E., *Glioblastoma: Overview of Disease and Treatment.* Clin J Oncol Nurs, 2016.

807   **20**(5 Suppl): p. S2-8.

808   22.   Lathia, J.D., et al., *Cancer stem cells in glioblastoma.* Genes Dev, 2015. **29**(12): p. 1203-

809   17.

810   23.   Kalkan, R., *Glioblastoma Stem Cells as a New Therapeutic Target for Glioblastoma.* Clin
811   Med Insights Oncol, 2015. **9**: p. 95-103.
812   24.   Thakkar, J.P., et al., *Epidemiologic and molecular prognostic review of glioblastoma.*

813   Cancer Epidemiol Biomarkers Prev, 2014. **23**(10): p. 1985-96.

814   25.   Mao, H., et al., *Deregulated signaling pathways in glioblastoma multiforme: molecular*
815   *mechanisms and therapeutic targets.* Cancer Invest, 2012. **30**(1): p. 48-56.
816   26.   Altaner, C., *Glioblastoma and stem cells.* Neoplasma, 2008. **55**(5): p. 369-74.

817   27.   Mourad, P.D., et al., *Why are systemic glioblastoma metastases rare? Systemic and*
818   *cerebral growth of mouse glioblastoma.* Surg Neurol, 2005. **63**(6): p. 511-9; discussion
819   519.
820   28.   Zhang, Y., et al., *The p53 Pathway in Glioblastoma.* Cancers (Basel), 2018. **10**(9).

821   29.   Raney, B.J., et al., *Track data hubs enable visualization of user-defined genome-wide*
822   *annotations on the UCSC Genome Browser.* Bioinformatics, 2014. **30**(7): p. 1003-5.
823   30.   Kent, W.J., et al., *The human genome browser at UCSC.* Genome Res, 2002. **12**(6): p.
824   996-1006.
825   31.   Quinlan, A.R., *BEDTools: The Swiss-Army Tool for Genome Feature Analysis.* Curr Protoc
826   Bioinformatics, 2014. **47**: p. 11 12 1-34.
827   32.   Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing genomic*
828   *features.* Bioinformatics, 2010. **26**(6): p. 841-2.
829   33.   Cancer Genome Atlas Research, N., et al., *The Cancer Genome Atlas Pan-Cancer*
830   *analysis project.* Nat Genet, 2013. **45**(10): p. 1113-20.
831   34.   Stanfill, A.G. and X. Cao, *Enhancing Research Through the Use of the Genotype-Tissue*
832   *Expression (GTEx) Database.* Biol Res Nurs, 2021. **23**(3): p. 533-540.

833   35.   Pertea, M., et al., *CHESS: a new human gene catalog curated from thousands of*
834   *largescale RNA sequencing experiments reveals extensive transcriptional noise.* Genome
835   Biol, 2018. **19**(1): p. 208.
836   36.   Langfelder, P. and S. Horvath, *WGCNA: an R package for weighted correlation network*
837   *analysis.* BMC Bioinformatics, 2008. **9**: p. 559.
838   37.   Hanif, F., et al., *miR-3174 Is a New Tumor Suppressor MicroRNA That Inhibits Several*
839   *Tumor-Promoting Genes in Glioblastoma.* Int J Mol Sci, 2023. **24**(11).
840   38.   Mulcahy, E.Q.X., et al., *MicroRNA 3928 Suppresses Glioblastoma through*

841   *Downregulation of Several Oncogenes and Upregulation of p53.* Int J Mol Sci, 2022. **23**(7).

842   39.   Cruickshanks, N., et al., *Discovery and Therapeutic Exploitation of Mechanisms of*
843   *Resistance to MET Inhibitors in Glioblastoma.* Clin Cancer Res, 2019. **25**(2): p. 663-673.
844   40.   Kumar, P., A. Nagarajan, and P.D. Uchil, *Analysis of Cell Viability by the alamarBlue*
845   *Assay.*

846   Cold Spring Harb Protoc, 2018. **2018**(6).

847   41.   Voytik-Harbin, S.L., et al., *Application and evaluation of the alamarBlue assay for cell*
848   *growth and survival of fibroblasts.* In Vitro Cell Dev Biol Anim, 1998. **34**(3): p. 239-46.
849   42.   Stoellinger, H.M. and A.R. Alexanian, *Modifications to the Transwell Migration/Invasion*
850   *Assay Method That Eases Assay Performance and Improves the Accuracy.* Assay Drug
851   Dev Technol, 2022. **20**(2): p. 75-82.
852   43.   Liu, X. and X. Wu, *Utilizing Matrigel Transwell Invasion Assay to Detect and Enumerate*
853   *Circulating Tumor Cells.* Methods Mol Biol, 2017. **1634**: p. 277-282.
854   44.   Justus, C.R., et al., *In vitro cell migration and invasion assays.* J Vis Exp, 2014(88).

855   45.   Guessous, F., et al., *Cooperation between c-Met and Focal Adhesion Kinase Family*
856   *Members in Medulloblastoma and Implications for Therapy.* Mol Cancer Ther, 2012. **11**(2):
857   p. 288-97.
858   46.   Sonoda, Y., et al., *Akt pathway activation converts anaplastic astrocytoma to glioblastoma*
859   *multiforme in a human astrocyte model of glioma.* Cancer Res, 2001. **61**(18): p. 6674-8.
860   47.   Won, J., et al., *Membrane frizzled-related protein is necessary for the normal development*
861   *and maintenance of photoreceptor outer segments.* Vis Neurosci, 2008. **25**(4): p. 563-74.
862   48.   Katoh, M., *Molecular cloning and characterization of MFRP, a novel gene encoding a*
863   *membrane-type Frizzled-related protein.* Biochem Biophys Res Commun, 2001. **282**(1):
864   p. 116-23.
865   49.   Slaby, O., R. Laga, and O. Sedlacek, *Therapeutic targeting of non-coding RNAs in cancer.*

866   Biochem J, 2017. **474**(24): p. 4219-4251.

867   50.   Saleembhasha, A. and S. Mishra, *Long non-coding RNAs as pan-cancer master gene*
868   *regulators of associated protein-coding genes: a systems biology approach.* PeerJ, 2019.
869   **7**: p. e6388.

870   51.   Li, L., et al., *HOX cluster-embedded antisense long non-coding RNAs in lung cancer.*

871   Cancer Lett, 2019. **450**: p. 14-21.

52. Jiang, D., et al., *Long Chain Non-Coding RNA (lncRNA) HOTAIR Knockdown Increases*
miR-454-3p to Suppress Gastric Cancer Growth by Targeting STAT3/Cyclin D1. Med Sci
Monit, 2019. **25**: p. 1537-1548.

53. Soler, M., et al., *The transcribed ultraconserved region uc.160+ enhances processing and*
A-to-I editing of the miR-376 cluster: hypermethylation improves glioma prognosis. Mol
Oncol, 2022. **16**(3): p. 648-664.

54. Team, R., *RStudio: Integrated Development for R.* 2020.

55. Team., R.D.C. *R: A Language and Environment for Statistical Computing.* 2012; Available
from: http://www.r-project.org/.