1 **An association between *Dnmt1* and *Wnt* in the production of oocytes in the whitefly *Bemisia tabaci***
2

3 **Christopher B. Cunningham[1], Emily A. Shelby[1], Elizabeth C. McKinney[1], Alvin M. Simmons[2], Allen**
4 **J. Moore[1], Patricia J. Moore[1]**
5

6 [1] Department of Entomology, University of Georgia, Athens, Georgia, USA
7

8 [2] U.S. Department of Agriculture, Agricultural Research Service, U.S. Vegetable Laboratory, Charleston,
9 South Carolina, USA
10

11 **Correspondence**
12 Christopher B. Cunningham, Department of Entomology, University of Georgia, Athens, GA, 30646, USA.
13 Email: cbc83@uga.edu
14
15

16 Running title: ***Dnmt1* and *Wnt* in the production of oocytes**
17
18

19
20   **Abstract**
21
The function of methylation in insects and the DNA methyltransferase (*Dnmt*) genes that influence
methylation remains uncertain. We used RNAi to reduce the gene expression of *Dnmt1* within the whitefly
Bemisia tabaci, a hemipteran species that relies on *Dnmt1* for proper gametogenesis. We then used RNA-
seq to test an *a priori* hypothesis that meiosis related genetic pathways would be perturbed. We generally
did not find an overall effect on meiosis related pathways. However, we found that genes in the *Wnt*
pathway, genes associated with the entry into meiosis in vertebrates, were differentially expressed. Our
results are consistent with *Dnmt1* knockdown influencing specific pathways and not causing general
transcriptional response. This is a finding that is also seen with other insect species. We also characterized
the methylome of *B. tabaci* and assessed the influence of *Dnmt1* knockdown on cytosine methylation. This
species has methylome characteristics comparable to other hemipterans regarding overall level, enrichment
within gene bodies, and bimodal distribution of methylated/non-methylated genes. Very little differential
methylation was observed, and difference of methylation were not associated with differences of gene
expression. The effect on *Wnt* presents an interesting new candidate pathway for future studies.

35
36   **KEYWORDS**
37   DNA methylation, epigenetics, gametogenesis, reproduction
38
39

Introduction

5-methylcytosine, commonly referred to as DNA methylation, is a critical epigenetic modification for many plant and animal species, including insects (Gladstad et al., 2014; Schmitz et al., 2019). DNA methylation has been linked to regulation of gene expression, transposon defense, and imprinting in vertebrates and plants (Schmitz et al., 2019). For insects, however, the role of DNA methylation is less clear. Many but not all insects possess DNA methylation and a full complement of the three DNA methylation enzyme, DNA methyltransferase 1-3 (*Dnmt1*, *Dnmt2*, *Dnmt3*; Gladstad et al., 2011; Bewick et al., 2017; Provataris et al., 2018). As with vertebrates, the genomic methylation of cytosines almost exclusively occurs within a CpG context for insects (Gladstad et al., 2016; Schmitz et al., 2019). However, insects diverge from vertebrates in that they have generally lower levels of genomic methylation and it is enriched within gene bodies and not gene promoters or transposons. Differences in cytosine methylation level are not associated with differences in gene expression within a locus across insects, another contrast to vertebrates (Moradin and Brendel, 2021; Dixon and Matz, 2022; Duncan et al., 2022; Malezka and Kucharski, 2022). There is also considerable variation among both the genomic methylation level and the gene copy number of each *Dnmt* among insect species (Gladstad et al., 2011; Bewick et al., 2017; Provataris et al., 2018; Duncan et al., 2022). Some insects have little to no detectable cytosine methylation, *Drosophila melanogaster* and *Tribolium castaneum* being prominent examples (Zemach et al., 2010; Bewick et al., 2017; Provataris et al., 2018). Others have zero annotated copies of DNA methyltransferases, as found in several species of Diptera and Strepsiptera (Gladstad et al., 2011; Bewick et al., 2017; Provataris et al., 2018; Duncan et al., 2022). There isn't a perfect correlation between the presence of *Dnmt* genes and cytosine methylation. *Tribolium castaneum* requires *Dnmt1* function, despite not having detectible cytosine methylation in its genome (Schulz et al. 2018). Despite the general conservation of genomic cytosine methylation and its machinery, no general function of DNA methylation has been established across insects (Dixon and Matz, 2022; Duncan et al., 2022; Malezka and Kucharski, 2022).

While no general function of cytosine methylation has been established, some functions of the DNA methylation enzymes have been characterized for several species, particularly involving *DNA methyltransferase 1*. As with vertebrates *Dnmt1* is responsible for maintaining methylation patterns after a cell division (Law & Jacobsen, 2010; Gladstad et al., 2016; Schmitz et al., 2019). Reduction of *Dnmt1* expression reduces genomic cytosine methylation level following mitosis, as expected (Bewick et al., 2019; Amukamara et al., 2020; Ventos-Alfonso et al., 2020; Washington et al., 2021; Ivasyk et al. 2023). In addition to its canonical function, several studies have established that *Dnmt1* has additional functions for insects. When *Dnmt1* expression is reduced, females show reduced egg production, males show reduced sperm production, and embryos show reduced viability after oviposition for several insect species (*Bemisia tabaci*, Shelby et al., 2023; *Blattella germanica*, Ventos-Alfonso et al., 2020; *Bombyx mori*, Li et al., 2019; *Harmonia axyridis,* Gegner et al., 2020; *Oncopeltus fasciatus*, Bewick et al., 2019, Amukamara et al., 2020, Washington et al., 2021, Cunningham et al., 2023; *Ooceraea biroi*, Ivasyk et al. 2023; *Nasonia vitripennis*, Zwier et al., 2012, Arsala et al., 2022; *Nilaparvata lugens*, Zhang et al., 2015; *Tribolium castaneum*, Schulz et al., 2018); an effect also observed for mammals (*Mus musculus*, Takada et al., 2021). Somewhat unexpectedly, this effect appears to be specific to gametes as the gonads themselves do not show any gross signs of apoptosis (Amukamara et al., 2020, Washington et al., 2021; Shelby et al., 2023; Ivasyk et al. 2023). Lifespan is generally not reduced despite a reduction of *Dnmt1* expression (Amukamara et al., 2020; Shelby et al., 2023; Ivasyk et al. 2023). Gamete-specific effects are even present for *Tribolium castaneum*, a species that has functioning *Dnmt1* but no cytosine methylation (Schulz et al., 2018). These phenotypes

and the apparent cell type-specific effects lead us to further investigate the genetic mechanisms that
underpin the deficits, focusing on potential genetic networks.
In this study we used RNA interference (RNAi) to test the function of *Dnmt1* of the whitefly
Bemisia tabaci (Gennadius). *Bemisia tabaci* is a global pest species (Stanley & Naranjo, 2012). As such,
work has been done to understand its reproduction and the molecular pathways that underpin its
reproduction with an eye towards more efficient control of this pest (Upadhyay et al., 2016; Luo et al.,
2017; Grover et al., 2018; Shelby et al., 2020; Hu et al., 2022; Shelby et al., 2023). As with other insect
species, reduction of *Dnmt1* expression of this species leads to greatly reduce oocyte production and embryo
inviability (Shelby et al., 2023). From previous studies, gametogenesis at the mitosis-meiosis transition is
arrested (Amukamara et al., 2020; Washington et al., 2021; Cunningham et al., 2023; Ivasyk et al. 2023).
This led us to hypothesize that reducing *Dnmt1* expression targets specific gene networks. To that end our
first aim was to test if reducing *Dnmt1* expression preferentially perturbs meiosis-related pathways, but not
mitotic or general cell cycling pathways. We tested this in two ways. First, we tested a panel of *a priori*
candidate genes canonically associated with each process to contrast; general cell proliferation, cell cycling,
mitosis, and meiosis. Second, we tested a panel *a priori* Gene Ontology Terms representing these processes
and assessed if they were enriched among the most differentially expressed genes. Finally, we assessed the
general patterns of differential gene expression to explore the dataset for perturbed pathways. Our analysis
showed that reducing *Dnmt1* expression does not cause a general transcriptional response but uncovered
Wnt as a novel candidate pathway. Our second aim for this study was to characterize the general patterns
of genomic cytosine methylation of this species, the effect of *Dnmt1* reduction on cytosine methylation,
and the association between cytosine methylation and gene expression. We found that *B. tabaci* has cytosine
methylation at a level comparable to other hemiptera. However, reducing *Dnmt1* expression did not cause
pervasive reduction of cytosine methylation, but cell type-specific effects remain to be assessed for future
studies. The differences of cytosine methylation that are detectable are not associated with difference of
gene expression. Future experiments assessing more tissue-specific level expression is now needed to
determine if and how *Dnmt1* interacts with meiotic pathways.
110
111
112 **Results**
113
114 ***Dnmt1* RNAi knockdown**
115
116 Consistent with Shelby et al. (2023), our fed RNAi treatment reduced the expression of *B. tabaci*'s single
117 copy of *Dnmt1* within adult females (LOC109040312; $\log_2$ fold change: -0.0216, P = 0.0276). This is the
118 same magnitude of knockdown that was seen with Shelby et al. (2023) when we first characterized the
119 reproductive deficits that *Dnmt1* RNAi knockdown produced.
120
121 ***a priori* candidate gene screen**
122
123 Our hypothesis that meiotic-specific candidate genes would be preferentially perturbed was not strongly
124 supported; none of the 5 meiosis-specific candidate genes were differentially expressed (Table 1). Four of
125 the 22 candidate genes were statistically differentially expressed after FDR thresholding (Table 1). These
126 genes were *Wnt*-like genes, along with one *CDC* and one *SMC3*. All gene assessed both here and by Shelby
127 et al. (2023) had the same directional change and rank order of effect sizes.

Table 1. *a priori* candidate genes tested for differences between Control and *Dnmt1* knockdown treatments. Expression value and fold changes are from DESeq2. Bolded values are statisitcally significant after adaptive linear two-step procedure (Benjamini et al., 2006). Gene orthology and nomeclature is from OrthoDB.

Gene ID	Accession	Pathway	Standardized Expression	Fold Change	P value	*a priori* Category
Wingless (*Wnt* ortholog)	LOC109044616	Cell fate & proliferation	7.641	-0.001	0.463	General Cell Proliferation: 2/9
Wnt-1-like	LOC109042201	Cell fate & proliferation	10.265	-0.005	0.560	
Wnt-2	LOC109041803	Cell fate & proliferation	8.729	0.001	0.512	
Wnt-4-like	LOC109044333	Cell fate & proliferation	**10.343**	**0.012**	**0.009**	
Wnt-6-like	LOC109044640	Cell fate & proliferation	**8.916**	**0.005**	**0.015**	
Wnt-10a-like	LOC109044612	Cell fate & proliferation	7.642	0.001	0.424	
Frizzled-2-like	LOC109035614	Cell fate & proliferation	8.536	-0.004	0.222	
Frizzled-4-like	LOC109030198	Cell fate & proliferation	9.002	-0.004	0.222	
Frizzled-5-like	LOC109039814	Cell fate & proliferation	12.458	-0.001	0.933	
CDC20	LOC109035963	Cell cycle control	**12.465**	**-0.012**	**0.034**	Cell Cycle: 1/3
CYCD2	LOC109039503	Cell cycle control - G1/S-specific	10.915	0.005	0.533	
CYCD3	LOC109034725	Cell cycle control - G1/S-specific	8.745	-0.001	0.594	
Vasa	LOC109042479	Germ cell Development	15.291	0.001	0.856	Mitosis Specific: 1/4
Vasa	LOC109030586	Germ cell Development	15.052	-0.009	0.172	
CYCB3	LOC109043986	Cell cycle control - G2/mitotic-specific	12.196	-0.003	0.494	
SMC3-like	LOC109038740	Chromosomal Structure Maintenance	**11.516**	**-0.025**	**0.050**	
Pumillio	LOC109041486	Repressor of Mitosis	13.453	0.004	0.425	Meiosis Specific: 0/9
Decapentaplegic	LOC109031148	Repressor of Meiosis	8.575	0.001	0.787	
Decapentaplegic-like	LOC109032526	Repressor of Meiosis	7.903	-0.002	0.393	
Glass bottom boat	LOC109036141	Repressor of Meiosis	11.289	0.000	0.968	
Boule	LOC109034894	meiotic G2/M transition	11.156	0.002	0.233	
SPO11	LOC109044039	Meiotic Recombination	7.689	-0.001	0.545	
MSH5	LOC109042904	Meiotic Recombination	8.962	-0.002	0.618	
MND1	LOC109040052	Meiotic Recombination	9.436	-0.001	0.857	
HOP2	LOC109030819	Meiotic Recombination	9.824	0.001	0.856	

129
130

131 ***a priori* GO term enrichment**

133 Our hypothesis that the meiosis pathway would be the most affected by *Dnmt1* knockdown was supported in a minority of our *a priori* GO term
134 enrichment analysis. In particular, the meiosis GO Term was not enriched. However, five of the six *a priori* GO terms were statistically significantly.

135 These were among the genes that showed the least differences of expression between the control and RNAi-treatments, the opposite pattern predicted
136 (Table 2).
137

Table 2. *a priori* GO term enrichment. Loss of *Dnmt1* does not preferentially target our predcited pathways. NES = normalized enrichment score.

Pathway	GO Term	Number of Annotated Genes	NES	FDR-adjusted P-value
Cell Cycle	GO:0007049	384	-4.742	**< 0.001**
Gametogenesis	GO:0007276	681	-4.260	**< 0.001**
Mitotic Cell Cycle	GO:0000278	255	-4.315	**< 0.001**
Meiotic Cell Cycle	GO:0051321	144	-3.385	**< 0.001**
Oogenesis	GO:0048477	491	-3.644	**< 0.001**
Apoptotic Process	GO:0006915	61	0.676	8.605E-01

138
139
140

Differential gene expression

We assessed the impact of *Dnmt1* knockdown to identify genes that were differentially expressed (11 biological replicates of *Dnmt1* and 12 of *eGFP*). The individual samples of each treatment clustered together, but the treatments were differentiated using a PCA and not statistically significantly different as the 95% confidence intervals were highly overlapping (Fig. 1). There were 96 genes (49 up-regulated, 47 down-regulated) between the treatments.

Fig 1. There was little differentiation of samples reflecting how few genes were differentially expressed between treatments. The percent of variation capture by each axis is in parenthesis. Smaller points represent individual samples and are colored by their treatment category (n = 12 for *eGFP*, 11 for *Dnmt1*). Large points represent the centroid (i.e., mean) of each treatment. Circles represent 95% confidence intervals.

GO term overrepresentation

The differentially expressed genes between *Dnmt1* knockdown and controls produce 87 overrepresented GO terms. The top three Biological Process terms are female meiosis I – GO:0007144, sperm axoneme

assembly – GO:0007288, and negative regulation of imaginal disc growth – GO:0045571 (P = 0.00021. P=
0.00044, P = 0.00075, respectively). Although meiosis is a top GO term, the overall list does not contain
other terms related to meiosis. Specifically, the list contains only one meiosis specific term, but five terms
specific to mitosis and five terms specific to general cell development (Supplementary File 3). The top three
Molecular Function terms with more than one annotated gene are mRNA 3'-UTR binding - GO:000373,
ribonucleoside-diphosphate reductase activity - GO:000474, and coenzyme A transmembrane transporter
activity - GO:0015228 (all P = 0.0101). The top three Cellular Compartments terms with more than one
annotated gene are MKS complex - GO:0036038, mitotic spindle pole - GO:0097431, and mitochondrial
respiratory chain complex IV - GO:0005751 (P = 0.0108, P = 0.0162, P = 0.0162, respectively). The
remainder of the terms can be found in Supplementary File 3.
171
172   **Characterization of cytosine methylation**
173
We assessed genomic distribution and abundance of cytosine methylation to try to understand its potential
function for *B. tabaci*. Cytosine methylation is found almost exclusively within a CpG context, which is
consistent with other hemipterans and insects in general. Cytosines are methylated symmetrically (i.e., both
strands of a CpG are methylated or not; Fig. 2), which is an indication that *Dnmt1* functions as expected.
GC content of genes is higher than the background intragenic genome. This is expected because *B. tabaci*
is an AT biased genome overall (genome: 39% GC, genes: 44% GC; Fig. 3A). Methylated cytosines within
genic regions mirror the GC content of genes (Fig. 3B). Genes have a very bimodal distribution of
methylation status (i.e., genes are either highly methylated or not; Fig. 3C). Transposons did not have
consistent or high levels of methylation (Fig. 3D). Methylated cytosines are enriched within genic and
exonic regions but are seen at genomic levels within introns (Fig. 4).

184
185
186 **Fig 2. Cytosine methylation of *Bemisia tabaci* is highly symmetric, consistent with many other insects.**
187 Density plot of CpG's methylation within different contexts. Lighter colors represent denser value regions.
188

189
190
Fig 3. Cytosine methylation of *Bemisia tabaci* is mostly consistent with many observed trends of other
insects. (A) The GC content increases over gene bodies. CDS = coding sequence, TSS = transcriptional
start site, TES = transcriptional end site. (B) CpG methylation is mostly found in gene bodies. (C) Genes
are either highly methylated or non-methylated. This is a more bimodal distribution than is seen for other
insects. (D) There is little cytosine methylation over the most frequent transposon within the *B. tabaci*
genome. Each colored line represents the four most populous families of transposons within the *B. tabaci*
genome (orange = BEL, green = Gypsy, blue = hAT, navy = Mariner).
198

199
200
201 **Fig 4. Cytosine methylation of *Bemisia tabaci* is not detectably influenced by knockdown of *Dnmt1*.**
202 Cytosine methylation is enriched with genic and exonic regions but is not detectably influenced by *Dnmt1*
203 RNAi. Each point represents the mean ± SEM of each treatment within the labeled category.
204
205 **Cytosine methylation is associated with overall gene expression (between loci differences)**
206
207 High levels of gene methylation are associated with high levels and low variances of gene expression (Fig.
208 5). This is a well-established pattern that suggests if cytosine methylation is performing a function, it is
209 likely similar between *B. tabaci* and other insects.
210
211

212
213
Fig 5. Gene body methylation is weakly, positively associated with overall gene expression levels and
negatively associated with gene expression variance. (A) Plot of mean standardized gene expression
values against decile of the percent of CpG's methylated within a gene body. (B) Plot of gene-wise standard
deviation against the decile of the percent of CpG's methylated within a gene body. Both panels have values
that represent the mean ± SEM.

Differential gene methylation

After assessing the general trends of cytosine methylation within *B. tabaci*, we assessed the effect of *Dnmt1*
knockdown on gene methylation level between *Dnmt1* and *eGFP* samples (five biological replicates of each
treatment). Very few gene loci differed at all between the treatments; only 2,391 differed by 1% or more
(Fig. 6). No gene was statistically significantly different after FDR correction.

226
227
Fig 6. Very few genes differ for methylation status between treatments. The inner and outer diagonal
lines represent 5% and 20% differentiation of *Dnmt1* vs *eGFP* treatments, respectively. Each circle
represents an individual gene.

Differential gene methylation is not associated with differential gene expression (within locus differences)

Lastly, we assessed if the difference of gene methylation that were present influenced gene expression
within (and not between as above) a gene. There was no indication that difference of methylation influenced
difference of gene expression (Fig. 7). This is consistent with almost all other insects.

239
240
Fig 7. Differential methylation is not associated with differential gene expression. Scatterplot of
differential expression against differential methylation. The pattern would approximate a 1:1 relationship
if cytosine methylation directly influenced gene expression. Each circle represents an individual gene.

Discussion

DNA methylation and its machinery is present within many insect species but there is variation among
species for both genomic level and copy number of the genes influencing DNA methylation, *Dnmt1* and
Dnmt3 (Bewick et al. 2017). While DNA methylation itself does not have a universal function among
insects, one of the genes, *Dnmt1*, consistently perturbs gametogenesis when its expression is reduced (Zwier
et al., 2012; Zhang et al., 2015; Schulz et al., 2018; Bewick et al., 2019; Li et al. 2019; Amukamara et al.,
2020; Gegner et al., 2020; Ventós-Alfonso et al., 2020; Takada et al., 2021; Washington et al., 2021; Arsala
et al., 2022; Xu et al. 2022; Cunningham et al., 2023; Ivasky et al., 2023; Shelby et al. 2023). Here, we
investigated the genetic underpinning of the severe reproductive deficits after *Dnmt1* expression was
reduced using RNAi fed to adult female *B. tabaci*. This treatment, at the reduced expression level observed
here, causes females to produce both fewer and less viable eggs (Shelby et al., 2023). We used RNA-seq
of whole-body samples to specifically test an *a priori* hypothesis that this treatment would impact meiotic
pathways preferentially, the developmental point at which many gametes appear to arrest (Amukamara et
al., 202; Washington et al., 2021; Cunningham et al., 2023). We also explored this dataset to examine if

other genetic pathways associated with germ cell maturation were impacted. Relatively few genes overall
were differentially expressed. This work could have detected strong effects, we believe the lack of pervasive
differential gene expression given the severe reproductive deficit phenotype is likely due to using whole
bodies. While our hypothesis of meiosis-specific pathways was not supported, we did find differential
expression of *Wnt*-like genes. From this, we think that *Wnt* and its pathways become an interesting candidate
pathway for additional study.
We also performed a baseline characterization of genomic cytosine methylation of *B. tabaci* and
the impact of *Dnmt1* knockdown on the level of methylation. Cytosine methylation was found at
comparable level to other hemipteran, within a CpG context, and was enriched within genic regions across
the genome. Cytosine methylation differences between genes were positively associated with gene
expression level. However, differences of cytosine methylation level due to treatment within the same gene
were not associated with gene expression differences across all genes. No statistically significant
differences in gene body methylation were detected. This is likely attributable to overall few cellular
divisions happening at the whole-body level where *Dnmt1*'s reduced expression is likely to manifest its
effect. Moving forward, there is a need to take a tissue specific approach to understanding the genetics of
this phenotype. Our analysis also provides a new candidate pathway for the study of reproductive deficits
of female insects and provide a resource for studies of genomic methylation of *B. tabaci*.
Despite the dramatic reproductive phenotype, we did not find large numbers of differentially
expressed genes, a lack also observed with other studies (Bewick et al., 2019; Cunningham et al., 2023).
This suggests that *Dnmt1* knockdown is affecting specific genes or gene networks, rather than producing
the reproductive deficit by instigating a global transcriptional response. Our *a priori* hypothesis that meiotic
specific pathways would be more affected compared with mitosis or general cell cycling pathways did not
receive strong support overall. None of our meiosis candidate genes were differentially expressed compared
with one or two in the other categories – general cell proliferation, cell cycle, and mitosis. Genes that were
annotated with the GO term of meiosis were not enriched among the genes ranked highest by expression
differences, regardless of whether they differentially expressed or not. Among only the differentially
expressed genes themselves (96 genes), one meiosis GO term was overrepresented. However, the list also
contained several mitosis specific and development specific GO terms. We believe this general
overrepresentation of development genes is an overall reflection of assaying a perturbed reproductive tissue
and not specific of any pathway being preferentially affected. Thus, we did not find support for *Dnmt1*
being directly involved in the control of meiosis with this study.
However, we found intriguing evidence that a conserved genetic pathway, the *Wnt* pathway, might
be affected by *Dnmt1* knockdown. *Wnt*'s of insects are usually viewed as general development genes;
involved with cell proliferation, migration, patterning, and identity (Murat et al., 2010; Waghmare and
Page-McCaw, 2018). In human colon cancer cells, *Dnmt1* influences the *Wnt* pathway via its members *β-*
catenin and *Snail1* through a DNA methylation-independent mechanism (Espada et al., 2011). While
mammalian DNA methylation operates differently than it does for insects, these results generate specific,
testable hypotheses about how the expression of *Dnmt1* is required for proper regulation of gametogenesis.
The regulation of the *Wnt* pathway by *Dnmt1* being methylation independent also aligns with much of the
evidence produced by insect systems (Bewick et al., 2019; Amukamara et al., 2020; Ventós-Alfonso et al.,
2020; Washington et al., 2021; Arsala et al., 2022; Cunningham et al., 2023; Ivasky et al., 2023; Shelby et
al., 2023). This pathway has also been shown to be involved in entry into meiosis of mice (Chassot et al.,
2011, Le Rolle et al., 2021). Our future experiments will explore the interaction between the *Wnt/β-catenin*
and *Dnmt1* in *B. tabaci*.

One possible explanation for detecting few differential expressed genes is that we sampled not only
the tissue in which *Dnmt1* knockdown has its observed effect, the ovaries, but whole bodies. However,
Dnmt1 expression was reduced, and the magnitude of reduction observed here has been shown to reduce
the presence its protein, DNMT1, within the nucleus of ovarian cells (Shelby et al., 2023). The observed
amount of differentially expressed genes is reasonable from whole bodies of this species when the
experimental effects are regional or tissue specific (Tadmor et al., 2022). A second possible explanation is
that we did not detect high levels of differentially expressed genes because we did not appreciably reduce
DNA cytosine methylation. The reduction in DNA methylation with loss of *Dnmt1* requires cell division
and DNA synthesis. High turnover tissues exist within adult insects, such as digestive and reproductive
tissues. However, they would be the minority of the cell pollution sampled here. Thus, our bisulfite
sequencing data might not detect differential DNA methylation within the small proportion of cells that
divided between treatment and sampling (24 hours). However, given the consistent result that changes of
cytosine methylation pattern are not associated with differential expression in *O. fasciatus* gonads (Bewick
et al., 2019, Cunningham et al., 2023), our data support the hypothesis that *Dnmt1* knockdown is affecting
targeted genetic pathways in the gonads to specifically affect gametogenesis. This adds to the evidence that
Dnmt1 or lack thereof is acting through a cytosine methylation independent pathway. The gametes/eggs
that were affected were post-meiotic and such cells would have similar cytosine methylation patterns to
control samples; however, they lacked nuclear DNMT1 (Shelby et al., 2023). Recently, an ancestrally
duplicated domain of DNMT3 of honey bees showed binding to histone residues, a previously
uncharacterized interaction within insects (Kucharski et al., 2023). Reanalysis of the alternative splicing of
this gene during development also showed many isoforms produced with the duplicated domain and without
the catalytic methyltransferase domain (Kucharski et al., 2023). This supports the general idea that these
Dnmt gene families are more diverse than previously appreciated and that they have the capacity to act via
different pathways. This would suggest that *Dnmt1* knockdown effects are not likely being mediated
through difference of cytosine methylation.
Our final aim of the study was to characterize the cytosine methylation of *B. tabaci*. This was both
a general characterization and how *Dnmt1* knockdown affected the patterns. Cytosine methylation was
found within a CpG context and enriched within genic regions. Genes have a bimodal distribution whereby
gene are either highly methylated or not and methylation level is associated with gene expression level
between loci. There was a positive association between methylation level and absolute level of gene
expression and a negative association with gene expression variability. All of these patterns are common
among insects and suggest that cytosine methylation is comparable among hempiterans (Bewick et al.,
2017; Provataris et al., 2018). There was little cytosine methylation found across transposons, which
suggests that this nucleotide modification is not serving as a defense against those elements (Bewick et al.,
2019; Gladstad et al., 2014). The RNAi *Dnmt1* treatment produced no statistically significant differentially
methylated genes. We believe the same explanation for few differentially expressed genes applies to this
result as well – whole body samples were not sensitive enough to detect changes of gene methylation within
a minority of cells within our samples. Although methylation level between treatments were not statistically
significantly different, some loci did show differences; these differences were not associated with gene
expression levels. This also makes *B. tabaci* comparable to other insects where an association between
difference of gene methylation and expression is rare (Moradin and Brendel, 2021; Dixon and Matz, 2022;
Duncan et al., 2022; Malezka and Kucharski, 2022). Overall, *B. tabaci* shows many of the conserved
patterns of cytosine methylation of insects but a more tissue specific approach is warranted before making
definitive conclusions.

Pest management of *B. tabaci* is one of the primary reasons for any work with this species. Our results here are add to the understanding of its reproduction and will inform experimental design moving forward. Clearly, *Dnmt1* knockdown produces severe reproductive deficit for female *B. tabaci* (Shelby et al., 2023). We also now know that *B. tabaci* has cytosine methylation patterns and association like other hemipteran and insects meaning we can use those species to help inform hypotheses, predictions, and conclusions. The work here is an initial attempt to fill in the mechanistic understanding of how *Dnmt1* might work. This current work also suggests that we need to focus on disrupted tissues themselves, which should facilitate a more complete understanding of the pathways and mechanisms by which *Dnmt1* influences insect reproduction.

Experimental Procedures

To understand the functional role that *Dnmt1* plays during gamete formation of female *B. tabaci*, we used RNAi to knockdown *Dnmt1* gene expression replicating the protocol of Shelby and colleagues (2023). After the treatment, we tested two sets of *a priori* predictions after gene expression quantification using RNA-seq; one using candidate genes and the other using Gene Ontology terms. The same RNA-seq data were also used to perform exploratory analyses to assess how *Dnmt1* knockdown perturbed the transcriptome of females. We also assessed cytosine methylation of both control and RNAi-treated females. We first characterized the level *B. tabaci* has genomic methylation, in what genomic context it is found, how gene methylation related to gene expression globally. We then assessed if *Dnmt1* expression reduction caused differential gene methylation and if differential gene methylation was associated with differential gene expression.

Animal colony and husbandry

This experiment used the same laboratory population and husbandry protocols as Shelby and colleagues (2023). Briefly, these *B. tabaci* colonies were started from population collected from a cotton (*Gossypium hirsutum*) field site in Tift County, Georgia, USA in 2018 (McKenzie et al., 2020) belonging to the MEAM1 *B. tabaci*. The whiteflies were maintained on collard (*Brassica oleracea*) plants using individual pots and grown to a minimum height of 15 cm. Both whiteflies and collards were maintained at 26°C with a 14:10 hour light:dark photoperiod in 51 cm × 28 cm × 28 cm aluminum insect cages (Bioquip Products, Rancho Dominguez, CA, USA) within a Percival Incubator (Percival Scientific, Perry, Iowa, USA) during the entire experiment.

RNA interference (RNAi) synthesis, administration, and quality control

We created RNAi constructs targeting *Dnmt1* following Shelby and colleagues (20123). Briefly, RNAi constructs targeting *Dnmt1* were prepared via PCR using 500 ng of RNA from unmated females. The construct was targeted towards the Replication Fork Domain (RFD) motif the *Dnmt1* gene. Afterwards, double stranded RNA was synthesized with an Ambion MEGAscript kit (ThermoFisher Scientific, Waltham, MA, USA) following the manufacturer's protocol. This reaction was purified with a phenol:chloroform:IAA extraction followed by a sodium acetate precipitation. We measured concentration using Invitrogen's Qubit system with the ssRNA kit following the manufacturer's protocol. We used *eGFP* as an exogenous control construct.

Newly-eclosed females were removed from colonies within 24 hours post-eclosion to drastically
reduce the opportunity to mate. Thereafter, adult female at 3-6 days post-eclosion were fed 200 µl of RNAi
solution of either *Dnmt1* or *eGFP* each at a concentration of 0.05 µg/mL within a 10% w/v sucrose solution
contained within a cap of a 1.7 mL microcentrifuge tube using an artificial feeding apparatus made from
microcentrifuge tube caps. Females were haphazardly allocated to treatment group. The RNAi solution was
colored with 5 µl of green food coloring. Parafilm was placed over the cap to provide a membrane for
feeding. 75–100 female whiteflies were placed into each prepared acetate tube, which was then wrapped
with black construction paper to promote movement to the top of the tube. Females were allowed to feed
for 24 hours, and consumption was confirmed by observing a green abdomen before collection. *eGFP*
control and *Dnmt1* females were collected after this feeding period and immediately flash frozen with liquid
nitrogen and stored at − 80°C until nucleotide extraction. Twenty-five females were pooled into each
biological replicate.

This treatment does not impact lifespan between control and *Dnmt1* treated females or gross
morphological structures (Shelby et al., 2023). This treatment statistically significantly reduces the
expression of *Dnmt1* of treated females (Shelby et al., 2023).

407
408   **RNA & DNA extraction**
409
410   Total RNA and genomic DNA was extracted using a Qiagen Allprep DNA/RNA Mini Kit (Qiagen, Venlo,
411   The Netherlands) following the manufacturer's protocol. Homogenization of the females was performed
412   using a Bead Bug (Benchmark Scientific, Sayreville, New Jersey, USA) at 4000 rpm for 60 seconds.
413   Quantification of DNA and RNA was done with a Qubit fluorometer using the DNA HS and RNA BR kits,
414   respectively, following the manufacturers protocol.

415
416   **RNA-seq High-throughput library preparation and sequencing**
417
418   Twelve biological replicates each of *eGFP* and *Dnmt1* were used for the RNA-seq experiment. A sample
419   of 150 ng of extracted total RNA of *B. tabaci* was used to construct poly-A selected, non-stranded Illumina
420   compatible libraries by Novogene Corporation (Sacramento, CA, USA). We targeted 20M 2 x 150 bp read
421   pairs per biological replicate using an Illumina NovaSeq by Novogene.

422
423   **RNA-seq quality control and mapping**
424
Reads were initially assessed for quality with fastQC (v0.11.9; default settings; Andrews, 2010). Reads had
adapters trimmed with cutadapt (v2.8; --trim-n -O 3 -u -2 -U -2 -q 10,10 -m 30; Martin, 2011) using the
TruSeq adapter sequences. Afterward, reads were reassessed with fastQC with default settings. Overlapping
reads were merged with FLASh (v 1.2.11; default settings; Magoc and Salzberg, 2011). As a final QC step,
reads were mapped to rRNA genes and were removed with SortMeRNA (v4.3.3; s rRNA: NC_006279, l
rRNA: NC_006279.1; Kopylova et al. 2012). We used the NCBI RefSeq assembly (ASM185493v1; Chen
et al., 2016) and annotation (GCF_001854935.1). HISAT2 (v2.2.1; no soft-clipping; Kim et al. 2019) was
used to map reads to the genome. Mappings were converted to read counts by StringTie (v2.1.7; Pertea et
al. 2015) following the manual instructions for export to DESeq2.

434
Functional annotation

We added functional annotation to the *B. tabaci* proteome/transcriptome using eggNOG-mapper webserver
using the Auto taxonomic scope setting (http://eggnog-mapper.embl.de; v 2.19; Cantalapiedra et al. 2021).
This annotated 16,813 of 22,737 protein/transcript models (74.1%) with Gene Ontology terms, which was
the database to perform our GO term enrichment analysis and GO term overrepresentation analysis
(Supplementary File 1).

a priori **candidate gene screen**

We tested our hypothesis that *Dnmt1* knockdown would preferentially influence meiotic, not mitotic, or
general cell cycle, pathways by assessing a series of candidate genes associated with each pathway. These
genes were chosen as canonical members based on literature searches (Cunningham et al., 2023). These
include cell fate and proliferation genes that we did not expect to be differentially expressed (members of
the *Wnt*, *Frizzled*, and *Pumillio* families), cell cycle control genes (some of which we did expect to be
differentially expressed (*Vasa*) and some of which we did not expect to be differentially expressed (*Cyclin*
B3, *Cyclin D2*, *Cyclin D3*, *Cell Division Cycle 20*, *Cell Division Cycle 25* family members)), maintenance
of chromosome genes that we did expect to be differentially expressed (*Structural Maintenance of*
Chromosome 3 family members), meiotic transition gene that we did expect to be differentially expressed
(*Boule, Decapentaplegic*, and *Glass boat bottom* families), and meiotic recombination genes that we
expected to be differentially expressed (*SPO11 Initiator of Meiotic Double Stranded Breaks*, *MutS*
Homolog 5, *Meiotic Nuclear Divisions 1*, and *Homologous-Pairing Protein 2* families). We did not predict
specific directional changes, except decreased expression of *Vasa* and *Boule* within the *Dnmt1* knockdowns.
We extracted raw P values for differential gene expression from DESeq2 and compared them with Adaptive
Two-Step B-H Thresholding test, which is less conservative than standard Benjamini-Hockberg correction
(Benjamini et al., 2006).

a priori **Gene Ontology (GO) term enrichment analysis**

We also tested our hypothesis using a very broad test of pathway enrichment. We selected five high-level
GO terms representing the genetic pathways that we predicted would or would not be after *Dnmt1*
knockdown. These were GO:0007049 Cell Cycle, GO:0000278 Mitotic Cell Cycle, GO:0051321 Meiotic
Cell Cycle, GO:0007276 Gametogenesis, and GO:0007283 Spermatogenesis. With our previous observed
phenotypes, we expect Meiotic Cell Cycle, Gametogenesis, and Spermatogenesis to be enriched. We
expected Cell Cycle and Mitotic Cell Cycle to show no enrichment. We included GO:0070265 Necrotic
Cell Death and GO:0006915 Apoptotic Process to assess if our samples were undergoing these processes.
We used the S values from DESeq2 to rank order genes. We used fgsea R package (Korotkevich et al. 2021)
with default parameters to test if genes annotated with these GO terms were enriched at the head of these
lists. The same R package was used to visualize the results.

Differential gene expression

Read counts were imported into R using tximport (Bioconductor v1.20.0; Soneson et al. 2015) following
the manual's instructions. We used R (v4.1.0; R Core Team, 2021) within an Rstudio IDE (build 492;
Rstudio Team, 2022) for the analysis.

DESeq2 (Bioconductor v1.32.0; default settings; Love et al. 2014) was used for the differential
gene expression analyze following the manuals suggestions for exploratory data analysis and
sample/analysis quality control. *eGFP* was set and used as the comparison group. We checked the model
matrix specification to ensure correct specification (i.e., that program was contrasting the samples in the
correct way). After importing and initial analysis, samples were plotted with a PCA to visually check for
outliers according to the manual's recommendation. After inspection, we removed one *Dnmt1* sample that
was both an intra-group PCA outlier and having very high expression of *Dnmt1* that suggested the RNAi
had not worked properly. This left 12 biological replicates of *eGFP* and 11 of *Dnmt1* for the final
differential gene expression analysis.

After removal of the outliers the analysis was repeated using the same settings of DESeq2 and the
results were again quality controlled. We used the default dispersion estimator and shrinkage method,
apeglm (Zhu et al. 2019). We used s-values to estimate statistical significance after false discovery rate
correction at the level of 0.05 (Stephens 2016). Results were visualized with the fviz_pca_ind function of
the factoextra R package (Kassambara and Mundt 2020).

494
495   **Gene Ontology (GO) term overrepresentation analysis**

496
We found overrepresented GO terms among the differential expressed gene to provide us with biological
processes that might be perturbed using the topGO package of R (v2.48.0; Alexa et al. 2006). Often termed
an enrichment analysis (Huang et al. 2009), the analysis specifically tests for overrepresentation of a GO
term from a cohort of provided genes (usually those that are differentially expressed) compared with the
expectation of the same number of random picks from all expressed genes. We performed these GO term
tests using Fisher's exact test with the weighted algorithm.

503
504   **Whole-genome bisulfite sequencing (WGBS) library preparation and sequencing**

505
Five biological replicates each of *eGFP* and *Dnmt1* were used for the differential methylation analysis. A
sample of 150 ng of genomic DNA was used to construct bisulfite treated genomic DNA Illumina
compatible libraries by the Novogene Corporation (Sacramento, CA, USA). Lambda phage gDNA, which
is known to be completely unmethylated, was added to these libraries to provide a negative control and
estimate conversion efficiency. We targeted 30X coverage of the genome using 2 x 150 bp pair-end reads.
Sequencing was performed on an Illumina NovaSeq.

512
513   **WGBS quality control, mapping, and cytosine methylation analysis**

514
As described above, reads were quality assessed with fastQC, trimmed with cutadapt using the TruSeq
adapter sequences, and reassessed with fastQC with default settings. Reads were mapped to the NCBI
RefSeq assembly (ASM185493v1) & annotation (GCF_001854935.1) using HISAT-3N (v2.2.1; C to T
base change, uniquely mapped non-clonal reads; Zhang et al., 2021; Supplementary File 2). We then used
HISAT-3N to generate a per nucleotide results table for all cytosines and a separate one for cytosines within
a CpG context, which is were almost all of cytosine methylation is found for insects (Bewick et al., 2017).
Using the HISAT-3N results tables per sample, we first filtered out cytosines that had fewer than five
mapped reads. We then used Python and the numpy, scipy. stats, statsmodel.stats, and pandas libraries to
generate per nucleotide result tables of percent reads that indicated methylation, total mapped reads, and

one-tailed binomial tests with FDR-corrected P values for each sample (P < 0.016, which was the average
non-conversion rate of reads mapped to the lambda phage controls). Methylation status of each cytosine
was collected into a per gene results table for all samples using bedtools (Quinlan and Hall, 2010) and
pandas. We also generated a results table that contained only CpGs that were symmetrically mapped above
our threshold. To be able to characterize cytosine methylation over repeats and transposons, we identified
transposable elements using RepeatMasker (v4.1.2; Smit et al., 2013) which was consistent with the
invertebrate repeat library from Repbase (Jurka, 1998) following Bewick and colleagues (2019). Cytosine
methylation over these genomic regions was collated with bedtools.

532
533   **Characterization of cytosine methylation**

534
535   We characterized the genomic context of cytosine methylation of the *B. tabaci* in several ways. We first
536   characterized the GC content of the genome using bedtools. We then calculated the percent of cytosines
537   that were methylated for several genomic regions: the genome as a whole, intragenic regions, genes, exons,
538   and introns. Lastly, we characterized percent of cytosines methylated across genes and transposons with
539   bedtools and plotted with deepTools2 (v3.5.1; Ramirez et al., 2016).

540
541   **Differential gene methylation and analysis**

542
After understanding the genomic context of cytosine methylation, we focused on understanding its
association with gene expression. First, we expected to see an association between gene methylation and
gene expression such that highly methylated genes were also the highest expressed genes (Duncan et al.,
2022). To that end, we divided genes into deciles and plotted those values against the mean standardized
expression of the gene within each decile using R. We also plotted how differences of gene methylation
were associated with gene expression standard deviations.
To understand the effect of *Dnmt1* RNAi on gene methylation, we tested for statistically
significantly differences of gene methylation levels between the two treatments using the t.test function of
R. Before this test, we filtered the results for genes that showed at least a 1% difference for percent of
cytosines methylated (2,391/13,901 genes) so the *t* test would not return an error due to data showing no
variation. We plotted the mean percent cytosines methylated for each gene between the two treatments.
Lastly, we assessed how differences of gene methylation were associated with difference of gene
expression by plotting log2 Fold Changes of gene expression by difference of mean percent methylated
cytosines.

Acknowledgements

559
560   We appreciate the discussions and contributions from other members of our laboratories, especially Ahva
561   Potticary and Bazgha Zia. The mention of a proprietary product does not constitute an endorsement or a
562   recommendation for its use by USDA or University of Georgia.

563
564   **Funding**

565
566   This work was funded by the USDA-ARS Non-Assistance Cooperative Agreement "Managing whiteflies
567   and whitefly-transmitted viruses in vegetable crops in the southeastern U.S." (#58-6080-9-006) to AJM.

Author Contribution Statement

CBC, PJM, AJM, EAS, and AS designed the study. EAS proposed examining the *Wnt* genetic pathway. EAS and ECM performed the experiments. CBC analyzed the data. CBC, PJM, AJM, and EAS interpreted the data. All authors contributed to drafting the manuscript.

Conflicts of Interest

The authors have no conflicts of interest to declare.

Data Availability

All high-throughput data is available under NCBI BioProject # XXXX. Analysis scripts are available at github.com/cbcunningham/Bt_RNA-seq_methylC-seq.

References

Alexa A, Rahnenfuhrer J, Lengauer T (2006) Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. Bioinformatics 22:1600–1607.

Arsala D, Wu X, Yi SV, Lynch JA (2022) *Dnmt1a* is essential for gene body methylation and the regulation of the zygotic genome in a wasp. PLoS Genetics 18:e1010181.

Amukamara AU, Washington JT, Sanchez Z, McKinney EC, Moore AJ, Schmitz RJ et al. (2020) More than DNA methylation: does pleiotropy drive the complex pattern of evolution of *Dnmt1*? Front Ecol Evol 8:10.3389.

Andrews S (2010) FastQC: A quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Benjamini Y, Krieger AM, Yekutelli D (2006) Adaptive linear step-up false discovery rate controlling procedures. Biometrika 93:491-507.

Bewick AJ, Vogel KJ, Moore AJ, Schimitz RJ (2017) Evolution of DNA methylation across insects. Mol Biol Evol 34:654-665.

Bewick AJ, Sanchez Z, McKinney EC, Moore AJ, Moore PJ, Schmitz RJ (2019) *Dnmt1* is essential for egg production and embryo viability in the large milkweed bug *Oncopeltus fasciatus*. Epigenet Chrom 12:1-14.

Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J (2021) eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. Mol Biol Evol 38:5825–5829.

Chassot AA, Gregoire EP, Lavery R, Taketo MM, de Rooij DG, Adams IR, Chanoissier M (2011) RSPO1/β-Catenin signaling pathway regulates oogonia differentiation and entry into meiosis in the mouse fetal ovary. PLoS ONE 6:e25641.

Chen W, et al. (2016) The draft genome of whitefly *Bemisia tabaci* MEAM1, a global crop pest, provides novel insights into virus transmission, host adaptation, and insecticide resistance. BMC Biol 14:110.

Cunningham CB, Shelby EA, McKinney EC, Schmitz RJ, Moore AJ, Moore PJ (2023) The role of *Dnmt1* during spermatogenesis of the insect *Oncopeltus fasciatus*. Epigenetics & Chromatin 16:28.

Dixon G, Matz M (2022) Changes in gene body methylation do not correlate with changes in gene expression in Anthozoa or Hexapoda. BMC Genomics 23:234.

Duncan EJ, Cunningham CB, Dearden PK (2022) Phenotypic plasticity: what has DNA methylation got to do with it? Insects 13:110.

Espada J et al. (2011) Regulation of SNAIL1 and E-cadherin function by DNMT1 in a DNA methylation-independent context. Nucleic Acids Res 39: 9194–9205.

Gegner J, Gegner T, Vogel H, Vilcinskas A (2020) Silencing of the *DNA methyltransferase 1 associated protein 1* (*DMAP1*) gene in the invasive ladybird *Harmonia axyridis* implies a role of the *DNA methyltransferase 1*complex in female fecundity. Insect Mol Biol 29:148–159.

Gladstad KM, Hunt BG, Yi SV, Goodisman MAD (2011) DNA methylation in insects: on the brink of the epigenomic era. Insect Mol Biol 20:553-565.

Gladstad KM, Hunt BG, Goodisman MAD (2014) Evolutionary insights into DNA methylation in insects. Curr Opin Insect Sci 1:25-30.

Glastad KM, Goodisman MA, Yi SV, Hunt BG (2016) Effects of DNA methylation and chromatin state on rates of molecular evolution in insects. G3-Genes Genomes Genet 6:357–363.

Grover S (2018) Potential of RNA interference in the study and management of the whitefly, *Bemisia tabaci*. Insect Biochem Physiol 100:e21522.

Hu Y, Yang Z, Gong C, Wu Q, Wang S, Li C, Zhang Y, Guo Z (2022) Molecular characterization of insulin-like peptide genes and their important roles in *Bemisia tabaci* reproduction. Entomol Gen 42:977-986.

Huang DW, Sherman BT, Lempicki RA (2009) Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res 37:1–13.

Ivasyk I, Olivos-Cisneros L, Valdés-Rodríguez S, Doual M, Jang H, Schmitz RJ, Kronauer DJC (2023) DNMT1 mutant ants develop normally but have disrupted oogenesis. Nature Commun 14:2201.

Jurka, J. (1998) Repeats in genomic DNA: mining and meaning. Curr Opin Struct Biol 8:333-337.

Kassambara A, Mundt F (2020) Factoextra: Extract and visualize the results of multivariate data analyses.
v1.0.7. CRAN.R-project.org/package=factoextra

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and genotyping
with HISAT2 and HISAT-genotype. Nat Biotechnol 37:907–915.

Kopylova E, Noé, Touzet H (2012) SortMeRNA: Fast and accurate filtering of ribosomal RNAs in
metatranscriptomic data. Bioinformatics 28:3211-3217.

Korotkevich F, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A (2021) Fast gene set
enrichment analysis. bioRxiv, doi.org/10.1101/060012.

Kuchari R, Ellis N, Jurkowshi TP, Hurd PJ, Maleszka R (2023) The PWWP domain and the evolution of
unique DNA methylation toolkits of Hymenoptera. SSRN doi.org/10.2139/ssrn.4471672

Law JA, Jacobsen, SE (2010) Establishing, maintaining, and modifying DNA methylation patterns in plants
and animals. Nat Rev Genet 11:204–220.

Le Rolle M et al. (2021) Arrest of WNT/β-catenin signaling enables the transition from pluripotent to
differentiated germ cells in mouse ovaries. Proc Natl Acad Sci USA 118: e2023376118.

Li B, Hu P, Zhang SZ, Toufeeq S, Wang J, Zhao K, Xu X, Xu JP, Huang SJ (2019) DNA methyltransferase
BmDnmt1 and *BmDnmt2* in silkworm (*Bombyx mori*) and the regulation of silkworm embryonic
development. Arch Insect Biochem Physiol 100: e21529

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biol 15:550.

Luo Y, Chen Q, Luan J, Chung S, Van Eck J, Turgeon R, Douglas AE (2017) Towards an understanding
of the molecular basis of effective RNAi against a global insect pest, the whitefly *Bemisia tabaci*. Insect
Biochem Mol Biol 88:21-29.

McKenzie CL, Sparks Jr AN, Roberts P, Oetting RD, Osborne LS (2020) Survey of *Bemisia tabaci*
(Hemiptera: Aleyrodidae) in agricultural ecosystems in Georgia. J Entomol Sci 55:163-170

Magoc T, Salzberg S (2011) FLASH: fast length adjustment of short reads to improve genome assemblies.
Bioinformatics 27:2957–2963

Maleszka R, Kucharski R (2022) Without mechanism, theories and models in insect epigenetics remain a
black box. Trends Genet 38:1108-1111.

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet j
17:10-12.

Morandin C, Brendel VP (2021) Tools and applications for integrative analysis of DNA methylation in
social insects. Mole Ecol Resour 22:1656-1674

Murat S, Hopfen C, McGregot AP (2010) The function and evolution of *Wnt* genes in arthropods. Arthropod
Struct Develop 39, 446-452.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg S (2015) StringTie enables
improved reconstruction of a transcriptome from RNA-seq reads. Nature Biotechnol 33:290-295.

Provataris P, Meusemann K, Neihuis O, Garth S, Misof B (2018) Signatures of DNA Methylation across
insects suggest reduced DNA methylation levels in holometabola. Genome Biol Evol 10:1185-1197.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics 26:841-842.

R Core Team (2021) R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. https://www.R-project.org/.

RStudio Team (2022) RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL
http://www.rstudio.com/.

Ramírez, Fidel, Devon P. Ryan, Björn Grüning, Vivek Bhardwaj, Fabian Kilpert, Andreas S. Richter,
Steffen Heyne, Friederike Dündar, and Thomas Manke. (2016) deepTools2: A next Generation Web Server
for Deep-Sequencing Data Analysis. Nuc Acids Res 44, W160-W165.

Schmitz RJ, Lewis ZA, Goll MG (2019) DNA methylation: Shared divergent features across eukaryotes.
Trends Genet 35:818-827.

Schulz NKE, Wagner CI, Ebeling J, Raddatz G, Diddens-de Buhr M, Lyko F et al. (2018) *Dnmt1* has an
essential function despite the absence of CpG DNA methylation in the red flour beetle *Tribolium*
castaneum. Sci Rep 8:16462.

Shelby EA, Moss JB, Andreason SA, Simons AM, Moore AJ, Moore PJ (2020) Debugging: Strategies and
considerations for efficient RNAi-mediated control of the whitefly *Bemisia tabaci*. Insects 11:723.

Shelby EA, McKinney EC, Cunningham CB, Simmons AM, Moore AJ, Moore PJ (2023) The role of *Dnmt1*
in oocyte development. J Insect Physiol 147:104507.

Smit, AFA, Hubley, R & Green, P. RepeatMasker Open-4.0. 2013-2015 <http://www.repeatmasker.org>

Soneson C, Love MI, Robinson MD (2015) Differential analyses for RNA-seq: transcript-level estimates
improve gene-level inferences. F1000Research, f1000research.7563.1.

Bemisia: bionomics and management of a global pest (2010) Eds., Stanley PA, Naranjo SE. Spinger, New
York. 544 pg.

Stephens M (2016) False discovery rates: a new deal. Biostatistics, 18:2.

Tadmor E, Juravel K, Morin S, Santos-Garcia D (2022) Evolved transcriptional responses and their trade-
offs after long-term adaptation of *Bemisia tabaci* to a marginally suitable host. Genome Biol Evol 14,
evac118.

Takada Y, (2021) Maintenance DNA methylation in pre-meiotic germ cells regulates meiotic prophase by
facilitating homologous chromosome pairing. Develop 148, dev194605.

Upadhyay SK, Singh H, Dixit S, Mendu V, Verma PC (2016). Molecular characterization of vitellogenin and
vitellogenin receptor of *Bemisia tabaci*. PLOS One, 11(5), e0155306.

Ventos-Alfonso A, Ylla G, Montanes J, Belles X (2020) DNMT1 promotes genome methylation and early
embryo development in cockroaches. iScience 23:101778.

Waghmare I, Page-McCaw A (2018) *Wnt* signaling in stem cell maintenance and differentiation in the
Drosophila germarium. genes 9:127.

Washington JT, Cavender KR, Amukamara AU, McKinney EC, Schmitz RJ, Moore PJ (2021) The essential
role of *Dnmt1* in gametogenesis in the large milkweed bug *Oncopeltus fasciatus*. eLife 10:e62202.

Xu GF, Gong CC, Tian YL, Fu TY, Lin YG, Lyu H, Peng YL, Tong CM, Feng QL, Song QS, Zheng SC
(2022) DNA methylation-mediated expression of zinc finger protein 615 affects embryonic development
in *Bombyx mori*. Zool Res 43: 552.

Zemach A, McDaniel IE, Silva P, Zilberman D (2010) Genome-wide evolutionary analysis of eukaryotic
DNA methylation. Science 328: 916-919.

Zhang Y, Park C, Bennett C, Thornton M, Kim D (2021) Rapid and accurate alignment of nucleotide
conversion sequencing reads with HISAT-3N. Genome Res 31:1290-1295.

Zhang J, Xing Y, Li Y, Yin C, Ge C, Li F (2015) DNA methyltransferases have an essential role in female
fecundity in brown planthopper, *Nilaparvata lugens* Biochem Biophys Res Comm 464:83-88.

Zhu A, Ibrahim JG, Love MI (2018) Heavy-tailed prior distributions for sequence count data: removing the
noise and preserving large differences. Bioinformatics 35:2084-2092.

Zwier MV, Verhulst EC, Zwahlen RD, Beukeboom LW, van de Zande L (2012) DNA methylation plays a
crucial role during early *Nasonia* development. Insect Mol Biol 21:129–138.
788