Unraveling variant-to-function relationships by integrating SNP-specific gene

expression networks and metabolic models

Srijith Sasikumar[1,2,3], Pavan Kumar S[2,3,4,5], Nirav Bhatt[2,3,4,5,6], Himanshu Sinha[1,2,3],*

[1] Systems Genetics Lab, Department of Biotechnology, Bhupat and Jyoti Mehta School of Biosciences, Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

[2] Centre for Integrative Biology and Systems Medicine (IBSE), Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

[3] Robert Bosch Centre for Data Science and Artificial Intelligence (RBCDSAI), Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

[4] BioSystems Engineering and Control (BiSECt) Lab, Department of Biotechnology, Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

[5] pCoE for Network Systems Learning, Control and Evolution, Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

[6] School of Data Science and Artificial Intelligence, Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

*Corresponding author: sinha@iitm.ac.in

ABSTRACT

Elucidating genotype-phenotype or variant-to-function relationships remains a challenge in genetics. For quantitative traits, causal SNPs act either additively or epistatically, resulting in complex interactions that are difficult to dissect molecularly. Here, we developed gene co-expression networks and genome-scale metabolic models for all combinations of causal SNPs of yeast sporulation efficiency, a quantitative trait, to determine how genetic interactions drive phenotypic variation. Analysis of expression networks identified SNP-SNP interaction-dependent changes in the connectivity of crucial metabolic regulators. These regulators revealed how some SNP combinations added to higher phenotypic value. Using IMAT and Localgini thresholding algorithm, we integrated the gene expression data of all SNP combinations in the Yeast genome-scale metabolic model, resulting in 16 SNP-specific metabolic models. Genome-scale differential flux analysis was used to identify differentially activated metabolic reactions across multiple metabolic pathways in each SNP-specific model. This analysis revealed causal reactions in six major metabolic pathways explaining the observed sporulation efficiency differences. Finally, the differential regulation of the pentose phosphate pathway in specific SNP combinations suggested autophagy as a pentose phosphate pathway-dependent compensatory mechanism for increasing sporulation efficiency. Our study presents a modelling framework that has the potential to discover causal metabolic pathways or reactions regulated by combinations of SNPs, providing insights into Genome-wide association studies (GWAS) variants of metabolic diseases.

AUTHOR SUMMARY

Understanding how genetic changes influence traits that vary in degree, such as human height, can be difficult to identify. In this study with yeast, we have developed an approach to understand how different genetic variations work together to affect a specific trait: sporulation efficiency. We used a network of genes that are co-expressed, or turned on and off together, to identify how different genetic variations interact with one another. We found that certain combinations of variations can make key genes in metabolic pathways more or less active, which can ultimately affect the overall efficiency of sporulation. To further understand how these genetic changes affect metabolism, we used a genome-scale model to simulate the metabolic pathways in yeast. By comparing the metabolic pathways of different genetic combinations, we identified specific reactions that were responsible for the observed changes in sporulation efficiency. Overall, this study provides a new way to understand how variations in our genes can affect our metabolism and could be particularly useful for identifying the genetic causes of metabolic diseases in humans.

KEYWORDS

Genome-scale metabolic models, Yeast sporulation, gene expression networks, quantitative trait, metabolism, genetic interactions, Single nucleotide polymorphisms.

INTRODUCTION

Genome-wide association studies (GWAS) have significantly advanced our understanding of complex traits by identifying causal single nucleotide polymorphisms (SNPs) and their intricate interactions [1,2]. These studies reveal that multiple SNPs, each exerting a modest impact on a trait, contribute to the complexity of genetic effects through additive and epistatic interactions. Unravelling this complexity within the broader genetic and biological network poses a considerable challenge, necessitating the development of robust computational tools and strategies to explore SNP interactions [3–6].

It is crucial to examine intermediate phenotypes to address the challenge of understanding how SNPs and their interactions influence complex traits, particularly the variant-to-function (V2F) relationship [7]. Various techniques, such as expression quantitative trait loci (e-QTLs), protein expression loci (p-QTLs), and metabolite loci (m-QTLs), impact diverse phenotypes [8,9]. In addition, SNPs influence gene co-expression and interaction networks, particularly in complex metabolic traits where coding and non-coding region variants can alter enzymatic reactions and metabolic flux [10].

Genome-scale metabolic models (GSMMs) provide a systems biology framework to describe cellular metabolism [11]. While previous studies have explored the direct impact of SNPs on enzyme coding regions [12,13], understanding how SNPs collectively reshape cellular metabolism in both coding and non-coding regions, especially in transcription factors, remains a gap to be addressed. Integrating SNP-specific gene expression alterations into GSMMs becomes essential to precisely decipher the effects of variants and their interactions on metabolic flux variations.

This study aims to fill this gap by leveraging GSMMs and integrating gene expression data of

causal SNPs and their combinations. Focusing on yeast sporulation as a model known for its

metabolic nature, the analysis investigates how SNPs and their combinations modulate

metabolic fluxes to influence sporulation efficiency [14–16]. The findings reveal that SNP

combinations uniquely drive divergent sets of differentially expressed genes and alter the

connectivity of critical metabolic regulators. Context-specific GSMMs for each SNP and their

combinations illustrate changes in flux distribution patterns and identify synergistic and

antagonistic metabolic pathway interactions in the yeast sporulation process. Additionally, a

new role for autophagy in yeast sporulation, dependent on the pentose phosphate pathway,

is predicted in an SNP-specific manner. Hence, integrating gene expression data into GSMMs

is an effective technique for deciphering the intricate relationships between SNPs and cellular

metabolism, paving the way for a more nuanced approach to therapeutic intervention.

RESULTS

To investigate how SNPs and their combinations contribute to phenotypic variation, we

conducted a comprehensive analysis using gene expression data from an allele replacement

panel of *Saccharomyces cerevisiae* [16] during sporulation. The SNPs studied here included

coding variants in sporulation genes *IME1*^{L325M}, *RSF1*^{D181G}, and non-coding variants in

IME1nc^{A-548G} and *RME1*$^{indel-308A}$) that conferred high sporulation efficiency phenotype. Here,

we present the results of our study, where we followed a multi-step approach. First, we

created SNP-specific differential gene expression networks highlighting sporulation,

metabolic processes, and regulators. Secondly, we employed a genome-scale metabolic

modeling approach to understand how these SNPs influence intracellular metabolic fluxes.

These analyses identified causal metabolic reactions that could explain increased phenotype

by specific SNPs and their combination, providing insight into how the SNPs interact to

modulate a quantitative trait phenotype. The entire workflow, from the pre-processing of

RNA-Seq data to the application of genome-scale metabolic modeling techniques, is

described in Figure 1.

Transcriptomic analysis reveals SNP-specific differentially expressed genes

Based on the sporulation efficiency data of the 16 allele replacement strains containing causal

oak alleles in the vineyard background strains [16], we classified *IME1_IME1nc_RME1_RSF1*,

IME1_IME1nc_RME1, *IME1_IME1nc_RSF1*, *IME1_RME1_RSF1*, *IME1nc_RME1_RSF1* allelic

combination strains as 'high sporulating.' All two allele combination strains (*IME1_IME1nc,*

IME1_RME1, IME1_RSF1, IME1nc_RME1, IME1nc_RSF1, RME1_RSF1) were 'medium

sporulating strains', and all single SNP and wildtype were considered 'low sporulating strains.'

Differential gene expression analysis, after adjusting for the false discovery rate (< 10% using

the Benjamini-Hochberg method), identified a total of 380 significant differentially expressed

genes (DEGs, upregulated and downregulated) across 15 SNP combinations (Figure 2A, Table

S1). A comparison of the number of DEGs in each SNP combination showed that the strain

with *IME1* coding and *IME1nc* combination had the highest number of DEGs. Interestingly, of

all the SNP combinations, we also found a significant downregulation of several genes in the

strain with the two non-coding alleles - *RME1* and *IME1nc*.

Since we were interested in determining the number of sporulation-related genes among

DEGs between wildtype and allele replacement strains, we compared DEGs with 362 genes

known to influence sporulation efficiency phenotype as described in the literature [17]. Of

the 362 genes, we identified 48 variable DEGs across all SNP combinations. Each SNP

combination had a different set of sporulation-specific DEGs, with the strain with all four

alleles having the maximum number (Figure 2B, Figure S1). The presence of *RME1nc* resulted

in the downregulation of *RME1* expression in all combinations except for *RME1nc_IME1nc*

(Figure S1). This showed that the cis effect of the *RME1nc* allele downregulated its expression,

enhancing sporulation efficiency in oak strains. Upregulation of the *IME1,* whenever the *RSF1*

allele was in combination with either *IME1* or *IME1nc* (Figure S1), might be due to a trans-

effect of the *RSF1* variation on the expression of *RIM4*, a major regulator of early sporulation

genes and regulates *IME1* expression.

Interestingly, no common genes were found to be upregulated across all five strains

(*IME1_IME1nc_RME1_RSF1*, *IME1_IME1nc_RME1*, *IME1_IME1nc_RSF1*, *IME1_RME1_RSF1*,

IME1nc_RME1_RSF1), even though several genes were uniquely activated in each SNP

combination strain (Figure 2C). However, *IME2*, a meiosis initiator gene, was activated and

uniquely upregulated when all four SNP were combined. We identified five upregulated

genes, *HSP26*, *IME1*, *REC8*, *RIM4*, and *YIRO16W*, common across the subset of strains with

IME1 and *RSF1* variants in combination (Figure 2D). In particular, *REC8*, a gene that mediates

sister chromatid cohesion, homologous recombination, and chromosome synapsis [18], was

uniquely upregulated only in this subset of strains, while *RIM4* was upregulated in an *RSF1*-

dependent manner (Figure 2E).

Since yeast sporulation is a metabolically driven process, we considered all metabolic genes

in the Yeast8 metabolic model and investigated if they were differentially expressed across

different SNP combinations. Of the 1150 metabolic genes, 164 were differentially expressed

across all SNP combinations (Figure S2). The highest number of differentially expressed

metabolic genes were observed in *IME1_IME1nc* (Figure 2B). Interestingly, we also identified

that SRT1, a gene involved in dolichol biosynthesis and whose null mutant showed reduced

sporulation efficiency, was differentially expressed whenever the RSF1 allele was in

combination with *IME1* and *IME1nc*, i.e., in *RSF1_IME1_IME1nc* and

RSF1_IME1_IME1nc_RME1 (Figure S2).

GO analysis, using ClueGO, of the upregulated DEGs revealed that meiosis nuclear division

was significantly enriched in the strains with SNP combinations of *RME1_RSF1_IME1_IME1nc*,

RME1_RSF1_IME1, and *RSF1_IME1nc_IME1* indicating that these strains have already been

undergoing meiosis during the second hour of sporulation (Figure S3). Further, we observed

that two non-coding SNPs (*RME1* and *IME1nc*), when combined, could activate genes

enriched for cellular respiration (Figure S3). These findings suggested that SNP interactions

could affect the timing and progression of meiosis during sporulation.

Connectivity of critical metabolic regulators changes as SNP interact

Given that the strain with all SNP combinations has the highest sporulation efficiency, we

examined how the connectivity of regulators that control the expression of differentially

expressed genes in this strain varies across other combinations of SNPs. Regulatory impact

factor (RIF1 and RIF2) analyses helped determine the TFs that control the expression of DEGs

in the strain with all 4 SNP combinations and also assigned a score to these TFs. In brief, the

RIF1 metric classified TFs that were differentially co-expressed with highly abundant DEGs,

and RIF2 was used to identify the TFs with the altered ability to predict the abundance of

DEGs. Thus, the RIF1 metric identified TFs most differentially co-expressed with highly

abundant DE genes. On the other hand, the RIF2 measure was used to determine the TFs with

the most significant altered capacity to predict the abundance of DEGs. Using the RIF1 and

RIF2 *z*-score cut-offs, we obtained 15 TFs as regulators of DEG expression (Table S2). The RIF1

analysis identified *GCR1* (*z*-score = 2.40), a transcriptional activator of glycolysis [19] gene

with the highest value. The top two candidate transcription factors from RIF2 analysis were

ROF1 (z-score = 2.22), a putative TF, and *BAS1* (*z*-score = 2.18), a Myb-related TF involved in

regulating basal and induced expression of genes of the purine and histidine biosynthesis

pathways [20]. Of these 15 regulators, we were able to identify seven metabolic regulators:

UPC2 involved in sterol biosynthesis, *GCR1, GCR2* in glycolysis, *BAS1* in histidine biosynthesis,

and *ARG80, LYS14,* and *MET32* in amino acid metabolism. We then constructed fifteen

networks to identify differences in gene co-expression profiles among the wild type and 14

allelic combinations using the partial correlation and information theory (PCIT) algorithm. The

correlation of 6,307 genes with a cut-off greater than |0.95| retrieved varying numbers of

correlated pairs across wildtype and allele-specific networks (Table 1).

Filtering for co-expressed gene pairs that were DEGs between wildtype (null model) and all

oak allele combination groups (143 genes) along with the 15 TFs from the RIF analyses ($R^2 >$

|0.95|) allowed the construction of 15 allele-specific sub-networks with varying numbers of

nodes and edges (Table S3). These allele-specific sub-networks permitted the determination

of how these TFs regulated gene expression by varying their connectivity across the SNPs and

their combinations (Figure 3A). Further, we found that the connectivity of *UPC2* differed

between high, moderate, and low sporulating strains (Figure 3B). These findings confirm that

SNP combinations significantly impact overall transcriptional activity and highlight the

possibility that critical metabolic alterations could be responsible for sporulation efficiency

variation between allele replacement strains.

SNP-specific metabolic models show metabolic heterogeneity

As previously established, numerous genes and pathways undergo deregulation during yeast

sporulation [21,22]. Additionally, our analysis showed that the SNPs studied here could

activate different genes (Figure 2A). Hence, extracting only the active genes and pathways from the generalized yeast metabolic model is crucial to reconstructing SNP-specific metabolic models. We utilized Localgini thresholding and iMAT to integrate gene expression data from each allele replacement strain with the Yeast8 genome-scale metabolic model following the procedure outlined in Figure 1. Each resulting SNP-specific model (M1 to M16) corresponded to its genotype and varied in the number of genes, reactions, and metabolites (Table 1). The metabolic heterogeneity among the SNP-specific models was captured by calculating the Hamming distance between each model (Figure S4). For example, it was clear from this analysis that M15 with *IME1nc* had a unique set of metabolic reactions compared to the rest of the models.

Table 1: Number of reactions, metabolites, and genes in each SNP-specific model generated using iMAT+Localgini thresholding. 'O' denotes the oak allele, and 'W' indicates the vineyard allele.

Models	Genes	Reactions	Metabolites	Genotypes
M1	603	1726	1227	$RME1^O RSF1^O IME1^O IME1nc^O$
M2	595	1745	1246	$RME1^W RSF1^O IME1^O IME1nc^O$
M3	598	1737	1240	$RME1^O RSF1^W IME1^O IME1nc^O$
M4	557	1737	1242	$RME1^O RSF1^O IME1^W IME1nc^O$
M5	569	1696	1209	$RME1^O RSF1^O IME1^O IME1nc^W$
M6	579	1717	1222	$RME1^W RSF1^W IME1^O IME1nc^O$
M7	595	1742	1238	$RME1^W RSF1^O IME1^W IME1nc^O$
M8	570	1604	1139	$RME1^W RSF1^O IME1^O IME1nc^W$
M9	602	1750	1247	$RME1^O RSF1^W IME1^W IME1nc^O$
M10	515	1560	1088	$RME1^O RSF1^W IME1^O IME1nc^W$
M11	594	1733	1233	$RME1^O RSF1^O IME1^W IME1nc^W$
M12	565	1644	1155	$RME1^O RSF1^W IME1^W IME1nc^W$
M13	591	1731	1237	$RME1^W RSF1^O IME1^W IME1nc^W$
M14	571	1720	1219	$RME1^W RSF1^W IME1^O IME1nc^W$
M15	504	1247	1012	$RME1^W RSF1^W IME1^W IME1nc^O$

M16	596	1728	1231	$RME1^W RSF1^W IME1^W IME1nc^W$

A set of fundamental reactions consists of the core meiosis-specific network shared by all the

extracted models and is crucial for the survival and sporulation of yeast strains in an acetate

medium. We identified 401 reactions common in all extracted models and hypothesized that

these reactions form the core meiosis network. As expected, these reactions are significantly

enriched with pathways that play essential roles in sporulation (Figure S5). Since each SNP-

specific model had different metabolic reactions, we hypothesized that this could be either

due to the activation of novel reactions in each model or variations in intracellular flux within

the meiosis core network.

SNPs activate unique subsystems to modulate sporulation efficiency variation

Using the wildtype (M16) model as a baseline, a subnetwork topology analysis on each SNP

model helped investigate the contribution of each SNP combination to the variation in

sporulation efficiency. This analysis revealed that each SNP model had uniquely enriched

subsystems (a set of reactions involved sharing a common function, Table S4). For example,

the M13 model with *RSF1* SNP showed enrichment of valine, leucine, and isoleucine

degradation pathways, glycolysis, gluconeogenesis, pyruvate metabolism pathways, and the

TCA cycle (Figure 4A). Identifying metabolic pathways required during respiration in the M13

model (*RFS1*, Figure 4A) using the subnetwork topology analysis was consistent with the

function of the *RSF1* gene as a gene responsible for respiration, particularly during the early

sporulation [23,24]. Furthermore, purine metabolism is uniquely enriched only in the M12

model (*RME1nc*, Figure 4A).

The *IME1* coding variant, which affects the interactions between Rim11 and Ume6, is crucial

for initiating the sporulation [25,26]. The study by Sudarsanam and Cohen [16] revealed that

IME1 coding and non-coding variants contribute to sporulation efficiency variation, with the

coding variant explaining a more significant proportion. A comparison of SNP-specific

metabolic models, *IME1* (M14) and *IME1nc* (M15), revealed significant differences in the

metabolic activity of the two variants. Enrichment analysis of active reactions indicated that

the *IME1* coding variant activated autophagy, one carbon pool by folate, phagosome,

glycerolipid metabolism, and propionate metabolism pathways (Figure S6A). In contrast, the

IME1nc did not trigger any of these metabolic pathways. This highlighted the metabolic

plasticity of the functional and non-functional forms of the Ime1 protein.

Next, we investigated whether, in comparison to their individual effects, interactions

between SNPs had the potential to activate or deactivate particular cellular pathways

selectively. In other words, we investigated whether these interactions could result in

synergistic or antagonistic effects on specific metabolic reactions. To demonstrate this idea,

we examined the combination of two coding SNPs (*RSF1_IME1* (M8)) and two non-coding

SNPs *RME1_IME1nc* (M9) as well as their single SNP models – *RSF1* (M13), *IME1* (M14),

RME1 (M12), and *IME1nc* (M15). By comparing reactions that were activated exclusively in

the M8 (*RSF1_IME1*) model but not in the M13 (*RSF1*), M14 (*IME1*), and M16 (wildtype)

models, we identified 113 synergistic reactions. These reactions were notably enriched in

glycerophospholipid, glycerolipid, and pyrimidine metabolism pathways (Figure 4B, Table S5).

Conversely, when identifying reactions that were inactive in the M8 model but active in the

M13, M14, and M16 models, we determined 271 antagonistic reactions. These antagonistic

reactions were enriched in various subsystems, including fatty acid biosynthesis, lipid

metabolism, the phosphatidyl inositol signalling pathway, steroid biosynthesis, etc. (Table S5,

Figure S7). Similarly, when comparing the *RME1* (M12), *IME1nc* (M15), and *RME1_IME1nc*

(M9) models, we uncovered the role of *RME1_IME1nc* in synergistically modulating the "one

carbon pool by folate" subsystems, encompassing the "glycine cleavage complex" and

"tetrahydrofolate aminomethyl transferase" (Figure S8, Table S6).

GS-DFA analysis reveals the impact of SNP-SNP interactions on intracellular flux variation

The analysis of the unique sub-network topology showed that combinations of SNPs could

activate shared metabolic pathways. However, whether SNPs could modify the flux

distribution patterns of these pathways to modulate phenotypic variation was unclear.

Therefore, flux variations in common reactions among SNP and wildtype models were

analyzed to investigate flux distribution patterns. While the flux balance analysis (FBA, [27]),

for a given constraint, can find multiple potential flux values for reactions in a metabolic

model, it cannot ensure the uniqueness of a solution. To overcome this limitation in FBA, we

used flux sampling to assess variations in metabolic flux distribution using the genome-scale

differential flux analysis (GS-DFA) described in the Methods section. Using GS-DFA analysis,

we identified upregulated and downregulated reactions in each SNP model compared to the

wildtype model. The complete list of upregulated and downregulated reactions is provided in

Table S7 and Table S8, respectively.

GS-DFA analysis showed that the upregulated reactions were primarily enriched in the

following six subsystems involved in the biosynthesis of amino acids, antibiotics, secondary

metabolites, glycerophospholipid metabolism, glycerolipid metabolism, and pentose

phosphate pathway in a strain with all four SNP combinations (M1, Figure 5B, Table S9).

As M1 had the highest sporulation efficiency, we asked if these six subsystems were present

in the other high sporulation efficiency strains (M2-M5). We found that in M2-M5, the six

subsystems were variable subsets of the M1 model, highlighting intracellular flux variation

among these pathways as significant contributors to sporulation efficiency variation among

these strains (Figure 5B). The two subsystems, glycerophospholipid and glycerolipid

metabolism, present in all models suggested that they were essential for the overall fitness

of the yeast strains during sporulation.

Further, we explored how the functional and non-functional forms of Ime1 protein could

change the flux distribution patterns to modulate sporulation efficiency variation. Using the

GS-DFA analysis, we found that reactions involved in glycerolipid metabolism,

glycerophospholipid metabolism, and biosynthesis of secondary metabolism were

upregulated in *IME1_IME1nc* (M6) and *IME1* (M14) models but not in *IME1nc* (M15) model

(Figure S6B). Interestingly, we could also predict the metabolic consequence of the non-

functional form of *IME1* (M15), which was to upregulate the reactions involved in the

peroxisome, fatty acid metabolism, fatty acid degradation, and fatty acid biosynthesis (Figure

S6B).

We found that models were clustered into two groups upon hierarchically clustering the Flux

Change (FC) values of upregulated reactions in the M1 model across other models (Figure S9).

Hence, we were interested in exploring differentially upregulated subsystems in each SNP-

specific model. We found that SNP combinations without the *IME1nc* variant, i.e., M5, M8,

M10, M11, M12, M14, and the M1 models (which contains all four SNPs), led to an increased

flux through reactions involved in the pentose phosphate pathway (Table 2, Table S9). This

result was consistent with our flux variability analysis (see Supplementary Methods section,

[28]), where we found that the PPP reactions were upregulated in these models (Figure S10).

The increased flux through PPP reactions was inversely coupled with the glucose-6-phosphate

isomerase reaction. This indicated that the flux was redirected through the pentose

phosphate pathway for biosynthesis rather than glycolysis for energy generation (Figure 5C).

In highly sporulating SNP-specific models M2, M3, and M4, which had *IME1nc* variant and

where the pentose phosphate pathway was not upregulated (Figure 5C), showed enrichment

of reactions involved in autophagy and phagosomes (Figure 5A, 5B). These findings were

interesting since previous studies have established a correlation between autophagy and

yeast sporulation, and the deletion of several autophagy-related genes has shown reduced

sporulation [29,30]. Moreover, we found that the upregulation of reactions involved in amino

acid biosynthesis was observed only in the M1, M4, and M11 models (Table S5), highlighting

the effect of SNP combinations *RME1_RSF1* and their interaction with *IME1nc* variants on flux

distribution patterns. These findings demonstrate the significant impact of SNPs and their

interactions on the metabolic flux distribution, explaining how these interactions affected

sporulation efficiency.

Table 2: Differential regulation of Pentose phosphate pathway subsystem across SNP models

in an *IME1nc*-dependent manner (highlighted in bold).

PPP normal	*RME1_**IME1nc***
	*IME1_**IME1nc***
	*RME1_IME1_**IME1nc***
	*RME1_RSF1_**IME1nc***
	*IME1_RSF1_**IME1nc***
	RSF1
	IME1nc
	*RSF1_**IME1nc***
PPP upregulation	*RME1*
	IME1
	RME1_IME1
	RME1_RSF1

	IME1_RSF1
	RME1_RSF1_IME1
	RME1_RSF1_IME1_IME1nc

DISCUSSION

Here, we attempted to bridge the V2F challenge for a well-studied complex trait in yeast –

sporulation efficiency. By harnessing the gene expression data from an allele replacement

panel of four SNPs, we showed how SNPs interacted and modulated the metabolic pathways

to drive sporulation efficiency variation. We showed that certain SNP combinations could

activate or deactivate the flux through a particular reaction. We identified nucleotide

metabolism (purine and pyrimidine metabolism) was synergistically modulated while 'steroid

biosynthesis' was antagonistically modulated when coding variants of *RSF1* and *IME1* were in

combination. This insight gave us a deeper understanding of how SNP-SNP interactions could

redirect metabolic flux, prioritizing nucleotide synthesis over steroid synthesis in the early

stages of meiotic progression during sporulation. Previous studies have underscored the

pivotal role of pyrimidine metabolism in facilitating efficient meiotic progression [15].

Further, the SNP-specific modulation of the glycine cleavage system was established in this

study, where the SNPs of *RME1* and *IME1nc* were synergistic.

We characterised the intracellular metabolic flux states of each SNP-specific model without

requiring an objective function as previously described [31,32]. Using GS-DFA analysis, we

identified flux variation in six major pathways as significant contributors to sporulation

efficiency variation. Interestingly, our GS-DFA analysis did not identify the fundamental

reactions that were known to constitute the meiosis-specific metabolic model [14]. These

included the TCA cycle, glyoxylate cycle, and acetate uptake, which have been shown to influence sporulation at an early stage through gene-knockout approaches. However, we were able to capture these pathways in our core meiosis-specific network by comparing the common reactions shared among SNP-specific models. These findings suggested that SNPs and their interactions might modulate sporulation efficiency variation by influencing metabolic reactions beyond the essential ones required for sporulation, highlighting the specific effects of genetic variations at the SNP level instead of the broader gene deletion effects. Our analysis also predicted the differential regulation of the pentose phosphate pathway in an *IME1nc*-specific manner. The pentose phosphate pathway is critical in supplying the cell with the necessary building blocks and reducing equivalents needed for the biosynthesis of nucleotides and amino acids [33]. Our hypothesis suggested that autophagy could produce the essential precursors for nucleotide and amino acid biosynthesis by degrading and recycling cellular components compensating for PPP, particularly in models where PPP upregulation was absent (Figure S11). Previous studies have shown autophagy as a compensatory mechanism for the PPP in tumour cells [34]. It would be interesting to experimentally validate whether autophagosomes are induced when the PPP reaction is downregulated. Thus, integrating SNP and their combination-specific gene expression data in GSMM has provided several novel mechanisms by which phenotypic variation in a well-studied trait is regulated at the molecular level.

Several studies have reported complex genetic interactions between identified SNPs in metabolic diseases [35,36]. A recent study has used GWAS variant data to develop personalized organ-specific metabolic models for 524,615 individuals of the INTERVAL and UK Biobank cohorts to clarify the impact of genetic variations on the metabolic processes

implicated in coronary artery disease [37]. Studying the population-level data by clustering

based on variant combinations rather than phenotype along with GS-DFA analysis can provide

additional novel targets for therapeutic interventions in complex human metabolic diseases.

Applying SNP-specific genome-scale metabolic models that integrate transcriptomic and

metabolomic data will help identify known and novel pathways perturbed by SNPs and their

combinations. With scRNASeq becoming common and generated for diseases such as

cancers, single-cell GSMM [38,39] can open a new level of metabolic heterogeneity

corresponding to cellular heterogeneity. Combining this with the power of scRNASeq with

context-specific GSMM can provide novel insights into cellular heterogeneity and uncover

causes of disease, drug response, drug resistance, etc. Therefore, this can be a robust

framework to provide candidate genes for further validation in the molecular processes and

phenotype-genotype relationship, leading to clearer variant-to-function pathways.

METHODS

RNA-Seq data, pre-processing, and differential expression analysis

Raw RNA sequence read data for single SNP and their combinations was downloaded from

NCBI GEO (accession number GSE55409, [16]) for 16 isogenic allele replacement strains of the

oak alleles of four SNPs *IME1*, *RME1, RSF1,* and *IME1nc*. The overview of the dataset is

discussed in Supplementary Methods.

The reads were converted to Fastq format using the command 'Fastqdump' in the SRA toolkit.

The FastQC and MutiQC were performed for quality control [40]. The reads were mapped to

reference *Saccharomyces cerevisiae* R64-1-1.105 downloaded from the Ensembl database

(https://ftp.ensembl.org/pub/release110/fasta/saccharomyces_cerevisiae/cdna/Saccharom

yces_cerevisiae.R64-1-1.cdna.all.fa.gz) using Salmon (an alignment-free tool), accounting for

GC and sequence bias [41]. The transcript TPM values obtained from Salmon were converted

to gene-level counts using Tximport v1.4.0 [42]. The effect of day on the gene expression

values was removed using ComBatseq [43].

Differential gene expression analysis compared each allele replacement strain with the

wildtype strain using DESeq2 [44]. The batch-corrected untransformed raw counts were given

as input to the DESeq2. After filtering for low counts with less than 100 counts in all 63

samples, 6305 genes out of 6571 were selected for further processing. The effect of the day

was included in the design model of DESeq2 to account for the contribution of covariates to

the total gene expression. log2 fold changes (LFC) shrinkage using the alpegm method [45],

with a false discovery rate of 10% (p-value adjusted using the Benjamini-Hochberg method),

was used to identify differentially expressed genes (DEGs). Genes with LFC values greater than

0.5 and less than -0.5 are considered upregulated and down-regulated, respectively. The

upregulated DEGs obtained for each allele replacement strain are given as input to ClueGO

[46] from Cytoscape [47] for gene ontology and functional annotations analysis.

Regulatory impact factor (RIF) analysis

CeTF package in R was utilized to perform the Regulatory Impact Factors analysis [48,49]. The

Variance stabilizing transformation (VST) was performed on the gene expression data of

wildtype and strain with four SNPs. It was given as input for the RIF analysis as it required two

contrasting conditions. The Transcription factors (TFs), downloaded from the Yeastract

database (http://www.yeastract.com/, [50]), were contrasted to the list of DEGs identified by

comparing the wildtype strain (i.e., a strain with no SNPs) to the strain with all SNP

combinations. The TFs identified using the RIF analysis with scores deviating ± 1.96 SD from

the mean were considered significant and termed 'critical regulators.'

PCIT and connectivity analysis

The partial correlation and information theory (PCIT) approach [51] was used to arrive at co-

expression networks for wildtype and 14 SNP combinations except for strain with *IME1*, which

had only three replicates. PCIT evaluated all possible triplets of genes, identifying the strongly

associated pair after screening all the network's genes. The correlation of |0.95| and based

on the p-value < 0.05 was used as a threshold for arriving at a significant association (this

threshold was chosen to arrive at a maximum number of significant edges, [52]). Further, the

DEGs (specific for wild type vs. four SNP combination strains) and critical regulators identified

using the RIF analysis are filtered to arrive at sub-networks for each SNP-specific network. The

individual co-expression sub-networks for wild type and 14 allele replacement strains were

visualized using Cytoscape [47] version (3.9.1). Each subnetwork was analyzed for network

properties using the Cytoscape plugin NetworkAnalyser 4.4.8. The connectivity for each

critical regulator was identified by dividing the degree of that regulator in any specific

network with the highest degree in that network [52,53]. The connectivity of each critical

regulator was then compared across each SNP combination and was visualized using the

Pheatmap package in R (v1.0.12, https://CRAN.R-project.org/package=pheatmap).

Reconstruction of SNP-specific metabolic models

Yeast 8.4.2 [54], a commonly used yeast Genome-scale metabolic model, was used in this

study. It is accessible via GitHub (https://github.com/sysbiochalmers/yeast-gem). This GSMM

had 1,150 genes, 4,058 reactions, and 2,742 metabolites. The bounds of exchange reactions

of the GSMM were changed to reflect the early stages of the sporulation state appropriately.

For this, we restricted the intake of glucose and nitrogen (Lower bound=upper bound=0

[mmol/(g DW h)]) while allowing an unrestricted supply of oxygen and acetate (lower bound=

-1000 [mmol/(g DW h)], [14]).

The gene expression data (TPM values) averaged across replicates of wild type and 15 allele

replacement strains were pre-processed using Localgini thresholding [55] and integrated into

Yeast 8.4.2 using the model extraction method, iMAT [56]. The detailed methodology for

reconstructing an SNP-specific model using Localgini thresholding and iMAT is discussed in

Supplementary Methods. This led us to reconstruct 15 SNP-specific models (M1 to M15) and

a wildtype model (M16).

To understand the metabolic heterogeneity of each extracted model, we created a Boolean

reaction vector for each SNP-specific model. In brief, for each model, we made a Boolean

vector of length 4058 (corresponding to all the reactions in the Yeast8 metabolic model) with

1s and 0s, indicating whether a reaction was included in the SNP-specific model. Further, we

identified reactions in each SNP-specific model that were not in the wildtype model. These

reactions, active in SNP models while not in wildtype, were searched for significantly enriched

subsystems using the flux enrichment analysis function $(FEA.m)$ provided in the COBRA

toolbox [57]. In brief, $FEA.m$ used a hypergeometric 1-sided test (α = 0.05) and FDR

correction (adjusted p-value < 0.05) to obtain significantly enriched subsystems.

Genome-scale differential flux analysis (GS-DFA) using a flux sampling approach

To discern how each SNP and their combinations can modulate the intracellular fluxes to

show sporulation efficiency variation, we leveraged the random flux sampling approach,

which acquires a significant number of evenly distributed solutions over the whole solution

space and does not require an objective function. We adopted a previously developed protocol [32] to identify the most significantly dysregulated reactions in the presence of a particular SNP and their combination by comparing the wildtype (M16) and each SNP model (M1 to M15). However, we modified the pipeline to use an optGpSampler [58] instead of an ACHR sampler. This modification was based on previous studies that demonstrated the optGpSampler's advantage in faster convergence and reduced run time compared to the ACHR sampler [59]. The optGpSampler in COBRApy version 0.10.1 [60] was used in this study to sample 10,000 flux solutions, and the thinning factor was set to 100 for all models. This was to ensure the spanning of the whole flux space and to obtain sparse and un-correlated fluxes [32]. Let X_{SNP} and $X_{Wildtype}$ be the flux distributions of a particular reaction in SNP and wildtype models, respectively. We identified the significantly deregulated reactions in each SNP model compared to the wild type using a two-sided Kolmogorov-Smirnov test with a significance level of 0.05 (adjusted p-value using the Benjamini Hochberg method). Upregulated and downregulated reactions were identified from these dysregulated reactions using the 'Flux change' value obtained. The reactions with flux change greater than 0.82 were considered upregulated, and flux change less than -0.82 were considered downregulated, indicating 10 times higher or lower fold change in flux in the SNP model compared to the wildtype model.

The flux change was calculated for each reaction as follows:

$$Flux\ change = \frac{\bar{X}_{SNP} - \bar{X}_{Wildtype}}{\left|\bar{X}_{SNP} + \bar{X}_{Wildtype}\right|}$$

where, $\bar{X}_{SNP}$ and $\bar{X}_{Wildtype}$, the arithmetic mean of the flux distributions for a given reaction in SNP and wildtype conditions, respectively.

COBRA toolbox and COBRApy setup

COBRA Toolbox version 3.0.0 and COBRApy version 0.10.1 on MATLAB R2021b and Python version 3.5.2, respectively. The optimizer used for linear programming and mixed-integer linear programming optimization was Gurobi, version 8.0.0. optGpSampler was run using parallel processes. We used customized Python scripts for statistical analysis and visualization of flux sampling results.

CODE AVAILABILITY

The codes used in this paper are available in the following GitHub repository (https://github.com/HimanshuLab/Modelling-and-transcriptomics-analysis-of-QTNs), along with all generated models.

ACKNOWLEDGEMENTS

S.S. and P.K.S. are supported by a fellowship from the Indian Institute of Technology Madras; P.K.S. was also supported by a grant from the Centre for Integrative Biology and Systems Medicine (IBSE), IIT Madras (BIO/18-19/304/ALUM/KARH) to H.S. We thank Dr Karthik Raman for critically reading our manuscript.

AUTHOR CONTRIBUTIONS

Conceptualization: S.S., and H.S.; Formal Analysis: S.S., P.K.S., and N.B.; Investigation: S.S., P.K.S., N.B., and H.S.; Writing original draft: S.S. and H.S.; Supervision: H.S.; All authors reviewed and approved the final paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1. Visscher PM, Wray NR, Zhang Q, et al. 10 years of GWAS discovery: Biology, function, and translation. *Am J Hum Genet*. 2017;101(1):5-22. doi:10.1016/j.ajhg.2017.06.005

2. Tam V, Patel N, Turcotte M, Bossé Y, Paré G, Meyre D. Benefits and limitations of genome-wide association studies. *Nat Rev Genet*. 2019;20(8):467-484. doi:10.1038/s41576-019-0127-1

3. Cordell HJ. Detecting gene–gene interactions that underlie human diseases. *Nat Rev Genet*. 2009;10(6):392-404. doi:10.1038/nrg2579

4. Martin MP, Gao X, Lee JH, et al. Epistatic interaction between KIR3DS1 and HLA-B delays the progression to AIDS. *Nat Genet*. 2002;31(4):429-434. doi:10.1038/ng934

5. Wan X, Yang C, Yang Q, et al. BOOST: A fast approach to detecting gene-gene interactions in genome-wide case-control studies. *Am J Hum Genet*. 2010;87(3):325-340. doi:10.1016/j.ajhg.2010.07.021

6. Fang G, Wang W, Paunic V, et al. Discovering genetic interactions bridging pathways in genome-wide association studies. *Nat Commun*. 2019;10(1):4274. doi:10.1038/s41467-019-12131-7

7. Cano-Gamez E, Trynka G. From GWAS to function: Using functional genomics to identify the mechanisms underlying complex diseases. *Front Genet*. 2020;11:424. doi:10.3389/fgene.2020.00424

8. Battle A, Khan Z, Wang SH, et al. Genomic variation. Impact of regulatory variation from RNA to protein. *Science*. 2015;347(6222):664-667. doi:10.1126/science.1260793

9. Olayinka OA, O'Neill NK, Farrer LA, Wang G, Zhang X. Molecular quantitative trait locus mapping in human complex diseases. *Curr Protoc*. 2022;2(5):e426. doi:10.1002/cpz1.426

10. Tong H, Küken A, Razaghi-Moghadam Z, Nikoloski Z. Characterization of effects of genetic variants via genome-scale metabolic modelling. *Cell Mol Life Sci*. 2021;78(12):5123-5138. doi:10.1007/s00018-021-03844-4

11. Gu C, Kim GB, Kim WJ, Kim HU, Lee SY. Current status and applications of genome-scale metabolic models. *Genome Biol*. 2019;20(1):121. doi:10.1186/s13059-019-1730-3

12. Øyås O, Borrell S, Trauner A, et al. Model-based integration of genomics and metabolomics reveals SNP functionality in *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A*. 2020;117(15):8494-8502. doi:10.1073/pnas.1915551117

13. Sarkar D, Maranas CD. SNPeffect: identifying functional roles of SNPs using metabolic networks. *Plant J*. 2020;103(2):512-531. doi:10.1111/tpj.14746

14. Ray D, Ye P. Characterization of the metabolic requirements in yeast meiosis. *PLoS One*.
2013;8(5):e63707. doi:10.1371/journal.pone.0063707

15. Walther T, Létisse F, Peyriga L, et al. Developmental stage dependent metabolic regulation
during meiotic differentiation in budding yeast. *BMC Biol*. 2014;12(1):60.
doi:10.1186/s12915-014-0060-x

16. Sudarsanam P, Cohen BA. Single nucleotide variants in transcription factors associate
more tightly with phenotype than with gene expression. *PLoS Genet*.
2014;10(5):e1004325. doi:10.1371/journal.pgen.1004325

17. Deutschbauer AM, Williams RM, Chu AM, Davis RW. Parallel phenotypic analysis of
sporulation and postgermination growth in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci*
U S A. 2002;99(24):15530-15535. doi:10.1073/pnas.202604399

18. Yoon SW, Lee MS, Xaver M, et al. Meiotic prophase roles of Rec8 in crossover
recombination and chromosome structure. *Nucleic Acids Res*. 2016;44(19):9296-9314.
doi:10.1093/nar/gkw682

19. Holland MJ, Yokoi T, Holland JP, Myambo K, Innis MA. The *GCR1* gene encodes a positive
transcriptional regulator of the enolase and glyceraldehyde-3-phosphate dehydrogenase
gene families in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 1987;7(2):813-820.
doi:10.1128/mcb.7.2.813-820.1987

20. Daignan-Fornier B, Fink GR. Coregulation of purine and histidine biosynthesis by the
transcriptional activators *BAS1* and *BAS2*. *Proc Natl Acad Sci U S A*. 1992;89(15):6746-
6750. doi:10.1073/pnas.89.15.6746

21. Primig M, Williams RM, Winzeler EA, et al. The core meiotic transcriptome in budding
yeasts. *Nat Genet*. 2000;26(4):415-423. doi:10.1038/82539

22. Enyenihi AH, Saunders WS. Large-scale functional genomic analysis of sporulation and
meiosis in *Saccharomyces cerevisiae*. *Genetics*. 2003;163(1):47-54.
doi:10.1093/genetics/163.1.47

23. Lu L, Roberts G, Simon K, Yu J, Hudson AP. Rsf1p, a protein required for respiratory growth
of *Saccharomyces cerevisiae*. *Curr Genet*. 2003;43(4):263-272. doi:10.1007/s00294-003-
0398-z

24. Roberts GG 3rd, Hudson AP. Rsf1p is required for an efficient metabolic shift from
fermentative to glycerol-based respiratory growth in *S. cerevisiae*. *Yeast*. 2009;26(2):95-
110. doi:10.1002/yea.1655

25. Rubin-Bejerano I, Mandel S, Robzyk K, Kassir Y. Induction of meiosis in *Saccharomyces*
cerevisiae depends on conversion of the transcriptional repressor Ume6 to a positive
regulator by its regulated association with the transcriptional activator Ime1. *Mol Cell Biol*.
1996;16(5):2518-2526. doi:10.1128/MCB.16.5.2518

26. Gertz J, Gerke JP, Cohen BA. Epistasis in a quantitative trait captured by a molecular model
of transcription factor interactions. *Theor Popul Biol*. 2010;77(1):1-5.
doi:10.1016/j.tpb.2009.10.002

27. Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? *Nat Biotechnol*.
2010;28(3):245-248. doi:10.1038/nbt.1614

28. Gudmundsson S, Thiele I. Computationally efficient flux variability analysis. *BMC
Bioinformatics*. 2010;11(1):489. doi:10.1186/1471-2105-11-489

29. Barve G, Manjithaya R. Cross-talk between autophagy and sporulation in *Saccharomyces
cerevisiae*. *Yeast*. 2021;38(7):401-413. doi:10.1002/yea.3556

30. Wen FP, Guo YS, Hu Y, et al. Distinct temporal requirements for autophagy and the
proteasome in yeast meiosis. *Autophagy*. 2016;12(4):671-688.
doi:10.1080/15548627.2016.1149659

31. Scott WT Jr, Smid EJ, Block DE, Notebaart RA. Metabolic flux sampling predicts strain-
dependent differences related to aroma production among commercial wine yeasts.
Microb Cell Fact. 2021;20(1):204. doi:10.1186/s12934-021-01694-0

32. Nanda P, Ghosh A. Genome Scale-Differential Flux Analysis reveals deregulation of lung
cell metabolism on SARS-CoV-2 infection. *PLoS Comput Biol*. 2021;17(4):e1008860.
doi:10.1371/journal.pcbi.1008860

33. Bertels LK, Fernández Murillo L, Heinisch JJ. The pentose phosphate pathway in yeasts-
more than a poor cousin of glycolysis. *Biomolecules*. 2021;11(5):725.
doi:10.3390/biom11050725

34. Parkhitko AA, Priolo C, Coloff JL, et al. Autophagy-dependent metabolic reprogramming
sensitizes TSC2-deficient cells to the antimetabolite 6-aminonicotinamide. *Mol Cancer
Res*. 2014;12(1):48-57. doi:10.1158/1541-7786.MCR-13-0258-T

35. Lin E, Kuo PH, Liu YL, Yang AC, Kao CF, Tsai SJ. Association and interaction of APOA5,
BUD13, CETP, LIPA and health-related behavior with metabolic syndrome in a Taiwanese
population. *Sci Rep*. 2016;6(1). doi:10.1038/srep36830

36. Cadena-López RO, Hernández-Rodríguez LV, Aguilar-Galarza A, et al. Association between
SNPs in Leptin pathway genes and anthropometric, biochemical, and dietary markers
related to obesity. *Genes (Basel)*. 2022;13(6):945. doi:10.3390/genes13060945

37. Foguet C, Xu Y, Ritchie SC, et al. Genetically personalised organ-specific metabolic models
in health and disease. *Nat Commun*. 2022;13(1):7356. doi:10.1038/s41467-022-35017-7

38. Gustafsson J, Anton M, Roshanzamir F, et al. Generation and analysis of context-specific
genome-scale metabolic models derived from single-cell RNA-Seq data. *Proc Natl Acad
Sci U S A*. 2023;120(6):e2217868120. doi:10.1073/pnas.2217868120

39. Huang Y, Mohanty V, Dede M, et al. Characterizing cancer metabolism from bulk and
single-cell RNA-seq data using METAFlux. *Nat Commun*. 2023;14(1):4883.
doi:10.1038/s41467-023-40457-w

40. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics*. 2016;32(19):3047-3048.
doi:10.1093/bioinformatics/btw354

41. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. *Nat Methods*. 2017;14(4):417-419.
doi:10.1038/nmeth.4197

42. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. *F1000Res*. 2015;4:1521.
doi:10.12688/f1000research.7563.1

43. Zhang Y, Parmigiani G, Johnson WE. ComBat-seq: batch effect adjustment for RNA-seq
count data. *NAR Genom Bioinform*. 2020;2(3):lqaa078. doi:10.1093/nargab/lqaa078

44. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-0550-8

45. Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. *Bioinformatics*. 2019;35(12):2084-
2092. doi:10.1093/bioinformatics/bty895

46. Bindea G, Mlecnik B, Hackl H, et al. ClueGO: a Cytoscape plug-in to decipher functionally
grouped gene ontology and pathway annotation networks. *Bioinformatics*.
2009;25(8):1091-1093. doi:10.1093/bioinformatics/btp101

47. Shannon P, Markiel A, Ozier O, et al. Cytoscape: a software environment for integrated
models of biomolecular interaction networks. *Genome Res*. 2003;13(11):2498-2504.
doi:10.1101/gr.1239303

48. Oliveira de Biagi CA Jr, Nociti RP, Brotto DB, et al. CeTF: an R/Bioconductor package for
transcription factor co-expression networks using regulatory impact factors (RIF) and
partial correlation and information (PCIT) analysis. *BMC Genomics*. 2021;22(1):624.
doi:10.1186/s12864-021-07918-2

49. Reverter A, Hudson NJ, Nagaraj SH, Pérez-Enciso M, Dalrymple BP. Regulatory impact
factors: unraveling the transcriptional regulation of complex traits from expression data.
Bioinformatics. 2010;26(7):896-904. doi:10.1093/bioinformatics/btq051

50. Teixeira MC, Viana R, Palma M, et al. YEASTRACT+: a portal for the exploitation of global
transcription regulation and metabolic model data in yeast biotechnology and
pathogenesis. *Nucleic Acids Res*. 2023;51(D1):D785-D791. doi:10.1093/nar/gkac1041

51. Watson-Haigh NS, Kadarmideen HN, Reverter A. PCIT: an R package for weighted gene co-
expression networks based on partial correlation and information theory approaches.
Bioinformatics. 2010;26(3):411-413. doi:10.1093/bioinformatics/btp674

52. Banerjee P, Balraj P, Ambhore NS, et al. Network and co-expression analysis of airway
smooth muscle cell transcriptome delineates potential gene signatures in asthma. *Sci Rep*.
2021;11(1):14386. doi:10.1038/s41598-021-93845-x

53. Lau LY, Reverter A, Hudson NJ, Naval-Sanchez M, Fortes MRS, Alexandre PA. Dynamics of
gene co-expression networks in time-series data: A case study in *Drosophila melanogaster*
embryogenesis. *Front Genet*. 2020;11:517. doi:10.3389/fgene.2020.00517

54. Lu H, Li F, Sánchez BJ, et al. A consensus S. cerevisiae metabolic model Yeast8 and its
ecosystem for comprehensively probing cellular metabolism. *Nat Commun*.
2019;10(1):3586. doi:10.1038/s41467-019-11581-3

55. Pavan Kumar S, Bhatt NP. Localgini: A method for harnessing inequality in gene expression
to improve the quality of context-specific models. *bioRxiv*. Published online 17 September
2023. doi:10.1101/2023.09.14.557840

56. Zur H, Ruppin E, Shlomi T. iMAT: an integrative metabolic analysis tool. *Bioinformatics*.
2010;26(24):3140-3142. doi:10.1093/bioinformatics/btq602

57. Heirendt L, Arreckx S, Pfau T, et al. Creation and analysis of biochemical constraint-based
models using the COBRA Toolbox v.3.0. *Nat Protoc*. 2019;14(3):639-702.
doi:10.1038/s41596-018-0098-2

58. Megchelenbrink W, Huynen M, Marchiori E. optGpSampler: an improved tool for
uniformly sampling the solution-space of genome-scale metabolic networks. *PLoS One*.
2014;9(2):e86587. doi:10.1371/journal.pone.0086587

59. Herrmann HA, Dyson BC, Vass L, Johnson GN, Schwartz JM. Flux sampling is a powerful
tool to study metabolism under changing environmental conditions. *NPJ Syst Biol Appl*.
2019;5(1):32. doi:10.1038/s41540-019-0109-0

60. Ebrahim A, Lerman JA, Palsson BO, Hyduke DR. COBRApy: COnstraints-Based
Reconstruction and analysis for Python. *BMC Syst Biol*. 2013;7(1):74. doi:10.1186/1752-
0509-7-

    FIGURE LEGENDS

Figure 1: Schematic representation outlines the key steps and methodologies employed in

our study

The raw transcriptomic data of 16 allele replacement panels were pre-processed and

corrected for batch effects using CombatSeq. Differentially expressed genes were obtained

using DeSeq2, and the critical regulators were obtained from RIF analysis and further studied

for connectivity using PCIT analysis. Integration of transcriptomic data into the genome-scale

metabolic model of yeast was performed using Localgini and IMAT to reconstruct SNP-specific

metabolic models. The change in intracellular fluxes between the wildtype and the SNP

models was studied using GS-DFA analysis. Figure 1 was created with BioRender.com.

Figure 2: Transcriptional differences between the allele replacement strains

(A) Number of upregulated and downregulated genes (LFC>0.5 and LFC<-0.5) identified using

DeSeq2 between wildtype vs. each SNP combination. (B) The number of differentially

expressed ((LFC > 0.5 and LFC < 0.5) sporulation-specific and metabolic genes identified using

DeSeq2 between wildtype vs. each SNP. (C) Venn diagram plot for upregulated genes

identified using DeSEq2 with LFC > 0.5 and p-adj > 0.1 for three SNP combination strains

(*RME1_RSF1_IME1*, *RME1_IME1_IME1nc*, *RSF1_IME1_IME1nc* and *RME1_RSF1_IME1nc*) and

four SNP combination strain (*RME1_RSF1_IME1_IME1nc*) strains in comparison with the

wildtype. (D) Venn diagram analysis for *RSF1* and *IME1* SNP combinations. (E) Heatmap

representing the LFC values of *IME1*, *RIM4*, and *REC8* genes across all SNP combination strains

compared to the wildtype strain

Figure 3: Analysis of connectivity of critical metabolic regulators

(A) Connectivity of critical metabolic regulators identified using RIF analysis across

combinations of SNPs. (B) Comparing the connectivity of critical metabolic regulators based

on the sporulation efficiency of the strains. Significance was calculated using a t-test (ns

represents non-significant).

Figure 4: SNPs and their interaction can activate unique reactions and subsystems

(A) Metabolic subsystems enrichment analysis for reactions that are over-represented in each

comparison that are present exclusively in single SNP models compared with the wildtype

model. (B) Synergistic effect of *RSF1* and *IME1* in modulating nucleotide metabolism. Here,

the violin plots represent the flux distribution obtained by optGpSampler for key reactions

involved in purine and pyrimidine metabolism compared across wildtype (M16), *RSF1+IME1*

(M8), *RSF1* (M13) and *IME1* (M14) models.

Figure 5: Pentose phosphate pathway is differentially regulated in SNP models

(A) Flux enrichment plots for unique reactions in highly sporulating strains compared to the

wildtype strain. (B) Flux enrichment plots for upregulated reactions in highly sporulating

strains compared to the wildtype strain obtained after GF-DFA analysis. (C) Box plots

representing the flux distribution of key reactions involved in the pentose phosphate pathway

obtained from optGpSampler for three and four SNP models (M1 to M5) compared to the

wildtype (M16) model.

 SUPPLEMENTARY METHODS

Supplementary methods accompany this manuscript.

SUPPLEMENTARY TABLE LEGENDS

Table S1: Differentially expressed genes identified using DESeq2 between wildtype and 15

allele replacement strains.

Table S2: The critical regulators identified by regulatory impact factor analysis. The metabolic

regulators are shown in bold.

Table S3: Number of nodes and edges in wildtype and SNP-specific networks before and after

filtering DEGs and RIFs. We could not construct networks for the strain with the IME1 variant

due to the limited number of replicates to perform PCIT analysis.

Table S4: Subsystems enriched for unique reactions in each SNP model compared to wildtype.

Table S5: Synergistic and antagonistic reactions and their subsystem enrichment for

RSF1+IME1 (M8) SNP combination.

Table S6: Synergistic and antagonistic reactions and their subsystem enrichment for

RME1+IME1nc (M9) SNP combination.

Table S7: List of upregulated reactions identified by GS-DFA analysis for all SNP models

compared with wildtype model.

Table S8: List of downregulated reactions identified by GS-DFA analysis for all SNP models

compared with wildtype model.

Table S9: Subsystems enriched for upregulated reactions in each SNP model compared to wildtype identified using GS-DFA analysis.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1: Clustering of differentially expressed sporulation-related genes based on LFC values for each allele replacement strain comparison with the wildtype strain.

Figure S2: Clustering of differentially expressed metabolic genes based on the LFC values for each allele replacement strain comparison with the wildtype strain.

Figure S3: Over-represented pathways of upregulated differentially expressed genes between wildtype and (A) *RME1_RSF1_IME1_IME1nc*, (B) *RME1_RSF1_IME1*, (C) *RSF1_IME1_IME1nc*, (D) *RME1_IME1nc* and (E) *IME1_IME1nc* strains. The circle size for a specific GO Term (pathway) varies with the order of significance (p-value) level. The most prominent node represents the most significant pathway. The pathways were constructed using Cytoscape v3.8.0 (https://cytoscape.org/) using the plugin ClueGO v2.5.41.

Figure S4: The Hamming distance between all the generated context-specific models based on the differences in reactions in the model.

Figure S5: Enriched subsystems (pathways) for the core reactions in all extracted SNP-specific models.

Figure S6: Metabolic subsystems enrichment analysis reveals the metabolic role of *IME1* functional and non-functional forms. (A) Flux enrichment analysis for reactions that are exclusively in each model in comparison with the wildtype (B) Flux enrichment analysis of upregulated reactions obtained using GS-DFA analysis between wildtype and *IME1*, *IME1nc*

and *IME1_IME1nc* strains. The presented reactions have an FC value (Flux Change) greater

than 0.8 (as described in materials and methods) and have an adjusted p-value less than 0.05.

Figure S7: Antagonistic effect of *RSF1* and *IME1* in modulating 'Steroid biosynthesis'. The

violin plots represent the flux distribution obtained by optGpSampler for key reactions

involved in **'Steroid biosynthesis'** compared across wildtype (M16), *RSF1_IME1* (M8), *RSF1*

(M13) and *IME1* (M14) models.

Figure S8: Synergistic effect of *RME1* and *IME1nc* in modulating 'one carbon pool by folate'

metabolism. The violin plots represent the flux distribution obtained by optGpSampler for

key reactions involved in **'one carbon pool by folate' metabolism** compared across wildtype

(M16), *RME1_IME1nc* (M9), *RME1* (M12) and *IME1c* (M15) models.

Figure S9: Hierarchical clustering of FC values of upregulated reactions in the M1 model across

other models shows two specific groups. The presented reactions have an FC value greater

than 0.8 in M1 models compared to the wildtype model (M16). The FC values have been

scaled for better representation.

Figure S10: (A) Enriched pathways (subsystems) in each SNP model were revealed by Flux

Variability analysis. The presented reactions have a flux span ratio (as described in materials

and methods) greater than two and have an adjusted p-value less than 0.05. (B) Clustering of

FSR values of the upregulated reactions specific for the M1 model that are compared across

all SNP models.

Figure S11: Proposed mechanism of autophagy as a pentose-phosphate pathway-dependent

compensatory mechanism during the early phase of yeast sporulation.

Figure 1: Schematic representation outlines the key steps and methodologies employed in our study

The raw transcriptomic data of 16 allele replacement panels were pre-processed and corrected for batch effects using CombatSeq. Differentially expressed genes were obtained using DeSeq2, and the critical regulators were obtained from RIF analysis and further studied for connectivity using PCIT analysis. Integration of transcriptomic data into the genome-scale metabolic model of yeast was performed using Localgini and IMAT to reconstruct SNP-specific metabolic models. The change in intracellular fluxes between the wildtype and the SNP models was studied using GS-DFA analysis. (created with BioRender.com)

Figure 2

Figure 2: Transcriptional differences between the allele replacement strains
(A) Number of upregulated and downregulated genes (LFC>0.5 and LFC<-0.5) identified using DeSeq2 between wildtype vs. each SNP combination. (B) Number of differentially expressed ((LFC > 0.5 and LFC < 0.5) sporulation-specific and metabolic genes identified using DeSeq2 between wildtype vs. each SNP. (C) Venn diagram plot for upregulated genes identified using DeSEq2 with LFC > 0.5 and p-adj > 0.1 for three SNP combination strains (*RME1_RSF1_IME1*, *RME1_IME1_IME1nc*, *RSF1_IME1_IME1nc* and *RME1_RSF1_IME1nc*) and four SNP combination strain (*RME1_RSF1_IME1_IME1nc*) strains in comparison with the wildtype. (D) Venn diagram analysis for *RSF1* and *IME1* SNP combinations. (E) Heatmap representing the LFC values of *IME1*, *RIM4*, and *REC8* genes across all SNP combination strains compared to the wildtype strain.

Figure 3

A

B

Figure 3: Analysis of connectivity of critical metabolic regulators
(A) Connectivity of critical metabolic regulators identified using RIF analysis across combinations of SNPs. (B) Comparing the connectivity of critical metabolic regulators based on the sporulation efficiency of the strains. Significance was calculated using a t-test (ns represents non-significant).

Figure 4

Figure 4: SNPs and their interaction can activate unique reactions and subsystems
(A) Metabolic subsystems enrichment analysis for reactions that are over-represented in each comparison that are present exclusively in single SNP models compared with the wildtype model. (B) Synergistic effect of *RSF1* and *IME1* in modulating nucleotide metabolism. Here, the violin plots represent the flux distribution obtained by optGpSampler for key reactions involved in purine and pyrimidine metabolism compared across wildtype (M16), *RSF1+IME1* (M8), *RSF1* (M13) and *IME1* (M14) models.

Figure 5

Differential enrichment of autophagy related reactions

Differential enrichment of Pentose phosphate related reactions

Figure 5: Pentose phosphate pathway is differentially regulated in SNP models (A) Flux enrichment plots for unique reactions in highly sporulating strains compared to the wildtype strain (B) Flux enrichment plots for upregulated reactions in highly sporulating strains compared to the wildtype strain obtained after GF-DFA analysis. (C) Box plots representing the flux distribution of key reactions involved in the pentose phosphate pathway obtained from optGpSampler for three and four SNP models (M1 to M5) compared to the wildtype (M16) model.