Title: Reinforcement learning during locomotion

Authors and affiliations:

Jonathan M Wood[1,2]

Hyosub E Kim[1-4]

Susanne M Morton[1,2]

1. Department of Physical Therapy, University of Delaware, Newark, DE 19713, United

States

2. Interdisciplinary Graduate Program in Biomechanics & Movement Science, University of

Delaware, Newark, DE 19713, United States

3. Department of Psychological and Brain Sciences, University of Delaware, Newark, DE

19716, United States

4. School of Kinesiology, University of British Columbia, Vancouver, BC, Canada

Corresponding Author: Susanne M Morton

Email: smmorton@udel.edu

Keywords: Reinforcement learning, motor learning, gait, reward, motor memory, variability

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: This work was supported by the National Institutes of Health, S10-

RR028114 (SMM), K12 HD055931(HEK), and the University of Delaware Graduate College

through the University Dissertation Fellowship (JMW). We also thank Saunders Penn for his

help with data collections and making Figure 1A and Maci Keaton for her help with data

collections.

Abstract

When learning a new motor skill, people often must use trial and error to discover which movement is best. In the reinforcement learning framework, this concept is known as exploration and has been observed as increased movement variability in motor tasks. For locomotor tasks, however, increased variability decreases upright stability. As such, exploration during gait may jeopardize balance and safety, making reinforcement learning less effective. Therefore, we set out to determine if humans could acquire and retain a novel locomotor pattern using reinforcement learning alone. Young healthy male and female humans walked on a treadmill and were provided with binary reward feedback (success or failure only) to learn a novel stepping pattern. We also recruited a comparison group who walked with the same novel stepping pattern but did so by correcting for target error, induced by providing real time veridical visual feedback of steps and a target. In two experiments, we compared learning, motor variability, and two forms of motor memories between the groups. We found that individuals in the binary reward group did, in fact, acquire the new walking pattern by exploring (increased variability). Additionally, while reinforcement learning did not increase implicit motor memories, it resulted in more accurate explicit motor memories compared to the target error group. Overall, these results demonstrate that humans can acquire new walking patterns with reinforcement learning and retain much of the learning over 24 hours.

Significance Statement

Humans can learn some novel movements by independently discovering the actions that lead to success. This discovery process, exploration, requires increased motor variability to determine the best movement. However, in bipedal locomotion especially, increasing motor variability decreases stability, heightening the risk of negative outcomes such as a trip, injury, or fall. Despite this stability constraint, the current study shows that individuals do use exploration to find the most rewarding walking patterns. This form of learning led to improved explicit retention

53   but not implicit aftereffects. Thus, the reinforcement learning framework can explain findings

54   across a wide range of motor and cognitive tasks, including locomotion.

Introduction

In the 2020 Summer Olympics, U.S. athlete Sydney McLauchlin broke the world record and won

the gold medal in the 400-meter hurdles, aided by her proficiency in jumping over the hurdles

equally well with both her dominant and non-dominant legs. While reward (clearing the hurdle)

is ostensibly a critical learning signal for this and other tasks, its specific role in locomotor

learning remains unknown. Rewards influence learning via reward prediction errors, the

difference between expected and actual rewards (Rescorla and Wagner, 1972). These error

signals form the basis for reinforcement learning theory, a widely used framework describing

how animals and machines learn the values of actions (Sutton and Barto, 2017). A core

component of reinforcement learning is exploration, defined as sampling from a range of

possible actions to update their values. In movement-based reinforcement learning, exploration

can be characterized as motor variability, where greater motor variability indicates more

exploration (Dhawale et al., 2017). However, in locomotion, exploration could be problematic,

since more variable gait patterns reduce stability (McAndrew Young and Dingwell, 2012), and, if

left unchecked, could lead to catastrophic consequences (e.g., loss of balance, injuries, falls).

During motor tasks, reinforcement learning is often probed by providing only binary reward

feedback, and has been demonstrated in reaching (Izawa and Shadmehr, 2011), speech

production (Parrell, 2021), and saccadic eye movements (Madelain et al., 2011). Since binary

reward feedback does not provide information about the direction or distance from a task goal,

the only signal by which to learn the correct movement is a reward prediction error, resulting in

exploratory behavior. In sharp contrast, learning signals such as target error (i.e., the difference

between a task goal and movement feedback) and sensory prediction error (i.e., the difference

between the expected and actual sensory consequences of movements), convey both

magnitude and directional information. As a result, they produce comparatively less exploration

than reward prediction errors (Izawa and Shadmehr, 2011; Therrien et al., 2016; Cashaback et

al., 2019). To date, no studies have determined if binary reward feedback can produce learning,

and the expected exploratory behavior, during locomotion.

Reinforcement learning also impacts motor memories. Implicit aftereffects, defined as motor

memories not under conscious control (Krakauer et al., 2019), are strengthened when

reinforcement is combined with sensorimotor adaptation (Huang et al., 2011; Shmuelof et al.,

2012; Galea et al., 2015), or use-dependent learning (Mawase et al., 2017; Bao and Lei, 2022;

c.f., Tsay et al., 2022). Reinforcement learning may also strengthen explicit retention, defined as

the ability to consciously remember and reproduce a previously learned movement (Schmidt

and Lee, 2005). However, previous studies examining explicit retention in reinforcement

learning either did not induce learning with purely binary reward feedback (Hasson et al., 2015;

Codol et al., 2018), or did not include a comparison group (Holland et al., 2018). Therrien et al.

(2016) found retention was better after binary reward feedback than sensory prediction and

target errors, but this probe into retention may have also included a strong implicit component.

Therefore, the extent to which reinforcement learning alone contributes to implicit and explicit

motor memory formation remains unclear.

The primary objective of this study was to determine if individuals can acquire a novel gait

pattern using reinforcement learning. We tested this by providing binary reward feedback to

participants learning to walk with a longer left step length than baseline. For comparison, we

recruited a group that performed the same walking pattern using explicit, target error feedback,

which does not induce significant exploration. We previously showed that use-dependent

learning is the predominant source of implicit aftereffects in this target error paradigm (Wood et

al., 2020, 2021), and neither implicit aftereffects nor explicit retention is contaminated by

sensory prediction errors. We hypothesized that the reward prediction error group would learn

the new walking pattern using greater exploration, and that they would have larger implicit

aftereffects and better explicit retention compared to the target error group.

Material and Methods

Participants

A total of 59 (34 female, 25 male) young, healthy adults were recruited from the University of

Delaware community. Within 2 separate experiments, participants were randomized into 1 of 2

groups: a reward prediction error (RPE) group and a target error (TE) group. Participants were

excluded if they had any orthopedic or neurologic conditions that might interfere with their ability

to learn a new walking pattern. Five participants were removed from the analysis (2 due to

technical errors, 2 due to not following instructions, and 1 who was provided an erroneous

verbal reminder of the instructions during learning). Thus, thirty participants were included in

experiment 1 and 24 in experiment 2. Before participating, all individuals signed a written,

informed consent document. This study was approved by the University of Delaware

Institutional Review Board.

Experimental design

Feedback

Participants walked on a dual belt treadmill (Figure 1A; Bertec, Columbus OH) at a self-

selected, comfortable but brisk pace, constrained between 1.0 and 1.2 m/s. The belts moved at

the same speed throughout the experiment. Participants wore a ceiling mounted harness, which

did not provide body weight support, and lightly held onto a handrail for safety while walking. A

computer monitor placed in front of the treadmill provided real-time feedback of participants' left

step lengths (The Motion Monitor Toolbox, Innovative Sports Training Inc., Chicago IL). We

chose to focus on step length because it is a robustly modifiable gait parameter that has

frequently been the focus of locomotor learning paradigms (Reisman et al., 2005; Roemmich et

al., 2016; French et al., 2018, 2021; Wood et al., 2020, 2021), and because both implicit and

explicit forms of motor memory have been observed after learning an altered step length pattern

(French et al., 2018, 2021; Wood et al., 2020, 2021).

137
Figure 1: Experimental paradigm. (A) All participants walked on a dual-belt treadmill, at a
comfortable self-selected pace with a computer monitor in front of them. **(B)** The RPE and TE
groups received different feedback during the learning phase. The RPE group received only
binary reward feedback, with a check mark and money added to a total when they performed a
correct step length. The TE group received real-time feedback of their left step length related to
the pink, horizontal target line. **(C)** All participants walked in 3 different phases: 1) baseline, where
individuals were asked to walk normally; 2) learning, where they gradually learned to walk with a
longer left step length with the feedback and instructions depending on group assignment; 3) post-
learning, where implicit aftereffects (experiment 1) or explicit retention (experiment 2) were
probed, both without visual feedback. Participants in experiment 2 also were tested for explicit
retention 24 hours later (not shown).
149

We compared a group that received reward prediction error feedback (RPE group) to a group

that received explicit task instructions and visual target errors (TE group). The TE group

received real-time visual feedback of their step length relative to a step length target (Figure 1B,

top row). The left step length appeared as a blue bar, starting with a height of '0' and growing

vertically during the swing phase, then holding position on the screen at its final height at the

time of heel strike, until the next swing phase began. The TE group was instructed in the

definition of a step length and that they needed to hit a pink horizontal target line exactly with

the blue bar that represented their left step length. They were also informed how the target left

step length would change during learning. Although they could see if they hit the target line, no

external reward was provided either on the screen or verbally by the experimenter when they

successfully hit the target. Thus, since the TE group knew exactly where to step and could

correct for any target errors throughout the task, they had no need to explore. Furthermore,

since the feedback was veridical, sensory prediction errors could not contaminate either implicit

aftereffects or explicit retention.

The RPE group was provided with only binary reward feedback (Figure 1B, bottom row). This

feedback took the form of a large green check mark and money added to a total when a left step

length fell within the target window, the same target as the TE group, but here, invisible to

participants. If successful, the check mark appeared on the screen at heel strike and remained

on the screen until the next swing phase began. When the left step length fell outside of the

target window, no check mark appeared, and no money was added to the total. A running tally

of the total reward earned was consistently displayed at the top of the screen during the learning

phase. The RPE group was told that their goal was to gain as much money as possible and that

their compensation for participation in the study would depend on their performance, but in

reality, all individuals were compensated the same. As awareness of the rewarded parameter

may be important for movement based reinforcement learning (Manley et al., 2014), individuals

in the RPE group were told that the reward was contingent on their left step length. The

experimenter then explained and demonstrated a left step length until the participant could

explain the concept themselves. Based on pilot testing, we also told the participants in the RPE

group to start off by walking normally so they could achieve some success at the beginning of

the learning phase, but critically, no information was provided regarding if or how the rewarded

step length would change. Thus, to learn the longer left step length, the RPE group was

provided with no other learning signal except a reward prediction error, the difference between

the expected and actual reward. Participants in both groups were provided with a brief, verbal

reminder of their instructions every 100 strides (TE: "hit the target"; RPE: "gain as much money

as possible").

Experimental schedules

Participants in both groups walked under three different phases: baseline, learning and post-

learning (Figure 1C). During baseline walking (250 strides) participants were asked to walk

normally and no visual feedback was provided. Next, participants performed a learning phase

(900 strides), where they gradually learned to lengthen their left step length by 10% from their

baseline using the feedback displayed (step length bars and target lines for the TE group or

check marks and tally of money earned for the RPE group) on the computer monitor. While the

feedback was different between the groups, the learning schedule was the same. For the first

50 strides of the learning phase, the target was centered on each individual's baseline left step

length. The target step length then increased by 1% every 10 strides until it reached its final

10% change from baseline where it remained for the rest of learning (strides 141 to 900). The

target maintained a ±2 cm margin of error throughout the learning phase.

The post-learning phases were used to probe either implicit aftereffects (experiment 1) or

explicit retention (experiment 2). Therefore, the visual feedback was turned off during this phase

for both experiments, but the instructions were slightly different. In experiment 1, participants

completed a washout phase (900 strides) where they were asked to look forward and "walk

normally". Any left step length differences from baseline during this phase represented the

implicit aftereffect, a motor memory not under conscious control. In experiment 2, participants

completed two explicit retention tests, one immediately after learning (25 strides) and one 24-

hours later (250 strides). For both tests, participants were instructed to remember how they

walked at the end of the learning phase and repeat that walking pattern exactly. Thus, explicit

retention measured the ability to consciously remember and reproduce the previously learned

step length pattern. For both experiments, participants in the RPE group were told that no

additional money could be earned during these periods.

Data collection

Step length was calculated in real time using The Motion Monitor which received kinematic and

kinetic data from Nexus software (Vicon Motion Systems Inc., London, UK). Kinetic data,

sampled at 1000 Hz, was acquired from the instrumented force plates through each treadmill

belt (Bertec, Columbus OH), and kinematic data, sampled at 100 Hz, was acquired from an 8-

camera Vicon MX40 motion capture system with Nexus software. To capture the kinematic

data, we used a custom marker set with seven retroreflective markers, one on each heel, lateral

malleolus, and fifth metatarsal head, and one on the left first metatarsal. The heel markers were

used to determine left step length by calculating the difference along the laboratory's y-axis

(anterior-posterior dimension) between the left and the right heel markers at the moment of left

heel strike. Left heel strike was determined in real time by detecting when there was a ground

reaction force on the left belt and the velocity of the left heel marker first transitioned from

positive to negative (Zeni et al., 2008).

Data analyses

After data were collected, we used custom written MATLAB scripts (v2022a The MathWorks,

Natick, MA) to calculate and plot the outcome variables. The primary outcome measure was

percent left step length change from baseline (ΔLSL) calculated on each stride (s):

$\Delta LSL_s = \frac{LSL_s - LSL_{baseline}}{LSL_{baseline}} * 100\%$ (1)

$LSL_{baseline}$ was calculated as the mean left step length during the final 50 strides of the baseline

phase. Key time periods used for analysis during learning (for both experiments) included: early

learning, the mean of the first 50 strides after the target reached its plateau (10% change) and

late learning, the mean of the last 50 strides of the learning phase.

We tested the hypothesis that the RPE group could learn a new gait pattern by averaging ΔLSL

during late learning. Learning in the RPE group would be reflected in a ΔLSL reliably greater

than 0. In addition, to observe how learning emerged in both groups over time, we calculated

two secondary outcome measures during early and late learning. Error was calculated as the

absolute difference between ΔLSL and the center of the ΔLSL target (10%), and success was

calculated as the percentage of left step lengths that fell within the target step length window.

In stable reward environments (in the current study, once the target reached its plateau),

exploration should decrease as the values of actions are learned (Daw et al., 2006; Sutton and

Barto, 2017). Based on this theory and our assumption that the TE group correcting for target

error required very little exploring, we made two predictions: 1) During early learning, the RPE

group would demonstrate greater ΔLSL variability than the TE group; and 2) The RPE group

would reduce their ΔLSL variability by a greater magnitude from early to late learning compared

to the TE group. We defined variability of ΔLSL as the standard deviation of ΔLSL, normalized

253 by the standard deviation of baseline ΔLSL. We denote this measure as $\sigma_{\Delta LSL}$. These two

254 predictions served as our test of the hypothesis that the RPE group would demonstrate greater

255 exploration during learning compared to the TE group.

257 Exploration can also be characterized using a win-stay / lose-shift model. When an individual

258 performs unsuccessful actions, the trial-to-trial changes tend to be more variable than after

259 successful actions (Pekny et al., 2015; Cashaback et al., 2019; Uehara et al., 2019). Therefore,

260 using all of the learning phase data except for the first 50 strides (i.e., before the target started

261 shifting), we performed this trial-to-trial change analysis secondarily to our main motor variability

262 analysis. We calculated the difference in ΔLSL values between subsequent strides separately

263 after successful steps and unsuccessful steps. The trial-to-trial change measure was calculated

264 as the standard deviation of these changes, normalized by baseline trial-to-trial change

265 variability (Cashaback et al., 2019; Roth et al., 2023). We denote this measure as $\sigma_{trial-to-trial}$.

267 For implicit aftereffects and explicit retention, we controlled for any potential differences in

268 learning by calculating both as a percentage of the learned gait pattern for each individual

269 participant on each stride. For experiment 1, we calculated immediate percent implicit aftereffect

270 as the mean of the first 5 strides of the washout phase (immediate washout), and early percent

271 implicit aftereffect as the mean of the next 25 strides (early washout). Therefore, we compared

272 the implicit aftereffects at both timepoints between groups to test the hypothesis that the RPE

273 group would show larger implicit aftereffects than the TE group. For experiment 2, we calculated

274 the percent explicit retention at two different timepoints: the mean of all 25 strides during

275 immediate retention and the mean of the first 25 strides during 24-hour retention. Here, we

276 considered 25 strides so that individuals had time to achieve a consistent level of performance

277 that reflected their explicit motor memory. For experiment 2, it was also necessary to calculate

278 retention error, the absolute difference between ΔLSL at late learning and ΔLSL at immediate

and 24-hour retention. This additional measurement of error, or accuracy, was necessary

because, unlike for implicit motor memories, we expected that explicit motor memories could

either overshoot or undershoot the remembered target location, but the goal was to reproduce

the previously learned stepping pattern as accurately as possible. Thus, retention error at both

timepoints served as our test for the hypothesis that explicit retention would be more accurate

for the RPE group compared to the TE group.

Statistical analysis

We used Bayesian statistics to make inferences regarding differences within and between

groups. This allowed us to estimate the specific probability that our hypotheses were supported

or not, given the data we collected as opposed to the classical p value, which provides the

probably of obtaining the data, assuming the null hypothesis was true. We assumed data were

generated from student's t distributions with a mean and standard deviation that varied based

on both between (group) and within (time) subject factors. We estimated the mean and standard

deviation parameters by combining our prior assumptions regarding their values with evidence

from the data we collected. We set wide and uninformative priors for each of the model

parameters similar to the approached described by Kruschke (2014). We calculated the joint

posterior probability distribution with the Python package bayes-toolbox (Kim, 2023) which uses

the Pymc library (v4.3 Salvatier et al., 2016) in Python (version 4.3.0). The joint posterior

distribution was estimated using Markov Chain Monte Carlo (MCMC) sampling, with 10,000

draws in 4 chains (i.e., 40,000 total samples) and 2,000 tuning samples in each chain. We

performed diagnostics for each model, ensuring parameter values were consistent across

chains and checking posterior estimates for possible errors (Kruschke, 2014; McElreath, 2016).

We made inferences regarding differences between groups based on posterior contrast

distributions: the difference between two posterior distributions of interest. When these contrasts

represent a between group difference at a specific timepoint we use the term "group difference",

and when they represent differences across time between groups (analogous to an interaction

effect) we use the term "slope difference". We report the mean difference of each posterior

contrast distribution along with the 95% high density interval (HDI), defined as the narrowest

span of credible values that contain 95% of the posterior contrast distribution (Kruschke, 2014).

We also report the probability of a difference given the data we collected as a percentage of the

posterior contrast distribution that lies on one side of 0 (e.g., 85.6% > 0).

Code Accessibility

The data analysis code for individual participants and figure generation, as well as the full

statistical models and code, including detailed information about the priors and diagnostics are

available online at Open Science Framework [https://osf.io/tcwu6/].

Results

In experiment 1, 15 individuals (9 females; mean age ± 1 SD, 22.8 ± 4.2 years) were

randomized into the RPE group and 15 (10 females; 22.5 ± 3.7 years) were randomized into the

TE group. In experiment 2, a new sample of individuals was recruited, with 12 (6 females; 23.5

± 4.6 years) randomized into the RPE group and 12 (7 females; 22.3 ± 4.2 years) randomized

into the TE group.

Experiment 1

Individual performance in the RPE group was idiosyncratic

Qualitatively, individuals in the TE group performed consistently across the learning phase

(Figure 2, top row), which is not surprising given they received accurate visual feedback of their

step length and the target. Individuals in the RPE group, however, were highly variable at

different points of the learning phase (Figure 2, bottom row). For example, some RPE

individuals (bottom left) were highly variable after the target reached its plateau, then became

less variable as the learning phase progressed. Others (bottom middle) were more variable at

the beginning of the learning phase and struggled to be consistently successful until the very

end of the learning phase. Still others (bottom right) were quite consistent and even appeared to

perform similarly to individuals in the TE group, despite having no visual feedback of their step

length or the target. Overall, the individual performance of the RPE groups was more

idiosyncratic than the TE group.

339
340   **Figure 2: Individual participant data.** ΔLSL data for all strides of the learning phase from 6
341   exemplar subjects; 3 each from the TE and RPE groups. Blue dots represent unsuccessful steps,
342   orange dots represent successful steps. The dashed lines represent the target window, centered
343   on 10% ΔLSL (the width of the window was ±2 cm for all participants).
344

The RPE group learned the novel locomotor pattern

Despite the idiosyncratic behavior of the RPE group, overall, they learned the new walking

pattern by the end of the 900-stride learning phase (Figure 3A). The RPE group's ΔLSL was

much larger than baseline (Figure 3B; RPE late learning, mean [95% HDI] = 9.16% [8.36 9.95],

100% > 0), providing a high degree of certainty that the RPE group learned the new walking

pattern. The TE group's ΔLSL at late learning was slightly greater by comparison (group

difference = -0.94% [-1.99 0.02], 96.8% < 0). Additionally, the RPE group's performance was

greatly improved from early to late learning compared to the TE group (Figure 3C and D).

Specifically, compared to the TE group, the RPE group had much larger improvements from

early to late learning in terms of error (slope difference = 1.22% [-0.07 2.47], 97.1% > 0) and

percent success (slope difference = -12.0% [-23.6 0.8], 97.1% < 0). Together, these findings

indicate that the RPE group learned the novel gait pattern, though not as completely as the TE

group, despite the RPE group demonstrating larger reductions in error and greater increases in

percent success from early to late learning.

360
Figure 3. Experiment 1 learning and implicit aftereffects. (A) Group average ΔLSL data shown
over all strides of experiment 1. Solid colored lines represent group means and shading
represents 1 SEM. Gray regions denote the times when no feedback was provided and individuals
were asked to "walk normally". The dashed line represents the center of the target. Rectangular
boxes denote the key timepoints of early learning, late learning, and initial and early washout. **(B)**
Group averaged ΔLSL at late learning for both groups. Thick horizontal lines represent group
means; dots represent individuals; error bars represent ±1 SEM. **(C)** Group average ΔLSL error
at early and late learning. Thick lines represent group means; thin lines represent individuals. **(D)**

Group average percent success at early and late learning. **(E)** Group average implicit aftereffects.
Thick horizontal lines represent group means; dots represent individuals; error bars represent ±1
SEM.
372

The RPE group explored to learn the novel gait pattern

The variability analysis supported the hypothesis that individuals in the RPE group explored to

learn the new gait pattern. We found differences in $\sigma_{\Delta LSL}$ between the RPE and TE groups

during the learning phase (Figure 4A and B). Specifically, $\sigma_{\Delta LSL}$ was much greater during early

learning for individuals in the RPE group compared to individuals in the TE group (group

difference = 1.20 [-0.02 2.57], 97.8% > 0), and the $\sigma_{\Delta LSL}$ decreased more from early to late

learning for the RPE group compared to the TE group (slope difference = 0.99 [-0.27 2.31],

94.1% > 0). These two results provide strong evidence for the hypothesis that the RPE group

explored to learn the new walking pattern.

383
Figure 4. Motor variability. (A) Motor variability across the learning phase, measured as the
standard deviation of ΔLSL. For visualization purposes only, we calculated each individual's motor
variability in 18 bins of 50 strides each across the learning phase, then averaged motor variability
among individuals in each group (solid lines) with the shading representing 1 SEM. The dashed
rectangle represents the bins when the target was gradually shifting toward 10%. **(B)** Early and
late motor variability. We calculated ΔLSL variability at early and late learning timepoints. Thick
lines represent group means; thin lines represent individuals. **(C)** Motor variability measured as

the standard deviation of trial-to-trial changes after successful and unsuccessful steps ($\sigma_{trial-to-trial}$).
Thick lines represent group means; thin lines represent individuals.
393

In addition to overall motor variability at the two specific timepoints, we also performed a

secondary analysis of exploration, measured as the variability of trial-to-trial changes after

unsuccessful trials across most of the learning phase (Figure 4C). The RPE group had slightly

greater $\sigma_{trial-to-trial}$ after unsuccessful steps compared to the TE group (group difference = 0.37 [-

0.17 1.01], 90.8% > 0). And the difference between $\sigma_{trial-to-trial}$ for successful vs unsuccessful

steps was also slightly larger for the RPE group compared to the TE group (slope difference =

0.31 [-0.25 0.97], 84.6% > 0). Overall, while less reliable than the motor variability differences

(probabilities of ~91% and 85%), these findings provide additional support for the hypothesis

that the RPE group was exploring during locomotion.

Reinforcement learning did not increase the magnitude of implicit aftereffects

We probed implicit aftereffects by asking individuals to walk normally during the no-feedback

washout phase. Implicit aftereffects were greater for the TE group during initial washout, (group

difference = -10.7% [-23.8 2.8], 94.9% < 0), and during early washout (group difference = -9.6%

[-17.4 -1.9], 98.9% < 0). Therefore, contrary to our hypothesis, learning from reward prediction

errors did not increase implicit aftereffects more than learning from target error, rather, the

evidence strongly supported the opposite. This finding could reflect a reduced amount of use-

dependent learning for the RPE group caused by the increased variability (Wood et al., 2020,

2021).

<u>Experiment 2</u>

Learning from reward prediction errors versus target errors resulted in unique signatures of

explicit retention

In experiment 2, we compared explicit retention after learning from reward prediction errors and

target errors. This was done with a new sample of individuals who experienced a different post-

learning phase but the same baseline and learning phases. We replicated the main findings

from experiment 1 (see code for the remainder of the analyses). The RPE group learned (RPE

ΔLSL at late learning = 8.74% [7.82 9.66], 100% > 0), and they did so by exploring (group $\sigma_{\Delta LSL}$

difference at early learning = 0.87 [0.07 1.65], 98.6% > 0; $\sigma_{\Delta LSL}$ slope difference = 0.54 [-0.24

1.37], 89.9% > 0). After learning, both groups performed a test of explicit retention where they

were instructed to repeat the same walking pattern they performed at the end of learning without

feedback, both immediately after learning and 24 hours later (Figure 5A).

427
Figure 5. Experiment 2 explicit retention. (A) Group average ΔLSL data shown over all
strides of experiment 2. Solid colored lines represent group means and shading represents 1
SEM. Gray regions denote the times when no feedback was provided. During retention testing,
participants were instructed to "walk like you did at the end of the previous phase". The dashed
line represents the center of the target during the learning phase. **(B)** Group average ΔLSL
percent error data for each stride during the immediate and 24-hour retention timepoints
(rectangles represent the 25 strides of each epoch). 0 represents perfect retention. The inset
shows the group average (horizontal lines) and individual (dots) retention levels for the two

timepoints. Error bars represent 1 SEM. **(C)** Group average percent retention data for each
stride during the immediate and 24-hour retention timepoints. All shown in the same manner as
in **(B)**. The dashed line at 100% represents perfect retention.
439

First, we assessed the accuracy of the explicit motor memories by comparing the absolute error

between the ΔLSL behavior at the end of learning and during the retention phases (Figure 5B).

Recall that this is the primary outcome measure of explicit retention because it reflects the true

accuracy of retention, or how far each individual's steps were from the previous day's

remembered target, regardless of direction. The RPE group had less error (better accuracy) at

immediate retention compared to the TE group (group difference = -1.36% [-2.94 0.20], 95.8% <

0), but after 24-hours, the RPE group was very similar to the TE group (group difference = -

0.13% [-2.01 1.68], 56.5% < 0). Thus, the experiment 2 results revealed that learning a new gait

pattern using reward prediction errors produces a unique pattern of explicit retention compared

to correcting for target error that is, at least initially, more accurate.

We found large between group differences in the specific walking pattern that was remembered

both immediately and 24-hours after learning (Figure 5C). Specifically, percent explicit retention

was much larger for the TE group at both immediate (group difference = -44.4% [-63.3 -24.9],

100% < 0) and 24-hour retention (group difference = -45.8% [-71.4 -20.0], 99.8% < 0) compared

to the RPE group. Recall that here, "larger" retention does not equate to "better" retention

because we asked individuals to walk "exactly" like they did at the end of learning. Rather, this

result reflects the fact that the TE group performed a step length that was longer than what they

performed at the end of learning (i.e., over-stepping), resulting in percent retention values

mostly >100% (9/12 and 8/12 for Immediate and 24-hour retention, respectively). On the other

hand, the RPE group performed a step length that was shorter than what they performed at the

end of learning (i.e., under-stepping), creating in percent retention values mostly <100% (8/12

and 10/12 for Immediate and 24-hour retention, respectively).

Explicit retention resulted from differences in declarative knowledge of the successful step

length

Why did the RPE and TE groups have such distinct patterns of percent explicit retention? One

possibility is that declarative knowledge of the successful step length impacted explicit retention.

To investigate this, we used our post-session interviews after the 24-hour retention phase to

probe declarative knowledge of successful step lengths. We asked each individual to rate the

length of the successful step length compared to their baseline left step into one of three

categories: "not longer", "slightly longer", or "longer". The proportions of individuals in each

category were as follows: Not longer: TE 0%, RPE 25%; Slightly longer: TE 50, RPE 50%;

Longer: TE 50%, RPE 25%. On average, the TE group rated the successful step length as

longer (by about 1 category) than the RPE group (group difference = -1.00 [-2.00 -0.02], 97.7%

< 0). Thus, the different feedback and instructions led to differences in declarative knowledge of

the successful step length.

Discussion

This works shows, for the first time, that despite the stability constraints inherent in gait, young

healthy individuals can learn a new walking pattern using reward prediction errors. Consistent

with the reinforcement learning framework, individuals in the RPE group learned the most

valuable action by initially exploring the action space, assigning values to possible step lengths,

then as learning progressed, exploiting the learned action/value pairs with more consistent,

correct step lengths. We also found that reinforcement learning produced smaller magnitudes of

implicit aftereffects, but more accurate explicit retention, than individuals provided with target

errors and a specific task strategy, an effect that waned by 24 hours.

<u>Reinforcement learning during locomotion is accomplished by exploring</u>

Prior studies of locomotor learning have not isolated reward prediction errors by providing only

binary reward feedback. Rather, they either added rewards to other teaching signals like target

error (Sato et al., 2022) or sensory prediction errors (Bakkum and Marigold, 2022), or used

graded reward feedback, increasing the reward as movements became closer to a target

(Hasson et al., 2015). While graded reward feedback provides a reward prediction error, having

any knowledge about distance or direction from the target adds another learning signal similar

to target error. We therefore provided only binary reward feedback in the current study to isolate

reinforcement learning as much as possible. The RPE group learned the new walking pattern, a

finding consistent with prior studies in reaching (Izawa and Shadmehr, 2011; Therrien et al.,

2016; Cashaback et al., 2017; Holland et al., 2018), eye movements (Madelain et al., 2011),

and speech production (Parrell, 2021). Unlike a prior study in reaching (Wu et al., 2014),

baseline variability was not related to the rate of learning for the RPE groups across both

experiments (slope = 0.5 [-2.8 3.8], 63.9% > 0), but our ability to assess the rate of learning

could be hindered by the gradual nature of the perturbation in the current study.

To learn the new walking pattern, the RPE group explored the action space. We observed a

classic signature of reinforcement learning behavior, the exploration / exploitation trade-off, in

the RPE group's step length variability. Interestingly, prior reinforcement-based motor learning

studies in reaching have not observed a reduction in exploratory behavior over time in stable

reward environments. We attribute this to the relatively short (360 – 400 trials, Pekny et al.,

2015; Uehara et al., 2019), or absent (Izawa and Shadmehr, 2011; Therrien et al., 2016;

Cashaback et al., 2017) steady state learning phases in those studies. However, in the current

study, steady state learning was 760 strides which allowed the RPE group adequate time to

learn the reward landscape well enough to exploit the rewarding step length.

While parsimonious, motor variability is not the only way to characterize exploration in

movement-based reinforcement learning experiments. Some methods have characterized

exploration as a random walk (Haith and Krakauer, 2014; Roth et al., 2023), but most methods

have focused on the magnitude or variability of trial-to-trial of changes after successful vs

unsuccessful movements (i.e., win-stay lose shift methods; Pekny et al., 2015; Therrien et al.,

2016; Cashaback et al., 2019; Uehara et al., 2019; van Mastrigt et al., 2020, 2021). However,

no studies have compared the latter measure of exploration while learning from reward

prediction error versus correcting for target errors. While we found that the RPE group had

slightly greater trial-to-trial variability after unsuccessful steps compared to the TE group, the

effect was relatively small compared to the effect in overall motor variability. It is possible other

models of exploration, for example, random walk behavior (Roth et al., 2023), may better

distinguish between reward prediction error learning and target error correction. Or perhaps

these methods of exploration could be task dependent. Future work could elucidate different

methods of movement-based exploration similar to the distinct modes of exploration described

in decision making tasks (Wilson et al., 2014).

<u>Reinforcement learning does not increase the magnitude of implicit aftereffects</u>

Reinforcement learning did not increase implicit aftereffects as we expected, and the RPE group

actually produced smaller implicit aftereffects compared to the TE group which was not provided

with any overt reward. We suspect the implicit aftereffects in the TE group were caused by use-

dependent learning because the feedback was similar to prior studies in our lab and the implicit

aftereffect persisted throughout the 900-stride washout phase (Wood et al., 2020, 2021). It is

possible that implicit aftereffects in the RPE group were also caused by use-dependent learning

as suggested previously (Codol et al., 2018; Holland et al., 2018). If so, the current results

contrast somewhat with findings in sensorimotor adaptation (Huang et al., 2011; Shmuelof et al.,

2012; Galea et al., 2015; Therrien et al., 2016) and use dependent learning (Floel et al., 2008;

Mawase et al., 2017) in which reward boosts implicit aftereffects caused by other learning

processes. However, the current results align with work suggesting that increased movement

variability attenuates use-dependent learning (Verstynen and Sabes, 2011; Marinovic et al.,

2017; Wood et al., 2021; Tsay et al., 2022). Indeed, when we match individuals across groups

based on motor variability, the difference in implicit aftereffects disappears (n = 6 per group;

initial washout difference = -2.3% [-62.1 61.0], 53.9% < 0; early washout difference = -6.1% [-

59.0 47.8], 60.5% < 0). Therefore, we suggest implicit aftereffects in both groups were due to

use-dependent learning since, after controlling for motor variability, both groups demonstrated a

similar magnitude of implicit aftereffects.

<u>Reinforcement learning improves the accuracy of explicit retention</u>

Despite less explicit knowledge of the task itself, including the mapping between step length and

target position, the RPE group had more accurate immediate explicit retention compared to the

TE group. Better explicit retention accuracy for the RPE group could have been caused by the

reward itself (money and green checkmark) by solidifying the memory of the walking pattern

better than simply hitting a target without overt reward. This explanation is similar to Thorndike's

Law of Effect (Thorndike, 1927), and is consistent with studies showing that rewarding

successful movements during skill learning and sensorimotor adaptation result in better motor

memories than without reward (Abe et al., 2011; Shmuelof et al., 2012; Therrien et al., 2016;

Mawase et al., 2017). Another possibility is that the act of exploring produces a clearer picture

of the successful action(s) needed within the workspace, which then leads to better, more

accurate explicit retention. Studies across various learning tasks suggest that greater variability

produces worse performance during training but better memories/generalization (Raviv et al.,

2022). A recent study in a grid-sailing, key press task suggested that more variable training

leads to better knowledge of novel action outcome pair mappings (Velázquez-Vargas et al.,

2023). Therefore, it is possible that by exploring the space, individuals in the RPE group

developed better knowledge of the action outcome pair.

<u>Reinforcement learning and correcting for target errors produce distinct explicit motor memories</u>

Interestingly, we found consistent and unique biases in explicit motor memories between groups

when assessed as percent explicit retention, a measure that takes the direction of error into

account. During both explicit retention phases, the RPE group was biased to step shorter than

they did at the end of the learning phase, while the TE group was biased to step longer than

they did at the end of the learning phase. Participants' declarative memories of the successful

step length followed a similar pattern (i.e., more individuals in the TE group reported stepping

"longer"), suggesting their bias was likely cognitive in nature. While the link between declarative

knowledge and different forms of motor learning and memory has been made previously

(Shanks and Johnstone, 1999; Keisler and Shadmehr, 2010; Stanley and Krakauer, 2013; Haith

et al., 2015; Wong et al., 2015; McDougle et al., 2022), we are not aware of any previous such

stark display of distinct biases in declarative motor memories. We speculate that these biases

were caused by heuristics, computationally inexpensive mental shortcuts, which not only

produce biases in cognitive decisions (Tversky and Kahneman, 1974), but can also explain

biases in visual and auditory perception (Raviv et al., 2012; Rahnev and Denison, 2018;

Gardner, 2019; Branch et al., 2022).

<u>Conclusions</u>

Here, we determined that reinforcement learning can drive the acquisition and retention of new

gait patterns. Consistent with the reinforcement learning framework, individuals do so by

exploring the action space until the most rewarding movement is discovered. Then, the most

rewarding action is exploited by repetition. Thus, the reinforcement learning framework can

describe performance across a wide range of species and behavior, and bipedal human

locomotion is no exception, as our environment is rife with risks (e.g., a trip over a cracked

sidewalk) and rewards (e.g., clearing a hurdle on the way to winning a race) that we must learn

to pair with specific actions to successfully navigate in the world.

References

Abe M, Schambra H, Wassermann EM, Luckenbaugh D, Schweighofer N, Cohen LG (2011)
Reward Improves Long-Term Retention of a Motor Memory through Induction of Offline
Memory Gains. Current Biology 21:557–562.

Bakkum A, Marigold DS (2022) Learning from the Physical Consequences of Our Actions
Improves Motor Memory. eNeuro 9 Available at:
https://www.eneuro.org/content/9/3/ENEURO.0459-21.2022 [Accessed June 7, 2022].

Bao S, Lei Y (2022) Memory decay and generalization following distinct motor learning
mechanisms. Journal of Neurophysiology 128:1534–1545.

Branch F, Park E, Hegdé J (2022) Heuristic Vetoing: Top-Down Influences of the Anchoring-
and-Adjustment Heuristic Can Override the Bottom-Up Information in Visual Images.
Front Neurosci 16:745269.

Cashaback JGA, Lao CK, Palidis DJ, Coltman SK, McGregor HR, Gribble PL (2019) The
gradient of the reinforcement landscape influences sensorimotor learning Haith AM, ed.
PLoS Comput Biol 15:e1006839.

Cashaback JGA, McGregor HR, Mohatarem A, Gribble PL (2017) Dissociating error-based and
reinforcement-based loss functions during sensorimotor learning. PLOS Computational
Biology 13:e1005623.

Codol O, Holland PJ, Galea JM (2018) The relationship between reinforcement and explicit
control during visuomotor adaptation. Scientific Reports 8 Available at:
http://www.nature.com/articles/s41598-018-27378-1 [Accessed October 11, 2018].

Daw ND, O'Doherty JP, Dayan P, Seymour B, Dolan RJ (2006) Cortical substrates for

exploratory decisions in humans. Nature 441:876–879.

Dhawale AK, Smith MA, Ölveczky BP (2017) The Role of Variability in Motor Learning. Annu

Rev Neurosci 40:479–498.

Floel A, Garraux G, Xu B, Breitenstein C, Knecht S, Herscovitch P, Cohen LG (2008) Levodopa

increases memory encoding and dopamine release in the striatum in the elderly.

Neurobiol Aging 29:267–279.

French MA, Morton SM, Charalambous CC, Reisman DS (2018) A locomotor learning paradigm

using distorted visual feedback elicits strategic learning. J Neurophysiol 120:1923–1931.

French MA, Morton SM, Reisman DS (2021) Use of explicit processes during a visually guided

locomotor learning task predicts 24-h retention after stroke. Journal of Neurophysiology

125:211–222.

Galea JM, Mallia E, Rothwell J, Diedrichsen J (2015) The dissociable effects of punishment and

reward on motor learning. Nature Neuroscience 18:597–602.

Gardner JL (2019) Optimality and heuristics in perceptual neuroscience. Nat Neurosci 22:514–

523.

Haith A, Krakauer J (2014) Motor Learning by Sequential Sampling of Actions. In, pp 2.

Washington D.C.

Haith AM, Huberdeau DM, Krakauer JW (2015) The Influence of Movement Preparation Time

on the Expression of Visuomotor Learning and Savings. Journal of Neuroscience

35:5109–5117.

Hasson CJ, Manczurowsky J, Yen S-C (2015) A reinforcement learning approach to gait training

improves retention. Frontiers in Human Neuroscience 9 Available at:

http://journal.frontiersin.org/Article/10.3389/fnhum.2015.00459/abstract [Accessed

October 11, 2018].

Holland P, Codol O, Galea JM (2018) Contribution of explicit processes to reinforcement-based

motor learning. Journal of Neurophysiology 119:2241–2255.

Huang VS, Haith A, Mazzoni P, Krakauer JW (2011) Rethinking motor learning and savings in

adaptation paradigms: model-free memory for successful actions combines with internal

models. Neuron 70:787–801.

Izawa J, Shadmehr R (2011) Learning from sensory and reward prediction errors during motor

adaptation. PLoS Comput Biol 7:e1002012.

Keisler A, Shadmehr R (2010) A Shared Resource between Declarative Memory and Motor

Memory. J Neurosci 30:14817–14823.

Kim HE (2023) bayes-toolbox. Available at: https://zenodo.org/record/7855456 [Accessed

August 1, 2023].

Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM (2019) Motor Learning. In:

Comprehensive Physiology, pp 613–663. American Cancer Society. Available at:

http://onlinelibrary.wiley.com/doi/abs/10.1002/cphy.c170043 [Accessed August 30,

2021].

Kruschke J (2014) Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan. Academic

Press.

Madelain L, Paeye C, Wallman J (2011) Modification of saccadic gain by reinforcement. Journal

of Neurophysiology 106:219–232.

Manley H, Dayan P, Diedrichsen J (2014) When Money Is Not Enough: Awareness, Success,

and Variability in Motor Learning van Beers RJ, ed. PLoS ONE 9:e86580.

Marinovic W, Poh E, de Rugy A, Carroll TJ (2017) Action history influences subsequent

movement via two distinct processes. eLife 6:e26713.

Mawase F, Uehara S, Bastian AJ, Celnik P (2017) Motor Learning Enhances Use-Dependent

Plasticity. The Journal of Neuroscience 37:2673–2685.

McAndrew Young PM, Dingwell JB (2012) Voluntarily changing step length or step width affects

dynamic stability of human walking. Gait & Posture 35:472–477.

McDougle SD, Wilterson SA, Turk-Browne NB, Taylor JA (2022) Revisiting the Role of the

Medial Temporal Lobe in Motor Learning. Journal of Cognitive Neuroscience 34:532–

549.

McElreath R (2016) Statistical Rethinking: A Bayesian Course with Examples in R and Stan.

CRC Press.

Parrell B (2021) A Potential Role for Reinforcement Learning in Speech Production. Journal of

Cognitive Neuroscience:1–17.

Pekny SE, Izawa J, Shadmehr R (2015) Reward-Dependent Modulation of Movement

Variability. Journal of Neuroscience 35:4015–4024.

Rahnev D, Denison RN (2018) Suboptimality in perceptual decision making. Behavioral and

Brain Sciences 41:e223.

Raviv L, Lupyan G, Green SC (2022) How variability shapes learning and generalization.

Trends in Cognitive Sciences 26:462–483.

Raviv O, Ahissar M, Loewenstein Y (2012) How Recent History Affects Perception: The

Normative Approach and Its Heuristic Approximation. PLoS Comput Biol 8:e1002731.

Reisman DS, Block HJ, Bastian AJ (2005) Interlimb coordination during locomotion: what can

be adapted and stored? J Neurophysiol 94:2403–2415.

Rescorla R, Wagner A (1972) A theory of Pavlovian conditioning: The effectiveness of

reinforcement and non-reinforcement. Classical Conditioning: Current Research and

Theory.

Roemmich RT, Long AW, Bastian AJ (2016) Seeing the Errors You Feel Enhances Locomotor

Performance but Not Learning. Current Biology 26:2707–2716.

Roth AM, Calalo JA, Lokesh R, Sullivan SR, Grill S, Jeka JJ, van der Kooij K, Carter MJ,

Cashaback JGA (2023) Reinforcement-Based Processes Actively Regulate Motor

Exploration Along Redundant Solution Manifolds. Neuroscience. Available at:

http://biorxiv.org/lookup/doi/10.1101/2023.02.08.527668 [Accessed March 1, 2023].

Salvatier J, Wiecki TV, Fonnesbeck C (2016) Probabilistic programming in Python using

PyMC3. PeerJ Comput Sci 2:e55.

Sato S, Cui A, Choi JT (2022) Visuomotor errors drive step length and step time adaptation

during 'virtual' split-belt walking: the effects of reinforcement feedback. Exp Brain Res

240:511–523.

Schmidt RA, Lee TD (2005) Motor control and learning: A behavioral emphasis, 4th ed.

Champaign, IL, US: Human Kinetics.

Shanks DR, Johnstone T (1999) Evaluating the relationship between explicit and implicit

knowledge in a sequential reaction time task. Journal of Experimental Psychology:

Learning, Memory, and Cognition 25:1435–1451.

Shmuelof L, Huang VS, Haith AM, Delnicki RJ, Mazzoni P, Krakauer JW (2012) Overcoming

motor "forgetting" through reinforcement of learned actions. J Neurosci 32:14617–

14621a.

Stanley J, Krakauer JW (2013) Motor skill depends on knowledge of facts. Front Hum Neurosci

7 Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3756281/ [Accessed

March 26, 2020].

Sutton R, Barto A (2017) Reinforcement Learning: An Introduction, 2nd ed. Cambridge,

Massachusetts: MIT Press.

Therrien AS, Wolpert DM, Bastian AJ (2016) Effective reinforcement learning following

cerebellar damage requires a balance between exploration and motor noise. Brain

139:101–114.

Thorndike EL (1927) The Law of Effect. The American Journal of Psychology 39:212–222.

Tsay JS, Kim HE, Saxena A, Parvin DE, Verstynen T, Ivry RB (2022) Dissociable use-

dependent processes for volitional goal-directed reaching. Proc R Soc B 289:20220415.

Tversky A, Kahneman D (1974) Judgment under Uncertainty: Heuristics and Biases. Science

185:1124–1131.

Uehara S, Mawase F, Therrien AS, Cherry-Allen KM, Celnik P (2019) Interactions between

motor exploration and reinforcement learning. Journal of Neurophysiology 122:797–808.

van Mastrigt NM, Smeets JBJ, van der Kooij K (2020) Quantifying exploration in reward-based
motor learning Worthy DA, ed. PLoS ONE 15:e0226789.

van Mastrigt NM, van der Kooij K, Smeets JBJ (2021) Pitfalls in quantifying exploration in
reward-based motor learning and how to avoid them. Biol Cybern 115:365–382.

Velázquez-Vargas CA, Daw ND, Taylor JA (2023) Learning generalizable visuomotor mappings
for de novo skills. :2023.07.18.549179 Available at:
https://www.biorxiv.org/content/10.1101/2023.07.18.549179v1 [Accessed July 24, 2023].

Verstynen T, Sabes PN (2011) How each movement changes the next: an experimental and
theoretical study of fast adaptive priors in reaching. J Neurosci 31:10050–10059.

Wilson RC, Geana A, White JM, Ludvig EA, Cohen JD (2014) Humans use directed and
random exploration to solve the explore–exploit dilemma. Journal of Experimental
Psychology: General 143:2074–2081.

Wong AL, Lindquist MA, Haith AM, Krakauer JW (2015) Explicit knowledge enhances motor
vigor and performance: motivation versus practice in sequence tasks. Journal of
Neurophysiology 114:219–232.

Wood JM, Kim HE, French MA, Reisman DS, Morton SM (2020) Use-dependent plasticity
explains aftereffects in visually guided locomotor learning of a novel step length
asymmetry. Journal of Neurophysiology 124:32–39.

Wood JM, Morton SM, Kim HE (2021) The Consistency of Prior Movements Shapes Locomotor
Use-Dependent Learning. eNeuro 8 Available at:
https://www.eneuro.org/content/8/5/ENEURO.0265-20.2021 [Accessed September 16,
2021].

Wu HG, Miyamoto YR, Castro LNG, Ölveczky BP, Smith MA (2014) Temporal structure of

motor variability is dynamically regulated and predicts motor learning ability. Nat

Neurosci 17:312–321.

Zeni JA, Richards JG, Higginson JS (2008) Two simple methods for determining gait events

during treadmill and overground walking using kinematic data. Gait & Posture 27:710–

714.

751