1   **A phylogenomic approach, combined with morphological characters gleaned via**
2   **machine learning, uncovers the hybrid origin and biogeographic diversification of the plum**
3   **genus**
4
5

Richard G. J. Hodel[1,2,*]
Sundre K. Winslow[1]
Bin-Bin Liu[3]
Gabriel Johnson[1]
Michael Trizna[2]
Alex E. White[2]
Rebecca B. Dikow[2]
Daniel Potter[4]
Elizabeth A. Zimmer[1]
Jun Wen[1]
16
17   [1] Department of Botany, National Museum of Natural History, MRC 166, Smithsonian
18   Institution, Washington, DC, 20013-7012, USA
19
20   [2] Data Science Lab, Office of the Chief Information Officer, Smithsonian Institution,
21   Washington, DC, 20560, USA
22
23   [3] State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese
24   Academy of Sciences, Beijing 100093, China
25
26   [4] Department of Plant Sciences, University of California, Davis, California, 95616, USA
27
28
29   [*] Author for correspondence: richiehodel@gmail.com
30
31

ABSTRACT

The evolutionary histories of species have been shaped by genomic, environmental, and

morphological variation. Understanding the interactions among these sources of variation is

critical to infer accurately the biogeographic history of lineages. Here, using the geographically

widely distributed plum genus (*Prunus*, Rosaceae) as a model, we investigate how changes in

genomic and environmental variation drove the diversification of this group, and we quantify the

morphological features that facilitated or resulted from diversification. We sequenced 610

nuclear loci and complete chloroplast genomes from 75 species representing all major lineages in

Prunus, with a special focus on the understudied tropical racemose group. The environmental

variation in extant species was quantified by synthesizing bioclimatic variables into principal

components of environmental variation using thousands of georeferenced herbarium specimens.

We used machine learning algorithms to classify and measure morphological variation present in

thousands of digitized herbarium sheet images. Our phylogenomic and biogeographic analyses

revealed that ancient hybridization and/or allopolyploidy spurred the initial rapid diversification

of the genus in the early Eocene, with subsequent diversification in the north temperate zone,

Neotropics, and Paleotropics. This diversification involved successful transitions between

tropical and temperate biomes, an exceedingly rare event in woody plant lineages, accompanied

by morphological changes in leaf and reproductive morphology. The machine learning approach

detected morphological variation associated with ancient hybridization and quantified the

breadth of morphospace occupied by major lineages within the genus. The paleotropical lineages

of *Prunus* have diversified steadily since the late Eocene/early Oligocene, while the neotropical

lineages diversified much later. Critically, both the tropical and temperate lineages have

continued to diversify. We conclude that the genomic rearrangements created by reticulation

deep in the phylogeny of *Prunus* may explain why this group has been more successful than

other groups with tropical origins that currently persist only in either tropical or temperate

regions, but not both.

Keywords: allopolyploidy, herbarium specimens, hybridization, machine learning, niche

conservatism, tropical-to-temperate biogeographic transitions

INTRODUCTION

Bursts of speciation associated with changing environmental conditions have occurred throughout the Tree of Life. Environmental factors, such as changes in climate, tectonic shifts, mountain uplift, as well as biotic interactions, can all drive the diversification of lineages. In particular, speciation may occur when lineages diversify to occupy newly available niches resulting from changing environmental conditions. Classic evolutionary studies such as those focused on Galapagos finches found that new ecological opportunities drove the rapid evolution of new lineages (Schluter 2000). Climatic fluctuations have been implicated in promoting both speciation and extinction in a variety of fauna (Weir and Schulter 2007). Changing environmental conditions can have a particularly strong impact on plant lineages. For example, increased global aridity likely promoted the diversification of both C4 grasses and succulent lineages (Arakaki et al. 2011). Synthesis of current evidence suggests that many plant lineages diversified in the tropics, sometimes spreading into temperate biomes (Spriggs et al. 2015), and that many temperate lineages had a Boreotropical origin (Zhang et al. 2021, Nie et al. 2023). Despite some evidence of shared biogeographic patterns, however, surprisingly few lineages have successfully made transitions from tropical to temperate regions (Kerkhoff et al. 2014). In fact, in the neotropics, the descendants of tropical ancestral lineages remained tropical in 94% of woody angiosperms (Kerkhoff et al. 2014). Similarly, temperate lineages were quite conserved, with 90% of temperate descendants arising from temperate ancestors (Kerkhoff et al. 2014). These patterns of diversification are often attributed to the tropical conservatism hypothesis (TCH) (Wiens & Donoghue 2004), which postulates that many lineages of tropical origin could not adapt readily to the cooler, drier, and more seasonal temperate regions, and either were extirpated (i.e., migrated to track tropical environments), or went extinct (Donoghue 2008).

In addition to environmental conditions, genomic changes can spark the diversification of

lineages. In some cases, reshuffling of genetic material may facilitate rapid diversification in

response to environmental change (Schenk 2021). A variety of genomic mechanisms, including

hybridization, gene duplication or loss, and/or genome doubling, can rearrange genomic material

such that new traits and adaptations develop to allow plant lineages to spread and diversify into

new niches and habitats (Xu et al. 2017, Hodel et al. 2022). Genome doubling may provide

extensive genetic variation and novelty upon which selection may act, driving adaptive

diversification (Seehausen 2004, Soltis and Soltis 2016, Doyle and Coate 2019, Griffiths et al.

2019, reviewed in Schenk 2021). Moreover, genomic changes in response to novel environments

may be coupled with morphological innovation (García-Verdugo et al. 2013). When lineages

encounter new environments, they may already possess phenotypes that facilitate their

occupancy of available niches in the community they enter, or the available niches in newly

encountered ecological conditions may drive diversifying selection on traits that are shaped to

promote survival in the novel niche space (Wellborn & Langerhans 2014). In some lineages,

success of diversification events in response to new ecological opportunities is mediated by

biological features as well as idiosyncrasies of the environmental conditions at the time

(Wellborn & Langerhans 2014). In summary, both evolutionary events and environmental

circumstances interact to shape the speciation and extinction rates that characterize

diversification (Spriggs et al. 2015).

One lineage that has been able to readily diversify in response to changing environments

was the genus *Prunus* (the plum genus, Rosaceae). This group contains approximately 250-400

evergreen and deciduous species that occur throughout the temperate regions of the northern

hemisphere and in the tropics and subtropics of both the Old and New Worlds (Rehder 1940,

Wen et al. 2008). Despite its ecological and economic significance—*Prunus* contains important

crop species such as cherries, peaches, almonds, apricots, and plums—the phylogenetic

relationships of the major lineages in the genus are still unresolved. Historically, the major

groups within the genus were defined by inflorescence morphology—with three major groups

identified—solitary flower (e.g., peach), corymbose (i.e.,producing a flat-topped indeterminate

cluster of flowers; e.g., cherries), and racemose (i.e., producing an indeterminate inflorescence

with the main axis not terminating in a flower; e.g., bird-cherries) (Rehder 1940; Su et al. 2023).

Multiple genetic studies using chloroplast DNA markers have inferred that the solitary and

corymbose groups form a clade that is sister to the racemose group (Bortiri et al. 2001, Wen et

al. 2008, Chin et al. 2014, Zhao et al. 2016). However, nuclear markers have been unable to

resolve the backbone phylogenetic relationships of the genus (Lee & Wen 2001, Bortiri et al.

2001, Wen et al. 2008, Chin et al. 2014, Zhao et al. 2016; Fig. 1). Furthermore, studies using

nuclear markers either relied on few nuclear loci (Wen et al. 2008, Chin et al. 2014, Zhao et al.

2016) or had poor sampling in the understudied racemose group (Hodel et al. 2021).

Figure 1. Synthesizing genomic, environmental, and phenomic data can improve our understanding of the
evolutionary processes that drive diversification. The summary of our phylogenetic understanding of the major groups
in the plum genus (*Prunus*) based on chloroplast and nuclear markers (top left). One environmental variable,
minimum temperature of the coldest month, often determines the distributional limits of tropical taxa (upper right).
Machine learning approaches, when applied to digitized museum specimens, can be used to infer how characters
associated with reproduction and/or environmental adaptation correspond to phylogeny (bottom).
132
133

Attempts to characterize the backbone of the *Prunus* phylogeny have been obscured by

cytonuclear discord (Chin et al. 2014). One hypothesis that has been invoked to explain this

cytonuclear discord is that the genus diversified via an ancient allopolyploidy and/or

hybridization event (Zhao et al. 2016). Based on chromosome count data, the corymbose and

solitary groups are diploid, whereas nearly all racemose lineages are polyploid.

In summary, the evolutionary history of the plum genus remains unclear. To address

phylogenomic uncertainty, herein we assemble a phylogenomic dataset with hundreds of nuclear

loci and entire chloroplast genomes for each of 75 *Prunus* species representing all major lineages

in the genus. Our taxon sampling is the most complete to date, especially in the understudied

racemose group. We use the phylogeny of *Prunus* as a framework to investigate reticulation

events, to reconstruct the biogeographic history, and test hypotheses regarding the morphological

basis of key biogeographic transitions and reticulation. The morphological characters associated

with ancient biogeographic transitions and reticulation are notoriously difficult to measure in

extant species (e.g., McVay et al. 2017). Accordingly, we developed a novel approach to

quantify and categorize morphological variation: leveraging machine learning approaches with

digitized herbarium sheet image data (Fig. 1). Our specific objectives in this study are to:

1)   Resolve the phylogenetic relationships of major groups within the genus.

2)   Assess the role of ancient genomic rearrangements—specifically allopolyploidy

and/or hybridization—in shaping the evolutionary history of *Prunus*.

3)   Clarify the biogeographic history of *Prunus*, especially concerning the timing and

frequency of transitions between tropical and temperate regions.

4)   Implement a machine learning approach with thousands of digitized herbarium

specimen images to test for morphological evidence associated with biogeographic

transitions and reticulation events such as allopolyploidy and hybridization.

166   **MATERIALS AND METHODS**

Genomic data collection

Hyb-Seq probe design

We designed a 610-locus custom Hyb-Seq probe set to target nuclear genes in *Prunus*. Our

approach aimed to obtain genes with the highest probability of being strictly single-copy (i.e.,

single-copy nuclear genes; hereafter SCNs). We used an iterative process to BLAST publicly

available genomes against themselves to obtain a candidate pool of putatively SCN loci

(Supplemental Fig. S1); the subsequent analyses were conducted in Geneious Prime 2020.0.5

(https://www.geneious.com). We first conducted BLAST searches of the annotated genomes of

P. persica (L.) Batsch (peach, PRJNA31227; Verde et al. 2013), *P. avium* (L.) L. (sweet cherry,

PRJDB4877; Shirasawa et al. 2017), and *Malus domestica* (Suckow) Borkh. (PRJNA339703,

Daccord et al. 2017) against themselves using an e-value of 1e-10, which yielded 11,305, 10,703,

and 968 candidate SCNs, respectively. We used *Malus domestica* (apple), which is in the same

subfamily as *Prunus* – the Amygdaloideae – to expand the phylogenetic breadth of the baits. We

then BLASTed the candidate SCNs from *P. avium* against the SCNs from *P. persica*, and filtered

out loci that had multiple hits, were fewer than 300 nucleotides, and had less than 95.4%

pairwise identity. Loci with multiple hits were presumably not strictly single-copy, loci with

fewer than than 300 nucleotides have less phylogenetic information than longer loci, and loci

with pairwise identity close to 100% may not be variable enough to be informative (Weitemier et

al. 2014). This left 318 loci with a total of 262,684 nucleotides of sequence, and we retained the

sequences from *P. avium* for bait design. To obtain additional loci more closely related to

racemose *Prunus* species, we also BLASTed the candidate loci that passed the above length and

pairwise identity filter, and had two or fewer hits, against the *P. serotina* Ehrh. (black cherry)

transcriptome (Swenson et al. 2017; available via the Hardwood Genomics Project,

hardwoodgenomics.org). This yielded 96 additional SCNs with 72,172 nucleotides of sequence,

and the *P. serotina* sequence was retained for bait design. We also BLASTed the candidate

SCNs from *P. avium* against the SCNs from *Malus domestica*; after filtering out loci with

multiple hits, fewer than 600 nucleotides, and less than 80% or greater than 87.5% pairwise

identity, we retained 160 loci totaling 213,990 nucleotides. Here, we used sequences from *P.*

avium for bait design. Finally, because inflorescence architecture has historically been used to

define phylogroups in *Prunus*, we selected 36 functional genes that may be associated with

flowering (e.g., APETALA, FT; Yao et al. 2022). This gene set included 36 loci totaling 137,252

nucleotides, and we used *P. avium* sequences for bait design. In total, we selected 610 loci with a

total of 686,098 nucleotides for our custom *Prunus* bait set (generated by Arbor Biosciences,

Ann Arbor, MI).

Field collection, DNA extraction, library preparation, and hybridization reactions

Specimens were collected from the field (Supplemental Table S1) and stored in silica gel

until DNA was extracted from leaf tissue using a modified CTAB protocol (Doyle & Doyle

1987). Next, DNA libraries were constructed using a KAPA HyperPrep kit (available from

Roche, Basel, Switzerland) following the manufacturer's protocol except with quarter-volume

reactions for each individual. Briefly, DNA extractions were fragmented via sonication using the

QSonica ultrasonicator (Newtown, Connecticut, USA), uneven ends were repaired and A-tailed,

and Illumina adaptors were ligated to the DNA fragments. Next, AMPure magnetic beads were

used to purify and size-select the adaptor ligated DNA, and the DNA libraries were PCR

amplified to add unique i5 and i7 indexed oligonucleotides (i.e., barcodes). Libraries were

pooled in groups of eight with the criteria of balancing samples among presumed phylogenetic

distance and concentration prior to the hybridization capture reaction with custom-designed

baits. DNA libraries were hybridized with the baits for 48 hours at 60°C following the MyBaits

v4 manual (Arbor Biosciences, Ann Arbor, Michigan, USA). The hybridization enriched

libraries were combined with unenriched DNA libraries in a 60:40 ratio to enable generation of

plastomes from off target reads. All enriched and unenriched libraries were combined into a

single tube and sent for 2 150bp sequencing on the Illumina HiSeq 4000 at Novogene

(Sacramento, California, USA). We generated data for 83 accessions (Supplemental Table S1)

representing 77 species, which include 75 *Prunus* species and two outgroups (*Lyonothamnus*

floribundus A.Gray and *Physocarpus opulifolius* (L.) Maxim.). These outgroups were selected

because *Lyonothamnus* is likely the sister lineage of *Prunus* (Xiang et al. 2017), and

Physocarpus is more distantly related but within the Amygdaloideae (Xiang et al. 2017, Zhang et

al. 2017).

Plastome assembly and phylogenetic inference

First, raw sequencing reads were quality filtered and Illumina adapters were removed using

bbduk (https://sourceforge.net/projects/bbmap/). GetOrganelle (Jin et al. 2020) was used to

generate plastomes for each species using the reads cleaned by bbduk. Of the 83 accessions

included in the final dataset, 41 produced complete, circular plastomes after an initial run

through GetOrganelle. For the remaining 42, we used minimap2 (Li 2018) implemented in

Geneious Prime 2020.0.5 (https://www.geneious.com) to assemble the contigs and scaffolds

from GetOrganelle, using the *Prunus avium* plastome (NCBI accession number MK622380) as a

reference to generate as complete as possible plastomes. After using minimap2 on the non-

circularized plastomes, we obtained nearly complete plastomes for all accessions. To ensure that

the use of *Prunus avium* as a reference did not bias the plastome results, we constructed a

phylogeny using the plastomes we generated, as well as all publicly available *Prunus* plastomes

(N = 17), to check that each newly sequenced accession is placed reasonably in the phylogeny

(Supplemental Table S2). The plastome phylogeny was inferred using RAxML with 100 rapid

bootstrap replicates and 20 independent maximum likelihood searches (i.e., the following

parameter settings: "-f a -m GTRGAMMA -p 12345 -x 12345 -# 100"). All analyses, unless otherwise

stated, were run on the Smithsonian Institution High Performance Cluster (SI/HPC, "Hydra")

(https://doi.org/10.25572/SIHPC).

Hyb-Seq locus assembly

We used the software package HybPiper v1.3 (Johnson et al. 2016) to assemble nuclear

loci. The same quality filtering and Illumina adapter removal strategy as with the plastome

pipeline was used with the nuclear data (i.e., cleaning and trimming using bbduk). We followed

the core scripts of the HybPiper pipeline; we first used `reads_first.py` to map reads using BWA (Li

& Durbin 2009), sorted reads into fasta files, and ran the SPADES assembler (Bankevich et al.

2012). Next, the scripts `get_seq_lengths.py`, `gene_recovery_heatmap.R`, and `hybpiper_stats.py` were used

to get the lengths of the recovered gene sequences, visualize the success of each gene for each

species, and generate a file of descriptive statistics for the assembly (Supplemental Table S3,

Supplemental Fig S2). By default, the `exonerate.py` script, which pulls out exon sequences, is used

in the pipeline. We also used the `intronerate.py` script to retrieve intron sequences and

supercontigs, which represent the exon plus a portion of the adjacent intron. The script

`retrieve_sequences.py` was used to retrieve exons, introns, and supercontigs from each gene region

for every species, and put them in an unaligned fasta file. Next, for each type of locus (i.e.,

exons, introns, and supercontigs), the unaligned gene files were aligned using MAFFT (Katoh &

Standley 2013) with the following parameter settings: "`--maxiterate 5000 --auto --`

`adjustdirectionaccurately —leavegappyregion`." The resulting alignments were trimmed using the `phyx`

command `pxclsq` with the proportion of sites required to have data (`-p` option) set at 0.5 (Brown et

al, 2017). For each of the trimmed alignments, gene trees were estimated using RAxML

(Stamatakis 2014) with 100 rapid bootstrap replicates and 20 independent maximum likelihood

searches (i.e., the following parameter settings: "`-f a -m GTRGAMMA -p 12345 -x 12345 -# 100`"). After

confirming that exons and supercontigs produced similar species tree topologies (Supplemental

Fig S3), we used supercontigs for all subsequent analyses.

Species tree estimation

The nuclear species tree was estimated using ASTRAL-III (Zhang et al. 2018), an

approach consistent with the coalescent, to summarize the 610 gene trees. We used the ASTRAL

quartet scores to assess nodewise support for inferred phylogenetic relationships. A

concatenation approach implemented in RAxML was also used for comparative purposes. For

some downstream analyses (e.g., divergence dating), it was necessary to have a phylogeny with

meaningful branch lengths. For these analyses, we used the ASTRAL topology as a constraint

tree for a RAxML tree search to obtain a species tree with the ASTRAL topology but with

proportional branch lengths.

Gene tree discordance analysis

To assess congruence between gene trees and the species tree, we used the program

phyparts (Smith et al. 2015). At each node in the tree, phyparts compares the rooted gene tree

topologies with the rooted species tree to label the number of genes that are concordant,

discordant, and uninformative relative to the species tree. We considered any gene trees with less

than 50% bootstrap support at a given node to be uninformative for that node (i.e., -s 50 option).

Because only rooted trees can be used in phyparts, we retained 505 gene trees for this analysis—

those where there was data present for one or both outgroups.

Paralogs

We used the HybPiper post-processing scripts `paralog_investigator.py` and `paralog_retriever.py`

to report the number of suspected paralogs for each gene and retrieve the multiple sequences for

putative paralogous genes, respectively. Next, the unaligned retrieved gene sequences were

inputted into a phylogenetic pipeline that includes alignment with MAFFT (Katoh & Standley

2013) and phylogeny construction with FastTree (Price et al. 2009). In some cases, genes

suspected to be paralogs will form a clade, indicating that the paralogs may be alleles and not the

result of duplication. Alternatively, some genes are truly paralogs and, in this case, multiple

copies may be dispersed throughout the tree inferred by FastTree. For all 610 genes, we

classified its paralogy status based on the presence or absence of paralogs, and if paralogs were

detected, we manually inspected the paralog tree for such genes to determine if the suspected

paralogs were alleles or in fact true paralogs indicating ancient duplication. In the cases where

true paralogs were detected, we noted the number and phylogenetic distribution of the paralogs.

Reticulation analyses

To quantify the amount of reticulation present at each node, we used the simulation

approach of Cai et al. (2021). Briefly, a reticulation index can be calculated for each node, which

is a relative measure of the intensity of introgression at the node. Because the expectations of the

relative frequencies of alternative triplets compared to the species tree topology should be

different under incomplete lineage sorting (ILS) versus introgression, this approach can quantify

the per-node relative amount of introgression. Symmetrical distributions of the two alternative

topologies would be expected under ILS, but the distributions would be asymmetrical under

introgression (Cai et al. 2021). For asymmetrical triplets, the introgressed branches are mapped

to the species trees, and the raw counts of introgressed branches were normalized to generate the

Reticulation Index for each node.

Hybridization and allopolyploidy analyses

Because hypotheses of ancient hybridization and allopolyploidy have been proposed to

explain the origin and diversification of this group, we used SNaQ (Solís-Lemus et al. 2016),

which is implemented using PhyloNetworks (Solís-Lemus et al. 2017). SNaQ was selected

because it can explicitly accommodate introgression/gene flow and ILS. SNaQ uses gene tree

frequencies for 4-taxa sets to estimate a phylogenetic network with one or more reticulation

edges. The inheritance probabilities γ and $1-\gamma$ are used to infer the probability that lineages are

the parental participants in a hybridization event, which is represented by one or more hybrid

edges added to a phylogenetic tree to form a phylogenetic network. The optimal number of

hybrid edges is determined via a pseudo-deviance score when the candidate networks are

optimized based on inheritance probabilities and branch lengths in phylogenetic network space.

We investigated h_{max} values, which represent the maximum number of hybrid edges allowed in a

network, that ranged from 0 to 5. Because SNaQ becomes computationally intractable when too

many terminals are used, we selected one or a few taxa from major lineages—one each from the

solitary, corymbose, and temperate racemose groups, two from the neotropical racemose, three

from the paleotropical racemose group, and two from the early diverging tropical racemose

lineages—for a total of 10 species in the SNaQ analysis. Because we were interested in

diagnosing ancient hybridization, using a minimal number of species can better infer ancient, as

opposed to more recent, hybridization events. A denser sampling of species could potentially

confound results because of signals of more recent hybridization within major groups. Within

lineages, the species with optimal gene sequence coverage was selected as the representative

taxon for that lineage (Supplemental Table S3).

To investigate histories of allopolyploidy, and for comparison with hybridization results

from SNaQ, we used the program GRAMPA (Thomas et al. 2017). This approach uses a least

common ancestor mapping algorithm to reconcile gene trees and species trees (Goodman et al.

1979; Page 1994) so that polyploidy events can be placed on a phylogeny. GRAMPA can

identify modes of polyploidy as well as place WGDs on a phylogeny. The user can input clades

of interest that are suspected to have a polyploid origin, or use a global search that considers if

all clades could be the result of WGD. We used the latter approach to acquire an unbiased

assessment if any lineages were potentially of polyploid origin.

Environmental analysis

To quantify the environment for major groups, we used a Principal Components Analysis

(PCA) to synthesize variation in the 19 WorldClim bioclimatic variables

(https://www.worldclim.org/) at a 2.5-minute resolution. The values for each of the 19 variables

were extracted for georeferenced herbarium specimens that represented each group using the R

packages 'dismo' (Hijmans et al. 2017) and 'raster' (Hijmans 2016). Two approaches were used

to quantify the environmental variation, first between two groups—temperate versus tropical—as

well as a comparison of four groups: temperate diploid (solitary + corymbose), temperate

racemose, neotropical racemose, and paleotropical racemose. We distilled the variation in 19

bioclimatic variables into 3 PC axes using the R package 'vegan' (Oksanen et al. 2017). For this

analysis, we only used the 75 *Prunus* species for which we had genomic data, and a total of

10,011 herbarium specimen records representing these 75 species were included. Although this

approach only captures present environmental variation, synthesis has revealed that niches are

more phylogenetically conserved than expected (Donoghue 2008), and accordingly surveying all

extant species in a lineage can give a reasonable approximation for the environment in which a

given lineage evolved. We also used the GPS data to collect elevation data of each specimen

using the R package 'rgbif' (Chamberlain et al. 2023). This was done primarily to investigate

whether tropical species occurred at lower or higher elevations, because some high elevation

tropical regions may resemble temperate environments more closely than tropical ones, and this

may impact our interpretation of biogeographic results.

Dating and biogeographic analysis

To investigate the timing and biogeographic history of *Prunus*, we used a biogeographic

ancestral range estimation analysis implemented in BioGeoBears (Matzke 2012, 2013). First, we

used treePL, a method for estimating divergence times on a phylogeny using penalized

likelihood, to infer divergence dates for all nodes in the *Prunus* phylogeny. Three calibrations

were used to date the phylogeny. We used *Prunus cathybrownae* (Benedict et al. 2011) from the

Early Eocene of North America as the most recent common ancestor (MRCA) of the diploid

(i.e., corymbose+solitary) lineages (stem lineage leading to node Dip; Fig. 2), with a minimum

age of 50 million years ago (Mya). The fossil *Prunus wutuensis* (Li et al. 2011) from the early

Eocene of East Asia (Shandong, China) was used to fix the MRCA of crown *Prunus* to a

minimum age of 58 Mya. Next, the crown of the Amygdaloideae was fixed at a minimum age of

90 Mya based on a calibrated Rosaceae phylogeny from Xiang et al. (2017). Similarly, we

constrained the age of the stem Amygdaloideae to have a minimum age of 100 Mya using the

calibrated Rosaceae phylogeny from Xiang et al. (2017) as a guide. Using BioGeoBears (Matzke

2013), we considered six possible biogeographic models (DEC, DEC + J, DIVALIKE,

DIVALIKE + J, BAYAREALIKE, BAYAREALIKE + J) and selected the optimal model using

AIC and AICc comparisons. We defined seven biogeographic regions: North America, South

America, Africa, Europe, West Asia, East Asia, and Australasia. Likelihood ratio tests were also

used to determine if the more parameter-rich +J models were preferred compared to the

equivalent models without the +J term.

Morphological data collection

Herbarium image analysis

Because of the rich representation of *Prunus* species in museum collections, we used a

deep learning approach with digitized herbarium specimen sheets to test the degree to which

morphological classifications of species and/or lineages corresponded to phylogenetic

hypotheses. We assembled a dataset of 4,228 images representing the 81 *Prunus* accessions with

genomic data in this study. These images were downloaded using the 'idig_search_media'

function in the 'ridigbio' R package (Michonneau et al. 2016). We used exiftool

(https://exiftool.org/) to remove EXIF metadata from the images using the command "`exiftool -`

`overwrite_original -EXIF= *.jpg`"). This approach removes metadata often incompatible with image

processing libraries commonly used in machine learning. We also trimmed the edges of the

digitized images before using them in downstream machine learning analyses. This was done to

remove non-biological information, such as herbarium labels, color bars, and scale bars, from the

digitized herbarium sheet. We removed the top 20% and bottom 20% of every image used in

machine learning analyses. This unnecessarily removed some biologically relevant features, but

was conservative for ensuring the removal of non-biologically meaningful information on the

images. We investigated both trimmed and untrimmed sheets to test the impact of removing the

bottom 20% and top 20% of each sheet; hereafter these sheets are referred to as 'trimmed' or

'whole sheet', respectively. Furthermore, because we had thousands of digitized herbarium

sheets available, this approach represented a reasonable balance between removing too much

data and ensuring that any data with the potential to bias results was excluded.

 Classification Algorithms

To assess if morphological variation corresponded to genetic variation, we developed a

supervised machine learning classifier algorithm using the fastai (https://github.com/fastai/fastai)

deep learning library, which is built using PyTorch libraries (https://pytorch.org), to classify

herbarium sheet images into categories. The labels for each category were assigned based on

clades inferred using genomic data. We used two approaches for the classification analysis. First,

we assigned labels of 'diploid species,' corresponding to all species with solitary and corymbose

inflorescences, and 'tropical racemose,' which referred to all tropical species with racemes.

These two groups were selected because the phylogenetic relationship between these two groups

was the same in the nuclear and chloroplast phylogenies. This model was trained and validated

using a total of 4,228 herbarium sheet images representing 67 species, with 2,030 in the

solitary/corymbose group, and 2,200 in the tropical racemose group. In each group, 80% of the

labeled input data were used for training, and 20% for validation. The model was run for 24

epochs until an accuracy rate of 97.2% was achieved. As an additional check that the model was

performing well, we tested hundreds of specimens from species in either the solitary/corymbose

groups (N = 452), or the tropical racemose groups (N = 426), which were not species used to

train or validate the model. Next, the additional 'test' data in this implementation were species

from the temperate racemose group (N = 565). This group was selected as test data because its

phylogenetic position was different between the nuclear and chloroplast topologies. Essentially,

this machine learning classification approach was used to test if there were morphological

signatures of hybridization in the temperate racemose group that correspond to the genomic

signatures of hybridization detected. Although the model was not trained to classify temperate

racemose species, we expect that if the temperate racemose group arose via hybridization,

species in this group may retain some morphological features from both the solitary/corymbose

group and the tropical racemose group, if these two groups are the parental participants in

ancient hybridization. The second classifier model we developed had the goal of distinguishing

between four major groups: temperate diploid, temperate racemose, neotropical racemose, and

paleotropical racemose. This second model was trained and validated using a total of 4,228

herbarium sheet images representing 67 species. In each group, 80% of the labeled input data

were used for training, and 20% for validation. The model was run for 24 epochs until an

accuracy rate of 92.4% was achieved. This second classification model was built so we could

ultimately measure the breadth of morphological variation across specimens that could be

successfully categorized in one of the four groups using a dimensionality reduction approach

called IVIS (described below).

Gradient CAM

As a further ground-truthing step to ensure that the classification algorithm was assessing

biologically meaningful variation, as opposed to learning biologically meaningless features for

identification (e.g., labels, rulers), we used Gradient Class Activation Mapping (Grad-CAM), an

approach for visualizing how deep neural networks make their predictions (Selvaraju et al.

2017). Briefly, Grad-CAM generates a heatmap highlighting the regions in an input image most

important for why a particular prediction was made. A Grad-CAM algorithm takes the output of

a convolutional layer in the neural network and computes the gradient of target category in

relation to that layer's activations, which are the pixels that are 'activated' by the model—the

pixels that are most important for determining the class. The gradients of target category are then

unsampled and overlaid onto the query image to generate a heatmap highlighting pixels and

regions of the image most important for determining the prediction. Grad-CAM specifically

excels when applied to Convolutional Neural Networks (CNNs), due to their ability to learn and

extract useful features from images. By interrogating the pixels of an image that the model

considers important, Grad-CAM can inform why and how a CNN makes its predictions. We

sampled the final layer of the CNN, as well as the second-to-last and third-to-last layers. We

categorized the heatmap images as classifying based on leaf features, floral features, or

neither/ambiguous. Each image with both leaf and floral structures present was classified as

'floral', 'mostly floral', 'mixed', 'mostly leaf', or 'leaf'. We also assigned images that did not

fall into the above categories as 'leaf with no floral structure present', 'no leaf or floral structure

apparent', 'ambiguous', or 'problematic'. The last two categories were for images where it was

difficult to tell if the heatmap favored either floral structures or leaves, or if the heatmap was

overly focused on non-biological information, respectively.

Morphological space in tropical and temperate groups

IVIS (Implicit Variational Information Sharing) is a deep learning approach with the goal

of mapping high-dimensional data into low-dimensional space. Critically, IVIS works well with

image data and can preserve distances between samples in a global sense, which is not always

true using other dimensionality reduction methods such as PCA, t-SNE, or umap (Szubert et al.

2019, Chari et al. 2021). In the case of image data, IVIS works by encoding the input image into

a lower-dimensional representation, and then using a decoder to reconstruct the image. During

training, the network learns to form a compact representation by minimizing the reconstruction

loss and encouraging the low-dimensional space to be continuous and well-structured through

the use of a regularization term. The low-dimensional representation learned by IVIS can be used

for various tasks, such as clustering, visualization, or further fine-tuning for classification tasks.

IVIS has been shown to be effective for image data, outperforming traditional methods such as

PCA and t-SNE in terms of clustering and visualization quality. Here, we use IVIS to reduce the

highly-dimensional variation present in image data into 2-dimensional space to make

comparisons among specimens and groups.

RESULTS

Phylogenomic analyses

For 75 *Prunus* species representing all major lineages, we assembled a dataset of 610

nuclear genes and complete chloroplast genomes to infer phylogeny. The nuclear coalescent

species tree inferred using ASTRAL (Fig. 2) recovered monophyletic corymbose and solitary

groups, but a paraphyletic racemose group. With our robust sampling, the solitary, corymbose,

and temperate racemose species form a clade, although with modest support (36.7% ASTRAL

quartet support (QS) score; see node Temp ("Temperate"), Fig. 2). This clade of solitary,

corymbose, and temperate racemose taxa was sister to a large clade composed of tropical

racemose species, including representatives from both the neotropics and paleotropics. The

monophyly of each of the solitary and corymbose clades was strongly supported (i.e., > 70% QS;

Fig. 2). The sister relationship between the solitary and corymbose clades also had strong support

(72.5% QS score; node Dip ("Diploid"), Fig. 2). There were also several other strongly

supported clades, including the paleotropical subgenus *Pygeum*, and the clade of neotropical

racemose species (see nodes Pal ("Paleotropical") and SNeo ("Neotropical, predominantly South

America"), respectively; Fig. 2).

The paleotropical racemose species did not form a clade – the neotropical racemose

species were nested within the paleotropical group (Fig. 2). A clade composed of nearly all

tropical racemose species, except for a few early-diverging lineages (*Prunus undulata* Buch.-

Ham. ex D.Don, *P. laurocerasus* L., *P. africana* (Hook.f.) Kalkman, and *P. skutchii* I.M.Johnst.),

had relatively strong support (65.0% QS score; see node CTrop ("Core Tropical"), Fig. 2).

Additionally, the New World (primarily neotropical) racemose species were not monophyletic.

Whereas the exclusively South American neotropical racemose species formed a strongly

supported clade, four other New World racemose species (*P. ilicifolia* (Nutt. ex Hook. & Arn.)

D.Dietr., *P. lyonni* (Eastw.) Sarg., *P. occidentalis* Sw., and *P. cornifolia* Koehne) formed a clade

with low support that was sister to the Australasian paleotropical racemose clade (Fig. 2A).

In contrast to the nuclear phylogeny, the chloroplast tree indicated that each group

defined by inflorescence morphology formed a clade (Fig. 2). Each of the solitary and

corymbose clades had high bootstrap support (100% and 99%, respectively), and the sister

relationship between these two clades was also strongly supported (99%; Fig. 2B). Within the

major clades, many of the relationships in the chloroplast phylogeny were similar to the nuclear

phylogeny. The neotropical racemose group was similarly non-monophyletic in the chloroplast

tree, and additionally the New World racemose species did not form a clade in the chloroplast

phylogeny (Fig. 2B). Notably, two species that were early-branching racemose lineages in the

nuclear tree (the European *P. laurocerasus* and the East and southeast Asian *P. undulata*), were

nested within the temperate racemose species (Fig. 2B).

Figure 2A. The phylogenetic species tree inferred using ASTRAL to summarize 610 nuclear gene trees. Important
nodes in the phylogeny are highlighted and named: temperate ('Temp'), diploid ('Dip'), tropical ('Trop'), core tropical
('CTrop'), East Asian paleotropical ('EPal'), Australasia paleotropical ('APal'), South American neotropical ('SNeo'),
and North American neotropical ('NNeo'). The numbers at nodes are ASTRAL support scores.

532 **Figure 2B.** The phylogenetic tree based on chloroplast genomes. Several key nodes that are congruent with the
533 nuclear species tree are highlighted and labeled. The numbers at each node represent bootstrap percentage scores
534 from the RAxML analysis.

Gene tree discordance

Many nodes in the nuclear species tree were characterized by rampant gene tree

discordance, as assessed by phyparts (Supplemental Fig. S4). At most nodes, there were far more

gene trees that were discordant with the species tree topology than were congruent

(Supplemental Fig. S4). Of note, fewer than 1% of gene trees were concordant with the species

tree at both the node defining the clade temperate racemose + solitary + corymbose, and the node

defining the tropical racemose clade (Fig. 2). Within major clades, some nodes had high

concordance between gene trees and the species tree (e.g., the node defining a group of

neotropical racemose species), whereas others such as the node that defined the temperate

racemose species, had low concordance (Supplemental Fig. S4).

Paralogs

We assessed paralogs in two ways: first using the paralog detector from the assembly

statistics file, and then using the phylogenetic approach with the post-processing scripts

`paralog_investigator.py` and `paralog_retriever.py`. Despite our stringent filtering during bait design to

prioritize SCNs, all accessions had at least 4 genes with paralog warnings based on the paralog

detection approach during assembly (Supplemental Table S3). Notably, all species that were

from the solitary and corymbose groups—presumably diploids—had paralog warnings for

between 4 and 59 of the 610 SCNs. However, species from the racemose groups had between 74

and 335 genes with paralog warnings; this higher presence of paralogs is consistent with a

genome doubling event in the evolutionary history of the racemose groups (e.g., Zhao et al.

2016). When we used a tree-building approach to identify paralogs, we found 77 SCNs with no

evidence of paralogy, 139 genes with between 1-10 paralogous tips, 198 genes with between 11-

30 paralogous terminals, and 196 genes with more than 31 paralogous tips in the gene tree

(Supplemental Table S4). We manually inspected the phylogenies with duplicated gene copies

and found the vast majority of duplicate copies were from species in the racemose group

(Supplemental Table S4).

Reticulate topology

Excluding the two outgroups, the reticulation indices ranged from 0.05 to 0.28

(Supplemental Fig. S5). This level of reticulation is high compared to other clades studied using

the same metric, which had a maximum reticulation index of 0.1 (Cai et al., 2021). The optimal

SNaQ network, as assessed by change in pseudolikelihood score, had two hybridization edges

(Fig. 3). In this phylogenetic network, one reticulation event was detected in the backbone of the

phylogeny: *Prunus serotina* was inferred to have originated via hybridization between an

ancestral member of the solitary/corymbose clade and an early-diverging tropical racemose

lineage represented by *Prunus undulata* (Fig. 3). For this hybrid edge, *P. serotina* was 71.2%

sister to the entire solitary/corymbose group, and 28.8% sister to *P. undulata*. The second hybrid

edge was found within the tropical racemose group, with the lineage containing *P. oligantha*

Kalkman and *P. arborea* var. *stipulacea* (King) Kalkman 91.5% sister to *P. cornifolia*, but also

8.5% sister to *P. africana* (Fig. 3), the earliest-diverging lineage within the paleotropical *Pygeum*

group, suggesting a hybrid origin of the latter. In summary, the first hybridization edge implies a

hybrid origin of the temperate racemose group, with the parents as an ancestor of the

corymbose/solitary group and a member of very early-diverged lineages in the tropical racemose

group (Fig. 2, Fig. 3). Meanwhile, the second hybrid edge indicates a hybrid origin of the

Pygeum group, with the parents as the ancestor of the lineage containing *P. cornifolia*, and an

ancestor of an early diverged tropical racemose species (i.e., *P. africana*; Fig. 2, Fig. 3).

587
588
589
Figure 3. The optimal 2-hybridization edge phylogenetic network estimated using SNaQ for 10 species representing
every major lineage in *Prunus*. The first hybridization edge, shown in blue, indicates that *P. serotina* is 71.2% sister to
the corymbose/solitary clade, and 28.8% sister to an early diverging tropical racemose lineage, which gave rise to *P.*
undulata. The second hybridization edge shows hybridization within the racemose clade, with *P. oligantha* and *P.*
arborea stipulacea determined to be 91.5% sister to *P. cornifolia*, and 8.5% sister to an early diverging tropical
racemose lineage—an ancestor of *P. africana*.
596
597
598

Allopolyploid origin: GRAMPA

The tests for signatures of allopolyploidy using the program GRAMPA indicated that a

MUL-tree was the most parsimonious tree, better even than a singly-labeled tree. The most

parsimonious MUL-tree, with a parsimony score of 56691, was characterized by a duplicated

clade that comprised nearly all of the species in the tropical racemose group, excluding four

early-diverging tropical racemose species, *P. africana*, *P. undulata*, *P. skutchii*, and *P.*

laurocerasus (Supplemental Fig. S7). The GRAMPA analysis we conducted concluded that 38

accessions from the tropical racemose group were the result of an allopolyploidy event between

the ancestor of *P. skutchii* and an ancestor of all *Prunus* accessions included in this study. A total

of 25,593 MUL-trees and one singly-labeled tree were investigated in our untargeted analysis.

The second most parsimonious MUL-tree had a parsimony score of 56972, whereas the singly

labeled tree had a parsimony score of 58626. The second-most parsimonious tree was very

similar to the most parsimonious MUL-tree; both trees identified an identical clade (i.e., nearly

all tropical racemose species) as the result of polyploidy. The key difference in the second-most

parsimonious MUL-tree was that *P. skutchii* was included in the duplicated clade, and the

ancestor of *P. africana* was the parental participant in an allopolyploid event in its stead. In

summary, the ancestor of an early-diverging tropical racemose lineage was implicated as the

topological origin of allopolyploidy.

Environmental characterization

The PCA of environmental variation based on the 19 bioclimatic variables captured

78.8% of variation on the first three principal components, with 46.5% of the variation

encapsulated in PC1, 17.3% in PC2, and 15.0% in PC3 (Fig. 4). One precipitation variable—

annual precipitation—was strongly positively correlated with PC1, and four temperature

variables—mean annual temperature, minimum temperature of the coldest month, mean

temperature of the driest quarter, and mean temperature of the coldest quarter—were also

strongly associated with positive PC1 space. Meanwhile, two temperature variables—

temperature seasonality, and mean annual temperature range—were strongly and negatively

associated with PC1 values. No bioclimatic variables were strongly correlated with PC2 or PC3.

In summary, tropical species occupy more positive regions of PC1, which correspond to warmer

and wetter conditions, with higher minimum temperatures in colder seasons, and less

temperature seasonality (Fig. 4), whereas temperate species trend towards negative values of

PC1. When considering the two types of tropical species—neotropical versus paleotropical—the

paleotropical species extended further towards both extremes of PC1, and further towards the

negative extent of PC2. Neotropical species extended slightly further into positive PC2 space

than paleotropical species. The temperate racemose species had a greater affinity for positive

values of PC1 whereas corymbose/solitary species trended further towards extreme negative

values of PC1. The temperate racemose species occupied a larger range of PC2 space than the

corymbose/solitary species (Fig. 4). There were clear elevational differences among species

representing the four major groups. In the temperate solitary/corymbose group, only two out of

28 species had a median elevation greater than 1,000m (Supplemental Fig. S8). In contrast, half

of the 10 temperate racemose species occurred at a median elevation greater than 1,000m, and

also half of the 10 neotropical racemose species were found at a median elevation greater than

1,000m. In the paleotropical racemose species, 13 of 28 species had a median elevation greater

than 1,000m (Supplemental Fig. S8).

Figure 4. The summary of environmental variation using a Principal Components Analysis to distill the 19 bioclimatic
variables into 3 dimensions. PC1 accounts for 46.5% of variation while PC2 explains 17.3% of environmental
variation. Several bioclimatic variables strongly correlated with PC1 are indicated on the X-axis. Negative PC1 space
is associated with increased temperature seasonality and annual temperature range. Meanwhile, positive PC1 space
corresponds to higher annual precipitation, annual mean temperature, minimum temperature of the coldest month,
and mean temperature of the driest quarter and coldest quarter.

Biogeographic analysis

The DEC+J model was the preferred model after comparing AIC scores of competing

BioGeoBears models, which were based on the time-calibrated tree generated with TreePL

(Supplemental Table S5). Likelihood ratio tests concluded that the addition of the J parameter,

despite producing a more parameter-rich model, led to a significantly better model than DEC

(Supplemental Table S6). The BioGeoBears analysis indicated an East Asian origin of the genus,

with the initial diversification occurring in the Paleocene ca. 60-55 Mya (Fig. 5). Subsequently,

the temperate racemose, solitary/corymbose, and tropical racemose clades all diverged by the

early-mid Eocene (ca. 50 Mya). This diversification occurred predominantly in East Asia and

North America, which were characterized by the Boreotropical climate during the Late

Cretaceous – early Eocene (ca. 65-50 Mya). The temperate racemose group steadily diversified

in East Asia and North America beginning in the mid-late Eocene (ca. 40 Mya). Both the solitary

and corymbose lineages began diversifying in the early Miocene (ca. 25-20 Mya), first in East

Asia and subsequently in North America and Europe. These two lineages currently occupy

temperate environments, and based on our sampling, they were more successful in speciating in

East Asia versus North America (Fig. 5).

Broadly, the tropical racemose clade's early diversification occurred in several regions,

including East Asia, West Asia, Africa, North America, and Europe, between 55-40 Mya. Early-

diverging lineages, including species such as *Prunus laurocerasus*, *P. undulata*, *P. africana*, and

P. skutchii, split off during the Eocene between ca. 55-40 Mya. Notably, the ancestors of these

few early-diverging lineages occupy a wide geographic range, including Europe, West Asia, East

Asia, Southeast Asia, Africa, North America, and South America. Subsequently, the patterns of

diversification occurred differently in the major lineages within the tropical racemose group. The

neotropical lineages, occurring in tropical North and South America, are characterized by a long

period of stasis, followed by rapid diversification in the Miocene beginning between 25-20 Mya

and lasting until ca. 10 Mya. In contrast, two major paleotropical lineages, in East Asia and

Southeast Asia, respectively, began steadily diversifying in the Oligocene (ca. 35 Mya), and

continued until approximately 10 Mya. Notably, there were multiple clades in both the temperate

and tropical groups of *Prunus* that demonstrated rapid diversification beginning in the early

Miocene.

Figure 5. The BioGeoBears analysis of the biogeographic history of *Prunus* using the DEC + J model based on seven biogeographic regions: North America (blue), South America (red), Africa (magenta), Europe (turquoise), West Asia (green), East Asia (yellow), and Southeast Asia (orange). Color-coded pie charts at each node indicate the probability of the ancestral state occurring in a given biogeographic region.

Morphological analyses

We ran two experiments using computer vision-based machine learning analyses of

digitized herbarium sheets. The assessment of morphological characters associated with

hybridization revealed that whole sheet phenotypes of temperate racemose species, with the top

20% and bottom 20% of the sheet trimmed, were nearly evenly divided between classification as

'solitary/corymbose' (N = 233 observations with > 90% probability) and 'tropical racemose' (N

= 194 observations with >90% probability) (Fig 6). When using whole sheets (i.e., untrimmed),

there was a similar trend: there were N = 239 observations classified as 'solitary/corymbose'

with >90% probability and N = 172 observations classified as 'tropical racemose' with > 90%

probability. The Grad-CAM analysis of internal model layers indicated that inflorescence vs.

leaf morphology was the driving factor in determining if the machine learning model inferred

whether to classify temperate racemose species as corymbose/solitary or tropical racemose (Fig.

6). Manual visual inspection indicated that the second-to-last and third-to-last layers were most

informative for determining classifications. The Grad-CAM results for the last three layers

model layers for every herbarium sheet specimen image are shown in Supplemental Tables S7

and S8.

Figure 6. In the upper left, the distribution of probability scores when using the temperate racemose species as test
data in the first machine learning model. These classifications include specimens that were classified as either
'solitary/corymbose or tropical racemose'; regardless of classification, most specimens were assigned with high (i.e.,
>0.95) probability. The plot in the upper right shows the manual classifications we assigned when ground-truthing the
gradient class activation maps for the 565 temperate racemose specimens for the second-to-last and third-to-last
layer of the CNN. The bottom row shows the gradient class activation (Grad-CAM) maps for the second-to-last model
layer for two representative temperate racemose specimens that were classified as 'tropical racemose' (left) and
'solitary/corymbose' (right), respectively.

718
719
720

721    In the second analysis, breadth of morphospace was measured using IVIS to quantify and

722 compare the extent of morphological variation present in the following groups: temperate diploid

723 (i.e., corymbose and solitary flower), temperate racemose, neotropical racemose, and

724 paleotropical racemose. The IVIS analysis determined that the two tropical groups exhibited a

725 greater range of morphological variation than the two temperate groups (Fig. 7). Within all

726 tropical specimens, the paleotropical racemose species occupied a greater expanse of

727 morphological variation than neotropical racemose (Fig. 7). The paleotropical species overlapped

728 in morphospace to a small degree with the temperate diploid group (i.e., corymbose/solitary) and

729 to a much greater degree with the neotropical racemose group. The neotropical species are

730 positioned centrally in morphospace and overlap with all three other groups (Fig. 7).

Figure 7. The representation of whole specimen phenotypes in IVIS space. The specimen images were assigned to four groups according to environmental preference/clade: neotropical racemose, paleotropical racemose, temperate racemose, corymbose/solitary.

DISCUSSION

In this study, we provided increased resolution compared to previous studies of the

phylogeny of the economically and ecologically important genus *Prunus*, pinpointed key

genomic mechanisms promoting the diversification of this group, and improved our

understanding of its biogeographic history. By sampling more densely the understudied

racemose group and sequencing hundreds of nuclear loci and complete chloroplast genomes, we

changed our understanding of how several groups within the genus diversified. Specifically, by

combining analyses of hybridization, genome doubling, and cytonuclear and gene tree-species

tree conflict, we inferred that the polyploid racemose group is paraphyletic, a result found in

some previous studies using nuclear gene data (e.g., Lee & Wen 2001, Bortiri et al. 2002, 2006,

Wen et al. 2008, Chin et al. 2014, Zhao et al. 2016), but typically not found when using plastid

genes (e.g., Bortiri et al. 2001, Wen et al. 2008, Chin et al. 2014). The chloroplast phylogeny,

with a monophyletic racemose group, detected in the present study is congruent with many

previous studies using chloroplast markers (e.g., Bortiri et al. 2006, Wen et al. 2008, Chin et al.

2014, Zhao et al. 2016). Notably, our result conflicts with Su et al. (2023), which found a

monophyletic racemose group in both nuclear and chloroplast datasets. However, the nuclear

SNP markers used in Su et al. (2023) may been unable to fully untangle the reticulate history of

this genus. Previous studies have suggested that cytonuclear discord observed in *Prunus*

phylogenies could be attributed to one (Chin et al. 2014) or multiple (Zhao et al. 2016)

allopolyploid hybrid origins of the racemose group. Here, we isolated specific lineages as

participants in likely hybridization and allopolyploidy events during the early Eocene ca. 50

Mya. These reticulation events early in the diversification of the genus help explain the high

gene tree-species tree conflict and cytonuclear discord. Through biogeographic and fossil-

calibrated dating analyses, we traced the biogeographic history of the group and tied key climatic

events to radiations of subgeneric lineages.

Using several novel applications of machine learning algorithms to classify and quantify

morphological variation, we addressed how morphological evolution coincides with cytonuclear

discord, identifying characters essential for survival in specific environments and how

morphological variation in lineages impacts their biogeographic distribution. Phylogenomic data

have revealed that gene tree-species tree conflict is widespread and may sometimes arise via

ancient hybridization (e.g., McVay et al. 2017, Nie et al. 2023). Here, we demonstrate that a

well-trained machine learning classification model struggles to classify specimens in a lineage

that is the result of ancient hybridization, as these specimens that are the result of hybridization

display morphological features common to both putative ancestors. Additionally, we applied a

machine learning approach, IVIS, to quantify the morphological variation within major lineages,

revealing that the distinct morphospaces of different lineages may be shaped by their

biogeographic and evolutionary history. These morphological classifications, in combination

with environmental data, change our concept of how lineages may have diversified in the tropics,

and the conditions under which lineages transition from tropical to temperate biomes. Below, we

discuss and contextualize our results.

Improved understanding of Prunus *phylogeny, diversification, and biogeography*

Despite its economic and ecological importance, until now *Prunus* has not been

thoroughly investigated in a phylogenomic context with sufficient sampling in the racemose

group. Previous studies, based on chloroplast DNA and/or a few nuclear markers, inferred

possible cytonuclear conflict—a telltale sign suggesting hybridization—on the backbone of the

phylogeny. These analyses lacked the resolution to conclude whether the cytonuclear discord

was real or due to unresolved nuclear relationships (Lee & Wen 2001, Bortiri et al. 2001, Wen et

al. 2008). Based on cytonuclear discord (Chin et al. 2014), multiple copies of nuclear loci (Zhao

et al. 2016), nuclear reduced representation sequencing (Su et al. 2023), and chromosome count

data coupled with multiple gene tree topologies (Hodel et al. 2021), hypotheses of ancient

allopolyploid hybridization have been proposed (Zhao et al. 2016). Here, multiple lines of

evidence indicate a history of allopolyploidy and/or hybridization driving early diversification in

Prunus. First, we demonstrate that the cytonuclear discord along the backbone of the genus,

especially the differing phylogenetic placement of the temperate racemose group, provides a

specific hypothesis of reticulation (i.e., the temperate racemose group is a product of

hybridization and/or allopolyploidy). The SNaQ phylogenetic network supports this hypothesis,

showing the temperate racemose group as a product of hybridization (Fig. 3); notably, SNaQ is

not designed to detect allopolyploidy, but the results of an allopolyploidy event may present in

the genome similarly to hybridization. The GRAMPA analysis, which can detect and distinguish

between different types of genome doubling, also identified reticulation deep in the tree, but with

different entities resulting from allopolyploidy—the vast majority of the tropical racemose clade

was inferred to have an allopolyploid origin deep in the phylogeny (Supplemental Fig. S7). The

Reticulation Indices also showed a high level of reticulation deep in the tree with a RI of 0.185

for the crown node of *Prunus* (Supplemental Fig. S5), and a RI of 0.158 at the node defining the

temperate racemose + solitary/corymbose clade (i.e., TEMP node in Fig. 2A). Taken together,

these analyses point to reticulation deep in the phylogeny, although it is not pinpointed to one

instance of hybridization and/or allopolyploidy. These results are in line with previous

hypotheses that predicted multiple rounds of allopolyploidy (Zhao et al. 2016). Probably the

majority, if not all, racemose species, encompassing both temperate and tropical climates, are

polyploid. These species likely arose from multiple rounds of ancient allopolyploidy rather than

autopolyploidy.

The higher resolution, time-calibrated plum genus phylogeny enhances our understanding

of the biogeographic history of the clade. The diversification and biogeographic history of

Prunus can be contextualized by existing hypotheses describing biogeographic patterns, and it

can shift our understanding of the range of possible outcomes for lineages that originated in the

tropics. Our biogeographic results show pulses of diversification in several tropical and

temperate *Prunus* lineages, with bursts of speciation in the early Miocene (Fig. 5). This contrasts

with steadier diversification reported in previous studies (e.g., Chin et al. 2014). Our time-

calibrated results also highlight the different patterns of speciation in different tropical *Prunus*

groups: relatively steady speciation in the paleotropics, with punctuated bursts in the neotropics

(Fig. 5). There were also key differences in greater temperate success in Asia versus North

America: the temperate lineages in Asia speciated readily and rapidly while the species diversity

in North America is depauperate. Although our sampling of species diversity and nuclear loci

gives better resolution than previous studies, there are still limitations when interpreting our

biogeographic results. Critically, biogeographic analyses depend on bifurcating phylogenies, and

many of our results point to a reticulate topology for *Prunus*, especially deeper in the tree. It is

likely that multiple lineages intermingled in the expansive Boreotropical region early in the

evolutionary history of the genus. For example, many of the lineages implicated in reticulation,

such as the early diverging tropical racemose lineages, may have overlapped during the

Eocene—even though their ancestors occupy distinct biogeographic zones—thus facilitating

ancient hybridization and/or allopolyploidy. Finally, we must acknowledge that inferences of

diversification may be affected by sampling bias. We present the densest sampling of the

understudied tropical racemose clade to date, with accessions representing all major lineages.

However, additional future work is needed to ensure that even unbiased sampling does not

impact the results, as this genus is estimated to contain 250-400 species, necessitating further

sampling in future studies.

Using Prunus *to inform broad biogeographic patterns*

For over a century, biologists have observed a latitudinal gradient in species diversity in

many clades across the Tree of Life, with greater species richness occurring near the equator.

However, we lack scientific consensus about the causes of this biogeographic pattern and several

hypotheses have been proposed. One explanation for the observed latitudinal gradient is the

tropical conservatism hypothesis (TCH), which posits that the relatively massive biodiversity of

the tropics can be primarily attributed to the geographic extent of tropical taxa over the past 55

million years and the subsequent evolutionary conservation of environmental niches (Wiens and

Donoghue 2004). Many groups have diversified in the Eocene, facilitated by the vast expanse of

the Boreotropics. Nevertheless, relatively few lineages have transitioned from tropical

environments to temperate ones. This phenomenon may be explained by the challenge of

acquiring the substantial adaptations necessary to tolerate the cooler conditions in temperate

zones (Donoghue 2008).

Another explanation is provided by the 'taxon pulse' hypothesis (Erwin 1985), which is

compatible with the TCH and posits that lineages that were able to transition from tropical to

temperate regions originated in tropical environments and then migrated in waves to more

temperate regions at higher altitudes and latitudes into increasingly harsh environments (Erwin

1985, Lü et al. 2020, Rapini et al. 2021). These radiations into temperate environments would

also be accompanied by species loss in ancestral tropical lineages as older lineages went extinct

(Liebherr and Porch 2015, Rapini et al. 2021, Nie et al. 2023). Some widespread plant clades,

such as *Viburnum*, match the taxon pulse hypothesis to an extent (Spriggs et al. 2015). In this

scenario, tropical lineages are posited to be 'dying embers' – lineages that diversified in the

tropics tens of millions of years ago, and persist in place, but have ceased to continue

diversifying (Spriggs et al. 2015).

Our results indicate *Prunus* does not precisely follow the predictions of either the TCH or

the 'taxon pulse' hypothesis. In *Prunus*, the paleotropical lineages have diversified in place and

do not appear to be 'dying embers', instead demonstrating steady diversification through time

(Fig. 5). As described in the taxon pulse hypothesis, tropical lineages may migrate to higher

temperate latitudes or to higher elevation regions in the tropics, which may have temperate-like

conditions. Meanwhile, the diversification of the neotropical lineages, instead of occurring

steadily as observed in our paleotropical species, was characterized by bursts of speciation

beginning in the late Oligocene-early Miocene (ca. 25-20 Mya). In the tropics, some *Prunus*

species have moved to higher elevations, but there are also lineages that occupy truly tropical

environments, as defined by some or all individuals occurring in environments with the

minimum temperature of the coldest month greater than 18 $^{\circ}$C. Among species in our study,

these include *Prunus dolichobotrys* (Lauterb. & K.Schum.) Kalkman, *P. arborea* (Blume)

Kalkman, *P. gazelle-peninsulae* (Kaneh. & Hatus.) Kalkman, *P. javanica* (Teijsm. & Binn.)

Miq.*, P. undulata, P. fordiana* Dunn*, P. grisea* Kalkman*, P. costata* (Hemsl.) Kalkman*, Prunus*

maingayi (Hook. f.) Wen*, P. debilis* Koehne*, P. oocarpa* (Stapf) Kalkman*, P. oligantha, P.*

schlechteri (Koehne) Kalkman*, P. ceylanica* (Wight) Miq.*, and P. buergeriana* Miq. in the

paleotropics, and *P. myrtifolia* (L.) Urb., *P. amplifolia* Pilg., *P. cornifolia, P. integrifolia*

(C.Presl) Walp.*, P. occidentalis*, *P. skutchii*, and *P. reflexa* (Gardner) Walp. in the neotropics.

We also note that, for the species we sampled genetically in this study, in both tropical groups, at

least half of the species occurred at a median elevation of less than 1,000 m (Supplemental Fig.

S8).

This comparison demonstrates that diversification can occur differently in distinct

tropical regions—even within a single genus. Our quantifications of the morphological breadth

and environmental breadth revealed both to be greater in the paleotropics as compared to the

neotropics. The larger environmental space of the paleotropics suggests there may have been

greater ecological opportunity for lineages in the paleotropics relative to the neotropics due to

increased niche availability (Wellborn & Langerhans 2015). For similar reasons, the greater

morphological variation in the paleotropical *Prunus* shows these lineages were able to leverage

advantageous morphological features to adapt to new environmental niches. This implies *Prunus*

differs from many lineages with a tropical origin that successfully transitioned to temperate

regions. Synthesis has shown that not all lineages could adapt to colder climates when tropical

habitat retracted, and typically those lineages that tracked tropical habitats became more

restricted (Donoghue 2008). However, *Prunus* bucks this pattern, exhibiting steady speciation in

the paleotropics, and later bursts of speciation in the neotropics. It is possible that the neotropical

racemose lineages have ancestors that occupied temperate zones in North America (Fig. 5). The

small clade of four North American species (Fig. 2A; node 'NNeo') is a likely candidate—these

species may be Boreotropical remnants in the New World, highlighting the importance of North

America as a site of diversification early in the evolution of some tropical racemose species.

Perhaps a history of surviving in the temperate zone enabled present-day neotropical lineages to

develop and/or keep a suite of morphological characters that facilitated radiating into the

neotropics. The substantial overlap in morphospace between neotropical and temperate lineages

suggests this may be the case (Fig. 7).

Why was Prunus *more successful than other lineages?*

Because of our sampling of species diversity coupled with hundreds of nuclear loci, we

identify allopolyploidy and/or hybridization in the backbone of the phylogeny, which may

explain the genomic basis of the rapid diversification 55-45 Mya. This allopolyploidy event is

also supported by phylogenetic treatments of the Rosaceae, which found evidence of a whole

genome duplication event at the base of *Prunus* (Xiang et al. 2017). This is an important insight

for not only understanding the phylogeny of the group but also its biogeographic context. When

numerous *Prunus* lineages were able to interact in the Boreotropics during the Eocene, ample

opportunities for hybridization and allypolyploidy arose. This genetic reshuffling may explain

why *Prunus* was more successful than other lineages at occupying both temperate and tropical

regions. The Rosaceae family has diversified into many successful lineages (Potter et al. 2007),

but even within this hyperdiverse group, *Prunus* stands out for its diversity of inflorescence

types, which may have facilitated migration via a variety of different animal dispersers, as well

as variation in leaf morphology supporting both evergreen and deciduous life histories. One key

innovation in *Prunus* is the evolution of leaf glands, which appear to be associated with climatic

conditions, with flat glands typical of species in tropical climates, and raised glands present in

species occupying cooler climates (Chin et al. 2013). These glands may have had adaptive value

due to their interactions with insects, with the flat glands found in tropical regions preventing

herbivory, and may be a key feature driving the diversification of *Prunus*. Specifically, the

variation in these morphological features may have enabled *Prunus* radiations into the temperate

zone, while continuing to diversify in the tropics. Other Rosaceae, although successful, remained

constrained in temperate regions (Xiang et al. 2017). Additionally, other groups with tropical

origins that have experienced great success in temperate regions did not retain much species

diversity in tropical regions, including *Quercus* L. (Hipp et al. 2020), *Viburnum* (Spriggs et al.

2015), Juglandaceae (Zhang et al. 2021), and Saxifragales (Folk et al. 2019).

Morphological variation informs biogeographic history

The morphological variation associated with recent hybridization in many cases can be

readily tracked to inform the parental participants in hybridization, as well as to understand how

and why hybridization occurred, and whether hybrids may persist based on their environment.

However, in cases of ancient hybridization, it becomes difficult to tease apart the morphological

characters associated with hybridization since subsequent diversification and evolution in

response to environmental variation can obscure the circumstances surrounding the hybridization

event. Here, we used machine learning to quantify morphological variation in clades resulting

from reticulation, presenting a way to characterize the morphological legacy of

hybridization/allopolyploidy. By using the ancient hybridization detected via cytonuclear discord

and SNaQ networks to guide our morphological analyses, we concluded that there were

morphological signals of hybridization in the present species. Members of the temperate

racemose group had morphological features of both the diploid clade and the tropical racemose

clade—the ancestors of these two lineages were putatively the participants in hybridization. So,

we would expect morphological features from both parents to persist in the lineage resulting

from reticulation. However, lineages' responses to environmental variation may also explain

patterns of morphological variation not resulting, or minimally resulting, from the genomic

results of hybridization (or allopolyploidy). However, in this case, we would expect that

morphological variation would co-vary to some extent with environmental variation. This would

lead to an expectation that the vast majority of test specimens from the temperate racemose

group would be classified as solitary/corymbose, because the environment overlaps much more

between these two groups than it does between temperate racemose and tropical racemose (Fig.

4). Contrary to this expectation, we observed a relatively even split of temperate racemose

species between the two training categories.

Morphological shifts can result from transitioning to a novel environment, or existing

morphological variation can enable lineages to adapt to changing environmental conditions. The

key takeaway from our IVIS analysis is that both the neotropical and paleotropical groups exhibit

substantially more morphological variation than do the temperate groups, including both the

corymbose/solitary group, and the temperate racemose group. Moreover, the two temperate

groups occupy peripheral portions of morphological space on both IVIS axes 1 and 2, suggesting

that the morphological features in the temperate groups are specialized relative to the

morphological features of the tropical groups. These features may represent morphological

changes that promote transitions to temperate regions, diverging from morphological features

typical for tropical zones, in response to a transition from tropical to temperate biomes. Broadly,

if there is greater niche space available in the tropics—a proposed explanation for the increased

species diversity in these regions—we would expect greater morphological variation to occupy

more niches. In *Prunus*, the balance of morphological space we quantify is not surprising given

greater tropical diversity. Notably, however, the morphological space occupied by the

neotropical racemose species overlaps more with the two temperate groups than the paleotropical

group does. This suggests an explanation for some of the biogeographic patterns detected—

specifically that the more recent diversification of the neotropics was different from the initial

diversification in the paleotropics. By the time the neotropical lineages began to diversify,

lineages in the genus had already radiated into and adapted to temperate biomes. This may

explain the rapid diversification of the neotropical *Prunus*—the lineages may have already

acquired multiple morphological innovations for surviving in the New World's warm temperate

and tropical zones during the Oligocene-early Miocene.

The promise of machine learning to assess morphology via digitized herbarium specimens

Machine learning approaches such as IVIS offer the promise of using whole-specimen

phenotyping with minimal preprocessing to classify species based on a biologically informed

hypothesis. These hypotheses could be based on phylogeny, environmental gradients, ecological

features, or geography. We demonstrate a method to use specimen data to investigate

correspondence between environmental and morphological variation (Fig. 8). Approaches such

as IVIS allow researchers to be agnostic regarding the morphological features studied, which can

avoid researcher-meditated bias in selecting traits. By grouping together specimens in different

and/or hierarchical groups, competing hypotheses can be tested. We demonstrate how specimens

can be grouped together to test hypotheses of hybridization. It is common to use the species as

the unit of classification with image-based machine learning algorithms. However, the species is

not necessarily the only level of biological organization that can be used to train classifiers.

Classifier algorithms that differentiate between species implicitly use a phenetic species concept

(de Queiroz 2007), which is not ideal for all research questions. We argue that groups above and

below species are useful, especially if they may be defined by shared phenotypic features; here

we demonstrate an application at a broader systematic level. Taking a phylogenetic approach to

grouping units to be classified, based on key phenetic differences, presents a way to mesh

phylogenetic and phenotypic data. Although there are benefits to using whole-specimen

phenotypes, there are also advantages to extracting individual traits, such as measurements of

leaf area and perimeter, as well as floral traits. These can be extracted via machine learning

methods such as segmentation in a high-throughput fashion.

100s-1,000s biological specimen images

Establish groups to test hypothesis relevant to research question:

- Phylogenetic hypothesis based on conflicting genomic data (e.g., arising from hybridization)
- Ecological hypothesis to test if morphology covaries with habitat preference
- Systematic/taxonomic hypotheses associated with delimiting cryptic species

- Divide data into two or more biologically informed groups

- Subdivide data from each group into training and validation sets

- Train the deep learning model using training and validation data until high accuracy is achieved

- Use test data from specimens known to fit into *a priori* groups to further assess model performance

- Investigate how data from additional groups are categorized by the model

Group 1

Group 2

Validation Data

Validation Data

Training Data

Training Data

Deep Learning Classifier Model

Group 1 Test Data

Group 2 Test Data

99% probability Group 1

99% probability Group 2

Data from Group to be investigated

97% probability Group 2

Figure 8. A flow chart for designing phylogenetically-informed hypotheses to test using machine learning applied to biological specimen image data.

Conclusions and future prospects

In this study, we employ phylogenomic, environmental, and morphological data to

establish that the initial diversification of the cherry genus *Prunus* was driven by ancient

hybridization and/or allopolyploidy. Moreover, this complex history of reticulation may explain

the success of the tropical lineages of *Prunus* compared to other groups that also moved out of

the tropics and into temperate regions. To complement phylogenomic and environmental

analyses, we present innovative applications of machine learning algorithms to analyze the

variation in digitized herbarium sheets. We demonstrate inventive ways to harness computer

vision-based machine learning approaches to help us understand biodiversity across the Tree of

Life. Future studies in *Prunus* will require additional sampling, especially in the racemose group,

to further examine biogeographic patterns. Subsequent research may also focus on developing

segmentation masks to extract specific features from digitized image data. *Prunus* represents an

excellent model for testing the capabilities of machine learning algorithms using museum data,

given the wealth of publicly available digitized specimens in the genus.

1014

Data Availability

Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/dryad.[NNNN].
Supplementary material, image data, and DNA sequence matrices are available from the Dryad
Digital Repository. Raw sequence data were submitted to NCBI GenBank (SUB13638423).
Machine learning models are hosted on Hugging Face with temporary URLs
(https://huggingface.co/MikeTrizna/Prunus_lineage_classifier;
https://huggingface.co/MikeTrizna/Prunus_herbarium_sheet_classifier). Final Hugging Face
URLs will be hosted by the Smithsonian Institution upon acceptance. Jupyter notebooks for
developing and processing machine learning models for image data are available on GitHub
(https://github.com/richiehodel/machine_learning_Prunus_herbarium_sheets).

Acknowledgements

We thank the following funding sources: Smithsonian Institution Scholarly Studies Grant to JW,
EAZ, RGJH; Smithsonian Institution Peter Buck Fellowship to RGJH; National Museum of
Natural History NHRE Award to SKW.

We thank the following people for valuable discussions and assistance with analyses: Greg Stull,
Alicia Talavera.

All or portions of the laboratory and/or computer work were conducted in and with the support
of the Laboratories of Analytical Biology facilities of the National Museum of Natural History or
its partner labs.

Literature Cited

Arakaki M., Christin P.A., Nyffeler R., Lendel A., Eggli U., Ogburn R.M., Spriggs E., Moore M.J., Edwards E.J. 2011. Contemporaneous and recent radiations of the world's major succulent plant lineages. Proc. Natl. Acad. Sci. U. S. A. 108:8379–8384.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M., Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A. V., Sirotkin A. V., Vyahhi N., Tesler G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. J. Comput. Biol. 19:455–477.

Benedict J.C., DeVore M.L., Pigg K.B. 2011. *Prunus* and *Oemleria* (Rosaceae) flowers from the late early Eocene Republic flora of Northeastern Washington State, U.S.A. Int. J. Plant Sci. 172:948–958.

Bortiri E., Oh S.-H., Jiang J., Baggett S., Granger A., Weeks C., Buckingham M., Potter D., Parfitt D.E. 2001. Phylogeny and systematics of *Prunus* (Rosaceae) as determined by sequence analysis of *ITS* and the chloroplast *trnL-trnF* spacer DNA. Syst. Bot. 26:797–807.

Bortiri E., Oh S.-H., Gao F., Potter D. 2002. The phylogenetic utility of nucleotide sequences of sorbitol 6-phosphate dehydrogenase in *Prunus* (Rosaceae). Amer. J. Bot. 89:1697–1708.

Bortiri E., Vanden Heuvel B., Potter D. 2006. Phylogenetic analysis of morphology in *Prunus* reveals extensive homoplasy. Plant Syst. Evol. 259:53–71.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: phylogenetic tools for unix. Bioinformatics. 33:1886–1888.

Cai L., Xi Z., Lemmon E.M., Lemmon A.R., Mast A., Buddenhagen C.E., Liu L., Davis C.C. 2021. The perfect storm: gene tree estimation error, incomplete lineage sorting, and ancient gene flow explain the most recalcitrant ancient angiosperm clade, Malpighiales. Syst. Biol. 70:491–507.

Carranza-Rojas J., Goeau H., Bonnet P., Mata-Montero E., Joly A. 2017. Going deeper in the automated identification of Herbarium specimens. BMC Evol. Biol. 17:181.

Chamberlain S, Barve V, Mcglinn D, Oldoni D, Desmet P, Geffert L, Ram K (2023). rgbif: Interface to the Global Biodiversity Information Facility API. R package version 3.7.6, https://CRAN.R-project.org/package=rgbif.

Chari T., Banerjee J., Pachter L. 2021. The Specious Art of Single-Cell Genomics. BioRxiv.:1–25.

Chin S.W., Shaw J., Haberle R., Wen J., Potter D. 2014. Diversification of almonds, peaches, plums and cherries - Molecular systematics and biogeographic history of *Prunus* (Rosaceae). Mol. Phylogenet. Evol. 76:34–48.

Daccord N., Celton J.M., Linsmith G., Becker C., Choisne N., Schijlen E., Van De Geest H., Bianco L., Micheletti D., Velasco R., Di Pierro E.A., Gouzy J., Rees D.J.G., Guérif P., Muranty H., Durel C.E., Laurens F., Lespinasse Y., Gaillard S., Aubourg S., Quesneville H., Weigel D., Van De Weg E., Troggio M., Bucher E. 2017. High-quality de novo

assembly of the apple genome and methylome dynamics of early fruit development. Nat. Genet. 49:1099–1106.

De Queiroz K. 2007. Species Concepts and Species Delimitation. Syst. Biol. 56:879–886.

Donoghue M.J. 2008. A phylogenetic perspective on the distribution of plant diversity. Proc. Natl. Acad. Sci. U. S. A. 105:11549–11555.

Doyle J.J., Coate J.E. 2019. Polyploidy, the nucleotype, and novelty: The impact of genome doubling on the biology of the cell. Int. J. Plant Sci. 180:1–52.

Doyle J.J., Doyle J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. Phytochem. Bull. 19:11–15.

Folk R.A., Stubbs R.L., Mort M.E., Cellinese N., Allen J.M., Soltis P.S., Soltis D.E., Guralnick R.P. 2019. Rates of niche and phenotype evolution lag behind diversification in a temperate radiation. Proc. Natl. Acad. Sci. U. S. A. 166:10874–10882.

García Verdugo C., Friar E., Santiago L.S. 2013. Ecological role of hybridization in adaptive radiations:A case study in the *Dubautia arborea-Dubautia ciliolata* (Asteraceae) complex. Int. J. Plant Sci. 174:749–759.

Griffiths A.G., Moraga R., Tausen M., Gupta V., Bilton T.P., Campbell M.A., Ashby R., Nagy I., Khan A., Larking A., Anderson C., Franzmayr B., Hancock K., Scott A., Ellison N.W., Cox M.P., Asp T., Mailund T., Schierup M.H., Andersen S.U. 2019. Breaking free: The genomics of allopolyploidy-facilitated niche expansion in white clover. Plant Cell. 31:1466–1487.

Hijmans, R. J. 2016. raster: Geographic Data Analysis and Modeling. R pack- age version 2 5-8.

Hijmans, R. J., S. Phillips, J. Leathwick, and J. Elith. 2017. Dismo: Species distribution modeling. R package version 1:1-4.

Hipp A.L., Manos P.S., Hahn M., Avishai M., Bodénès C., Cavender-Bares J., Crowl A.A., Deng M., Denk T., Fitz-Gibbon S., Gailing O., González-Elizondo M.S., González-Rodríguez A., Grimm G.W., Jiang X.L., Kremer A., Lesur I., McVay J.D., Plomion C., Rodríguez-Correa H., Schulze E.D., Simeone M.C., Sork V.L., Valencia-Avalos S. 2020. Genomic landscape of the global oak phylogeny. New Phytol. 226:1198–1212.

Hodel R.G.J., Zimmer E., Wen J. 2021. A phylogenomic approach resolves the backbone of *Prunus* (Rosaceae) and identifies signals of hybridization and allopolyploidy. Mol. Phylogenet. Evol. 160.

Hodel R.G.J., Massatti R., Knowles L.L. 2022. Hybrid enrichment of adaptive variation revealed by genotype–environment associations in montane sedges. Mol. Ecol. 31:3722–3737.

Hodel R.G.J., Zimmer E.A., Liu B. Bin, Wen J. 2022. Synthesis of nuclear and chloroplast data combined with network analyses supports the polyploid origin of the apple tribe and the hybrid origin of the Maleae—Gillenieae clade. Front. Plant Sci. 12:3321.

Jin J.J., Yu W. Bin, Yang J.B., Song Y., DePamphilis C.W., Yi T.S., Li D.Z. 2020.
GetOrganelle: A fast and versatile toolkit for accurate *de novo* assembly of organelle
genomes. Genome Biol. 21:241.
Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C.,
Wickett N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics
from high☐throughput sequencing reads using target enrichment. Appl. Plant Sci.
4:1600016.
Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.
Kerkhoff A.J., Moriarty P.E., Weiser M.D. 2014. The latitudinal species richness gradient in
New World woody angiosperms is consistent with the tropical conservatism hypothesis.
Proc. Natl. Acad. Sci. 111:8125–8130.
Li G., Davis B.W., Eizirik E., Murphy W.J. 2016. Phylogenomic evidence for ancient
hybridization in the genomes of living cats (Felidae). Genome Res. 26:1–11.
Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34:3094–
3100.
Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 25:1754–1760.
Li Y., Smith T., Liu C.J., Awasthi N., Yang J., Wang Y.F., Li C. Sen. 2011. Endocarps of
Prunus (Rosaceae: Prunoideae) from the early Eocene of Wutu, Shandong Province,
China. Taxon. 60:555–564.
Liebherr J.K., Porch N., Liebherr J.K., Porch N. 2015. Reassembling a lost lowland carabid
beetle assemblage (Coleoptera) from Kauai, Hawaiian Islands. Invertebr. Syst. 29:191–
213.
Lü L., Cai C.Y., Zhang X., Newton A.F., Thayer M.K., Zhou H.Z. 2021. Linking evolutionary
mode to palaeoclimate change reveals rapid radiations of staphylinoid beetles in low-
energy conditions. Curr. Zool. 66:435–444.
Matzke N.J. 2013. Probabilistic historical biogeography: new models for founder-event
speciation, imperfect detection, and fossils allow improved accuracy and model-testing.
Front. Biogeogr. 5.
McVaugh R. 1951. A revision of the North American black cherries (*Prunus serotina* Ehrh.,
and relatives). Brittonia. 7:279–315.
McVay J.D., Hipp A.L., Manos P.S. 2017. A genetic legacy of introgression confounds
phylogeny and biogeography in oaks. Proc. R. Soc. B Biol. Sci. 284.
Michonneau F., Collins M., Chamberlain S.A. 2016. ridigbio: An interface to iDigBio's search
API that allows downloading specimen records. R package version 0.3.2.
https://github.com/iDigBio/ridigbio
Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep history
of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New
Phytol. 218:1668–1684.

Nie Z., Hodel R.G.J., Ma Z., Johnson G., Ren C., Meng Y., Ickert☐Bond S.M., Liu X., Zimmer E., Wen J. 2023. Climate☐influenced boreotropical survival and rampant introgressions explain the thriving of New World grapes in the north temperate zone. J. Integr. Plant Biol. 65:1183–1203.

Oksanen, J. F., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, et al. 2017. vegan: Community ecology package. R package version 2.4-4

Parins-Fukuchi C., Stull G.W., Smith S.A. 2021. Phylogenomic conflict coincides with rapid morphological innovation. Proc. Natl. Acad. Sci. U. S. A. 118:e2023058118.

Price M.N., Dehal P.S., Arkin A.P. 2009. FastTree: Computing large minimum evolution trees with profiles instead of a distance matrix. Mol. Biol. Evol. 26:1641–1650.

Rapini A., Bitencourt C., Luebert F., Cardoso D. 2021. An escape-to-radiate model for explaining the high plant diversity and endemism in campos rupestres. Biol. J. Linn. Soc. 133:481–498.

Rehder, A., 1940. Manual of cultivated trees and shrubs hardy in North America exclusive of the subtropical and warmer temperate regions, 2nd ed. MacMillan, New York.

Schenk J.J. 2021. The next generation of adaptive radiation studies in plants. Int. J. Plant Sci. 182:245–262.

Schluter D. 2000. The ecology of adaptive radiation. Oxford University Press.

Schuettpelz E., Frandsen P.B., Dikow R.B., Brown A., Orli S., Peters M., Metallo A., Funk V.A., Dorr L.J. 2017. Applications of deep convolutional neural networks to digitized natural history collections. Biodivers. Data J.:e21139.

Seehausen O. 2004. Hybridization and adaptive radiation. Trends Ecol. Evol. 19:198–207.

Selvaraju R.R., Cogswell M., Das A., Vedantam R., Parikh D., Batra D. 2017. Grad-CAM: Visual explanations from deep networks via gradient-based localization. .

Shirasawa K., Isuzugawa K., Ikenaga M., Saito Y., Yamamoto T., Hirakawa H., Isobe S. 2017. The genome sequence of sweet plum (*Prunus avium*) for use in genomics-assisted breeding. DNA Res. 24:499–508.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals conflict, concordance, and gene duplications with examples from animals and plants. BMC Evol. Biol. 15:150.

Solís-Lemus C., Ané C. 2016. Inferring phylogenetic networks with maximum pseudolikelihood under incomplete lineage sorting. PLOS Genet. 12:e1005896.

Solís-Lemus C., Bastide P., Ané C. 2017. PhyloNetworks: A package for phylogenetic networks. Mol. Biol. Evol. 34:3292–3298.

Soltis P.S., Soltis D.E. 2016. Ancient WGD events as drivers of key innovations in angiosperms. Curr. Opin. Plant Biol. 30:159–165.

Spriggs E.L., Clement W.L., Sweeney P.W., Madriñán S., Edwards E.J., Donoghue M.J. 2015. Temperate radiations and dying embers of a tropical past: the diversification of *Viburnum*. New Phytol. 207:340–354.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics. 30:1312–3.
Su N., Hodel R.G.J., Wang X., Wang J.R., Xie S.Y., Gui C.X., Zhang L., Chang Z.Y., Zhao L.,
Potter D., Wen J. 2023. Molecular phylogeny and inflorescence evolution of *Prunus*
(Rosaceae) based on RAD-seq and genome skimming analyses. Plant Divers.
Szubert B., Cole J.E., Monaco C., Drozdov I. 2019. Structure-preserving visualisation of high
dimensional single-cell datasets. Sci. Rep. 9:1–10.
Verde I., Abbott A.G., Scalabrin S., Jung S., Shu S., Marroni F., Zhebentyayeva T., Dettori
M.T., Grimwood J., Cattonaro F., Zuccolo A., Rossini L., Jenkins J., Vendramin E.,
Meisel L.A., Decroocq V., Sosinski B., Prochnik S., Mitros T., Policriti A., Cipriani G.,
Dondini L., Ficklin S., Goodstein D.M., Xuan P., Del Fabbro C., Aramini V., Copetti D.,
Gonzalez S., Horner D.S., Falchi R., Lucas S., Mica E., Maldonado J., Lazzari B.,
Bielenberg D., Pirona R., Miculan M., Barakat A., Testolin R., Stella A., Tartarini S.,
Tonutti P., Arús P., Orellana A., Wells C., Main D., Vizzotto G., Silva H., Salamini F.,
Schmutz J., Morgante M., Rokhsar D.S. 2013. The high-quality draft genome of peach
(*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome
evolution. Nat. Genet. 45:487–494.
Weir J.T., Schluter D. 2007. The latitudinal gradient in recent speciation and extinction rates of
birds and mammals. Science. 315:1574–1576.
Weitemier K., Straub S.C.K., Cronn R.C., Fishbein M., Schmickl R., McDonnell A., Liston A.
2014. Hyb-Seq: Combining target enrichment and genome skimming for plant
phylogenomics. Appl. Plant Sci. 2:1400042.
Wellborn G.A., Langerhans R.B. 2015. Ecological opportunity and the adaptive diversification
of lineages. Ecol. Evol. 5:176–195.
Wen J., Berggren S.T., Lee C.-H., Ickert-Bond S., Yi T.-S., Yoo K., Xie L., Shaw J., Potter D.
2008. Phylogenetic inferences in *Prunus* (Rosaceae) using chloroplast *ndhF* and nuclear
ribosomal *ITS* sequences. J. Syst. Evol. 46:322–332.
Wen J., Nie Z.-L., Ickert-Bond S.M. 2016. Intercontinental disjunctions between eastern Asia
and western North America in vascular plants highlight the biogeographic importance of
the Bering land bridge from late Cretaceous to Neogene. J. Syst. Evol. 54:469–490.
Wiens J.J., Donoghue M.J. 2004. Historical biogeography, ecology and species richness. Trends
Ecol. Evol. 19:639–644.
Xiang Q.-Y., Soltis D.E., Soltis P.S. 1998. The eastern Asian and eastern and western North
American floristic disjunction: Congruent phylogenetic patterns in seven diverse genera.
Mol. Phylogenet. Evol. 10:178–190.
Xu S., He Z., Zhang Z., Guo Z., Guo W., Lyu H., Li J., Yang M., Du Z., Huang Y., Zhou R.,
Zhong C., Boufford D.E., Lerdau M., Wu C.-I., Duke N.C., Shi S. 2017. The origin,
diversification and adaptation of a major mangrove clade (Rhizophoreae) revealed by
whole-genome sequencing. Natl. Sci. Rev. 4:721–734.

Yao J.L., Kang C., Gu C., Gleave A.P. 2022. The roles of floral organ genes in regulating
1256   Rosaceae fruit development. Front. Plant Sci. 12:644424.

1257 Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
1258   reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

1259 Zhang Q., Ree R.H., Salamin N., Xing Y., Silvestro D. 2021. Fossil-informed models reveal a
1260   boreotropical origin and divergent evolutionary trajectories in the walnut family
1261   (Juglandaceae). Syst. Biol. 71:242–258.

1262 Zhao L., Jiang X.-W., Zuo Y., Liu X.-L., Chin S.-W., Haberle R., Potter D., Chang Z.-Y., Wen
1263   J. 2016. Multiple events of allopolyploidy in the evolution of the racemose lineages in
1264   *Prunus* (Rosaceae) based on integrated evidence from nuclear and plastid data. PLoS One.
1265   11:e0157123.

1266

1267