A phylogenomic approach, combined with morphological characters gleaned via machine learning, uncovers the hybrid origin and biogeographic diversification of the plum genus

Richard G. J. Hodel[1,2,3*]

Sundre K. Winslow[2]

Si-Yu Xie[4]

Bin-Bin Liu[5]

Liang Zhao[4]

Gabriel Johnson[2]

Michael Trizna[3]

Alex E. White[3]

Rebecca B. Dikow[3]

Daniel Potter[6]

Elizabeth A. Zimmer[1]

Jun Wen[1]

[1] Department of Biological Sciences, Northern Arizona University, 617 S. Beaver St., Flagstaff AZ 86011, USA

[2] Department of Botany, National Museum of Natural History, MRC 166, Smithsonian Institution, Washington, DC, 20013-7012, USA

[3] Data Science Lab, Office of the Chief Information Officer, Smithsonian Institution, Washington, DC, 20560, USA

[4] College of Life Sciences, Northwest A&F University, Yangling 712100, China

[5] State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

[6] Department of Plant Sciences, University of California, Davis, California, 95616, USA

[*] Author for correspondence: richiehodel@gmail.com

36   **ABSTRACT**

The evolutionary histories of species have been shaped by genomic, environmental, and

morphological variation. Understanding the interactions among these sources of variation is

critical to infer accurately the biogeographic history of lineages. Here, using the geographically

widely distributed plum genus (*Prunus*, Rosaceae) as a model, we investigate how changes in

genomic and environmental variation drove the diversification of this group, and we quantify the

morphological features that facilitated or resulted from diversification. We sequenced 587

nuclear loci and complete chloroplast genomes from 99 species representing all major lineages in

Prunus, with a special focus on the understudied tropical racemose group. The environmental

variation in extant species was quantified by synthesizing bioclimatic variables into principal

components of environmental variation using thousands of georeferenced herbarium specimens.

We used machine learning algorithms to classify and measure morphological variation present in

thousands of digitized herbarium sheet images. Our phylogenomic and biogeographic analyses

revealed that ancient hybridization and/or allopolyploidy spurred the initial rapid diversification

of the genus in the early Eocene, with subsequent diversification in the north temperate zone,

neotropics, and paleotropics. This diversification involved successful transitions between tropical

and temperate biomes, an exceedingly rare event in woody plant lineages, accompanied by

morphological changes in leaf and reproductive morphology. The machine learning approach

detected morphological variation associated with ancient hybridization and quantified the

breadth of morphospace occupied by major lineages within the genus. The paleotropical lineages

of *Prunus* have diversified steadily since the late Eocene/early Oligocene, while the neotropical

lineages diversified much later. Critically, both the tropical and temperate lineages have

continued to diversify. We conclude that the genomic rearrangements created by reticulation

deep in the phylogeny of *Prunus* may explain why this group has been more successful than

other groups with tropical origins that currently persist only in either tropical or temperate

regions, but not both.

Keywords: allopolyploidy, herbarium specimens, hybridization, machine learning, niche

conservatism, tropical-to-temperate biogeographic transitions

INTRODUCTION

Bursts of speciation associated with changing environmental conditions have occurred

throughout the Tree of Life. Environmental factors, such as changes in climate, tectonic shifts,

mountain uplift, as well as biotic interactions, can all drive the diversification of lineages. In

particular, speciation may occur when lineages diversify to occupy newly available niches

resulting from changing environmental conditions. Classic evolutionary studies such as those

focused on Galapagos finches found that new ecological opportunities drove the rapid evolution

of new lineages (Schluter 2000). Climatic fluctuations have been implicated in promoting both

speciation and extinction in a variety of fauna (Weir and Schulter 2007). Changing

environmental conditions can have a particularly strong impact on plant lineages. For example,

increased global aridity likely promoted the diversification of both C4 grasses and succulent

lineages (Arakaki et al. 2011). Synthesis of current evidence suggests that many plant lineages

diversified in the tropics, sometimes spreading into temperate biomes (Spriggs et al. 2015), and

that many temperate lineages had a Boreotropical origin (Zhang et al. 2021, Nie et al. 2023).

Despite some evidence of shared biogeographic patterns, however, surprisingly few lineages

have successfully made transitions from tropical to temperate regions (Kerkhoff et al. 2014). In

fact, in the neotropics, the descendants of tropical ancestral lineages remained tropical in 94% of

woody angiosperms (Kerkhoff et al. 2014). Similarly, temperate lineages were quite conserved,

with 90% of temperate descendants arising from temperate ancestors (Kerkhoff et al. 2014).

These patterns of diversification are often attributed to the tropical conservatism hypothesis

(TCH) (Wiens & Donoghue 2004), which postulates that many lineages of tropical origin could

not adapt readily to the cooler, drier, and more seasonal temperate regions, and either were

extirpated (i.e., migrated to track tropical environments), or went extinct (Donoghue 2008).

91 In addition to environmental conditions, genomic changes can spark the diversification of

92 lineages. In some cases, reshuffling of genetic material may facilitate rapid diversification in

93 response to environmental change (Schenk 2021). A variety of genomic mechanisms, including

94 hybridization, gene duplication or loss, and/or genome doubling, can rearrange genomic material

95 such that new traits and adaptations develop to allow plant lineages to spread and diversify into

96 new niches and habitats (Xu et al. 2017, Hodel et al. 2022). Genome doubling may provide

97 extensive genetic variation and novelty upon which selection may act, driving adaptive

98 diversification (Seehausen 2004, Soltis and Soltis 2016, Doyle and Coate 2019, Griffiths et al.

99 2019, reviewed in Schenk 2021). Moreover, genomic changes in response to novel environments

100 may be coupled with morphological innovation (García-Verdugo et al. 2013). When lineages

101 encounter new environments, they may already possess phenotypes that facilitate their

102 occupancy of available niches in the community they enter, or the available niches in newly

103 encountered ecological conditions may drive diversifying selection on traits that are shaped to

104 promote survival in the novel niche space (Wellborn & Langerhans 2014). In some lineages,

105 success of diversification events in response to new ecological opportunities is mediated by

106 biological features as well as idiosyncrasies of the environmental conditions at the time

107 (Wellborn & Langerhans 2014). In summary, both evolutionary events and environmental

108 circumstances interact to shape the speciation and extinction rates that characterize

109 diversification (Spriggs et al. 2015).

110 One lineage that has been able to readily diversify in response to changing environments

111 was the genus *Prunus* (the plum genus, Rosaceae). This group contains approximately 250-400

112 evergreen and deciduous species that occur throughout the temperate regions of the northern

113 hemisphere and in the tropics and subtropics of both the Old and New Worlds (Rehder 1940,

Wen et al. 2008, Perez Zabala 2022). Species in this genus are key elements of both temperate

and tropical broadleaf forests. Despite its ecological and economic significance—*Prunus*

contains important crop species such as cherries, peaches, almonds, apricots, and plums—the

phylogenetic relationships of the major lineages in the genus are still unresolved. Historically,

the major groups within the genus were defined by inflorescence morphology—with three major

groups identified—solitary flower (e.g., peach), corymbose (i.e., producing a flat-topped

indeterminate cluster of flowers; e.g., cherries), and racemose (i.e., producing an indeterminate

inflorescence with the main axis not terminating in a flower; e.g., bird-cherries) (Rehder 1940;

Su et al. 2023). Multiple genetic studies using chloroplast DNA markers have inferred that the

solitary and corymbose groups form a clade that is sister to the racemose group (Bortiri et al.

2001, Wen et al. 2008, Chin et al. 2014, Zhao et al. 2016). However, nuclear markers have been

unable to resolve the backbone phylogenetic relationships of the genus (Lee & Wen 2001, Bortiri

et al. 2001, Wen et al. 2008, Chin et al. 2014, Zhao et al. 2016; Fig. 1). Furthermore, studies

using nuclear markers either relied on few nuclear loci (Wen et al. 2008, Chin et al. 2014, Zhao

et al. 2016) or had poor sampling in the understudied racemose group (Hodel et al. 2021).

Figure ... nic dat
evolutio ... ry of o
in the p ... r markers (left). One environmental variable, minimum
tempera ... butional limits of tropical taxa (upper right). Machine
learning ... cimens, can be used to infer how characters associated
with rep ... d to phylogeny (lower right).

he *Prunus* phylogeny have been obscured by

cytonu ... nesis that has been invoked to explain this

cytonu ... a an ancient allopolyploidy and/or

hybric ... hromosome count data, the corymbose and

solitar ... mose lineages are polyploid.

In summary, the evolutionary history of the plum genus remains unclear. To address

phylogenomic uncertainty, herein we assemble a phylogenomic dataset with hundreds of nuclear

loci and entire chloroplast genomes for each of 99 *Prunus* species representing all major lineages

in the genus. Our taxon sampling is the most complete to date, especially in the understudied

racemose group. We use the phylogeny of *Prunus* as a framework to investigate reticulation

events, to reconstruct the biogeographic history, and test hypotheses regarding the morphological

basis of key biogeographic transitions and reticulation. The morphological characters associated

with ancient biogeographic transitions and reticulation are notoriously difficult to measure in

extant species (e.g., McVay et al. 2017). Accordingly, we developed a novel approach to

quantify and categorize morphological variation: leveraging machine learning approaches with

digitized herbarium sheet image data (Fig. 1). Our specific objectives in this study are to:

1)   Resolve the phylogenetic relationships of major groups within the genus.

2)   Assess the role of ancient genomic rearrangements—specifically allopolyploidy

and/or hybridization—in shaping the evolutionary history of *Prunus*.

3)   Clarify the biogeographic history of *Prunus*, especially concerning the timing and

frequency of transitions between tropical and temperate regions.

4)   Implement a machine learning approach with thousands of digitized herbarium

specimen images to test for morphological evidence associated with biogeographic

transitions and reticulation events such as allopolyploidy and hybridization.

163
164
165
MATERIALS AND METHODS

Genomic data collection

Hyb-Seq probe design

We designed a 610-locus custom Hyb-Seq probe set to target nuclear genes in *Prunus*. Our

approach aimed to obtain genes with the highest probability of being strictly single-copy (i.e.,

single-copy nuclear genes; hereafter SCNs). We used an iterative process to BLAST publicly

available genomes against themselves to obtain a candidate pool of putatively SCN loci

(Supplemental Fig. S1); the subsequent analyses were conducted in Geneious Prime 2020.0.5

(https://www.geneious.com). We first conducted BLAST searches of the annotated genomes of

P. persica (L.) Batsch (peach, PRJNA31227; Verde et al. 2013), *P. avium* (L.) L. (sweet cherry,

PRJDB4877; Shirasawa et al. 2017), and *Malus domestica* (Suckow) Borkh. (PRJNA339703,

Daccord et al. 2017) against themselves using an e-value of 1e-10, which yielded 11,305, 10,703,

and 968 candidate SCNs, respectively. We used *Malus domestica* (apple), which is in the same

subfamily as *Prunus* – the Amygdaloideae – to expand the phylogenetic breadth of the baits. We

then BLASTed the candidate SCNs from *P. avium* against the SCNs from *P. persica*, and filtered

out loci that had multiple hits, were fewer than 300 nucleotides, and had less than 95.4%

pairwise identity. Loci with multiple hits were presumably not strictly single-copy, loci with

fewer than than 300 nucleotides have less phylogenetic information than longer loci, and loci

with pairwise identity close to 100% may not be variable enough to be informative (Weitemier et

al. 2014). This left 318 loci with a total of 262,684 nucleotides of sequence, and we retained the

sequences from *P. avium* for bait design. To obtain additional loci more closely related to

racemose *Prunus* species, we also BLASTed the candidate loci that passed the above length and

pairwise identity filter, and had two or fewer hits, against the *P. serotina* Ehrh. (black cherry)

transcriptome (Swenson et al. 2017; available via the Hardwood Genomics Project,

hardwoodgenomics.org). This yielded 96 additional SCNs with 72,172 nucleotides of sequence,

and the *P. serotina* sequence was retained for bait design. We also BLASTed the candidate

SCNs from *P. avium* against the SCNs from *Malus domestica*; after filtering out loci with

multiple hits, fewer than 600 nucleotides, and less than 80% or greater than 87.5% pairwise

identity, we retained 160 loci totaling 213,990 nucleotides. Here, we used sequences from *P.*

avium for bait design. Finally, because inflorescence architecture has historically been used to

define phylogroups in *Prunus*, we selected 36 functional genes that may be associated with

flowering (e.g., APETALA, FT; Yao et al. 2022). This gene set included 36 loci totaling 137,252

nucleotides, and we used *P. avium* sequences for bait design. In total, we selected 610 loci with a

total of 686,098 nucleotides for our custom *Prunus* bait set (Arbor Biosciences, Ann Arbor, MI).

In the Rosaceae, a custom bait set may perform better than a universal set (Ufimov et al. 2021).

Field collection, DNA extraction, library preparation, and hybridization reactions

Specimens were collected from the field (Supplemental Table S1) and stored in silica gel

until DNA was extracted from leaf tissue using a modified CTAB protocol (Doyle & Doyle

1987). Next, DNA libraries were constructed using a KAPA HyperPrep kit (available from

Roche, Basel, Switzerland) following the manufacturer's protocol except with quarter-volume

reactions for each individual. Briefly, DNA extractions were fragmented via sonication using the

QSonica ultrasonicator (Newtown, Connecticut, USA), uneven ends were repaired and A-tailed,

and Illumina adaptors were ligated to the DNA fragments. Next, AMPure magnetic beads were

used to purify and size-select the adaptor ligated DNA, and the DNA libraries were PCR

amplified to add unique i5 and i7 indexed oligonucleotides (i.e., barcodes). Libraries were

pooled in groups of eight with the criteria of balancing samples among presumed phylogenetic

distance and concentration prior to the hybridization capture reaction with custom-designed

baits. DNA libraries were hybridized with the baits for 48 hours at 60°C following the MyBaits

v4 manual (Arbor Biosciences, Ann Arbor, Michigan, USA). The hybridization enriched

libraries were combined with unenriched DNA libraries in a 60:40 ratio to enable generation of

plastomes from off target reads. All enriched and unenriched libraries were combined into a

single tube and sent for 2x150bp sequencing on the Illumina HiSeq 4000 at Novogene

(Sacramento, California, USA). We also acquired sequence data for one species from NCBI

GenBank (Supplemental Table S1). In total, we generated or obtained data for 119 accessions

(Supplemental Table S1) representing 101 species, which include 99 *Prunus* species and two

outgroups (*Lyonothamnus floribundus* A.Gray and *Physocarpus opulifolius* (L.) Maxim.). These

outgroups were selected because *Lyonothamnus* is likely the sister lineage of *Prunus* (Xiang et

al. 2017), and *Physocarpus* is more distantly related but within the Amygdaloideae (Xiang et al.

2017, Zhang et al. 2017).

Plastome assembly and phylogenetic inference

First, raw sequencing reads were quality filtered and Illumina adapters were removed using

bbduk (https://sourceforge.net/projects/bbmap/). GetOrganelle (Jin et al. 2020) was used to

generate plastomes for each species using the reads cleaned by bbduk. Of the 119 accessions

included in the final dataset, 75 produced complete, circular plastomes after an initial run

through GetOrganelle. For the remaining 44, we used minimap2 (Li 2018) implemented in

Geneious Prime 2020.0.5 (https://www.geneious.com) to assemble the contigs and scaffolds

from GetOrganelle, using the *Prunus avium* plastome (NCBI accession number MK622380) as a

reference to generate as complete as possible plastomes. After using minimap2 on the non-

circularized plastomes, we obtained nearly complete plastomes for all accessions. To ensure that

the use of *Prunus avium* as a reference did not bias the plastome results, we constructed a

phylogeny using the plastomes we generated, as well as all publicly available *Prunus* plastomes

(N = 17), to check that each newly sequenced accession is placed reasonably in the phylogeny

(Supplemental Table S2). The plastome phylogeny was inferred using RAxML with 100 rapid

bootstrap replicates and 20 independent maximum likelihood searches (i.e., the following

parameter settings: "-f a -m GTRGAMMA -p 12345 -x 12345 -# 100"). All analyses, unless otherwise

stated, were run on the Smithsonian Institution High Performance Cluster (SI/HPC, "Hydra")

(https://doi.org/10.25572/SIHPC).

Hyb-Seq locus assembly

We used the software package HybPiper v2.16 (Johnson et al. 2016) to assemble nuclear

loci. The same quality filtering and Illumina adapter removal strategy as with the plastome

pipeline was used with the nuclear data (i.e., cleaning and trimming using bbduk). We followed

the core scripts of the HybPiper pipeline; we first used hybpiper assemble –run_intronerate to map

reads using BWA (Li & Durbin 2009), sorted reads into fasta files, and ran the SPADES

assembler (Bankevich et al. 2012). Next, the command hybpiper stats was used to get the lengths

of the recovered gene sequences, visualize the success of each gene for each species, and

generate a file of descriptive statistics for the assembly (Supplemental Table S3, Supplemental

Fig. S2). The command hybpiper retrieve_sequences was used to retrieve exons, introns, and

supercontigs from each gene region for every species, and put them in an unaligned fasta file.

Next, for each type of locus (i.e., exons, introns, and supercontigs), the unaligned gene files were

aligned using MAFFT (Katoh & Standley 2013) with the following parameter settings: "--

maxiterate 5000 --auto --adjustdirectionaccurately –leavegappyregion." The resulting alignments were

trimmed using the phyx command pxclsq with the proportion of sites required to have data (-p

option) set at 0.5 (Brown et al, 2017). For each of the trimmed alignments, gene trees were

estimated using RAxML (Stamatakis 2014) with 100 rapid bootstrap replicates and 20

independent maximum likelihood searches (i.e., the following parameter settings: "-f a -m

GTRGAMMA -p 12345 -x 12345 -# 100"). After confirming that exons and supercontigs produced

similar species tree topologies (Supplemental Fig. S3), we used supercontigs for all subsequent

analyses.

Species tree estimation

The nuclear species tree was estimated using ASTRAL-III (Zhang et al. 2018), an

approach consistent with the coalescent, to summarize the 587 gene trees. We used the ASTRAL

quartet scores to assess nodewise support for inferred phylogenetic relationships. A

concatenation approach implemented in RaXML, with 100 rapid bootstrap replicates and 20

independent maximum likelihood searches, was also used for comparative purposes. For some

downstream analyses (e.g., divergence dating), it was necessary to have a phylogeny with

meaningful branch lengths. For these analyses, we used the ASTRAL topology as a constraint

tree for a RAxML tree search to obtain a species tree with the ASTRAL topology but with

proportional branch lengths.

Gene tree discordance analysis

To assess congruence between gene trees and the species tree, we used the program

phyparts (Smith et al. 2015). At each node in the tree, phyparts compares the rooted gene tree

topologies with the rooted species tree to label the number of genes that are concordant,

discordant, and uninformative relative to the species tree. Gene trees were rooted for the phyparts

analysis and we considered any gene trees with less than 50% bootstrap support at a given node

to be uninformative for that node (i.e., -s 50 option).

Paralogs

We used the tree-based paralog detection implemented in HypPiper 2.0+ to distinguish

between true orthologs and paralogs. The HybPiper post-processing command hybpiper

paralog_retriever was used to retrieve the multiple sequences for putative paralogous genes. Next,

the unaligned retrieved gene sequences were inputted into a phylogenetic pipeline that included

alignment with MAFFT (Katoh & Standley 2013) and phylogeny construction with FastTree

(Price et al. 2009). We manually examined the resulting phylogeny for each gene. In some cases,

genes suspected to be paralogs formed a clade, indicating that the putative paralogs may be

alleles and not the result of duplication. Alternatively, genes that were truly paralogs would

present as multiple copies dispersed throughout the phylogeny inferred by FastTree. For all 610

genes, we classified paralogy status based on the presence or absence of paralogs. If paralogs

were detected, we manually inspected the paralog tree for such genes to determine if the

suspected paralogs clustered, or were in fact true paralogs dispersed throughout the gene tree. In

the cases where true paralogs were detected, we discarded that gene from the analysis. In the

cases where the main contig and additional contigs clustered and formed a clade, we retained the

primary copy outputted by HybPiper. We discarded 23 genes that had true paralogs, and retained

587 genes for downstream phylogenetic analyses.

Hybridization and allopolyploidy analyses

Because hypotheses of ancient hybridization and allopolyploidy have been proposed to

explain the origin and diversification of this group, we used phylonet v3.8.2 (Than et al. 2008) to

investigate potential reticulation. We assembled 20 haphazardly sampled datasets each consisting

of seven taxa, representing the major lineages—one species each from the corymbose, solitary-

flower, and temperate racemose clades, three from the tropical racemose clade, and an outgroup

(i.e., *Physocarpus opulifolius*). We constructed rooted gene trees for each gene in each of the 20

datasets using RAxML (Stamatakis 2014) with 100 rapid bootstrap replicates and 20 independent

maximum likelihood searches. We used the maximum pseudolikelihood approach (Yu &

Nakhleh 2015) to infer networks with maximum reticulations set to 1, 2, and 3. We ran multiple

replicates (n=10) for each dataset with each of the three maximum reticulation values. For each

dataset, the networks with the optimal pseudolikelihood scores were retained and visualized

using Dendroscope (Huson & Scornavacca 2012).

To investigate histories of allopolyploidy, and for comparison with reticulation results

from phylonet, we used the program GRAMPA (Thomas et al. 2017). This approach uses a least

common ancestor mapping algorithm to reconcile gene trees and species trees (Goodman et al.

1979; Page 1994) so that polyploidy events can be placed on a phylogeny. GRAMPA can

identify modes of polyploidy as well as place whole genome duplications (WGDs) on a

phylogeny, and infer parental participants in instances of polyploidy. The user can input clades

of interest that are suspected to have a polyploid origin, or use a global search that considers if

all clades could be the result of WGD. We used the former approach to target specific major

clades that were hypothesized to have an ancient polyploid origin. Specifically, we tested the

nodes defining the following clades: temperate racemose, solitary-flower, corymbose, all

temperate ('Temp'), tropical ('Trop'), core tropical ('CTrop'), Australasia paleotropical ('APal'),

South American neotropical ('SNeo'), and North American neotropical ('NNeo') (see Fig. 2A

for clade definitions). In the analysis, we allowed any other node in the tree to be a parental

lineage to any clades determined to be of polyploid origin. For hypothesized instance of

polyploidy, GRAMPA assesses if a standard singly-labeled species tree or a multi-labeled tree is

the most parsimonious explanation of the data.

To complement the GRAMPA analysis and investigate all nodes in the phylogeny, we

used the approach of Yang et al. (2017) to map duplications to nodes in the species tree. Briefly,

we mapped duplication events within orthogroups to the ASTRAL species tree. For a given

subclade, if two or more taxa overlapped between daughter clades, a gene duplication event was

counted at the node which was defined as the most recent common ancestor of the subclade on

the ASTRAL species tree. We required the average bootstrap percentage for each orthogroup to

be >50.

Environmental characterization

To quantify the environment for major groups, we used a Principal Components Analysis

(PCA) to synthesize variation in the 19 WorldClim bioclimatic variables

(https://www.worldclim.org/) at a 2.5-minute resolution. The values for each of the 19 variables

were extracted for georeferenced herbarium specimens that represented each group using the R

packages 'dismo' (Hijmans et al. 2017) and 'raster' (Hijmans 2016). Two approaches were used

to quantify the environmental variation, first between two groups—temperate versus tropical—as

well as a comparison of four groups: temperate diploid (solitary + corymbose), temperate

racemose, neotropical racemose, and paleotropical racemose. We distilled the variation in 19

bioclimatic variables into 3 PC axes using the R package 'vegan' (Oksanen et al. 2017). For each

of the four groups, we defined 95% confidence interval ellipses, which indicate the region with

95% probability that the centroid is contained within the ellipse. For this analysis, we only used

the species for which we had genomic data, and a total of 10,011 herbarium specimen records

were included. Although this approach only captures present environmental variation, synthesis

has revealed that niches are more phylogenetically conserved than expected (Donoghue 2008),

and accordingly surveying all extant species in a lineage can give a reasonable approximation for

the environment in which a given lineage evolved. We also used the GPS data to collect

elevation data of each specimen using the R package 'rgbif' (Chamberlain et al. 2023). This was

done primarily to investigate whether tropical species occurred at lower or higher elevations,

because some high elevation tropical regions may resemble temperate environments more

closely than tropical ones, and this may impact our interpretation of biogeographic results.

Dating and biogeographic analysis

To investigate the timing and biogeographic history of *Prunus*, we used a biogeographic

ancestral range estimation analysis implemented in BioGeoBears (Matzke 2012, 2013). First, we

used treePL, a method for estimating divergence times on a phylogeny using penalized

likelihood, to infer divergence dates for all nodes in the *Prunus* phylogeny. Three calibrations

were used to date the phylogeny. We used *Prunus cathybrownae* (Benedict et al. 2011) from the

Early Eocene of North America as the most recent common ancestor (MRCA) of the diploid

(i.e., corymbose+solitary) lineages (stem lineage leading to node Dip; Fig. 2A), with a minimum

age of 50 million years ago (Mya). The fossil *Prunus wutuensis* (Li et al. 2011) from the early

Eocene of East Asia (Shandong, China) was used to fix the MRCA of crown *Prunus* to a

minimum age of 58 Mya. Next, the crown of the Amygdaloideae was fixed at a minimum age of

90 Mya based on a calibrated Rosaceae phylogeny from Xiang et al. (2017). Similarly, we

constrained the age of the stem Amygdaloideae to have a minimum age of 100 Mya using the

calibrated Rosaceae phylogeny from Xiang et al. (2017) as a guide. Using BioGeoBears (Matzke

2013), we considered six possible biogeographic models (DEC, DEC + J, DIVALIKE,

DIVALIKE + J, BAYAREALIKE, BAYAREALIKE + J) and selected the optimal model for the

data using AIC and AICc comparisons. Likelihood ratio tests were also used to determine if the

more parameter-rich +J models were preferred compared to the equivalent models without the +J

term.

We defined seven biogeographic regions based on geography: North America, South

America, Africa, Europe, West Asia, East Asia, and Australasia. In this analysis, the maximum

areas was set to two. We also used a biome-based approaches to delimit biogeographic regions.

Using the georeferenced herbarium data from the previous section ('*Environmental analysis*'),

we coded each species' presence in each of the 14 biomes delineated in the Ecoregions-17

dataset (Dinerstein et al. 2017). We combined similar biomes such that the BioGeoBears analysis

would be computationally feasible and the results readily interpretable. We defined four biome-

based biogeographic regions: 1) Dry (Desert & Xeric Shrubland/Mediterranean Forests,

Woodlands & Scrub); 2) Cold (Boreal Forests & Taiga/Montane Grasslands & Shrublands); 3)

Temperate (Temperate Broadleaf & Mixed Forests/Temperate Conifer Forests/Temperate

Grasslands, Savannas & Shrublands); 4) Tropical (Tropical & Subtropical Coniferous

Forests/Tropical & Subtropical Dry Broadleaf Forests/Tropical & Subtropical Grasslands,

Savannas & Shrublands/Tropical & Subtropical Moist Broadleaf Forests). In this analysis, we set

the maximum areas to four. The frequency and type of biogeographic events were estimated

using Biogeographic Stochastic Mapping (BSM) in BioGeoBears (Matzke 2013). For each of the

sets of biogeographic regions (i.e., continent-based, biome-based), we selected the optimal model

based on AIC and AICc comparisons and ran the BSM analysis on each biogeographic model,

with 50 independent mappings. The BSM approach can differentiate between biogeographic

events that occur at speciation nodes (i.e., cladogenesis events such as sympatric speciation,

vicariance, or founder events) versus transitions occurring along branches (e.g., anagenetic

dispersal).

Diversification analysis

To investigate the variation in diversification rates across the phylogeny, we implemented

an episodic birth-death model (EBD; Höhna 2015) and a branch-specific diversification model

(LSBDS; Höhna et al. 2019) in RevBayes (Höhna et al. 2016). The EBD model assumes

speciation and extinction rates are constant within each time interval but are allowed to vary

between time intervals. We used log-transformed rates following a Horseshoe Markov random

field prior distribution; this approach assumes that rates are autocorrelated. The LSBDS model

assumes rate heterogeneity across branches and does not require *a priori* assignment of shifts. In

other words, this model allows for a birth-death process with diversification rates that can vary

among branches. We used exponential priors for both diversification and extinction rates and

used 50,000 generations of reversible-jump Markov chain Monte Carlo, with 25% burnin, to

sample models with a variety of rate shift placements.

The dated species tree inferred from ASTRAL, with node ages calibrated using penalized

likelihood in *treePL*, was used as input for the EBD and LSBDS models. We report the

speciation rate, extinction rate, relative extinction rate, and net diversification rate. The software

CRABS (Congruent Rate Analyses in Birth–death Scenarios; Höhna et al. 2022) was used to

assess heterogeneity in diversification rates, and if rate patterns were robust to the non-

identifiability of the birth–death model. Specifically, we assessed the estimated posterior median

from the EBD model to compare speciation and extinction rate over time. The posterior median

is considered a robust estimate for such rates (Magee et al. 2020).

Morphological data collection

Herbarium image analysis

428   Because of the rich representation of *Prunus* species in museum collections, we used a

deep learning approach with digitized herbarium specimen sheets to test the degree to which

morphological classifications of species and/or lineages corresponded to phylogenetic

hypotheses. We assembled a dataset of 4,228 images representing the 99 *Prunus* species with

genomic data in this study. These images were downloaded using the 'idig_search_media'

function in the 'ridigbio' R package (Michonneau et al. 2016). We used exiftool

(https://exiftool.org/) to remove EXIF metadata from the images using the command "exiftool -

overwrite_original -EXIF= *.jpg"). This approach removes metadata often incompatible with image

processing libraries commonly used in machine learning. We also trimmed the edges of the

digitized images before using them in downstream machine learning analyses. This was done to

remove non-biological information, such as herbarium labels, color bars, and scale bars, from the

digitized herbarium sheet. We removed the top 20% and bottom 20% of every image used in

machine learning analyses. This unnecessarily removed some biologically relevant features, but

was conservative for ensuring the removal of non-biologically meaningful information on the

images. We investigated both trimmed and untrimmed sheets to test the impact of removing the

bottom 20% and top 20% of each sheet; hereafter these sheets are referred to as 'trimmed' or

'whole sheet', respectively. Furthermore, because we had thousands of digitized herbarium

sheets available, this approach represented a reasonable balance between removing too much

data and ensuring that any data with the potential to bias results was excluded.

Classification Algorithms

To assess if morphological variation corresponded to genetic variation, we developed a

supervised machine learning classifier algorithm using the fastai (https://github.com/fastai/fastai)

deep learning library, which is built using PyTorch libraries (https://pytorch.org), to classify

herbarium sheet images into categories. The labels for each category were assigned based on

clades inferred using genomic data. We used two approaches for the classification analysis. First,

we assigned labels of 'diploid species,' corresponding to all species with solitary and corymbose

inflorescences, and 'tropical racemose,' which referred to all tropical species with racemes.

These two groups were selected because the phylogenetic relationship between these two groups

was the same in the nuclear and chloroplast phylogenies. This model was trained and validated

using a total of 5,045 herbarium sheet images representing 81 species, with 2,746 in the

solitary/corymbose group, and 2,301 in the tropical racemose group. In each group, 80% of the

labeled input data were used for training, and 20% for validation. The model was run for 24

epochs until an accuracy rate of 97.2% was achieved. As an additional check that the model was

performing well, we tested hundreds of specimens from species in either the solitary/corymbose

groups (N = 452), or the tropical racemose groups (N = 426), which were not species used to

train or validate the model. Next, the additional 'test' data in this implementation were species

from the temperate racemose group (N = 676). This group was selected as test data because its

phylogenetic position was different between the nuclear and chloroplast topologies. Essentially,

this machine learning classification approach was used to test if there were morphological

signatures of hybridization in the temperate racemose group that correspond to the genomic

signatures of hybridization detected. Although the model was not trained to classify temperate

racemose species, we expect that if the temperate racemose group arose via hybridization,

species in this group may retain some morphological features from both the solitary/corymbose

group and the tropical racemose group, if these two groups are the parental participants in

ancient hybridization. The second classifier model we developed had the goal of distinguishing

between four major groups: temperate diploid, temperate racemose, neotropical racemose, and

paleotropical racemose. This second model was trained and validated using a total of 4,117

herbarium sheet images representing 81 species. In each group, 80% of the labeled input data

were used for training, and 20% for validation. The model was run for 24 epochs until an

accuracy rate of 91.3% was achieved. This second classification model was built so we could

ultimately measure the breadth of morphological variation across specimens that could be

successfully categorized in one of the four groups using a dimensionality reduction approach

called IVIS (described below).

Gradient CAM

As a further ground-truthing step to ensure that the classification algorithm was assessing

biologically meaningful variation, as opposed to learning biologically meaningless features for

identification (e.g., labels, rulers), we used Gradient Class Activation Mapping (Grad-CAM), an

approach for visualizing how deep neural networks make their predictions (Selvaraju et al.

2017). Briefly, Grad-CAM generates a heatmap highlighting the regions in an input image most

important for why a particular prediction was made. A Grad-CAM algorithm takes the output of

a convolutional layer in the neural network and computes the gradient of target category in

relation to that layer's activations, which are the pixels that are 'activated' by the model—the

pixels that are most important for determining the class. The gradients of target category are then

unsampled and overlaid onto the query image to generate a heatmap highlighting pixels and

regions of the image most important for determining the prediction. Grad-CAM specifically

excels when applied to Convolutional Neural Networks (CNNs), due to their ability to learn and

extract useful features from images. By interrogating the pixels of an image that the model

considers important, Grad-CAM can inform why and how a CNN makes its predictions. We

sampled the final layer of the CNN, as well as the second-to-last and third-to-last layers. We

categorized the heatmap images as classifying based on leaf features, floral features, or

neither/ambiguous. Each image with both leaf and floral structures present was classified as

'floral', 'mixed', or 'leaf'. We also assigned images that did not fall into the above categories as

'leaf with no floral structure present', 'ambiguous', or 'problematic'. The last two categories

were for images where it was difficult to tell if the heatmap favored either floral structures or

leaves, or if the heatmap was overly focused on non-biological information, respectively.

Morphological space in tropical and temperate groups

IVIS (Implicit Variational Information Sharing) is a deep learning approach with the goal

of mapping high-dimensional data into low-dimensional space. Critically, IVIS works well with

image data and can preserve distances between samples in a global sense, which is not always

true using other dimensionality reduction methods such as PCA, t-SNE, or umap (Szubert et al.

2019, Chari et al. 2021). In the case of image data, IVIS works by encoding the input image into

a lower-dimensional representation, and then using a decoder to reconstruct the image. During

training, the network learns to form a compact representation by minimizing the reconstruction

loss and encouraging the low-dimensional space to be continuous and well-structured through

the use of a regularization term. The low-dimensional representation learned by IVIS can be used

for various tasks, such as clustering, visualization, or further fine-tuning for classification tasks.

IVIS has been shown to be effective for image data, outperforming traditional methods such as

PCA and t-SNE in terms of clustering and visualization quality. Here, we use IVIS to reduce the

highly-dimensional variation present in image data into 2-dimensional space to make

comparisons among specimens and groups.

522   **RESULTS**

Phylogenomic analyses

For 99 *Prunus* species representing all major lineages, we assembled a dataset of 587

nuclear genes and complete chloroplast genomes to infer phylogeny. The nuclear coalescent

species tree inferred using ASTRAL (Fig. 2A) recovered monophyletic corymbose and solitary

groups, but a paraphyletic racemose group. With our robust sampling, the solitary, corymbose,

and temperate racemose species form a clade, although with modest support (37.9% ASTRAL

quartet support (QS) score; see node Temp ("Temperate"), Fig. 2A). This clade of solitary,

corymbose, and temperate racemose taxa was sister to a large clade composed of tropical

racemose species, including representatives from both the pics and paleotropics. The monophyly

of each of the solitary and corymbose clades was strongly supported (i.e., > 70% QS; Fig. 2A).

The sister relationship between the solitary and corymbose clades also had strong support (73.9%

QS score; node Dip ("Diploid"), Fig. 2A). There were also several other strongly supported

clades, including the paleotropical subgenus *Pygeum*, and the clade of neotropical racemose

species (see nodes APal ("Australasian paleotropical"), EPal ("East Asian paleotropical") and

SNeo ("Neotropical, predominantly South America"), respectively; Fig. 2A).

The paleotropical racemose species did not form a clade – the neotropical racemose

species were nested within the paleotropical group (Fig. 2A). A clade composed of nearly all

tropical racemose species, except for a few early-diverging lineages (*Prunus undulata* Buch.-

Ham. ex D.Don, *P. laurocerasus* L., *P. africana* (Hook.f.) Kalkman, and *P. skutchii* I.M.Johnst.),

had relatively strong support (63.0% QS score; see node CTrop ("Core Tropical"), Fig. 2A).

Additionally, the New World (primarily neotropical) racemose species were not monophyletic.

Whereas the exclusively South American neotropical racemose species formed a strongly

supported clade, four other New World racemose species (*P. ilicifolia* (Nutt. ex Hook. & Arn.)

D.Dietr., *P. lyonni* (Eastw.) Sarg., *P. occidentalis* Sw., and *P. cornifolia* Koehne) formed a clade

with low support that was sister to the Australasian paleotropical racemose clade (Fig. 2A). The

concatenation tree also indicated that all temperate species formed a clade, albeit with low

support (42% bootstrap support; Supplemental Fig. S4). The concatenation tree was largely

congruent with the ASTRAL tree; one notable exception is that the early-diverging racemose

species *P. laurocerasus* and *P. undulata* were sister to the temperate clade in the concatenation

phylogeny (Supplemental Fig. S4).

In contrast to the nuclear phylogeny, the chloroplast tree indicated that each group

defined by inflorescence morphology formed a clade (Fig. 2B). Each of the solitary-flower and

corymbose clades had high bootstrap support (100% and 99%, respectively), and the sister

relationship between these two clades was also strongly supported (99%; Fig. 2B). Within the

major clades, many of the relationships in the chloroplast phylogeny were similar to the nuclear

phylogeny. The neotropical racemose group was similarly non-monophyletic in the chloroplast

tree, and additionally the New World racemose species did not form a clade in the chloroplast phylogeny (Fig. 2B). Notably, two species that were early-branching racemose lineages in the nuclear tree (the European *P. laurocerasus* and the East and southeast Asian *P. undulata*), were nested within the temperate racemose species (Fig. 2B).

Figure 2A. The phylogenetic species tree inferred using ASTRAL to summarize 587 nuclear gene trees. Important nodes in the phylogeny are highlighted and named: temperate ('Temp'), diploid ('Dip'), tropical ('Trop'), core tropical ('CTrop'), East Asian paleotropical ('EPal'), Australasia paleotropical ('APal'), South American neotropical ('SNeo'), and North American neotropical ('NNeo'). The numbers at nodes are ASTRAL quartet support scores.

Figure 2B. The phylogenetic tree based on chloroplast genomes. Several key nodes that are congruent with the nuclear species tree are highlighted and labeled. The numbers at each node represent bootstrap percentage scores from the RAxML analysis.

Gene tree discordance

Many nodes in the nuclear species tree were characterized by rampant gene tree

discordance, as assessed by *phyparts* (Supplemental Fig. S5). At most nodes, there were far more

gene trees that were discordant with the species tree topology than were congruent

(Supplemental Fig. S5). Of note, fewer than 1% of gene trees were concordant with the species

tree at both the node defining the clade temperate racemose + solitary + corymbose, and the node

defining the tropical racemose clade (Fig. 2A). Within major clades, some nodes had high

concordance between gene trees and the species tree (e.g., the node defining a group of

neotropical racemose species), whereas others such as the node that defined the temperate

racemose species, had low concordance (Supplemental Fig. S5).

Reticulate topology

Given existing hypotheses of multiple hybridization events in *Prunus*, we investigated

multiple sets of species representing all major clades to better understand the variation in

reticulation events in this clade. The 20 networks with one reticulation often demonstrated

hybridization edges originating with the outgroup (Supplemental Fig. S6). The 20 networks with

two reticulation events frequently indicated hybridization with the outgroup, but also reticulation

within the focal clade (Fig. 3, Supplemental Fig. S6). Finally, the 20 networks with three

reticulations all indicated at least two reticulation events among the major lineages of *Prunus*

(Supplemental Fig. S6). Six of the 20 networks with one reticulation supported a hybrid origin of

the temperate racemose group. The other 14 one-reticulation networks had reticulation events

leading to the corymbose clade, tropical racemose clade, or corymbose/solitary-flower clade

(Supplemental Fig. S6). In contrast, 16/20 and 17/20 networks with two and three reticulations,

respectively, displayed reticulation implying a hybrid origin of the temperate racemose group

(Supplemental Fig. S6).

602
603

Figure 3. Two representative optimal 2-hybridization edge phylogenetic networks estimated using phylonet for seven species representing every major lineage in *Prunus* (top). In the network on the left, reticulation edges from the outgroup connect to the species representing the temperate racemose clade (*P. himalaica*) and the tropical racemose clade (*P. andersonii*, *P. caroliniana*, *P. polystachya*). In the network on the right, reticulation edges connect the temperate racemose species (*P. wilsonii*) to the outbroup, and also indicate that the temperate racemose species is sister to both the tropical racemose group (*P. henryi*, *P. brittoniana*, *P. pullei*) and the solitary-corymbose group (*P. persica*, *P. subhirtella*), suggesting hybridization leading to the temperate racemose group. The bottom portion shows the two most parsimonious GRAMPA results, which suggest an allopolyploid origin of the entire temperate clade (i.e., temperate racemose, solitary-flower, corymbose groups) (left; most parsimonious). The second-most parsimonious GRAMPA tree indicates an allopolyploid origin of the tropical racemose clade (right).

Allopolyploid origin: GRAMPA

The tests for signatures of allopolyploidy using the program GRAMPA indicated that at many of the nodes we investigated, a MUL-tree was more parsimonious than a singly-labeled

tree. The most parsimonious MUL-tree, with a parsimony score of 93,501, was characterized by

a duplicated clade that comprised all species in the temperate group (Fig. 3). The second most

parsimonious MUL-tree (score = 107,277) indicated an allopolyploidy event gave rise to the

tropical racemose clade (Fig. 3). We also detected evidence of allopolyploidy leading to the

temperate racemose clade (score = 115,580), tropical racemose clade (excluding early-diverging

species; score = 118,993), to the East Asian paleotropical clade (score = 126,618), and to the

Australasia paleotropical clade (score = 117,855) (Supplemental Fig. S7). Notably, when we

separately investigated the nodes defining the South American neotropical clade, the solitary-

flower clade, and the corymbose clade, the singly-labeled tree was most parsimonious in each

case, implying no evidence of allopolyploidy was detected at these nodes (parsimony scores of

126,710, 124,649, and 125,303, respectively; Supplemental Fig. S7).

The analysis to map WGDs to the species tree identified several nodes with relatively

high percentages of duplications. The crown *Prunus* node indicated a duplication percentage of

79.66% (Supplemental Fig. S8). The node with the second highest duplication percentage is the

one defining the tropical racemose species excluding the early diverging species (i.e., 'CTrop'

node from Fig. 2A) at 8.47% (Supplemental Fig. S8). Three other nodes within the tropical

racemose clade had duplication percentages >4%: the South American neotropical species

('SNeo' node from Fig. 2A), the Australasian paleotropical species ('APal' node from Fig. 2A),

and a small clade comprised of *P. australis*, *P. hypotricha*, *P. zippeliana*, and *P. spinulosa*

(Supplemental Fig. S8). Within the temperate clade, the node defining the solitary-flower group

had a duplication percentage of 4.24% (Supplemental Fig. S8).

Environmental characterization

The PCA of environmental variation based on the 19 bioclimatic variables captured

78.8% of variation on the first three principal components, with 46.5% of the variation

encapsulated in PC1, 17.3% in PC2, and 15.0% in PC3 (Fig. 4). One precipitation variable—

annual precipitation—was strongly positively correlated with PC1, and four temperature

variables—mean annual temperature, minimum temperature of the coldest month, mean

temperature of the driest quarter, and mean temperature of the coldest quarter—were also

strongly associated with positive PC1 space. Meanwhile, two temperature variables—

temperature seasonality, and mean annual temperature range—were strongly and negatively

associated with PC1 values. No bioclimatic variables were strongly correlated with PC2 or PC3.

In summary, tropical species occupy more positive regions of PC1, which correspond to warmer

and wetter conditions, with higher minimum temperatures in colder seasons, and less

temperature seasonality (Fig. 4), whereas temperate species trend towards negative values of

PC1. When considering the two types of tropical species—neotropical versus paleotropical—the

paleotropical species extended further towards both extremes of PC1, and further towards the

negative extent of PC2. Neotropical species extended slightly further into positive PC2 space

than paleotropical species. The temperate racemose species had a greater affinity for positive

values of PC1 whereas corymbose/solitary species trended further towards extreme negative

values of PC1. The temperate racemose species occupied a larger range of PC2 space than the

corymbose/solitary species (Fig. 4). There were clear elevational differences among species

representing the four major groups. In the temperate solitary/corymbose group, only two out of

28 species had a median elevation greater than 1,000m (Supplemental Fig. S9). In contrast, half

of the 10 temperate racemose species occurred at a median elevation greater than 1,000m, and

also half of the 10 neotropical racemose species were found at a median elevation greater than

1,000m. In the paleotropical racemose species, 13 of 28 species had a median elevation greater

tha

Figure 4. The summary of environmental variation using a Principal Components Analysis to distill the 19 bioclimatic variables into 3 dimensions. PC1 accounts for 46.5% of variation while PC2 explains 17.3% of environmental variation. Several bioclimatic variables strongly correlated with PC1 are indicated on the X-axis. Negative PC1 space is associated with increased temperature seasonality and annual temperature range. Meanwhile, positive PC1 space corresponds to higher annual precipitation, annual mean temperature, minimum temperature of the coldest month, and mean temperature of the driest quarter and coldest quarter.

Biogeographic analysis

The DEC+J model was the preferred model after comparing AIC scores of competing

geography-based BioGeoBears models, which were based on the time-calibrated tree generated

with TreePL (Supplemental Table S4). Likelihood ratio tests concluded that the addition of the J

parameter, despite producing a more parameter-rich model, led to a significantly better model

than DEC (Supplemental Table S5). In contrast, the BAYAREALIKE model was favored by

AIC scores for the biome-based BioGeoBears model (Supplemental Table S4). Here, likelihood

ratio tests did not indicate that the +J model was significantly better (Supplemental Table S5).

The geography-based analysis indicated an East Asian origin of the genus, with the initial

diversification occurring in the Paleocene ca. 60-55 Mya (Fig. 5). Subsequently, the temperate

racemose, solitary/corymbose, and tropical racemose clades all diverged by the early-mid

Eocene (ca. 50 Mya). This diversification occurred predominantly in East Asia and North

America, which were characterized by the Boreotropical climate during the Late Cretaceous –

early Eocene (ca. 65-50 Mya). The temperate racemose group steadily diversified in East Asia

and North America beginning in the mid-late Eocene (ca. 40 Mya). Both the solitary and

corymbose lineages began diversifying in the early Miocene (ca. 25-20 Mya), first in East Asia

and subsequently in North America and Europe. These two lineages currently occupy temperate

environments, and based on our sampling, they were more successful in speciating in East Asia

versus North America (Fig. 5).

Broadly, the tropical racemose clade's early diversification occurred in several regions,

including East Asia, West Asia, Africa, North America, and Europe, between 55-40 Mya. Early-

diverging lineages, including species such as *Prunus laurocerasus*, *P. undulata*, *P. africana*, and

P. skutchii, split off during the Eocene between ca. 55-40 Mya. Notably, the ancestors of these

few early-diverging lineages occupy a wide geographic range, including Europe, West Asia, East

Asia, Southeast Asia, Africa, North America, and South America. Subsequently, the patterns of

diversification occurred differently in the major lineages within the tropical racemose group. The

neotropical lineages, occurring in tropical North and South America, are characterized by a long

period of stasis, followed by rapid diversification in the Miocene beginning between 25-20 Mya

and lasting until ca. 10 Mya. In contrast, two major paleotropical lineages, in East Asia and

Southeast Asia, respectively, began steadily diversifying in the Oligocene (ca. 35 Mya), and

continued until approximately 10 Mya. Notably, there were multiple clades in both the temperate

and tropical groups of *Prunus* that demonstrated rapid diversification beginning in the early

Miocene.

The biome-based biogeographic analysis indicated that the ancestral biome state of the

clade consisted of both tropical and temperate biomes (Supplemental Fig. S10). In the portion of

the clade that presently occupies temperate environments (top clade; Supplemental Fig. S10),

there were transitions to other biome categories, including temperate (many corymbose species)

and dry/temperate (many solitary-flower species) beginning approximately 20-15 Mya

(Supplemental Fig. S10). In the temperate racemose clade, the biome-based analysis indicated

that part of this lineage retained its ancestral biome-affinity (green pies), whereas other species

transitioned from the tropical/temperate biomes to dry/temperate/tropical, dry/temperate,

temperate, or cold/temperate beginning around 15 Mya (Supplemental Fig. S10). In the tropical

racemose clade, the ancestral state of temperate/tropical transitioned to predominately tropical

environments from approximately 50 to 30 Mya (Supplemental Fig. S10). Then, portions of the

paleotropical racemose group in subgenus *Laurocerasus* shifted back to temperate/tropical

environments (top portion of tropical racemose clade; Supplemental Fig. S10). Meanwhile,

neotropical species either remained tropical, or transitioned to dry/temperate,

dry/temperate/tropical, or cold/tropical between 25-10 Mya (Supplemental Fig. S10). In the

bottom portion of the clade, lineages either remained tropical, or transitioned to cold/tropical or

temperate/tropical approximately 25-10 Mya (Supplemental Fig. S10).

The stochastic mapping analysis clarified the timing of transitions between geographic

regions and biomes (Supplemental Fig. S11). The vast majority of transitions have taken place

since 25 Mya—either as cladogenesis events at nodes—or anagenetic shifts along branches

(Supplemental Fig. S11). The anagenetic transitions in particular frequently occurred in the past

15 million years (Supplemental Fig. S11). In the geography-based analysis, the majority of

biogeographic transitions between regions was sympatric speciation, with smaller proportions of

founder events and anagenetic dispersal (Supplemental Table S6). Most biogeographic events in

the biome-based analysis were sympatric, meaning that there were speciation events without a

corresponding biome shift. Approximately 30% of the biogeographic events were anagenetic

dispersal—or transitions between biomes along a branch (Supplemental Table S6). In the

geography-based biogeographic analysis, East Asia acted as a major source, with especially

strong dispersal to N. America, Europe, and Oceania (Supplemental Table S7). Europe and West

Asia underwent substantial reciprocal dispersal, and North America acted as a source to South

America's sink (Supplemental Table S7). In the biome-based biogeographic analysis, both the

temperate and tropical biomes acted as key sources for other biomes (dry, cold). The dispersal

rate from tropical to temperate was larger than the rate from temperate to tropical (Supplemental

Table S7).

Figure 5. The BioGeoBears analysis (right) of the biogeographic history of *Prunus* using the DEC + J model based on seven biogeographic regions shown in the upper left: North America (blue), South America (red), Africa (magenta), Europe (turquoise), West Asia (green), East Asia (yellow), and Southeast Asia (orange). Color-coded pie charts at each node indicate the probability of the ancestral state occurring in a given biogeographic region. The episodic birth-death (EBD) model diversification analysis is shown in the lower left. The speciation rate and net diversification rate of the clade are highest during the Miocene, between approximately 20-10 Mya.

Diversification analysis

To investigate the variation in diversification rates across the phylogeny, we implemented an episodic birth-death model (EBD; Höhna 2015) and a branch-specific diversification model (LSBDS; Höhna et al. 2019) in RevBayes (Höhna et al. 2016). The EBD model estimated that the net diversification rate peaked approximately 20-10 Mya; this was driven by the speciation rate, which also was at its maximum between approximately 20-10 Mya (Fig. 5). The LSBDS

model, which shows branch-specific diversification, generally showed lower net diversification

in the early-diverging tropical racemose species, with similar net diversification in most other

lineages (Supplemental Fig. S12). The CRABS analysis confirmed results from the EBD model;

the extinction rate remained low, whereas the speciation rate increased to a peak approximately

10 Mya, followed by a decline (Supplemental Fig. S12).

Morphological analyses

We ran two experiments using computer vision-based machine learning analyses of

digitized herbarium sheets. The assessment of morphological characters associated with

hybridization revealed that whole sheet phenotypes of temperate racemose species, with the top

20% and bottom 20% of the sheet trimmed, were nearly evenly divided between classification as

'solitary/corymbose' (N = 310 observations with >90% probability) and 'tropical racemose' (N =

229 observations with >90% probability) (Fig 6). When using whole sheets (i.e., untrimmed),

there was a similar trend: there were N = 279 observations classified as 'solitary/corymbose'

with >90% probability and N = 223 observations classified as 'tropical racemose' with >90%

probability. The Grad-CAM analysis of internal model layers indicated that inflorescence vs.

leaf morphology was the driving factor in determining if the machine learning model inferred

whether to classify temperate racemose species as corymbose/solitary or tropical racemose (Fig.

6). Manual visual inspection indicated that the second-to-last and third-to-last layers were most

informative for determining classifications. The Grad-CAM results for the last three model

layers for every herbarium sheet specimen image are shown in Supplemental Tables S8 and S9.

Training and test data for all models are listed in Supplemental Tables S10 and S11.

Figure 6. In the upper left, the distribution of probability scores when using the temperate racemose species as test data in the first machine learning model. These classifications include specimens that were classified as either 'solitary/corymbose or tropical racemose'; regardless of classification, most specimens were assigned with high (i.e., >0.95) probability. The plot in the upper right shows the manual classifications we assigned when ground-truthing the gradient class activation maps for the 676 temperate racemose specimens for the second-to-last and third-to-last layer of the CNN. The bottom row shows the gradient class activation (Grad-CAM) maps for the second-to-last model layer for two representative temperate racemose specimens that were classified as 'tropical racemose' (left) and 'solitary/corymbose' (right), respectively.

In the second analysis, breadth of morphospace was measured using IVIS to quantify and

compare the extent of morphological variation present in the following groups: temperate diploid

(i.e., corymbose and solitary flower), temperate racemose, neotropical racemose, and

paleotropical racemose. The IVIS analysis determined that the two tropical groups exhibited a

greater range of morphological variation than the two temperate groups (Fig. 7). Within all

tropical specimens, the paleotropical racemose species occupied a greater expanse of

morphological variation than neotropical racemose (Fig. 7). The paleotropical species overlapped

in morphospace to a small degree with the temperate diploid group (i.e., corymbose/solitary) and

to a much greater degree with the neotropical racemose group. The neotropical species are

po   groups (Fig. 7).

Figure 7. The representation of whole specimen phenotypes in IVIS space. The specimen images were assigned to four groups according to environmental preference/clade: neotropical racemose, paleotropical racemose, temperate racemose, corymbose/solitary.

DISCUSSION

In this study, we provided increased resolution compared to previous studies of the

phylogeny of the economically and ecologically important genus *Prunus*, pinpointed key

genomic mechanisms promoting the diversification of this group, and improved our

understanding of its biogeographic history. By sampling more densely the understudied

racemose group and sequencing hundreds of nuclear loci and complete chloroplast genomes, we

changed our understanding of how several groups within the genus diversified. Specifically, by

combining analyses of hybridization, genome doubling, and cytonuclear and gene tree-species

tree conflict, we inferred that the polyploid racemose group is paraphyletic, a result found in

some previous studies using nuclear gene data (e.g., Lee & Wen 2001, Bortiri et al. 2002, 2006,

Wen et al. 2008, Chin et al. 2014, Zhao et al. 2016), but typically not found when using plastid

genes (e.g., Bortiri et al. 2001, Wen et al. 2008, Chin et al. 2014). The chloroplast phylogeny,

with a monophyletic racemose group, detected in the present study is congruent with many

previous studies using chloroplast markers (e.g., Bortiri et al. 2006, Wen et al. 2008, Chin et al.

2014, Zhao et al. 2016). Notably, our result conflicts with Su et al. (2023), which found a

monophyletic racemose group in both nuclear and chloroplast datasets. However, the nuclear

SNP markers used in Su et al. (2023) may have been unable to fully untangle the reticulate

history of this genus. Previous studies have suggested that cytonuclear discord observed in

Prunus phylogenies could be attributed to one (Chin et al. 2014) or multiple (Zhao et al. 2016)

allopolyploid hybrid origins of the racemose group. Here, we isolated specific lineages as

participants in likely hybridization and allopolyploidy events during the early Eocene ca. 50

Mya. These reticulation events early in the diversification of the genus help explain the high

gene tree-species tree conflict and cytonuclear discord. Through biogeographic, diversification,

and fossil-calibrated dating analyses, we traced the biogeographic history of the group and tied

key climatic events to radiations of subgeneric lineages.

Using several novel applications of machine learning algorithms to classify and quantify

morphological variation, we addressed how morphological evolution coincides with cytonuclear

discord, identifying characters essential for survival in specific environments and how

morphological variation in lineages impacts their biogeographic distribution. Phylogenomic data

have revealed that gene tree-species tree conflict is widespread and may sometimes arise via

ancient hybridization (e.g., McVay et al. 2017, Nie et al. 2023). Here, we demonstrate that a

well-trained machine learning classification model struggles to classify specimens in a lineage

that is the result of ancient hybridization, as these specimens that are the result of hybridization

display morphological features common to both putative ancestors. Additionally, we applied a

machine learning approach, IVIS, to quantify the morphological variation within major lineages,

revealing that the distinct morphospaces of different lineages may be shaped by their

biogeographic and evolutionary history. These morphological classifications, in combination

with environmental data, change our concept of how lineages may have diversified in the tropics,

and the conditions under which lineages transition from tropical to temperate biomes. Below, we

discuss and contextualize our results.

Improved understanding of Prunus *phylogeny, diversification, and biogeography*

Despite its economic and ecological importance, until now *Prunus* has not been

thoroughly investigated in a phylogenomic context with sufficient sampling in the racemose

group. Previous studies, based on chloroplast DNA and/or a few nuclear markers, inferred

possible cytonuclear conflict—a telltale sign suggesting hybridization—on the backbone of the

phylogeny. These analyses lacked the resolution to conclude whether the cytonuclear discord

was real or due to unresolved nuclear relationships (Lee & Wen 2001, Bortiri et al. 2001, Wen et

al. 2008). Based on cytonuclear discord (Chin et al. 2014), multiple copies of nuclear loci (Zhao

et al. 2016), nuclear reduced representation sequencing (Su et al. 2023), and chromosome count

data coupled with multiple gene tree topologies (Hodel et al. 2021), hypotheses of ancient

allopolyploid hybridization have been proposed (Zhao et al. 2016). Here, multiple lines of

evidence indicate a history of allopolyploidy and/or hybridization driving early diversification in

Prunus. First, we demonstrate that the cytonuclear discord along the backbone of the genus,

especially the differing phylogenetic placement of the temperate racemose group, provides a

specific hypothesis of reticulation (i.e., the temperate racemose group is a product of

hybridization and/or allopolyploidy). Phylonet analyses with 2- and 3-reticulations supported a

likely hybrid origin of the temperate racemose group. When 1-reticulation networks were also

considered, the reticulation analysis suggested possible hybrid and/or polyploid origins of the

tropical racemose group, solitary-flower clade, and corymbose clade (Fig. 3, Supplemental Fig.

S6). The GRAMPA analysis, which can detect and distinguish between different types of

genome doubling, also identified reticulation deep in the tree. The temperate clade was inferred

to have the best-supported allopolyploid origin deep in the phylogeny (Fig. 3, Supplemental Fig.

S7). However, additional clades, including the tropical racemose clade, also were inferred to

likely be the result of allopolyploidy (Fig. 3, Supplemental Fig. S7). Furthermore, mapping

WGDs to nodes in the phylogeny identify several major clades with evidence of duplication

(e.g., tropical racemose clade, excluding early-diverging species; Supplemental Fig. S8). Taken

together, these analyses point to reticulation deep in the phylogeny, although it is not constrained

to one instance of hybridization and/or allopolyploidy. These results are in line with previous

hypotheses that predicted multiple rounds of allopolyploidy (Zhao et al. 2016). Probably the

majority, if not all, racemose species, encompassing both temperate and tropical climates, are

polyploid. These species likely arose from multiple rounds of ancient allopolyploidy rather than

autopolyploidy. Only one of the GRAMPA analyses could possibly be autopolyploidy—the

Australasian paleotropical group (Supplemental Fig. S7C). All other WGD events demonstrated

that the duplicated clades were spread throughout the tree, necessarily implying allopolyploidy

(Supplemental Fig S7).

The higher resolution, time-calibrated plum genus phylogeny enhances our understanding

of the biogeographic history of the clade. The diversification and biogeographic history of

Prunus can be contextualized by existing hypotheses describing biogeographic patterns, and it

can shift our understanding of the range of possible outcomes for lineages that originated in the

tropics. Our biogeographic and diversification results show pulses of diversification in several

tropical and temperate *Prunus* lineages, with bursts of speciation in the early Miocene (Fig. 5,

Supplemental Fig. S10). This contrasts with steadier diversification reported in previous studies

(e.g., Chin et al. 2014). Our time-calibrated results also highlight the different patterns of

speciation in different tropical *Prunus* groups: relatively steady speciation in the paleotropics,

with punctuated bursts in the neotropics (Fig. 5). There were also key differences in greater

temperate success in Asia versus North America: the temperate lineages in Asia speciated readily

and rapidly while the species diversity in North America is depauperate. Although our sampling

of species diversity and nuclear loci gives better resolution than previous studies, there are still

limitations when interpreting our biogeographic results. Critically, biogeographic analyses

depend on bifurcating phylogenies, and many of our results point to a reticulate topology for

Prunus, especially deeper in the tree. It is likely that multiple lineages intermingled in the

expansive Boreotropical region early in the evolutionary history of the genus. The biome-based

biogeographic analysis is consistent with this explanation, where the deeper portion of the tree

showed all nodes as either tropical or temperate/tropical (Supplemental Fig. S10). Furthermore,

the biogeographic stochastic mapping analysis identified speciation in sympatry as the majority

of biogeographic events in both the geography- and biome-based analyses (Supplemental Table

S6). Many of the lineages implicated in reticulation, such as the early diverging tropical

racemose lineages, may have overlapped during the Eocene, thus facilitating ancient

hybridization and/or allopolyploidy. We must acknowledge that inferences of diversification

may be affected by sampling bias. We present the densest sampling of the understudied tropical

racemose clade to date, with accessions representing all major lineages. However, additional

future work is needed to ensure that even unbiased sampling does not impact the results, as this

genus is estimated to contain 250-400 species, depending on taxonomic treatments and sampling

of tropical species, necessitating further sampling in future studies.

Using Prunus *to inform broad biogeographic patterns*

For over a century, biologists have observed a latitudinal gradient in species diversity in

many clades across the Tree of Life, with greater species richness occurring near the equator.

However, we lack scientific consensus about the causes of this biogeographic pattern and several

hypotheses have been proposed. One explanation for the observed latitudinal gradient is the

tropical conservatism hypothesis (TCH), which posits that the relatively massive biodiversity of

the tropics can be primarily attributed to the geographic extent of tropical taxa over the past 55

million years and the subsequent evolutionary conservation of environmental niches (Wiens and

Donoghue 2004). Many groups have diversified in the Eocene, facilitated by the vast expanse of

the Boreotropics. Nevertheless, relatively few lineages have transitioned from tropical

environments to temperate ones. This phenomenon may be explained by the challenge of

acquiring the substantial adaptations necessary to tolerate the cooler conditions in temperate

zones (Donoghue 2008).

Another explanation is provided by the 'taxon pulse' hypothesis (Erwin 1985), which is

compatible with the TCH and posits that lineages that were able to transition from tropical to

temperate regions originated in tropical environments and then migrated in waves to more

temperate regions at higher altitudes and latitudes into increasingly harsh environments (Erwin

1985, Lü et al. 2020, Rapini et al. 2021). These radiations into temperate environments would

also be accompanied by species loss in ancestral tropical lineages as older lineages went extinct

(Liebherr and Porch 2015, Rapini et al. 2021, Nie et al. 2023). Some widespread plant clades,

such as *Viburnum*, match the taxon pulse hypothesis to an extent (Spriggs et al. 2015). In this

scenario, tropical lineages are posited to be 'dying embers' – lineages that diversified in the

tropics tens of millions of years ago, and persist in place, but have ceased to continue

diversifying (Spriggs et al. 2015).

Our results indicate *Prunus* does not precisely follow the predictions of either the TCH or

the 'taxon pulse' hypothesis. In *Prunus*, the paleotropical lineages have diversified in place and

do not appear to be 'dying embers', instead demonstrating steady diversification through time

(Fig. 5, Supplemental Fig. S12). As described in the taxon pulse hypothesis, tropical lineages

may migrate to higher temperate latitudes or to higher elevation regions in the tropics, which

may have temperate-like conditions. Biogeographic stochastic mapping suggested that in *Prunus*

there was substantial dispersal from the tropics to all other biome types (Supplemental Table S7).

Meanwhile, the diversification of the neotropical lineages, instead of occurring steadily as

observed in our paleotropical species, was characterized by bursts of speciation beginning in the

late Oligocene-early Miocene (ca. 25-20 Mya). In the tropics, some *Prunus* species have moved

to higher elevations, but there are also lineages that occupy truly tropical environments, as

defined by some or all individuals occurring in environments with the minimum temperature of

the coldest month greater than 18 ºC. Among species in our study, these include *Prunus*

dolichobotrys (Lauterb. & K.Schum.) Kalkman, *P. arborea* (Blume) Kalkman, *P. gazelle-*

peninsulae (Kaneh. & Hatus.) Kalkman, *P. javanica* (Teijsm. & Binn.) Miq*., P. undulata, P.*

fordiana Dunn*, P. grisea* Kalkman*, P. costata* (Hemsl.) Kalkman*, Prunus maingayi* (Hook. f.)

Wen*, P. oocarpa* (Stapf) Kalkman*, P. oligantha, P. schlechteri* (Koehne) Kalkman*, P. ceylanica*

(Wight) Miq*., and P. buergeriana* Miq. in the paleotropics, and *P. myrtifolia* (L.) Urb., *P.*

amplifolia Pilg., *P. cornifolia, P. integrifolia* (C.Presl) Walp*., P. occidentalis*, *P. skutchii*, and *P.*

reflexa (Gardner) Walp. in the neotropics. We also note that, for the species we sampled

genetically in this study, in both tropical groups, at least half of the species occurred at a median

elevation of less than 1,000 m (Supplemental Fig. S8). The biome-based biogeographic analysis

showed that some lineages and species in the tropics clear remained tropical over the past 20

million years (e.g., *P. andersonii, P. oleifolia, P. reflexa, P. amplifolia* in the neotropics, *P.*

dolichobotrys, P. gazelle-peninsulae, P. polystachya in the paleotropics; Supplemental Fig. S10).

At the same time, the analysis demonstrated partial or complete biome shifts in many tropical

species, such as from tropical to temperate/tropical, tropical to dry/temperate, and tropical to

cold/tropical (Supplemental Fig. S10).

The comparison of the neotropics and paleotropics demonstrates that diversification can

occur differently in distinct tropical regions—even within a single genus. Our quantifications of

the morphological breadth and environmental breadth revealed both to be greater in the

paleotropics as compared to the neotropics. The larger environmental space of the paleotropics

suggests there may have been greater ecological opportunity for lineages in the paleotropics

relative to the neotropics due to increased niche availability (Wellborn & Langerhans 2015). For

similar reasons, the greater morphological variation in the paleotropical *Prunus* shows these

lineages were able to leverage advantageous morphological features to adapt to new

environmental niches. This implies *Prunus* differs from many lineages with a tropical origin that

successfully transitioned to temperate regions. Synthesis has shown that not all lineages could

adapt to colder climates when tropical habitat retracted, and typically those lineages that tracked

tropical habitats became more restricted (Donoghue 2008). However, *Prunus* bucks this pattern,

exhibiting steady speciation in the paleotropics, and later bursts of speciation in the neotropics. It

is possible that the neotropical racemose lineages have ancestors that occupied temperate zones

in North America (Fig. 5). The small clade of four North American species (Fig. 2A; node

'NNeo') is a likely candidate—these species may be Boreotropical remnants in the New World,

highlighting the importance of North America as a site of diversification early in the evolution of

some tropical racemose species. Perhaps a history of surviving in the temperate zone enabled

present-day neotropical lineages to develop and/or keep a suite of morphological characters that

facilitated radiating into the neotropics. The substantial overlap in morphospace between

neotropical and temperate lineages suggests this may be the case (Fig. 7).

Why was Prunus *more successful than other lineages?*

Because of our sampling of species diversity coupled with hundreds of nuclear loci, we

identify allopolyploidy and/or hybridization in the backbone of the phylogeny, which may

explain the genomic basis of the rapid diversification 55-45 Mya. This allopolyploidy event is

supported by phylogenetic treatments of the Rosaceae, which found evidence of a WGD event at

the base of *Prunus* (Xiang et al. 2017); this also is corroborated by the high duplication

percentage at the base of the genus in our WGD mapping (Supplemental Fig. S8). This is an

important insight for not only understanding the phylogeny of the group but also its

biogeographic context. When numerous *Prunus* lineages were able to interact in the Boreotropics

during the Eocene, ample opportunities for hybridization and allypolyploidy arose. This genetic

reshuffling may explain why *Prunus* was more successful than other lineages at occupying both

temperate and tropical regions. The Rosaceae family has diversified into many successful

lineages (Potter et al. 2007), but even within this hyperdiverse group, *Prunus* stands out for its

diversity of inflorescence types, which may have facilitated migration via a variety of different

animal dispersers, as well as variation in leaf morphology supporting both evergreen and

deciduous life histories. One key innovation in *Prunus* is the evolution of leaf glands, which

appear to be associated with climatic conditions, with flat glands typical of species in tropical

climates, and raised glands present in species occupying cooler climates (Chin et al. 2013).

These glands may have had adaptive value due to their interactions with insects, with the flat

glands found in tropical regions preventing herbivory, and may be a key feature driving the

diversification of *Prunus*. Specifically, the variation in these morphological features may have

enabled *Prunus* radiations into the temperate zone, while continuing to diversify in the tropics.

Other Rosaceae, although successful, remained constrained in temperate regions (Xiang et al.

2017). Additionally, other groups with tropical origins that have experienced great success in

temperate regions did not retain much species diversity in tropical regions, including *Quercus* L.

(Hipp et al. 2020), *Viburnum* (Spriggs et al. 2015), Juglandaceae (Zhang et al. 2021), and

Saxifragales (Folk et al. 2019).

Morphological variation informs biogeographic history

The morphological variation associated with recent hybridization in many cases can be

readily tracked to inform the parental participants in hybridization, as well as to understand how

and why hybridization occurred, and whether hybrids may persist based on their environment.

However, in cases of ancient hybridization, it becomes difficult to tease apart the morphological

characters associated with hybridization since subsequent diversification and evolution in

response to environmental variation can obscure the circumstances surrounding the hybridization

event. Here, we used machine learning to quantify morphological variation in clades resulting

from reticulation, presenting a way to characterize the morphological legacy of

hybridization/allopolyploidy. By using the ancient hybridization detected via cytonuclear discord

and phylogenetic networks to guide our morphological analyses, we concluded that there were

morphological signals of hybridization in the present species. Members of the temperate

racemose group had morphological features of both the diploid clade and the tropical racemose

clade—the ancestors of these two lineages were putatively the participants in hybridization. So,

we would expect morphological features from both parents to persist in the lineage resulting

from reticulation. However, lineages' responses to environmental variation may also explain

patterns of morphological variation not resulting, or minimally resulting, from the genomic

results of hybridization (or allopolyploidy). In this case, we would expect that morphological

variation would co-vary to some extent with environmental variation. This would lead to an

expectation that the vast majority of test specimens from the temperate racemose group would be

classified as solitary/corymbose, because the environment overlaps much more between these

two groups than it does between temperate racemose and tropical racemose (Fig. 4). Contrary to

this expectation, we observed a relatively even split of temperate racemose species between the

two training categories.

Morphological shifts can result from transitioning to a novel environment, or existing

morphological variation can enable lineages to adapt to changing environmental conditions. The

key takeaway from our IVIS analysis is that both the neotropical and paleotropical groups exhibit

substantially more morphological variation than do the temperate groups, including both the

corymbose/solitary group, and the temperate racemose group. Moreover, the two temperate

groups occupy peripheral portions of morphological space on both IVIS axes 1 and 2, suggesting

that the morphological features in the temperate groups are specialized relative to the

morphological features of the tropical groups. These features may represent morphological

changes that promote transitions to temperate regions, diverging from morphological features

typical for tropical zones, in response to a transition from tropical to temperate biomes. Broadly,

if there is greater niche space available in the tropics—a proposed explanation for the increased

species diversity in these regions—we would expect greater morphological variation to occupy

more niches. In *Prunus*, the balance of morphological space we quantify is not surprising given

greater tropical diversity. Notably, however, the morphological space occupied by the

neotropical racemose species overlaps more with the two temperate groups than the paleotropical

group does. This suggests an explanation for some of the biogeographic patterns detected—

specifically that the more recent diversification of the neotropics was different from the initial

diversification in the paleotropics. By the time the neotropical lineages began to diversify,

lineages in the genus had already radiated into and adapted to temperate biomes. This may

explain the rapid diversification of the neotropical *Prunus*—the lineages may have already

acquired multiple morphological innovations for surviving in the New World's warm temperate

and tropical zones during the Oligocene-early Miocene.

1071

1072   *The promise of machine learning to assess morphology via digitized herbarium specimens*

1073   Machine learning approaches such as IVIS offer the promise of using whole-specimen

phenotyping with minimal preprocessing to classify species based on a biologically informed

hypothesis. These hypotheses could be based on phylogeny, environmental gradients, ecological

features, or geography. We demonstrate a method to use specimen data to investigate

correspondence between environmental and morphological variation (Fig. 8). Approaches such

as IVIS allow researchers to be agnostic regarding the morphological features studied, which can

avoid researcher-meditated bias in selecting traits. By grouping together specimens in different

and/or hierarchical groups, competing hypotheses can be tested. We demonstrate how specimens

can be grouped together to test hypotheses of hybridization. It is common to use the species as

the unit of classification with image-based machine learning algorithms. However, the species is

not necessarily the only level of biological organization that can be used to train classifiers.

Classifier algorithms that differentiate between species implicitly use a phenetic species concept

(de Queiroz 2007), which is not ideal for all research questions. We argue that groups above and

below species are useful, especially if they may be defined by shared phenotypic features; here

we demonstrate an application at a broader systematic level. Taking a phylogenetic approach to

grouping units to be classified, based on key phenetic differences, presents a way to mesh

phylogenetic and phenotypic data. Although there are benefits to using whole-specimen

phenotypes, there are also advantages to extracting individual traits, such as measurements of

leaf area and perimeter, as well as floral traits. These can be extracted via machine learning

methods such as segmentation in a high-throughput fashion.

1093

1094

Figure 8. A flow chart for designing phylogenetically-informed hypotheses to test using machine learning applied to
biological specimen image data.

Conclusions and future prospects

In this study, we employ phylogenomic, environmental, and morphological data to

establish that the initial diversification of the plum genus *Prunus* was driven by ancient

hybridization and/or allopolyploidy. Moreover, this complex history of reticulation may explain

the success of the tropical lineages of *Prunus* compared to other groups that also moved out of

the tropics and into temperate regions. To complement phylogenomic and environmental

analyses, we present innovative applications of machine learning algorithms to analyze the

variation in digitized herbarium sheets. We demonstrate inventive ways to harness computer

vision-based machine learning approaches to help us understand biodiversity across the Tree of

Life. Future studies in *Prunus* will require additional sampling, especially in the racemose group,

to further examine biogeographic patterns. Subsequent research may also focus on developing

segmentation masks to extract specific features from digitized image data. *Prunus* represents an

excellent model for testing the capabilities of machine learning algorithms using museum data,

given the wealth of publicly available digitized specimens in the genus.

1111

Data Availability

Data available from the Dryad Digital Repository: DOI: 10.5061/dryad.x95x69pwr; Reviewer

URL: http://datadryad.org/share/L-QUcxrgnpTr6l0Td3MxdfJDCUT1iRbPmT4SzA2LwMA.

Supplementary material, image data, and DNA sequence matrices are available from the Dryad

Digital Repository. Raw sequence data were submitted to NCBI GenBank (SUB13638423).

Machine learning models are hosted on Hugging Face with temporary URLs

(https://huggingface.co/richiehodel/Prunus_lineage_classifier;

https://huggingface.co/richiehodel/Prunus_herbarium_sheet_classifier). Final Hugging Face

URLs will be hosted by the Smithsonian Institution upon acceptance.

Jupyter notebooks for developing and processing machine learning models for image data are

available on GitHub

(https://github.com/richiehodel/machine_learning_Prunus_herbarium_sheets).

Acknowledgements

We thank the following funding sources: Smithsonian Institution Scholarly Studies Grant to JW,

EAZ, RGJH; Smithsonian Institution Peter Buck Fellowship to RGJH; National Museum of

Natural History NHRE Award to SKW.

We thank the following people for valuable discussions and assistance with analyses: Lauren

Ayers, Louis Cox, Greg Stull, Alicia Talavera.

All or portions of the laboratory and/or computer work were conducted in and with the support

of the Laboratories of Analytical Biology facilities of the National Museum of Natural History or

its partner labs.

Literature Cited

Arakaki M., Christin P.A., Nyffeler R., Lendel A., Eggli U., Ogburn R.M., Spriggs E., Moore M.J., Edwards E.J. 2011. Contemporaneous and recent radiations of the world's major succulent plant lineages. Proc. Natl. Acad. Sci. U. S. A. 108:8379–8384.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M., Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A. V., Sirotkin A. V., Vyahhi N., Tesler G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. J. Comput. Biol. 19:455–477.

Benedict J.C., DeVore M.L., Pigg K.B. 2011. *Prunus* and *Oemleria* (Rosaceae) flowers from the late early Eocene Republic flora of Northeastern Washington State, U.S.A. Int. J. Plant Sci. 172:948–958.

Bortiri E., Oh S.-H., Jiang J., Baggett S., Granger A., Weeks C., Buckingham M., Potter D., Parfitt D.E. 2001. Phylogeny and systematics of *Prunus* (Rosaceae) as determined by sequence analysis of *ITS* and the chloroplast *trnL-trnF* spacer DNA. Syst. Bot. 26:797–807.

Bortiri E., Oh S.-H., Gao F., Potter D. 2002. The phylogenetic utility of nucleotide sequences of sorbitol 6-phosphate dehydrogenase in *Prunus* (Rosaceae). Amer. J. Bot. 89:1697–1708.

Bortiri E., Vanden Heuvel B., Potter D. 2006. Phylogenetic analysis of morphology in *Prunus* reveals extensive homoplasy. Plant Syst. Evol. 259:53–71.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: phylogenetic tools for unix. Bioinformatics. 33:1886–1888.

Carranza-Rojas J., Goeau H., Bonnet P., Mata-Montero E., Joly A. 2017. Going deeper in the automated identification of Herbarium specimens. BMC Evol. Biol. 17:181.

Chamberlain S, Barve V, McGlinn D, Oldoni D, Desmet P, Geffert L, Ram K (2023). rgbif: Interface to the Global Biodiversity Information Facility API. R package version 3.7.6, https://CRAN.R-project.org/package=rgbif.

Chari T., Banerjee J., Pachter L. 2021. The Specious Art of Single-Cell Genomics. BioRxiv.:1–25.

Chin S.W., Shaw J., Haberle R., Wen J., Potter D. 2014. Diversification of almonds, peaches, plums and cherries - Molecular systematics and biogeographic history of *Prunus* (Rosaceae). Mol. Phylogenet. Evol. 76:34–48.

Daccord N., Celton J.M., Linsmith G., Becker C., Choisne N., Schijlen E., Van De Geest H., Bianco L., Micheletti D., Velasco R., Di Pierro E.A., Gouzy J., Rees D.J.G., Guérif P., Muranty H., Durel C.E., Laurens F., Lespinasse Y., Gaillard S., Aubourg S., Quesneville H., Weigel D., Van De Weg E., Troggio M., Bucher E. 2017. High-quality de novo assembly of the apple genome and methylome dynamics of early fruit development. Nat. Genet. 49:1099–1106.

De Queiroz K. 2007. Species Concepts and Species Delimitation. Syst. Biol. 56:879–886.

Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., Hahn, N., Palminteri, S., Hedao, P., Noss, R., Hansen, M., Locke, H., Ellis, E. C., Jones, B., Barber,

C. V., Hayes, R., Kormos, C., Martin, V., Crist, E., … Saleem, M. 2017. An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm. *BioScience*, 67: 534–545.

Donoghue M.J. 2008. A phylogenetic perspective on the distribution of plant diversity. Proc. Natl. Acad. Sci. U. S. A. 105:11549–11555.

Doyle J.J., Coate J.E. 2019. Polyploidy, the nucleotype, and novelty: The impact of genome doubling on the biology of the cell. Int. J. Plant Sci. 180:1–52.

Doyle J.J., Doyle J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. Phytochem. Bull. 19:11–15.

Folk R.A., Stubbs R.L., Mort M.E., Cellinese N., Allen J.M., Soltis P.S., Soltis D.E., Guralnick R.P. 2019. Rates of niche and phenotype evolution lag behind diversification in a temperate radiation. Proc. Natl. Acad. Sci. U. S. A. 166:10874–10882.

García Verdugo C., Friar E., Santiago L.S. 2013. Ecological role of hybridization in adaptive radiations:A case study in the *Dubautia arborea-Dubautia ciliolata* (Asteraceae) complex. Int. J. Plant Sci. 174:749–759.

Griffiths A.G., Moraga R., Tausen M., Gupta V., Bilton T.P., Campbell M.A., Ashby R., Nagy I., Khan A., Larking A., Anderson C., Franzmayr B., Hancock K., Scott A., Ellison N.W., Cox M.P., Asp T., Mailund T., Schierup M.H., Andersen S.U. 2019. Breaking free: The genomics of allopolyploidy-facilitated niche expansion in white clover. Plant Cell. 31:1466–1487.

Hijmans, R. J. 2016. raster: Geographic Data Analysis and Modeling. R pack- age version 2 5-8.

Hijmans, R. J., S. Phillips, J. Leathwick, and J. Elith. 2017. Dismo: Species distribution modeling. R package version 1:1-4.

Hipp A.L., Manos P.S., Hahn M., Avishai M., Bodénès C., Cavender-Bares J., Crowl A.A., Deng M., Denk T., Fitz-Gibbon S., Gailing O., González-Elizondo M.S., González-Rodríguez A., Grimm G.W., Jiang X.L., Kremer A., Lesur I., McVay J.D., Plomion C., Rodríguez-Correa H., Schulze E.D., Simeone M.C., Sork V.L., Valencia-Avalos S. 2020. Genomic landscape of the global oak phylogeny. New Phytol. 226:1198–1212.

Hodel R.G.J., Zimmer E., Wen J. 2021. A phylogenomic approach resolves the backbone of *Prunus* (Rosaceae) and identifies signals of hybridization and allopolyploidy. Mol. Phylogenet. Evol. 160.

Hodel R.G.J., Massatti R., Knowles L.L. 2022. Hybrid enrichment of adaptive variation revealed by genotype–environment associations in montane sedges. Mol. Ecol. 31:3722–3737.

Hodel R.G.J., Zimmer E.A., Liu B. Bin, Wen J. 2022. Synthesis of nuclear and chloroplast data combined with network analyses supports the polyploid origin of the apple tribe and the hybrid origin of the Maleae—Gillenieae clade. Front. Plant Sci. 12:3321.

Höhna, S. 2015. The time-dependent reconstructed evolutionary process with a key-role for mass-extinction events. *Journal of Theoretical Biology*, 380, 321–331.

Höhna S., Freyman W.A., Nolen Z., Huelsenbeck J.P., May M.R., Moore B.R. 2019. A Bayesian Approach for Estimating Branch-Specific Speciation and Extinction Rates. bioRxiv. 10.1101/555805

Höhna S., Landis M.J., Heath T.A., Boussau B., Lartillot N., Moore B.R., Huelsenbeck J.P., Ronquist F. 2016. RevBayes: Bayesian phylogenetic inference using graphical models and an interactive model-specification language. *Systematic Biology*, 65:726-736.

Höhna, S., Kopperud, B. T., & Magee, A. F. 2022. CRABS: Congruent rate analyses in birth–death scenarios. *Methods in Ecology and Evolution*, 13(12), 2709–2718.

Huson, D. H., & Scornavacca, C. 2012. Dendroscope 3: An Interactive Tool for Rooted Phylogenetic Trees and Networks. *Systematic Biology*, 61: 1061–1067.

Jin J.J., Yu W. Bin, Yang J.B., Song Y., DePamphilis C.W., Yi T.S., Li D.Z. 2020. GetOrganelle: A fast and versatile toolkit for accurate *de novo* assembly of organelle genomes. Genome Biol. 21:241.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics from high-throughput sequencing reads using target enrichment. Appl. Plant Sci. 4:1600016.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kerkhoff A.J., Moriarty P.E., Weiser M.D. 2014. The latitudinal species richness gradient in New World woody angiosperms is consistent with the tropical conservatism hypothesis. Proc. Natl. Acad. Sci. 111:8125–8130.

Li G., Davis B.W., Eizirik E., Murphy W.J. 2016. Phylogenomic evidence for ancient hybridization in the genomes of living cats (Felidae). Genome Res. 26:1–11.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34:3094–3100.

Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 25:1754–1760.

Li Y., Smith T., Liu C.J., Awasthi N., Yang J., Wang Y.F., Li C. Sen. 2011. Endocarps of *Prunus* (Rosaceae: Prunoideae) from the early Eocene of Wutu, Shandong Province, China. Taxon. 60:555–564.

Liebherr J.K., Porch N., Liebherr J.K., Porch N. 2015. Reassembling a lost lowland carabid beetle assemblage (Coleoptera) from Kauai, Hawaiian Islands. Invertebr. Syst. 29:191–213.

Lü L., Cai C.Y., Zhang X., Newton A.F., Thayer M.K., Zhou H.Z. 2021. Linking evolutionary mode to palaeoclimate change reveals rapid radiations of staphylinoid beetles in low-energy conditions. Curr. Zool. 66:435–444.

Magee, A. F., Höhna, S., Vasylyeva, T. I., Leaché, A. D., & Minin, V. N. 2020. Locally adaptive Bayesian birth-death model successfully detects slow and rapid rate shifts. *PLOS Computational Biology*, 16: e1007999.

Matzke N.J. 2013. Probabilistic historical biogeography: new models for founder-event
speciation, imperfect detection, and fossils allow improved accuracy and model-testing.
Front. Biogeogr. 5.

McVaugh R. 1951. A revision of the North American black cherries (*Prunus serotina* Ehrh.,
and relatives). Brittonia. 7:279–315.

McVay J.D., Hipp A.L., Manos P.S. 2017. A genetic legacy of introgression confounds
phylogeny and biogeography in oaks. Proc. R. Soc. B Biol. Sci. 284.

Michonneau F., Collins M., Chamberlain S.A. 2016. ridigbio: An interface to iDigBio's search
API that allows downloading specimen records. R package version 0.3.2.
https://github.com/iDigBio/ridigbio

Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep history
of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New
Phytol. 218:1668–1684.

Nie Z., Hodel R.G.J., Ma Z., Johnson G., Ren C., Meng Y., Ickert-Bond S.M., Liu X., Zimmer
E., Wen J. 2023. Climate-influenced boreotropical survival and rampant introgressions
explain the thriving of New World grapes in the north temperate zone. J. Integr. Plant
Biol. 65:1183–1203.

Oksanen, J. F., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin,
R. B. O'Hara, G. L. Simpson, P. Solymos, et al. 2017. vegan: Community ecology
package. R package version 2.4-4

Parins-Fukuchi C., Stull G.W., Smith S.A. 2021. Phylogenomic conflict coincides with rapid
morphological innovation. Proc. Natl. Acad. Sci. U. S. A. 118:e2023058118.

Perez Zabala, J. 2022. Taxonomic Diversity, Phylogeny, and Diversification of the
Environmental Niche of the Genus Prunus L. with emphasis on the New World Tropics.
UC Davis. Retrieved from https://escholarship.org/uc/item/3xw4b21k

Price M.N., Dehal P.S., Arkin A.P. 2009. FastTree: Computing large minimum evolution trees
with profiles instead of a distance matrix. Mol. Biol. Evol. 26:1641–1650.

Rapini A., Bitencourt C., Luebert F., Cardoso D. 2021. An escape-to-radiate model for
explaining the high plant diversity and endemism in campos rupestres. Biol. J. Linn. Soc.
133:481–498.

Rehder, A., 1940. Manual of cultivated trees and shrubs hardy in North America exclusive of
the subtropical and warmer temperate regions, 2nd ed. MacMillan, New York.

Schenk J.J. 2021. The next generation of adaptive radiation studies in plants. Int. J. Plant Sci.
182:245–262.

Schluter D. 2000. The ecology of adaptive radiation. Oxford University Press.

Schuettpelz E., Frandsen P.B., Dikow R.B., Brown A., Orli S., Peters M., Metallo A., Funk
V.A., Dorr L.J. 2017. Applications of deep convolutional neural networks to digitized
natural history collections. Biodivers. Data J.:e21139.

Seehausen O. 2004. Hybridization and adaptive radiation. Trends Ecol. Evol. 19:198–207.

Selvaraju R.R., Cogswell M., Das A., Vedantam R., Parikh D., Batra D. 2017. Grad-CAM: Visual explanations from deep networks via gradient-based localization. .

Shirasawa K., Isuzugawa K., Ikenaga M., Saito Y., Yamamoto T., Hirakawa H., Isobe S. 2017. The genome sequence of sweet plum (*Prunus avium*) for use in genomics-assisted breeding. DNA Res. 24:499–508.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals conflict, concordance, and gene duplications with examples from animals and plants. BMC Evol. Biol. 15:150.

Soltis P.S., Soltis D.E. 2016. Ancient WGD events as drivers of key innovations in angiosperms. Curr. Opin. Plant Biol. 30:159–165.

Spriggs E.L., Clement W.L., Sweeney P.W., Madriñán S., Edwards E.J., Donoghue M.J. 2015. Temperate radiations and dying embers of a tropical past: the diversification of *Viburnum*. New Phytol. 207:340–354.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 30:1312–3.

Su N., Hodel R.G.J., Wang X., Wang J.R., Xie S.Y., Gui C.X., Zhang L., Chang Z.Y., Zhao L., Potter D., Wen J. 2023. Molecular phylogeny and inflorescence evolution of *Prunus* (Rosaceae) based on RAD-seq and genome skimming analyses. Plant Divers.

Szubert B., Cole J.E., Monaco C., Drozdov I. 2019. Structure-preserving visualisation of high dimensional single-cell datasets. Sci. Rep. 9:1–10.

Than, C., Ruths, D., & Nakhleh, L. 2008. PhyloNet: A software package for analyzing and reconstructing reticulate evolutionary relationships. *BMC Bioinformatics* 9:1–16.

Ufimov, R., Zeisek, V., Píšová, S., Baker, W. J., Fér, T., van Loo, M., Dobeš, C., & Schmickl, R. 2021. Relative performance of customized and universal probe sets in target enrichment: A case study in subtribe Malinae. *Applications in Plant Sciences* 9(7).

Verde I., Abbott A.G., Scalabrin S., Jung S., Shu S., Marroni F., Zhebentyayeva T., Dettori M.T., Grimwood J., Cattonaro F., Zuccolo A., Rossini L., Jenkins J., Vendramin E., Meisel L.A., Decroocq V., Sosinski B., Prochnik S., Mitros T., Policriti A., Cipriani G., Dondini L., Ficklin S., Goodstein D.M., Xuan P., Del Fabbro C., Aramini V., Copetti D., Gonzalez S., Horner D.S., Falchi R., Lucas S., Mica E., Maldonado J., Lazzari B., Bielenberg D., Pirona R., Miculan M., Barakat A., Testolin R., Stella A., Tartarini S., Tonutti P., Arús P., Orellana A., Wells C., Main D., Vizzotto G., Silva H., Salamini F., Schmutz J., Morgante M., Rokhsar D.S. 2013. The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. Nat. Genet. 45:487–494.

Weir J.T., Schluter D. 2007. The latitudinal gradient in recent speciation and extinction rates of birds and mammals. Science. 315:1574–1576.

Weitemier K., Straub S.C.K., Cronn R.C., Fishbein M., Schmickl R., McDonnell A., Liston A. 2014. Hyb-Seq: Combining target enrichment and genome skimming for plant phylogenomics. Appl. Plant Sci. 2:1400042.

Wellborn G.A., Langerhans R.B. 2015. Ecological opportunity and the adaptive diversification
of lineages. Ecol. Evol. 5:176–195.
Wen J., Berggren S.T., Lee C.-H., Ickert-Bond S., Yi T.-S., Yoo K., Xie L., Shaw J., Potter D.
2008. Phylogenetic inferences in *Prunus* (Rosaceae) using chloroplast *ndhF* and nuclear
ribosomal *ITS* sequences. J. Syst. Evol. 46:322–332.
Wen J., Nie Z.-L., Ickert-Bond S.M. 2016. Intercontinental disjunctions between eastern Asia
and western North America in vascular plants highlight the biogeographic importance of
the Bering land bridge from late Cretaceous to Neogene. J. Syst. Evol. 54:469–490.
Wiens J.J., Donoghue M.J. 2004. Historical biogeography, ecology and species richness. Trends
Ecol. Evol. 19:639–644.
Xiang Q.-Y., Soltis D.E., Soltis P.S. 1998. The eastern Asian and eastern and western North
American floristic disjunction: Congruent phylogenetic patterns in seven diverse genera.
Mol. Phylogenet. Evol. 10:178–190.
Xu S., He Z., Zhang Z., Guo Z., Guo W., Lyu H., Li J., Yang M., Du Z., Huang Y., Zhou R.,
Zhong C., Boufford D.E., Lerdau M., Wu C.-I., Duke N.C., Shi S. 2017. The origin,
diversification and adaptation of a major mangrove clade (Rhizophoreae) revealed by
whole-genome sequencing. Natl. Sci. Rev. 4:721–734.
Yang, Y., Moore, M. J., Brockington, S. F., Mikenas, J., Olivieri, J., Walker, J. F., & Smith, S.
A. 2017. Improved transcriptome sampling pinpoints 26 ancient and more recent
polyploidy events in Caryophyllales, including two allopolyploidy events. *New
Phytologist* 217(2): 855–870.
Yao J.L., Kang C., Gu C., Gleave A.P. 2022. The roles of floral organ genes in regulating
Rosaceae fruit development. Front. Plant Sci. 12:644424.
Yu, Y., & Nakhleh, L. 2015. A maximum pseudo-likelihood approach for phylogenetic
networks. *BMC Genomics*, 16: 1–10.
Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.
Zhang Q., Ree R.H., Salamin N., Xing Y., Silvestro D. 2021. Fossil-informed models reveal a
boreotropical origin and divergent evolutionary trajectories in the walnut family
(Juglandaceae). Syst. Biol. 71:242–258.
Zhao L., Jiang X.-W., Zuo Y., Liu X.-L., Chin S.-W., Haberle R., Potter D., Chang Z.-Y., Wen
J. 2016. Multiple events of allopolyploidy in the evolution of the racemose lineages in
Prunus (Rosaceae) based on integrated evidence from nuclear and plastid data. PLoS One.
11:e0157123.
1389
1390