2 Machine-Learning Identifies a Strategy for Differentiation Therapy in Solid Tumors

Authors: Saptarshi Sinha, Ph.D[1], Joshua Alcantara, B.S.[1, 2], Kevin Perry, B.S.[1, 2], Vanessa Castillo, B.S.[1], Celia R.

Espinoza, Ph.D. [1], Sahar Taheri, B.S.[3], Eleadah Vidales, B.S.[1, 2], Courtney Tindle, M.S.[1, 2], Adel Adel, B.S.[1], Siamak

Amirfakhri, D.V.M, Ph.D.[5], Joseph R. Sawires, B.S.[6], Jerry Yang, Ph.D. [6], Michael Bouvet, M.D.[5,7], Debashis Sahoo,

Ph.D. [3,4,7] and Pradipta Ghosh, M.D.[1,2,7,8‡]

Affiliations:

[1]Department of Cellular and Molecular Medicine, School of Medicine, University of California San Diego, La Jolla, CA

92093.

[2]HUMANOID™ Center of Research Excellence (CoRE), University of California San Diego, La Jolla, CA 92093.

[3]Department of Computer Science and Engineering, Jacob's School of Engineering, University of California San

Diego, CA 92093.

[4]Department of Paediatrics, School of Medicine, University of California San Diego, La Jolla, CA 92093.

[5]Department of Surgery, University of California San Diego, San Diego, CA 92093, USA; Department of Surgery, VA

San Diego Healthcare System, San Diego, CA, USA.

[6]Department of Chemistry and Biochemistry, University of California San Diego, San Diego, CA, 92093, USA.

[7]Moores Cancer Center, University of California San Diego, San Diego, CA, 92093, USA.

[8]Department of Medicine, University of California San Diego, San Diego, CA, 92093, USA.

25 **‡Corresponding author and Lead Contact**: Pradipta Ghosh, M.D.; Professor, Departments of Medicine and

26 Cellular and Molecular Medicine, University of California San Diego; 9500 Gilman Drive (MC 0651), George E. Palade

27 Building, Rm 232, 239; La Jolla, CA 92093. Phone: 858-822-7633. Email: prghosh@ucsd.edu.

Abstract

Background: Although differentiation therapy can 'cure' some hematologic malignancies, its curative potential remains unrealized in solid tumors. This is because conventional computational approaches succumb to the thunderous noise of inter-/intratumoral heterogeneity. Using colorectal cancers (CRCs) as an example, here we outline a machine learning(ML)-based approach to track, differentiate, and selectively target cancer stem cells (CSCs).

Methods: A transcriptomic network was built and validated using healthy colon and CRC tissues in diverse gene expression datasets (~5,000 human and >300 mouse samples). Therapeutic targets and perturbation strategies were prioritized using ML, with the goal of reinstating the expression of a transcriptional identifier of the differentiated colonocyte, CDX2, whose loss in poorly differentiated (CSC-enriched) CRCs doubles the risk of relapse/death. The top candidate target was then engaged with a clinical-grade drug and tested on 3 models: CRC lines *in vitro*, xenografts in mice, and in a prospective cohort of healthy (n = 3) and CRC (n = 23) patient-derived organoids (PDOs).

Results: The drug shifts the network predictably, induces *CDX2* and crypt differentiation, and shows cytotoxicity in all 3 models, with a high degree of selectivity towards all *CDX2*-negative cell lines, xenotransplants, and PDOs. The potential for effective pairing of therapeutic efficacy (IC50) and biomarker (*CDX2*-low state) is confirmed in PDOs using multivariate analyses. A 50-gene signature of therapeutic response is derived and tested on 9 independent cohorts (~1700 CRCs), revealing the impact of *CDX2*-reinstatement therapy could translate into a ~50% reduction in the risk of mortality/recurrence.

Conclusions: Findings not only validate the precision of the ML approach in targeting CSCs, and objectively assess its impact on clinical outcome, but also exemplify the use of ML in yielding clinical directive information for enhancing personalized medicine.

Introduction

Poor differentiation (increased stemness) is a hallmark of cancers; differentiation therapy that can target the cancer stem cells is expected to improve outcomes. A poster child of differentiation therapy is all-trans-retinoic acid (ATRA), which has achieved a 'cure' for the otherwise fatal acute promyelocytic leukemia (APML)[1-10]. Although differentiation therapy with a few other agents has been achieved in leukemia[11,12], it is yet to be realized in carcinomas. One of the reasons often cited for such failure is the thunderous intra- and inter-tumoral heterogeneity that limits our ability to track the identity the few cancer stem cells[13] (CSCs) that lie within the tumor. To look past such heterogeneity, computational algorithms have emerged as popular approaches to sort through 'big data' (such as transcriptomics), build networks, understand complex human diseases, and even prioritize targets. For example, mapping the network of robust relationships between genes has been a popular tool for understanding human diseases[14-19]. Pair-wise gene connections are usually established by performing correlation[20-25], mutual information[17] or other symmetric computational frameworks such as linear regression[26], dimension reduction[27], and clustering[28,29]. These methods often succumb to the heterogeneity in the real-world. More recently, a network transcriptomics-based approach, in which gene clusters relate to each other via directed edges of Boolean invariant relationships[30,31], has been successfully used to extract insights from heterogeneous big data and create maps of evolving cellular states along any disease continuum. This methodology has been used to identify translationally relevant cell states with high degrees of precision in diverse contexts and tissues[32-45]; most recently it has been used to identify a first-in-class therapeutic target to protect the vulnerable gut barrier in inflammatory bowel disease[30].

As for the therapeutic goal of the network, we set it as reinstatement of CDX2 which is a nuclear homeobox transcription factor and a tumor suppressor that belongs to the caudal-related family of CDX homeobox genes[46-49]. In 2016, an unbiased search (using the same Boolean logic approach[50,51]) looking for a biomarker of tumor stemness/differentiation status identified CDX2 as the statistically top ranked candidate biomarker of mature colon epithelial tissues, and by that token, of well-differentiated tumors [52]. The study predicted *CDX2* as the top developmentally regulated pro-differentiation candidate gene whose expression in colorectal cancer (CRCs) was inversely related to the "activated leukocyte cell adhesion molecule" (*ALCAM*/CD166); the latter is a marker of stem-like colon epithelial cells at the bottom of colon crypts[53,54] and on human colon cancer cells with high tumorigenic capacity in mouse xenograft models[55]. Validation studies using immunohistochemistry (IHC) showed that compared to patients with CDX2-pos tumors, those with a loss of CDX2 (accounting for approximately 9% of patients) had lower disease-free survival (DFS) and benefited from an adjuvant chemotherapy, even in stage II CRC.

Numerous groups have since reproduced the prognostic claims [56-69]. Thirty-two independent studies, involving >13,000 unique subjects, most that used IHC and one that used IHC in combination with quantitative mass spectrometry [70], have determined that CDX2-neg CRCs are associated with lower overall survival (OS) and DFS independent of ethnicity, DNA mismatch repair (MMR) status or stage [52,65,69,71] [**Figure 1**-*Step 1*]. Absence of CDX2 in tumors is associated with several adverse prognostic features, such as poor differentiation, advanced stage, vascular invasion, right-sided location, CpG island methylator phenotype (CIMP), and *BRAF* mutation [71-73]. Presence of CDX2 lowers the risk of death and recurrence by 50% and

52%, respectively[69]; this finding is more pronounced in stage II and III CRCs, with up to 70% reduction in the

risk of death. These findings drove consensus that pharmacologic reinstatement of CDX2 in CRCs is a

clinically important therapeutic goal [see **Figure 1**-*Step 1*], one that is also challenging because promising

candidates have not emerged.

Here we adopted the same Boolean-logic-based network transcriptomic approach[50,51] to screen for

therapeutic targets that can induce CDX2 and promote tumor differentiation.

Methods: Detailed computational and experimental methods are presented in *Supplementary Materials* and mentioned briefly here.

Bioinformatics analysis of gene-expression array databases. A complete catalogue of all the NCBI-GEO sample number identifiers (GSEIDs) of individual gene-expression array experiments used for model training and validation are provided (Supplemental Information 1).

Patient derived organoids. To enhance reproducibility and rigor, all CRC PDOs used in this work were acquired from the American Type Culture Collection (ATCC), which acquired these from diverse clinical sources [The Human Cancer Models Initiative (HCMI) US National Cancer Institute (NCI), Cancer Research UK (CRUK), the Wellcome Sanger Institute (WSI), and the foundation Hubrecht Organoid Technology (HUB)]. Relevant demographic, clinical and genetic information is provided (Supplemental Information 3).

Statistical tests. Optimal cut-off values of gene expression levels are derived using a threshold defined by *StepMiner* algorithm in the individual dataset[74]. Gene signature is used to classify sample categories and the performance of the multi-class classification is measured by ROC-AUC (receiver operating characteristics area under the curve) values. Standard t-tests were performed using python scipy.stats.ttest_ind package (version 0.19.0) with Welch's two sample t-test (unpaired, unequal variance (equal_var = False), and unequal sample size) parameters. Multiple hypothesis correction were performed by adjusting p-values with statsmodels.stats.multitest.multipletests (fdr_bh: Benjamini/Hochberg principles). Differential expression analysis was performed in DESeq2 in R (version 1.16.1). Kaplan–Meier analysis and determination of statistical significance by Log Rank test was performed using GraphPad Prism 9.1. Violin and Bubble plots are created using python seaborn package version 0.12. BooleanNet algorithm is used to statistically determine Boolean Implication relationships between genes (statistic > 3 and error-rate < 0.1). The source code for Boolean analysis framework is https://github.com/sinha7290/Prodiff[32]. The best estimate of effect size (ratio between the difference in mean and standard deviation) is 2.5 in PDO and 1.0 in animal experiment for CDX2 induction using PRKAB1-agonist. The sample size in each group of at least 4 PDOs and 17 animals should suffice (α = 0.05 and power = 0.80).

Study Design. The study was designed by the first and last author. All authors contributed to the gathering, analysis, and interpretation of the data. The first draft of the manuscript was written by the first and last author and was subsequently edited based on comments from all authors.

Ethics statement. This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. All animal studies were approved by the San Diego Veterans Administration Medical Center Institutional Animal Care and Use Committee (protocol A17-020; PI Bouvet). All colorectal cancer PDOs were obtained from the American Type Culture Collection (ATCC). The latter is a nonprofit, global biological resource centre and a distributor of organoid models curated as part of Human Cancer Models Initiative (HCMI) for enhancing reproducibility. Deidentified case-associated data, including that of molecular characterization of the models, the tissues from which they were derived were used here, whereas available harmonized data are accessible through NCI's Genomic Data Commons. All healthy colon PDOs were obtained from individuals presenting for routine colonoscopy as part of colon cancer screening at UC San Diego, with their informed consent to participate in a study protocol approved by an Institutional Review Board (IRB; protocol #190105; PI Ghosh).

Role of funders

Funders of the study had no role in study design, data collection, data analyses, interpretation, or writing of report.

Results

A network-rationalized approach

We used a machine learning-based computational platform, Boolean Network Explorer (*BoNE*)[30], and *CDX2* as a 'seed' gene to create a model of progressive gene regulatory events that occur during differentiation in the colon (**Figure 1**-*Step 2;* **Figure S1A-B**). The model was trained on a pooled heterogeneous dataset that included maximum possible diversity (1662 CRCs, 68 adenomas and 170 normal tissues) (**Figure S1C-D**). Statistical learning on the *CDX2*-proximal immediate gene clusters (clusters #2 and #3, **Figure S1D**) helped statistically rank 5 genes as top targets (**Figure 1**-*Step 2;* **Figure S1E**). By virtue of how these genes relate to *CDX2* within the network, upregulation of any of these 5 genes (presumably via target/pathway agonists) is predicted to also upregulate *CDX2*. Validation studies on an additional 2,937 human (**Figure S1F**) and 318 mouse (**Figure S2**) colon samples showed that these 5 genes remain relevant across species and diverse cohorts.

In search of potent and specific drugs to modulate the 5 targets, we searched the PDB [Protein Data Bank; https://www.rcsb.org] for structure resolved selective agonistic modulators and found 25 molecules/structures for PRKAB1, but none for the remaining targets. To determine the most clinically advanced candidate, the PDB-derived hits were cross-searched through the Clinical Trials.gov portal, which led to the identification of a clinical-grade drug, PF-06409577 (henceforth, PF). Although lacking clinical indication at present, the drug was deemed safe in a randomized, double-blind, placebo-controlled Phase I trial [*ClinicalTrials.gov* ID: NCT02286882]. Furthermore, PF was recently shown to demonstrate target specificity and the ability to upregulate the *PRKAB1* transcript and the translated protein in the colonic epithelium ex-vivo and mouse colon in vivo[30].

The network model predicted that the agonism of the protein that is translated by the *PRKAB1* gene, i.e., Protein Kinase AMP-activated Non-Catalytic Subunit Beta-1 will-- (i) upregulate *PRKAB1* expression and, by virtue of their "*equivalent*" relationship, also upregulate the other genes that co-cluster with it (**Figure S1D**); (ii) upregulate *CDX2* and the *CDX2*-proximal gene clusters (#2 and #3) by virtue of "hi=>hi" relationship (**Figure S1D**); and (iii) downregulate stemness genes by virtue of "hi=>lo" relationship (*CCDC88A* and others in C#9; **Figure S1D**). Notably, *PRKAB1* has a universally conserved relationship with *CDX2*, across diverse human and mouse CRC datasets, including early onset CRCs (**Figure S2A-D**). That PRKAB1 (but not the other β-subunit, PRKAB2) is strongly expressed in the GI tract[30] and that PRKAB1-agonists trigger cell polarity in the gut epithelium[75-78], served as additional justifications for its rationalized selection as target.

For target validation, we prioritized three models, based on their *CDX2*-low (target tumor type) and high (expected to be negative control) status. Alongside the traditional readouts of anti-cancer effects, a set of additional end points were *a priori* formulated as metrics of success; therapeutic goal was set such that all three models must pass all three end points (see **Figure 1**-*Step 3*). Prior attempts that successfully induced *CDX2* temporarily[79], but ultimately failed to emerge as differentiation therapy, had mixed impact on these metrics of success (**Figure S3**). Finally, to track successful *CDX2*-reinstatement and assess its impact on survival, a 50-gene signature of therapeutic response is derived and tested on numerous independent cohorts (see **Figure 1**-*Step 4*).

Target validation on CRC cell lines and xenograft models

Next, we prioritized two poorly differentiated CRC cell lines (HCT116 and Sw480 cells; both exhibited the

lowest levels of *CDX2* transcripts among a panel of 26 sequenced CRC cell lines that were profiled by

microarrays; GSE10843) as the optimal models for hypothesis testing, and one of the two well-differentiated

cells with high *CDX2* levels (DLD1) as the negative control (**Figure 2A**). PF was capable of inducing cell

death within 48 h in *CDX2*-low HCT116 and Sw480 cells but not the *CDX2*-high DLD1 cells (IC50's in 6-7

µM range vs > 40 µM, respectively; **Figure 2B**; **Figure S4A**). In both *CDX2*-low cell lines, PF-induced cell

death was preceded by induction of differentiation and suppression of stemness markers, at both transcript

(**Figure 2C**) and protein (**Figure S4B-E**) levels in a dose-dependent manner. These effects were associated

also with dose-dependent increase in apoptosis (**Figure S5A-H**). PF failed to reinstate *CDX2,* change other

markers of stemness and differentiation (**Figure 2D**) or induce cell death (**Figure 2E**) in *PRKAB1*-depleted

cells, thereby demonstrating that its anti-cancer action is target specific. Finally, Metformin which is a global,

non-specific and indirect agonist of all forms of the trimeric AMP-kinase (regardless of whether PRKAB1 or

PRKAB2 is the β-subunit within the αβγ trimer) failed to trigger cell death or induce *CDX2* expression even

at very high doses (**Figure 2B**; **Figure S5I-K**), suggesting that PRKAB1-specific effect is required for

therapeutic potency.

When implanted subcutaneously as xenografts in nude mice (**Figure 2F**), HCT116 tumors revealed

a 68% reduction in tumor volume (total AUC for vehicle and PF time series were 51280.23 and 26676.77,

respectively; **Figure 2G-I**) and an ~62.48% increase in survival (**Figure 2J**; Mantel-Haenszel Hazard Ratio

of 0.09 for reduction in death rates). PF promoted glandular structures within tumor (see legend **Figure S6A**),

reinstated *CDX2*, induced differentiation markers and suppressed stemness markers, as determined by

qPCR (**Figure 2K**). Reinstatement of CDX2 protein and its nuclear localization, alongside upregulation of

PRKAB1 protein were confirmed also by immunohistochemistry (IHC; **Figure 2L-M; Figure S6B**). Consistent

with the notion that colon stem cells divide very slowly [80], PF triggered a ~3-fold increase in mitotic bodies

(**Figure S6C**; **Figure 2M**). Conventional markers of tumor differentiation were also observed, such as a

reduction of nuclear β-Catenin (**Figure S6D**; **Figure 2M**) with concomitant increase in junctional β-Catenin,

where it co-localized with E-Cadherin (**Figure 2N**; **Figure S6E**).

205   *Target validation in patient-derived organoids*

We next assessed the impact of PF on 23 CRC PDOs (recruited as two prospective independent cohorts)

and 3 healthy colon PDOs (see **Table** and *Methods*; **Figure 3A**). The anti-cancer action (as determined by

low IC50 values) was limited only to *CDX2*-low CRC PDOs, whereas the drug was virtually inert (with IC50>40

µM) against *CDX2*-high PDOs as well as healthy colon PDOs (**Figure 3B**). Both *CDX2* ('seed' gene in the

network) and *PRKAB1* (network-derived therapeutic target) showed significant positive correlations with IC50

(**Figure 3B**). By contrast, *CCDC88A* (a network-derived stemness marker; see **Figure S1D**), showed a

significant inverse correlation with IC50 (**Figure 3B**). Hierarchical unsupervised clustering performed using

just 3 network-derived genes (*CDX2*, *PRKAB1* and *CCDC88A*) perfectly classified sensitive from resistant

PDOs (**Figure 3C**). PF treatment enhanced differentiation and suppressed stemness (**Figure 3D**), as

observed in CRC cell lines (**Figure 2C**) and in xenografts (**Figure 2K**) previously, but these impacts were

restricted only to *CDX2*-low CRC PDOs. A multivariate analysis performed on the cohort of CRC PDOs (see

Supplemental Information 3 for all variables included) showed that *CDX2*-status was the strongest variable

that determined IC50 (**Figure 3E**). In addition, two additional co-variates emerged: (i) histological grade and *KRAS*[mut] status, which indicates that PF was less likely to work on moderate/well-differentiated tumors that tend to frequently harbor oncogenic *KRAS* mutations (**Figure 3E**; **Table**).

The relationship between CDX2-induction therapy and KRAS-status suggested that CDX2 may not be prognostic among *KRAS*[mut] group because its expression remains high (i.e., not lost). A univariate analysis on a large cohort of CRCs with known driver mutations and patient outcome confirmed that *CDX2* was indeed a determinant of outcome in tumors with *KRAS*[WT] (**Figure 3F**-*left*), but not in those with *KRAS*[mut] (**Figure 3F**-*right*). These findings suggest that although relatively small, our PDO cohort is sufficiently diverse and representative of the heterogeneity in the clinic to have revealed the interplay between *CDX2* and *KRAS*[mut].

Indicators of therapeutic efficacy and target cell specificity

RNA seq on all 3 models— CRC cell lines, xenografts, and PDOs—confirmed that PRKAB1 agonism met 3 *a priori* formulated metrics of success: (i) *CDX2* was reinstated, which was the primary therapeutic goal (bubble plots in **Figure 4A**-*top row*; **4B**; violin plots in **Figure S7A-C**)*,* (ii) markers of crypt top enterocytes and Goblet cells, which are established surrogate markers of differentiation in the colon crypt [51,81] were induced (**Figure 4A**-*middle row*; **Figure S7D-F**), and (iii) the network was perturbed predictably, i.e., genes in clusters #2 and #3 were induced (**Figure 4A**-*bottom row*; **Figure S7G-I**). Of note, all three metrics were fulfilled selectively in *CDX2*-low CRC cell lines, xenografts and PDOs, but not in *CDX2*-high PDOs or in healthy PDOs (**Figure 4A**-*columns #5 and #6*; **Figure S7C, S7F, S7I**).

A recent effort, using an innovative machine-learning algorithm extracted transcriptomic and epigenetic feature sets derived from healthy (normal) stem cells and their differentiated progeny; the work released novel gene signatures of oncogenic dedifferentiation[82]. Those signatures revealed that *CDX2*-reinstatement therapy had no effect on normal stem cell-related genes in healthy and CRC PDOs (**Figure 4C**) but specifically reduced the expression of genes that impart cancer-related stemness (**Figure 4D**), those that enable somatic cell reprogramming (i.e., the Yamanaka factors which have been implicated in the origin of CSCs[83] (**Figure 4E**) and those that define CRC-specific CSCs (**Figure 4F**).

Taken together these findings show that *CDX2*-reinstatement therapy selectively downregulates CSC-programs that are unique to *CDX2*-low tumors, thereby sparing normal stem cells in the healthy colon tissue as well as *CDX2*-high tumor cells.

Potential for synergy with other therapeutic modalities

Lack of CDX2 expression in Stage II-III CRCs has been previously shown to be associated with benefits from adjuvant chemotherapy [52]; CDX2-positive tumors lack such benefit. We asked how therapeutic reinstatement of CDX2 in *CDX2*-low tumors may impact response to chemotherapy. PF treatment consistently altered a panel of patient-derived gene signatures across all models tested; it induced signatures of favourable response and suppressed signatures of non-response to diverse chemotherapeutic agents (e.g., 5-FU, Oxaliplatin, FOLFIRI) (**Figure 4G-i**)*.* Consistent with these gene expression changes, PF showed synergy with a panel of chemotherapeutic agents (see legend, **Figure 4H**).

Similar observations were made when we extended the analysis to include other treatment modalities;
PF consistently induced gene signatures of response to anti-EGFR (Cetuximab; **Figure 4G-ii**) and other
tyrosine kinase inhibitors (TKIs; **Figure 4G-iii**), with a few notable exceptions (Afatinib and Gefitinib; **Figure**
4G-iii). PF also induced gene signatures of response and suppressed signatures of non-response to
immunotherapy (**Figure 4G-iv**). The latter is consistent with the observation that PF also robustly induced a
consensus molecular subtype (cms)-1 in all three model systems tested (**Figure 4G-v**); the immune
landscape in cms1 subtype is known to favour response to immune checkpoint inhibitors[84].
263
A molecular signature of therapeutic response and selectivity
To define the nature of the therapeutic response, we leveraged RNA seq data from all 3 models to conduct
an integrated differential expression analysis (see *Methods*). Briefly, differentially expressed genes (DEGs)
from two independent models (cell lines and xenografts; **Supplemental Information 4-5**) were further refined
on a third model, the CRC PDOs (**Figure 5A**). A set of 50 DEGs emerged; these genes were expressed in
CDX2-low CRC PDOs at levels significantly higher than healthy or *CDX2*-high CRC PDOs (**Figure 5B**;
Figure S8A-B). We used this 50-gene signature as a computational tool to track the changes along the
stemness→differentiation axis that occurs due to reinstatement of *CDX2* and found that PF suppressed these
genes (**Figure 5B**; see **Supplemental Information 6**) to levels seen in healthy PDOs and *CDX2*-high CRC
PDOs (**Figure S8A-B**).
This 50-gene signature was notable for the presence of several unique CSC-associated genes, such
as, *AGR2* [85], *ASCL2* [86,87], *ALDH3A1* [88], and *TM4SF1* [89,90], that were not parts of CSC signatures identified by
machine-learning (**Figure 4C-F**). The 50 genes were enriched for pathways that are known to augment CSC
survival [91-94], e.g., PTK6-dependent JAK/STAT signaling, numerous growth factor receptor and non-receptor
TK and the Hippo/YAP pathways and loss of TGFβ signaling (**Figure 5C**). The selective upregulation of these
pro-survival pathways in *CDX2*-low CRCs (but not healthy PDOs or *CDX2*-high CRCs) suggests that their
concomitant and catastrophic withdrawal in the setting of *CDX2* reinstatement could be the reason for the
observed target-cell selectivity.
282
Impact of CDX2 reinstatement therapy on survival
To assess the impact of differentiation therapy on disease outcome, we analyzed the ability of the 50-gene
signature of therapeutic response to stratify relapse/disease-free survival (RFS/DFS) in diverse cohorts of
CRC patients. Compared to patients whose tumor had a high 50-gene score (like the untreated *CDX2*-low
PDOs), those with a low 50-gene score consistently had a better outcome across all datasets analyzed
(**Figure 5D, G-I**). The odds ratio (O.R) was ~2.0 (**Supplemental Information 7-*tab #1***), implying that high
expression of the 50-gene signature carries a double risk of death/recurrence and that therapeutic
reinstatement of *CDX2* will reduce that risk by 50%. A univariate analysis revealed that the 50-gene signature
and *CDX2* were significant deterministic covariates of RFS (**Figure 5E-*left***) but emerged as the better of the
two in *KRAS*[WT] (**Figure 5E-*middle***). The 50-gene score also outperformed *CDX2* as a prognostic indicator,
consistently across all cohorts tested (**Figure S8C-F**; **Supplemental Information 7**). These findings show
that the positive impact of CDX2 reinstatement is better represented in the 50-gene score (compared to
CDX2 alone). Consistent with our observations earlier in the case of *CDX2* (**Figure 3E**), the 50-gene

signature was not a factor in *KRAS*^{mut} tumors (**Figure 5E**-*right*). These results provide objective evidence that

therapeutic suppression of the 50-gene signature is a desirable therapeutic goal that could translate to

improved survival.

Discussion

The major discovery we report here is a network-rationalized strategy to push differentiation of CRC stem cells by reinstating *CDX2* expression and define the cellular, molecular, and translational impact of such reinstatement. Much like immunotherapy acts by reinvigorating a physiologic response, we show that *CDX2*-reistatement acts by reinvigorating yet another physiologic program, that of differentiation. We identify and validate one target (PRKAB1) and confirm the utility of a first-in-class agent with proven safety (PF-06409577) to exert a target- and cell-type specific pro-differentiation and anti-tumor effect. *CDX2*-reinstatement therapy also fulfilled a set of rigorous and objective metrics of success that were formulated *a priori,* and its efficacy can be objectively tracked using a set of companion biomarkers (*CDX2* expression and a 50-gene signature of therapeutic response). Mechanistically, we show that *CDX2*-reinstatement is associated with the suppression of key stemness genes and numerous pro-survival pathways to which *CDX2*-low CRC stem cells (but not normal stem cells) may be uniquely addicted. Finally, we provide evidence that *CDX2*-reinstatement therapy is expected to synergize with most existing therapeutic modalities and translate to a reduction in death/relapse by ~50%.

Translational potential and key considerations: We advocate for the findings to be translated into clinical trials, prioritized as follows. The study pinpointed *CDX2*-low CRCs, and more specifically, those with a high 50-gene signature, as the optimal target group that will maximally benefit from this form of therapy. That *CDX2*-high tumors were found to be more differentiated and more likely to harbor mutant *KRAS* (but not *BRAF*) is consistent with the fact that $KRAS^{mut}$ promotes hyperplasia and mucinous differentiation[95], but not stemness[96] and that loss of CDX2 (as determined by IHC) is positively associated with $BRAF^{mut97}$ but inversely associated with $KRAS^{mut97}$. This intriguing distinction could be attributed to cell-type specific roles of $KRAS/BRAF^{mut}$ reported recently[98]. $KRAS^{mut}$ status, however, does not appear to have any bearing on the determination of therapeutic candidacy because the two cell lines that were prioritized here based on low *CDX2* expression levels are also $KRAS^{mut}$, [HCT116 (1 mutant allele; G13D) and Sw480 (both alleles mutated; G12V)], and are sensitive to PF. These findings are consistent with a population-based study on metastatic CRCs (mCRCs; n = 796) that showed loss of CDX2 (as determined by IHC) defines a small subgroup of $KRAS^{mut}$ cases (~9%) with a worse prognosis[97], and hence, could benefit from differentiation therapy. We conclude that $KRAS^{mut}$ status is a confounding co-variate that is associated with the differentiation status of a tumor; however, it should not override low-CDX2 and/or high-50-gene signature in the determination of who may benefit from differentiation therapy.

Because loss of *CDX2* is an adverse prognostic factor in stage IV and stage III/$BRAF^{mut}$ CRCs[71,99], these subgroups can be prioritized for *CDX2*-reinstatement therapy with PRKAB1 agonist as single agent or together with other modalities.

Although *CDX2* is prognostic in stage II(MSS)[52,65,69] and stage III CRCs[52,100], the use of *CDX2*-reinstatement therapy in these subgroups deserves additional considerations, and even caution. In stage III disease, adjuvant chemotherapy has been a standard of care since the 90's based on the Moertel trial[101]; however, many in this subgroup relapse despite chemotherapy (variable ~15-75% based on T-N substage[102]) and die of disease[103]. Because CDX2 expression is associated with increased expression of the multi-drug resistance gene, MDR1, which is implicated in the chemoresistance of *CDX2*-pos tumors[104,105], the use of

PRKAB1 agonist in combination with conventional chemotherapeutic agents must wait despite demonstrable

synergy in our studies. Considering these factors, PRKAB1 agonist as a single agent or with other modalities

can instead be prioritized for those with *CDX2*-low tumors who relapse and/or progress to metastatic disease.

This is feasible because CDX2 expression status in CRCs is not affected by metastasis or chemotherapy[106].

If those studies show benefit, PRKAB1 agonist as single agent could be considered as a first-line option for

patients with *CDX2*-low tumors. Because stage II/III early-onset CRC is a subgroup that show worse outcome

(poorer DFS) than its late-onset counterparts[107], and the *CDX2-PRKAB1* relationship is strongly conserved

in this subgroup (**Figure S2A**), we advocate the inclusion of this group in future studies considering PRKAB1

agonists in stage II/III CRCs.

Finally, because the degree of differentiation dictates the susceptibility to *BRAF*-mut-driven

tumorigenesis[108,109], early-stage patients with histopathologic criteria not qualifying for chemotherapy might

be reconsidered for *CDX2*-reinstatement therapy if their tumor has focal or global loss of CDX2 and

mutated *BRAF*.

Expected impact on pre-neoplastic lesions of the colon:

There is sufficient literature on how *CDX2*-loss (and hence reduced differentiation) is permissive to CRC

initiation, especially along the serrated pathway[108-110] and in the setting of inflammatory bowel disease-

associated CRCs[111-113]. A recent study using mosaic *CDX2*-/- mouse models showed that *CDX2* serves as a

tumor suppressor gene in a non–cell-autonomous manner; *CDX2*-/- metaplastic pre-cancerous cells could

induce tumorigenesis from neighbouring non-metaplastic cells without themselves becoming cancerous.

Focused studies on PDOs derived from such high-risk population are expected to reveal if PRKAB1 agonists

such as PF can serve as prophylactic therapy for those at highest risk of CRCs.

Study limitations:

How PRKAB1-agonism translates to *CDX2*-reinstatement and/or crypt differentiation were not studied. Future

efforts will assess whether aberrant methylation at the CDX2 promoter reported previously[114,115] is reversed

or if the loss of the highly specialized, junction-localized, tumor-suppressive stress-polarity signaling

pathway[76,77] is reinstated. Given the relatively small size of PDO cohorts, these results will need to be further

validated. We advocate for these findings to be confirmed within the framework of randomized clinical trials,

in conjunction with RNA sequencing studies.

Conclusions

Despite being at the forefront of biomedical research, therapies that can reprogram CSCs in carcinomas had not emerged. We have addressed this unmet need using an network-based approach that differs from the current practice in three fundamental ways: 1) Target identification and prediction modeling that is guided by a Boolean implication network; 2) Target validation in network-rationalized animal models that most accurately recapitulate the CSC-specific *CDX2*-low state; 3) Target validation in human preclinical organoid models that have the potential to test target-biomarker pairing and thereby, personalize the choice of therapies. The combined synergy of these approaches validates a first-in-class agent in tracking and killing CSCs, while sparing normal stem cells. As for the impact of these findings, it is noteworthy that despite level 1 evidence showing the prognostic role of CDX2, its independent validation through numerous studies[52,65,66,68-70,79,116,117], and a compelling pharmacoeconomic justification (i.e., cost $ 50 000 per QALY) for a CDX2-stratified decision to offer chemotherapy[118], prospective trials demonstrating the superiority of a CDX2-stratified decision has not emerged. Consequently, current guidelines do not recommend CDX2 be used in prognostication in CRCs[119-123]. Our finding that CDX2 is a companion biomarker for differentiation therapy with PF [a drug that has been deemed as clinically safe in Phase 1 on healthy volunteers (NCT02286882) and can be immediately repurposed], provides a compelling impetus for setting up prospective trials that can rigorously test both claims.

Data availability statement

All transcriptomic datasets generated in this work has been deposited at NCBI GEO [GSE237623, GSE237624, GSE237625]. The data underlying all the figures and tables are available in the article and its online supplemental materials.

Acknowledgements

We are grateful to the staffs of the UC San Diego HUMANOID™ Center of Research Excellence, for support in the logistical support in conducting the studies on human PDOs. This work was supported by the National Institutes for Health (NIH) grants UG3TR002968 and UH3 TR002968 (to P.G), R01-GM138385 (to DS), and R01-CA238042, R01-CA100768 and R01-CA160911 (to P.G) and VA Merit awards 1 I01 BX003856-01A1 and 1 I01 BX004494-01 (M.B). S.S was supported in part by a American Association of Immunologists (AAI) Intersect Fellowship Program for Computational Scientists and Immunologists. Other sources of support include: R01-AI155696 (P.G and D.S), R01-AI141630 (P.G), and UG3TR003355 (D.S. and P.G), a Torey Coast Foundation Award (P.G, D.S, M.B, J.Y), and by three Padres Pedal the Cause awards #PTC2017, #PTC2021 and #PTC2022. D.S and P.G were also supported by the Leona M. and Harry B. Helmsley Charitable Trust. Funders had no role in study design or conclusions.

Author Contributions

SS, DS and PG conceptualized the study; SS and PG designed and analysed all parts of the study; JA, KP, EV and CT acquired, expanded, biobanked and conducted all experiments with patient-derived organoids; VC conducted immunocytochemistry studies; CRE conducted qPCR studies and assisted SS, VC and PG on key parts of the study; ST and DS built and validated the network model; SS and PG conducted all other computational and bioinformatic analyses; SA and MB conducted the xenotransplantation studies; JRS and JY synthesized the pharmacophore; MB, JY, DS and PG secured funding; SS and PG prepared all display items; PG wrote the first draft of the manuscript; all authors edited and approved the final version.

Conflicts of Interest Statement: The authors have declared that no conflict of interest exists.

Figures and Legends:

Figure 1. Study Design: Identification and validation of a therapeutic strategy to reinstate CDX2 using ML.

Step 1: Flow charts presents the emerging evidence, in chronological order, which rationalized reinstatement of CDX2 as a desirable end goal for differentiating cancer stem cells in the most the aggressive subtypes of CRCs.

Step 2. The key steps towards the network-based identification of PRKAB1 as a potent and actionable target, whose agonism is predicted to reinstate CDX2 expression in those CRCs where its expression is lost. **Supplemental Information 1** for a catalogue of all datasets used for model training and validation and **Supplemental Information 2** for gene clusters, beyond *CDX2*, that are predicted to be co-induced upon PRKAB1 agonism.

Step 3. Three model systems and objective metrics of success that are used to validate PRKAB1 agonism as a strategy to reinstate CDX2 expression and induce differentiation.

Step 4. A 50-gene signature of therapeutic response derived from all 3 models is used to objectively measure the impact of CDX2 reinstatement therapy and identify patients who are most likely to benefit from the same.

Figure 2. Target validation in CRC cell lines and in xenotransplantation studies in mice. A. Violin plot
displays *CDX2* expression (transcripts/million; tpm) in 4 common CRC cell lines. p-values are calculated

using Welch's t-test. **B**. Table shows IC50 values for specific (PF) or non-specific (Metformin) PRKAB1-
agonists on CRC cell lines, as determined by MTT assays at 48 h. **C**. RT-qPCR on viable cells (>95%, at 16
h) of differentiation and stemness related markers in the colon is displayed as a heatmap of Z-score
normalized values. **D-E**. Immunoblots (D) and MTT based survival assays (E) show the impact of PRKAB1
depletion (in HCT116 cells by shRNA) on the anti-cancer effect of PF. p-values are calculated using one-way
ANOVA. Error bars indicate S.E.M of 3 independent biological experiments. See **Figure S4** and **S5** for
additional studies on HCT116 and Sw480 cells challenged with increasing doses of PF (**S4** and **S5A-H**) or
Metformin (**S5I-K**). **F**. Schematic showing experimental design and workflow. **G**. Graphs display growth
kinetics of vehicle (Veh; n = 13) and PF-treated (n = 15) tumors, measured using callipers (2/wk), collated
from 3 independent experimental cohorts. p-values are calculated using multiple paired t-test following
Benjamini, Krieger, and Yekutieli method. **H**. Images of tumor xenografts at 8-wk post-implantation. **I**. Bar
plots compare the area under the curve (AUC) of treatment and control arms [95% Confidence Interval was
31788.29 to 70772.18 (Veh) and 16949.29 to 36404.26 (PF); Error bars represent S.E.M., and p-value was
determined by unpaired 2-tailed t-test using GraphPad]. **J**. Kaplan-Meier analysis shows time to the fulfilment
of IACUC-mandated criteria for euthanasia (death and/or signs of discomfort). p-value was determined by
Log Rank test. H.R., hazard ratio. **K**. RT-qPCR on xenografts displayed as a heatmap of Z-score normalized
values. **L**. Representative fields from CDX2-stained FFPE-xenografts are shown. Scale bar = 100 μm. **M**.
Quantification of CDX2, PRKAB1, and nuclear β-catenin staining by ImageJ (IHC profiler application) and
mitotic bodies (manually counted in H&E-stained sections). p-values are calculated by unpaired 2-tailed t-
test. See **Figure S6B-D** for images from representative fields. **N**. Representative fields from FFPE tumors
co-stained for E-cadherin (red), β-catenin (green) and DAPI (nuclei; blue) are shown. Scale bar = 20 μm. See
also **Figure S6K** for colocalization analysis by ImageJ. **O**. Immunoblots of lysates of viable tumor tissues.
See **Figure S6F** for quantification.

Figure 3. Differentiation and anti-cancer effects of PRKAB1 agonism are restricted to *CDX2*-low PDOs. A. Schematic of workflow for sequential analyses in two independent patient cohorts. See **Supplemental Information 3** for patient demographics, tumor characteristics and mutation status of key genes. **B**. Correlation plots between IC50 and the levels of expression of 3 genes, as determined by qPCR. **C**. Heatmap displays the results of unsupervised clustering of the CRC PDOs using Z-score normalized qPCR estimation of 3 genes. **D**. Heatmap generated as above of the vehicle (Veh) and PF-treated CRC PDOs, using a panel of stemness and differentiation markers. **E**. Multivariate analysis of IC50 with PF as a linear combination of all variables in the PDOs used in this study. The coefficient of each variable (at the centre) with their upper and lower bounds of 95% confidence interval (as error bars) and the p-values from t-tests are illustrated in the bar plot. The p-value for each term tests the null hypothesis that the coefficient is equal to zero (no effect). Red = significant co-variates. *p ≤ 0.05; **p ≤ 0.01. See also **Supplemental Information 3** for source data. **F**. Univariate analysis model of relapse-free survival (RFS) in KRAS-WT vs KRAS-mutant tumors in GSE39582. *p ≤ 0.05; **p ≤ 0.01; ***p ≤ 0.001.

Figure 4. Indicators of success, target-cell specificity, and the potential for synergy with other treatment modalities. A. Expression of *CDX2* (top row), crypt top and Goblet cell markers (middle row) and genes in clusters #2-3 in the network (bottom row) gene/gene signatures in HCT116, Sw480, xenografts and PDOs is visualized as bubble plots of ROC-AUC values (radius of circles are based on the ROC-AUC) for

the classification of vehicle (Veh) vs PF-treated samples. Colors indicate the direction of gene regulation (Up,

red; Down, blue) in the treated samples. The significance of AUC is determined based on Welch's t-test and

displayed using standard codes (''p>0.1, '.'p<=0.1, *p<=0.05, **p<=0.01, ***p<=0.001) next to the ROC-AUC

values. See **Figure S7** for the display of the same data as violin plots with actual p-values. **B**. Violin plots

show *CDX2* expression in PDOs. p-values are based on Welch's t-test compared to vehicle (Veh)-treated

CDX2-low PDOs. **C-F.** Violin plots display the composite score of stemness markers recently reported in a

pan-cancer effort that used machine learning. p-values were determined by Welch's t-test, comparing each

sample against vehicle (Veh)-treated *CDX2*-low PDOs. **G**. Expression of gene signatures of response

(default or indicated with 'R') or non-response (N.R.) to multiple therapeutic modalities in CRC cell lines,

xenografts and PDO is visualized as bubble plots of ROC-AUC values (radius of circles are based on the

ROC-AUC) for the classification of vehicle (Veh) vs PF-treated samples. Colors indicate the direction of gene

regulation (Up, red; Down, blue). p-values displayed are based on significance of AUC using Welch's t-test

and using standard code (*p<=0.05, **p<=0.01, ***p<=0.001) next to the ROC-AUC. **H**. IC50 values for

combinations of 5 µM PF (constant) with increasing conc. of the indicated chemotherapeutic agents, as

determined on HCT116 and sw480 cell lines by MTT assays were used to derive a combination index (CI)

using a previously established formula [124]. Results are displayed as a heatmap.

Figure 5. A gene signature for tracking therapeutic response and estimating impact on survival.

A. Computational workflow used in the derivation of a 50-gene signature of therapeutic response using an integrated differential expression analysis (DEA) approach. Differentially expressed genes (DEGs) from CRC cell lines and xenografts (see Supplemental Information 4-6) were further refined based on their ability to accurately classify PDO samples with/without PF treatment (see Supplemental Information 6 for gene list rank-ordered by ROC-AUC score). **B**. Heatmap displays the results of unsupervised clustering of PDOs

based on the 50-gene signature. Veh, vehicle. See **Figure S8A-B** for visualization of findings as violin plots.

C. Reactome pathway analysis of the 50-gene signature. **D**, **F-H**. Kaplan-Meier (KM) plots show recurrence-

free (RFS) or disease-free (DFS) survival over time on various cohorts of CRC patients, stratified by high vs

low expression scores of the 50-gene signature. p-values were determined by Log Rank test. See **Figure**

S8C for KM plots on the same cohorts based on *CDX2*-based stratification. **E**. Univariate analysis models of

RFS as a linear combination of all variables in a large CRC cohort, including the 50-gene signature. The

coefficient of each variable (at the centre) with 95% CI (as error bars) and the p values are illustrated. The p-

value for each term tests the null hypothesis that the coefficient is equal to zero (no effect). Left, entire cohort;

middle, *KRAS*-WT CRCs; right, *KRAS*-mut CRCs. *$p \leq 0.05$; **$p \leq 0.01$.

References

1. Stone RM, Maguire M, Goldberg MA, Antin JH, Rosenthal DS, Mayer RJ. Complete remission in acute promyelocytic leukemia despite persistence of abnormal bone marrow promyelocytes during induction therapy: experience in 34 patients. Blood 1988;71(3):690-6.
2. Kantarjian HM, Keating MJ, Walters RS, et al. Acute promyelocytic leukemia. M.D. Anderson Hospital experience. Am J Med 1986;80(5):789-97.
3. Cunningham I, Gee TS, Reich LM, Kempin SJ, Naval AN, Clarkson BD. Acute promyelocytic leukemia: treatment results during a decade at Memorial Hospital. Blood 1989;73(5):1116-22.
4. Tallman MS, Andersen JW, Schiffer CA, et al. All-trans-retinoic acid in acute promyelocytic leukemia. N Engl J Med 1997;337(15):1021-8. DOI: 10.1056/NEJM199710093371501.
5. Tallman MS, Andersen JW, Schiffer CA, et al. All-trans retinoic acid in acute promyelocytic leukemia: long-term outcome and prognostic factor analysis from the North American Intergroup protocol. Blood 2002;100(13):4298-302. DOI: 10.1182/blood-2002-02-0632.
6. De Botton S, Dombret H, Sanz M, et al. Incidence, clinical features, and outcome of all trans-retinoic acid syndrome in 413 cases of newly diagnosed acute promyelocytic leukemia. The European APL Group. Blood 1998;92(8):2712-8.
7. Testi AM, Biondi A, Lo Coco F, et al. GIMEMA-AIEOPAIDA protocol for the treatment of newly diagnosed acute promyelocytic leukemia (APL) in children. Blood 2005;106(2):447-53. DOI: 10.1182/blood-2004-05-1971.
8. Fenaux P, Chevret S, Guerci A, et al. Long-term follow-up confirms the benefit of all-trans retinoic acid in acute promyelocytic leukemia. European APL group. Leukemia 2000;14(8):1371-7.
9. Sanz MA, Martin G, Rayon C, et al. A modified AIDA protocol with anthracycline-based consolidation results in high antileukemic efficacy and reduced toxicity in newly diagnosed PML/RARalpha-positive acute promyelocytic leukemia. PETHEMA group. Blood 1999;94(9):3015-21.
10. Mandelli F, Diverio D, Avvisati G, et al. Molecular remission in PML/RAR alpha-positive acute promyelocytic leukemia by combined all-trans retinoic acid and idarubicin (AIDA) therapy. Gruppo Italiano-Malattie Ematologiche Maligne dell'Adulto and Associazione Italiana di Ematologia ed Oncologia Pediatrica Cooperative Groups. Blood 1997;90(3):1014-21.
11. Bots M, Verbrugge I, Martin BP, et al. Differentiation therapy for the treatment of t(8;21) acute myeloid leukemia using histone deacetylase inhibitors. Blood 2014;123(9):1341-52. DOI: 10.1182/blood-2013-03-488114.
12. Yen K, Travins J, Wang F, et al. AG-221, a First-in-Class Therapy Targeting Acute Myeloid Leukemia Harboring Oncogenic IDH2 Mutations. Cancer Discov 2017;7(5):478-493. DOI: 10.1158/2159-8290.CD-16-1034.
13. Medema JP. Targeting the Colorectal Cancer Stem Cell. N Engl J Med 2017;377(9):888-890. (In eng). DOI: 10.1056/NEJMcibr1706541.
14. Butte AJ, Kohane IS. Unsupervised knowledge discovery in medical databases using relevance networks. Proc AMIA Symp 1999:711-5.
15. Shameer K, Readhead B, Dudley JT. Computational and experimental advances in drug repositioning for accelerated therapeutic stratification. Curr Top Med Chem 2015;15(1):5-20.
16. Shen Y, Alvarez MJ, Bisikirska B, et al. Systematic, network-based characterization of therapeutic target inhibitors. PLoS Comput Biol 2017;13(10):e1005599. DOI: 10.1371/journal.pcbi.1005599.
17. Margolin AA, Wang K, Lim WK, Kustagi M, Nemenman I, Califano A. Reverse engineering cellular networks. Nat Protoc 2006;1(2):662-71. DOI: 10.1038/nprot.2006.106.
18. van Someren EP, Wessels LF, Backer E, Reinders MJ. Genetic network modeling. Pharmacogenomics 2002;3(4):507-25. DOI: 10.1517/14622416.3.4.507.
19. Kusonmano K. Gene Expression Analysis Through Network Biology: Bioinformatics Approaches. Adv Biochem Eng Biotechnol 2017;160:15-32. DOI: 10.1007/10_2016_44.
20. Allocco DJ, Kohane IS, Butte AJ. Quantifying the relationship between co-expression, co-regulation and gene function. BMC Bioinformatics 2004;5:18. DOI: 10.1186/1471-2105-5-18.
21. Arkin A, Ross J. Statistical Construction of Chemical Reaction Mechanisms from Measured Time-Series. The Journal of Physical Chemistry 1995;99(3):970-979. DOI: 10.1021/j100003a020.
22. Jordan IK, Marino-Ramirez L, Wolf YI, Koonin EV. Conservation and coevolution in the scale-free human gene coexpression network. Mol Biol Evol 2004;21(11):2058-70. DOI: 10.1093/molbev/msh222.
23. Lee HK, Hsu AK, Sajdak J, Qin J, Pavlidis P. Coexpression analysis of human genes across many microarray data sets. Genome Res 2004;14(6):1085-94. DOI: 10.1101/gr.1910904.

566   24.   Tavazoie S, Hughes JD, Campbell MJ, Cho RJ, Church GM. Systematic determination of genetic
567   network architecture. Nat Genet 1999;22(3):281-5. DOI: 10.1038/10343.
568   25.   Butte AJ, Tamayo P, Slonim D, Golub TR, Kohane IS. Discovering functional relationships between
569   RNA expression and chemotherapeutic susceptibility using relevance networks. Proc Natl Acad Sci
570   U S A 2000;97(22):12182-6. DOI: 10.1073/pnas.220392197.
571   26.   Paik S, Shak S, Tang G, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-
572   negative breast cancer. N Engl J Med 2004;351(27):2817-26. DOI: 10.1056/NEJMoa041588.
573   27.   Witten DM, Tibshirani R. Survival analysis with high-dimensional covariates. Stat Methods Med Res
574   2010;19(1):29-51. DOI: 10.1177/0962280209105024.
575   28.   Alizadeh AA, Eisen MB, Davis RE, et al. Distinct types of diffuse large B-cell lymphoma identified by
576   gene expression profiling. Nature 2000;403(6769):503-11. DOI: 10.1038/35000501.
577   29.   Zhao H, Ljungberg B, Grankvist K, Rasmuson T, Tibshirani R, Brooks JD. Gene expression profiling
578   predicts survival in conventional renal cell carcinoma. PLoS Med 2006;3(1):e13. DOI:
579   10.1371/journal.pmed.0030013.
580   30.   Sahoo D, Swanson L, Sayed IM, et al. Artificial intelligence guided discovery of a barrier-protective
581   therapy in inflammatory bowel disease. Nat Commun 2021;12(1):4246. (In eng). DOI:
582   10.1038/s41467-021-24470-5.
583   31.   Ghosh P, Sinha S, Katkar GD, et al. Machine learning identifies signatures of macrophage reactivity
584   and tolerance that predict disease outcomes. EBioMedicine 2023;94:104719. (In eng). DOI:
585   10.1016/j.ebiom.2023.104719.
586   32.   Sahoo D, Seita J, Bhattacharya D, et al. MiDReG: a method of mining developmentally regulated
587   genes using Boolean implications. Proc Natl Acad Sci U S A 2010;107(13):5732-7. (In eng). DOI:
588   10.1073/pnas.0913635107.
589   33.   Inlay MA, Bhattacharya D, Sahoo D, et al. Ly6d marks the earliest stage of B-cell specification and
590   identifies the branchpoint between B-cell and T-cell development. Genes Dev 2009;23(20):2376-81.
591   (In eng). DOI: 10.1101/gad.1836009.
592   34.   Chen JY, Miyanishi M, Wang SK, et al. Hoxb5 marks long-term haematopoietic stem cells and
593   reveals a homogenous perivascular niche. Nature 2016;530(7589):223-7. (In eng). DOI:
594   10.1038/nature16943.
595   35.   Chan CK, Seo EY, Chen JY, et al. Identification and specification of the mouse skeletal stem cell.
596   Cell 2015;160(1-2):285-98. (In eng). DOI: 10.1016/j.cell.2014.12.002.
597   36.   Dimov IK, Lu R, Lee EP, et al. Discriminating cellular heterogeneity using microwell-based RNA
598   cytometry. Nat Commun 2014;5:3451. (In eng). DOI: 10.1038/ncomms4451.
599   37.   Chan CK, Lindau P, Jiang W, et al. Clonal precursor of bone, cartilage, and hematopoietic niche
600   stromal cells. Proc Natl Acad Sci U S A 2013;110(31):12643-8. (In eng). DOI:
601   10.1073/pnas.1310212110.
602   38.   Seita J, Sahoo D, Rossi DJ, et al. Gene Expression Commons: an open platform for absolute gene
603   expression profiling. PLoS One 2012;7(7):e40321. (In eng). DOI: 10.1371/journal.pone.0040321.
604   39.   Cheah MT, Chen JY, Sahoo D, et al. CD14-expressing cancer cells establish the inflammatory and
605   proliferative tumor microenvironment in bladder cancer. Proc Natl Acad Sci U S A
606   2015;112(15):4725-30. (In eng). DOI: 10.1073/pnas.1424795112.
607   40.   Volkmer JP, Sahoo D, Chin RK, et al. Three differentiation states risk-stratify bladder cancer into
608   distinct subtypes. Proc Natl Acad Sci U S A 2012;109(6):2078-83. (In eng). DOI:
609   10.1073/pnas.1120605109.
610   41.   Shin K, Lim A, Zhao C, et al. Hedgehog signaling restrains bladder cancer progression by eliciting
611   stromal production of urothelial differentiation factors. Cancer Cell 2014;26(4):521-33. (In eng). DOI:
612   10.1016/j.ccell.2014.09.001.
613   42.   Sin MLY, Mach KE, Sinha R, et al. Deep Sequencing of Urinary RNAs for Bladder Cancer Molecular
614   Diagnostics. Clin Cancer Res 2017;23(14):3700-3710. (In eng). DOI: 10.1158/1078-0432.CCR-16-
615   2610.
616   43.   Sahoo D, Wei W, Auman H, et al. Boolean analysis identifies CD38 as a biomarker of aggressive
617   localized prostate cancer. Oncotarget 2018;9(5):6550-6561. (In eng). DOI:
618   10.18632/oncotarget.23973.
619   44.   Bhamre S, Sahoo D, Tibshirani R, Dill DL, Brooks JD. Temporal changes in gene expression
620   induced by sulforaphane in human prostate cancer cells. Prostate 2009;69(2):181-90. (In eng). DOI:
621   10.1002/pros.20869.
622   45.   Ghosh P, Campos VJ, Vo DT, et al. AI-assisted discovery of an ethnicity-influenced driver of cell
623   transformation in esophageal and gastroesophageal junction adenocarcinomas. JCI Insight
624   2022;7(18) (In eng). DOI: 10.1172/jci.insight.161334.

46. Bonhomme C, Duluc I, Martin E, et al. The Cdx2 homeobox gene has a tumour suppressor function in the distal colon in addition to a homeotic role during gut development. Gut 2003;52(10):1465-1471.

47. Silberg DG, Swain GP, Suh ER, Traber PG. Cdx1 and cdx2 expression during intestinal development. Gastroenterology 2000;119(4):961-971.

48. Chawengsaksophak K, James R, Hammond V, Köntgen F, Beck F. Homeosis and intestinal tumours in Cdx2 mutant mice. Nature 1997;386(6620):84-87.

49. Hryniuk A, Grainger S, Savory JG, Lohnes D. Cdx1 and Cdx2 function as tumor suppressors. Journal of Biological Chemistry 2014;289(48):33343-33354.

50. Sahoo D, Dill DL, Gentles AJ, Tibshirani R, Plevritis SK. Boolean implication networks derived from large scale, whole genome microarray datasets. Genome biology 2008;9(10):R157. (In eng).

51. Dalerba P, Kalisky T, Sahoo D, et al. Single-cell dissection of transcriptional heterogeneity in human colon tumors. Nat Biotechnol 2011;29(12):1120-7. (In eng).

52. Dalerba P, Sahoo D, Paik S, et al. CDX2 as a prognostic biomarker in stage II and stage III colon cancer. New England Journal of Medicine 2016;374(3):211-222.

53. Levin TG, Powell AE, Davies PS, et al. Characterization of the intestinal cancer stem cell marker CD166 in the human and mouse gastrointestinal tract. Gastroenterology 2010;139(6):2072-2082 e5. DOI: 10.1053/j.gastro.2010.08.053.

54. Weichert W, Knosel T, Bellach J, Dietel M, Kristiansen G. ALCAM/CD166 is overexpressed in colorectal carcinoma and correlates with shortened patient survival. Journal of clinical pathology 2004;57(11):1160-4. (In eng).

55. Dalerba P, Dylla SJ, Park IK, et al. Phenotypic characterization of human colorectal cancer stem cells. Proc Natl Acad Sci U S A 2007;104(24):10158-63. (In eng).

56. Derangère V, Lecuelle J, Lepage C, et al. Combination of CDX2 H-score quantitative analysis with CD3 AI-guided analysis identifies patients with a good prognosis only in stage III colon cancer. Eur J Cancer 2022;172:221-230. (In eng). DOI: 10.1016/j.ejca.2022.05.040.

57. Bennedsen ALB, Cai L, Hasselager RP, et al. An exploration of immunohistochemistry-based prognostic markers in patients undergoing curative resections for colon cancer. BMC Cancer 2022;22(1):62. (In eng). DOI: 10.1186/s12885-022-09169-0.

58. Azcue P, Guerrero Setas D, Encío I, et al. A Novel Prognostic Biomarker Panel for Early-Stage Colon Carcinoma. Cancers (Basel) 2021;13(23) (In eng). DOI: 10.3390/cancers13235909.

59. den Uil SH, de Wit M, Slebos RJC, et al. Quantitative analysis of CDX2 protein expression improves its clinical utility as a prognostic biomarker in stage II and III colon cancer. Eur J Cancer 2021;144:91-100. (In eng). DOI: 10.1016/j.ejca.2020.10.029.

60. Tarazona N, Gimeno-Valiente F, Gambardella V, et al. Detection of postoperative plasma circulating tumour DNA and lack of CDX2 expression as markers of recurrence in patients with localised colon cancer. ESMO Open 2020;5(5):e000847. (In eng). DOI: 10.1136/esmoopen-2020-000847.

61. Hestetun KE, Aasebø K, Rosenlund NB, Müller Y, Dahl O, Myklebust MP. Mismatch repair phenotype determines the implications of tumor grade and CDX2 expression in stage II-III colon cancer. Mod Pathol 2021;34(1):161-170. (In eng). DOI: 10.1038/s41379-020-0634-9.

62. Asgari-Karchekani S, Karimian M, Mazoochi T, Taheri MA, Khamehchian T. CDX2 Protein Expression in Colorectal Cancer and ItsCorrelation with Clinical and Pathological Characteristics, Prognosis, and Survival Rate of Patients. J Gastrointest Cancer 2020;51(3):844-849. (In eng). DOI: 10.1007/s12029-019-00314-w.

63. Ma C, Olevian D, Miller C, et al. SATB2 and CDX2 are prognostic biomarkers in DNA mismatch repair protein deficient colon cancer. Mod Pathol 2019;32(8):1217-1231. (In eng). DOI: 10.1038/s41379-019-0265-1.

64. Ben-Aharon I, Goshen-Lago T, Sternschuss M, et al. Sidedness Matters: Surrogate Biomarkers Prognosticate Colorectal Cancer upon Anatomic Location. Oncologist 2019;24(8):e696-e701. (In eng). DOI: 10.1634/theoncologist.2018-0351.

65. Hansen TF, Kjær-Frifeldt S, Eriksen AC, et al. Prognostic impact of CDX2 in stage II colon cancer: results from two nationwide cohorts. British Journal of Cancer 2018;119(11):1367-1373.

66. Pilati C, Taieb J, Balogoun R, Marisa L, De Reyniès A, Laurent-Puig P. CDX2 prognostic value in stage II/III resected colon cancer is related to CMS classification. Annals of Oncology 2017;28(5):1032-1035.

67. Nolte S, Zlobec I, Lugli A, et al. Construction and analysis of tissue microarrays in the era of digital pathology: a pilot study targeting CDX1 and CDX2 in a colon cancer cohort of 612 patients. J Pathol Clin Res 2017;3(1):58-70. (In eng). DOI: 10.1002/cjp2.62.

68.   Zhang BY, Jones JC, Briggler AM, et al. Lack of caudal-type homeobox transcription factor 2
expression as a prognostic biomarker in metastatic colorectal cancer. Clinical colorectal cancer
2017;16(2):124-128.
69.   Tomasello G, Barni S, Turati L, et al. Association of CDX2 expression with survival in early
colorectal cancer: a systematic review and meta-analysis. Clinical colorectal cancer 2018;17(2):97-
103.
70.   Sjoerd H, de Wit M, Slebos RJ, et al. Quantitative analysis of CDX2 protein expression improves its
clinical utility as a prognostic biomarker in stage II and III colon cancer. European Journal of Cancer
2021;144:91-100.
71.   Bruun J, Sveen A, Barros R, et al. Prognostic, predictive, and pharmacogenomic assessments of
CDX 2 refine stratification of colorectal cancer. Molecular oncology 2018;12(9):1639-1655.
72.   Baba Y, Nosho K, Shima K, et al. Relationship of CDX2 loss with molecular features and prognosis
in colorectal cancer. Clinical cancer research 2009;15(14):4665-4673.
73.   Olsen J, Eiholm S, Kirkeby L, et al. CDX2 downregulation is associated with poor differentiation and
MMR deficiency in colon cancer. Experimental and molecular pathology 2016;100(1):59-66.
74.   Sahoo D, Dill DL, Tibshirani R, Plevritis SK. Extracting binary signals from microarray time-course
data. Nucleic Acids Res 2007;35(11):3705-12. DOI: 10.1093/nar/gkm284.
75.   Ghosh P. The stress polarity pathway: AMPK 'GIV'-es protection against metabolic insults. Aging
(Albany NY) 2017;9(2):303-314. DOI: 10.18632/aging.101179.
76.   Ghosh P, Swanson L, Sayed IM, et al. The stress polarity signaling (SPS) pathway serves as a
marker and a target in the leaky gut barrier: implications in aging and cancer. Life Sci Alliance
2020;3(3) (In eng). DOI: 10.26508/lsa.201900481.
77.   Aznar N, Patel A, Rohena CC, et al. AMP-activated protein kinase fortifies epithelial tight junctions
during energetic stress via its effector GIV/Girdin. Elife 2016;5 (In eng). DOI: 10.7554/eLife.20795.
78.   Gough NR. Maintaining Polarity in the Face of Energetic Stress. Science's STKE
2007;2007(386):tw166-tw166. DOI: doi:10.1126/stke.3862007tw166.
79.   Graule J, Uth K, Fischer E, et al. CDX2 in colorectal cancer is an independent prognostic factor and
regulated by promoter methylation and histone deacetylation in tumors of the serrated pathway.
Clinical epigenetics 2018;10(1):1-12.
80.   Boman BM, Huang E. Human colon cancer stem cells: a new paradigm in gastrointestinal oncology.
J Clin Oncol 2008;26(17):2828-38. (In eng). DOI: 10.1200/JCO.2008.17.6941.
81.   Kalisky T, Rajendran PS, Sahoo D, et al. Analysis of human colon tissue cell composition using
single-cell gene-expression PCR. Journal of biomolecular techniques: JBT 2013;24(Suppl):S11.
82.   Malta TM, Sokolov A, Gentles AJ, et al. Machine Learning Identifies Stemness Features Associated
with Oncogenic Dedifferentiation. Cell 2018;173(2):338-354 e15. DOI: 10.1016/j.cell.2018.03.034.
83.   Friedmann-Morvinski D, Verma IM. Dedifferentiation and reprogramming: origins of cancer stem
cells. EMBO Rep 2014;15(3):244-53. (In eng). DOI: 10.1002/embr.201338254.
84.   Guinney J, Dienstmann R, Wang X, et al. The consensus molecular subtypes of colorectal cancer.
Nat Med 2015;21(11):1350-6. (In eng). DOI: 10.1038/nm.3967.
85.   Valladares-Ayerbes M, Blanco-Calvo M, Reboredo M, et al. Evaluation of the adenocarcinoma-
associated gene AGR2 and the intestinal stem cell marker LGR5 as biomarkers in colorectal
cancer. Int J Mol Sci 2012;13(4):4367-4387. (In eng). DOI: 10.3390/ijms13044367.
86.   Li Y, Yuan Y, Yang L, et al. AT7867 Inhibits the Growth of Colorectal Cancer Stem-Like Cells and
Stemness by Regulating the Stem Cell Maintenance Factor Ascl2 and Akt Signaling. Stem Cells Int
2023;2023:4199052. (In eng). DOI: 10.1155/2023/4199052.
87.   Jubb AM, Chalasani S, Frantz GD, et al. Achaete-scute like 2 (ascl2) is a target of Wnt signalling
and is upregulated in intestinal neoplasia. Oncogene 2006;25(24):3445-57. (In eng). DOI:
10.1038/sj.onc.1209382.
88.   Durinikova E, Kozovska Z, Poturnajova M, et al. ALDH1A3 upregulation and spontaneous
metastasis formation is associated with acquired chemoresistance in colorectal cancer cells. BMC
Cancer 2018;18(1):848. (In eng). DOI: 10.1186/s12885-018-4758-y.
89.   Chen G, She X, Yin Y, et al. Targeting TM4SF1 exhibits therapeutic potential via inhibition of cancer
stem cells. Signal Transduct Target Ther 2022;7(1):350. (In eng). DOI: 10.1038/s41392-022-01177-
7.
90.   Tang Q, Chen J, Di Z, et al. TM4SF1 promotes EMT and cancer stemness via the Wnt/β-
catenin/SOX2 pathway in colorectal cancer. J Exp Clin Cancer Res 2020;39(1):232. (In eng). DOI:
10.1186/s13046-020-01690-z.
91.   Matsui WH. Cancer stem cell signaling pathways. Medicine (Baltimore) 2016;95(1 Suppl 1):S8-S19.
DOI: 10.1097/MD.0000000000004765.

92. Yang L, Shi P, Zhao G, et al. Targeting cancer stem cell pathways for cancer therapy. Signal
Transduct Target Ther 2020;5(1):8. DOI: 10.1038/s41392-020-0110-5.
93. Gierut J, Zheng Y, Bie W, et al. Disruption of the mouse protein tyrosine kinase 6 gene prevents
STAT3 activation and confers resistance to azoxymethane. Gastroenterology 2011;141(4):1371-80,
1380.e1-2. (In eng). DOI: 10.1053/j.gastro.2011.06.071.
94. Liu C, Pan Z, Chen Q, et al. Pharmacological targeting PTK6 inhibits the JAK2/STAT3 sustained
stemness and reverses chemoresistance of colorectal cancer. J Exp Clin Cancer Res
2021;40(1):297. (In eng). DOI: 10.1186/s13046-021-02059-6.
95. Li X, Sun K, Liao X, Gao H, Zhu H, Xu R. Colorectal carcinomas with mucinous differentiation are
associated with high frequent mutation of KRAS or BRAF mutations, irrespective of quantity of
mucinous component. BMC Cancer 2020;20(1):400. DOI: 10.1186/s12885-020-06913-2.
96. Feng Y, Bommer GT, Zhao J, et al. Mutant KRAS promotes hyperplasia and alters differentiation in
the colon epithelium but does not expand the presumptive stem cell pool. Gastroenterology
2011;141(3):1003-1013 e1-10. DOI: 10.1053/j.gastro.2011.05.007.
97. Aasebø K, Dragomir A, Sundström M, et al. CDX2: A Prognostic Marker in Metastatic Colorectal
Cancer Defining a Better. Front Oncol 2020;10:8. (In eng). DOI: 10.3389/fonc.2020.00008.
98. Brandt R, Sell T, Luthen M, et al. Cell type-dependent differential activation of ERK by oncogenic
KRAS in colon cancer and intestinal epithelium. Nat Commun 2019;10(1):2919. DOI:
10.1038/s41467-019-10954-y.
99. Aasebø K, Dragomir A, Sundström M, et al. CDX2: a prognostic marker in metastatic colorectal
cancer defining a better BRAF mutated and a worse KRAS mutated subgroup. Frontiers in oncology
2020;10:8.
100. Slik K, Turkki R, Carpén O, et al. CDX2 loss with microsatellite stable phenotype predicts poor
clinical outcome in stage II colorectal carcinoma. The American journal of surgical pathology
2019;43(11):1473-1482.
101. Moertel CG, Fleming TR, Macdonald JS, et al. Levamisole and fluorouracil for adjuvant therapy of
resected colon carcinoma. New England Journal of Medicine 1990;322(6):352-358.
102. Sobrero AF, Puccini A, Shi Q, et al. A new prognostic and predictive tool for shared decision making
in stage III colon cancer. European Journal of Cancer 2020;138:182-188.
103. Taieb J, André T, Auclin E. Refining adjuvant therapy for non-metastatic colon cancer, new
standards and perspectives. Cancer Treatment Reviews 2019;75:1-11.
104. Delhorme J-B, Bersuder E, Terciolo C, et al. CDX2 controls genes involved in the metabolism of 5-
fluorouracil and is associated with reduced efficacy of chemotherapy in colorectal cancer.
Biomedicine & Pharmacotherapy 2022;147:112630.
105. Takakura Y, Hinoi T, Oue N, et al. CDX2 regulates multidrug resistance 1 gene expression in
malignant intestinal epithelium. Cancer research 2010;70(17):6767-6778.
106. Shigematsu Y, Inamura K, Mise Y, et al. CDX2 expression is concordant between primary colorectal
cancer lesions and corresponding liver metastases independent of chemotherapy: a single-center
retrospective study in Japan. Oncotarget 2018;9(24):17056.
107. Fontana E, Meyers J, Sobrero A, et al. Early-Onset Colorectal Adenocarcinoma in the IDEA
Database: Treatment Adherence, Toxicities, and Outcomes With 3 and 6 Months of Adjuvant
Fluoropyrimidine and Oxaliplatin. Journal of Clinical Oncology 2021;39(36):4009-4019.
108. Tong K, Pellón-Cárdenas O, Sirihorachai VR, et al. Degree of tissue differentiation dictates
susceptibility to BRAF-driven colorectal cancer. Cell reports 2017;21(13):3833-3845.
109. Sakamoto N, Feng Y, Stolfi C, et al. BRAFV600E cooperates with CDX2 inactivation to promote
serrated colorectal tumorigenesis. Elife 2017;6:e20331.
110. Felipe De Sousa EM, Wang X, Jansen M, et al. Poor-prognosis colon cancer is defined by a
molecularly distinct subtype and develops from serrated precursor lesions. Nature medicine
2013;19(5):614-618.
111. Coskun M. The role of CDX2 in inflammatory bowel disease. Danish medical journal
2014;61(3):B4820-B4820.
112. Coskun M, Olsen AK, Holm TL, et al. TNF-α-induced down-regulation of CDX2 suppresses MEP1A
expression in colitis. Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease
2012;1822(6):843-851.
113. Yin F, Xie H, Lai J, et al. Double negativity for expression of YAP1 and CDX2 defines an aggressive
type of colitis-associated cancer. Anticancer research 2020;40(10):5411-5416.
114. Kawai H, Tomii K, Toyooka S, et al. Promoter methylation downregulates CDX2 expression in
colorectal carcinomas. Oncology reports 2005;13(3):547-551.

115.   Sun X, Yang Q, Rogers CJ, Du M, Zhu M-J. AMPK improves gut epithelial differentiation and barrier
function via regulating Cdx2 expression. Cell Death & Differentiation 2017;24(5):819-831.
116.   Bae JM, Lee TH, Cho N-Y, Kim T-Y, Kang GH. Loss of CDX2 expression is associated with poor
prognosis in colorectal cancer patients. World journal of gastroenterology: WJG 2015;21(5):1457.
117.   Reggiani Bonetti L, Lionti S, Vitarelli E, Barresi V. Prognostic significance of CDX2
immunoexpression in poorly differentiated clusters of colorectal carcinoma. Virchows Archiv
2017;471(6):731-741.
118.   Alarid-Escudero F. CDX2 testing+ FOLFOX for CDX2-negative colon cancer patients cost-effective
strategy. PharmacoEconomics & Outcomes News 2021;892:9-27.
119.   Argilés G, Tabernero J, Labianca R, et al. Localised colon cancer: ESMO Clinical Practice
Guidelines for diagnosis, treatment and follow-up. Annals of Oncology 2020;31(10):1291-1305.
120.   Benson AB, Venook AP, Al-Hawary MM, et al. Colon cancer, version 2.2021, NCCN clinical practice
guidelines in oncology. Journal of the National Comprehensive Cancer Network 2021;19(3):329-
359.
121.   Committee ACNCCGR. Guidelines for the prevention, early detection and management of
colorectal cancer. Sydney: The Cancer Council Australia and Australian Cancer Network 2005.
122.   Van Cutsem E, Cervantes A, Adam R, et al. ESMO consensus guidelines for the management of
patients with metastatic colorectal cancer. Annals of Oncology 2016;27(8):1386-1422.
123.   Baum M, Fallowfield L, Farewell V, Macbeth F, Treasure T. NICEGuidelines: management of
colorectal cancer metastases. British Journal of Surgery 2020;107(9):e357-e357.
124.   Foucquier J, Guedj M. Analysis of drug combinations: current methodological landscape. Pharmacol
Res Perspect 2015;3(3):e00149. (In eng). DOI: 10.1002/prp2.149.
322