Boosting biodiversity monitoring using smartphone-driven, rapidly accumulating community-sourced data

Keisuke Atsumi,[1] Yuusuke Nishida[1], Masayuki Ushio[2,3,4], Hirotaka Nishi[5], Takanori Genroku[1], Shogoro Fujiki[1*]

[1]*Biome Inc., 134 Chudoji Minamimachi, Shimogyo-ku, Kyoto 600-8813, Japan*

[2]*Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR*

[3]*Hakubi Center, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto 606-8501, Japan*

[4]*Center for Ecological Research, Kyoto University, 2-509-3 Hirano, Otsu, Shiga 520-2113, Japan*

[5]*Toyohashi Museum of Natural History, 1-238 Oana, Oiwa, Toyohashi, Aichi 441-3147, Japan*

*Correspondence: fujiki@biome.co.jp

Abstract

Comprehensive biodiversity data is crucial for ecosystem protection. The '*Biome*' mobile app, launched in Japan, efficiently gathers species observations from the public using species identification algorithms and gamification elements. The app has amassed >6M observations since 2019. Nonetheless, community-sourced data may exhibit spatial and taxonomic biases. Species distribution models (SDMs) estimate species distribution while accommodating such bias. Here, we investigated the quality of *Biome* data and its impact on SDM performance. Species identification accuracy exceeds 95% for birds, reptiles, mammals, and amphibians, but seed plants, mollusks, and fishes scored below 90%. Our SDMs for 132 terrestrial plants and animals across Japan revealed that incorporating *Biome* data into traditional survey data improved accuracy. For endangered species, traditional survey data required >2,000 records for accurate models (Boyce index ≥ 0.9), while blending the two data sources reduced this to around 300. The uniform coverage of urban-natural gradients by *Biome* data, compared to traditional data biased towards natural areas, may explain this improvement. Combining multiple data sources better estimates species distributions, aiding in protected area designation and ecosystem service assessment. Establishing a platform for accumulating community-sourced distribution data will contribute to conserving and monitoring natural ecosystems.

35 **KEY WORDS:**

36 **Community science, Citizen science, Habitat suitability model, MaxEnt, Big data,**
37 **Taskforce on Nature-related Financial Disclosures: TNFD, Other Effective area-based**
38 **Conservation Measures: OECMs, Natural capital**

41 ──

Introduction

Nature underpins human society, and the conservation of ecosystems and associated
ecosystem services contributes to the sustainable development of human society, yet
these services have been rapidly declining in recent years (IPBES, 2019; Loh et al., 2005;
Newbold et al., 2016; Scholes and Biggs, 2005). Kunming-Montreal Global Biodiversity
Framework (KM-GBF) by the United Nations envisions reversing the nature loss by 2030.
As direct means for nature conservation, KM-GBF targeted making 30% of Earth's land
and ocean area as protected areas by 2030 (i.e. 30by30). As an indirect but influential
way, KM-GBF requires companies to "*monitor, assess, and transparently disclose their risks,*
dependencies and impacts on biodiversity through their operations, supply and value chains
and portfolios," which is guided by Taskforce on Nature-related Financial Disclosures
(TNFD) (TNFD, 2023). To achieve these goals, it is imperative to assess the state of
biodiversity with a sufficient spatiotemporal resolution to support conservation
planning, adaptive management, and companies' annual nature-related financial
disclosures. The basis for such assessments lies in our knowledge of species
distributions (Gonzalez et al., 2023; Newbold et al., 2016). Traditionally, distribution data
was acquired through on-site surveys by experts (people have expertise about
biodiversity), but collecting distribution data with sufficient spatiotemporal resolution is
challenging if we rely only on such limited human resources (Miya et al., 2022; Mori et al.,
2023; Pocock et al., 2018).

Since the emergence of digital devices and the internet, people have been able to share
their observations through various media, such as images and video/audio recordings.
Such community-sourced data have significantly contributed to the accumulation of
ecosystem information. These datasets have been instrumental in assessing the
impacts of climate change and urbanisation on phenology (Fuccillo Battle et al., 2022;
Klinger et al., 2023), detecting distribution changes including invasive alien species
(Larson et al., 2020; Roy et al., 2023; Wallace and Bargeron, 2014), exploring large-scale

geographic variations in traits (Atsumi and Koizumi, 2017; Leighton et al., 2016), and
estimating species distributions (Chandler et al., 2017; Feldman et al., 2021; Johnston et
al., 2018; Steen et al., 2019). Moreover, the utilisation of machine learning to describe
population trends based on community-sourced data (Fink et al., 2023) offers
opportunities for conducting time-series analyses. These analyses can help us
understand community assembly processes, unravel species interaction networks, and
assess ecosystem stability (Cornwell and Ackerly, 2009; Tilman et al., 2006; Ushio et al.,
2018), capitalising on the spatio-temporally dense sampling effort facilitated by
community-sourced data (Chandler et al., 2017; Kobori et al., 2016; Pocock et al., 2017).
Such analytical approaches enable us to make informed predictions about changes in
species distribution, population dynamics, and ecosystem stability in the face of climate
change (Bury et al., 2021; Pennekamp et al., 2019; Urban et al., 2016). In essence,
community-sourced data, owing to its extensive sampling across time and space, has
the potential to test existing ecological theories, expand our comprehension of
ecosystems and the underlying processes, eventually allowing us to forecast ecological
dynamics in the context of climate change.

When people photograph organisms using digital devices with GPS capabilities, the
images often contain timestamps and location details. Such images, when accompanied
by species identifications, serve as evidence for tracking phenology and species
occurrences. This crowdsourcing approach has been particularly successful on web- or
mobile-based platforms such as eBird and iNaturalist (Chandler et al., 2017; Wood et al.,
2011). Individuals submit records to these platforms for various reasons, including a
desire to contribute to science and engage with cutting-edge technologies (Herodotou
et al., 2023; Kaplan Mintz et al., 2023). By making the process more enjoyable (i.e.
gamification), we can potentially gather even more biological data from the public
(Bowser et al., 2013; Ponti et al., 2015). Yet, the collection process of Community-
sourced data is usually not well-designed (e.g., spatially biassed "presence-only" data)
(Feldman et al., 2021; Steen et al., 2019) and its interpretation is challenging without
proper statistical modelling. Thus, although much effort has been invested in
developing effective monitoring and modelling methods for biodiversity assessment,
current approaches can be further improved by incorporating (i) more enjoyable
community-based survey platforms using mobile applications and (ii) employing an
advanced statistical modelling framework in estimating species distribution.

To fuel communities' engagement in biodiversity surveys and environmental education,
we launched the mobile application '*Biome*' in 2019 in Japan (Fujiki and Tatsuno, 2021).
For supporting species identification, *Biome* implements artificial intelligence (AI)
algorithms that generate lists of potential species and enable users to seek
help/suggestions from others for species identification (Figure 1) as in other
applications such as iNaturalist and eBird. The unique feature of *Biome* is gamification

which offers enjoyable experiences and facilitates communication among users (Fujiki
and Tatsuno, 2021; Koide et al., 2023). For example, users can earn "points" by
contributing in various ways such as submitting records and suggesting species
identifications to others, and their levels are determined based on the total points
earned. The inclusion of networking and gamification elements can attract a wider user
base, including those who may not typically engage in community science (Bowser et al.,
2013; Groom et al., 2021). Consequently, *Biome* has accumulated data rapidly. Since its
launch, 6 million records have been collected through the app (by 17 October 2023).
This is more than four times greater than the number of records accumulated by GBIF
(Global Biodiversity Information Facility) from any data sources including iNaturalist and
eBird during the same period in Japan (ca. 1.3 million). The data gathered through the
app has been used for conservation planning and facilitating companies' financial
disclosures by supplying and analysing species occurrence records.

Species distribution models (SDMs) are effective statistical tools for assessing
biodiversity at specific sites while accounting for biases in survey efforts. SDMs use
species occurrence records and environmental conditions to estimate the potential
geographic ranges and suitable habitats for species (Booth et al., 2014; Box, 1981; Elith
et al., 2011; Hutchinson, 1957; Phillips et al., 2006). These models play a crucial role in
conservation and restoration planning by helping predict how changes in land use and
climate impact species distributions (Kindt, 2023; Porfirio et al., 2014; Urban et al., 2016).
While species presence/absence data—which needs extensive surveys by experts—is
limited, presence-only data—which can be obtained from communities' observations—
is much more available. MaxEnt (Phillips et al., 2006; Phillips and Dudík, 2008) is one of
the most popular SDM methods due to its computational efficiency and estimation
accuracy (Valavi et al., 2022). It can estimate species distribution from presence-only
data by maximising the entropy of the probability distribution while satisfying
constraints based on the available information (Elith et al., 2011; Phillips and Dudík,
2008). Since MaxEnt only requires occurrence records, it is well-suited for empowering
community-based observations to predict species distributions. Also, while community-
sourced data often suffer from spatially-biassed sampling efforts (i.e., sampling tends to
concentrate in densely populated or touristic areas (Kendal et al., 2020; Reddy and
Dávalos, 2003)), SDMs such as MaxEnt can account for such spatial biases by
considering the spatial distribution of sampling efforts when selecting pseudo-absence
(background) locations (Milanesi et al., 2020; Phillips et al., 2009). When sampling efforts
are adequately controlled, adding community-sourced data improves the accuracy of
SDMs (Johnston et al., 2018; Robinson et al., 2020; Steen et al., 2019). This implies that
SDMs may be substantially improved by utilising rapidly accumulating *Biome*'s species
occurrence records if we adequately control the sampling efforts.

Here, we show the quality of community-based data gathered through the smartphone
app *Biome*, and how the data improves the prediction accuracy of species distribution.
First, we assess the quality of occurrence records by investigating the fractions of non-
wild and misidentified records. Second, we built SDMs based on two types of data: (i)
traditional survey data (e.g. forest inventory census, museum specimens and records
extracted from published researches) only and (ii) a mixture of traditional survey and
Biome data. We then compare the performance of the two SDMs. We modelled the
distributions of 132 terrestrial animals and seed plants in the Japanese archipelago
which covers subtropical to boreal areas. We finally discuss how our SDMs relying on
community-sourced data may contribute to meeting the goals of GBF.

____

Results

The amount and quality of Biome data

By 7 July 2023, *Biome* had accumulated 5,275,457 occurrence records of 40,957 species
across the Japanese archipelago (Figure 2A). The amount of occurrence records
submitted to *Biome* has increased across the years (Figure 2B). On average in 2022,
users submitted 5,407 records per day. The distribution of data along environmental
gradients somewhat differs between *Biome* and Traditional survey data. To elucidate
this distinction, we employed principal component (PC) analysis to summarise all
environmental variables. The two datasets demonstrated divergent distribution
patterns along PC1 (Figure 2C). This component, accounting for 6.1% of the total
variation, is primarily influenced by land use, topography, and climate (Supplementary
File 1). Among the environmental variables, a notable contrast between the datasets
was observed in relation to the natural-urban gradient. The *Biome* data exhibited a
relatively uniform distribution encompassing the entire gradient, while Traditional
survey data was substantially biassed towards natural areas (Figure 2C). The majority of
records are attributed to insects (31.2%) and to seed plants (41.8%), which are relatively
accessible and can be easily photographed using smartphones (Figure 2D).

Out of all the records submitted to *Biome*, a total of 2,373,303 records (45.0%)
successfully passed through the automatic filtering process. This dataset, referred to as

the *Biome* data, is utilised for subsequent investigations. The quality of *Biome* data
varied across taxa and the rarities of species (Table 1). The fraction of the records of
wild individuals exceeded 97% in insects and birds, while it was lower than 90% in
molluscs, seed plants, mammals and fishes. Among the records of wild individuals, at
the species level, identification accuracy was higher than 95% in birds, reptiles,
mammals and amphibians but less than 90% in insects, fishes and seed plants. At the
genus level, identification accuracy was higher than 90% in all taxa except for insects. In
the case of fishes and seed plants, identifications became 5-6% more accurate at the
genus level compared to the species level. The family was correctly identified in more
than 94% of records in all taxa examined. Common species had higher identification
accuracy than rare species (average value, 95% vs. 87%). This tendency was prominent
in insects and seed plants, but less in the other taxa. These results suggest that
identifying rare species in taxonomically diverse taxa (i.e. seed plants and insects) is a
challenging task.

The performance of species distribution models

SDMs using *Biome*+Traditional data, including *Biome* data at 50%, were more accurate
than those modelled only using Traditional survey data when the two datasets have the
same amount of occurrence records (Figure 3). Our analysis revealed that although the
intercept of the Boyce Index (BI, model accuracy metric that ranges between -1 to 1) did
not differ between the two datasets (generalised linear mixed model, see Methods:
β=0.02±0.03, t=0.60, P=0.55), *Biome*+Traditional data consistently led to a more rapid
increase in SDM accuracy as the amount of data increased, comparing to models solely
relying on Traditional survey data (β=0.02±0.01, t=3.72, P<0.001).

When compared to SDMs using Traditional survey data, those using *Biome*+Traditional
data achieved a high level of accuracy with a much smaller amount of data. For instance,
BI which ranges from –1 to 1, exceeds 0.9 with 294±471 records (mean±SD across all
species) in the *Biome*+Traditional data, whereas the Traditional survey data requires
2,129±4,157 records to achieve the same accuracy. This was also true in endangered
species (included in Japanese national or prefectural red lists); although 2,336±3,718
Traditional survey records were required to exceed 0.9 of BI, only 338±571 were
required for *Biome*+Traditional data.

Because we controlled the proportion of *Biome* data within the *Biome*+Traditional data
as 50%, the amount of records of the *Biome*+Traditional data is often limited. In cases
where a species had less *Biome* data compared to Traditional survey data, the total
amount of records of *Biome*+Traditional data ends up being smaller than that of

Traditional survey data alone. Therefore, the two datasets did not differ in the best
model performances in each species (BIs of *Biome*+Traditional data: 0.81±0.20;
Traditional survey data: 0.83±0.20).

Discussion

Biome: The amount and quality of submitted data

Since its launch in 2019, the app *Biome* accumulates species occurrence data rapidly
(Figure 2). Despite our concerted efforts to engage non-expert users through
gamification features, it is important to acknowledge that an excessive influx of non-
expert users could potentially compromise the quality of the collected data. This could
manifest in misidentifications or incomplete documentation, such as failing to
appropriately label non-wild individuals. We thus have developed algorithms to exclude
such suspicious records based on the features of records and users' behaviour on the
app. The implementation of automatic data filtering techniques is expected to enhance
the quality of the data, although further refinement is necessary. Notably, for insects
and birds, which encompass numerous species that can be kept in captivity, the
majority of records that underwent filtering procedures were restricted to observations
of wild individuals. Yet, the fraction of non-wild individuals is high in several taxa such as
fishes and seed plants. In response, we have updated the posting flow in the app to
prompt users to differentiate between non-wild and wild individuals. Further analysis is
warranted to evaluate the impact of this update on data quality.

Once we could exclude non-wild individuals, species identification accuracy exceeded
95% in taxa with moderate species diversity (amphibians, reptiles, birds and mammals).
In seed plants, *Biome'*s species Identification accuracy was 90%, which is higher than the
accuracy of auto-suggest identification by commonly used apps for plants (69%,
PlantNet, PlantSnap, LeafSnap, iNaturalist and Google Lens: (Hart et al., 2023)). During
the invasive plants survey in the US, the reports by non-professional volunteers were
72% correct (Crall et al., 2011). The higher accuracy of species identification in *Biome*
data can be attributed to two key factors. Firstly, the vigilant oversight of the user
community through the "suggest identification" feature plays a crucial role. *Biome*
encourages users to participate in suggesting identifications by offering "points" as
rewards for their contributions. Secondly, the species identification AI algorithm

leverages past occurrence data from nearby areas, resulting in increasingly accurate
automatic identifications as the data accumulates. Given these, as a community science
app, the data quality of *Biome* is decent. Yet, rare species generally showed lower
identification accuracy, which would require identification by experts and further
improvement of species identification AI algorithm.

Species distribution modelling

The inclusion of *Biome* data resulted in improved accuracy of SDMs (Figure 3). The most
accurate model predictions were obtained when the training data consisted of 50-70%
Biome data (Appendix 1), highlighting the necessity of incorporating both traditional
surveys and citizen observations for a comprehensive understanding of species
distributions (Miller et al., 2019; Pacifici et al., 2017; Robinson et al., 2020).

The improvement can be attributed to introducing data with different biases compared
to the Traditional survey data. Indeed, when controlling for the number of occurrence
records, the model performance was higher in the *Biome*+Traditional data compared to
the Traditional survey data. The variation in performance can be attributed to the
distribution of data in relation to environmental conditions. Traditional survey data
exhibits a strong bias towards natural areas, whereas *Biome* data is well balanced
across the natural-urban habitat gradients (Figure 2C). Therefore, incorporating *Biome*
data could significantly enhance modelling accuracy in urban and suburban landscapes,
which are typically underrepresented in traditional survey data. As pseudo-absences are
selected based on search effort, our models utilise numerous pseudo-absences from
these areas. Consequently, this might lead to better estimation of species absence in
such areas, not just presence, resulting in an overall increase in model accuracy across
a wider range of species. A balanced distribution along with the natural-urban gradient
is noteworthy because community science data is typically biassed towards human
population centres (Kendal et al., 2020; Reddy and Dávalos, 2003). This could be
influenced by the distribution of users' residencies, although we do not have specific
information about the users' locations. The app has collaborated with numerous local
governments across Japan, including nine prefectures and 29 local municipalities such
as cities and towns. Through these collaborations, the user base may be widely
dispersed, enriching the geographical coverage of *Biome* data.

The *Biome* data also can improve SDM accuracy by simply increasing the overall amount
of data. Essentially, SDM accuracy is enhanced with an increased amount of data (Figure
3) (Erickson and Smith, 2023; Stockwell and Peterson, 2002). In our analysis, we
maintained a fixed proportion of 50% for *Biome* data within the *Biome*+Traditional

dataset, which in turn restricted the amount of available *Biome*+Traditional data.
However, our preliminary analysis (Appendix 1) demonstrates that the enhancement of
SDM accuracy occurs across a range of proportion variations for *Biome* data blending.
This implies that the proportion of *Biome* data does not necessarily need to be
controlled. Therefore, in practical application scenarios, the incorporation of *Biome* data
predominantly serves to augment the overall volume of training data.

The impact of community-sourced data on SDMs has primarily been investigated using
birds, with a limited focus on plants (Feldman et al., 2021). In our investigation, we
observed that incorporating *Biome* data improved SDM accuracy for seed plants and
insects, while the impact on birds remained unclear (Figure 3). This ambiguity is likely
because community-sourced data from platforms such as eBird are already
incorporated in Traditional data through GBIF. In comparison to other taxonomic
groups, our results indicate that seed plants exhibited lower model accuracy when
evaluated against both *Biome*+Traditional survey data (Figure 3) and Traditional survey
data alone (Figure 3–figure supplement 1). The variation in model accuracy among
taxonomic groups may be attributed to data quality issues in both *Biome* and
Traditional survey data. For instance, in *Biome* data, while the fractions of wild
individuals were high in birds and insects, it was lower for seed plants (Table 1).
Compared with other taxa, distinguishing between wild and non-wild individuals can be
particularly difficult in plants when they are planted outside. In addition, identifying
plant species may be challenging in certain taxa, primarily due to the absence of key
identification traits on leaves and stems. This becomes especially problematic when
flowers are not present. These difficulties could potentially impact the quality of
Traditional data as well. Although few studies have simultaneously assessed the quality
of community-sourced data and its impact on SDMs across different taxa, it is important
to recognize that data quality can vary among taxa.

Importantly, SDMs for endangered species, which often suffer from data deficit
(Erickson and Smith, 2023; Wisz et al., 2008), became accurate in a much fewer amount
of records by blending *Biome* data (Figure 3). Specifically, a threshold of >0.9 Boyce
index could be reached with only around 300 records when using *Biome* data, whereas
over 6 times of data is required when using Traditional survey data only. This finding
highlights the importance of community-sourced data not only for monitoring the
dynamics of endangered species (Chandler et al., 2017; Zapponi et al., 2017) but also for
modelling purposes. Considering the rapid accumulation of *Biome* data, *Biome* data
would make a significant contribution to the more effective distribution modelling of
endangered species.

Limitations of this study

In assessing data quality, reidentification was impossible for records that did not photograph key traits for species identification. To address this limitation, further app improvements can include allowing users to submit multiple images. Encouraging users to document various body parts of organisms through multiple images would make capturing key identification traits much easier. This will make reidentification easier, and possibly improve automatic species identification accuracy.

Given the absence of a comprehensive, environmentally unbiased occurrence dataset spanning a wide range of taxa, we assessed SDM accuracy not relying on an independent test dataset. In this evaluation, the test data was meticulously crafted to include 25% *Biome* data, serving as an intermediary proportion between *Biome*+Traditional (50%) and Traditional survey data (0%). By leveraging the distinct distribution patterns of *Biome* and Traditional survey data along environmental variables (Figure 2C), the test data would better encapsulate the actual species distribution, compared to datasets composed solely of either *Biome* or Traditional survey data. It is noteworthy that, even when the test data exclusively consisted of Traditional survey data (i.e., unfavourable conditions for *Biome*+Traditional data SDMs), the accuracy of SDMs derived from *Biome*+Traditional and Traditional survey data did not differ (Figure 3–figure supplement 1). This result further supports our conclusions that *Biome* provides valuable data for SDM in terms of the amount and quality, and that blending *Biome* data improves SDM accuracy.

We evaluated SDMs based on spatial transferability using the central Japan region, which encompasses a range of environmental conditions. However, the evaluation results may not necessarily indicate transferability across the entire Japanese archipelago. Instead, in the near future, we anticipate that we can evaluate SDM accuracy using temporal transferability. The rapid accumulation of *Biome* data will allow us to evaluate the temporal transferability using the occurrence dataset from different time periods, and thus enable assessing their performance in much wider regions. In addition, limited data availability for certain taxa hindered the assessment in those taxa (e.g., molluscs, amphibians, reptiles, and mammals), but *Biome* would be a platform to overcome the data limitation for many taxa.

Finally, our SDMs do not directly indicate the species' presence probability. The output from presence-only SDMs usually deviates from the probability of presence when species prevalence (i.e. the proportion of area where the species occupied, requiring presence/absence data throughout the area) is unavailable (Elith et al., 2011; Ward et al., 2009). Due to the unavailability of absence data, SDM outputs in this work are indirect measures of species presence and thus are not directly comparable across different

363 species. Nonetheless, they are comparable within a species, providing useful
364 information for understanding species distributions.

Future directions

By blending data from traditional surveys and communities, we improved the accuracy
of species distribution estimates. This enhanced estimation lays the groundwork for
more precise subsequent analyses. For instance, estimated distributions will be useful
in selecting new protected areas or areas with OECMs (Other Effective area-based
Conservation Measures: allowing a wider range of land use as long as biodiversity and
ecosystem services are sustained/improved). Using estimated distributions of each
species, hotspots of species or evolutionary diverse taxa can be inferred. Such sites will
be good candidates for protected areas (Jones et al., 2016) or OECMs (Shiono et al.,
2021). Further, estimated distributions can be used as input for spatial conservation
prioritisation tools (e.g. Marxan (Ball et al., 2009)).

In our experience, stakeholders—including corporate social responsibility managers
and conservation practitioners—often seek the list of species potentially inhabiting their
locations. Due to the uncertainty of SDMs and their thresholding into presence/absence,
on-site surveys remain essential for assessing biodiversity status. Yet, SDMs can make
such surveys cost-effective by screening important locations for on-site assessment (e.g.,
Locate phase in TNFD framework) and narrowing down the target species for surveying.
Improved estimation through SDMs can mitigate risks associated with their use in
society and enable more informed decision-making for conservation efforts.

The rapid accumulation of data from diverse locations holds the potential to unveil
valuable ecological patterns. The accumulated data enables early detection capabilities
for range expansions of invasive species (Sakai et al., in prep). For instance, *Biome* data
has hinted at potential range expansions in several insect species, including butterflies,
dragonflies, and stink bugs, as well as changes in wintering areas for birds (Biome Inc.,
2023). Given the diverse taxonomic coverage of *Biome* data (Figure 2D), detecting
phenological changes across various taxa may be possible. This, in turn, is useful in
uncovering phenological mismatches exacerbated by climate change, which can
significantly change the dynamics of interacting species (Renner and Zohner, 2018;
Visser and Gienapp, 2019). Moreover, *Biome* data is well-suited for assessing the effects
of urbanisation on ecosystems since it comprehensively spans both urban and natural
habitats (Figure 2C). The benefit of rapidly accumulating data, combined with recent
advancements in machine learning methods, opens up opportunities for conducting
time-series analyses. Community science data has rarely been used for time-series

population analysis due to its notable spatio-temporal bias in sampling efforts (Feldman
et al., 2021; Zhang et al., 2021). However, the two-step machine learning approach, as
demonstrated by Fink and colleagues in estimating bird population trends using eBird
data (Fink et al., 2023), sets a precedent. In the future, *Biome* data may facilitate the
inference of population dynamics for multiple taxa. This will enable various time-series
analyses to unveil ecosystem stability and interaction strength, which holds potential for
forecasting ecosystem dynamics (Laubmeier et al., 2020; Pennekamp et al., 2019; Ushio
et al., 2018).

For financial disclosures, companies will assess how their activities rely on ecosystem
services and their opportunities for protecting/recovering nature (TNFD, 2023). By
incorporating taxon-specific ecosystem services, multifaceted ecosystem services can
be preliminarily screened (Kass et al., 2023). For example, based on estimated
distributions of bumblebees or insectivorous animals, the functioning of pollination
services or pest regulation services might be inferred. Using counts of "likes" or records
from *Biome* data, the charismatic species can be determined. By identifying places with
a high estimated richness of charismatic species, potential areas for ecotourism can be
screened. Because SDMs allow us to simulate the impacts of changes in landuse and
climate (Porfirio et al., 2014; Urban et al., 2016), we will be able to forecast how those
changes may influence local biodiversity and/or ecosystem functioning. Hence,
estimated distributions provide the basis of nature-related financial disclosures.

Our platform facilitates collaboration among diverse stakeholders, including local
communities, landowners, and employees from both private companies and
government agencies. Engaging a broader spectrum of stakeholders is crucial for
effective biodiversity assessment, nature management planning, and nature-related
financial disclosures: this inclusivity allows for the incorporation of traditional
knowledge into planning processes, mitigates conflicts among stakeholders, and
ultimately supports more seamless and informed decision-making (Chan et al., 2021;
Keough and Blahna, 2006; Linsley et al., 2023; Roy et al., 2023; TNFD, 2023). Supporting
natural experiences for a wide range of people is also expected to contribute to
changing people's minds towards nature. Through experiencing nature, people become
familiar with it and subsequently make pro-nature decisions (Soga and Gaston, 2023).
We believe that community science can significantly contribute to KM-GBF and create a
sustainable society by fostering nature-positive awareness in society and providing data
tools that enable effective action.

────────────────────────────────────

Methods

Occurrence record accumulation through mobile app *Biome*

In April 2019, a free smartphone app called *Biome* was launched for the Japanese
markets. The app has been downloaded 839,844 times by September 13, 2023. The app
allows users to collect data on the distribution of plants and animals using their mobile
devices. Users can post photographs of the plants and animals they find, and the app
automatically records the location and timestamp from EXIF data. If the EXIF data is
unavailable, users can manually input the locality and timestamp.

To support species identification, the app provides users with two options. First, the app
provides a list of candidate species based on the image and metadata (e.g., location and
timestamp). *Biome* employs a synergistic approach that integrates image recognition
technology and geospatial data to facilitate species identification. The image recognition
algorithm, constructed upon convolutional neural networks, classifies species at higher
taxonomic levels. Subsequently, these candidates are refined based on their frequency
of recent occurrences in the geographical area. Consequently, as the correctly identified
records accumulate for a given area, species identification AI will improve the accuracy.
Second, users can seek help from other users. If a user selects the "ask Biomers" button,
their occurrence record is added to a waiting list that appears on the home screen.
Other users can suggest possible identifications for the records, as in other records of
which species was already identified.

Users can view and comment on other users' records. However, for conservation
purposes, *Biome* automatically conceals the geolocations of endangered species that
are listed on the Japanese national or prefectural red lists. This feature sets it apart from
iNaturalist, where users must manually choose to hide the location of endangered
species (Koide et al., 2023). The social networking function provides opportunities for
communication among users, including non-experts (Fujiki and Tatsuno, 2021). Users
earn "points" through their contributions, including record submissions and
identification suggestions to other users, and progress to higher levels based on their
total points. The points awarded depend on the rarity, conservation status, and societal
impact of the species submitted, meaning that users earn more points when submitting
records of rare, endangered, or invasive species. The app occasionally offers "Quests"
events that provide users with an opportunity to earn additional points by submitting
records from specific locations or of particular species, crucial for monitoring phenology.
Through the variety of gamification features, we stimulate people to participate in
biological surveys as a fun activity.

We obtained occurrence records submitted to *Biome* by 7 July, 2023. The raw data
collected through *Biome* contains invalid presence records which we defined in the
present study as unclear images, documenting non-wild individuals and
misidentifications, and images including some privacy issues. To improve data quality,
we excluded records deemed to be invalid mainly based on location metadata and
users' reactions to the record is as detailed below. This filtered *Biome* data is used in the
subsequent investigations.

Filtering suspicious occurrence record in *Biome* data

Occurrence records of non-wild individuals were eliminated as much as possible by
using information provided by users and location of records. *Biome* users sometimes
report inappropriate records (e.g., unclear images and images from websites or books),
and we excluded all of those reported records. All private records were excluded
because they can harbour inappropriate and misidentified records not being screened
by other users. We also excluded occurrence records that users had marked as non-
wild individuals: users have an option to label their records as photographing bred or
cultivated individuals, or specimens. Records from cultural centres (i.e. zoos, botanical
gardens, museums, and aquariums) and large pet stores were removed as well. During
the data correction process, we prioritise the suggestions provided by *certified users* (see
below for the definition), regardless of the decisions made by the users who originally
created the record. Furthermore, we exclude records that have not been posted by
certified users or have not received identification suggestions from *certified users*.

Certified users are defined as users who achieved the higher accuracy of species
identification (<15% of public occurrence records were suggested as misidentification by
other users), submitted few inappropriate records (<0.5% of public records), and have
created >20 public records. We also defined *specialist users*, a subset of *certified users*
identified in each taxa (see Figure 2 for the classification), who made a total of >30
records or identification suggestions with high identification accuracy (the fraction of
suggested records is less than the average of *certified users* in the taxa). *Specialist users*
are used in determining pseudo-absence for SDMs.

Assessing the accuracy of records

We investigated the proportion of occurrence records within the *Biome* data that were
suitable for SDMs. Since SDMs are influenced by invalid presence records, we assessed

the quality of *Biome* data based on a total of 1420 records from rare and common
species of seed plants, molluscs, insects (including Arachnid and Insecta), fishes,
mammals, birds, reptiles and amphibians (Figure 4). We defined rare species as those
with fewer than or equal to 10 occurrences in *Biome* data, and common species as
those with the highest 15% of records in each taxonomic category. In each of the seed
plant and insect species which account for the majority of *Biome* data (Figure 2D), we
randomly selected 145 records of each rare and common species. For the other
taxonomic categories, we chose each of the 70 records from rare and common species.

Records were first screened whether they targeted organisms (images with no
organisms were discarded) and contained wild individuals. To assess the accuracy of
species identification, species in the records documented wild individuals were
manually reidentified by experts with taxonomic knowledge (Figure 4). These experts
have professional backgrounds, serving as a technician at a prefectural research
institute (fish), highly-experienced field survey conductors (plants and insects,
respectively), a post-doctoral researchers (amphibians and reptiles, and mammals,
respectively), and a museum curator (mollusks) specialising in the focal taxa. Then, by
comparing species identifications by the experts and on *Biome* data, the results were
classified into two categories: (1) correct based on the image and locality—based on the
image, identification was probably correct, and the image locality matches with
habitat/range of the species; (2) misidentification—records were reidentified by experts
if possible. We also examined if the identification was correct at genus and family levels.

Species distribution models

Occurrence data

To evaluate the impact of *Biome* data on SDM prediction accuracy, we compiled two
datasets: "Traditional survey data" and "*Biome*+Traditional data". The Traditional survey
data comprised records collected through conventional survey techniques (e.g. riverine
census, forest inventory census, and museum specimens) primarily sourced from The
National Census on River and Dam Environments (NCRE) and GBIF. In contrast, the
Biome+Traditional data encompassed records submitted to *Biome* that passed filtering
methods, in addition to the Traditional survey data. To control the relative proportion of
Biome data, we constrained the fraction of *Biome* data within the *Biome*+Traditional data
to 50% for each species. Our preliminary results showed that blending 50-70% of *Biome*
data in training data improved prediction accuracy (Appendix 1).
540
For traditional survey data, we downloaded occurrence records of relevant taxa from

GBIF between April 20th, 2023. To prevent significant differences between the sampling
periods of the GBIF records and environmental data, we used the GBIF sampled after
1970. The clean_coordinates function of the R package 'CoordinateCleaner' was used to
remove records with erroneous coordinates such as records from country capitals and
centroids, and biodiversity institutions. We obtained occurrence data from the large
occurrence datasets such as National Census on River and Dam Environments (NCRE)
and Forest Ecosystem Diversity Basic Survey. For the areas or taxa where occurrences
were scarce, we further compiled the literature with detailed locality information, such
as local species inventories. The amount of occurrence records in the modelled species
and species coverage of each dataset are summarised in Table 2. For the species
analysed (S9 Table), traditional survey data contains a negligible portion of community-
sourced data (5.5%) because GBIF contains community-sourced data from iNaturalist
and eBird.

Predictor variables

Predictors encompass a range of environmental variables recognized to impact species
distribution (Table 3): land use (Newbold et al., 2015), climate (bioclim variables (Booth
et al., 2014)), vegetation (Abe, 2018), lithology (Ott, 2020) and elevational range (Udy et
al., 2021). Additionally, categorical variables representing known biogeographic regions,
reflecting geological history, were included. We applied Blakiston's Line —Tsugaru
straits dividing the northern and main islands of Japan (i.e., Hokkaido and Honshu
islands)— reflecting a significant historical migration barrier for mammals and birds
(Dobson, 1994; Saitoh et al., 2015). Due to the distinct fauna (Wepfer et al., 2016;
Yamasaki, 2017), we also specified oceanic islands (i.e. Ogasawara and Daito isles) which
have never been connected with the Asiatic continents. Continuous environmental
variables were transformed into linear, quadratic and hinge feature classes to illustrate
nonlinear associations between environments and species occurrence (Phillips et al.,
2017). The regularisation multiplier was set at 2.5, falling within the established optimal
range of 1.5 to 4 (Elith et al., 2010; Moreno-Amat et al., 2015).

Pseudo-absence reflecting search effort

We considered sampling efforts when selecting a total of 10,000 pseudo-absence
locations. To accommodate biases in sampling efforts, we assigned picking probabilities
as an increasing function of the amount of occurrence records of all and relevant taxa
at the grid cell (an index of sampling efforts) (Milanesi et al., 2020; Phillips et al., 2009).
That is, grid cells with rich occurrence records of relevant taxa are more likely to be

chosen as pseudo-absences than cells with few records, as detailed below (see also
Figure 5).

To generate pseudo-absence (i.e. background) data, we employed two approaches
considering different sampling efforts. The first approach incorporated all observers
and taxa, while the second approach focused on experts and relevant taxa (Figure4). In
both cases, pseudo-absences were selected from grid cells that lacked any occurrence
records of the species being modelled. However, due to variations in sampling efforts
across locations, it was important to address potential bias. To mitigate this bias, we
adjusted the picking probability based on the number of occurrences of other species in
each grid cell (Milanesi et al., 2020; Phillips et al., 2009).

In the first approach, we assumed that the users of *Biome* submit records of any taxon
without specifically selecting species from particular taxa. The picking probability was
simply determined by the total number of records from all taxa in the *Biome* data in
every grid. In the second approach, we considered the expertise of observers (Milanesi
et al., 2020) and the sampling effort for relevant taxa (Phillips et al., 2009). We also
assumed that Traditional surveys targeted particular taxa. Under this approach, we
selected records from *Biome* data contributed by *specialist users* and all records from
the Traditional survey data. From this subset of data, we calculated the number of
records for the taxa (e.g. seed plant, insect and amphibian) to which the modelled
species belonged. This information was then used to calculate the picking probability for
each grid cell. To account for the variability in record counts among locations, we
applied a logarithmic transformation to the number of records. We also added a value
of 1.2 before taking logarithms to allow for the selection of pseudo-absences with low
probabilities, particularly in locations with only one or no records of other species.
Pseudo-absences were not chosen from the spatial block used as test data, but
otherwise, there were no geographical restrictions on their selection.

Using the described approaches, we obtained a total of 10,000 pseudo-absences for our
analyses. The amount of pseudo-absences follows the default setting of MaxEnt (Elith et
al., 2011). For the models using *Biome*+Traditional dataset (also in *Biome*-blended
dataset in Appendix 1), pseudo-absences were generated by merging each of the 5,000
points identified through the two approaches. Meanwhile, for SDMs using the
Traditional survey data only, we obtained 10,000 pseudo-absences by exclusively using
the second approach without incorporating *Biome* data.

Modelling

We modelled distributions of terrestrial seed plants and animals at a scale of 1 x 1 km
grid cell, based on Traditional survey data and *Biome*+Traditional data. To model species
distributions from presence-only data, several algorithms have been utilised, including
generalised additive models, random forest, and neural networks (Norberg et al., 2019;
Valavi et al., 2022). In our study, we opted for MaxEnt (Phillips and Dudík, 2008) due to
its high estimation accuracy and relatively low computational burden (Valavi et al., 2022).
We perfomed MaxEnt via ENMeval 2.0 package (Kass et al., 2021) on R 4.1.3 (R Core
Team, 2021).

Model evaluation

We evaluated the model by examining spatial transferability because we could not find
occurrence data that are environmentally unbiased and independent from training data.
To minimise spatial autocorrelation between training and test data, we set a spatial
block for splitting data (Araújo et al., 2019; Santini et al., 2021). As the spatial block, we
chose the central Japan region (latitude, 33.7˚–37.7˚ N; longitude, 136.2˚–137.6˚ E: Figure
6) which covers various environments—alpine to coastal lowlands, metropolis to highly
intact areas.

To ensure a fair and balanced assessment of the accuracy of SDMs built from
Traditional survey data (0% *Biome* data) and *Biome*+Traditional data (50% *Biome* data),
we compiled a test dataset that embodies characteristics intermediate between these
two datasets. This composite test dataset encompasses 25% *Biome* data and 75%
Traditional data, effectively bridging the differences between the two original datasets
and providing a comprehensive basis for evaluating SDM accuracy.

Due to the presence of invalid records, *Biome* records were used as test data only when
multiple users recorded the same species within an identical 1km grid cell. Although
Biome data may include invalid records (i.e. non-wild individuals or misidentification), if
multiple users recorded the same species at the same place, any one of the records
from the place is likely to be valid. As we know the fraction of valid records within the
Biome dataset in each taxon (see Results), we can calculate the probability of the true
presence in a given location as follows, by assuming that records made by different
users were independent:

$$= 1 - (1 - \quad)$$

The probability of valid records at a given taxon is shown as   , and the number of
users reported given species at the place is indicated as   . If   exceeds 99%, we
deemed that the species occurred in the location.

To reduce spatial sampling bias, we downsampled a dataset within Traditional survey
data, NCRE with massive records from freshwaters, to match the number of records
from the remaining Traditional survey data. This procedure is applied to all test datasets
in both the main analysis and preliminary analyses documented in Figure 3–figure
supplement 1 and Appendix 1.

Boyce index (BI) was used to measure model performance because it was designed to
evaluate presence-only SDMs (Hirzel et al., 2006). In short, BI measures the correlation
between estimated habitat preference and the frequency of actual presence, and
ranges from -1 to 1. A high BI indicates high SDM accuracy that presence data points
tend to be located in grids with higher habitat suitability values. To reliably calculate BI,
at least 50 occurrences should be needed in test data (Hirzel et al., 2006). Thus, we used
132 species that have more than 50 occurrences in test data for calculating BI
(Supplementary File 3).

Examining influences of blending *Biome* data on SDM accuracy

Given that the accuracy of SDMs is affected by the amount and quality of data (Araújo et
al., 2019; Erickson and Smith, 2023; Stockwell and Peterson, 2002), blending *Biome* data
in SDMs may affect the model performances in two possible ways: by increasing the
overall amount of data, and/or by introducing data with different information than the
original data. We analysed to distinguish between these effects. We prepared two
different datasets: "Traditional survey data" and "*Biome*+Traditional data". Then, we
separately trained SDMs using these two datasets. We further varied the data size by
performing random downsampling, ranging from a minimum of 20 to a maximum of
20,000 records, in order to evaluate its impact on the model. As for the
"*Biome*+Traditional data" category, the proportion of *Biome* data was kept at 50%. For
each condition, we conducted three iterations of training and testing to reduce the
impact of random sampling stochasticity. Because the modelling was performed for
each species, we obtained BI for each species, amount of records, and dataset (i.e., two
datasets consisted of 132 species, each with a maximum of 123 conditions for the
amount of records, and the models were replicated three times, resulting in a total of
12,351 individual model runs).

After obtaining BIs for each run, we evaluated the effects of data type (i.e.,
Biome+Traditional data or Traditional survey data) and species on BI while accounting

for the amount of records. For each species and under each amount of records, the
mean BI was calculated across the three iterations. Given that BI is a correlation
coefficient, we applied the Fisher z-transformation to these BIs to approximate their
distribution as a normal distribution. To the transformed BIs, we fitted a generalised
linear mixed model that accounted for both the fixed and interaction effects of data
type and amount of records. This model accommodated species identity as a random
effect. The model was implemented and tested using R packages lme4 (Bates et al.,
2015) and lmerTest (Kuznetsova et al., 2017), respectively.

——

Acknowledgments

We are grateful to Jamie M. Kass for advice on the construction and evaluation of SDMs.
We thank Midzuho Tatsuno, Kazumichi Morishita, Hironori Tanaka, Kotaro Takai and
Shuhei Tochino for species identification; Akira Sawada and Yuko Maegawa for their
advice on the analysis; Trevor H Booth and Dalelan Anderson for comments on the
manuscript. We appreciate the invaluable contributions of *Biome* users. This research
was partly supported by the collaborative research agreement between MU at Kyoto
University and Biome Inc. from 2020 to 2021.

Conflict of Interest

702 KA and YN are employed by Biome Inc., of which the CEO is SF and the CTO is TG. SF and
703 TG are inventors of the species-identification-AI-algorithm JPN patents 6590417 and US
704 patents 11048969. MU and HN declare that they have no competing interests. All
705 authors will not financially benefit directly from the publication of this paper.

707 ——

References

Abe S. 2018. Habitat classification for 69 near threatened plants based on national vegetation survey data. *Veg Sci* **35**:67–88. doi:https://doi.org/10.15031/vegsci.35.67

Araújo MB, Anderson RP, Márcia Barbosa A, Beale CM, Dormann CF, Early R, Garcia RA, Guisan A, Maiorano L, Naimi B, O'Hara RB, Zimmermann NE, Rahbek C. 2019. Standards for distribution models in biodiversity assessments. *Sci Adv* **5**:eaat4858. doi:10.1126/sciadv.aat4858

Atsumi K, Koizumi I. 2017. Web image search revealed large-scale variations in breeding season and nuptial coloration in a mutually ornamented fish, Tribolodon hakonensis. *Ecol Res* **32**:567–578. doi:10.1007/s11284-017-1466-z

Ball IR, Possingham HP, Watts ME. 2009. Marxan and relatives: Software for spatial conservation prioritizationSpatial Conservation Prioritisation: Quantitative Methods and Computational Tools. Oxford University Press. p. 320.

Bates DM, Maechler M, Bolker B, Walker S. 2015. lme4: linear mixed-effects models using S4 classes. *J Stat Softw* **67**:1–48.

Biome Inc. 2023. The report of Climate Change Biosurvey in 2022.

Booth TH, Nix HA, Busby JR, Hutchinson MF. 2014. bioclim: the first species distribution modelling package, its early applications and relevance to most current MaxEnt studies. *Divers Distrib* **20**:1–9. doi:10.1111/ddi.12144

Bowser A, Hansen D, He Y, Boston C, Reid M, Gunnell L, Preece J. 2013. Using gamification to inspire new citizen science volunteersProceedings of the First International Conference on Gameful Design, Research, and Applications, Gamification '13. New York, NY, USA: Association for Computing Machinery. pp. 18–25. doi:10.1145/2583008.2583011

Box EO. 1981. Macroclimate and Plant Forms, Tasks for Vegetation Science. Dordrecht: Springer Netherlands. doi:10.1007/978-94-009-8680-0

Bury TM, Sujith RI, Pavithran I, Scheffer M, Lenton TM, Anand M, Bauch CT. 2021. Deep learning for early warning signals of tipping points. *Proc Natl Acad Sci U S A* **118**. doi:10.1073/pnas.2106140118

Chan L, Hillel O, Werner P, Holman N, Coetzee I, Galt R, Elmqvist T. 2021. Handbook on the Singapore Index on Cities' Biodiversity (also known as the City Biodiversity Index). Montreal: Secretariat of the Convention on Biological Diversity and Singapore: National Parks Board, Singapore: The Secretariat of the Convention on Biological Diversity.

Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, Legind JK, Masinde S, Miller-Rushing AJ, Newman G, Rosemartin A, Turak E. 2017. Contribution of citizen science towards international biodiversity monitoring. *Biol Conserv*, SI:Measures of biodiversity **213**:280–294. doi:10.1016/j.biocon.2016.09.004

Cornwell WK, Ackerly DD. 2009. Community assembly and shifts in plant trait distributions across an environmental gradient in coastal California. *Ecol Monogr* **79**:109–126. doi:10.1890/07-1134.1

Crall AW, Newman GJ, Stohlgren TJ, Holfelder KA, Graham J, Waller DM. 2011. Assessing citizen science data quality: an invasive species case study. *Conserv Lett* **4**:433–442. doi:10.1111/j.1755-263X.2011.00196.x

Dobson M. 1994. Patterns of distribution in Japanese land mammals. *Mammal Rev* **24**:91–111. doi:10.1111/j.1365-2907.1994.tb00137.x

Elith J, Kearney M, Phillips S. 2010. The art of modelling range-shifting species. *Methods Ecol Evol* **1**:330–342. doi:10.1111/j.2041-210X.2010.00036.x

Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ. 2011. A statistical explanation of MaxEnt for ecologists. *Divers Distrib* **17**:43–57. doi:10.1111/j.1472-4642.2010.00725.x

Erickson KD, Smith AB. 2023. Modeling the rarest of the rare: a comparison between multi-species distribution models, ensembles of small models, and single-species models

at extremely low sample sizes. *Ecography* **2023**:e06500. doi:10.1111/ecog.06500

Feldman MJ, Imbeau L, Marchand P, Mazerolle MJ, Darveau M, Fenton NJ. 2021. Trends
and gaps in the use of citizen science derived data as input for species distribution
models: A quantitative review. *PLOS ONE* **16**:e0234587.
doi:10.1371/journal.pone.0234587

Fink D, Johnston A, Strimas-Mackey M, Auer T, Hochachka WM, Ligocki S, Oldham
Jaromczyk L, Robinson O, Wood C, Kelling S, Rodewald AD. 2023. A Double
machine learning trend model for citizen science data. *Methods Ecol Evol* **14**:2435–
2448. doi:10.1111/2041-210X.14186

Fuccillo Battle K, Duhon A, Vispo CR, Crimmins TM, Rosenstiel TN, Armstrong-Davies LL,
de Rivera CE. 2022. Citizen science across two centuries reveals phenological
change among plant species and functional groups in the Northeastern US. *J Ecol*
110:1757–1774. doi:10.1111/1365-2745.13926

Fujiki S, Tatsuno M. 2021. Practice of citizen science for developing biodiversity monitoring
methods using mobile devices. *Jpn J Ecol* **71**:85–90.
doi:https://doi.org/10.18960/seitai.71.2_85

Gonzalez A, Chase JM, O'Connor MI. 2023. A framework for the detection and attribution of
biodiversity change. *Philos Trans R Soc B Biol Sci* **378**:20220182.
doi:10.1098/rstb.2022.0182

Groom Q, Pernat N, Adriaens T, de Groot M, Jelaska SD, Marčiulynienė D, Martinou AF,
Skuhrovec J, Tricarico E, Wit EC, Roy HE. 2021. Species interactions: next-level
citizen science. *Ecography* **44**:1781–1789. doi:10.1111/ecog.05790

Hart AG, Bosley H, Hooper C, Perry J, Sellors-Moore J, Moore O, Goodenough AE. 2023.
Assessing the accuracy of free automated plant identification applications. *People*
Nat **5**:929–937. doi:10.1002/pan3.10460

Herodotou C, Ismail N, I. Benavides Lahnstein A, Aristeidou M, Young AN, Johnson RF,
Higgins LM, Ghadiri Khanaposhtani M, Robinson LD, Ballard HL. 2023. Young
people in iNaturalist: a blended learning framework for biodiversity monitoring. *Int J*
Sci Educ Part B **0**:1–28. doi:10.1080/21548455.2023.2217472

Hirzel AH, Le Lay G, Helfer V, Randin C, Guisan A. 2006. Evaluating the ability of habitat
suitability models to predict species presences. *Ecol Model*, Predicting Species
Distributions **199**:142–152. doi:10.1016/j.ecolmodel.2006.05.017

Hutchinson EG. 1957. Concluding remarksCold Spring Harbor Symposia on Quantitative
Biology. Cold Spring Harbor Laboratory Press.

IPBES. 2019. Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
Bonn, Germany: IPBES secretariat.

Johnston A, Fink D, Hochachka WM, Kelling S. 2018. Estimates of observer expertise
improve species distributions from citizen science data. *Methods Ecol Evol* **9**:88–97.
doi:10.1111/2041-210X.12838

Jones KR, Watson JEM, Possingham HP, Klein CJ. 2016. Incorporating climate change into
spatial conservation prioritisation: A review. *Biol Conserv* **194**:121–130.
doi:10.1016/j.biocon.2015.12.008

Kaplan Mintz K, Arazy O, Malkinson D. 2023. Multiple forms of engagement and motivation
in ecological citizen science. *Environ Educ Res* **29**:27–44.
doi:10.1080/13504622.2022.2120186

Kass JM, Fukaya K, Thuiller W, Mori AS. 2023. Biodiversity modeling advances will improve
predictions of nature's contributions to people. *Trends Ecol Evol* **0**.
doi:10.1016/j.tree.2023.10.011

Kass JM, Muscarella R, Galante PJ, Bohl CL, Pinilla-Buitrago GE, Boria RA, Soley-Guardia
M, Anderson RP. 2021. ENMeval 2.0: Redesigned for customizable and reproducible
modeling of species' niches and distributions. *Methods Ecol Evol* **12**:1602–1608.
doi:10.1111/2041-210X.13628

Kendal D, Egerer M, Byrne JA, Jones PJ, Marsh P, Threlfall CG, Allegretto G, Kaplan H,
Nguyen HKD, Pearson S, Wright A, Flies EJ. 2020. City-size bias in knowledge on

the effects of urban nature on people and biodiversity. *Environ Res Lett* **15**:124035.
doi:10.1088/1748-9326/abc5e4

Keough HL, Blahna DJ. 2006. Achieving Integrative, Collaborative Ecosystem Management.
Conserv Biol **20**:1373–1382. doi:10.1111/j.1523-1739.2006.00445.x

Kindt R. 2023. TreeGOER: A database with globally observed environmental ranges for
48,129 tree species. *Glob Change Biol*. doi:10.1111/gcb.16914

Klinger YP, Eckstein RL, Kleinebecker T. 2023. iPhenology: Using open-access citizen
science photos to track phenology at continental scale. *Methods Ecol Evol* **14**:1424–
1431. doi:10.1111/2041-210X.14114

Kobori H, Dickinson JL, Washitani I, Sakurai R, Amano T, Komatsu N, Kitamura W,
Takagawa S, Koyama K, Ogawara T, Miller-Rushing AJ. 2016. Citizen science: a
new approach to advance ecology, education, and conservation. *Ecol Res* **31**:1–19.
doi:10.1007/s11284-015-1314-y

Koide D, Tsujimoto S, Kumagai N, Ikegami M, Nishihiro J. 2023. Species' spatiotemporal
distribution platform based on citizen science through desirable circulation between
the real and digital worlds. *Jpn J Conserv Ecol*.
doi:https://doi.org/10.18960/hozen.2217

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest package: Tests in linear
mixed effects models. *J Stat Softw* **82**. doi:10.18637/jss.v082.i13

Larson ER, Graham BM, Achury R, Coon JJ, Daniels MK, Gambrell DK, Jonasen KL, King
GD, LaRacuente N, Perrin-Stowe TI, Reed EM, Rice CJ, Ruzi SA, Thairu MW,
Wilson JC, Suarez AV. 2020. From eDNA to citizen science: emerging tools for the
early detection of invasive species. *Front Ecol Environ* **18**:194–202.
doi:10.1002/fee.2162

Laubmeier AN, Cazelles B, Cuddington K, Erickson KD, Fortin MJ, Ogle K, Wikle CK, Zhu K,
Zipkin EF. 2020. Ecological Dynamics: Integrating Empirical, Statistical, and
Analytical Methods. *Trends Ecol Evol* **35**:1090–1099. doi:10.1016/j.tree.2020.08.006

Leighton GRM, Hugo PS, Roulin A, Amar A. 2016. Just Google it: assessing the use of
Google Images to describe geographical variation in visible traits of organisms.
Methods Ecol Evol **7**:1060–1070. doi:10.1111/2041-210X.12562

Linsley P, Abdelbadie Roba, Abdelbadie Rasha. 2023. The Taskforce on Nature-related
Financial Disclosures must engage widely and justify its market-led approach. *Nat*
Ecol Evol **7**:1343–1346. doi:10.1038/s41559-023-02113-w

Loh J, Green RE, Ricketts T, Lamoreux J, Jenkins M, Kapos V, Randers J. 2005. The Living
Planet Index: using species population time series to track trends in biodiversity.
Philos Trans R Soc B Biol Sci **360**:289–295. doi:10.1098/rstb.2004.1584

Milanesi P, Mori E, Menchetti M. 2020. Observer-oriented approach improves species
distribution models from citizen science data. *Ecol Evol* **10**:12104–12114.
doi:10.1002/ece3.6832

Miller DAW, Pacifici K, Sanderlin JS, Reich BJ. 2019. The recent past and promising future
for data integration methods to estimate species' distributions. *Methods Ecol Evol*
10:22–37. doi:10.1111/2041-210X.13110

Miya M, Sado T, Oka S, Fukuchi T. 2022. The use of citizen science in fish eDNA
metabarcoding for evaluating regional biodiversity in a coastal marine region: A pilot
study. *Metabarcoding Metagenomics* **6**:e80444. doi:10.3897/mbmg.6.80444

Moreno-Amat E, Mateo RG, Nieto-Lugilde D, Morueta-Holme N, Svenning J-C, García-
Amorena I. 2015. Impact of model complexity on cross-temporal transferability in
Maxent species distribution models: An assessment using paleobotanical data. *Ecol*
Model **312**:308–317. doi:10.1016/j.ecolmodel.2015.05.035

Mori AS, Suzuki KF, Hori M, Kadoya T, Okano K, Uraguchi A, Muraoka H, Sato T, Shibata
H, Suzuki-Ohno Y, Koba K, Toda M, Nakano S, Kondoh M, Kitajima K, Nakamura M.
2023. Perspective: sustainability challenges, opportunities and solutions for long-term
ecosystem observations. *Philos Trans R Soc B Biol Sci* **378**:20220192.
doi:10.1098/rstb.2022.0192

Newbold T, Hudson LN, Arnell AP, Contu S, De Palma A, Ferrier S, Hill SLL, Hoskins AJ,

Lysenko I, Phillips HRP, Burton VJ, Chng CWT, Emerson S, Gao D, Hale GP, Hutton
J, Jung M, Sanchez-Ortiz K, Simmons BI, Whitmee S, Zhang H, Scharlemann JPW,
Purvis A. 2016. Has land use pushed terrestrial biodiversity beyond the planetary
boundary? A global assessment. *Science* **353**:291–288.
doi:10.1126/science.aaf2201
Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, Börger L, Bennett DJ,
Choimes A, Collen B, Day J, De Palma A, Díaz S, Echeverria-Londoño S, Edgar MJ,
Feldman A, Garon M, Harrison MLK, Alhusseini T, Ingram DJ, Itescu Y, Kattge J,
Kemp V, Kirkpatrick L, Kleyer M, Correia DLP, Martin CD, Meiri S, Novosolov M, Pan
Y, Phillips HRP, Purves DW, Robinson A, Simpson J, Tuck SL, Weiher E, White HJ,
Ewers RM, Mace GM, Scharlemann JPW, Purvis A. 2015. Global effects of land use
on local terrestrial biodiversity. *Nature* **520**:45–50. doi:10.1038/nature14324
Norberg A, Abrego N, Blanchet FG, Adler FR, Anderson BJ, Anttila J, Araújo MB, Dallas T,
Dunson D, Elith J, Foster SD, Fox R, Franklin J, Godsoe W, Guisan A, O'Hara B, Hill
NA, Holt RD, Hui FKC, Husby M, Kålås JA, Lehikoinen A, Luoto M, Mod HK, Newell
G, Renner I, Roslin T, Soininen J, Thuiller W, Vanhatalo J, Warton D, White M,
Zimmermann NE, Gravel D, Ovaskainen O. 2019. A comprehensive evaluation of
predictive performance of 33 species distribution models at species and community
levels. *Ecol Monogr* **89**:e01370. doi:10.1002/ecm.1370
Ott RF. 2020. How Lithology Impacts Global Topography, Vegetation, and Animal
Biodiversity: A Global-Scale Analysis of Mountainous Regions. *Geophys Res Lett*
47:e2020GL088649. doi:10.1029/2020GL088649
Pacifici K, Reich BJ, Miller DAW, Gardner B, Stauffer G, Singh S, McKerrow A, Collazo JA.
2017. Integrating multiple data sources in species distribution modeling: a framework
for data fusion*. *Ecology* **98**:840–850. doi:10.1002/ecy.1710
Pennekamp F, Iles AC, Garland J, Brennan G, Brose U, Gaedke U, Jacob U, Kratina P,
Matthews B, Munch S, Novak M, Palamara GM, Rall BC, Rosenbaum B, Tabi A,
Ward C, Williams R, Ye H, Petchey OL. 2019. The intrinsic predictability of ecological
time series and its potential to guide forecasting. *Ecol Monogr* **89**:e01359.
doi:10.1002/ecm.1359
Phillips SJ, Anderson RP, Dudík M, Schapire RE, Blair ME. 2017. Opening the black box: an
open-source release of Maxent. *Ecography* **40**:887–893. doi:10.1111/ecog.03049
Phillips SJ, Anderson RP, Schapire RE. 2006. Maximum entropy modeling of species
geographic distributions. *Ecol Model* **190**:231–259.
doi:10.1016/j.ecolmodel.2005.03.026
Phillips SJ, Dudík M. 2008. Modeling of species distributions with Maxent: new extensions
and a comprehensive evaluation. *Ecography* **31**:161–175. doi:10.1111/j.0906-
7590.2008.5203.x
Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S. 2009. Sample
selection bias and presence-only distribution models: implications for background
and pseudo-absence data. *Ecol Appl* **19**:181–197. doi:10.1890/07-2153.1
Pocock MJO, Chandler M, Bonney R, Thornhill I, Albin A, August T, Bachman S, Brown
PMJ, Cunha DGF, Grez A, Jackson C, Peters M, Rabarijaon NR, Roy HE, Zaviezo T,
Danielsen F. 2018. Chapter Six - A Vision for Global Biodiversity Monitoring With
Citizen Science In: Bohan DA, Dumbrell AJ, Woodward G, Jackson M, editors.
Advances in Ecological Research, Next Generation Biomonitoring: Part 2. Academic
Press. pp. 169–223. doi:10.1016/bs.aecr.2018.06.003
Pocock MJO, Tweddle JC, Savage J, Robinson LD, Roy HE. 2017. The diversity and
evolution of ecological and environmental citizen science. *PLoS ONE* **12**:1–17.
doi:10.1371/journal.pone.0172579
Ponti M, Hillman T, Stankovic I. 2015. Science and Gamification: The Odd
Couple?Proceedings of the 2015 Annual Symposium on Computer-Human
Interaction in Play, CHI PLAY '15. New York, NY, USA: Association for Computing
Machinery. pp. 679–684. doi:10.1145/2793107.2810293
Porfirio LL, Harris RMB, Lefroy EC, Hugh S, Gould SF, Lee G, Bindoff NL, Mackey B. 2014.

Improving the Use of Species Distribution Models in Conservation Planning and
Management under Climate Change. *PLOS ONE* **9**:e113749.
doi:10.1371/journal.pone.0113749
R Core Team. 2021. R: A language and environment for statistical computing. *R Found Stat*
Comput Vienna Austria.
Reddy S, Dávalos LM. 2003. Geographical sampling bias and its implications for
conservation priorities in Africa. *J Biogeogr* **30**:1719–1727. doi:10.1046/j.1365-
2699.2003.00946.x
Renner SS, Zohner CM. 2018. Climate Change and Phenological Mismatch in Trophic
Interactions Among Plants, Insects, and Vertebrates. *Annu Rev Ecol Evol Syst*
49:165–182. doi:10.1146/annurev-ecolsys-110617-062535
Robinson OJ, Ruiz-Gutierrez Viviana, Reynolds MD, Golet GH, Strimas-Mackey M, Fink D.
2020. Integrating citizen science data with expert surveys increases accuracy and
spatial extent of species distribution models. *Divers Distrib* **26**:976–986.
doi:10.1111/ddi.13068
Roy HE, Pauchard A, Stoett P, Renard Truong T, Bacher S, Galil BS, Hulme PE, Ikeda T,
Sankaran KV, McGeoch MA, Meyerson LA, Nuñez MA, Ordonez A, Rahlao SJ,
Schwindt E, Seebens H, Sheppard AW, Vandvik V. 2023. IPBES Invasive Alien
Species Assessment: Summary for Policymakers. Zenodo.
doi:10.5281/zenodo.8314303
Saitoh T, Sugita N, Someya S, Iwami Y, Kobayashi S, Kamigaichi H, Higuchi A, Asai S,
Yamamoto Y, Nishiumi I. 2015. DNA barcoding reveals 24 distinct lineages as cryptic
bird species candidates in and around the Japanese Archipelago. *Mol Ecol Resour*
15:177–186. doi:10.1111/1755-0998.12282
Santini L, Benítez-López A, Maiorano L, Čengić M, Huijbregts MAJ. 2021. Assessing the
reliability of species distribution projections in climate change research. *Divers Distrib*
27:1035–1050. doi:10.1111/ddi.13252
Schmitt S, Pouteau R, Justeau D, de Boissieu F, Birnbaum P. 2017. ssdm: An r package to
predict distribution of species richness and composition based on stacked species
distribution models. *Methods Ecol Evol* **8**:1795–1803. doi:10.1111/2041-210X.12841
Scholes RJ, Biggs R. 2005. A biodiversity intactness index. *Nature* **434**:45–49.
doi:10.1038/nature03289
Shiono T, Kubota Y, Kusumoto B. 2021. Area-based conservation planning in Japan: The
importance of OECMs in the post-2020 Global Biodiversity Framework. *Glob Ecol*
Conserv **30**:e01783. doi:10.1016/j.gecco.2021.e01783
Soga M, Gaston KJ. 2023. Nature benefit hypothesis: Direct experiences of nature predict
self-reported pro-biodiversity behaviors. *Conserv Lett* **n/a**:e12945.
doi:10.1111/conl.12945
Steen VA, Elphick CS, Tingley MW. 2019. An evaluation of stringent filtering to improve
species distribution models from citizen science data. *Divers Distrib* **25**:1857–1869.
doi:10.1111/ddi.12985
Stockwell DRB, Peterson AT. 2002. Effects of sample size on accuracy of species
distribution models. *Ecol Model* **148**:1–13. doi:10.1016/S0304-3800(01)00388-X
Tilman D, Reich PB, Knops JMH. 2006. Biodiversity and ecosystem stability in a decade-
long grassland experiment. *Nature* **441**:629–632. doi:10.1038/nature04742
TNFD. 2023. Taskforce on Nature-related Financial Disclosures (TNFD) Recommendations
version 1.0.
Udy K, Fritsch M, Meyer KM, Grass I, Hanß S, Hartig F, Kneib T, Kreft H, Kukunda CB,
Pe'er G, Reininghaus H, Tietjen B, Tscharntke T, van Waveren C-S, Wiegand K.
2021. Environmental heterogeneity predicts global species richness patterns better
than area. *Glob Ecol Biogeogr* **30**:842–851. doi:10.1111/geb.13261
Urban MC, Bocedi G, Hendry AP, Mihoub J-B, Pe'er G, Singer A, Bridle JR, Crozier LG, De
Meester L, Godsoe W, Gonzalez A, Hellmann JJ, Holt RD, Huth A, Johst K, Krug CB,
Leadley PW, Palmer SCF, Pantel JH, Schmitz A, Zollner PA, Travis JMJ. 2016.
Improving the forecast for biodiversity under climate change. *Science* **353**:aad8466.

doi:10.1126/science.aad8466

Ushio M, Hsieh CH, Masuda R, Deyle ER, Ye H, Chang CW, Sugihara G, Kondoh M. 2018.
Fluctuating interaction network and time-varying stability of a natural fish community.
Nature **554**:360–363. doi:10.1038/nature25504

Valavi R, Guillera-Arroita G, Lahoz-Monfort JJ, Elith J. 2022. Predictive performance of
presence-only species distribution models: a benchmark study with reproducible
code. *Ecol Monogr* **92**:e01486. doi:10.1002/ecm.1486

Visser ME, Gienapp P. 2019. Evolutionary and demographic consequences of phenological
mismatches. *Nat Ecol Evol* **12**. doi:10.1038/s41559-019-0880-8

Wallace RD, Bargeron CT. 2014. Identifying invasive species in real time: early detection
and distribution mapping system (EDDMapS) and other mapping tools. *Invasive*
Species Glob Clim Change, CABI Invasives Series 219–231.
doi:10.1079/9781780641645.0219

Ward G, Hastie T, Barry S, Elith J, Leathwick JR. 2009. Presence-Only Data and the EM
Algorithm. *Biometrics* **65**:554–563. doi:10.1111/j.1541-0420.2008.01116.x

Wepfer PH, Guénard B, Economo EP. 2016. Influences of climate and historical land
connectivity on ant beta diversity in East Asia. *J Biogeogr* **43**:2311–2321.
doi:10.1111/jbi.12762

Wisz MS, Hijmans RJ, Li J, Peterson AT, Graham CH, Guisan A, Group NPSDW. 2008.
Effects of sample size on the performance of species distribution models. *Divers*
Distrib **14**:763–773. doi:10.1111/j.1472-4642.2008.00482.x

Wood C, Sullivan B, Iliff M, Fink D, Kelling S. 2011. eBird: Engaging Birders in Science and
Conservation. *PLOS Biol* **9**:e1001220. doi:10.1371/journal.pbio.1001220

Yamasaki T. 2017. Biogeographic Pattern of Japanese Birds: A Cluster Analysis of Faunal
Similarity and a Review of Phylogenetic Evidence In: Motokawa M, Kajihara H,
editors. Species Diversity of Animals in Japan, Diversity and Commonality in
Animals. Tokyo: Springer Japan. pp. 117–134. doi:10.1007/978-4-431-56432-4_4

Zapponi L, Cini A, Bardiani M, Hardersen S, Maura M, Maurizi E, Redolfi De Zan L, Audisio
P, Bologna MA, Carpaneto GM, Roversi PF, Sabbatini Peverieri G, Mason F,
Campanaro A. 2017. Citizen science data as an efficient tool for mapping protected
saproxylic beetles. *Biol Conserv*, The role of citizen science in biological conservation
208:139–145. doi:10.1016/j.biocon.2016.04.035

Zhang W, Sheldon BC, Grenyer R, Gaston KJ. 2021. Habitat change and biased sampling
influence estimation of diversity trends. *Curr Biol* **31**:3656-3662.e3.
doi:10.1016/j.cub.2021.05.066

1018

1019

Figure, tables and supporting information

1021

Figure 1. Workflow of submitting records to *Biome*. (1) Users can upload images that were taken by the smartphone camera or import existing images from the storage, including those imported from external devices. (2) Users select whether the image is about animals or plants to activate the species identification AI. (3) The AI analyses the image and its metadata to generate a candidate species list. (4) Alternatively, users can input the taxon name manually and obtain a list of candidate species. To submit the occurrence record, users can either (5) seek identification assistance from other users through the "ask Biomers" feature, or (6) identify the species from the list. To the records, users can add memos and tags indicating phenology, life-stage, sex, and whether the individual is wild or captive.

1031

Figure 2. Description of data accumulated by *Biome*. Data distributions are shown based on
all records submitted to *Biome* by 7 July 2023 (*N*=5,275,457). **A** Spatial distribution of records
across Japan. **B** Accumulation of records through time. The barplot represents the number of
records each month and the line shows the cumulative amount of records. **C** Distributions of
records along with PC1 of all environmental variables and standardised area occupancy of
urban-type land uses. Grey and green represent distributions of Traditional and *Biome* data,
respectively. **D** Taxonomic composition of records is shown as the area sizes. 'Other plant'
consists of non-seed terrestrial plants; 'insects' include Arachnids and Insects; 'arthropods' cover
any Arthropod not included in insects; 'other animals' covers all invertebrates not included in the
taxa above.

1042

Figure 3. The accuracy of species distribution models. Accuracy of SDMs using Traditional
survey data (grey dots and lines) and *Biome*+Traditional data (i.e. 50% of *Biome* data: green). Each
SDM was performed with a specific dataset, species, and the amount of records. For each species
and amount of records, we computed the average model accuracy (Boyce index) from three
replicated runs. Subsequently, we calculated the median model accuracy across species for each
amount of records. These medians were then illustrated for each taxon in the strip of each respective
panel. The "Endangered" category includes species that are listed as endangered on Japan's national
or prefectural red lists.

1051

1052 **Figure 4. The workflow of checking accuracy of *Biome* data.**

1053

1054 **Figure 5. The workflow for selecting pseudo-absence (background) grid cells for SDMs**
1055 **using the *Biome*-Traditional dataset.** In this process, both *Biome* data and Traditional dataset
1056 are utilised to determine the suitable locations for pseudo-absence grid cells. However, when
1057 constructing SDMs using the Traditional dataset exclusively, *Biome* data is not involved in the
1058 selection of pseudo-absence points.

1059

1060 **Figure 6. Japanese archipelago, coloured by altitude.** Shaded area shows spatial block of test
1061 data. Retrieved from Wikipedia (2023, May 30), licensed under Creative Commons Attribution-
1062 ShareAlike 3.0 Unported (CC BY-SA 3.0).

1063

1064

1065 **Table 1. Data quality of *Biome*.** The fraction of records documenting wild individuals, and
1066 identification accuracy at species, genus and family levels among the records documenting wild
1067 individuals are shown. Species were identified only for records documenting wild individuals.

Species group	Species rarity	N	Wild / total(%)	Species correct / wild (%)	Genus correct / wild (%)	Family correct / wild (%)
TOTAL	TOTAL	1420	81.6	91	93.6	96.9
Seed plant	TOTAL	290	86.2	89.6	94.4	97.2
Mollusca	TOTAL	140	87.9	90.2	91.1	96.7
Insect	TOTAL	290	100	83.4	86.9	94.1
Fish	TOTAL	140	73.6	87.4	93.2	96.1
Amphibian	TOTAL	140	93.6	96.2	96.2	98.5
Reptile	TOTAL	140	91.4	97.7	100	100
Bird	TOTAL	140	98.6	98.6	99.3	99.3
Mammal	TOTAL	140	80.7	95.6	95.6	96.5
TOTAL	Rare	710	88.7	87	91	95.6
TOTAL	Common	710	91	95	96.3	98.3
Seed plant	Rare	145	80.7	82.9	91.5	94.9
Seed plant	Common	145	91.7	95.5	97	99.2
Mollusca	Rare	70	82.9	86.2	87.9	96.6
Mollusca	Common	70	92.9	93.8	93.8	96.9
Insect	Rare	145	100	75.2	80	91.7
Insect	Common	145	100	91.7	93.8	96.6
Fish	Rare	70	74.3	88.5	94.2	94.2

Fish	**Common**	70	72.9	86.3	92.2	98
Amphibian	**Rare**	70	95.7	95.5	95.5	98.5
Amphibian	**Common**	70	91.4	96.9	96.9	98.4
Reptile	**Rare**	70	94.3	95.5	100	100
Reptile	**Common**	70	88.6	100	100	100
Bird	**Rare**	70	97.1	98.5	100	100
Bird	**Common**	70	100	98.6	98.6	98.6
Mammal	**Rare**	70	81.4	91.2	91.2	93
Mammal	**Common**	70	80	100	100	100

1068

1069 **Table 2. List of species occurrence datasets used for constructing SDMs.** To compare *Biome*
1070 dataset with the other datasets, iNaturalist and eBird data based on community science were
1071 classified as 'Traditional survey' data. *For the list of GBIF download doi and literatures, see
1072 Supplementary File 2.

Original dataset	Occurrence records of modelled species		Species coverage among modelled species	Survey method	Data group in SDM	Down-load date	Availability
	N	Occu-pancy					
Biome (filtering applied)	201114	8.6	132/132	Citizen science through smartphone app	*Biome*	7 July 2023	https://biome.co.jp/
National Census on River and Dam Environments (NCRE)	1413541	60.2	126/132	Traditional survey on freshwater and its adjacent ecosystems	Traditional survey	10 Jan. 2023	http://www.nilim.go.jp/lab/fbg/ksnkankyo/

Institute records registered at GBIF	530952	22.6	116/132	Traditional survey and museum specimens	Traditional survey	7 July 2023	GBIF*
iNaturalist and eBird	118050	5	110/132	Citizen science through smartphone app and web service	Traditional survey*	7 July 2023	GBIF*
Forest Ecosystem Diversity Basic Survey	80929	3.4	42/132	Traditional survey on forest trees	Traditional survey	30 March 2023	http://forestbio.jp/
Literatures	3293	0.1	130/132	Traditional survey	Traditional survey	31 March 2023	Refs*

1073

1074 **Table 3. Environmental data used for constructing SDMs.** Years indicate the data collection
1075 period. Usage in the SDM shows how the variables were converted before using in the species
1076 distribution modelling.

Data	Variables	Year	Usage in the SDM	Available at
Land use	The area sizes of forests, rice fields, farms, wastelands, inland waters, beaches, ocean, golf courses, urbanised areas, and others	2016	Extracted six principal components (PCA) explained ≧ 80% of total variation. PCs were converted into linear, quadratic and hinge terms	The Ministry of Land, Infrastructure, Transport and Tourism of Japan (MLIT) https://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-L03-a.html
Forest type	Forest type (planted and natural)	1994-1998	Converted into linear, quadratic and hinge terms	The Biodiversity Centre of Japan; http://gis.biodic.go.jp/webgis/index.html
Climate	Monthly average, minimum and maximum temperature and precipitation	1981-2010	Transformed into 19 bioclimatic variables (Booth et al., 2014), then extracted three PCs explained ≧ 80% of	MLIT, https://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-G02-v3_0.html

			total variation. Converted into linear, quadratic and hinge terms	
Elevation-al range	Differences between maximum and minimum elevation, and maximum slope	1981	Converted into linear, quadratic and hinge terms	MLIT, https://nlftp.mlit.go.jp/ksj/jpgis/datalist/KsjTmplt-G04-a.html
Vege-tation	The area sizes	1998	Transformed into 37 PCs of which total variation explained was more than 80%. Converted into linear, quadratic and hinge terms	MOE, http://gis.biodic.go.jp/webgis/index.html
Geology	The area sizes of limestone and serpentinite	2022	Converted into linear, quadratic and hinge terms	The Research Institute of Geology and Geoinformation, https://gbank.gsj.jp/seamless/use.html
Geo-history	Blakiston's Line (Dobson, 1994; Saitoh et al., 2015), oceanic islands (Wepfer et al., 2016; Yamasaki, 2017)		Categorical variables	

1077

1078

Figure 3–figure supplement 1. Accuracy of SDMs using Traditional survey data
(grey dots and lines) and *Biome*+Traditional data (i.e. 50% of *Biome* data: green),
evaluated against test data only consisting of Traditional survey data.

Supplementary File 1. Distributions of occurrence records along with
environmental variables.

Supplementary File 2. List of GBIF data doi and literatures compiled in occurrence

data.

Supplementary File 3. List of species for constructed species distribution models.
1087
1088
1089

1090

1091

Appendix 1. Determine the best blend of Traditional survey and *Biome* data

Methods

In this investigation, we aimed to determine the optimal proportion of *Biome* data within the training dataset of SDMs in order to enhance the accuracy of SDMs. To conduct this assessment, we initially selected a subset of species for which sufficient test data was available (as detailed below).

For each of the selected species, we generated training datasets by combining Traditional survey data with *Biome* data at different proportions: 15%, 30%, 40%, 50%, 60%, 70%, 85%, and 100%, referred to as *Biome*-blended datasets.

To compare the accuracy of SDMs, we created and evaluated models using both the Traditional survey dataset and each of the *Biome*-blended datasets. SDMs were created by following the methodology employed in the main analysis. To ensure equitable comparison, we equalised the amount of data in each pair of blended and Traditional survey datasets. This equalisation was achieved by randomly downsampling the larger dataset to match the size of the smaller one.

We assessed the accuracy of the models using the Boyce index (BI), which follows the same methodology as employed in the main analysis. In this specific investigation, we did not control the proportion of *Biome* data within the test data. We selected a set of species for which the test dataset contained at least 50 locations and randomly chose 20 species from each of the seed plants, insects, and birds (see Supplementary File 4).

Results

The analysis revealed that the relative model accuracy becomes high positive values when training data comprises 50-70% of *Biome* data (Appendix 1–figure 1). This indicates that SDM accuracy is substantially enhanced when the training data incorporates 50-70% of *Biome* data. The relative model accuracy remained positive in the 15-70% *Biome*-blended datasets, but decreased to negative values in the 85% and 100% *Biome*-blended datasets (S3 Fig). This suggests that blending *Biome* data generally enhances the accuracy of SDMs, but it is important to include at least 30% Traditional survey data to maintain accuracy. Based on the high performance observed and simplicity, we selected the 50% *Biome*-blended dataset as the *Biome*+Traditional data for comparing model accuracy with the Traditional survey data in the main text.

1124

1125   **Appendix 1–figure 1. The violin plots of relative model accuracy between SDMs using**
1126   ***Biome*-blended data and Traditional survey data.** The median values are shown as grey dots.
1127   The positive relative model accuracy indicates that SDMs that used *Biome* data outperformed
1128   models that used Traditional survey data.
1129

A Spatial distribution

Density of records (/10km)

8000
400
20
0

B Temporal accumulation

Occurrence records in each month

Cumulative amount of occurrence records

300000
6000000

200000
4000000

100000
2000000

0
0

2019-07 2019-07 2020-07 2021-01 2021-07 2022-01 2022-07 2023-01

C Environmental conditions

Occurrence records

All variables

4000000
3000000
2000000
1000000

Traditional survey data

Biome data

-5   0   5   10
PC1 value

10000000
9500000

2000000
1500000
1000000
500000

Urbanized area occupancy

0   1   2   3   4   5   6
Standardized value

D Taxonomic composition

Insect

Seed plant

Fish

Bird

Other plant
Mollusca
Crustacea
Amphibian
Reptile
Mammal
Other animal

Amount of occurrence records

Species groups examined

Seed plants | Insect | Mollusca | Fish

Amphibian | Reptile | Bird | Mammal

Common species
Species with
highest 15% of records
in the *species group*

Rare species
Species with
≤10 records

Randomly chose records from common/rare cateogories
in each *species group*:
each of 145 records in seed plants and insects,
each of 70 records in the rest of *species groups* were chosen

Data check

Does the record document wild individuals?
Clear and original images
photographed wild individuals

Yes

No

Identification
by experts

Finish

Strictly correct
based on
a single image

Correct based on
an image
and locality

Misidentification

Valid records