# Boosting biodiversity monitoring using smartphone-driven, rapidly accumulating community-sourced data

Keisuke Atsumi,[1] Yuusuke Nishida[1], Masayuki Ushio[2,3,4], Hirotaka Nishi[5],
Takanori Genroku[1], Shogoro Fujiki[1*]

[1]*Biome Inc., 134 Chudoji Minamimachi, Shimogyo-ku, Kyoto 600-8813, Japan*

[2]*Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR*

[3]*Hakubi Center, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto 606-8501, Japan*

[4]*Center for Ecological Research, Kyoto University, 2-509-3 Hirano, Otsu, Shiga 520-2113, Japan*

[5]*Toyohashi Museum of Natural History, 1-238 Oana, Oiwa, Toyohashi, Aichi 441-3147, Japan*

*Correspondence: fujiki@biome.co.jp

**Abstract**

Ecosystem services, which derive in part from biological diversity, are a fundamental support for human society. However, human activities are causing harm to biodiversity, ultimately endangering these critical ecosystem services. Halting nature loss and mitigating these impacts necessitates comprehensive biodiversity distribution data, a requirement for implementing the Kunming-Montreal Global Biodiversity Framework. To efficiently collect species observations from the public, we launched the '*Biome*' mobile application in Japan. By employing species identification algorithms and gamification elements, the app has gathered >6M observations since its launch in 2019. However, community-sourced data often exhibit spatial and taxonomic biases. Species distribution models (SDMs) enable inferring species distribution while accommodating such bias. We investigated *Biome* data's quality and how incorporating the data influences the performance of SDMs. Species identification accuracy exceeds 95% for birds, reptiles, mammals, and amphibians, but seed plants, molluscs, and fishes scored below 90%. The distributions of 132 terrestrial plants and animals across Japan were modelled, and their accuracy was improved by incorporating our data into traditional survey data. For endangered species, traditional survey data required >2,000 records to build accurate models (Boyce index ≥ 0.9), though only ca.300 records were required when the two data sources were blended. The unique data distributions may explain this improvement: *Biome* data covers urban-natural gradients uniformly, while traditional data is biassed

**towards natural areas. Combining multiple data sources offers insights into species**
**distributions across Japan, aiding protected area designation and ecosystem service**
**assessment. Providing a platform to accumulate community-sourced distribution data**
**and improving data processing protocol will contribute to not only conserving natural**
**ecosystems but also detecting species distribution changes and testing ecological**
**theories.**

**KEY WORDS:**

**Community science, Citizen science, Habitat suitability model, MaxEnt, Big data,**
**Taskforce on Nature-related Financial Disclosures: TNFD, Other Effective area-based**
**Conservation Measures: OECMs, Natural capital**

―――――――――――――――――――――――――――――――――――――――

# Introduction

Nature underpins human society, and the conservation of ecosystems and associated
ecosystem services contributes to the sustainable development of human society, yet
these services have been rapidly declining in recent years (IPBES, 2019; Loh et al., 2005;
Newbold et al., 2016; Scholes and Biggs, 2005). Kunming-Montreal Global Biodiversity
Framework (KM-GBF) by the United Nations envisions reversing the nature loss by 2030.
As direct means for nature conservation, KM-GBF targeted making 30% of Earth's land
and ocean area as protected areas by 2030 (i.e. 30by30). As an indirect but influential way,
KM-GBF requires companies to "*monitor, assess, and transparently disclose their risks,*
*dependencies and impacts on biodiversity through their operations, supply and value chains*
*and portfolios,*" which is guided by Taskforce on Nature-related Financial Disclosures
(TNFD) (TNFD, 2023). To achieve these goals, it is imperative to assess the state of
biodiversity with a sufficient spatiotemporal resolution to support conservation planning,
adaptive management, and companies' annual nature-related financial disclosures. The
basis for such assessments lies in our knowledge of species distributions (Gonzalez et al.,
2023; Newbold et al., 2016). Traditionally, distribution data was acquired through on-site
surveys by experts (people have expertise about biodiversity), but collecting distribution
data with sufficient spatiotemporal resolution is challenging if we rely only on such limited
human resources (Miya et al., 2022; Mori et al., 2023; Pocock et al., 2018).

Since the emergence of digital devices and the internet, people have been able to share
their observations through various media, such as images and video/audio recordings.
Such community-sourced data have significantly contributed to the accumulation of
ecosystem information. These datasets have been instrumental in assessing the impacts
of climate change and urbanisation on phenology (Fuccillo Battle et al., 2022; Klinger et
al., 2023), detecting distribution changes including invasive alien species (Larson et al.,
2020; Roy et al., 2023; Wallace and Bargeron, 2014), exploring large-scale geographic
variations in traits (Atsumi and Koizumi, 2017; Leighton et al., 2016), and estimating
species distributions (Chandler et al., 2017; Feldman et al., 2021; Johnston et al., 2018;
Steen et al., 2019). Moreover, the utilisation of machine learning to describe population
trends based on community-sourced data (Fink et al., 2023) offers opportunities for
conducting time-series analyses. These analyses can help us understand community
assembly processes, unravel species interaction networks, and assess ecosystem stability
(Cornwell and Ackerly, 2009; Tilman et al., 2006; Ushio et al., 2018), capitalising on the
spatio-temporally dense sampling effort facilitated by community-sourced data
(Chandler et al., 2017; Kobori et al., 2016; Pocock et al., 2017). Such analytical approaches
enable us to make informed predictions about changes in species distribution,
population dynamics, and ecosystem stability in the face of climate change (Bury et al.,
2021; Pennekamp et al., 2019; Urban et al., 2016). In essence, community-sourced data,
owing to its extensive sampling across time and space, has the potential to test existing
ecological theories, expand our comprehension of ecosystems and the underlying
processes, eventually allowing us to forecast ecological dynamics in the context of climate
change.

When people photograph organisms using digital devices with GPS capabilities, the
images often contain timestamps and location details. Such images, when accompanied
by species identifications, serve as evidence for tracking phenology and species
occurrences. This crowdsourcing approach has been particularly successful on web- or
mobile-based platforms such as eBird and iNaturalist (Chandler et al., 2017; Wood et al.,
2011). Individuals submit records to these platforms for various reasons, including a
desire to contribute to science and engage with cutting-edge technologies (Herodotou et
al., 2023; Kaplan Mintz et al., 2023). By making the process more enjoyable (i.e.
gamification), we can potentially gather even more biological data from the public
(Bowser et al., 2013; Ponti et al., 2015). Yet, the collection process of Community-sourced
data is usually not well-designed (e.g., spatially biassed "presence-only" data) (Feldman et
al., 2021; Steen et al., 2019) and its interpretation is challenging without proper statistical
modelling. Thus, although much effort has been invested in developing effective
monitoring and modelling methods for biodiversity assessment, current approaches can
be further improved by incorporating (i) more enjoyable community-based survey
platforms using mobile applications and (ii) employing an advanced statistical modelling
framework in estimating species distribution.

To fuel communities' engagement in biodiversity surveys and environmental education,
we launched the mobile application '*Biome*' in 2019 in Japan (Fujiki and Tatsuno, 2021).
For supporting species identification, *Biome* implements artificial intelligence (AI)
algorithms that generate lists of potential species and enable users to seek
help/suggestions from others for species identification (Figure 1) as in other applications
such as iNaturalist and eBird. The unique feature of *Biome* is gamification which offers
enjoyable experiences and facilitates communication among users (Fujiki and Tatsuno,
2021; Koide et al., 2023). For example, users can earn "points" by contributing in various
ways such as submitting records and suggesting species identifications to others, and
their levels are determined based on the total points earned. The inclusion of networking
and gamification elements can attract a wider user base, including those who may not
typically engage in community science (Bowser et al., 2013; Groom et al., 2021).
Consequently, *Biome* has accumulated data rapidly. Since its launch, 6 million records
have been collected through the app (by 17 October 2023). This is more than four times
greater than the number of records accumulated by GBIF (Global Biodiversity Information
Facility) from any data sources including iNaturalist and eBird during the same period in
Japan (ca. 1.3 million). The data gathered through the app has been used for conservation
planning and facilitating companies' financial disclosures by supplying and analysing
species occurrence records.

Species distribution models (SDMs) are effective statistical tools for assessing biodiversity
at specific sites while accounting for biases in survey efforts. SDMs use species
occurrence records and environmental conditions to estimate the potential geographic
ranges and suitable habitats for species (Booth et al., 2014; Box, 1981; Elith et al., 2011;
Hutchinson, 1957; Phillips et al., 2006). These models play a crucial role in conservation
and restoration planning by helping predict how changes in land use and climate impact
species distributions (Kindt, 2023; Porfirio et al., 2014; Urban et al., 2016). While species
presence/absence data—which needs extensive surveys by experts—is limited,
presence-only data—which can be obtained from communities' observations—is much
more available. MaxEnt (Phillips et al., 2006; Phillips and Dudík, 2008) is one of the most
popular SDM methods due to its computational efficiency and estimation accuracy (Valavi
et al., 2022). It can estimate species distribution from presence-only data by maximising
the entropy of the probability distribution while satisfying constraints based on the
available information (Elith et al., 2011; Phillips and Dudík, 2008). Since MaxEnt only
requires occurrence records, it is well-suited for empowering community-based
observations to predict species distributions. Also, while community-sourced data often
suffer from spatially-biassed sampling efforts (i.e., sampling tends to concentrate in
densely populated or touristic areas (Kendal et al., 2020; Reddy and Dávalos, 2003)), SDMs
such as MaxEnt can account for such spatial biases by considering the spatial distribution

of sampling efforts when selecting pseudo-absence (background) locations (Milanesi et
al., 2020; Phillips et al., 2009). When sampling efforts are adequately controlled, adding
community-sourced data improves the accuracy of SDMs (Johnston et al., 2018; Robinson
et al., 2020; Steen et al., 2019). This implies that SDMs may be substantially improved by
utilising rapidly accumulating *Biome*'s species occurrence records if we adequately control
the sampling efforts.

Here, we show the quality of community-based data gathered through the smartphone
app *Biome*, and how the data improves the prediction accuracy of species distribution.
First, we assess the quality of occurrence records by investigating the fractions of non-
wild and misidentified records. Second, we built SDMs based on two types of data: (i)
traditional survey data (e.g. forest inventory census, museum specimens and records
extracted from published researches) only and (ii) a mixture of traditional survey and
*Biome* data. We then compare the performance of the two SDMs. We modelled the
distributions of 132 terrestrial animals and seed plants in the Japanese archipelago which
covers subtropical to boreal areas. We finally discuss how our SDMs relying on
community-sourced data may contribute to meeting the goals of GBF.

# Results

## The amount and quality of Biome data

By 7 July 2023, *Biome* had accumulated 5,275,457 occurrence records of 40,957 species
across the Japanese archipelago (Figure 2A). The amount of occurrence records
submitted to *Biome* has increased across the years (Figure 2B). On average in 2022, users
submitted 5,407 records per day. The distribution of data along environmental gradients
somewhat differs between *Biome* and Traditional survey data. To elucidate this distinction,
we employed principal component (PC) analysis to summarise all environmental variables.
The two datasets demonstrated divergent distribution patterns along PC1 (Figure 2C).
This component, accounting for 6.1% of the total variation, is primarily influenced by land
use, topography, and climate (supplementary file 1). Among the environmental variables,
a notable contrast between the datasets was observed in relation to the natural-urban
gradient. The *Biome* data exhibited a relatively uniform distribution encompassing the
entire gradient, while Traditional survey data was substantially biassed towards natural
areas (Figure 2C). The majority of records are attributed to insects (31.2%) and to seed

plants (41.8%), which are relatively accessible and can be easily photographed using
smartphones (Figure 2D).

Out of all the records submitted to *Biome*, a total of 2,373,303 records (45.0%) successfully
passed through the automatic filtering process. This dataset, referred to as the *Biome*
data, is utilised for subsequent investigations. The quality of *Biome* data varied across
taxa and the rarities of species (Table 1). The fraction of the records of wild individuals
exceeded 97% in insects and birds, while it was lower than 90% in molluscs, seed plants,
mammals and fishes. Among the records of wild individuals, at the species level,
identification accuracy was higher than 95% in birds, reptiles, mammals and amphibians
but less than 90% in insects, fishes and seed plants. At the genus level, identification
accuracy was higher than 90% in all taxa except for insects. In the case of fishes and seed
plants, identifications became 5-6% more accurate at the genus level compared to the
species level. The family was correctly identified in more than 94% of records in all taxa
examined. Common species had higher identification accuracy than rare species (average
value, 95% vs. 87%). This tendency was prominent in insects and seed plants, but less in
the other taxa. These results suggest that identifying rare species in taxonomically diverse
taxa (i.e. seed plants and insects) is a challenging task.

## 203  The performance of species distribution models

SDMs using *Biome*+Traditional data, including *Biome* data at 50%, were more accurate
than those modelled only using Traditional survey data when the two datasets have the
same amount of occurrence records (Figure 3). Our analysis revealed that although the
intercept of the Boyce Index (BI, model accuracy metric that ranges between -1 to 1) did
not differ between the two datasets (generalised linear mixed model, see Methods:
$\beta$=0.02±0.03, *t*=0.60, *P*=0.55), *Biome*+Traditional data consistently led to a more rapid
increase in SDM accuracy as the amount of data increased, comparing to models solely
relying on Traditional survey data ($\beta$=0.02±0.01, *t*=3.72, *P*<0.001).

When compared to SDMs using Traditional survey data, those using *Biome*+Traditional
data achieved a high level of accuracy with a much smaller amount of data. For instance,
BI which ranges from –1 to 1, exceeds 0.9 with 294±471 records (mean±SD across all
species) in the *Biome*+Traditional data, whereas the Traditional survey data requires
2,129±4,157 records to achieve the same accuracy. This was also true in endangered
species (included in Japanese national or prefectural red lists); although 2,336±3,718
Traditional survey records were required to exceed 0.9 of BI, only 338±571 were required
for *Biome*+Traditional data.

Because we controlled the proportion of *Biome* data within the *Biome*+Traditional data as
50%, the amount of records of the *Biome*+Traditional data is often limited. In cases where
a species had less *Biome* data compared to Traditional survey data, the total amount of
records of *Biome*+Traditional data ends up being smaller than that of Traditional survey
data alone. Therefore, the two datasets did not differ in the best model performances in
each species (BIs of *Biome*+Traditional data: 0.81±0.20; Traditional survey data: 0.83±0.20).

────────────────────────────────────────────

# Discussion

## *Biome*: The amount and quality of submitted data

Since its launch in 2019, the app *Biome* accumulates species occurrence data rapidly
(Figure 2). Despite our concerted efforts to engage non-expert users through gamification
features, it is important to acknowledge that an excessive influx of non-expert users could
potentially compromise the quality of the collected data. This could manifest in
misidentifications or incomplete documentation, such as failing to appropriately label
non-wild individuals. We thus have developed algorithms to exclude such suspicious
records based on the features of records and users' behaviour on the app. The
implementation of automatic data filtering techniques is expected to enhance the quality
of the data, although further refinement is necessary. Notably, for insects and birds,
which encompass numerous species that can be kept in captivity, the majority of records
that underwent filtering procedures were restricted to observations of wild individuals.
Yet, the fraction of non-wild individuals is high in several taxa such as fishes and seed
plants. In response, we have updated the posting flow in the app to prompt users to
differentiate between non-wild and wild individuals. Further analysis is warranted to
evaluate the impact of this update on data quality.

Once we could exclude non-wild individuals, species identification accuracy exceeded
95% in taxa with moderate species diversity (amphibians, reptiles, birds and mammals).
In seed plants, *Biome*'s species Identification accuracy was 90%, which is higher than the
accuracy of auto-suggest identification by commonly used apps for plants (69%, PlantNet,
PlantSnap, LeafSnap, iNaturalist and Google Lens: (Hart et al., 2023)). During the invasive
plants survey in the US, the reports by non-professional volunteers were 72% correct
(Crall et al., 2011). The higher accuracy of species identification in *Biome* data can be
attributed to two key factors. Firstly, the vigilant oversight of the user community through

the "suggest identification" feature plays a crucial role. *Biome* encourages users to
participate in suggesting identifications by offering "points" as rewards for their
contributions. Secondly, the species identification AI algorithm leverages past occurrence
data from nearby areas, resulting in increasingly accurate automatic identifications as the
data accumulates. Given these, as a community science app, the data quality of *Biome* is
decent. Yet, rare species generally showed lower identification accuracy, which would
require identification by experts and further improvement of species identification AI
algorithm.

## 264   Species distribution modelling

The inclusion of *Biome* data resulted in improved accuracy of SDMs (Figure 3). The most
accurate model predictions were obtained when the training data consisted of 50-70%
*Biome* data (supplementary file 3), highlighting the necessity of incorporating both
traditional surveys and citizen observations for a comprehensive understanding of
species distributions (Miller et al., 2019; Pacifici et al., 2017; Robinson et al., 2020).

The improvement can be attributed to introducing data with different biases compared
to the Traditional survey data. Indeed, when controlling for the number of occurrence
records, the model performance was higher in the *Biome*+Traditional data compared to
the Traditional survey data. The variation in performance can be attributed to the
distribution of data in relation to environmental conditions. Traditional survey data
exhibits a strong bias towards natural areas, whereas *Biome* data is well balanced across
the natural-urban habitat gradients (Figure 2C). Therefore, incorporating *Biome* data
could significantly enhance modelling accuracy in urban and suburban landscapes, which
are typically underrepresented in traditional survey data. As pseudo-absences are
selected based on search effort, our models utilise numerous pseudo-absences from
these areas. Consequently, this might lead to better estimation of species absence in such
areas, not just presence, resulting in an overall increase in model accuracy across a wider
range of species. A balanced distribution along with the natural-urban gradient is
noteworthy because community science data is typically biassed towards human
population centres (Kendal et al., 2020; Reddy and Dávalos, 2003). This could be
influenced by the distribution of users' residencies, although we do not have specific
information about the users' locations. The app has collaborated with numerous local
governments across Japan, including nine prefectures and 29 local municipalities such as
cities and towns. Through these collaborations, the user base may be widely dispersed,
enriching the geographical coverage of *Biome* data.

The *Biome* data also can improve SDM accuracy by simply increasing the overall amount
of data. Essentially, SDM accuracy is enhanced with an increased amount of data (Figure
3) (Erickson and Smith, 2023; Stockwell and Peterson, 2002). In our analysis, we
maintained a fixed proportion of 50% for *Biome* data within the *Biome*+Traditional dataset,
which in turn restricted the amount of available *Biome*+Traditional data. However, our
preliminary analysis (supplementary file 3) demonstrates that the enhancement of SDM
accuracy occurs across a range of proportion variations for *Biome* data blending. This
implies that the proportion of *Biome* data does not necessarily need to be controlled.
Therefore, in practical application scenarios, the incorporation of *Biome* data
predominantly serves to augment the overall volume of training data.

The impact of community-sourced data on SDMs has primarily been investigated using
birds, with a limited focus on plants (Feldman et al., 2021). In our investigation, we
observed that incorporating *Biome* data improved SDM accuracy for seed plants and
insects, while the impact on birds remained unclear (Figure 3). This ambiguity is likely
because community-sourced data from platforms such as eBird are already incorporated
in Traditional data through GBIF. In comparison to other taxonomic groups, our results
indicate that seed plants exhibited lower model accuracy when evaluated against both
*Biome*+Traditional survey data (Figure 3) and Traditional survey data alone (Figure 3 -
Figure supplement 1). The variation in model accuracy among taxonomic groups may be
attributed to data quality issues in both *Biome* and Traditional survey data. For instance,
in *Biome* data, while the fractions of wild individuals were high in birds and insects, it was
lower for seed plants (Table 1). Compared with other taxa, distinguishing between wild
and non-wild individuals can be particularly difficult in plants when they are planted
outside. In addition, identifying plant species may be challenging in certain taxa, primarily
due to the absence of key identification traits on leaves and stems. This becomes
especially problematic when flowers are not present. These difficulties could potentially
impact the quality of Traditional data as well. Although few studies have simultaneously
assessed the quality of community-sourced data and its impact on SDMs across different
taxa, it is important to recognize that data quality can vary among taxa.

Importantly, SDMs for endangered species, which often suffer from data deficit (Erickson
and Smith, 2023; Wisz et al., 2008), became accurate in a much fewer amount of records
by blending *Biome* data (Figure 3). Specifically, a threshold of >0.9 Boyce index could be
reached with only around 300 records when using *Biome* data, whereas over 6 times of
data is required when using Traditional survey data only. This finding highlights the
importance of community-sourced data not only for monitoring the dynamics of
endangered species (Chandler et al., 2017; Zapponi et al., 2017) but also for modelling
purposes. Considering the rapid accumulation of *Biome* data, *Biome* data would make a
significant contribution to the more effective distribution modelling of endangered
species.

## Limitations of this study

In assessing data quality, reidentification was impossible for records that did not
photograph key traits for species identification. To address this limitation, further app
improvements can include allowing users to submit multiple images. Encouraging users
to document various body parts of organisms through multiple images would make
capturing key identification traits much easier. This will make reidentification easier, and
possibly improve automatic species identification accuracy.

Given the absence of a comprehensive, environmentally unbiased occurrence dataset
spanning a wide range of taxa, we assessed SDM accuracy not relying on an independent
test dataset. In this evaluation, the test data was meticulously crafted to include 25%
*Biome* data, serving as an intermediary proportion between *Biome*+Traditional (50%) and
Traditional survey data (0%). By leveraging the distinct distribution patterns of *Biome* and
Traditional survey data along environmental variables (Figure 2C), the test data would
better encapsulate the actual species distribution, compared to datasets composed
solely of either *Biome* or Traditional survey data. It is noteworthy that, even when the test
data exclusively consisted of Traditional survey data (i.e., unfavourable conditions for
*Biome*+Traditional data SDMs), the accuracy of SDMs derived from *Biome*+Traditional and
Traditional survey data did not differ (Figure 3 - Figure supplement 1). This result further
supports our conclusions that *Biome* provides valuable data for SDM in terms of the
amount and quality, and that blending *Biome* data improves SDM accuracy.

We evaluated SDMs based on spatial transferability using the central Japan region, which
encompasses a range of environmental conditions. However, the evaluation results may
not necessarily indicate transferability across the entire Japanese archipelago. Instead, in
the near future, we anticipate that we can evaluate SDM accuracy using temporal
transferability. The rapid accumulation of *Biome* data will allow us to evaluate the
temporal transferability using the occurrence dataset from different time periods, and
thus enable assessing their performance in much wider regions. In addition, limited data
availability for certain taxa hindered the assessment in those taxa (e.g., molluscs,
amphibians, reptiles, and mammals), but *Biome* would be a platform to overcome the
data limitation for many taxa.

Finally, our SDMs do not directly indicate the species' presence probability. The output
from presence-only SDMs usually deviates from the probability of presence when species
prevalence (i.e. the proportion of area where the species occupied, requiring
presence/absence data throughout the area) is unavailable (Elith et al., 2011; Ward et al.,
2009). Due to the unavailability of absence data, SDM outputs in this work are indirect

measures of species presence and thus are not directly comparable across different
species. Nonetheless, they are comparable within a species, providing useful information
for understanding species distributions.

# Future directions

By blending data from traditional surveys and communities, we improved the accuracy
of species distribution estimates. This enhanced estimation lays the groundwork for more
precise subsequent analyses. For instance, estimated distributions will be useful in
selecting new protected areas or areas with OECMs (Other Effective area-based
Conservation Measures: allowing a wider range of land use as long as biodiversity and
ecosystem services are sustained/improved). Using estimated distributions of each
species, hotspots of species or evolutionary diverse taxa can be inferred. Such sites will
be good candidates for protected areas (Jones et al., 2016) or OECMs (Shiono et al., 2021).
Further, estimated distributions can be used as input for spatial conservation
prioritisation tools (e.g. Marxan (Ball et al., 2009)).

In our experience, stakeholders—including corporate social responsibility managers and
conservation practitioners—often seek the list of species potentially inhabiting their
locations. Due to the uncertainty of SDMs and their thresholding into presence/absence,
on-site surveys remain essential for assessing biodiversity status. Yet, SDMs can make
such surveys cost-effective by screening important locations for on-site assessment (e.g.,
Locate phase in TNFD framework) and narrowing down the target species for surveying.
Improved estimation through SDMs can mitigate risks associated with their use in society
and enable more informed decision-making for conservation efforts.

The rapid accumulation of data from diverse locations holds the potential to unveil
valuable ecological patterns. The accumulated data enables early detection capabilities
for range expansions of invasive species (Sakai et al., in prep). For instance, *Biome* data
has hinted at potential range expansions in several insect species, including butterflies,
dragonflies, and stink bugs, as well as changes in wintering areas for birds (Biome Inc.,
2023). Given the diverse taxonomic coverage of *Biome* data (Figure 2D), detecting
phenological changes across various taxa may be possible. This, in turn, is useful in
uncovering phenological mismatches exacerbated by climate change, which can
significantly change the dynamics of interacting species (Renner and Zohner, 2018; Visser
and Gienapp, 2019). Moreover, *Biome* data is well-suited for assessing the effects of
urbanisation on ecosystems since it comprehensively spans both urban and natural
habitats (Figure 2C). The benefit of rapidly accumulating data, combined with recent
advancements in machine learning methods, opens up opportunities for conducting

time-series analyses. Community science data has rarely been used for time-series
population analysis due to its notable spatio-temporal bias in sampling efforts (Feldman
et al., 2021; Zhang et al., 2021). However, the two-step machine learning approach, as
demonstrated by Fink and colleagues in estimating bird population trends using eBird
data (Fink et al., 2023), sets a precedent. In the future, *Biome* data may facilitate the
inference of population dynamics for multiple taxa. This will enable various time-series
analyses to unveil ecosystem stability and interaction strength, which holds potential for
forecasting ecosystem dynamics (Laubmeier et al., 2020; Pennekamp et al., 2019; Ushio
et al., 2018).

For financial disclosures, companies will assess how their activities rely on ecosystem
services and their opportunities for protecting/recovering nature (TNFD, 2023). By
incorporating taxon-specific ecosystem services, multifaceted ecosystem services can be
preliminarily screened (Kass et al., 2023). For example, based on estimated distributions
of bumblebees or insectivorous animals, the functioning of pollination services or pest
regulation services might be inferred. Using counts of "likes" or records from *Biome* data,
the charismatic species can be determined. By identifying places with a high estimated
richness of charismatic species, potential areas for ecotourism can be screened. Because
SDMs allow us to simulate the impacts of changes in landuse and climate (Porfirio et al.,
2014; Urban et al., 2016), we will be able to forecast how those changes may influence
local biodiversity and/or ecosystem functioning. Hence, estimated distributions provide
the basis of nature-related financial disclosures.

Our platform facilitates collaboration among diverse stakeholders, including local
communities, landowners, and employees from both private companies and government
agencies. Engaging a broader spectrum of stakeholders is crucial for effective biodiversity
assessment, nature management planning, and nature-related financial disclosures: this
inclusivity allows for the incorporation of traditional knowledge into planning processes,
mitigates conflicts among stakeholders, and ultimately supports more seamless and
informed decision-making (Chan et al., 2021; Keough and Blahna, 2006; Linsley et al.,
2023; Roy et al., 2023; TNFD, 2023).  Supporting natural experiences for a wide range of
people is also expected to contribute to changing people's minds towards nature.
Through experiencing nature, people become familiar with it and subsequently make
pro-nature decisions (Soga and Gaston, 2023). We believe that community science can
significantly contribute to KM-GBF and create a sustainable society by fostering nature-
positive awareness in society and providing data tools that enable effective action.

# Methods

## Occurrence record accumulation through mobile app *Biome*

In April 2019, a free smartphone app called *Biome* was launched for the Japanese markets. The app has been downloaded 839,844 times by September 13, 2023. The app allows users to collect data on the distribution of plants and animals using their mobile devices. Users can post photographs of the plants and animals they find, and the app automatically records the location and timestamp from EXIF data. If the EXIF data is unavailable, users can manually input the locality and timestamp.

To support species identification, the app provides users with two options. First, the app provides a list of candidate species based on the image and metadata (e.g., location and timestamp). *Biome* employs a synergistic approach that integrates image recognition technology and geospatial data to facilitate species identification. The image recognition algorithm, constructed upon convolutional neural networks, classifies species at higher taxonomic levels. Subsequently, these candidates are refined based on their frequency of recent occurrences in the geographical area. Consequently, as the correctly identified records accumulate for a given area, species identification AI will improve the accuracy. Second, users can seek help from other users. If a user selects the "ask Biomers" button, their occurrence record is added to a waiting list that appears on the home screen. Other users can suggest possible identifications for the records, as in other records of which species was already identified.

Users can view and comment on other users' records. However, for conservation purposes, *Biome* automatically conceals the geolocations of endangered species that are listed on the Japanese national or prefectural red lists. This feature sets it apart from iNaturalist, where users must manually choose to hide the location of endangered species (Koide et al., 2023). The social networking function provides opportunities for communication among users, including non-experts (Fujiki and Tatsuno, 2021). Users earn "points" through their contributions, including record submissions and identification suggestions to other users, and progress to higher levels based on their total points. The points awarded depend on the rarity, conservation status, and societal impact of the species submitted, meaning that users earn more points when submitting records of rare, endangered, or invasive species. The app occasionally offers "Quests" events that provide users with an opportunity to earn additional points by submitting records from specific locations or of particular species, crucial for monitoring phenology. Through the variety of gamification features, we stimulate people to participate in biological surveys as a fun activity.

We obtained occurrence records submitted to *Biome* by 7 July, 2023. The raw data
collected through *Biome* contains invalid presence records which we defined in the
present study as unclear images, documenting non-wild individuals and
misidentifications, and images including some privacy issues. To improve data quality, we
excluded records deemed to be invalid mainly based on location metadata and users'
reactions to the record is as detailed below. This filtered *Biome* data is used in the
subsequent investigations.

## 482   Filtering suspicious occurrence record in *Biome* data

Occurrence records of non-wild individuals were eliminated as much as possible by using
information provided by users and location of records. *Biome* users sometimes report
inappropriate records (e.g., unclear images and images from websites or books), and we
excluded all of those reported records. All private records were excluded because they
can harbour inappropriate and misidentified records not being screened by other users.
We also excluded occurrence records that users had marked as non-wild individuals:
users have an option to label their records as photographing bred or cultivated
individuals, or specimens. Records from cultural centres (i.e. zoos, botanical gardens,
museums, and aquariums) and large pet stores were removed as well. During the data
correction process, we prioritise the suggestions provided by *certified users* (see below for
the definition), regardless of the decisions made by the users who originally created the
record. Furthermore, we exclude records that have not been posted by *certified users* or
have not received identification suggestions from *certified users*.

*Certified users* are defined as users who achieved the higher accuracy of species
identification (<15% of public occurrence records were suggested as misidentification by
other users), submitted few inappropriate records (<0.5% of public records), and have
created >20 public records. We also defined *specialist users*, a subset of *certified users*
identified in each taxa (see Figure 2 for the classification), who made a total of >30 records
or identification suggestions with high identification accuracy (the fraction of suggested
records is less than the average of *certified users* in the taxa). *Specialist users* are used in
determining pseudo-absence for SDMs.

## 505   Assessing the accuracy of records

We investigated the proportion of occurrence records within the *Biome* data that were
suitable for SDMs. Since SDMs are influenced by invalid presence records, we assessed

the quality of *Biome* data based on a total of 1420 records from rare and common species
of seed plants, molluscs, insects (including Arachnid and Insecta), fishes, mammals, birds,
reptiles and amphibians (Figure 4). We defined rare species as those with fewer than or
equal to 10 occurrences in *Biome* data, and common species as those with the highest
15% of records in each taxonomic category. In each of the seed plant and insect species
which account for the majority of *Biome* data (Figure 2D), we randomly selected 145
records of each rare and common species. For the other taxonomic categories, we chose
each of the 70 records from rare and common species.

Records were first screened whether they targeted organisms (images with no organisms
were discarded) and contained wild individuals. To assess the accuracy of species
identification, species in the records documented wild individuals were manually
reidentified by experts with taxonomic knowledge (Figure 4). These experts have
professional backgrounds, serving as a technician at a prefectural research institute (fish),
highly-experienced field survey conductors (plants and insects, respectively), a post-
doctoral researchers (amphibians and reptiles, and mammals, respectively), and a
museum curator (mollusks) specialising in the focal taxa. Then, by comparing species
identifications by the experts and on *Biome* data, the results were classified into two
categories: (1) correct based on the image and locality—based on the image,
identification was probably correct, and the image locality matches with habitat/range of
the species; (2) misidentification—records were reidentified by experts if possible. We
also examined if the identification was correct at genus and family levels.

# Species distribution models

## Occurrence data

To evaluate the impact of *Biome* data on SDM prediction accuracy, we compiled two
datasets: "Traditional survey data" and "*Biome*+Traditional data". The Traditional survey
data comprised records collected through conventional survey techniques (e.g. riverine
census, forest inventory census, and museum specimens) primarily sourced from The
National Census on River and Dam Environments (NCRE) and GBIF. In contrast, the
*Biome*+Traditional data encompassed records submitted to *Biome* that passed filtering
methods, in addition to the Traditional survey data. To control the relative proportion of
*Biome* data, we constrained the fraction of *Biome* data within the *Biome*+Traditional data
to 50% for each species. Our preliminary results showed that blending 50-70% of *Biome*
data in training data improved prediction accuracy (supplementary file 3).
542
For traditional survey data, we downloaded occurrence records of relevant taxa from

GBIF between April 20th, 2023. To prevent significant differences between the sampling
periods of the GBIF records and environmental data, we used the GBIF sampled after
1970. The clean_coordinates function of the R package 'CoordinateCleaner' was used to
remove records with erroneous coordinates such as records from country capitals and
centroids, and biodiversity institutions. We obtained occurrence data from the large
occurrence datasets such as National Census on River and Dam Environments (NCRE) and
Forest Ecosystem Diversity Basic Survey. For the areas or taxa where occurrences were
scarce, we further compiled the literature with detailed locality information, such as local
species inventories. The amount of occurrence records in the modelled species and
species coverage of each dataset are summarised in Table 2. For the species analysed (S9
Table), traditional survey data contains a negligible portion of community-sourced data
(5.5%) because GBIF contains community-sourced data from iNaturalist and eBird.

## Predictor variables

Predictors encompass a range of environmental variables recognized to impact species
distribution (Table 3): land use (Newbold et al., 2015), climate (bioclim variables (Booth et
al., 2014)), vegetation (Abe, 2018), lithology (Ott, 2020) and elevational range (Udy et al.,
2021). Additionally, categorical variables representing known biogeographic regions,
reflecting geological history, were included. We applied  Blakiston's Line —Tsugaru straits
dividing the northern and main islands of Japan (i.e., Hokkaido and Honshu islands)—
reflecting a significant historical migration barrier for mammals and birds (Dobson, 1994;
Saitoh et al., 2015). Due to the distinct fauna (Wepfer et al., 2016; Yamasaki, 2017), we
also specified oceanic islands (i.e. Ogasawara and Daito isles) which have never been
connected with the Asiatic continents. Continuous environmental variables were
transformed into linear, quadratic and hinge feature classes to illustrate nonlinear
associations between environments and species occurrence (Phillips et al., 2017). The
regularisation multiplier was set at 2.5, falling within the established optimal range of 1.5
to 4 (Elith et al., 2010; Moreno-Amat et al., 2015).

## Pseudo-absence reflecting search effort

We considered sampling efforts when selecting a total of 10,000 pseudo-absence
locations. To accommodate biases in sampling efforts, we assigned picking probabilities
as an increasing function of the amount of occurrence records of all and relevant taxa at
the grid cell (an index of sampling efforts)  (Milanesi et al., 2020; Phillips et al., 2009). That
is, grid cells with rich occurrence records of relevant taxa are more likely to be chosen as
pseudo-absences than cells with few records, as detailed below (see also Figure 5).

To generate pseudo-absence (i.e. background) data, we employed two approaches
considering different sampling efforts. The first approach incorporated all observers and
taxa, while the second approach focused on experts and relevant taxa (Figure4). In both
cases, pseudo-absences were selected from grid cells that lacked any occurrence records
of the species being modelled. However, due to variations in sampling efforts across
locations, it was important to address potential bias. To mitigate this bias, we adjusted
the picking probability based on the number of occurrences of other species in each grid
cell (Milanesi et al., 2020; Phillips et al., 2009).

In the first approach, we assumed that the users of *Biome* submit records of any taxon
without specifically selecting species from particular taxa. The picking probability was
simply determined by the total number of records from all taxa in the *Biome* data in every
grid. In the second approach, we considered the expertise of observers (Milanesi et al.,
2020) and the sampling effort for relevant taxa (Phillips et al., 2009). We also assumed
that Traditional surveys targeted particular taxa. Under this approach, we selected
records from *Biome* data contributed by *specialist users* and all records from the
Traditional survey data. From this subset of data, we calculated the number of records
for the taxa (e.g. seed plant, insect and amphibian) to which the modelled species
belonged. This information was then used to calculate the picking probability for each
grid cell. To account for the variability in record counts among locations, we applied a
logarithmic transformation to the number of records. We also added a value of 1.2 before
taking logarithms to allow for the selection of pseudo-absences with low probabilities,
particularly in locations with only one or no records of other species. Pseudo-absences
were not chosen from the spatial block used as test data, but otherwise, there were no
geographical restrictions on their selection.

Using the described approaches, we obtained a total of 10,000 pseudo-absences for our
analyses. The amount of pseudo-absences follows the default setting of MaxEnt (Elith et
al., 2011). For the models using *Biome*+Traditional dataset (also in *Biome*-blended dataset
in supplementary file 3), pseudo-absences were generated by merging each of the 5,000
points identified through the two approaches. Meanwhile, for SDMs using the Traditional
survey data only, we obtained 10,000 pseudo-absences by exclusively using the second
approach without incorporating *Biome* data.

## 612     Modelling

We modelled distributions of terrestrial seed plants and animals at a scale of 1 x 1 km
grid cell, based on Traditional survey data and *Biome*+Traditional data. To model species
distributions from presence-only data, several algorithms have been utilised, including
generalised additive models, random forest, and neural networks (Norberg et al., 2019;

Valavi et al., 2022). In our study, we opted for MaxEnt (Phillips and Dudík, 2008) due to its
high estimation accuracy and relatively low computational burden (Valavi et al., 2022). We
perfomed MaxEnt via ENMeval 2.0 package (Kass et al., 2021) on R 4.1.3 (R Core Team,
2021).

## Model evaluation

We evaluated the model by examining spatial transferability because we could not find
occurrence data that are environmentally unbiased and independent from training data.
To minimise spatial autocorrelation between training and test data, we set a spatial block
for splitting data (Araújo et al., 2019; Santini et al., 2021). As the spatial block, we chose
the central Japan region (latitude, 33.7°–37.7° N; longitude, 136.2°–137.6° E: Figure 6) which
covers various environments—alpine to coastal lowlands, metropolis to highly intact
areas.

To ensure a fair and balanced assessment of the accuracy of SDMs built from Traditional
survey data (0% *Biome* data) and *Biome*+Traditional data (50% *Biome* data), we compiled
a test dataset that embodies characteristics intermediate between these two datasets.
This composite test dataset encompasses 25% *Biome* data and 75% Traditional data,
effectively bridging the differences between the two original datasets and providing a
comprehensive basis for evaluating SDM accuracy.

Due to the presence of invalid records, *Biome* records were used as test data only when
multiple users recorded the same species within an identical 1km grid cell. Although
*Biome* data may include invalid records (i.e. non-wild individuals or misidentification), if
multiple users recorded the same species at the same place, any one of the records from
the place is likely to be valid. As we know the fraction of valid records within the *Biome*
dataset in each taxon (see Results), we can calculate the probability of the true presence
in a given location as follows, by assuming that records made by different users were
independent:

644
$$p_{tp} = 1 - (1 - p_{valid})^{n_{users}}$$

The probability of valid records at a given taxon is shown as $p_{valid}$, and the number of
users reported given species at the place is indicated as $n_{users}$. If $p_{tp}$ exceeds 99%, we
deemed that the species occurred in the location.

To reduce spatial sampling bias, we downsampled a dataset within Traditional survey
data, NCRE with massive records from freshwaters, to match the number of records from
the remaining Traditional survey data. This procedure is applied to all test datasets in

both the main analysis and preliminary analyses documented in Figure 3 - Figure
supplement 1 and supplementary file 3.

Boyce index (BI) was used to measure model performance because it was designed to
evaluate presence-only SDMs (Hirzel et al., 2006). In short, BI measures the correlation
between estimated habitat preference and the frequency of actual presence, and ranges
from -1 to 1. A high BI indicates high SDM accuracy that presence data points tend to be
located in grids with higher habitat suitability values. To reliably calculate BI, at least 50
occurrences should be needed in test data (Hirzel et al., 2006). Thus, we used 132 species
that have more than 50 occurrences in test data for calculating BI (supplementary file 4).

# 661  Examining influences of blending *Biome* data on SDM accuracy

Given that the accuracy of SDMs is affected by the amount and quality of data (Araújo et
al., 2019; Erickson and Smith, 2023; Stockwell and Peterson, 2002), blending *Biome* data
in SDMs may affect the model performances in two possible ways: by increasing the
overall amount of data, and/or by introducing data with different information than the
original data. We analysed to distinguish between these effects. We prepared two
different datasets: "Traditional survey data" and "*Biome*+Traditional data". Then, we
separately trained SDMs using these two datasets. We further varied the data size by
performing random downsampling, ranging from a minimum of 20 to a maximum of
20,000 records, in order to evaluate its impact on the model. As for the "*Biome*+Traditional
data" category, the proportion of *Biome* data was kept at 50%. For each condition, we
conducted three iterations of training and testing to reduce the impact of random
sampling stochasticity. Because the modelling was performed for each species, we
obtained BI for each species, amount of records, and dataset (i.e., two datasets consisted
of 132 species, each with a maximum of 123 conditions for the amount of records, and
the models were replicated three times, resulting in a total of 12,351 individual model
runs).

After obtaining BIs for each run, we evaluated the effects of data type (i.e.,
*Biome*+Traditional data or Traditional survey data) and species on BI while accounting for
the amount of records. For each species and under each amount of records, the mean BI
was calculated across the three iterations. Given that BI is a correlation coefficient, we
applied the Fisher z-transformation to these BIs to approximate their distribution as a
normal distribution. To the transformed BIs, we fitted a generalised linear mixed model
that accounted for both the fixed and interaction effects of data type and amount of
records. This model accommodated species identity as a random effect. The model was

implemented and tested using R packages lme4 (Bates et al., 2015) and lmerTest
(Kuznetsova et al., 2017), respectively.

─────────────────────────────────────────────

## 692 Acknowledgments

We are grateful to Jamie M. Kass for advice on the construction and evaluation of SDMs.
We thank Midzuho Tatsuno, Kazumichi Morishita, Hironori Tanaka, Kotaro Takai and
Shuhei Tochino for species identification; Akira Sawada and Yuko Maegawa for their
advice on the analysis; Trevor H Booth and Dalelan Anderson for comments on the
manuscript. We appreciate the invaluable contributions of *Biome* users. This research was
partly supported by the collaborative research agreement between MU at Kyoto
University and Biome Inc. from 2020 to 2021.

## 701 Conflict of Interest

702 KA and YN are employed by Biome Inc., of which the CEO is SF and the CTO is TG. SF and
703 TG are inventors of the species-identification-AI-algorithm JPN patents 6590417 and US
704 patents 11048969. MU and HN declare that they have no competing interests. All authors
705 will not financially benefit directly from the publication of this paper.

707 ─────────────────────────────────────────────

## 710 References

Abe S. 2018. Habitat classification for 69 near threatened plants based on national
vegetation survey data. *Veg Sci* **35**:67–88. doi:https://doi.org/10.15031/vegsci.35.67
Araújo MB, Anderson RP, Márcia Barbosa A, Beale CM, Dormann CF, Early R, Garcia RA,
Guisan A, Maiorano L, Naimi B, O'Hara RB, Zimmermann NE, Rahbek C. 2019.
Standards for distribution models in biodiversity assessments. *Sci Adv* **5**:eaat4858.
doi:10.1126/sciadv.aat4858
Atsumi K, Koizumi I. 2017. Web image search revealed large-scale variations in breeding
season and nuptial coloration in a mutually ornamented fish, Tribolodon hakonensis.
*Ecol Res* **32**:567–578. doi:10.1007/s11284-017-1466-z

Ball IR, Possingham HP, Watts ME. 2009. Marxan and relatives: Software for spatial
conservation prioritizationSpatial Conservation Prioritisation: Quantitative Methods
and Computational Tools. Oxford University Press. p. 320.
Bates DM, Maechler M, Bolker B, Walker S. 2015. lme4: linear mixed-effects models using
S4 classes. *J Stat Softw* **67**:1–48.
Biome Inc. 2023. The report of Climate Change Biosurvey in 2022.
Booth TH, Nix HA, Busby JR, Hutchinson MF. 2014. bioclim: the first species distribution
modelling package, its early applications and relevance to most current MaxEnt
studies. *Divers Distrib* **20**:1–9. doi:10.1111/ddi.12144
Bowser A, Hansen D, He Y, Boston C, Reid M, Gunnell L, Preece J. 2013. Using
gamification to inspire new citizen science volunteersProceedings of the First
International Conference on Gameful Design, Research, and Applications,
Gamification '13. New York, NY, USA: Association for Computing Machinery. pp. 18–
25. doi:10.1145/2583008.2583011
Box EO. 1981. Macroclimate and Plant Forms, Tasks for Vegetation Science. Dordrecht:
Springer Netherlands. doi:10.1007/978-94-009-8680-0
Bury TM, Sujith RI, Pavithran I, Scheffer M, Lenton TM, Anand M, Bauch CT. 2021. Deep
learning for early warning signals of tipping points. *Proc Natl Acad Sci U S A* **118**.
doi:10.1073/pnas.2106140118
Chan L, Hillel O, Werner P, Holman N, Coetzee I, Galt R, Elmqvist T. 2021. Handbook on
the Singapore Index on Cities' Biodiversity (also known as the City Biodiversity
Index). Montreal: Secretariat of the Convention on Biological Diversity and
Singapore: National Parks Board, Singapore: The Secretariat of the Convention on
Biological Diversity.
Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, Legind JK, Masinde S,
Miller-Rushing AJ, Newman G, Rosemartin A, Turak E. 2017. Contribution of citizen
science towards international biodiversity monitoring. *Biol Conserv*, SI:Measures of
biodiversity **213**:280–294. doi:10.1016/j.biocon.2016.09.004
Cornwell WK, Ackerly DD. 2009. Community assembly and shifts in plant trait distributions
across an environmental gradient in coastal California. *Ecol Monogr* **79**:109–126.
doi:10.1890/07-1134.1
Crall AW, Newman GJ, Stohlgren TJ, Holfelder KA, Graham J, Waller DM. 2011. Assessing
citizen science data quality: an invasive species case study. *Conserv Lett* **4**:433–442.
doi:10.1111/j.1755-263X.2011.00196.x
Dobson M. 1994. Patterns of distribution in Japanese land mammals. *Mammal Rev* **24**:91–
111. doi:10.1111/j.1365-2907.1994.tb00137.x
Elith J, Kearney M, Phillips S. 2010. The art of modelling range-shifting species. *Methods*
*Ecol Evol* **1**:330–342. doi:10.1111/j.2041-210X.2010.00036.x
Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ. 2011. A statistical explanation of
MaxEnt for ecologists. *Divers Distrib* **17**:43–57. doi:10.1111/j.1472-
4642.2010.00725.x
Erickson KD, Smith AB. 2023. Modeling the rarest of the rare: a comparison between multi-
species distribution models, ensembles of small models, and single-species models
at extremely low sample sizes. *Ecography* **2023**:e06500. doi:10.1111/ecog.06500
Feldman MJ, Imbeau L, Marchand P, Mazerolle MJ, Darveau M, Fenton NJ. 2021. Trends
and gaps in the use of citizen science derived data as input for species distribution
models: A quantitative review. *PLOS ONE* **16**:e0234587.
doi:10.1371/journal.pone.0234587
Fink D, Johnston A, Strimas-Mackey M, Auer T, Hochachka WM, Ligocki S, Oldham
Jaromczyk L, Robinson O, Wood C, Kelling S, Rodewald AD. 2023. A Double
machine learning trend model for citizen science data. *Methods Ecol Evol* **14**:2435–
2448. doi:10.1111/2041-210X.14186
Fuccillo Battle K, Duhon A, Vispo CR, Crimmins TM, Rosenstiel TN, Armstrong-Davies LL,
de Rivera CE. 2022. Citizen science across two centuries reveals phenological
change among plant species and functional groups in the Northeastern US. *J Ecol*

**110**:1757–1774. doi:10.1111/1365-2745.13926

Fujiki S, Tatsuno M. 2021. Practice of citizen science for developing biodiversity monitoring
methods using mobile devices. *Jpn J Ecol* **71**:85–90.
doi:https://doi.org/10.18960/seitai.71.2_85

Gonzalez A, Chase JM, O'Connor MI. 2023. A framework for the detection and attribution of
biodiversity change. *Philos Trans R Soc B Biol Sci* **378**:20220182.
doi:10.1098/rstb.2022.0182

Groom Q, Pernat N, Adriaens T, de Groot M, Jelaska SD, Marčiulynienė D, Martinou AF,
Skuhrovec J, Tricarico E, Wit EC, Roy HE. 2021. Species interactions: next-level
citizen science. *Ecography* **44**:1781–1789. doi:10.1111/ecog.05790

Hart AG, Bosley H, Hooper C, Perry J, Sellors-Moore J, Moore O, Goodenough AE. 2023.
Assessing the accuracy of free automated plant identification applications. *People*
*Nat* **5**:929–937. doi:10.1002/pan3.10460

Herodotou C, Ismail N, I. Benavides Lahnstein A, Aristeidou M, Young AN, Johnson RF,
Higgins LM, Ghadiri Khanaposhtani M, Robinson LD, Ballard HL. 2023. Young
people in iNaturalist: a blended learning framework for biodiversity monitoring. *Int J*
*Sci Educ Part B* **0**:1–28. doi:10.1080/21548455.2023.2217472

Hirzel AH, Le Lay G, Helfer V, Randin C, Guisan A. 2006. Evaluating the ability of habitat
suitability models to predict species presences. *Ecol Model*, Predicting Species
Distributions **199**:142–152. doi:10.1016/j.ecolmodel.2006.05.017

Hutchinson EG. 1957. Concluding remarksCold Spring Harbor Symposia on Quantitative
Biology. Cold Spring Harbor Laboratory Press.

IPBES. 2019. Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
Bonn, Germany: IPBES secretariat.

Johnston A, Fink D, Hochachka WM, Kelling S. 2018. Estimates of observer expertise
improve species distributions from citizen science data. *Methods Ecol Evol* **9**:88–97.
doi:10.1111/2041-210X.12838

Jones KR, Watson JEM, Possingham HP, Klein CJ. 2016. Incorporating climate change into
spatial conservation prioritisation: A review. *Biol Conserv* **194**:121–130.
doi:10.1016/j.biocon.2015.12.008

Kaplan Mintz K, Arazy O, Malkinson D. 2023. Multiple forms of engagement and motivation
in ecological citizen science. *Environ Educ Res* **29**:27–44.
doi:10.1080/13504622.2022.2120186

Kass JM, Fukaya K, Thuiller W, Mori AS. 2023. Biodiversity modeling advances will improve
predictions of nature's contributions to people. *Trends Ecol Evol* **0**.
doi:10.1016/j.tree.2023.10.011

Kass JM, Muscarella R, Galante PJ, Bohl CL, Pinilla-Buitrago GE, Boria RA, Soley-Guardia
M, Anderson RP. 2021. ENMeval 2.0: Redesigned for customizable and reproducible
modeling of species' niches and distributions. *Methods Ecol Evol* **12**:1602–1608.
doi:10.1111/2041-210X.13628

Kendal D, Egerer M, Byrne JA, Jones PJ, Marsh P, Threlfall CG, Allegretto G, Kaplan H,
Nguyen HKD, Pearson S, Wright A, Flies EJ. 2020. City-size bias in knowledge on
the effects of urban nature on people and biodiversity. *Environ Res Lett* **15**:124035.
doi:10.1088/1748-9326/abc5e4

Keough HL, Blahna DJ. 2006. Achieving Integrative, Collaborative Ecosystem Management.
*Conserv Biol* **20**:1373–1382. doi:10.1111/j.1523-1739.2006.00445.x

Kindt R. 2023. TreeGOER: A database with globally observed environmental ranges for
48,129 tree species. *Glob Change Biol*. doi:10.1111/gcb.16914

Klinger YP, Eckstein RL, Kleinebecker T. 2023. iPhenology: Using open-access citizen
science photos to track phenology at continental scale. *Methods Ecol Evol* **14**:1424–
1431. doi:10.1111/2041-210X.14114

Kobori H, Dickinson JL, Washitani I, Sakurai R, Amano T, Komatsu N, Kitamura W,
Takagawa S, Koyama K, Ogawara T, Miller-Rushing AJ. 2016. Citizen science: a
new approach to advance ecology, education, and conservation. *Ecol Res* **31**:1–19.

doi:10.1007/s11284-015-1314-y

Koide D, Tsujimoto S, Kumagai N, Ikegami M, Nishihiro J. 2023. Species' spatiotemporal
distribution platform based on citizen science through desirable circulation between
the real and digital worlds. *Jpn J Conserv Ecol.*
doi:https://doi.org/10.18960/hozen.2217

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest package: Tests in linear
mixed effects models. *J Stat Softw* **82**. doi:10.18637/jss.v082.i13

Larson ER, Graham BM, Achury R, Coon JJ, Daniels MK, Gambrell DK, Jonasen KL, King
GD, LaRacuente N, Perrin-Stowe TI, Reed EM, Rice CJ, Ruzi SA, Thairu MW,
Wilson JC, Suarez AV. 2020. From eDNA to citizen science: emerging tools for the
early detection of invasive species. *Front Ecol Environ* **18**:194–202.
doi:10.1002/fee.2162

Laubmeier AN, Cazelles B, Cuddington K, Erickson KD, Fortin MJ, Ogle K, Wikle CK, Zhu K,
Zipkin EF. 2020. Ecological Dynamics: Integrating Empirical, Statistical, and
Analytical Methods. *Trends Ecol Evol* **35**:1090–1099. doi:10.1016/j.tree.2020.08.006

Leighton GRM, Hugo PS, Roulin A, Amar A. 2016. Just Google it: assessing the use of
Google Images to describe geographical variation in visible traits of organisms.
*Methods Ecol Evol* **7**:1060–1070. doi:10.1111/2041-210X.12562

Linsley P, Abdelbadie Roba, Abdelbadie Rasha. 2023. The Taskforce on Nature-related
Financial Disclosures must engage widely and justify its market-led approach. *Nat*
*Ecol Evol* **7**:1343–1346. doi:10.1038/s41559-023-02113-w

Loh J, Green RE, Ricketts T, Lamoreux J, Jenkins M, Kapos V, Randers J. 2005. The Living
Planet Index: using species population time series to track trends in biodiversity.
*Philos Trans R Soc B Biol Sci* **360**:289–295. doi:10.1098/rstb.2004.1584

Milanesi P, Mori E, Menchetti M. 2020. Observer-oriented approach improves species
distribution models from citizen science data. *Ecol Evol* **10**:12104–12114.
doi:10.1002/ece3.6832

Miller DAW, Pacifici K, Sanderlin JS, Reich BJ. 2019. The recent past and promising future
for data integration methods to estimate species' distributions. *Methods Ecol Evol*
**10**:22–37. doi:10.1111/2041-210X.13110

Miya M, Sado T, Oka S, Fukuchi T. 2022. The use of citizen science in fish eDNA
metabarcoding for evaluating regional biodiversity in a coastal marine region: A pilot
study. *Metabarcoding Metagenomics* **6**:e80444. doi:10.3897/mbmg.6.80444

Moreno-Amat E, Mateo RG, Nieto-Lugilde D, Morueta-Holme N, Svenning J-C, García-
Amorena I. 2015. Impact of model complexity on cross-temporal transferability in
Maxent species distribution models: An assessment using paleobotanical data. *Ecol*
*Model* **312**:308–317. doi:10.1016/j.ecolmodel.2015.05.035

Mori AS, Suzuki KF, Hori M, Kadoya T, Okano K, Uraguchi A, Muraoka H, Sato T, Shibata
H, Suzuki-Ohno Y, Koba K, Toda M, Nakano S, Kondoh M, Kitajima K, Nakamura M.
2023. Perspective: sustainability challenges, opportunities and solutions for long-term
ecosystem observations. *Philos Trans R Soc B Biol Sci* **378**:20220192.
doi:10.1098/rstb.2022.0192

Newbold T, Hudson LN, Arnell AP, Contu S, De Palma A, Ferrier S, Hill SLL, Hoskins AJ,
Lysenko I, Phillips HRP, Burton VJ, Chng CWT, Emerson S, Gao D, Hale GP, Hutton
J, Jung M, Sanchez-Ortiz K, Simmons BI, Whitmee S, Zhang H, Scharlemann JPW,
Purvis A. 2016. Has land use pushed terrestrial biodiversity beyond the planetary
boundary? A global assessment. *Science* **353**:291–288.
doi:10.1126/science.aaf2201

Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, Börger L, Bennett DJ,
Choimes A, Collen B, Day J, De Palma A, Díaz S, Echeverria-Londoño S, Edgar MJ,
Feldman A, Garon M, Harrison MLK, Alhusseini T, Ingram DJ, Itescu Y, Kattge J,
Kemp V, Kirkpatrick L, Kleyer M, Correia DLP, Martin CD, Meiri S, Novosolov M, Pan
Y, Phillips HRP, Purves DW, Robinson A, Simpson J, Tuck SL, Weiher E, White HJ,
Ewers RM, Mace GM, Scharlemann JPW, Purvis A. 2015. Global effects of land use
on local terrestrial biodiversity. *Nature* **520**:45–50. doi:10.1038/nature14324

Norberg A, Abrego N, Blanchet FG, Adler FR, Anderson BJ, Anttila J, Araújo MB, Dallas T, Dunson D, Elith J, Foster SD, Fox R, Franklin J, Godsoe W, Guisan A, O'Hara B, Hill NA, Holt RD, Hui FKC, Husby M, Kålås JA, Lehikoinen A, Luoto M, Mod HK, Newell G, Renner I, Roslin T, Soininen J, Thuiller W, Vanhatalo J, Warton D, White M, Zimmermann NE, Gravel D, Ovaskainen O. 2019. A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels. *Ecol Monogr* **89**:e01370. doi:10.1002/ecm.1370

Ott RF. 2020. How Lithology Impacts Global Topography, Vegetation, and Animal Biodiversity: A Global-Scale Analysis of Mountainous Regions. *Geophys Res Lett* **47**:e2020GL088649. doi:10.1029/2020GL088649

Pacifici K, Reich BJ, Miller DAW, Gardner B, Stauffer G, Singh S, McKerrow A, Collazo JA. 2017. Integrating multiple data sources in species distribution modeling: a framework for data fusion*. *Ecology* **98**:840–850. doi:10.1002/ecy.1710

Pennekamp F, Iles AC, Garland J, Brennan G, Brose U, Gaedke U, Jacob U, Kratina P, Matthews B, Munch S, Novak M, Palamara GM, Rall BC, Rosenbaum B, Tabi A, Ward C, Williams R, Ye H, Petchey OL. 2019. The intrinsic predictability of ecological time series and its potential to guide forecasting. *Ecol Monogr* **89**:e01359. doi:10.1002/ecm.1359

Phillips SJ, Anderson RP, Dudík M, Schapire RE, Blair ME. 2017. Opening the black box: an open-source release of Maxent. *Ecography* **40**:887–893. doi:10.1111/ecog.03049

Phillips SJ, Anderson RP, Schapire RE. 2006. Maximum entropy modeling of species geographic distributions. *Ecol Model* **190**:231–259. doi:10.1016/j.ecolmodel.2005.03.026

Phillips SJ, Dudík M. 2008. Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* **31**:161–175. doi:10.1111/j.0906-7590.2008.5203.x

Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S. 2009. Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecol Appl* **19**:181–197. doi:10.1890/07-2153.1

Pocock MJO, Chandler M, Bonney R, Thornhill I, Albin A, August T, Bachman S, Brown PMJ, Cunha DGF, Grez A, Jackson C, Peters M, Rabarijaon NR, Roy HE, Zaviezo T, Danielsen F. 2018. Chapter Six - A Vision for Global Biodiversity Monitoring With Citizen Science In: Bohan DA, Dumbrell AJ, Woodward G, Jackson M, editors. Advances in Ecological Research, Next Generation Biomonitoring: Part 2. Academic Press. pp. 169–223. doi:10.1016/bs.aecr.2018.06.003

Pocock MJO, Tweddle JC, Savage J, Robinson LD, Roy HE. 2017. The diversity and evolution of ecological and environmental citizen science. *PLoS ONE* **12**:1–17. doi:10.1371/journal.pone.0172579

Ponti M, Hillman T, Stankovic I. 2015. Science and Gamification: The Odd Couple?Proceedings of the 2015 Annual Symposium on Computer-Human Interaction in Play, CHI PLAY '15. New York, NY, USA: Association for Computing Machinery. pp. 679–684. doi:10.1145/2793107.2810293

Porfirio LL, Harris RMB, Lefroy EC, Hugh S, Gould SF, Lee G, Bindoff NL, Mackey B. 2014. Improving the Use of Species Distribution Models in Conservation Planning and Management under Climate Change. *PLOS ONE* **9**:e113749. doi:10.1371/journal.pone.0113749

R Core Team. 2021. R: A language and environment for statistical computing. *R Found Stat Comput Vienna Austria*.

Reddy S, Dávalos LM. 2003. Geographical sampling bias and its implications for conservation priorities in Africa. *J Biogeogr* **30**:1719–1727. doi:10.1046/j.1365-2699.2003.00946.x

Renner SS, Zohner CM. 2018. Climate Change and Phenological Mismatch in Trophic Interactions Among Plants, Insects, and Vertebrates. *Annu Rev Ecol Evol Syst* **49**:165–182. doi:10.1146/annurev-ecolsys-110617-062535

Robinson OJ, Ruiz-Gutierrez Viviana, Reynolds MD, Golet GH, Strimas-Mackey M, Fink D.

2020. Integrating citizen science data with expert surveys increases accuracy and spatial extent of species distribution models. *Divers Distrib* **26**:976–986. doi:10.1111/ddi.13068

Roy HE, Pauchard A, Stoett P, Renard Truong T, Bacher S, Galil BS, Hulme PE, Ikeda T, Sankaran KV, McGeoch MA, Meyerson LA, Nuñez MA, Ordonez A, Rahlao SJ, Schwindt E, Seebens H, Sheppard AW, Vandvik V. 2023. IPBES Invasive Alien Species Assessment: Summary for Policymakers. Zenodo. doi:10.5281/zenodo.8314303

Saitoh T, Sugita N, Someya S, Iwami Y, Kobayashi S, Kamigaichi H, Higuchi A, Asai S, Yamamoto Y, Nishiumi I. 2015. DNA barcoding reveals 24 distinct lineages as cryptic bird species candidates in and around the Japanese Archipelago. *Mol Ecol Resour* **15**:177–186. doi:10.1111/1755-0998.12282

Santini L, Benítez-López A, Maiorano L, Čengić M, Huijbregts MAJ. 2021. Assessing the reliability of species distribution projections in climate change research. *Divers Distrib* **27**:1035–1050. doi:10.1111/ddi.13252

Schmitt S, Pouteau R, Justeau D, de Boissieu F, Birnbaum P. 2017. ssdm: An r package to predict distribution of species richness and composition based on stacked species distribution models. *Methods Ecol Evol* **8**:1795–1803. doi:10.1111/2041-210X.12841

Scholes RJ, Biggs R. 2005. A biodiversity intactness index. *Nature* **434**:45–49. doi:10.1038/nature03289

Shiono T, Kubota Y, Kusumoto B. 2021. Area-based conservation planning in Japan: The importance of OECMs in the post-2020 Global Biodiversity Framework. *Glob Ecol Conserv* **30**:e01783. doi:10.1016/j.gecco.2021.e01783

Soga M, Gaston KJ. 2023. Nature benefit hypothesis: Direct experiences of nature predict self-reported pro-biodiversity behaviors. *Conserv Lett* **n/a**:e12945. doi:10.1111/conl.12945

Steen VA, Elphick CS, Tingley MW. 2019. An evaluation of stringent filtering to improve species distribution models from citizen science data. *Divers Distrib* **25**:1857–1869. doi:10.1111/ddi.12985

Stockwell DRB, Peterson AT. 2002. Effects of sample size on accuracy of species distribution models. *Ecol Model* **148**:1–13. doi:10.1016/S0304-3800(01)00388-X

Tilman D, Reich PB, Knops JMH. 2006. Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature* **441**:629–632. doi:10.1038/nature04742

TNFD. 2023. Taskforce on Nature-related Financial Disclosures (TNFD) Recommendations version 1.0.

Udy K, Fritsch M, Meyer KM, Grass I, Hanß S, Hartig F, Kneib T, Kreft H, Kukunda CB, Pe'er G, Reininghaus H, Tietjen B, Tscharntke T, van Waveren C-S, Wiegand K. 2021. Environmental heterogeneity predicts global species richness patterns better than area. *Glob Ecol Biogeogr* **30**:842–851. doi:10.1111/geb.13261

Urban MC, Bocedi G, Hendry AP, Mihoub J-B, Pe'er G, Singer A, Bridle JR, Crozier LG, De Meester L, Godsoe W, Gonzalez A, Hellmann JJ, Holt RD, Huth A, Johst K, Krug CB, Leadley PW, Palmer SCF, Pantel JH, Schmitz A, Zollner PA, Travis JMJ. 2016. Improving the forecast for biodiversity under climate change. *Science* **353**:aad8466. doi:10.1126/science.aad8466

Ushio M, Hsieh CH, Masuda R, Deyle ER, Ye H, Chang CW, Sugihara G, Kondoh M. 2018. Fluctuating interaction network and time-varying stability of a natural fish community. *Nature* **554**:360–363. doi:10.1038/nature25504

Valavi R, Guillera-Arroita G, Lahoz-Monfort JJ, Elith J. 2022. Predictive performance of presence-only species distribution models: a benchmark study with reproducible code. *Ecol Monogr* **92**:e01486. doi:10.1002/ecm.1486

Visser ME, Gienapp P. 2019. Evolutionary and demographic consequences of phenological mismatches. *Nat Ecol Evol* **12**. doi:10.1038/s41559-019-0880-8

Wallace RD, Bargeron CT. 2014. Identifying invasive species in real time: early detection and distribution mapping system (EDDMapS) and other mapping tools. *Invasive Species Glob Clim Change*, CABI Invasives Series 219–231.

doi:10.1079/9781780641645.0219

Ward G, Hastie T, Barry S, Elith J, Leathwick JR. 2009. Presence-Only Data and the EM Algorithm. *Biometrics* **65**:554–563. doi:10.1111/j.1541-0420.2008.01116.x

Wepfer PH, Guénard B, Economo EP. 2016. Influences of climate and historical land connectivity on ant beta diversity in East Asia. *J Biogeogr* **43**:2311–2321. doi:10.1111/jbi.12762

Wisz MS, Hijmans RJ, Li J, Peterson AT, Graham CH, Guisan A, Group NPSDW. 2008. Effects of sample size on the performance of species distribution models. *Divers Distrib* **14**:763–773. doi:10.1111/j.1472-4642.2008.00482.x

Wood C, Sullivan B, Iliff M, Fink D, Kelling S. 2011. eBird: Engaging Birders in Science and Conservation. *PLOS Biol* **9**:e1001220. doi:10.1371/journal.pbio.1001220

Yamasaki T. 2017. Biogeographic Pattern of Japanese Birds: A Cluster Analysis of Faunal Similarity and a Review of Phylogenetic Evidence In: Motokawa M, Kajihara H, editors. Species Diversity of Animals in Japan, Diversity and Commonality in Animals. Tokyo: Springer Japan. pp. 117–134. doi:10.1007/978-4-431-56432-4_4

Zapponi L, Cini A, Bardiani M, Hardersen S, Maura M, Maurizi E, Redolfi De Zan L, Audisio P, Bologna MA, Carpaneto GM, Roversi PF, Sabbatini Peverieri G, Mason F, Campanaro A. 2017. Citizen science data as an efficient tool for mapping protected saproxylic beetles. *Biol Conserv*, The role of citizen science in biological conservation **208**:139–145. doi:10.1016/j.biocon.2016.04.035

Zhang W, Sheldon BC, Grenyer R, Gaston KJ. 2021. Habitat change and biased sampling influence estimation of diversity trends. *Curr Biol* **31**:3656-3662.e3. doi:10.1016/j.cub.2021.05.066

# Figure, tables and supporting information

**Figure 1. Workflow of submitting records to *Biome*.** (1) Users can upload images that were taken by the smartphone camera or import existing images from the storage, including those imported from external devices. (2) Users select whether the image is about animals or plants to activate the species identification AI. (3) The AI analyses the image and its metadata to generate a candidate species list. (4) Alternatively, users can input the taxon name manually  and obtain a list of candidate species. To submit the occurrence record, users can either (5) seek identification assistance from other users through the "ask Biomers" feature, or (6) identify the species from the list. To the records, users can add memos and tags indicating phenology, life-stage, sex, and whether the individual is wild or captive.

**Figure 2. Description of data accumulated by *Biome*.** Data distributions are shown based on all records submitted to *Biome* by 7 July 2023 (*N*=5,275,457). **A** Spatial distribution of records across Japan. **B** Accumulation of records through time. The barplot represents the number of records each month and the line shows the cumulative amount of records. **C** Distributions of records along with PC1 of all environmental variables and standardised area occupancy of urban-type land uses. Grey and green represent distributions of Traditional and *Biome* data, respectively. **D** Taxonomic composition of records is shown as the area sizes. 'Other plant'

1039 consists of non-seed terrestrial plants; 'insects' include Arachnids and Insects; 'arthropods' cover
1040 any Arthropod not included in insects; 'other animals' covers all invertebrates not included in the
1041 taxa above.

1042

**Figure 3. The accuracy of species distribution models.** Accuracy of SDMs using Traditional
survey data (grey dots and lines) and *Biome*+Traditional data (i.e. 50% of *Biome* data: green). Each
SDM was performed with a specific dataset, species, and the amount of records. For each species
and amount of records, we computed the average model accuracy (Boyce index) from three
replicated runs. Subsequently, we calculated the median model accuracy across species for each
amount of records. These medians were then illustrated for each taxon in the strip of each respective
panel. The "Endangered" category includes species that are listed as endangered on Japan's national
or prefectural red lists.

1051

1052 **Figure 4. The workflow of checking accuracy of *Biome* data.**

1053

1054 **Figure 5. The workflow for selecting pseudo-absence (background) grid cells for SDMs**
1055 **using the *Biome*-Traditional dataset.** In this process, both *Biome* data and Traditional dataset
1056 are utilised to determine the suitable locations for pseudo-absence grid cells. However, when
1057 constructing SDMs using the Traditional dataset exclusively, *Biome* data is not involved in the
1058 selection of pseudo-absence points.

1059

1060 **Figure 6. Japanese archipelago, coloured by altitude.** Shaded area shows spatial block of test
1061 data. Retrieved from Wikipedia (2023, May 30), licensed under Creative Commons Attribution-
1062 ShareAlike 3.0 Unported (CC BY-SA 3.0).

1063

1064

1065 **Table 1. Data quality of *Biome*.** The fraction of records documenting wild individuals, and
1066 identification accuracy at species, genus and family levels among the records documenting wild
1067 individuals are shown. Species were identified only for records documenting wild individuals.

| Species group | Species rarity | *N* | Wild / total(%) | Species correct / wild (%) | Genus correct / wild (%) | Family correct / wild (%) |
|---|---|---|---|---|---|---|
| **TOTAL** | **TOTAL** | 1420 | 81.6 | 91 | 93.6 | 96.9 |
| **Seed plant** | **TOTAL** | 290 | 86.2 | 89.6 | 94.4 | 97.2 |

| | | | | | | |
|---|---|---|---|---|---|---|
| **Mollusca** | **TOTAL** | 140 | 87.9 | 90.2 | 91.1 | 96.7 |
| **Insect** | **TOTAL** | 290 | 100 | 83.4 | 86.9 | 94.1 |
| **Fish** | **TOTAL** | 140 | 73.6 | 87.4 | 93.2 | 96.1 |
| **Amphibian** | **TOTAL** | 140 | 93.6 | 96.2 | 96.2 | 98.5 |
| **Reptile** | **TOTAL** | 140 | 91.4 | 97.7 | 100 | 100 |
| **Bird** | **TOTAL** | 140 | 98.6 | 98.6 | 99.3 | 99.3 |
| **Mammal** | **TOTAL** | 140 | 80.7 | 95.6 | 95.6 | 96.5 |
| **TOTAL** | **Rare** | 710 | 88.7 | 87 | 91 | 95.6 |
| **TOTAL** | **Common** | 710 | 91 | 95 | 96.3 | 98.3 |
| **Seed plant** | **Rare** | 145 | 80.7 | 82.9 | 91.5 | 94.9 |
| **Seed plant** | **Common** | 145 | 91.7 | 95.5 | 97 | 99.2 |
| **Mollusca** | **Rare** | 70 | 82.9 | 86.2 | 87.9 | 96.6 |
| **Mollusca** | **Common** | 70 | 92.9 | 93.8 | 93.8 | 96.9 |
| **Insect** | **Rare** | 145 | 100 | 75.2 | 80 | 91.7 |
| **Insect** | **Common** | 145 | 100 | 91.7 | 93.8 | 96.6 |
| **Fish** | **Rare** | 70 | 74.3 | 88.5 | 94.2 | 94.2 |
| **Fish** | **Common** | 70 | 72.9 | 86.3 | 92.2 | 98 |
| **Amphibian** | **Rare** | 70 | 95.7 | 95.5 | 95.5 | 98.5 |
| **Amphibian** | **Common** | 70 | 91.4 | 96.9 | 96.9 | 98.4 |
| **Reptile** | **Rare** | 70 | 94.3 | 95.5 | 100 | 100 |
| **Reptile** | **Common** | 70 | 88.6 | 100 | 100 | 100 |

| Bird | Rare | 70 | 97.1 | 98.5 | 100 | 100 |
|---|---|---|---|---|---|---|
| Bird | Common | 70 | 100 | 98.6 | 98.6 | 98.6 |
| Mammal | Rare | 70 | 81.4 | 91.2 | 91.2 | 93 |
| Mammal | Common | 70 | 80 | 100 | 100 | 100 |

1068

1069 **Table 2. List of species occurrence datasets used for constructing SDMs.** To compare *Biome*
1070 dataset with the other datasets, iNaturalist and eBird data based on community science were
1071 classified as 'Traditional survey' data. *For the list of GBIF download doi and literatures, see
1072 supplementary file 2.

| Original dataset | Occurrence records of modelled species | | Species coverage among modelled species | Survey method | Data group in SDM | Down-load date | Availability |
|---|---|---|---|---|---|---|---|
| | *N* | Occu-pancy | | | | | |
| *Biome* (filtering applied) | 201114 | 8.6 | 132/132 | Citizen science through smartphone app | *Biome* | 7 July 2023 | https://biome.co.jp/ |
| National Census on River and Dam Environments (NCRE) | 1413541 | 60.2 | 126/132 | Traditional survey on freshwater and its adjacent ecosystems | Traditional survey | 10 Jan. 2023 | http://www.nilim.go.jp/lab/fbg/ksnkankyo/ |
| Institute records registered at GBIF | 530952 | 22.6 | 116/132 | Traditional survey and museum specimens | Traditional survey | 7 July 2023 | GBIF* |
| iNaturalist and eBird | 118050 | 5 | 110/132 | Citizen science through smartphone app and web service | Traditional survey* | 7 July 2023 | GBIF* |
| Forest | 80929 | 3.4 | 42/132 | Traditional survey | Traditional | 30 | http://fores |

| Ecosystem Diversity Basic Survey | | | | on forest trees | survey | March 2023 | tbio.jp/ |
|---|---|---|---|---|---|---|---|
| Literatures | 3293 | 0.1 | 130/132 | Traditional survey | Traditional survey | 31 March 2023 | Refs* |

1073

1074 **Table 3. Environmental data used for constructing SDMs.** Years indicate the data collection
1075 period. Usage in the SDM shows how the variables were converted before using in the species
1076 distribution modelling.

| Data | Variables | Year | Usage in the SDM | Available at |
|---|---|---|---|---|
| Land use | The area sizes of forests, rice fields, farms, wastelands, inland waters, beaches, ocean, golf courses, urbanised areas, and others | 2016 | Extracted six principal components (PCA) explained $\geqq$ 80% of total variation. PCs were converted into linear, quadratic and hinge terms | The Ministry of Land, Infrastructure, Transport and Tourism of Japan (MLIT) https://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-L03-a.html |
| Forest type | Forest type (planted and natural) | 1994-1998 | Converted into linear, quadratic and hinge terms | The Biodiversity Centre of Japan; http://gis.biodic.go.jp/webgis/index.html |
| Climate | Monthly average, minimum and maximum temperature and precipitation | 1981-2010 | Transformed into 19 bioclimatic variables (Booth et al., 2014), then extracted three PCs explained $\geqq$ 80% of total variation. Converted into linear, quadratic and hinge terms | MLIT, https://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-G02-v3_0.html |
| Elevation-al range | Differences between maximum and minimum elevation, and maximum slope | 1981 | Converted into linear, quadratic and hinge terms | MLIT, https://nlftp.mlit.go.jp/ksj/jpgis/datalist/KsjTmplt-G04-a.html |

| Vege-tation | The area sizes | 1998 | Transformed into 37 PCs of which total variation explained was more than 80%.<br>Converted into linear, quadratic and hinge terms | MOE, http://gis.biodic.go.jp/webgis/index.html |
|---|---|---|---|---|
| Geology | The area sizes of limestone and serpentinite | 2022 | Converted into linear, quadratic and hinge terms | The Research Institute of Geology and Geoinformation, https://gbank.gsj.jp/seamless/use.html |
| Geo-history | Blakiston's Line (Dobson, 1994; Saitoh et al., 2015), oceanic islands (Wepfer et al., 2016; Yamasaki, 2017) | | Categorical variables | |

1077

1078

**Figure 3 - Figure supplement 1. Accuracy of SDMs using Traditional survey data**
**(grey dots and lines) and *Biome*+Traditional data (i.e. 50% of *Biome* data: green),**
**evaluated against test data only consisting of Traditional survey data.**

**Supplementary file 1. Distributions of occurrence records along with**
**environmental variables.**

**Supplementary file 2. List of GBIF data doi and literatures compiled in occurrence**
**data.**

**Supplementary file 3. Determine the best blend of Traditional survey and *Biome***
**data.**

**Supplementary file 4. List of species for constructed species distribution models.**
1089

Ask other users

Use AI

Input
taxon name

Choose
species

**A** Spatial distribution

Density of records (/10km)

8000
400
20
0

**B** Temporal accumulation

Occurrence records in each month

Cumulative amount of occurrence records

300000 — — 6000000
— 4000000
200000 —
— 2000000
100000 —
0 — — 0

2019-07  2019-07  2020-07  2021-07  2021-07  2022-07  2023-01

**C** Environmental conditions

All variables

4000000
3000000
2000000
1000000
0

Traditional survey data

*Biome* data

Occurrence records

-5    0    5    10
PC1 value

Urbanized area occupancy

10000000
9500000
2000000
1500000
1000000
500000
0

0    1    2    3    4    5    6
Standardized value

**D** Taxonomic composition

Insect

Seed plant

Fish

Bird

Other plant
Mollusca
Crustacea
Amphibian
Reptile
Mammal
Other animal

*Species groups* examined

Seed plants | Insect | Mollusca | Fish
Amphibian | Reptile | Bird | Mammal

**Common species**
Species with
highest 15% of records
in the *species group*

**Rare species**
Species with
≤10 records

Randomly chose records from common/rare cateogories
in each *species group*:
each of 145 records in seed plants and insects,
each of 70 records in the rest of *species groups* were chosen

**Data check**

**Does the record document wild individuals?**
Clear and original images
photographed wild individuals

Yes

No

**Identification**
by experts

Finish

**Strictly correct**
based on
a single image

**Correct** based on
an image
and locality

**Misidentification**

**Valid records**