Characterization of sinking and suspended microeukaryotic communities in spring

Oyashio waters

Qingwei Yang[1], Yanhui Yang[2], Jun Xia[1], Hideki Fukuda[2], Yusuke Okazaki[1], Toshi Nagata[2], Hiroyuki

Ogata[1], Hisashi Endo[1*]

[1]Institute for Chemical Research, Kyoto University, Uji, Kyoto, Japan

[2]Atmosphere and Ocean Research Institute, The University of Tokyo, Chiba, Japan

Running title: Sinking and suspended microeukaryotic communities

Corresponding authors:

H. Endo, E-mail: endo@scl.kyoto-u.ac.jp, Phone: +81-774-38-3272

Abstract:

Microeukaryotes are important components of sinking particles contributing to carbon export from the

surface to deep oceans. The knowledge of the sinking microeukaryotic communities and their dynamics

is limited. We applied 18S rDNA metabarcoding method to investigate the microeukaryotic

communities in sinking and suspended particles distinguished by marine snow catchers (MSC) during

spring in the Oyashio region. Sinking particles displayed distinct communities and lower diversity than

did suspended particles. The community compositions of the sinking particles varied with depth,

suggesting that microeukaryotes were selectively removed through disaggregation or decomposition

during settling. Prymnesiophyceae and diatoms were effectively removed, as indicated by their

decreased abundance in the sinking particles at increasing depths. Conversely, phototrophic

dinoflagellates maintained a higher abundance in the sinking particles across depths, indicating

resistance to disaggregation and decomposition. Heterotrophic dinoflagellates and Spirotrichea were

enriched in sinking particles and marine stramenopiles (MAST) groups were enriched in suspended

particles. Sinking heterotrophic protist communities in the deep layers were similar to those in the

surface layers, whereas they differed from the suspended ones in the same layer. Therefore,

heterotrophic protists in surface layers were transported to deeper layers. Overall, our results

demonstrate the functional differences among microeukaryotes in the biological carbon pump.

Keywords: microeukaryotes, biological carbon pump, sinking particles, suspended particles, Marine

Snow Catcher, DNA metabarcoding

Introduction

The sinking of particulate organic carbon (POC) is a major driver of the oceanic biological carbon pump

(BCP), a process by which biologically fixed carbon is transferred from the euphotic layer to the deep

ocean (Boyd *et al.*, 2019). Global estimates of the sinking POC flux out of the surface mixed layer range

from 5 to 20 Gt C yr^{-1} (Zhang *et al.*, 2018); this influences the air-sea exchange of carbon dioxide and

the Earth's climate (Henson *et al.*, 2022). Among the various types of particulate constituents of sinking

POC, microeukaryotes, including photosynthetic and heterotrophic protists, play an important role in

determining the structure, sinking velocity, and chemical composition of sinking POC, thereby

influencing the magnitude and efficiency of the BCP (Guidi et al., 2009; Trudnowska et al., 2021).

Therefore, examining the taxonomic composition and variability of the microeukaryotic assemblages

in sinking POC is important.

Conventionally, microeukaryotes in sinking particles were collected using sediment traps and examined

using light microscopy and biomarker pigment analysis (Scharek *et al.*, 1999; Agusti *et al.*, 2015).

Silicified or calcified phytoplankton, such as diatoms and coccolithophorids, were often abundant in

sinking particles, suggesting that their mineral cell walls (frustules or plates) function as ballasts to

enhance the settlement of aggregates containing these cells or their remnants (Bauerfeind *et al.*, 2009).

However, these techniques have limited taxonomic resolutions and difficulty in identifying protists

lacking mineral cell walls, especially when small and non-pigmented (heterotrophic). Recent advances

in environmental DNA-based approaches, such as 18S rDNA metabarcoding method (Duret *et al.*, 2020;

Durkin *et al.*, 2022), have provided comprehensive information on community composition, improving

our understanding of the role of microeukaryotes in the export of carbon to the deep ocean. Small and

non-mineralizing species, such as non-calcifying haptophytes and dinoflagellates, could have been

underestimated in previous studies (Richardson, 2019; Lin *et al.*, 2022). Time-series sediment trap

studies at abyssal depths have documented the occurrence of various eukaryotic lineages of protists

originating from surface layers (Boeuf et al., 2019; Preston et al., 2020).

Despite the growing knowledge of microeukaryotic assemblages associated with sinking particles, some

important questions remain unresolved. One such question is whether certain microeukaryotes are

preferentially associated with sinking rather than suspended particles and are more efficiently

transferred from the sunlit layer to the abyssal depths. The sinking POC flux is strongly attenuated

through the mesopelagic layer (200−1000 m), leaving less than 10% of the exported POC available for

consumption in the bathypelagic layer and abyssal sediments (Zhang *et al.*, 2018). Several mechanisms

have been proposed to explain this depth-dependent POC flux attenuation, including microbial

degradation and solubilization (Anderson and Tang, 2010), zooplankton consumption, and

fragmentation due to the turbulence created by zooplankton swimming and feeding activities (Turner,

2015). The relative importance of the different mechanisms remains unclear; however, microeukaryotic

cells that are more resistant to POC flux attenuation processes (disaggregation or decomposition) could

contribute more strongly to the transfer of carbon to the deeper depths, thereby enhancing the efficiency

of BCP. Conversely, easily disaggregated, or degraded microeukaryotes could contribute to carbon

remineralization in the mesopelagic layer. Recently, large sedimentation chambers [marine snow

catcher (MSC), (Lampitt *et al.*, 1993)] were used to differentiate microeukaryotic assemblages in

sinking and suspended particles. The composition of microeukaryotic assemblages differed between

sinking and suspended particles in the Scotia Sea (Duret et al. 2020). In the upper mesopelagic layer,

a chain-forming diatoms dominated the sinking particle assemblages, whereas prymnesiophytes were

abundant in suspended particles, suggesting that diatom-enriched particles are more efficient in carbon

transfer to the upper mesopelagic layer than are those enriched in prymnesiophytes. This study provided

valuable insights into the functional differences among microeukaryotic taxa in the regulation of BCP;

the data were limited to the upper mesopelagic layer. This hampers the assessment of microeukaryotic

dynamics throughout the mesopelagic water column.

In this study, we used the MSC to investigate the characteristics of the microeukaryotic community

associated with sinking and suspended particles and their changes during the sedimentation through the

mesopelagic water column. Samples were collected from three depths: the subsurface chlorophyll

maximum (SCM, depth 11–30 m), 10 m below the pycnocline (PC, depth 65–250 m), and the bottom

boundary layer (BBL, depth 289–1489 m). This study was conducted during spring blooms in the

Oyashio waters off Hokkaido in the western North Pacific Ocean (Kuroda *et al.*, 2019). This oceanic

region is known for its strong POC flux (Takahashi et al., 2002) and high efficiency of BCP (Kawakami

and Honda, 2007) during blooms. Earlier sediment trap observations suggest that diatoms in this area

play a key role in transporting POC to deep oceans owing to their large size and high sedimentation

rates (Kawakami and Honda, 2007; Kawakami *et al.*, 2015). However, the detailed structures of sinking

particles and their decomposition processes during sedimentation from the surface to deep oceans

remain unexplored.

Results

Sequencing statistics and environmental conditions

A total of 3,913,663 18S rDNA (V4 region) paired-end reads were obtained from 32 suspended and

sinking particle fractions. After quality trimming, merging, and chimera and singleton removal, 2,416

amplicon sequence variants (ASVs) were generated. Among these, 242 ASVs taxonomically assigned

to multicellular organisms were excluded; in addition, 290 ASVs that were unique to BBL samples

(0.3–2% per BBL sample), potentially indigenous benthic organism, were removed (Fig. S1). The

remaining 1,875 protist ASVs were subjected to downstream analyses, and their relative frequencies

were normalized by rarefying the smallest number of sequences per sample (33,794 paired-end reads,

Fig. S2). The protist ASVs were classified into either phototrophs (686 ASVs, including mixotrophs)

or heterotrophs (1,189 ASVs) based on the taxonomic categories from PR2 (Tables S1, S2, and S3).

High chlorophyll a (Chl a) concentrations were detected at St3 (5.4 µg/L) and St4 (11.9 µg/L) in March,

and at St4 (7.9 µg/L) in May. Relatively higher POC fluxes (153.4–755.7 mg C m^{-2} d^{-1}) and PON fluxes

(30.9–129.7 mg N m^{-2} d^{-1}) were observed in these sites than in the other sites (89.7–385.3 mg C m^{-2} d^{-1}

and 18.4–64.8 mg N m^{-2} d^{-1}) (Table S4).

Microeukaryote community and its relationship with environmental variables

Diatoms (Bacillariophyta) were the most abundant group at St3 and St4 in March (50.5% ± 15.0% of

the protists 18S reads in suspended particles at SCM; Figs. 1-2). Considering the high Chl a

concentrations (> 5 µg/L) and high relative proportions of diatoms, we defined the St3 and St4 samples

in March as "diatom-dominant" samples (Kuroda et $al.$, 2019). At the genus level (level 7 category of

the PR2 database), two chain-forming diatoms, $Porosira$ and $Odontella$, were the most abundant at

SCM (26.0% ± 8.6% and 13.0% ± 4.5% in suspended particles, 32% ± 7.5% and 10.3% ± 3.4% in

sinking particles, respectively). Both genera have been previously classified as the spring-type diatoms

in the Oyashio region (Chiba et al., 2004). *Porosira* and *Odontella* consistently exhibited the higher

abundance at BBL in both suspended (24.2% ± 0% and 18.0% ± 1.3%, respectively) and sinking

particles (16.8% ± 10.5% and 4.8% ± 1.3% respectively), even at >1000 m depth (Fig. 2). In other

samples (St2 in March, and St1, 2, and 4 in May), the relative abundance of dinoflagellates was

relatively higher (24.6% ± 9.2%) than that in the diatom-dominant samples (10.6% ± 4.7%); these

samples are hereafter referred to as "dinoflagellate-abundant" samples. At St4, a substantial increase in

the relative abundance of dinoflagellates was observed from March (6.2% of suspended particles in the

SCM) to May (32.1%). Dinoflagellates constituted the group of protists with the highest relative

abundance in sinking microeukaryotic communities throughout the water column in May, which

showed a peak in the POC/PON fluxes (Figs. 2 and 3a; Table S1). In other stations, prymnesiophytes

such as *Phaeocystis* were the abundant phytoplankton in the SCM layer (22.4% ± 4.0% in suspended

particles), and they accounted for some proportions in the sinking particles at depth (2.8% ± 1.7%).

However, this was significantly lower (Wilcoxon test, $p < 0.01$) than that in the suspended particles at

the same depth (10.3% ± 6.2%) (Fig. 2). The relative abundance of sinking prymnesiophytes at SCM

(3.8% ± 1.7%) was higher than that at PC (3.2% ± 2.0%) and at BBL (2.5% ± 1.7%). Vector analysis

using vegan's "envfit" function showed that the surface Chl *a*, temperature, and PON flux were

significantly correlated with the variation in the eukaryotic community in the suspended particles ($p <$

0.05) (Fig. 3a).

Microeukaryotic communities differ between suspended and sinking fractions

Non-metric multidimensional scaling (NMDS) showed a clear and significant separation in the ASV

composition of the microeukaryotic community between suspended and sinking particles

(PERMANOVA, $p < 0.05$) (Fig. 3b, Table S5). Phytoplankton richness was lower in the sinking

particles than in the suspended particles (Wilcoxon test, $p < 0.05$; Figs. 3c and S3). Heterotrophic

protists exhibited significantly lower richness (Wilcoxon test, $p < 0.05$) and *Pielou*'s evenness

(Wilcoxon test, $p < 0.05$) in the sinking particles than in the suspended particles (Fig. 3c).

Phytoplankton communities exhibited consistently high similarity index between sinking and

suspended particle types with values between 0.54 and 0.63 at each depth, and between different depths

(the Bray-Curtis similarity: 0.58–0.60 for sinking particles and 0.56–0.59 for suspended particles) (Figs.

4 and S4). Communities of heterotrophic protists also showed higher similarities between sinking and

suspended particles at SCM (0.60 ± 0.12), but the values were lower at PC and BLL (0.40 ± 0.24 and

0.43 ± 0.21, respectively). Furthermore, the communities of sinking heterotrophic protists at depth

showed higher similarity to those in the sinking particles from the upper layer (0.55–0.66) than to those

in the suspended particles from the same layer (0.43–0.48) (Figs. 4b-c). Conversely, suspended

heterotrophic communities at depth showed lower similarity to their upper-layer counterparts (0.39–

0.51) (Figs. 5S and 4c).

The NMDS results also showed that the diatom-dominant samples were significantly different from the

dinoflagellate-abundant samples (PERMANOVA, $p < 0.001$; Fig. 3b, Table S5). Differential abundance

analyses revealed that the protist genera showed varying patterns in relative abundance between the

suspended and sinking particles in the samples under both conditions (Fig. 5). Under diatom-dominant

conditions, most of the diatom genera (except *Actinocyclus*), including the dominant *Porosira* and

Odontella, displayed no significant difference between the suspended and sinking particle fractions (Fig.

5a). Most phototrophic dinoflagellates (16 out of 18 genera) were more abundant in the sinking particles

than in suspended particles, with five genera showing significant differential abundances (Benjamini–

Hochberg (BH) adjusted p-value <0.05). Most genera belonging to Prymnesiophyceae (except

Coccolithus) (7 out of 9 genera), Chrysophyceae (2 out of 2 genera) and Pelagophyceae (4 out of 4

genera) were more enriched in the suspended particles than in sinking particles (Fig. 5a), with five

genera of which showed statistically significant differences (BH adjusted p-value < 0.05). These trends

in phytoplankton distribution across fractions were also evident under dinoflagellate-abundant

conditions (Fig. 5b). Regarding the heterotrophic protists in the diatom-dominant condition, several

Spirotrichea (a class of ciliates) genera and heterotrophic dinoflagellate genera were significantly (BH

adjusted p-value < 0.05) enriched in the sinking particles (Fig. 5c). In contrast, the MAST lineages

(MAST-1A, MAST-2, and MAST-7), which contain heterotrophic nanoflagellates, were primarily

found in the suspended particles (BH adjusted p-value < 0.05). Similar to phytoplankton communities,

heterotrophic protists showed divergent distribution patterns between the suspended and sinking

fractions in the dinoflagellate-abundant samples (Fig. 5d).

Depth-dependent attenuation of sinking particle-associated microeukaryotes

The composition of key protist classes (level 4 category of the PR2 database) changed with depth in the

sinking particles. The relative abundance of phototrophs decreased (from 64.8% ± 14.0% at SCM to

48.6% ± 2.2% at BBL), while that of heterotrophs increased with depth (Fig. 6a). In particular, the

dominant *Porosira* and *Odontella* genera displayed remarkable decreases in relative abundance with

depth, at St4 and St3 in March (Fig. 6b). Conversely, the relative abundances of Spirotrichea and

heterotrophic dinoflagellate increased with depth (Fig. 6a). This depth-dependent trend was more

pronounced for the relative abundances at the family level (level 5 category of the PR2 database) (Fig.

6c).

Discussion

We observed a clear separation of phytoplankton composition between the suspended and sinking

particles (Figs. 3b–c and 5a–b). The lower phytoplankton richness in the sinking particles, particularly

in the SCM (Fig. S3), could be attributed to selective aggregation and sinking processes. Namely, a

subset of the microeukaryotic taxa in the suspended particles is incorporated into the aggregates; this

contributes to the sinking processes. The selectivity for aggregation and sinking could be influenced by

the physical and physiological properties of the phytoplankton (size, shape, stickiness, and secretion of

exopolymers), which can affect the aggregate formation and sedimentation (Jackson, 1990).

Prymnesiophytes and pelagophytes, belonging to pico- and nanoplankton (0.2–20 μm), were enriched

more in the suspended particles (Figs. 2 and 5a-b), possibly due to their small cell size (Liu et al., 2009;

Wetherbee et al., 2015). These picophytoplankton groups were involved in sinking particles in the

surface layer; however, they were rarely detected in the sinking particles in the deep layer, indicating

that they were more prone to disaggregation or decomposition during the sinking process. This

observation is consistent with that of a previous study showing that prymnesiophyte-enriched particles

exported from the euphotic zone are likely to disintegrate into suspended particles in the upper

mesopelagic layer (Duret et al., 2020).

Phototrophic dinoflagellates, which were more abundant in sinking particles and showed stable relative

abundance with depth (Figs. 5a-b and 6a-b), play a significant role in the BCP across various ecosystems

(Guidi *et al.*, 2016; Durkin *et al.*, 2022; Lin *et al.*, 2022); Dinoflagellates are more carbon-dense than

diatoms, as evident from their carbon-to-cell volume ratio (Menden-Deuer and Lessard, 2000); their

sinking can sequester more carbon per volume of particles than that of diatoms. The formation of

massive carbon-rich and thick-walled resting cysts during dinoflagellate blooms makes them more

resistant to degradation, contributing to the intense sedimentation of organic matter (Heiskanen, 1993;

Cohen *et al.*, 2021). Thus, phototrophic dinoflagellates can act as important contributors to BCP in

Oyashio waters. This is supported by our results, which show a positive correlation between

dinoflagellate abundance and POC/PON fluxes (Fig. 3a).

Diatoms play a key role in POC sedimentation in the Oyashio waters, as suggested by previous studies

(Kawakami and Honda, 2007; Kawakami et al., 2015). In this study, centric diatom genera such as

Porosira and *Odontella* were dominant in the diatom-blooming sites near the coastal area (Figs. 1 and

2). These genera belong to microphytoplanktons, which have relatively large cells (20–200 μm)

(Olenina *et al.*, 2006). Diatoms are typically considered important contributors to BCP in various

ecosystems, as heavy silica shells serve as ballast for marine snow and fecal pellets (Belcher *et al.*, 2016;

Tilstone *et al.*, 2017). Previous observations of spring Oyashio water showed that *Porosira glacialis*

was the most dominant species during bloom period, contributing to 25% of the diatom carbon biomass

(Suzuki *et al.*, 2011). The genus *Porosira* is enriched in the sinking particles at BBL (16.8 ± 10.5%,

Fig. 2), indicating that the dominant diatom species would be the major contributors to vertical carbon

export. However, the relative abundance of *Porosira* sharply decreased with depth, especially from PC

(21.7%) to BBL (9.4%) at St3 in March (Figs. 6a-b). This indicates that the dominant diatoms displayed

high attenuation within the sinking communities during sedimentation. These findings are consistent

with that from previous field studies, which reported that the export efficiencies are lower in the diatom-

blooming sites than in the non-blooming sites (Francois *et al.*, 2002; Fischer and Karakaş, 2009; Henson

et al., 2012). Sinking aggregates composed of diatoms exhibit higher porosity, which reduces particle

density and sinking velocity despite high ballast availability, thus increasing their susceptibility to

remineralization (Francois *et al.*, 2002; Bach *et al.*, 2019). Furthermore, the transparent exopolymer

particles (TEPs) produced by diatoms (Engel, 2000), which facilitate aggregation, can reduce the

sinking velocities of aggregates owing to their low density (Mari et al., 2017). Indeed, the contribution

to vertical carbon export varies among diatom genera or species owing to differences in cell size, shape,

degree of silicification, and production of TEPs (Tréguer *et al.*, 2018).

The heterotrophic protist community structure also differed significantly between the sinking and

suspended particles (Figs. 3b-c and 5c-d). Lower richness and evenness of the heterotrophic protists in

the sinking particles than those in the suspended particles indicate that specific protist groups were

enriched in the sinking particles. The prominent heterotrophic protist groups enriched in sinking

particles were Spirotrichea and heterotrophic dinoflagellates (Figs. 5c-d). These protists are active

grazers of phytoplankton, including chain-forming diatoms (Montagnes and Lessard, 1999; Sherr et al.,

2013), suggesting that they contribute to the decomposition and transformation of diatom-enriched

aggregates in Oyashio waters.

The MAST groups of heterotrophic protists were more abundant in the suspended than in the sinking

particles. In addition, they were more abundant in the surface layers than in the deeper layers, which is

consistent with previous observations (Obiol et al., 2021). The MAST groups include consumers of

heterotrophic bacteria and picoeukaryotes in the water column (Lin et al., 2012; Orsi et al., 2018) and

parasites of diatoms in sediments (e.g., MAST-6, (Rodríguez-Martínez *et al.*, 2020)). Considering their

low abundance in sinking particles, it is less likely that the MAST groups in suspended particles

collected from the deeper layers of the Oyashio region were those transported from the upper layer.

MAST groups prefer a free-living, suspended lifestyle and consume picoeukaryotes and bacteria. The

distinct occurrences of heterotrophs in sinking and suspended particles reflects the differences in

lifestyle and feeding habits (Worden *et al.*, 2015).

High similarities in the sinking microeukaryote communities between the surface and deep layers were

recorded; this included both phytoplankton and heterotrophic protists (Fig. 4c). Heterotrophic protist

communities of sinking particles displayed more remarkable similarities across different depths (i.e.,

SCM, PC, and BBL) than did suspended communities in ambient waters. This suggests strong vertical

connectivity of heterotrophic protist communities with sinking particles, mediated by the association of

specific groups of heterotrophic protists (Spirotrichea and heterotrophic dinoflagellates).

In BBL samples collected approximately 10 m above the seafloor, the resuspension of bottom sediments,

indicated by high turbidity, could introduce benthic communities from the sediments into the sinking

and suspended particles collected with MSC. These benthic communities harbor highly diverse and

abundant indigenous eukaryotes that are unique to the environment (Cordier et al., 2022). However, the

BBL samples did not show an increased richness of microeukaryotes compared to that in the upper

samples (Fig. S4). The proportion of ASVs unique to the BBL samples was consistently low (0.3–2%).

These results suggest that resuspension of the benthic microeukaryotic community was minimal in the

studied region. Alternatively, the resuspended microeukaryotic communities mostly originate from

recently settled fresh particles (Almroth-Rosell et al., 2011; Rodil et al., 2020). In either case, our data

suggested that heterotrophs in the sinking particles mainly originate from the surface layers along with

the formation of sinking aggregates rather than through active colonization from ambient water.

Both the phototrophs and heterotrophs associated with sinking particles in the deeper layers were

derived from the surface; they displayed distinct depth-dependent attenuation patterns (Fig. 6). The

contribution of phototrophs decreased with depth, whereas that of the heterotrophic protists increased.

The increased abundance of heterotrophic protists relative to phototrophs with increasing depth could

reflect the consumption of phototrophs during the transit of sinking particles in the water column.

Therefore, heterotrophic protists such as Spirotrichea and heterotrophic dinoflagellates could contribute

to the decomposition of sinking particles by consuming phototrophic cells. The depth-dependent

attenuation of the relative abundance of phototrophs was more prominent in diatom-dominated

situations than in dinoflagellate-abundant situations, implying that diatoms are more rapidly consumed

by heterotrophic protists. These results indicate that the magnitude of sinking POC flux attenuation is

influenced by dominant phototrophic microeukaryotes (diatoms vs. dinoflagellates) and their

consumers (Spirotrichea and heterotrophic dinoflagellates), which comprise the microbial consortia of

sinking particles. Future studies are needed to elucidate the role of prey-predator interactions of

microeukaryotes, which are potentially highly selective in feeding (Lima-Mendez et al., 2015), in

regulating the POC flux attenuation with depth.

Conclusion

The use of MSC combined with DNA metabarcoding to investigate spring Oyashio waters allowed us

to characterize the microeukaryotic communities that contribute to the biological carbon sequestration

processes. This study provides profound insights into the composition and dynamics of microeukaryotic

communities within sinking and suspended particles in the Oyashio region of the western subarctic

Pacific. We present the distinct structures of the microeukaryotic communities in different particle types

and at varying depths. In addition, these findings highlight the differential patterns of depth-dependent

attenuation among different taxonomic groups; prymnesiophytes and diatoms were more rapidly

attenuated with depth than phototrophic dinoflagellates. We found strong vertical connectivity between

some heterotrophic protists (Spirotrichea and heterotrophic dinoflagellates) associated with sinking

particles, suggesting that they play a crucial role in the regulation of sinking POC fluxes through the

consumption of major phototrophic microeukaryotes associated with sinking particles.

Materials and methods

Cruises and sampling strategy

Sampling was conducted at four stations (Fig.1) during the KS-21-4 (March 11–21, 2021) and KS-21-

7 (May 3–11, 2021) cruises of the R/V *Shinsei-Maru* (JAMSTEC) in the Oyashio region off Hokkaido,

Japan. Samples for DNA analysis were collected using MSCs at three depths: the subsurface

chlorophyll maximum (SCM), 10 m below the pycnocline (PC), and 10 m above the bottom (BBL).

Detailed information on the sampling depths and environmental factors is presented in Table S4.

Particles of suspended and sinking particles were sampled from seawater in the upper and base parts of

MSC, respectively, then filtered using a 0.8 µm pore-size cellulose acetate membrane (47 mm diameter,

Millipore) with a gentle vacuum (<0.013MPa). The filters were stored in 1.5-mL cryotubes previously

filled with 600 µL of buffer RLT Plus (Qiagen), 6 µL of 2-mercaptoethanol (Sigma-Aldrich), and 0.2

g of glass beads; the samples were flash-frozen in liquid N_2 and stored in a deep-freezer until analysis

on land. The suspended fraction defined as the particles remaining in the upper part of the MSC, and

the sinking fraction referred to the particles that sunk to the base plate of the MSC after being placed

on the deck for 2 h. For detailed descriptions of MSC, refer to earlier reports (Duret *et al.*, 2019; Baumas

et al., 2021). The sinking fraction contained some suspended particles; therefore, after detaching the

upper part of the MSC, the base of the MSC was allowed to settle for another 10 min, and approximately

two-thirds of the seawater in the base plate of the MSC was siphoned from the top to remove as many

suspended particles as possible.

Measurements of environmental variables

Vertical profiles of temperature, dissolved oxygen (DO), salinity, Chl *a* fluorescence,

photosynthetically available radiation (PAR), and turbidity were measured at each site using an SBE

911-plus CTD system (Seabird Electronics, Bellevue, WA, USA) (Fig. S1). Seawater samples for Chl

a and macronutrient analysis were collected using Niskin bottles attached to a CTD-Carousel multiple-

sampler system. Samples for Chl *a* concentration measurement were collected on GF/F filters and

measured with a fluorometer (10-AU, Turner Designs) using the acidification method (Holm-Hansen

et al., 1965). The concentrations of macronutrients ($Si(OH)_4$, NO_2, NO_3, NH_4, and PO_4) were

determined using a flow injection analyzer (AACSIII, Bran+Luebbe). The detection limits of the

macronutrient measurements were defined as three times the deviation of the repeated measurements

from the blank. Detailed procedures for Chl *a* and macronutrient analyses have been described by

Fukuda *et al.* (2016). Samples for POC and PON concentrations were collected from the sinking fraction

of MSC and analyzed using an elemental analyzer (Elemental Analyzer-IRMS; FLASH 2000, Thermo

Fisher Scientific). The POC and PON sinking fluxes were calculated using the method described by

Yamada et al. (submitted).

DNA extraction and 18S rRNA gene amplification and sequencing

AllPrep DNA/RNA Mini Kits (Qiagen) were used for DNA extraction. The sample vials were agitated

thrice at 2,500 rpm for 50 s using a homogenizer (μT-12, TAITEC) before proceeding with the

manufacturer's protocol (BBL samples at St1 and St2 in May failed to yield sufficient DNA for

downstream analysis). For each sample, three PCR amplifications targeting the V4 region of the 18S

rRNA gene were performed using the following primer pairs: the forward E572F (5'-

CYGCGGTAATTCCAGCTC-3') and the reverse E1009R (5'-AYGGTATCTRATCRTCTTYG-3';

436 bp), as designed previously (Comeau et al., 2011). The PCR mixtures were prepared in a 25-μL

reaction volume containing 1 x KAPA HiFi HotStart ReadyMix, 1 μM each of forward and reverse

primers, and 2.5 ng of template DNA. The PCR consisted of a pre-denaturation step at 98°C for 30 s,

followed by 30 cycles of denaturation at 98°C for 10 s, annealing at 61°C for 30 s, and extension at

72°C for 30 s, and a final extension at 72°C for 5 min. The triplicate PCR products were mixed after

purification using Agencourt AMPure XP beads (Beckman Coulter). The quality and quantity of the

348 amplicons were checked using the D1000 ScreenTape Assay kit read on a 2100 TapeStation system

349 (Agilent Technologies) and a Qubit dsDNA BS assay kit (Invitrogen), respectively. Libraries were

350 prepared using the Nextera XT index kit V2 and sequenced on an Illumina MiSeq platform using a 300

351 bp paired-end sequencing. All raw sequencing data are available at the DDBJ Sequence Read Archive

352 under accession number of DRA015710.

354 **Analysis of sequencing data**

355 Raw sequencing data were preprocessed using QIIME2 (Bolyen *et al.*, 2019) (version 2021.11).

356 Trimming of low-quality reads and primer sequences, merging of paired-end reads, dereplication,

357 chimera removal, and ASV inference were performed using denoising algorithms in DADA2 (Callahan

358 *et al.*, 2016). A naïve Bayes classifier trained (Bokulich *et al.*, 2018) on the PR2 database (Guillou *et*

359 *al.*, 2013) (v 4.14.1) with the E572F/E1009R primer set was used to assign taxonomy to the ASV

360 sequences. The feature table of read counts and taxonomy assigned to ASVs was exported by

361 eliminating singleton ASVs. The ASVs that appeared only at the BBL samples were removed to reduce

362 the potential contamination by indigenous benthic organisms.

363 Potential multicellular eukaryotic ASVs, such as Embryophyceae, Fungi, Metazoa, and Rhodophyta,

364 were excluded; the unicellular eukaryotic (protists) ASV taxa were divided into photosynthetic and

365 heterotrophic groups (Table S3) (Simon *et al.*, 2009; Durkin *et al.*, 2022). We manually controlled the

366 detailed trophic groups of dinoflagellates based on existing literature; however, some of the

367 heterotrophic dinoflagellate sequences could have been erroneously assigned to photosynthetic groups

368 (Table S4) or vice versa. In addition, owing to the copy number variation in the 18S rRNA genes, the

369 relative abundance of the ribosomal marker genes does not correlate with cell abundance (Zhu et al.,

370 2005). This potentially affects the microeukaryote composition analysis results; however, it does not

371 affect the comparison of specific taxa between the suspended and sinking particle fractions.

373 **ASV-based diversity analyses**

Statistical analyses were performed using R v.4.4.1 and Python v.3.9.13. Samples were rarefied to

account for differences in library size, and rarefaction curves were used to investigate the degree of

sample saturation by calling the 'rrarefy' function in the vegan v.2.6.4 package (Oksanen et al., 2022).

NMDS analysis of the dissimilarity in the microeukaryotic communities between these two particle

fractions, and the relationship between the suspended microeukaryote composition and environmental

variables (NH_4 was omitted because 11 out of 16 samples had concentrations below the detection limit,

Table S4) were performed based on the Bray-Curtis dissimilarities with the rarefied abundance table.

ANOSIM with 9999 permutations was used to statistically test the significant dissimilarity between the

particle fractions (with the it'envfit' function in vegan). Alpha diversity indices (*Pielou*'s evenness and

richness) of phytoplankton and heterotrophs in each sample were evaluated by rarefying the read

numbers. The difference in alpha diversity between these two particle fractions was assessed using the

Wilcoxon signed-rank exact test to determine statistical significance.

Taxon-based differential abundance analysis

Differential analysis of protist genera between the two particle fractions was performed using DEseq2

v1.38.3 (Love et al., 2014). The samples were categorized into two conditions based on the NMDS

analysis (Fig. 3b): diatom-dominant regions, including St3 and St4 in March, and dinoflagellate-

abundant regions with a relatively higher abundance of dinoflagellates, including St2 in March and St1,

St2, and St4 in May. Protist genera with $\log_2$(fold change) (FC) > 1.5 or < -1.5 and an adjusted *p*-value

< 0.05 determined using the Benjamini-Hochberg method were regarded as "significant".

Acknowledgments

We gratefully acknowledge the captain, officers, and crew of the R/V Shinsei-Maru for their great

assistance during the cruise. We are thankful to Junyi Wu and Ruixuan Zhang for support with data

analysis and helpful discussions. This study was funded by JSPS KAKENHI (nos. 19H05667,

22H00384, and 22H02420). Computational work was performed at the SuperComputer System,

Institute for Chemical Research, Kyoto University.

References

Agusti, S., González-Gordillo, J.I., Vaqué, D., Estrada, M., Cerezo, M.I., Salazar, G., et al. (2015)

Ubiquitous healthy diatoms in the deep sea confirm deep carbon injection by the biological

pump. *Nat Commun* **6**: 7608.

Anderson, T.R. and Tang, K.W. (2010) Carbon cycling and POC turnover in the mesopelagic zone of

the ocean: Insights from a simple model. *Deep Sea Res Part II Top Stud Oceanogr* **57**: 1581–

1592.

Bach, L.T., Stange, P., Taucher, J., Achterberg, E.P., Algueró-Muñiz, M., Horn, H., et al. (2019) The

Influence of Plankton Community Structure on Sinking Velocity and Remineralization Rate

of Marine Aggregates. *Glob Biogeochem Cycles* **33**: 971–994.

Bauerfeind, E., Nöthig, E.-M., Beszczynska, A., Fahl, K., Kaleschke, L., Kreker, K., et al. (2009)

Particle sedimentation patterns in the eastern Fram Strait during 2000–2005: Results from the

Arctic long-term observatory HAUSGARTEN. *Deep Sea Res Part Oceanogr Res Pap* **56**:

1471–1487.

Baumas, C.M.J., Le Moigne, F.A.C., Garel, M., Bhairy, N., Guasco, S., Riou, V., et al. (2021)

Mesopelagic microbial carbon production correlates with diversity across different marine

particle fractions. *ISME J* **15**: 1695–1708.

Belcher, A., Iversen, M., Manno, C., Henson, S.A., Tarling, G.A., and Sanders, R. (2016) The role of

particle associated microbes in remineralization of fecal pellets in the upper mesopelagic of

the Scotia Sea, Antarctica. *Limnol Oceanogr* **61**: 1049–1064.

Boeuf, D., Edwards, B.R., Eppley, J.M., Hu, S.K., Poff, K.E., Romano, A.E., et al. (2019) Biological

composition and microbial dynamics of sinking particulate organic matter at abyssal depths in

the oligotrophic open ocean. *Proc Natl Acad Sci* **116**: 11824–11832.

Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., et al. (2018)

Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-

feature-classifier plugin. *Microbiome* **6**: 90.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., et al. (2019)

Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.

Nat Biotechnol **37**: 852–857.

Boyd, P.W., Claustre, H., Levy, M., Siegel, D.A., and Weber, T. (2019) Multi-faceted particle pumps

drive carbon sequestration in the ocean. *Nature* **568**: 327–335.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P. (2016)

DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**:

581–583.

Chiba, S., Ono, T., Tadokoro, K., Midorikawa, T., and Saino, T. (2004) Increased Stratification and

Decreased Lower Trophic Level Productivity in the Oyashio Region of the North Pacific: A

30-Year Retrospective Study. *J Oceanogr* **60**: 149–162.

Cohen, N.R., McIlvin, M.R., Moran, D.M., Held, N.A., Saunders, J.K., Hawco, N.J., et al. (2021)

Dinoflagellates alter their carbon and nutrient metabolic strategies across environmental

gradients in the central Pacific Ocean. *Nat Microbiol* **6**: 173–186.

Comeau, A.M., Li, W.K.W., Tremblay, J.-É., Carmack, E.C., and Lovejoy, C. (2011) Arctic Ocean

Microbial Community Structure before and after the 2007 Record Sea Ice Minimum. *PLOS*

ONE **6**: e27492.

Cordier, T., Angeles, I.B., Henry, N., Lejzerowicz, F., Berney, C., Morard, R., et al. (2022) Patterns

of eukaryotic diversity from the surface to the deep-ocean sediment. *Sci Adv* **8**: eabj9309.

Duret, M.T., Lampitt, R.S., and Lam, P. (2020) Eukaryotic influence on the oceanic biological carbon

pump in the Scotia Sea as revealed by 18S rRNA gene sequencing of suspended and sinking

particles. *Limnol Oceanogr* **65**: S49–S70.

Duret, M.T., Lampitt, R.S., and Lam, P. (2019) Prokaryotic niche partitioning between suspended and

sinking marine particles. *Environ Microbiol Rep* **11**: 386–400.

Durkin, C.A., Cetinić, I., Estapa, M., Ljubešić, Z., Mucko, M., Neeley, A., and Omand, M. (2022)

Tracing the path of carbon export in the ocean though DNA sequencing of individual sinking

particles. *ISME J* **16**: 1896–1906.

Engel, A. (2000) The role of transparent exopolymer particles (TEP) in the increase in apparent

particle stickiness (α) during the decline of a diatom bloom. *J Plankton Res* **22**: 485–497.

Fischer, G. and Karakaş, G. (2009) Sinking rates and ballast composition of particles in the Atlantic

Ocean: implications for the organic carbon fluxes to the deep ocean. *Biogeosciences* **6**: 85–

102.

Francois, R., Honjo, S., Krishfield, R., and Manganini, S. (2002) Factors controlling the flux of

organic carbon to the bathypelagic zone of the ocean. *Glob Biogeochem Cycles* **16**: 34-1-34–

20.

Fukuda, H., Katayama, R., Yang, Y., Takasu, H., Nishibe, Y., Tsuda, A., and Nagata, T. (2016)

Nutrient status of Otsuchi Bay (northeastern Japan) following the 2011 off the Pacific coast of

Tohoku Earthquake. *J Oceanogr* **72**: 39–52.

Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., et al. (2016) Plankton

networks driving carbon export in the oligotrophic ocean. *Nature* **532**: 465–470.

Guidi, L., Stemmann, L., Jackson, G.A., Ibanez, F., Claustre, H., Legendre, L., et al. (2009) Effects of

phytoplankton community on production, size, and export of large aggregates: A world-ocean

analysis. *Limnol Oceanogr* **54**: 1951–1963.

Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., et al. (2013) The Protist

Ribosomal Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA

sequences with curated taxonomy. *Nucleic Acids Res* **41**: D597-604.

Heiskanen, A.-S. (1993) Mass encystment and sinking of dinoflagellates during a spring bloom. *Mar*

Biol **116**: 161–167.

Henson, S., Lampitt, R., and Johns, D. (2012) Variability in phytoplankton community structure in

response to the North Atlantic Oscillation and implications for organic carbon flux. *Limnol*

Oceanogr **57**: 1591–1601.

Henson, S.A., Laufkötter, C., Leung, S., Giering, S.L.C., Palevsky, H.I., and Cavan, E.L. (2022)

Uncertain response of ocean biological carbon export in a changing world. *Nat Geosci* **15**:

248–254.

Holm-Hansen, O., Lorenzen, C.J., Holmes, R.W., and Strickland, J.D.H. (1965) Fluorometric

Determination of Chlorophyll. *ICES J Mar Sci* **30**: 3–15.

Jackson, G.A. (1990) A model of the formation of marine algal flocs by physical coagulation

processes. *Deep Sea Res Part Oceanogr Res Pap* **37**: 1197–1211.

Kawakami, H. and Honda, M.C. (2007) Time-series observation of POC fluxes estimated from 234Th

in the northwestern North Pacific. *Deep Sea Res Part Oceanogr Res Pap* **54**: 1070–1090.

Kawakami, H., Honda, M.C., Matsumoto, K., Wakita, M., Kitamura, M., Fujiki, T., and Watanabe, S.

(2015) POC fluxes estimated from 234Th in late spring–early summer in the western

subarctic North Pacific. *J Oceanogr* **71**: 311–324.

Kuroda, H., Toya, Y., Watanabe, T., Nishioka, J., Hasegawa, D., Taniuchi, Y., and Kuwata, A. (2019)

Influence of Coastal Oyashio water on massive spring diatom blooms in the Oyashio area of

the North Pacific Ocean. *Prog Oceanogr* **175**: 328–344.

Lampitt, R.S., Wishner, K.F., Turley, C.M., and Angel, M.V. (1993) Marine snow studies in the

Northeast Atlantic Ocean: distribution, composition and role as a food source for migrating

plankton. *Mar Biol* **116**: 689–702.

Lin, S., Li, T., Yuan, H., Li, H., Yu, L., Zhuang, Y., et al. (2022) Sediment Trap Study Reveals

Dominant Contribution of Metazoans and Dinoflagellates to Carbon Export and Dynamic

Impacts of Microbes in a Subtropical Marginal Sea. *J Geophys Res Biogeosciences* **127**:

e2021JG006695.

Lin, Y.-C., Campbell, T., Chung, C.-C., Gong, G.-C., Chiang, K.-P., and Worden, A.Z. (2012)

Distribution Patterns and Phylogeny of Marine Stramenopiles in the North Pacific Ocean.

Appl Environ Microbiol **78**: 3387–3399.

Liu, H., Probert, I., Uitz, J., Claustre, H., Aris-Brosou, S., Frada, M., et al. (2009) Extreme diversity in

noncalcifying haptophytes explains a major pigment paradox in open oceans. *Proc Natl Acad*

Sci **106**: 12803–12808.

Love, M.I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Mari, X., Passow, U., Migon, C., Burd, A.B., and Legendre, L. (2017) Transparent exopolymer

particles: Effects on carbon cycling in the ocean. *Prog Oceanogr* **151**: 13–37.

Menden-Deuer, S. and Lessard, E.J. (2000) Carbon to volume relationships for dinoflagellates,

diatoms, and other protist plankton. *Limnol Oceanogr* **45**: 569–579.

Montagnes, D.J.S. and Lessard, E.J. (1999) Population dynamics of the marine planktonic ciliate

Strombidinopsis multiauris: its potential to control phytoplankton blooms. *Aquat Microb Ecol*

20: 167–181.

Obiol, A., Muhovic, I., and Massana, R. (2021) Oceanic heterotrophic flagellates are dominated by a

few widespread taxa. *Limnol Oceanogr* **66**: 4240–4253.

Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., et al. (2022)

vegan: Community Ecology Package.

Olenina, Susanna, Hajdu, Lars, Edler, Agneta, Andersson, Norbert Wasmund, Susanne, Busch, et al.

(2006) Biovolumes and Size-Classes of Phytoplankton in the Baltic Sea.

Orsi, W.D., Wilken, S., Del Campo, J., Heger, T., James, E., Richards, T.A., et al. (2018) Identifying

protist consumers of photosynthetic picoeukaryotes in the surface ocean using stable isotope

probing. *Environ Microbiol* **20**: 815–827.

Preston, C.M., Durkin, C.A., and Yamahara, K.M. (2020) DNA metabarcoding reveals organisms

contributing to particulate matter flux to abyssal depths in the North East Pacific ocean. *Deep*

Sea Res Part II Top Stud Oceanogr **173**: 104708.

Richardson, T.L. (2019) Mechanisms and Pathways of Small-Phytoplankton Export from the Surface

Ocean. *Annu Rev Mar Sci* **11**: 57–74.

Rodríguez-Martínez, R., Leonard, G., Milner, D.S., Sudek, S., Conway, M., Moore, K., et al. (2020)

Controlled sampling of ribosomally active protistan diversity in sediment-surface layers

identifies putative players in the marine carbon sink. *ISME J* **14**: 984–998.

Scharek, R., Latasa, M., Karl, D.M., and Bidigare, R.R. (1999) Temporal variations in diatom

abundance and downward vertical flux in the oligotrophic North Pacific gyre. *Deep Sea Res*

Part Oceanogr Res Pap **46**: 1051–1075.

Sherr, E.B., Sherr, B.F., and Ross, C. (2013) Microzooplankton grazing impact in the Bering Sea

during spring sea ice conditions. *Deep Sea Res Part II Top Stud Oceanogr* **94**: 57–67.

Simon, N., Cras, A.-L., Foulon, E., and Lemée, R. (2009) Diversity and evolution of marine

phytoplankton. *C R Biol* **332**: 159–170.

Suzuki, K., Kuwata, A., Yoshie, N., Shibata, A., Kawanobe, K., and Saito, H. (2011) Population

dynamics of phytoplankton, heterotrophic bacteria, and viruses during the spring bloom in the

western subarctic Pacific. *Deep Sea Res Part Oceanogr Res Pap* **58**: 575–589.

Takahashi, T., Sutherland, S.C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., et al. (2002)

Global sea–air CO2 flux based on climatological surface ocean pCO2, and seasonal biological

and temperature effects. *Deep Sea Res Part II Top Stud Oceanogr* **49**: 1601–1622.

Tilstone, G.H., Lange, P.K., Misra, A., Brewin, R.J.W., and Cain, T. (2017) Micro-phytoplankton

photosynthesis, primary production and potential export production in the Atlantic Ocean.

Prog Oceanogr **158**: 109–129.

Tréguer, P., Bowler, C., Moriceau, B., Dutkiewicz, S., Gehlen, M., Aumont, O., et al. (2018)

Influence of diatom diversity on the ocean biological carbon pump. *Nat Geosci* **11**: 27–37.

Trudnowska, E., Lacour, L., Ardyna, M., Rogge, A., Irisson, J.O., Waite, A.M., et al. (2021) Marine

snow morphology illuminates the evolution of phytoplankton blooms and determines their

subsequent vertical export. *Nat Commun* **12**: 2816.

Turner, J.T. (2015) Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's biological

pump. *Prog Oceanogr* **130**: 205–248.

Wetherbee, R., Gornik, S.G., Grant, B., and Waller, R.F. (2015) Andersenia, a genus of filamentous,

sand-dwelling Pelagophyceae from southeastern Australia. *Phycologia* **54**: 35–48.

Worden, A.Z., Follows, M.J., Giovannoni, S.J., Wilken, S., Zimmerman, A.E., and Keeling, P.J.

(2015) Rethinking the marine carbon cycle: Factoring in the multifarious lifestyles of

microbes. *Science* **347**: 1257594.

Yamada, Y., Akiko, E., Hideki, F., Otosaka, S., Mitarai, S., and Nagata, T. (2023) Roles of

extracellular polymeric substances in partitioning suspended and sinking particles in the upper

oceans. *Unpubl Manuscr*.

Zhang, C., Dang, H., Azam, F., Benner, R., Legendre, L., Passow, U., et al. (2018) Evolving

paradigms in biological carbon cycling in the ocean. *Natl Sci Rev* **5**: 481–499.

Zhu, F., Massana, R., Not, F., Marie, D., and Vaulot, D. (2005) Mapping of picoeucaryotes in marine

ecosystems with quantitative PCR of the 18S rRNA gene. *FEMS Microbiol Ecol* **52**: 79–92.

Figures

Fig. 1 Location of sampling sites drawing by Ocean Data View.

Fig. 2 Overview of microeukaryote composition in Oyashio waters. The relative abundance of the
protists genus (level 7 category of the PR2 database, relative abundance ≥ 1%), colored based on major
divisions or class information at suspended and sinking particle samples across stations and depths; X-
axis indicates particle fractions of each depth: SCM_SS=Suspended particle at SCM, PC_SK=Sinking
Particle at PC.

Fig. 3 Comparison of microeukaryotic communities detected in different particle fractions. (a) NMDS plot of microeukaryote protists in suspended particles, including 10 fitted environmental variables: total nitrogen (NO2+NO3), phosphate (PO4), silicate (Si(OH)4), temperature (Temp), dissolved oxygen (DO), salinity (Sal), chlorophyll concentration (Chl a), and PON and POC fluxes.The variables with p < 0.05 were marked with red; (b) MNDS of Bray-Curtis dissimilarities of rarified read counts. Particle fractions were sinking particles (midnight blue) and suspended particles (golden yellow). Circle points represent the diatom-dominant condition, and triangle points represent the dinoflagellate-abundant condition. Letters indicate depth: S=SCM, P=PC, B=BBL; (c) Evenness and richness indices of different functional groups pooled by particle fractions. Stars indicate significant differences in indices between these two particle fractions, with * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (Wilcoxon test).

Fig. 4 Comparison of phytoplankton and heterotrophic protists communities. (a) Boxplot showing the Bray-Curtis similarity between suspended and sinking phytoplankton and heterotrophs across depth. (b) Boxplot showing the Bray-Curtis similarity of sinking phytoplankton and heterotrophic communities between different depths. Grey points in the plots represent the mean value of the observations. Outliers were removed from the boxplots. (c) Comparative schematic diagram of similarity of phytoplankton (green) and heterotrophic protists (purple) communities. Values indicate the Bray-Curtis similarity index.

Fig. 5 Differential abundance of functional groups at different conditions. The volcano plot depicts the differential abundance of phytoplankton and heterotrophs at diatom dominant condition (a&c) and dinoflagellate abundant condition (b&d). Log2 (fold change) (FC) is plotted against -log10 (BH adjusted p-value) mostly at the genus level. Each dot represents a genus, color-coded based on their division or class information. For those unassigned at the genus level, higher taxonomic information was retained up to the family level. Taxa unassigned at the higher than family level was removed. Significantly distinct genera between suspended and sinking particles were defined based on the absolute of log2 (FC) > 1.5 and BH adjusted *p*-value < 0.05. Log2 (FC) > 0 represents taxa enriched in sinking particles (triangle), while log2 (FC) < 0 indicates taxa enriched in suspended particles (circle).

610 Fig. 6 The vertical succession of sinking particle-associated (a) major microeukaryotes composition at

611 the class level, then further separated by phytoplankton at the genus level (relative abundance $\geq$ 5%) (b)

612 and heterotrophic protists at the family level (relative abundance $\geq$ 5%) (c) Stations without bottom

613 layers (St1 and St2 in May) were removed from the analysis.

614