A fourth locus in the *Plasmodium falciparum* genome associated with sickle haemoglobin

Authors

William L. Hamilton[1,2,3] *, Gavin Band[4], Yaw Aniweh[5], Victor Asoala[6], Eleanor Drury[1], Kirk A. Rockett[4], Gordon A. Awandare[5], Dominic P. Kwiatkowski[1], Lucas N. Amenga-Etego[5] *

Author affiliations

1. Wellcome Sanger Institute, Wellcome Trust Genome Campus, Hinxton, CB10 1RQ, United Kingdom
2. University of Cambridge, Department of Medicine, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, United Kingdom
3. Cambridge University Hospitals NHS Foundation Trust, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, United Kingdom
4. Wellcome Centre for Human Genetics, Roosevelt Dr, Oxford OX3 7BN, United Kingdom
5. West African Centre for Cell Biology of Infectious Pathogens (WACCBIP), University of Ghana, Legon, Accra, Ghana
6. Navrongo Health Research Centre (NHRC), Ghana Health Service, Navrongo, Upper East Region, Ghana

* Corresponding authors

Dr. William L. Hamilton: wh2@sanger.ac.uk

Dr. Lucas N. Amenga-Etego: lamenga-etego@ug.edu.gh

Key words

Malaria, *P. falciparum*, sickle, haemoglobin, HbS, GWAS, genome-wide association, host-parasite interaction

Author contributions

Conceptualization: LNA, WLH, DPK

Investigation: WLH, GB, ED, LNA

Formal analysis: WLH, GB, LNA

Resources: KAR

Data Curation: WLH, LNA

Writing – original draft preparation: WLH, GB, LNA

Writing – review and editing: WLH, GB, LNA, KAR, YA, VA

Visualization: WLH, GB

Supervision: LNA, DPK

Project administration: LNA, YA, VA

Funding acquisition: DPK, LNA, GAA

1 Abstract

Heterozygosity for sickle haemoglobin (HbS) confers protection against severe malaria caused by the

parasite *Plasmodium falciparum.* Recent work suggests that this protective effect can depend on the

parasite genotype. Here, we performed a genome-wide association analysis of *P. falciparum* against

human β-globin genotypes in a sample of 1,368 people with mild malaria in northern Ghana. We

replicated the previously identified associations with HbS at two parasite loci. However, a newly

identified locus within the serine/threonine kinase *FIKK4.2*, which we term *Pfsa4*, was also associated

with HbS; this finding replicated in a published sample from Mali. The *Pfsa1-4* mutations vary widely

in frequencies across Africa, are absent from Asia, and are highly correlated with each-other across

multiple populations. We found no strong associations with haemoglobin C. These findings add new

complexity to the emerging picture of association between human and co-evolving malaria parasite

genomes, suggesting new avenues for functional exploration.

1 Introduction

The World Health Organization (WHO) estimates that there were 247 million malaria cases and 619,000 deaths in 2021 [1]. 95% of cases and 96% of deaths occurred in Africa, and 76% of deaths were in children under 5 years old. Malaria has taken a major toll on human health for millennia, particularly for young children prior to the development of protective (though incomplete) immunity. Consequently, malaria has exerted a strong evolutionary selective pressure on the human genome [2]. Malaria can cause a wide spectrum of clinical disease, from asymptomatic parasitaemia to severe malaria syndromes and death. Understanding how human and parasite genetic variation interact with each-other and contribute to malaria disease severity is a key goal of malaria research.

Genetic variation at dozens of human loci have been proposed to provide protection against malaria [3–18]; although, some of these have not replicated in larger multi-centre studies [19,20]. It is likely that many host polymorphisms contribute to disease susceptibility, most with small effects [21]. Approximately 7% of the world's population are thought to be carriers for inherited haemoglobin disorders, making them the most common human monogenic diseases [22]. Natural variation in the human haemoglobin genes was hypothesised to provide protection against malaria since at least the 1950s [7–9], given the central role of red blood cells (RBCs) in the *Plasmodium* life cycle, and the overlay of certain haemoglobin variants with malaria prevalence. However, the full scope of genetic diversity in human haemoglobin, and how this relates to malaria disease severity, is highly complex, and remains incompletely understood.

The dominant form of adult haemoglobin (HbA) comprises a tetramer of proteins transcribed from two α-globin genes and two β-globin genes ($\alpha_2\beta_2$), responsible for oxygen transportation within RBCs. The α-globin and β-globin genes are located within the α- and β-globin loci on chromosomes 16 and 11, respectively. Of the haemoglobinopathies (altered haemoglobin makeup and/or structure), haemoglobin S (HbS, or sickle), haemoglobin C (HbC) and haemoglobin E (HbE) occur at significant population frequencies with distinct geographic distributions [23]. The sickle β-globin allele (β^S) is a single point non-synonymous polymorphism changing Glutamic acid (coded by GTG) to Valine (GAG) at the sixth codon; this remains the strongest known association with protection against malaria in the human genome [19,20], maintaining high frequencies in populations exposed to malaria, primarily in Africa [24]. Homozygosity for β^S is the most common cause of sickle cell disease (SCD) in Africa. The central pathophysiology of SCD involves polymerisation of deoxygenated HbS, causing RBCs to become inflexible and misshapen, leading to haemolysis, endothelial dysfunction, inflammation and occlusion within the microcirculation [25]. Malaria protection is driven by the heterozygous state (HbAS) and fixation of the sickle allele cannot be attained due to the deleterious nature of the

homozygote state (HbSS). The mechanisms by which heterozygous HbS provides malaria protection

are still not fully understood, but several have been proposed (summarised in [26]).

HbS is present throughout sub-Saharan Africa to varying degrees [23,24], but a second haemoglobin

type, HbC, occurs at high frequency only in central west Africa [27] (**Supplementary Figure 1**). HbC

arises due to an alternative single-point polymorphism in the sixth codon of β-globin ($β^C$) at the base

pair adjacent to the $β^S$ polymorphism, leading to a switch from Glutamatic acid to Lysine. In west

Africa, therefore, β-globin genotypes can include $β^{A/A}$ (HbAA wild-type), $β^{A/S}$ (HbAS sickle

heterozygote), $β^{S/S}$ (HbSS sickle homozygote), $β^{A/C}$ (HbAC heterozygote), $β^{C/C}$ (HbCC homozygote), and

$β^{S/C}$ (HbSC compound heterozygote). Many other polymorphisms in the β-globin gene have been

described, including those that change gene expression causing β-thalassaemias [23,28], but these are

less common in west Africa.

How variation in human and malaria parasite genomes interact with each-other and give rise to

disease phenotypes remains an active area of research. A recent study of around 4,000 samples from

children with severe malaria from Kenya and The Gambia identified three parasite loci that were

associated with sickle haemoglobin, termed *P. falciparum* sickle-associated loci or *Pfsa1*, *Pfsa2* and

Pfsa3 [29]. *Pfsa1* and *Pfsa2* are on different loci on chromosome 2, while *Pfsa3* is located in a region

of suspected structural variation on chromosome 11. The functional mechanism(s) underpinning these

associations is unclear. Here, we have tested for associations between the human β-globin variants $β^S$

and $β^C$, and genome-wide variation in *P. falciparum* parasites, in a sample of mild malaria infections

from a population in northern Ghana.

Results

Sample collection and β-globin genotypes

Blood samples were collected from people testing positive for *P. falciparum* with mild malaria who self-presented to community health centres in northern Ghana over a three year period, from 22nd July 2015 to 30th August 2018. *P. falciparum* whole genome sequencing was attempted on all 2,174 samples collected, as part of the MalariaGEN *P. falciparum* Community Project (Pf7 data release [30]). 1,555 (71.5%) samples successfully passed quality control (QC) filtering for the MalariaGEN Pf7 data resource. β-globin genotyping by PCR and capillary sequencing was attempted on all available samples following parasite sequencing, with 1,807 (83.1%) samples successfully assigned a β-globin genotype (**Figure 1**). 1,368 (62.9%) samples both passed Pf7 QC and had a β-globin genotype call and were used for genome-wide association (GWA) analysis, referred to as the study population.

Figure 1. Study flow diagram.

Keeping the same nomenclature as that used by Band *et al.*, the frequencies of *Pfsa1+* (chr2:631,190 T>A in the 3D7 v3 reference genome) and *Pfsa3+* (chr11:1,058,035 T>A) were 3.9% and 4.5% in the study population, respectively. The *Pfsa2+* variant (chr2:814,288 C>T) was not present in this population. At the human β-globin locus, $\beta^{A/A}$ was the most common genotype identified in the study population (1008/1368, 74%), followed by $\beta^{A/C}$ (312/1368, 23%) (**Table 1**). 36 samples (2.6%) included at least one β^{S} allele ($\beta^{A/S}$, $\beta^{S/C}$, or $\beta^{S/S}$ genotypes), and 12 samples were $\beta^{C/C}$.

HBB genotype	Samples with β-globin call	Samples with β-globin call (%)	Samples β-globin called & Pf7 pass	Samples β-globin called & Pf7 pass (%)
HbAA	1311	73	1008	74
HbAS	50	2.8	30	2.2
HbSS	1	0.055	1	0.073
HbSC	7	0.39	5	0.37
HbCC	21	1.2	12	0.88
HbAC	417	23	312	23
Total	1807	100	1368	100

1   **Table 1.** Counts of the different β-globin genotypes in all samples that had β-globin genotype calls
2   available (N=1,807) and those that also had *P. falciparum* genomic data available that passed
3   MalariaGEN Pf7 quality control filters (N=1,368).

A fourth locus in *P. falciparum* associated with sickle haemoglobin

To investigate associations between *P. falciparum* variants and the human β-globin genotype, we first

applied a logistic regression approach using the human genotype as predictor and parasite genotype

as outcome variable, after excluding a subset of 59 samples due to high relatedness (**Supplementary**

Methods and **Supplementary Figure 2A**). Principal components analysis suggested some population

structure was present (**Supplementary Figure 2B**) and we included three principal components as

covariates to account for this. We used an indicator of $\beta^{S/*}$ individuals (N=35; 34 $\beta^{A/S}$ and 1 $\beta^{S/S}$) versus

β^{no-S} (N=1,274) as predictor, and tested for association with genotypes at each parasite variant in turn.

Full results from this analysis can be found in **Supplementary Table 1** and **Figure 2**. This analysis

revealed convincing replication of the previously identified associations at the *Pfsa1* locus (chr2:

631,190 T>A; $BF_{HbS} = 6.0 \times 10^7$, $P = 3.6 \times 10^{-11}$) and the *Pfsa3* locus (chr11:1,058,035 T>A,

$BF_{HbS} = 4.9 \times 10^{11}, P = 2.6 \times 10^{-15}$). Observed effect sizes for these loci were comparable to those

estimated previously in severe malaria cases (e.g. estimated odds ratio (OR) = 23, 95% confidence

interval (CI) = 10 - 50 for the *Pfsa3* locus; compared to OR = 22, 95% CI = 10 - 47 across all data reported

previously [29]). The *Pfsa2+* allele (chr2: 814,288 C>T) is rare in this population and was not tested.

We observed evidence for association with a previously unreported locus on chromosome 4. The

strongest evidence was seen at chr4:1,121,472 T>A ($BF_{HbS} = 6.5 \times 10^4, P = 5.8 \times 10^{-8}; OR = 10.9$,

95% CI = 4.6 - 25.9 for the 'T' allele) (**Figure 2**). This non-synonymous polymorphism occurs within the

gene *FIKK4.2* (gene ID PF3D7_0424700 in the 3D7 reference), which has not previously been

associated with sickle. (Of note, the 3D7 reference clone carries the associated 'T' allele, i.e. it is the

‘reference allele’ that was positively associated with sickle). A second nearby SNP (chr4:1,122,147 A>C) also had modest evidence for association ($BF_{HbS} = 1.1 \times 10^3$) (**Supplementary Figure 3**).

We repeated an assessment for parasite associations with sickle using a second independent GWA approach, using the FaST-LMM algorithm implemented in Python [31], with the full dataset of 1,368 samples (**Supplementary Figure 4**, **Supplementary Methods**). This identified significant associations at the same three loci: *Pfsa1*, *Pfsa3*, and the newly reported locus on chromosome 4.

Figure 2. Evidence for association between haemoglobin variants and *P. falciparum* genetic variants in Ghanaian mild malaria cases. Plot shows the log10 Bayes factor for association with HbS (top row) and HbC (bottom row) (y axis), for SNPs across the parasite genome (x axis).

We attempted to replicate the putative new association by analysing two other datasets (**Figure 3**). The chr4:1,121,472 T>A SNP did not appear strongly associated in the previously reported set of severe malaria cases from The Gambia [29] (estimated OR = 1.68 for the 'T' allele, 95% CI = 0.61 - 4.64); an accurate estimate was not possible in Kenya where the T allele is at extremely low frequency. However, we observed convincing replication of this association in 32 previously reported mild malaria cases from Mali, for which RNA-seq data has previously been reported [32]. Excluding 5 samples which appeared to have mixed bases in RNA reads at this locus (requiring < 5% or > 95% of reads at the locus to support expression of the T allele), we found strong evidence for replication: 11 of 14 (79%) infections of HbAS children carried the associated 'T' allele; while 11 of 13 (85%) HbAA infections carried the alternative 'A' allele; OR = 20, 95% CI = 2.8-145; *P*=2.8x10⁻³. We saw no evidence in these

1   samples that total read counts (reflecting expression levels) were associated with β^S status or with the
2   chr4:1,121,472 T>A genotype.
3
4

5
6   **Figure 3**. Discovery and replication evidence for association between HbS and the chr4:1,122,147 'T'
7   allele. Points show point estimates and bars show 95% confidence intervals, estimated using logistic
8   regression. The populations are: Gambian severe cases from [29], Kenyan severe cases from [29],
9   Ghanaian cases from this study, and Malian cases from [32]. Text indicates the 2-sided *P*-value for
10 each point.
11
12
Taken together, these results provide compelling evidence that the chr4:1,121,472 T>A SNP is
associated with β^S in mild malaria infections in west Africa, but raise questions as to why the
association is not also seen in the severe malaria data from Kenya and The Gambia analysed
previously, discussed further below. Following previous work, we refer to this new locus as *P.*
falciparum sickle-associated locus 4 (*Pfsa4*), and to alleles that are associated or not associated with
sickle as *Pfsa4+* ('T' allele, present in the 3D7 reference genome) and *Pfsa4-* ('A' allele, alternative to
the 3D7 reference genome)*, respectively. In this Ghanaian cohort of mild malaria cases, the *Pfsa1+*,
Pfsa3+, and *Pfsa4+* allele frequencies differed markedly between sickle and non-sickle individuals
(**Figure 4**). For example, *Pfsa4+* was present in 12/36 (33%) of individuals carrying at least one sickle
allele, compared with 3% of people with HbAA (**Supplementary Table 4**). After excluding missing or
heterozygous calls for the *Pfsa* alleles, parasites from individuals carrying at least one sickle allele were
around 17x more likely to carry at least one of the *Pfsa1+*, *Pfsa3+*, or *Pfsa4+* polymorphisms than to
have none of these alleles present (22% vs 1.3%, respectively) (**Supplementary Figure 5**). Conversely,

individuals carrying at least one sickle allele were around 11x more likely to be infected by parasites

carrying at least one *Pfsa+* allele than by parasites that did not carry any of these alleles (53% vs 5.1%,

respectively). Thus, despite the prevalence of sickle carriage among this Ghanaian cohort being

approximately 3%, and the prevalence of *Pfsa1+, Pfsa3+* and *Pfsa4+* alleles in *P. falciparum* being

around 4%, over half of the malaria infections encountered by sickle individuals were caused by

parasites carrying at least one *Pfsa+* allele.

FIKK4.2 is a member of a larger family of FIKK serine/ threonine kinase proteins. It is exported to the

erythrocyte cytosol during blood-stage infections, and, along with other FIKKs, is thought to regulate

phosphorylation of erythrocyte and parasite proteins in the erythrocyte membrane. FIKK4.2 has also

been associated with specific compartments in the infected red cell (termed K-dots [33]). In the 3D7

genome assembly, chr4:1,121,472 lies in the second exon of the *FIKK4.2* gene, and encodes a Leucine

→ Isoleucine change at amino acid 961. This is within the kinase domain, and is downstream of a

lengthy segment of low complexity repeat sequences [33] (**Supplementary Figure 3**). There is

considerable variation between repeat copies, and the number of repeat copies also appears

substantially variable within currently available *P. falciparum* assemblies [34]. However, we were

unable to find evidence that *Pfsa4+* is linked to variation in this repeat unit.

Little evidence of association with HbC

One potential explanation for effects specific to central west African populations, could be the

presence of the haemoglobin C variant in these populations. We conducted a second GWAS to test

for association between parasite variants and individuals possessing at least one HbC allele. To avoid

confounding by the HbS association we excluded all individuals carrying HbS alleles, leading to an

analysis of $\beta^{C/no-S}$ (N=312; 301 $\beta^{A/C}$ and 11 $\beta^{C/C}$), vs $\beta^{A/A}$ individuals (N=962). Applying the same

methodology as with the sickle analysis, we performed logistic regression using HPTEST. In contrast to

HbS, we did not identify any strong signals of association with HbC (**Figure 2, Supplementary Table 2**).

Similarly, for the linear mixed model (FaST-LMM algorithm), no significant associations were identified

(**Supplementary Methods**).

Population genetics of the *Pfsa* loci

We investigated linkage disequilibrium (LD) between the *Pfsa* loci. As expected, adjacent SNPs from

the *Pfsa3* locus on chromosome 11 (chr11:1,058,035 T>A, chr11:1,057,437 T>C, and chr11:1,054,282

T>C) were all in high LD with each-other (r^2 0.76 – 0.95), as were the four SNPs identified at the *Pfsa1*

locus on chromosome 2 (chr2:630,545 C>T, chr2:631,190 T>A, chr2:630,290 A>T, and chr2:629,375

C>G; r^2 0.91 – 0.98). The *Pfsa4* SNP with strongest evidence of association with sickle (chr4:1,121,472

T>A) was in LD with the nearby SNP identified in the GWA analyses (chr4:1,122,147 A>C, r^2 = 0.84). As

noted by Band *et al.*, *Pfsa1* and *Pfsa3* loci were in high LD with each-other despite being located on

different chromosomes. For example, the top hit SNPs from the *Pfsa1* and *Pfsa3* loci had an r^2 of 0.62.

Pfsa4, on chromosome 4, was also in relatively high LD with both *Pfsa1* and *Pfsa3* loci. For example,

the top hit SNP at *Pfsa4* (chr4:1,121,472 T>A) had an r^2 of 0.22 and 0.24 for the top hit SNPs at the

Pfsa1 and *Pfsa3* loci, respectively. These data suggest that *Pfsa1+*, *Pfsa3+* and *Pfsa4+* alleles are

maintained together despite being located on different chromosomes, consistent with having linked

biological functions.

Next, we investigated the global distribution of *Pfsa1-4* alleles using the MalariaGEN Pf7 data resource

[30]. Of the 20,864 samples in Pf7, we limited the analysis to samples that passed MalariaGEN Pf7 QC

(N=16,203), and to countries with ≥100 QC pass samples, yielding an analysis set of 15,738 samples

from 22 countries. The *Pfsa1-4+* alleles were almost completely absent from Asia, but were present

at varying frequencies in Africa and the only South American country represented, Colombia (**Figure**

5). *Pfsa1+* and *Pfsa3+* were distributed widely across Africa but with substantial heterogeneity in

prevalence; for example, the frequency of *Pfsa1+* in Benin was 59%, compared with 2% in Malawi

(**Supplementary Table 4**). *Pfsa4+* was essentially absent from east African populations, and was at

very low frequency in the Democratic Republic of the Congo (DRC), but was found at much higher

frequencies in west African populations (7% - 66%). The inverse pattern was seen in *Pfsa2+*, which was

prevalent in east Africa and the DRC but almost completely absent from west Africa. Intriguingly,

Pfsa3+ and *Pfsa4+* were also present at high frequencies (63% and 50%, respectively), and *Pfsa2+* at

lower frequency (7%), in Colombia.

Figure 4. Global distribution of *Pfsa1-4+* alleles in the MalariaGEN Pf7 data resource. Data from 15,738 samples from 22 countries, which passed Pf7 QC filters and requiring ≥100 samples for each country. Countries are organised left to right approximately from West to East Africa, with Colombia as the only non-African country represented. The *Pfsa+* alleles were very low or absent from all Asian countries in Pf7, not shown on Figure.

 Globally, in the Pf7 dataset, patterns of LD were complex and varied between populations (illustrated by Ghana and Kenya in **Figure 5**; all countries shown in **Supplementary Figure 6**). In west Africa, where the combination of *Pfsa1+*, *Pfsa3+* and *Pfsa4+* was present, SNPs at these loci had relatively high LD with each-other in Guinea, Ghana and Mali, but weaker LD in Senegal and The Gambia. In east Africa, where the combination of *Pfsa1+*, *Pfsa2+*, and *Pfsa3+* is present, these loci were in high LD with each-other in Malawi, Kenya and Tanzania; but LD was lower for these loci in the DRC. In Colombia, *Pfsa3+* and *Pfsa4+* had high LD (r^2= 0.49).

Figure 5. Frequencies and linkage disequilibrium (LD) of *Pfsa1-4+* alleles in Ghana (top) and Kenya (bottom).

In summary, these data reveal complex patterns of *Pfsa1-4+* allele frequencies and associations between populations. *Pfsa1+* and *Pfsa3+* are present across Africa, *Pfsa2+* is more localised to east

1   Africa, and *Pfsa4+* more localised to west Africa; within these regions, the extent to which *Pfsa+* alleles

2   correlate with each-other is variable, ranging from very high (e.g. r^2 of 0.67 for *Pfsa1+* and *Pfsa3+* in

3   Ghana), to zero (e.g. *Pfsa3+* and *Pfsa4+* in Benin).

Discussion

We have investigated host-parasite interaction by searching for associations between variation in the *P. falciparum* genome with human variation in the β-globin gene in a population with mild malaria from northern Ghana. Two parasite loci previously associated with sickle haemoglobin in severe malaria cases from Kenya and The Gambia [29] were replicated in this cohort, termed *Pfsa1* and *Pfsa3*. A new locus of association on *P. falciparum* chromosome 4 was identified within the gene *FIKK4.2*, which we name *P. falciparum* sickle associated locus 4, or *Pfsa4*. This locus was not associated with sickle in the severe malaria cases from The Gambia, and the variant is absent in Kenya. However, the *Pfsa4+* association with sickle did replicate in a sample of mild malaria cases from Mali. There were no significant associations found between parasite SNPs and $β^C$ genotypes.

The high prevalence of $β^C$ and relatively low prevalence of $β^S$ observed in this study is broadly consistent with previous results. HbC has mainly been described in west Africa, and it has been hypothesised that the relatively more benign phenotype of HbCC compared with HbSS would lead to the gradual replacement of the $β^S$ allele by $β^C$ in west Africa [5]. The proportion of $β^S$ carriers we observed was lower than other studies in the region [35,36]. Around 2% (15,000) of Ghanaian newborns are diagnosed with sickle cell disease (SCD) annually [37,38]; the majority of patients with SCD attending a large teaching hospital in Accra were found to have HbSS (55.7%) or HbSC (39.6%) genotypes [37]. The relatively lower prevalence of $β^S$ alleles we observed may be partly because sickle-containing genotypes protect against malaria disease, so the prevalence of such genotypes in this population, who were selected for having mild malaria, would be lower than in an unselected population. Variation in HbS prevalence may also be due to resource limitations in available healthcare to support children born with SCD in rural populations versus urban centres in Accra.

The replication of the association between *Pfsa1+* and *Pfsa3+* alleles with sickle in people with mild malaria is consistent with the view that parasites possessing these variants can bypass the protective effects of sickle and cause both mild and severe malaria syndromes in sickle individuals more readily than parasites lacking these variants. On this view, the sickle-associated variants effectively open an ecological niche (individuals carrying $β^S$ alleles) that is otherwise relatively closed off (or restricted) for *Pfsa-* parasites; rather than the view that these variants increase severity in sickle individuals *per se*. The age of the $β^S$ mutation is uncertain, with estimates suggesting a single origin around 20,000 years ago (range: 13,000-70,000 years [39]). Selection for *Pfsa+* mutations may have occurred once $β^S$ reached appreciable frequencies. It is currently unclear what has prevented *Pfsa+* parasites from outcompeting *Pfsa-* parasites in regions where HbS is common. The frequency of *Pfsa+* polymorphisms may depend on the regional prevalence of HbS, and the relative fitness advantage of *Pfsa+* alleles in

HbS RBCs versus their disadvantage in HbA RBCs, compared with *Pfsa-* alleles. Moreover, sexual recombination during the mosquito phase would be expected to mix *Pfsa+* and *Pfsa-* alleles. The presence of high LD between the *Pfsa+* polymorphisms is consistent with an epistatic effect on parasite fitness counter-acting the effect of sexual recombination, to maintain the *Pfsa+* alleles together; however, this requires further functional studies to confirm.

The *Pfsa4* locus was found within the gene encoding the serine/ threonine kinase FIKK4.2. The FIKK kinase family (characterised by a phenylalanine (F) – isoleucine (I) – lysine (K) – lysine (K) motif) has expanded in the *Laverania* clade (which includes *P. falciparum* and other *Plasmodium* species that infect great apes), comprising around 19 proteins in *P. falciparum* distributed over 11 chromosomes. The majority of *P. falciparum* FIKK kinases are exported into the host Red Blood Cell (RBC), and each is associated with a unique 'phosphorylation fingerprint' of host RBC proteins [40]. There is evidence that FIKK kinases play a role in remodelling the RBC surface and altering the mechanical properties of infected RBCs [41,42]. RBCs infected by parasites lacking FIKK4.2 through experimental knockout were less rigid and less adhesive compared with RBCs infected by wild-type parasites despite unchanged levels of *P. falciparum* Erythrocyte Protein 1 (PfEMP1) expression, suggesting a role in malaria pathogenesis [33]. FIKK kinases have been proposed as potential targets for antimalarial drug design. Functional experimentation will be required to explore whether the main *Pfsa4+* variant identified in this study (Leucine → Isoleucine at amino acid 961 in FIKK4.2) plays a causal role in facilitating parasite infection of RBCs from sickle heterozygote individuals, and by what mechanisms.

The lack of association between β^C alleles and parasite variation may reflect the fact that the vast majority of individuals possessing a β^C allele were genotype $\beta^{A/C}$, which is thought to confer a reduced protective effect against *P. falciparum* relative to $\beta^{C/C}$ [5], and likely also less than $\beta^{A/S}$; the strength of selection acting on the parasites by $\beta^{A/C}$ individuals will therefore be accordingly weaker. With only 12 $\beta^{C/C}$ individuals in our cohort, the sample size is too small to analyse this sub-group. The effect specifically of the $\beta^{S/C}$ genotype on malaria protection and *P. falciparum* genetic associations is, to our knowledge, unknown, as is the effect of other compound heterozygotes such as β^S with thalassaemia.

This study has several limitations. First, the small number of individuals carrying β^C and β^S alleles significantly limits the statistical power of the GWAS, meaning only variants with large effect sizes could be detected. Associations with smaller effect sizes and/or specific to certain β-globin genotypes such as $\beta^{C/C}$ will require larger sample sizes for discovery. Second, the method of genotyping the human β-globin locus involved PCR amplification and capillary sequencing a small region at the start

of the β-globin gene and did not account for other forms of genetic diversity in the haemoglobin genes. For example, it is possible that some of the homozygous calls were in fact compound heterozygotes with a β-thalassaemia allele. Although, complete β-globin gene deletion is an uncommon mechanism of β-thalassaemia [43], and β-thalassaemia is less common in west Africa than in other populations [23]. Third, minimal population demographic data were available to control for potential confounders, such as participant ethnic group. However, the logistic regression analysis was performed after removing samples that appeared highly similar, and incorporated principal components as covariates to account for population structure. Fourth, this study does not have a matched control group who did not have clinical malaria against which to compare the β-globin genotype frequencies. This study therefore cannot directly assess the extent to which *Pfsa+* alleles reduce the protective effect of sickle against mild malaria relative to people without symptomatic malaria. However, their association with β^S alleles in mild malaria, combined with the fact that the same variants reduce the protective effect of sickle against severe malaria relative to population controls [29], supports the view that these variants allow parasites to access sickle individuals which wild-type parasites otherwise struggle to infect, and therefore reduce the protective effect of sickle against both mild and severe malaria. We also note that the MalariaGEN Pf7 data resource is collated from many separate studies, and this opportunistic sampling may not be entirely representative of unselected populations.

In conclusion, this study adds further evidence that several loci in the *P. falciparum* genome are associated with sickle haemoglobin, in both mild and severe malaria. Variants at these loci are maintained in linkage with each-other despite being on separate chromosomes. This suggests that in a population, subsets of *P. falciparum* parasites will inherit genetic traits that make them better suited to particular human genetic contexts, such as 'specialists' at infecting sickle RBCs. The frequencies of different host genotypes, and the relative fitness costs of the polymorphisms for the parasite in non-sickle hosts, will sculpt the evolutionary pressures acting on these loci in the parasite, which may in turn affect the selection pressures acting on the human genome in an ancient coevolutionary process. Future work could extend these results by recruiting a larger population to ensure robust statistical power to analyse different haemoglobin genotypes, incorporate more genetic variation from both human and *P. falciparum* genomes, and establish functional data to uncover the mechanisms of how these variants help the parasite to infect people carrying sickle haemoglobin alleles.

Methods

Study site and samples

Samples were collected from the Kassena-Nankana East Municipality and Kassena-Nankana West District, two adjoining administrative areas in the Upper East Region of northern Ghana, hereafter referred to as the KNDs. The population is primarily rural, and malaria transmission is seasonal, peaking during the rainy season from July to October. *P. falciparum* genetic diversity has been studied in this region previously [44] through a network of community health centres.

This study ran from 22/07/2015 to 30/08/2018, encompassing the rainy seasons for 2015, 2016 and 2017. Patients presenting to the Navrongo War Memorial Hospital (NWMH) and four satellite clinics in the KNDs with compatible symptoms for malaria were tested using an antigen Rapid Diagnostic Test (RDT) (CareStart™ malaria Pf (HRP2), Access Bio, NJ, USA). Participants were eligible if they tested positive for *P. falciparum* by RDT and informed consent was obtained and documented. Written consent forms were available in the predominant languages spoken in the region: Kasem, Nankam, and English. Four 50uL dried blood spots (DBS) and malaria blood smears for microscopy were collected from consenting participants. Parasitaemia was calculated by microscopy as asexual parasites per 200 white blood cells (WBC). DBS samples were transferred to the Wellcome Sanger Institute (WSI), UK, for DNA extraction and sequencing. DNA extraction from DBS was performed using the QIAamp Investigator Biorobot kit on the Qiagen Biorobot Universal instrument, protocol described in [45].

Parasite whole genome sequencing and genotype calling

P. falciparum whole genome sequencing from DBS extractions was performed using the MalariaGEN pipeline established at the Wellcome Sanger Institute (WSI), UK, described elsewhere [30]. Briefly, DBS samples undergo selective whole genome amplification (sWGA) to enrich for *P. falciparum* DNA [46], followed by short-read Illumina sequencing on the HiSeq platform. These samples are part of the broader MalariaGEN *P. falciparum* Community Project [30,47,48]. Genotype calling from the Illumina sequence data was performed at WSI using the MalariaGEN informatics pipeline [30]. All of the parasite genomes included in this study are available via the current iteration of the MalariaGEN *P. falciparum* Community Project, Pf7 [30].

Human haemoglobin genotyping

Samples with ethical approval for use in this study (a subset of the MalariaGEN *P. falciparum* Community Project collection) were transferred via Beckman robot to new 96-well plate layouts organised by sample concentration (measured by Qubit High Sensitivity (HS) assay) and remaining

volume after parasite sequencing. Human β-globin genotyping at codon 6 followed a PCR amplification and capillary sequencing strategy. Details on the primers and PCR conditions used can be found in the **Supplementary Methods**. The PCR products were capillary sequenced by the Eurofins company, using their PCR clean-up and Supreme Run service.

Capillary sequence chromatogram traces were visually inspected using the 4Peaks software tool (**Supplementary Methods**). The six callable β-globin genotypes from the sixth codon were: $\beta^{A/A}$, $\beta^{A/S}$, $\beta^{S/S}$, $\beta^{S/C}$, $\beta^{A/C}$, and $\beta^{C/C}$. Samples with ambiguous calls (eg. low amplitude chromatograph or chaotic uninterpretable signal) were repeat PCR-amplified and re-sequenced if available; if these samples remained ambiguous after the second attempt (or if there was no sample left for further attempts) then the sample was deemed to have failed β-globin genotyping. All $\beta^{A/S}$, $\beta^{S/S}$, $\beta^{S/C}$, and $\beta^{C/C}$ samples were repeat PCR-amplified and re-sequenced to confirm the result; there were no samples for which the second run disagreed with the first run if a call was successfully made on both runs.

P. falciparum genome variant filtering for study population

Parasite genomic analyses were conducted using the data structures and curated sample metadata of the MalariaGEN Pf7 data resource [30]. Full details of variant filtering steps are described in **Supplementary Methods**. Briefly, the approximately 10 million variants in Pf7 were initially filtered to create a 'working set' of variants, including only biallelic SNPs that passed Pf7 Quality Control (QC) and had at least 0.1% allele frequency in the Ghana study population, yielding 123,845 SNPs. Heterozygous genotypes (representing mixed infections in *P. falciparum*) were regarded as missing. The previously reported lead associations in the *Pfsa* regions were explicitly included. This 'working set' of 123,845 SNPs was then further filtered for different analyses; for example, the main logistic regression GWA analysis described in main text included 18,770 SNPs with a minor allele count of at least 10 among the 1,309 samples included in the analysis (after removing closely related samples). The FaST-LMM analysis was performed on 7,250 SNPs with $\geq$3% minor allele frequency and $\leq$20% missing or mixed genotype calls among the 1,368 samples analysed. All genomic data manipulations from the Pf7 data resource were performed in Python; genotype and haplotype manipulations and downstream population genetic analyses were performed using the *scikit-allel* package.

Population structure analysis

We used principal components analysis (PCA), as implemented in the QCTOOL package [49] to assess population structure in the 1,368 samples passing QC and with both parasite and β-globin genotypes in our data (further details in **Supplementary Methods**). For this analysis, we restricted attention to the set of 14,381 SNPs (selected from the 'working set' of 123,845 SNPs) with at least 1% minor allele

frequency and no more than 25% missing genotypes in these samples. Initial principal components (PCs) (**Supplementary Figure 3**) revealed a small number of sample pairs with high relatedness values (estimated r > 0.5) that dominated the first two principal components. We therefore implemented a greedy algorithm to remove one of each sample pair having $r_{ij} > 0.5$. In total we removed 59 samples, leaving 1,309. We then re-computed principal components and loadings (**Supplementary Figures 4-5**).

Genome Wide Association

We performed a logistic regression analysis using the program HPTEST [29], which fits a logistic regression model with the host genotype as predictor and the parasite genotype as outcome, treating each variant separately (further details in **Supplementary Methods**). HPTEST reports a Bayes Factor reflecting the evidence for association as well as other summaries including an effect size estimate, standard error, and *P*-value. Interpretation of HPTEST results was previously discussed in detail [29]. For this analysis, we used the 1,309 samples without closely related parasite samples identified through PCA, either with or without including the first 3 principal components.

 In a second approach, we used a Factored Spectrally Transformed (FaST) Linear Mixed Model (LMM) [31], implemented in Python, using the full set of 1,368 samples. LMMs tackle confounders by using measures of genetic similarity to capture the probabilities that pairs of individuals have causative alleles in common. FaST-LMM has previously been applied to GWAS in *P. falciparum* (eg. [50–52]).

Population genetics of the *Pfsa* loci

Allele frequencies and linkage disequilibrium (LD) for the *Pfsa* loci were computed in Python using the *scikit-allel* package. For the analysis of global patterns of *Pfsa* frequencies and LD, the total MalariaGEN Pf7 data resource (20,864 samples) was restricted to samples that passed Pf7 QC (N=16,203), and to countries with $\geq$100 QC pass samples, yielding 15,738 samples from 22 countries. One SNP was selected to represent each *Pfsa* locus for frequency and LD calculations: *Pfsa1* = chr2:631,190; *Pfsa2* = chr2:814,288; *Pfsa3* = chr11:1,058,035; *Pfsa4* (newly reported in this study) = chr4:1,121,472.

Ethics

Ethical approval for the study was granted by the Navrongo Health Research Centre Institutional Review Board (NHRCIRB203). All participants or their guardians (as appropriate) were provided detailed information sheets and gave informed consent prior to enrolment. Further approval was granted by the Wellcome Sanger Institute's Research Ethics Committee for the analysis of the samples.

Sample collection was conducted by LNA, who pseudonymised the samples prior to making them available to the Wellcome Sanger Institute. All patient-identifiable data are securely stored by LNA and only non-patient identifiable were provided to the Wellcome Sanger Institute. The study complies with all relevant ethical regulations.

Data availability statement

The *P. falciparum* variant calls (VCF file) are available to download open-access via the MalariaGEN *P. falciparum* Community Project Pf7 resource. (The VCF file contains variant calls for all samples in Pf7, which includes all of the samples analysed in this study).

Code availability statement

Source code for QCTOOL and HPTEST is available at: code.enkre.net/qctool, and on Zenodo (doi:10.5281/zenodo.5685581).

Inclusion and ethics statement

This research was conducted collaboratively between researchers based at the University of Ghana, the Navrongo Health Research Centre (NHRC), the Wellcome Sanger Institute, and University of Oxford. Study conceptualisation, data analysis and interpretation, and authorship of publications were shared jointly between personnel based in Ghana and the UK.

Acknowledgements

We thank the teams in the Navrongo Health Research Centre (NHRC) and the Wellcome Sanger Institute (WSI) who generated the *P. falciparum* genome sequence data as part of the MalariaGEN *P. falciparum* Community Project, and the study participants and their guardians who provided the samples. Thank you to Richard Pearson for his help accessing and using the Pf7 data resource, and to Christen Smith, Sonia Goncalves and Rachel Ogidan for support ensuring legal and ethical compliance. This work is dedicated to the memory of Professor Dominic Kwiatkowski, who died prior to its completion.

1 Bibliography

1. World Health Organization. World Malaria Report 2022 [Internet]. 2022 [cited 2022 Dec 9]. Available from: https://www.who.int/publications/i/item/9789240064898

2. Kwiatkowski DP. How malaria has affected the human genome and what human genetics can teach us about malaria. Am J Hum Genet. 2005;77:171–92.

3. Hamblin MT, Di Rienzo A. Detection of the signature of natural selection in humans: Evidence from the Duffy blood group locus. Am J Hum Genet. 2000;66:1669–79.

4. Williams TN, Maitland K, Bennett S, Ganczakowski M, Peto TEA, Newbold CI, et al. High incidence of malaria in α-thalassaemic children. Nature. 1996;383:522–5.

5. Modiano D, Luoni G, Sirima BS, Simporé J, Verra F, Konaté A, et al. Haemoglobin C protects against clinical Plasmodium falciparum malaria. Nature. 2001;414:305–8.

6. Siniscalco M, Bernini L, Filippi G, Latte B, Meera Khan P, Piomelli S, et al. Population genetics of haemoglobin variants, thalassaemia and glucose-6-phosphate dehydrogenase deficiency, with particular reference to the malaria hypothesis. Bull World Health Organ [Internet]. 1966;34:379–93. Available from: /pmc/articles/PMC2475975/?report=abstract

7. Allison AC. The discovery of resistance to malaria of sickle-cell heterozygotes. Biochem Mol Biol Educ. 2002;30:279–87.

8. Allison AC. The distribution of the sickle-cell trait in East Africa and elsewhere, and its apparent relationship to the incidence of subtertian malaria. Trans R Soc Trop Med Hyg [Internet]. 1954;48:312–8. Available from: https://doi.org/10.1016/0035-9203(54)90101-7

9. Allison AC. Protection Afforded by Sickle-cell Trait Against Subtertian Malarial Infection. Br Med J [Internet]. 1954;1:290 LP – 294. Available from: http://www.bmj.com/content/1/4857/290.abstract

10. Zimmerman PA, Woolley I, Masinde GL, Miller SM, McNamara DT, Hazlett F, et al. Emergence of FY*Anull in a Plasmodium vivax-endemic region of Papua New Guinea. Proc Natl Acad Sci U S A. 1999;96:13973–7.

11. Allen SJ, O'Donnell A, Alexander NDE, Alpers MP, Peto TEA, Clegg JB, et al. α+-Thalassemia protects children against disease caused by other infections as well as malaria. Proc Natl Acad Sci U S A. 1997;94:14736–41.

12. Weatherall D, Clegg J, Kwiatkowski D. The role of genomics in studying genetic susceptibility to infectious disease. Genome Res. 1997;7:967–73.

13. Flint J, Hill AVS, Bowden DK, Oppenheimer SJ, Sill PR, Serjeantson SW, et al. High frequencies of α-thalassaemia are the result of natural selection by malaria. Nature. 1986;321:744–50.

14. Yenchitsomanus P, Summers KM, Board PG, Bhatia KK, Jones GL, Johnston K, et al. Alpha-thalassemia in Papua New Guinea. Hum Genet. 1986;74:432–7.

15. Taylor SM, Parobek CM, Fairhurst RM. Haemoglobinopathies and the clinical epidemiology of malaria: A systematic review and meta-analysis. Lancet Infect Dis [Internet]. Elsevier Ltd; 2012;12:457–68. Available from: http://dx.doi.org/10.1016/S1473-3099(12)70055-5

16. Tishkoff SA, Varkonyi R, Cahinhinan N, Abbes S, Argyropoulos G, Destro-Bisol G, et al. Haplotype diversity and linkage disequilibrium at human G6PD: Recent origin of alleles that confer malarial resistance. Science (80-). 2001;293:455–62.

17. Carter R, Mendis KN. Evolutionary and historical aspects of the burden of malaria. Clin Microbiol Rev. 2002;15:564–94.

18. RAY RN, CHATTERJEA JB, CHAUDHURI RN. Observations on the Resistance in Hb E Thalassaemia Disease To Induced. Bull World Health Organ. 1964;30:51–5.

19. Rockett KA, Clarke GM, Fitzpatrick K, Hubbart C, Jeffreys AE, Rowlands K, et al. Reappraisal of known malaria resistance loci in a large multicenter study. Nat Genet. 2014;46:1197–204.

20. MalariaGEN. Insights into malaria susceptibility using genome-wide data on 17,000 individuals from Africa, Asia and Oceania. Nat Commun. 2019;10:5732.

21. Mackinnon MJ, Mwangi TW, Snow RW, Marsh K, Williams TN. Heritability of malaria in Africa. PLoS Med. 2005;2:1253–9.

22. Weatherall DJ. Phenotype-genotype relationships in monogenic disease: Lessons from the thalassaemias. Nat Rev Genet. 2001;2:245–55.

23. Williams TN, Weatherall DJ. World distribution, population genetics, and health burden of the hemoglobinopathies. Cold Spring Harb Perspect Med. 2012;2:1–14.

24. Piel FB, Patil AP, Howes RE, Nyangiri OA, Gething PW, Dewi M, et al. Global epidemiology of Sickle haemoglobin in neonates: A contemporary geostatistical model-based map and population estimates. Lancet [Internet]. Elsevier Ltd; 2013;381:142–51. Available from: http://dx.doi.org/10.1016/S0140-6736(12)61229-X

25. Sundd P, Gladwin MT, Novelli EM. Pathophysiology of Sickle Cell Disease. Annu Rev Pathol Mech Dis. 2019;14:263–92.

26. Archer NM, Petersen N, Clark MA, Buckee CO, Childs LM, Duraisingh MT. Resistance to Plasmodium falciparum in sickle cell trait erythrocytes is driven by oxygen-dependent growth inhibition. Proc Natl Acad Sci U S A. 2018;115:7350–5.

27. Piel FB, Howes RE, Patil AP, Nyangiri OA, Gething PW, Bhatt S, et al. The distribution of haemoglobin C and its prevalence in newborns in Africa. Sci Rep. 2013;3:1–8.

28. Weatherall DJ, Clegg JB. The thalassaemia syndromes. Oxford: Blackwell Scientific Publications.; 2001.

29. Band G, Leffler EM, Jallow M, Sisay-Joof F, Ndila CM, Macharia AW, et al. Malaria protection due to sickle haemoglobin depends on parasite genotype. Nature [Internet]. 2021; Available from: https://doi.org/10.1038/s41586-021-04288-3

30. MalariaGEN. Pf7: an open dataset of Plasmodium falciparum genome variation in 20,000 worldwide samples. Wellcome Open Res. 2023;8.

31. Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D. FaST linear mixed models for genome-wide association studies. Nat Methods. 2011;8:833–5.

32. Saelens JW, Petersen JE V., Freedman E, Moseley RC, Konaté D, Diakité SAS, et al. Impact of Sickle Cell Trait Hemoglobin on the Intraerythrocytic Transcriptional Program of Plasmodium falciparum. mSphere. American Society for Microbiology; 2021;6.

33. Kats LM, Fernandez KM, Glenister FK, Herrmann S, Buckingham DW, Siddiqui G, et al. An exported kinase (FIKK4.2) that mediates virulence-associated changes in Plasmodium falciparum-infected red blood cells. Int J Parasitol [Internet]. Australian Society for Parasitology Inc.; 2014;44:319–28. Available from: http://dx.doi.org/10.1016/j.ijpara.2014.01.003

34. Otto TD, Böhme U, Sanders M, Reid A, Bruske EI, Duffy CW, et al. Long read assemblies of geographically dispersed Plasmodium falciparum isolates reveal highly structured subtelomeres [version 1; referees: 3 approved]. Wellcome Open Res. 2018;3.

35. Ringelhann B, Dodu SRA, Konotey-Ahulu FID, Lehmann H. A survey for haemoglobin variants, thalassaemia and glucose-6-phosphate dehydrogenase deficiency in northern Ghana. Ghana Med J. 1968;7:120–4.

36. Mensah-Brown HE, Abugri J, Asante KP, Dwomoh D, Dosoo D, Atuguba F, et al. Assessing the impact of differences in malaria transmission intensity on clinical and haematological

indices in children with malaria. Malar J. BioMed Central; 2017;16:1–11.

37. Asare E V., Wilson I, Benneh-Akwasi Kuma AA, Dei-Adomakoh Y, Sey F, Olayemi E. Burden of Sickle Cell Disease in Ghana: The Korle-Bu Experience. Adv Hematol. 2018;2018.

38. Ohene-Frempong K, Oduro J, Tetteh H, Nkrumah F. Screening newborns for sickle cell disease in Ghana. Pediatrics [Internet]. 2008;121:S120–1. Available from: https://doi.org/10.1542/peds.2007-2022UUU

39. Laval G, Peyrégne S, Zidane N, Harmant C, Renaud F, Patin E, et al. Recent Adaptive Acquisition by African Rainforest Hunter-Gatherers of the Late Pleistocene Sickle-Cell Mutation Suggests Past Differences in Malaria Exposure. Am J Hum Genet. 2019;104:553–61.

40. Davies H, Belda H, Broncel M, Ye X, Bisson C, Introini V, et al. An exported kinase family mediates species-specific erythrocyte remodelling and virulence in human malaria. Nat Microbiol [Internet]. Springer US; 2020;5:848–63. Available from: http://dx.doi.org/10.1038/s41564-020-0702-4

41. Nunes MC, Goldring JPD, Doerig C, Scherf A. A novel protein kinase family in Plasmodium falciparum is differentially transcribed and secreted to various cellular compartments of the host cell. Mol Microbiol. 2007;63:391–403.

42. Nunes MC, Okada M, Scheidig-Benatar C, Cooke BM, Scherf A. Plasmodium falciparum FIKK kinase members target distinct components of the erythrocyte membrane. PLoS One. 2010;5:1–8.

43. Thein SL. Molecular basis of β thalassemia and potential therapeutic targets. Blood Cells, Mol Dis [Internet]. Elsevier; 2018;70:54–65. Available from: https://doi.org/10.1016/j.bcmd.2017.06.001

44. Amenga-Etego LN, Asoala V, Agongo G, Jacob C, Goncalves S, Awandare GA, et al. Temporal evolution of sulfadoxine-pyrimethamine resistance genotypes and genetic diversity in response to a decade of increased interventions against Plasmodium falciparum in northern Ghana. Malar J [Internet]. BioMed Central; 2021;20:1–11. Available from: https://doi.org/10.1186/s12936-021-03693-3

45. Girgis ST, Adika E, Nenyewodey FE, Jnr DKS, Ngoi JM, Lorenz O, et al. Nanopore sequencing for real-time genomic surveillance of Plasmodium falciparum. bioRxiv. 2022;2022.12.20.521122.

46. Oyola SO, Ariani C V., Hamilton WL, Kekre M, Amenga-Etego LN, Ghansah A, et al. Whole genome sequencing of Plasmodium falciparum from dried blood spots using selective whole genome amplification. Malar J. BioMed Central; 2016;15:597.

47. MalariaGEN. An open dataset of Plasmodium falciparum genome variation in 7,000 worldwide samples [version 1; peer review: 2 approved]. Wellcome Open Res [Internet]. 2021;6:42. Available from: https://doi.org/10.12688/wellcomeopenres.16168.1

48. Achidi EA, Agbenyega T, Allen S, Amodu O, Bojang K, Conway D, et al. A global network for investigating the genomic epidemiology of malaria. Nature. 2008;456:732–7.

49. Band G, Rockett KA, Spencer CCA, Kwiatkowski DP, Si Le Q, Clarke GM, et al. A novel locus of resistance to severe malaria in a region of ancient balancing selection. Nature. 2015;526:253–7.

50. Miotto O, Amato R, Ashley EA, Macinnis B, Almagro-Garcia J, Amaratunga C, et al. Genetic architecture of artemisinin-resistant Plasmodium falciparum. Nat Genet. 2015;47:226–34.

51. Amato R, Lim P, Miotto O, Amaratunga C, Dek D, Pearson RD, et al. Genetic markers associated with dihydroartemisinin–piperaquine failure in Plasmodium falciparum malaria in

Cambodia: a genotype–phenotype association study. Lancet Infect Dis [Internet]. The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license; 2017;17:164–73. Available from: http://dx.doi.org/10.1016/S1473-3099(16)30409-1

52. Zhu L, Tripathi J, Rocamora FM, Miotto O, van der Pluijm R, Voss TS, et al. The origins of malaria artemisinin resistance defined by a genetic and transcriptomic background. Nat Commun. 2018;9:1–13.