A fourth locus in the Plasmodium falciparum genome associated with sickle haemoglobin

Authors
William L. Hamilton[1,2,3 *], Annie J. Forster[4], Yaw Aniweh[5], Victor Asoala[6], Eleanor Drury[1], Kirk A. Rockett[4], Gordon A. Awandare[5], Dominic P. Kwiatkowski[1], Gavin Band[4], Lucas N. Amenga-Etego[5 *]

Author affiliations
1. Wellcome Sanger Institute, Wellcome Trust Genome Campus, Hinxton, CB10 1RQ, United Kingdom
2. University of Cambridge, Department of Medicine, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, United Kingdom
3. Cambridge University Hospitals NHS Foundation Trust, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, United Kingdom
4. Centre for Human Genetics, Roosevelt Dr, Oxford OX3 7BN, United Kingdom
5. West African Centre for Cell Biology of Infectious Pathogens (WACCBIP), University of Ghana, Legon, Accra, Ghana
6. Navrongo Health Research Centre (NHRC), Ghana Health Service, Navrongo, Upper East Region, Ghana

* Corresponding authors
Dr. William L. Hamilton: wh2@sanger.ac.uk
Dr. Lucas N. Amenga-Etego: lamenga-etego@ug.edu.gh

Key words
Malaria, *P. falciparum*, sickle, haemoglobin, HbS, GWAS, genome-wide association, host-parasite interaction

Author contributions
Conceptualization: LNA, WLH, DPK
Investigation: WLH, AJF, ED, GB, LNA
Formal analysis: WLH, AJF, GB, LNA
Resources: KAR
Data Curation: WLH, LNA
Writing – original draft preparation: WLH, GB, LNA
Writing – review and editing: WLH, GB, LNA, KAR, YA, VA
Visualization: WLH, AJF, GB
Supervision: LNA, GB, DPK
Project administration: LNA, YA, VA
Funding acquisition: DPK, LNA, GAA

Abstract

Heterozygosity for sickle haemoglobin (HbS) confers protection against severe malaria

caused by the parasite *Plasmodium falciparum.* Recent work suggests that this protective

effect can depend on the parasite genotype. Here, we performed a genome-wide association

analysis of *P. falciparum* against human β-globin genotypes in a sample of 1,368 people with

mild malaria in northern Ghana. We replicated the previously identified associations with

HbS at two parasite loci. However, a newly identified locus within the serine/threonine kinase

FIKK4.2, which we term *Pfsa4*, was also associated with HbS; this finding replicated in a

published sample from Mali. The *Pfsa1-4* mutations vary widely in frequencies across Africa,

are absent from Asia, and are highly correlated with each-other across multiple populations.

We found no strong associations with haemoglobin C. These findings add new complexity to

the emerging picture of association between human and co-evolving malaria parasite

genomes, suggesting new avenues for functional exploration.

1   Introduction

The World Health Organization (WHO) estimates that there were 247 million malaria cases

and 619,000 deaths in 2021 [1]. 95% of cases and 96% of deaths occurred in Africa, and

76% of deaths were in children under 5 years old. Malaria has taken a major toll on human

health for millennia, particularly for young children prior to the development of protective

(though incomplete) immunity. Consequently, malaria has exerted a strong evolutionary

selective pressure on the human genome [2]. Malaria can cause a wide spectrum of clinical

disease, from asymptomatic parasitaemia to severe malaria syndromes and death.

Understanding how human and parasite genetic variation interact with each-other and

contribute to malaria disease severity is a key goal of malaria research.

Genetic variation at dozens of human loci have been proposed to provide protection against

malaria [3–18]; although, some of these have not replicated in larger multi-centre studies

[19,20]. It is likely that many host polymorphisms contribute to disease susceptibility, most

with small effects [21]. Approximately 7% of the world's population are thought to be carriers

for inherited haemoglobin disorders, making them the most common human monogenic

diseases [22]. Natural variation in the human haemoglobin genes was hypothesised to

provide protection against malaria since at least the 1950s [15–17], given the central role of

red blood cells (RBCs) in the *Plasmodium* life cycle, and the overlay of certain haemoglobin

variants with malaria prevalence. However, the full scope of genetic diversity in human

haemoglobin, and how this relates to malaria disease severity, is highly complex, and

remains incompletely understood.

The dominant form of adult haemoglobin (HbA) comprises a tetramer of proteins transcribed

from two α-globin genes and two β-globin genes ($\alpha_2\beta_2$), responsible for oxygen

transportation within RBCs. The α-globin and β-globin genes are located within the α- and β-

globin loci on chromosomes 16 and 11, respectively. Of the haemoglobinopathies (altered

haemoglobin makeup and/or structure), haemoglobin S (HbS, or sickle), haemoglobin C

(HbC) and haemoglobin E (HbE) occur at significant population frequencies with distinct

geographic distributions [23]. The sickle β-globin allele (β^S) is a single point non-

synonymous polymorphism changing Glutamic acid (coded by GTG) to Valine (GAG) at the

sixth codon; this remains the strongest known association with protection against malaria in

the human genome [19,20], maintaining high frequencies in populations exposed to malaria,

primarily in Africa [24]. Homozygosity for β^S is the most common cause of sickle cell disease

(SCD) in Africa. The central pathophysiology of SCD involves polymerisation of

deoxygenated HbS, causing RBCs to become inflexible and misshapen, leading to

haemolysis, endothelial dysfunction, inflammation and occlusion within the microcirculation

[25]. Malaria protection is driven by the heterozygous state (HbAS) and fixation of the sickle

allele cannot be attained due to the deleterious nature of the homozygote state (HbSS). The

mechanisms by which heterozygous HbS provides malaria protection are still not fully

understood, but several have been proposed (summarised in [26]).

HbS is present throughout sub-Saharan Africa to varying degrees [23,24], but a second

haemoglobin type, HbC, occurs at high frequency only in central west Africa [27]. HbC arises

due to an alternative single-point polymorphism in the sixth codon of β-globin ($β^C$) at the

base pair adjacent to the $β^S$ polymorphism, leading to a switch from Glutamatic acid to

Lysine. In west Africa, therefore, β-globin genotypes can include $β^{A/A}$ (HbAA wild-type), $β^{A/S}$

(HbAS sickle heterozygote), $β^{S/S}$ (HbSS sickle homozygote), $β^{A/C}$ (HbAC heterozygote), $β^{C/C}$

(HbCC homozygote), and $β^{S/C}$ (HbSC compound heterozygote). Many other polymorphisms

in the β-globin gene have been described, including those that change gene expression

causing β-thalassaemias [23,28], but these are less common in west Africa.

How variation in human and malaria parasite genomes interact with each-other and give rise

to disease phenotypes remains an active area of research. A recent study of around 4,000

samples from children with severe malaria from Kenya and The Gambia identified three

parasite loci that were associated with sickle haemoglobin, termed *P. falciparum* sickle-

associated loci or *Pfsa1*, *Pfsa2* and *Pfsa3* [29]. *Pfsa1* and *Pfsa2* are on different loci on

chromosome 2, while *Pfsa3* is located in a region of suspected structural variation on

chromosome 11. The functional mechanism(s) underpinning these associations is unclear.

Here, we have tested for associations between the human β-globin variants $β^S$ and $β^C$, and

genome-wide variation in *P. falciparum* parasites, in a sample of mild malaria infections from

a population in northern Ghana.

Results

Sample collection and β-globin genotypes

Blood samples were collected from people testing positive for *P. falciparum* with mild malaria who self-presented to community health centres in northern Ghana over a three year period, from 22nd July 2015 to 30th August 2018. *P. falciparum* whole genome sequencing was attempted on all 2,174 samples collected, as part of the MalariaGEN *P. falciparum* Community Project (Pf7 data release [30]). 1,555 (71.5%) samples successfully passed quality control (QC) filtering for the MalariaGEN Pf7 data resource. β-globin genotyping by PCR and capillary sequencing was attempted on all available samples following parasite sequencing, with 1,807 (83.1%) samples successfully assigned a β-globin genotype (**Figure 1**). 1,368 (62.9%) samples both passed Pf7 QC and had a β-globin genotype call and were used for genome-wide association (GWA) analysis, referred to as the study population.

Keeping the same nomenclature as that used by Band *et al.*, the frequencies of *Pfsa1+* (chr2:631,190 T>A in the 3D7 v3 reference genome) and *Pfsa3+* (chr11:1,058,035 T>A) were 3.9% and 4.5% in the study population, respectively. The *Pfsa2+* variant (chr2:814,288 C>T) was not present in this population. At the human β-globin locus, β$^{A/A}$ was the most common genotype identified in the study population (1008/1368, 74%), followed by β$^{A/C}$ (312/1368, 23%) (**Table 1**). 36 samples (2.6%) included at least one β^{S} allele (β$^{A/S}$, β$^{S/C}$, or β$^{S/S}$ genotypes), and 12 samples were β$^{C/C}$.

A fourth locus in *P. falciparum* associated with sickle haemoglobin

To investigate associations between *P. falciparum* variants and the human β-globin genotype, we applied a logistic regression approach using the human genotype as predictor and parasite genotype as outcome variable, after excluding a subset of 59 samples due to high relatedness (**Supplementary Methods** and **Supplementary Figure 1A**). Principal components analysis suggested some population structure was present (**Supplementary Figure 1B**) and we included three principal components as covariates to account for this. We used an indicator of β$^{S/*}$ individuals (N=35; 34 β$^{A/S}$ and 1 β$^{S/S}$) versus β^{no-S} (N=1,274) as predictor, and tested for association with genotypes at each parasite variant in turn. Full results from this analysis can be found in **Supplementary Table 1** and **Figure 2**. This analysis revealed convincing replication of the previously identified associations at the *Pfsa1* locus (chr2: 631,190 T>A; $BF_{HbS} = 6.0 \times 10^{7}, \ P = 3.6 \times 10^{-11}$) and the *Pfsa3* locus (chr11:1,058,035 T>A, $BF_{HbS} = 4.9 \times 10^{11}, P = 2.6 \times 10^{-15}$). Observed effect sizes for these loci were comparable to those estimated previously in severe malaria cases (e.g. estimated odds ratio (OR) = 23, 95% confidence interval (CI) = 10 - 50 for the *Pfsa3* locus; compared

to OR = 22, 95% CI = 10 - 47 across all data reported previously [29]). The *Pfsa2+* allele

(chr2: 814,288 C>T) is rare in this population and was not tested.

We observed evidence for association with a previously unreported locus on chromosome 4.

The strongest evidence was seen at chr4:1,121,472 T>A ($BF_{HbS} = 6.5 \times 10^4, P = 5.8 \times 10^{-8}$;

$OR = 10.9$, 95% CI = 4.6 - 25.9 for the 'T' allele) (**Figure 2**). This non-synonymous

polymorphism occurs within the gene *FIKK4.2* (gene ID PF3D7_0424700 in the 3D7

reference), which has not previously been associated with sickle. (Of note, the 3D7

reference clone carries the associated 'T' allele, i.e. it is the 'reference allele' that was

positively associated with sickle). A second nearby SNP (chr4:1,122,147 A>C) also had

modest evidence for association ($BF_{HbS} = 1.1 \times 10^3$) (**Figure 3**).

We attempted to replicate the putative new association by analysing two other datasets

(**Figure 3**). The chr4:1,121,472 T>A SNP did not appear strongly associated in the

previously reported set of severe malaria cases from The Gambia [29] (estimated OR = 1.68

for the 'T' allele, 95% CI = 0.61 - 4.64); an accurate estimate was not possible in Kenya

where the T allele is at extremely low frequency. However, we observed convincing

replication of this association in 32 previously reported mild malaria cases from Mali, for

which RNA-seq data has previously been reported [32]. Excluding 5 samples which

appeared to have mixed bases in RNA reads at this locus (requiring < 5% or > 95% of reads

at the locus to support expression of the T allele), we found strong evidence for replication:

11 of 14 (79%) infections of HbAS children carried the associated 'T' allele; while 11 of 13

(85%) HbAA infections carried the alternative 'A' allele; OR = 20, 95% CI = 2.8-145;

P=2.8x10^{-3}. We saw no evidence in these samples that total read counts (reflecting

expression levels) were associated with β^S status or with the chr4:1,121,472 T>A genotype.

Taken together, these results provide compelling evidence that the chr4:1,121,472 T>A SNP

is associated with β^S in mild malaria infections in west Africa, but raise questions as to why

the association is not also seen in the severe malaria data from Kenya and The Gambia

analysed previously, discussed further below. Following previous work, we refer to this new

locus as *P. falciparum* sickle-associated locus 4 (*Pfsa4*), and to alleles that are associated or

not associated with sickle as *Pfsa4+* ('T' allele, present in the 3D7 reference genome) and

Pfsa4- ('A' allele, alternative to the 3D7 reference genome), respectively. In this Ghanaian

cohort of mild malaria cases, the *Pfsa1+*, *Pfsa3+*, and *Pfsa4+* allele frequencies differed

markedly between sickle and non-sickle individuals (**Supplementary Figure 2**). For

example, *Pfsa4+* was present in 12/36 (33%) of individuals carrying at least one sickle allele,

compared with 3% of people with HbAA (**Supplementary Table 3**). After excluding missing or heterozygous calls for the *Pfsa* alleles, parasites from individuals carrying at least one sickle allele were around 17x more likely to carry at least one of the *Pfsa1+*, *Pfsa3+*, or *Pfsa4+* polymorphisms than to have none of these alleles present (22% vs 1.3%, respectively). Conversely, individuals carrying at least one sickle allele were around 11x more likely to be infected by parasites carrying at least one *Pfsa+* allele than by parasites that did not carry any of these alleles (53% vs 5.1%, respectively). Thus, despite the prevalence of sickle carriage among this Ghanaian cohort being approximately 3%, and the prevalence of *Pfsa1+, Pfsa3+* and *Pfsa4+* alleles in *P. falciparum* being around 4%, over half of the malaria infections encountered by sickle individuals were caused by parasites carrying at least one *Pfsa+* allele.

Because notable linkage disequilibrium (LD) exists between the putative *Pfsa4* locus and the previously identified *Pfsa1* and *Pfsa3* loci (discussed further below), there is the possibility that the association at *Pfsa4* is driven by LD to truly associated signals and does not represent an independent signal. To assess this, we reperformed the association analysis within the Ghanaian, and previously published Gambian and Kenyan datasets, using genotypes at the *Pfsa1* and *Pfsa3* loci as covariates (**Supplementary Figure 3**). The effect of HbS on *Pfsa4* genotype is almost completely diminished by the inclusion of *Pfsa1* or *Pfsa3* as covariates in Ghana. This might suggest that *Pfsa4* does not represent an independent association to HbS. However, inspecting this relationship at the other identified *Pfsa* loci, we observe that the signals of association at the *Pfsa1*, *Pfsa2* and *Pfsa3* loci also diminish to a similar degree when conditioned on *Pfsa1* and *Pfsa3* genotype within severe cases in Kenya. This is not the case in severe cases from The Gambia, however, where the HbS-*Pfsa* association is largely unchanged by the inclusion of *Pfsa* genotypes as covariates. This is likely to be explained by lower levels of LD between the *Pfsa1* and *Pfsa3* loci within The Gambia compared to within Kenya and Ghana. Therefore, although HbS-*Pfsa* associations do not appear statistically independent of one another, they are connected through prominent LD, despite being situated far apart in the *P. falciparum* genome across multiple chromosomes. This is challenging to explain without a biological relationship between them and HbS.

To explore whether the effects of *Pfsa* loci were additive, we computed the Ratio of Relative Risk (RRR) for different combinations of *Pfsa* genotypes occurring in HbS-carrying individuals (**Supplementary Methods**). The RRR increased in infections carrying *Pfsa1+*, *Pfsa3+* and *Pfsa4+* alleles combined, compared to those carrying only *Pfsa1+* and *Pfsa3+*. This analysis suggests that *Pfsa4+* does have an effect in improving the ability of *P.*

falciparum parasites to infect HbS individuals, compared to *Pfsa1+* and *Pfsa3+* alone (**Supplementary Figure 4**).

The *Pfsa4* locus

FIKK4.2 is a member of a larger family of FIKK serine/ threonine kinase proteins. It is exported to the erythrocyte cytosol during blood-stage infections, and, along with other FIKKs, is thought to regulate phosphorylation of erythrocyte and parasite proteins in the erythrocyte membrane. FIKK4.2 has also been associated with specific compartments in the infected red cell (termed K-dots [33]). In the 3D7 genome assembly, chr4:1,121,472 lies in the second exon of the *FIKK4.2* gene, and encodes a Leucine → Isoleucine change at amino acid 961. This is within the kinase domain, and is downstream of a lengthy segment of low complexity repeat sequences [33].

Specifically, an amino acid hexamer repeat occupies a significant proportion of FIKK4.2 and is located nearby to the top candidate SNP. The majority of repeats take the form SD[HNS]NH[KM]. We investigated this region by multiple sequence alignment using 16 high-quality *P. falciparum* reference genomes [34] (**Supplementary Methods**). FIKK4.2 varied in length between these isolates from 1,079 amino acids (isolate PfKH01) to 1,223 amino acids (isolate Pf3D7). This variation reflected expansion/ contraction of the repeat sequence, which varied from 402 to 546 amino acids (67 - 91 hexamer repeats). Notably, the Pf3D7 sequence, with 91 repeats, was substantially longer than that in any other assembly; the next longest being Pf7G8 (protein length 1,181; 84 hexamer repeats). The major cause of this increased length was a triplication of a DNA sequence in Pf3D7 of approximately 170bp, at positions 1,120,534-1,120,005; this repeat was only present in one or two copies in each of the other 15 genome assemblies (**Supplementary Figure 5**). In 3D7, this repeat unit ends at position 1,121,413, which is only 59 bases away from the lead *Pfsa4* SNP. Intriguingly, the *Pfsa4+* allele is present in 4/16 reference genomes, all of which contain two or three copies of the repeat region; whereas of the 12/16 isolates with the non-associated allele at *Pfsa4*, 11/12 only have a single copy of the repeat section ($P<0.05$ for 11/12 isolates having one copy, if expectation was for 12/12 to have two or three copies as per *Pfsa4+* isolates, exact Binomial test). This raises the possibility that structural variation within the hexamer repeat segment of FIKK4.2 is causally involved in the sickle association signal, and the *Pfsa4+* SNP could be a bystander**.**

Little evidence of association with HbC

One potential explanation for effects specific to central west African populations, could be

the presence of the haemoglobin C variant in these populations. We conducted a second

GWA analysis to test for association between parasite variants and individuals possessing at

least one HbC allele. To avoid confounding by the HbS association we excluded all

individuals carrying HbS alleles, leading to an analysis (after removing highly related

individuals) of $\beta^{C/no-S}$ (N=312; 301 $\beta^{A/C}$ and 11 $\beta^{C/C}$), vs $\beta^{A/A}$ individuals (N=962). Applying the

same methodology as with the sickle analysis, we performed logistic regression using

HPTEST. In contrast to HbS, we did not identify any strong signals of association with HbC

(**Figure 2, Supplementary Table 2**). Our study was adequately powered to detect HbC-

associations in the *P. falciparum* genome of a similar effect size to the HbS-*Pfsa*

associations; however, smaller effect sizes could be missed particularly with rarer alleles

(**Supplementary Methods**).

Population genetics of the *Pfsa* loci

We investigated linkage disequilibrium (LD) between the *Pfsa* loci. As expected, nearby

SNPs from the *Pfsa3* locus on chromosome 11 (chr11:1,058,035 T>A, chr11:1,057,437

T>C, and chr11:1,054,282 T>C) were all in high LD with each-other (r^2 0.76 – 0.95), as were

the four SNPs identified at the *Pfsa1* locus on chromosome 2 (chr2:630,545 C>T,

chr2:631,190 T>A, chr2:630,290 A>T, and chr2:629,375 C>G; r^2 0.91 – 0.98). The *Pfsa4*

SNP with strongest evidence of association with sickle (chr4:1,121,472 T>A) was in LD with

the nearby SNP identified in the GWA analyses (chr4:1,122,147 A>C, $r^2 = 0.84$). As noted by

Band *et al.*, *Pfsa1* and *Pfsa3* loci were in high LD with each-other despite being located on

different chromosomes. For example, the top hit SNPs from the *Pfsa1* and *Pfsa3* loci had an

r^2 of 0.62. *Pfsa4*, on chromosome 4, was also in relatively high LD with both *Pfsa1* and

Pfsa3 loci. For example, the top hit SNP at *Pfsa4* (chr4:1,121,472 T>A) had an r^2 of 0.22 and

0.24 for the top hit SNPs at the *Pfsa1* and *Pfsa3* loci, respectively. These data suggest that

Pfsa1+, *Pfsa3+* and *Pfsa4+* alleles are maintained together despite being located on

different chromosomes, consistent with having linked biological functions.

Next, we investigated the global distribution of *Pfsa1-4* alleles using the MalariaGEN Pf7

data resource [30]. Of the 20,864 samples in Pf7, we limited the analysis to samples that

passed MalariaGEN Pf7 QC (N=16,203), and to countries with $\geq$100 QC pass samples,

yielding an analysis set of 15,738 samples from 22 countries. The *Pfsa1-4+* alleles were

almost completely absent from Asia, but were present at varying frequencies in Africa and

the only South American country represented, Colombia (**Figure 5**). *Pfsa1+* and *Pfsa3+*

were distributed widely across Africa but with substantial heterogeneity in prevalence; for
example, the frequency of *Pfsa1+* in Benin was 59%, compared with 2% in Malawi
(**Supplementary Table 3**). *Pfsa4+* was essentially absent from east African populations,
and was at very low frequency in the Democratic Republic of the Congo (DRC), but was
found at much higher frequencies in west African populations (7% - 66%). The inverse
pattern was seen in *Pfsa2+*, which was prevalent in east Africa and the DRC but almost
completely absent from west Africa. Intriguingly, *Pfsa3+* and *Pfsa4+* were also present at
high frequencies (63% and 50%, respectively), and *Pfsa2+* at lower frequency (7%), in
Colombia.

There was evidence of haplotype sharing at each of the *Pfsa1-4* loci in parasites carrying the
sickle-associated alleles in the MalariaGEN Pf7 data resource, consistent with each *Pfsa+*
allele having a common evolutionary origin at each locus (**Supplementary Methods,**
Supplementary Figure 6). However, we did not find evidence that any of the *Pfsa1-4* loci
were under strong recent positive selection, based on integrated haplotype scores (iHS)
(**Supplementary Table 4**, **Supplementary Methods**).

Globally, in the Pf7 dataset, patterns of LD were complex and varied between populations
(illustrated by Ghana and Kenya in **Figure 5**; all countries shown in **Supplementary Figure**
7). In west Africa, where the combination of *Pfsa1+*, *Pfsa3+* and *Pfsa4+* was present, SNPs
at these loci had relatively high LD with each-other in Guinea, Ghana and Mali, but weaker
LD in Senegal and The Gambia. In east Africa, where the combination of *Pfsa1+*, *Pfsa2+*,
and *Pfsa3+* is present, these loci were in high LD with each-other in Malawi, Kenya and
Tanzania; but LD was lower for these loci in the DRC. In Colombia, *Pfsa3+* and *Pfsa4+* had
high LD ($r^2 = 0.49$).

In summary, these data reveal complex patterns of *Pfsa1-4+* allele frequencies and
associations between populations. *Pfsa1+* and *Pfsa3+* are present across Africa, *Pfsa2+* is
more localised to east Africa, and *Pfsa4+* more localised to west Africa; within these regions,
the extent to which *Pfsa+* alleles correlate with each-other is variable, ranging from very high
(e.g. r^2 of 0.67 for *Pfsa1+* and *Pfsa3+* in Ghana), to zero (e.g. *Pfsa3+* and *Pfsa4+* in Benin).

Discussion

We have investigated host-parasite interaction by searching for associations between variation in the *P. falciparum* genome with human variation in the β-globin gene in a population with mild malaria from northern Ghana. Two parasite loci previously associated with sickle haemoglobin in severe malaria cases from Kenya and The Gambia [29] were replicated in this cohort, termed *Pfsa1* and *Pfsa3*. A new locus of association on *P. falciparum* chromosome 4 was identified within the gene *FIKK4.2*, which we name *P. falciparum* sickle associated locus 4, or *Pfsa4*. This locus was not associated with sickle in the severe malaria cases from The Gambia, and the variant is absent in Kenya. However, the *Pfsa4+* association with sickle did replicate in a sample of mild malaria cases from Mali. There were no significant associations found between parasite SNPs and $β^C$ genotypes. The geographic distributions of the *Pfsa1-4* alleles vary with distinct patterns observed in western and eastern Africa, explored further in [35]. Haplotype sharing indicates a common evolutionary origin for each *Pfsa* locus. The lack of strong positive selection at these loci, and their broad geographic distributions, suggests they have not emerged in the very recent evolutionary past.

The high prevalence of $β^C$ and relatively low prevalence of $β^S$ observed in this study is broadly consistent with previous results. HbC has mainly been described in west Africa, and it has been hypothesised that the relatively more benign phenotype of HbCC compared with HbSS would lead to the gradual replacement of the $β^S$ allele by $β^C$ in west Africa [13]. The proportion of $β^S$ carriers we observed was lower than other studies in the region [36,37]. Around 2% (15,000) of Ghanaian newborns are diagnosed with sickle cell disease (SCD) annually [38,39]; the majority of patients with SCD attending a large teaching hospital in Accra were found to have HbSS (55.7%) or HbSC (39.6%) genotypes [38]. The relatively lower prevalence of $β^S$ alleles we observed may be partly because sickle-containing genotypes protect against malaria disease, so the prevalence of such genotypes in this population, who were selected for having mild malaria, would be lower than in an unselected population. Variation in HbS prevalence may also be due to resource limitations in available healthcare to support children born with SCD in rural populations versus urban centres in Accra.

The replication of the association between *Pfsa1+* and *Pfsa3+* alleles with sickle in people with mild malaria is consistent with the view that parasites possessing these variants can bypass the protective effects of sickle and cause both mild and severe malaria syndromes in sickle individuals more readily than parasites lacking these variants. On this view, the sickle-associated variants effectively open an ecological niche (individuals carrying $β^S$ alleles) that

is otherwise relatively closed off (or restricted) for *Pfsa-* parasites; rather than the view that these variants increase severity in sickle individuals *per se*. The age of the β^S mutation is uncertain, with estimates suggesting a single origin around 20,000 years ago (range: 13,000-70,000 years [40]). Selection for *Pfsa+* mutations may have occurred once β^S reached appreciable frequencies. It is currently unclear what has prevented *Pfsa+* parasites from outcompeting *Pfsa-* parasites in regions where HbS is common. The frequency of *Pfsa+* polymorphisms may depend on the regional prevalence of HbS, and the relative fitness advantage of *Pfsa+* alleles in HbS RBCs versus their disadvantage in HbA RBCs, compared with *Pfsa-* alleles. Moreover, sexual recombination during the mosquito phase would be expected to mix *Pfsa+* and *Pfsa-* alleles. The presence of high LD between the *Pfsa+* polymorphisms is consistent with an epistatic effect on parasite fitness counter-acting the effect of sexual recombination, to maintain the *Pfsa+* alleles together. It is unclear whether all of the *Pfsa+* alleles play a direct and independent role in assisting parasites to survive in HbS-carrying hosts, or whether some only mediate their effect in the context of other *Pfsa* alleles, for example to compensate for fitness costs. Further experimental work is required to explore these effects functionally.

The *Pfsa4* locus was found within the gene encoding the serine/ threonine kinase FIKK4.2. The FIKK kinase family (characterised by a phenylalanine (F) – isoleucine (I) – lysine (K) – lysine (K) motif) has expanded in the *Laverania* clade (which includes *P. falciparum* and other *Plasmodium* species that infect great apes), comprising around 19 proteins in *P. falciparum* distributed over 11 chromosomes. The majority of *P. falciparum* FIKK kinases are exported into the host Red Blood Cell (RBC), and each is associated with a unique 'phosphorylation fingerprint' of host RBC proteins [41]. There is evidence that FIKK kinases play a role in remodelling the RBC surface and altering the mechanical properties of infected RBCs [42,43]. RBCs infected by parasites lacking FIKK4.2 through experimental knockout were less rigid and less adhesive compared with RBCs infected by wild-type parasites despite unchanged levels of *P. falciparum* Erythrocyte Protein 1 (PfEMP1) expression, suggesting a role in malaria pathogenesis [33]. FIKK kinases have been proposed as potential targets for antimalarial drug design [44]. Intriguingly, the lead *Pfsa4+* SNP identified in our GWA analysis sits very close to a lengthy repeat region. Analysis of 16 high-quality *P. falciparum* genome assemblies tentatively suggests that the *Pfsa4+* SNP may be associated with increased copy number in this repeat section. Calling copy number variation (CNV) from short read sequencing data following parasite selective whole genome amplification (sWGA) [45] is challenging, so further work is required to assess this locus at population scale and to investigate functional effects of FIKK4.2 variants including CNV.

The lack of association between β^C alleles and parasite variation may reflect the fact that the

vast majority of individuals possessing a β^C allele were genotype $\beta^{A/C}$, which is thought to

confer a reduced protective effect against *P. falciparum* relative to $\beta^{C/C}$ [13], and likely also

less than $\beta^{A/S}$; the strength of selection acting on the parasites by $\beta^{A/C}$ individuals will

therefore be accordingly weaker. With only 12 $\beta^{C/C}$ individuals in our cohort, the sample size

is too small to analyse this sub-group. The effect specifically of the $\beta^{S/C}$ genotype on malaria

protection and *P. falciparum* genetic associations is, to our knowledge, unknown, as is the

effect of other compound heterozygotes such as β^S with thalassaemia.

This study has several limitations. First, the small number of individuals carrying β^C and β^S

alleles significantly limits the statistical power of the GWA analysis, meaning only variants

with large effect sizes could be detected. Associations with smaller effect sizes and/or

specific to certain β-globin genotypes such as $\beta^{C/C}$ will require larger sample sizes for

discovery. Second, the method of genotyping the human β-globin locus involved PCR

amplification and capillary sequencing a small region at the start of the β-globin gene and

did not account for other forms of genetic diversity in the haemoglobin genes. For example, it

is possible that some of the homozygous calls were in fact compound heterozygotes with a

β-thalassaemia allele. Although, complete β-globin gene deletion is an uncommon

mechanism of β-thalassaemia [46], and β-thalassaemia is less common in west Africa than

in other populations [23]. Third, minimal population demographic data were available to

control for potential confounders, such as participant ethnic group. However, the logistic

regression analysis was performed after removing samples that appeared highly similar, and

incorporated principal components as covariates to account for population structure. Fourth,

this study does not have a matched control group who did not have clinical malaria against

which to compare the β-globin genotype frequencies. This study therefore cannot directly

assess the extent to which *Pfsa+* alleles reduce the protective effect of sickle against mild

malaria relative to people without symptomatic malaria. However, their association with β^S

alleles in mild malaria, combined with the fact that the same variants reduce the protective

effect of sickle against severe malaria relative to population controls [29], supports the view

that these variants allow parasites to access sickle individuals which wild-type parasites

otherwise struggle to infect, and therefore reduce the protective effect of sickle against both

mild and severe malaria. The GWA analysis was limited to high-confidence SNPs; CNV and

other more complex forms of genetic variation were not included, due to the risk of spurious

calls from short read post-sWGA data. Future work can explore these challenging forms of

genetic variation further, potentially leveraging long read sequencing platforms and

functional experimentation. This is particularly relevant given the potential link between CNV

and the lead *Pfsa4+* SNP identified in our study.

In conclusion, this study adds further evidence that several loci in the *P. falciparum* genome

are associated with sickle haemoglobin, in both mild and severe malaria. Variants at these

loci are maintained in linkage with each-other despite being on separate chromosomes. This

suggests that in a population, subsets of *P. falciparum* parasites will inherit genetic traits that

make them better suited to particular human genetic contexts, such as 'specialists' at

infecting sickle RBCs. The frequencies of different host genotypes, and the relative fitness

costs of the polymorphisms for the parasite in non-sickle hosts, will sculpt the evolutionary

pressures acting on these loci in the parasite, which may in turn affect the selection

pressures acting on the human genome in an ancient coevolutionary process. Future work

could extend these results by recruiting a larger population to ensure robust statistical power

to analyse different haemoglobin genotypes, incorporate more genetic variation from both

human and *P. falciparum* genomes, and establish functional data to uncover the

mechanisms of how these variants help the parasite to infect people carrying sickle

haemoglobin alleles.

16
17

Online methods

Study site and samples

Samples were collected from the Kassena-Nankana East Municipality and Kassena-Nankana West District, two adjoining administrative areas in the Upper East Region of northern Ghana, hereafter referred to as the KNDs. The population is primarily rural, and malaria transmission is seasonal, peaking during the rainy season from July to October. *P. falciparum* genetic diversity has been studied in this region previously [47] through a network of community health centres.

This study ran from 22/07/2015 to 30/08/2018, encompassing the rainy seasons for 2015, 2016 and 2017. Patients presenting to the Navrongo War Memorial Hospital (NWMH) and four satellite clinics in the KNDs with compatible symptoms for malaria were tested using an antigen Rapid Diagnostic Test (RDT) (CareStart™ malaria Pf (HRP2), Access Bio, NJ, USA). Participants were eligible if they tested positive for *P. falciparum* by RDT and informed consent was obtained and documented. Written consent forms were available in the predominant languages spoken in the region: Kasem, Nankam, and English. Four 50uL dried blood spots (DBS) and malaria blood smears for microscopy were collected from consenting participants. Parasitaemia was calculated by microscopy as asexual parasites per 200 white blood cells (WBC). DBS samples were transferred to the Wellcome Sanger Institute (WSI), UK, for DNA extraction and sequencing. DNA extraction from DBS was performed using the QIAamp Investigator Biorobot kit on the Qiagen Biorobot Universal instrument, protocol described in [48].

Parasite whole genome sequencing and genotype calling

P. falciparum whole genome sequencing from DBS extractions was performed using the MalariaGEN pipeline established at the Wellcome Sanger Institute (WSI), UK, described elsewhere [30]. Briefly, DBS samples undergo selective whole genome amplification (sWGA) to enrich for *P. falciparum* DNA [49], followed by short-read Illumina sequencing on the HiSeq platform. These samples are part of the broader MalariaGEN *P. falciparum* Community Project [30,50,51]. Genotype calling from the Illumina sequence data was performed at WSI using the MalariaGEN informatics pipeline [30]. All of the parasite genomes included in this study are available via the current iteration of the MalariaGEN *P. falciparum* Community Project, Pf7 [30].

Human haemoglobin genotyping

Samples with ethical approval for use in this study (a subset of the MalariaGEN *P. falciparum* Community Project collection) were transferred via Beckman robot to new 96-well

plate layouts organised by sample concentration (measured by Qubit High Sensitivity (HS)
assay) and remaining volume after parasite sequencing. Human β-globin genotyping at
codon 6 followed a PCR amplification and capillary sequencing strategy. Details on the
primers and PCR conditions used can be found in the **Supplementary Methods**. The PCR
products were capillary sequenced by the Eurofins company, using their PCR clean-up and
Supreme Run service.

Capillary sequence chromatogram traces were visually inspected using the 4Peaks software
tool (**Supplementary Methods**). The six callable β-globin genotypes from the sixth codon
were: $\beta^{A/A}$, $\beta^{A/S}$, $\beta^{S/S}$, $\beta^{S/C}$, $\beta^{A/C}$, and $\beta^{C/C}$. Samples with ambiguous calls (eg. low amplitude
chromatograph or chaotic uninterpretable signal) were repeat PCR-amplified and re-
sequenced if available; if these samples remained ambiguous after the second attempt (or if
there was no sample left for further attempts) then the sample was deemed to have failed β-
globin genotyping. All $\beta^{A/S}$, $\beta^{S/S}$, $\beta^{S/C}$, and $\beta^{C/C}$ samples were repeat PCR-amplified and re-
sequenced to confirm the result; there were no samples for which the second run disagreed
with the first run if a call was successfully made on both runs.

P. falciparum genome variant filtering for study population
Parasite genomic analyses were conducted using the data structures and curated sample
metadata of the MalariaGEN Pf7 data resource [30]. Full details of variant filtering steps are
described in **Supplementary Methods**. Briefly, the approximately 10 million variants in Pf7
were initially filtered to create a 'working set' of variants, including only biallelic SNPs that
passed Pf7 Quality Control (QC) and had at least 0.1% allele frequency in the Ghana study
population, yielding 123,845 SNPs. Heterozygous genotypes (representing mixed infections
in *P. falciparum*) were regarded as missing. The previously reported lead associations in the
Pfsa regions were explicitly included. This 'working set' of 123,845 SNPs was then further
filtered for different analyses; the main logistic regression GWA analysis described in main
text included 18,770 SNPs with a minor allele count of at least 10 among the 1,309 samples
included in the analysis (after removing closely related samples). Genomic data
manipulations from the Pf7 data resource were performed in Python; genotype and
haplotype manipulations and downstream population genetic analyses were performed using
the *scikit-allel* package.

Population structure analysis
We used principal components analysis (PCA), as implemented in the QCTOOL package
[52] to assess population structure in the 1,368 samples passing QC and with both parasite
and β-globin genotypes in our data (further details in **Supplementary Methods**). For this

analysis, we restricted attention to the set of 14,381 SNPs (selected from the 'working set' of

123,845 SNPs) with at least 1% minor allele frequency and no more than 25% missing

genotypes in these samples. Initial principal components (PCs) (**Supplementary Figure 1**)

revealed a small number of sample pairs with high relatedness values (estimated r > 0.5)

that dominated the first two principal components. We therefore implemented a greedy

algorithm to remove one of each sample pair having $r_{ij} > 0.5$. In total we removed 59

samples, leaving 1,309. We then re-computed principal components and loadings

(**Supplementary Figure 1)**.

Genome Wide Association

We performed a logistic regression analysis using the program HPTEST [29], which fits a

logistic regression model with the host genotype as predictor and the parasite genotype as

outcome, treating each variant separately (further details in **Supplementary Methods**).

HPTEST reports a Bayes Factor reflecting the evidence for association as well as other

summaries including an effect size estimate, standard error, and *P*-value. Interpretation of

HPTEST results was previously discussed in detail [29]. For this analysis, we used the

1,309 samples without closely related parasite samples identified through PCA, either with or

without including the first 3 principal components.

Population genetics of the *Pfsa* loci

Allele frequencies and linkage disequilibrium (LD) for the *Pfsa* loci were computed in Python

using the *scikit-allel* package. For the analysis of global patterns of *Pfsa* frequencies and LD,

the total MalariaGEN Pf7 data resource (20,864 samples) was restricted to samples that

passed Pf7 QC (N=16,203), and to countries with ≥100 QC pass samples, yielding 15,738

samples from 22 countries. One SNP was selected to represent each *Pfsa* locus for

frequency and LD calculations: *Pfsa1* = chr2:631,190; *Pfsa2* = chr2:814,288; *Pfsa3* =

chr11:1,058,035; *Pfsa4* (newly reported in this study) = chr4:1,121,472.

To analyse patterns of haplotype sharing at the *Pfsa* loci, we focussed on a 10kb section of

the *P. falciparum* genome centred on the lead sickle-associated variant (identified as the

variant with the highest Bayes factor) at each of the *Pfsa1-3* loci, using the MalariaGEN Pf7

data resource. At *Pfsa4*, we adjusted this to only include 7kb due to the higher density of

variants at this locus. Ancestral vs. derived allele status at the *Pfsa* loci was determined as

described in [35]. Further details on Pf7 variant filtering is described in **Supplementary**

Methods. Genotypes were parsed into python code using the package *scikit-allel* (version

1.3.11) and clustered using the package *scipy* (version 1.14.1). Initially, the haplotype matrix

was transposed so that each row represented a sample, and each column represented a

variant. Hierarchical clustering was then performed using the average linkage method, which calculated the distance between clusters as the average distance between all pairs of genotypes in the clusters. The function used the resulting dendrogram to determine the order of the samples, reordered the haplotype matrix and the samples array accordingly, and returned the reordered haplotype matrix and samples array.

Ethics

Ethical approval for the study was granted by the Navrongo Health Research Centre Institutional Review Board (NHRCIRB203). All participants or their guardians (as appropriate) were provided detailed information sheets and gave informed consent prior to enrolment. Further approval was granted by the Wellcome Sanger Institute's Research Ethics Committee for the analysis of the samples. Sample collection was conducted by LNA, who pseudonymised the samples prior to making them available to the Wellcome Sanger Institute. All patient-identifiable data are securely stored by LNA and only non-patient identifiable were provided to the Wellcome Sanger Institute. The study complies with all relevant ethical regulations.

Data availability statement

The *P. falciparum* variant calls (VCF file) are available to download open-access via the MalariaGEN *P. falciparum* Community Project Pf7 resource. (The VCF file contains variant calls for all samples in Pf7, which includes all of the samples analysed in this study). The β-globin genotype calls linked to Pf7 sample codes are available in **Supplementary Table 6**. The filtered parasite genotype calls derived from Pf7 and associated human beta-globin genotype calls can also be downloaded from FigShare, details below:

Hamilton, William (2025). Analysis datasets used for Pfsa4 genome-wide association analysis (Hamilton et al).. figshare. Dataset. https://doi.org/10.6084/m9.figshare.30705047

Code availability statement

Source code for QCTOOL and HPTEST is available at: code.enkre.net/qctool, and on Zenodo (doi:10.5281/zenodo.5685581).

Inclusion and ethics statement

This research was conducted collaboratively between researchers based at the University of Ghana, the Navrongo Health Research Centre (NHRC), the Wellcome Sanger Institute, and University of Oxford. Study conceptualisation, data analysis and interpretation, and authorship of publications were shared jointly between personnel based in Ghana and the UK.

Acknowledgements

We thank the teams in the Navrongo Health Research Centre (NHRC) and the Wellcome Sanger Institute (WSI) who generated the *P. falciparum* genome sequence data as part of the MalariaGEN *P. falciparum* Community Project, and the study participants and their guardians who provided the samples. Thank you to Richard Pearson for his help accessing and using the Pf7 data resource, and to Christen Smith, Sonia Goncalves and Rachel Ogidan for support ensuring legal and ethical compliance. Thank you to Kevin Esoh for contributions to discussions of *P. falciparum* population genetics. This work is dedicated to the memory of Professor Dominic Kwiatkowski, who died prior to its completion.

Bibliography

1. World Health Organization. World Malaria Report 2022 [Internet]. 2022 [cited 2022 Dec 9]. https://doi.org/ISBN 978 92 4 1564403

2. Kwiatkowski DP. How malaria has affected the human genome and what human genetics can teach us about malaria. Am J Hum Genet. 2005;77:171–92. https://doi.org/10.1086/432519

3. Hamblin MT, Di Rienzo A. Detection of the signature of natural selection in humans: Evidence from the Duffy blood group locus. Am J Hum Genet. 2000;66:1669–79. https://doi.org/10.1086/302879

4. Williams TN, Maitland K, Bennett S, Ganczakowski M, Peto TEA, Newbold CI, et al. High incidence of malaria in α-thalassaemic children. Nature. 1996;383:522–5. https://doi.org/10.1038/383522a0

5. Allen SJ, O'Donnell A, Alexander NDE, Alpers MP, Peto TEA, Clegg JB, et al. α+-Thalassemia protects children against disease caused by other infections as well as malaria. Proc Natl Acad Sci U S A. 1997;94:14736–41. https://doi.org/10.1073/pnas.94.26.14736

6. Weatherall D, Clegg J, Kwiatkowski D. The role of genomics in studying genetic susceptibility to infectious disease. Genome Res. 1997;7:967–73. https://doi.org/10.1101/gr.7.10.967

7. Flint J, Hill AVS, Bowden DK, Oppenheimer SJ, Sill PR, Serjeantson SW, et al. High frequencies of α-thalassaemia are the result of natural selection by malaria. Nature. 1986;321:744–50. https://doi.org/10.1038/321744a0

8. Yenchitsomanus P, Summers KM, Board PG, Bhatia KK, Jones GL, Johnston K, et al. Alpha-thalassemia in Papua New Guinea. Hum Genet. 1986;74:432–7.

9. Taylor SM, Parobek CM, Fairhurst RM. Haemoglobinopathies and the clinical epidemiology of malaria: A systematic review and meta-analysis. Lancet Infect Dis. Elsevier Ltd; 2012;12:457–68. https://doi.org/10.1016/S1473-3099(12)70055-5

10. Tishkoff SA, Varkonyi R, Cahinhinan N, Abbes S, Argyropoulos G, Destro-Bisol G, et al. Haplotype diversity and linkage disequilibrium at human G6PD: Recent origin of alleles that confer malarial resistance. Science (1979). 2001;293:455–62. https://doi.org/10.1126/science.1061573

11. Carter R, Mendis KN. Evolutionary and historical aspects of the burden of malaria. Clin Microbiol Rev. 2002;15:564–94. https://doi.org/10.1128/CMR.15.4.564-594.2002

12. RAY RN, CHATTERJEA JB, CHAUDHURI RN. Observations on the Resistance in Hb E Thalassaemia Disease To Induced. Bull World Health Organ. 1964;30:51–5.

13. Modiano D, Luoni G, Sirima BS, Simporé J, Verra F, Konaté A, et al. Haemoglobin C protects against clinical Plasmodium falciparum malaria. Nature. 2001;414:305–8. https://doi.org/10.1038/35104556

14. Siniscalco M, Bernini L, Filippi G, Latte B, Meera Khan P, Piomelli S, et al. Population genetics of haemoglobin variants, thalassaemia and glucose-6-phosphate dehydrogenase deficiency, with particular reference to the malaria hypothesis. Bull World Health Organ. 1966;34:379–93.

15. Allison AC. The discovery of resistance to malaria of sickle-cell heterozygotes. Biochemistry and Molecular Biology Education. 2002;30:279–87. https://doi.org/10.1002/bmb.2002.494030020034

16. Allison AC. The distribution of the sickle-cell trait in East Africa and elsewhere, and its apparent relationship to the incidence of subtertian malaria. Trans R Soc Trop Med Hyg. 1954;48:312–8. https://doi.org/10.1016/0035-9203(54)90101-7

17. Allison AC. Protection Afforded by Sickle-cell Trait Against Subtertian Malarial Infection. Br Med J. 1954;1:290 LP – 294. https://doi.org/10.1136/bmj.1.4857.290

18. Zimmerman PA, Woolley I, Masinde GL, Miller SM, McNamara DT, Hazlett F, et al. Emergence of FY*Anull in a Plasmodium vivax-endemic region of Papua New Guinea. Proc Natl Acad Sci U S A. 1999;96:13973–7. https://doi.org/10.1073/pnas.96.24.13973

19. Rockett KA, Clarke GM, Fitzpatrick K, Hubbart C, Jeffreys AE, Rowlands K, et al. Reappraisal of known malaria resistance loci in a large multicenter study. Nat Genet. 2014;46:1197–204. https://doi.org/10.1038/ng.3107

20. MalariaGEN. Insights into malaria susceptibility using genome-wide data on 17,000 individuals from Africa, Asia and Oceania. Nat Commun. 2019;10:5732. https://doi.org/10.1038/s41467-019-13480-z

21. Mackinnon MJ, Mwangi TW, Snow RW, Marsh K, Williams TN. Heritability of malaria in Africa. PLoS Med. 2005;2:1253–9. https://doi.org/10.1371/journal.pmed.0020340

22. Weatherall DJ. Phenotype-genotype relationships in monogenic disease: Lessons from the thalassaemias. Nat Rev Genet. 2001;2:245–55. https://doi.org/10.1038/35066048

23. Williams TN, Weatherall DJ. World distribution, population genetics, and health burden of the hemoglobinopathies. Cold Spring Harb Perspect Med. 2012;2:1–14. https://doi.org/10.1101/cshperspect.a011692

24. Piel FB, Patil AP, Howes RE, Nyangiri OA, Gething PW, Dewi M, et al. Global epidemiology of Sickle haemoglobin in neonates: A contemporary geostatistical model-based map and population estimates. The Lancet. Elsevier Ltd; 2013;381:142–51. https://doi.org/10.1016/S0140-6736(12)61229-X

25. Sundd P, Gladwin MT, Novelli EM. Pathophysiology of Sickle Cell Disease. Annual Review of Pathology: Mechanisms of Disease. 2019;14:263–92. https://doi.org/10.1146/annurev-pathmechdis-012418-012838

26. Archer NM, Petersen N, Clark MA, Buckee CO, Childs LM, Duraisingh MT. Resistance to Plasmodium falciparum in sickle cell trait erythrocytes is driven by oxygen-dependent growth inhibition. Proc Natl Acad Sci U S A. 2018;115:7350–5. https://doi.org/10.1073/pnas.1804388115

27. Piel FB, Howes RE, Patil AP, Nyangiri OA, Gething PW, Bhatt S, et al. The distribution of haemoglobin C and its prevalence in newborns in Africa. Sci Rep. 2013;3:1–8. https://doi.org/10.1038/srep01671

28. Weatherall DJ, Clegg JB. The thalassaemia syndromes. Oxford: Blackwell Scientific Publications.; 2001.

29. Band G, Leffler EM, Jallow M, Sisay-Joof F, Ndila CM, Macharia AW, et al. Malaria protection due to sickle haemoglobin depends on parasite genotype. Nature. 2021; https://doi.org/10.1038/s41586-021-04288-3

30. MalariaGEN. Pf7: an open dataset of Plasmodium falciparum genome variation in 20,000 worldwide samples. Wellcome Open Res. 2023;8. https://doi.org/https://doi.org/10.12688/wellcomeopenres.18681.1

31. Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D. FaST linear mixed models for genome-wide association studies. Nat Methods. 2011;8:833–5. https://doi.org/10.1038/nmeth.1681

32. Saelens JW, Petersen JE V., Freedman E, Moseley RC, Konaté D, Diakité SAS, et al. Impact of Sickle Cell Trait Hemoglobin on the Intraerythrocytic Transcriptional Program of Plasmodium falciparum. mSphere. American Society for Microbiology; 2021;6. https://doi.org/10.1128/msphere.00755-21

33. Kats LM, Fernandez KM, Glenister FK, Herrmann S, Buckingham DW, Siddiqui G, et al. An exported kinase (FIKK4.2) that mediates virulence-associated changes in Plasmodium falciparum-infected red blood cells. Int J Parasitol. Australian Society for Parasitology Inc.; 2014;44:319–28. https://doi.org/10.1016/j.ijpara.2014.01.003

34. Otto TD, Böhme U, Sanders M, Reid A, Bruske EI, Duffy CW, et al. Long read assemblies of geographically dispersed Plasmodium falciparum isolates reveal highly structured subtelomeres [version 1; referees: 3 approved]. Wellcome Open Res. 2018;3. https://doi.org/10.12688/wellcomeopenres.14571.1

35. Python A, Forster AJ, Todd P, Niaré K, Conrad MD, Amenga-Etego LN, et al. Geographical variation drives adaptive equilibrium of the P. falciparum sickle-associated mutations. 2025; https://doi.org/10.1101/2025.08.31.672853

36. Ringelhann B, Dodu SRA, Konotey-Ahulu FID, Lehmann H. A survey for haemoglobin variants, thalassaemia and glucose-6-phosphate dehydrogenase deficiency in northern Ghana. Ghana Med J. 1968;7:120–4.

37. Mensah-Brown HE, Abugri J, Asante KP, Dwomoh D, Dosoo D, Atuguba F, et al. Assessing the impact of differences in malaria transmission intensity on clinical and haematological indices in children with malaria. Malar J. BioMed Central; 2017;16:1–11. https://doi.org/10.1186/s12936-017-1745-8

38. Asare E V., Wilson I, Benneh-Akwasi Kuma AA, Dei-Adomakoh Y, Sey F, Olayemi E. Burden of Sickle Cell Disease in Ghana: The Korle-Bu Experience. Adv Hematol. 2018;2018. https://doi.org/10.1155/2018/6161270

39. Ohene-Frempong K, Oduro J, Tetteh H, Nkrumah F. Screening newborns for sickle cell disease in Ghana. Pediatrics. 2008;121:S120–1. https://doi.org/10.1542/peds.2007-2022UUU

40. Laval G, Peyrégne S, Zidane N, Harmant C, Renaud F, Patin E, et al. Recent Adaptive Acquisition by African Rainforest Hunter-Gatherers of the Late Pleistocene Sickle-Cell Mutation Suggests Past Differences in Malaria Exposure. Am J Hum Genet. 2019;104:553–61. https://doi.org/10.1016/j.ajhg.2019.02.007

41. Davies H, Belda H, Broncel M, Ye X, Bisson C, Introini V, et al. An exported kinase family mediates species-specific erythrocyte remodelling and virulence in human malaria. Nat Microbiol. Springer US; 2020;5:848–63. https://doi.org/10.1038/s41564-020-0702-4

42. Nunes MC, Goldring JPD, Doerig C, Scherf A. A novel protein kinase family in Plasmodium falciparum is differentially transcribed and secreted to various cellular compartments of the host cell. Mol Microbiol. 2007;63:391–403. https://doi.org/10.1111/j.1365-2958.2006.05521.x

43. Nunes MC, Okada M, Scheidig-Benatar C, Cooke BM, Scherf A. Plasmodium falciparum FIKK kinase members target distinct components of the erythrocyte membrane. PLoS One. 2010;5:1–8. https://doi.org/10.1371/journal.pone.0011747

44. Belda H, Bradley D, Christodoulou E, Nofal SD, Broncel M, Jones D, et al. The fast-evolving FIKK kinase family of Plasmodium falciparum can be inhibited by a single compound. Nat Microbiol. Nature Research; 2025;10:1463–83. https://doi.org/10.1038/s41564-025-02017-4

45. Oyola SO, Ariani C V., Hamilton WL, Kekre M, Amenga-Etego LN, Ghansah A, et al. Whole genome sequencing of Plasmodium falciparum from dried blood spots using selective whole genome amplification. Malar J. BioMed Central; 2016;15:597. https://doi.org/10.1186/s12936-016-1641-7

46. Thein SL. Molecular basis of β thalassemia and potential therapeutic targets. Blood Cells Mol Dis. Elsevier; 2018;70:54–65. https://doi.org/10.1016/j.bcmd.2017.06.001

47. Amenga-Etego LN, Asoala V, Agongo G, Jacob C, Goncalves S, Awandare GA, et al. Temporal evolution of sulfadoxine-pyrimethamine resistance genotypes and genetic diversity in response to a decade of increased interventions against Plasmodium falciparum in northern Ghana. Malar J. BioMed Central; 2021;20:1–11. https://doi.org/10.1186/s12936-021-03693-3

48. Girgis ST, Adika E, Nenyewodey FE, Jnr DKS, Ngoi JM, Lorenz O, et al. Nanopore sequencing for real-time genomic surveillance of Plasmodium falciparum. bioRxiv. 2022;2022.12.20.521122.

49. Oyola SO, Ariani C V., Hamilton WL, Kekre M, Amenga-Etego LN, Ghansah A, et al. Whole genome sequencing of Plasmodium falciparum from dried blood spots using selective whole genome amplification. Malar J. BioMed Central; 2016;15:597. https://doi.org/10.1186/s12936-016-1641-7

50. MalariaGEN. An open dataset of Plasmodium falciparum genome variation in 7,000 worldwide samples [version 1; peer review: 2 approved]. Wellcome Open Res. 2021;6:42.

51. Achidi EA, Agbenyega T, Allen S, Amodu O, Bojang K, Conway D, et al. A global network for investigating the genomic epidemiology of malaria. Nature. 2008;456:732–7. https://doi.org/10.1038/nature07632

52. Band G, Rockett KA, Spencer CCA, Kwiatkowski DP, Si Le Q, Clarke GM, et al. A novel locus of resistance to severe malaria in a region of ancient balancing selection. Nature. 2015;526:253–7. https://doi.org/10.1038/nature15390

53. Miotto O, Amato R, Ashley EA, Macinnis B, Almagro-Garcia J, Amaratunga C, et al. Genetic architecture of artemisinin-resistant Plasmodium falciparum. Nat Genet. 2015;47:226–34. https://doi.org/10.1038/ng.3189

54. Amato R, Lim P, Miotto O, Amaratunga C, Dek D, Pearson RD, et al. Genetic markers associated with dihydroartemisinin–piperaquine failure in Plasmodium falciparum malaria in Cambodia: a genotype–phenotype association study. Lancet Infect Dis. The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license; 2017;17:164–73. https://doi.org/10.1016/S1473-3099(16)30409-1

55. Zhu L, Tripathi J, Rocamora FM, Miotto O, van der Pluijm R, Voss TS, et al. The origins of malaria artemisinin resistance defined by a genetic and transcriptomic background. Nat Commun. 2018;9:1–13. https://doi.org/10.1038/s41467-018-07588-x

Figure legends

Figure 1. Study flow diagram.

Figure 2. Evidence for association between haemoglobin variants and *P. falciparum*
genetic variants in Ghanaian mild malaria cases. Plot shows the log10 Bayes factor for
association with HbS (top row) and HbC (bottom row) (y axis), for SNPs across the parasite
genome (x axis).

Figure 3. **Discovery and replication evidence for association between HbS and *Pfsa4*.**
The left hand side plot depicts the log10 Bayes factor for association with HbS (y axis) for
SNPs in the parasite genome zoomed onto the *Pfsa4* locus (x axis). The top hit is SNP
chr4:1,122,147, 'T' allele. The right hand plot shows point estimates with 95% confidence
intervals for the Pfsa4-HbS association, estimated using logistic regression from across
multiple studies. The populations are: Gambian severe cases from [29], Kenyan severe
cases from [29], Ghanaian cases from this study, and Malian cases from [32]. Text indicates
the 2-sided *P*-value for each point.

Figure 4. Global distribution of *Pfsa1-4+* alleles in the MalariaGEN Pf7 data resource.
Data from 15,738 samples from 22 countries, which passed Pf7 QC filters and requiring
$\geq$100 samples for each country. Countries are organised left to right approximately from
West to East Africa, with Colombia as the only non-African country represented. The *Pfsa+*
alleles were very low or absent from all Asian countries in Pf7, not shown on Figure.

Figure 5. Frequencies and linkage disequilibrium (LD) of *Pfsa1-4+* alleles in Ghana
(top) and Kenya (bottom). These well-sampled countries are shown to illustrate the distinct
patterns of *Pfsa1-4* population genetics observed in different African countries.

Tables

HBB genotype	Samples with β-globin call	Samples with β-globin call (%)	Samples β-globin called & Pf7 pass	Samples β-globin called & Pf7 pass (%)
HbAA	1311	73	**1008**	**74**
HbAS	50	2.8	**30**	**2.2**
HbSS	1	0.055	**1**	**0.073**
HbSC	7	0.39	**5**	**0.37**
HbCC	21	1.2	**12**	**0.88**
HbAC	417	23	**312**	**23**
Total	1807	100	**1368**	**100**

3   **Table 1.** Counts of the different β-globin genotypes in all samples that had β-globin
4   genotype calls available (N=1,807) and those that also had *P. falciparum* genomic data
5   available that passed MalariaGEN Pf7 quality control filters (N=1,368). A full breakdown of
6   sample counts by region, year and genotype is shown in **Supplementary Table 5.**
7