MTOR as a selectable genomic harbor for CRISPR-engineered CAR-T cell therapy

Sébastien Levesque[1,2,3], Gillian Carleton[4,5], Victoria Duque[1,2], Claudia Goupil[1,2], Jean-Philippe Fiset[1,2], Sarah Villeneuve[1,2], Eric Normandeau[6], Geneviève Morin[6], Nellie Dumont[7], Josée Laganière[7], Brian Boyle[6], Julian J. Lum[4,5], Yannick Doyon[1,2*]

[1]Centre Hospitalier Universitaire de Québec Research Center - Université Laval, Québec, QC, G1V 4G2, Canada.
[2]Université Laval Cancer Research Centre, Québec, QC, G1V 0A6, Canada.
[3]Present Address: Division of Hematology/Oncology, Boston Children's Hospital. Department of Pediatric Oncology, Dana-Farber Cancer Institute. Harvard Stem Cell Institute. Broad Institute of MIT and Harvard. Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA.
[4]Trev and Joyce Deeley Research Centre, BC Cancer, Victoria, BC, V8R 6V5, Canada.
[5]Department of Biochemistry and Microbiology, University of Victoria, Victoria, BC, V8P 5C2, Canada.
[6]Institut de Biologie Intégrative des Systèmes, Université Laval, Québec, QC, G1V 0A6.
[7]Medical Affairs and Innovation, Héma-Québec, Québec, Quebec, Canada

*Correspondence:
Yannick Doyon, Ph.D.
Centre de recherche du CHU de Québec – Université Laval
2705, boulevard Laurier, T-3-67
Québec, QC G1V 4G2
CANADA

Tel: 418-525-4444 ext. 46264
e-mail: Yannick.Doyon@crchudequebec.ulaval.ca
ORCID 0000-0002-3889-9563

ABSTRACT

Chimeric antigen receptors (CARs) reprogram T cells to recognize and target cancer cells. Despite remarkable responses observed with CAR-T cell therapy in patients with hematological malignancies, CAR-T cell engineering still relies mostly on randomly integrating vectors, limiting the possibilities of fine-tuning T cell function. Here, we designed a CRISPR-based marker-free selection strategy to simultaneously target a therapeutic transgene and a gain-of-function mutation to the *MTOR* locus to enrich cells resistant to rapamycin, a clinically used immunosuppressant. We readily engineered rapamycin-resistant (RapaR) CAR-T cells by targeting CAR expression cassettes to the *MTOR* locus. Using *in vitro* cytotoxicity assays, and a humanized mouse model of acute lymphoblastic leukemia, we show that RapaR-CAR-T cells can efficiently target CD19[+] leukemia cells in presence of rapamycin. Furthermore, our strategy allows multiplexed targeting of rapamycin-regulated immunoreceptors complexes (DARICs) to the *MTOR* and *TRAC* loci to pharmacologically control CAR-T cells' activity. We foresee that our approach could both facilitate the enrichment of CRISPR-engineered CAR-T cells *ex vivo* and *in vivo* while improving tumor eradication.

INTRODUCTION

Adoptive cell therapy with chimeric antigen receptor T (CAR-T) cells has shown remarkable potency against relapsed and refractory acute lymphoblastic leukemia (ALL) and non-Hodgkin lymphoma[1–5]. CAR-T cells are genetically engineered T cells that are able to recognize and target cancer cells expressing a specific antigen on their surface[6,7]. Randomly integrating vectors based on retroviruses, lentiviruses or *piggyBac* transposons are routinely used for cell therapy applications, but genome editing technologies offer a wide a range of opportunities to enhance CAR-T cells functionality[8,9]. The promising potential of CRISPR-based technologies to engineer CAR-T cells has been demonstrated by targeting a CAR to the *TRAC* locus to avert tonic CAR signaling and delay T cells differentiation and exhaustion[10], or by generating HLA-independent T cell receptors (HIT) to afford high antigen sensitivity[11]. CRISPR-based technologies also offer great potential to inactivate key genes involved in T cells dysfunction[12–14] or to generate allogeneic universal CAR-T cells[15–17].

While progress has been made towards high-yield targeted transgene integration in primary cells[10,11,18], enrichment strategies could further facilitate the development of CRISPR-engineered cell therapies. To enrich targeted transgene integration via homology-directed repair (HDR) in primary T cells, a selection method based on the expression of a methotrexate-resistant dihydrofolate reductase mutant has previously been reported[19]. A metabolic safety switch has also been developed via *UMPS* knockout to enrich T cells expressing reporter transgenes[20]. However, no marker-free selection strategy has been developed to simultaneously enrich CRISPR-engineered CAR-T cells and enhance their antitumor activity.

The serine/threonine protein kinase mTOR constitutes the catalytic subunit of two distinct complexes, known as mTORC1 and mTORC2, which coordinate eukaryotic cell growth and metabolism by integrating a diverse set of environmental inputs[21–23]. mTORC1 controls cellular growth and metabolism by sensing nutrient levels and growth factor signals to coordinate anabolic and catabolic metabolism[24]. While mTORC1 is sensitive to rapamycin, an immunosuppressant also used in oncology, mTORC2 remains functional upon acute rapamycin treatment[21–23]. Considering that mTOR signaling is commonly activated in tumor cells and alters the expression and the activity of key metabolic enzymes[21,25,26], combinatorial CAR-T cell therapy with rapamycin could synergistically enhance antitumor activity. Indeed, random integration of a rapamycin-resistant mTOR transgene and a CD19-CAR in primary T cells using *piggyBac* transposons has been shown to confer a selective growth advantage in the presence of rapamycin and enhanced antitumor activity when used in combination[27]. However, delivery of this large transgene is limiting, and product-derived lymphomas have been reported with *piggyBac*-modified CAR-T cells[28,29], so editing the endogenous *MTOR* locus to generate dominant cellular resistance to rapamycin using CRISPR could be a viable alternative. Since mTORC1 signaling lies downstream of several activation signals essential for T cell activation and expansion[21,30], this central signaling node is a prime target to develop a tissue-specific marker-free selection strategy for CRISPR-engineered CAR-T cells.

In this study, we performed saturation prime editing[31] at the *MTOR* locus to identify and characterize optimal mutations conferring rapamycin resistance. We then devised an intron nesting strategy[32,33] to enrich CRISPR-engineered cells *ex vivo* and *in vivo* without using an exogenous

selection marker. We demonstrate that targeting CAR expression cassettes to the *MTOR* locus

allows the selection and regulation of rapamycin-resistant CAR-T cells to perform combinatorial

immunotherapy with rapamycin. More broadly, we propose a versatile approach to facilitate

targeted therapeutic transgene integration for cell therapy applications.

RESULTS

Identification and characterization of mutations conferring rapamycin resistance via

saturation prime editing

After an extraordinary 18-month response to everolimus, a rapamycin analog, a patient with

metastatic anaplastic thyroid carcinoma relapsed with a resistant tumor harboring the *MTOR*-

F2108L mutation[34]. The same mutation was also found in spontaneous rapamycin-resistant yeast

mutants[35], and a clonal MCF-7 breast cancer cell line after three months of exposure to

rapamycin[36]. Of importance, this mutation does not hyperactivate mTOR[32,34,36] and confers

resistance to first-generation allosteric inhibitors of mTOR, but not to second- and third-generation

inhibitors[34,36]. F2108 lies within the FKBP12-rapamycin binding (FRB) domain of mTOR and is

an important residue involved in rapamycin interactions[37]. We performed saturation mutagenesis

of F2108 to install all possible amino acids substitutions at the endogenous *MTOR* locus to identify

and characterize alternative mutations causing rapamycin-resistance at this position. We devised

a saturation prime editing (PE) screen[31] using K562 cells stably expressing the mScarlet-I mTOR

signaling indicator (mSc-TOSI)[32,38] as a phenotypic readout to interrogate the functional impact of

these mutations (**Fig. 1a,b** and **Supplementary Fig. 1**). Under active mTORC1 signaling, the

fluorescent mSc-TOSI reporter is rapidly phosphorylated by S6K, ubiquitinated, and degraded by

the proteasome while mTOR inhibition stabilizes the reporter (**Fig. 1b**)[32,38].

We transfected our reporter cell line with a library of engineered pegRNA (epegRNA)[39] encoding

F2108 variants and performed rapamycin selection starting 3 days post-transfection until all non-

resistant cells were eliminated. Selection was performed at 0.5 µM rapamycin, the lowest

concentration of this drug that strongly abrogates K562 cell proliferation under our growth

conditions. We performed high-throughput sequencing prior and after FACS-based cell sorting

and analyzed the fold enrichment of all possible *MTOR*-F2108 permutations (**Fig. 1c,d** and

Supplementary Fig. 1). The most enriched mutation was F2108L[34,36] (4.8 fold enrichment), but

five additional mutations displaying 1.6 to 2.3 fold enrichments were selected for further
characterization (**Fig. 1d**). Of importance, cells modified at *MTOR* remained sensitive to the 2nd
generation ATP-competitive inhibitor AZD8055[40], which could be used as a safety switch in case
of overt toxicity (**Fig. 1c** and **Supplementary Fig. 1** and **2**).
129
To validate the top hits, we designed epegRNAs to test these variants individually. K562 cells
harboring these *MTOR*-F2108 substitutions grew robustly in the presence of 0.5 µM rapamycin
and we observed a marked increase in the percentage of alleles harboring the PE-specified edits
after rapamycin selection (**Supplementary Fig. 1**). As expected, no resistance and no enrichment
were observed with the silent F2108F mutation (**Supplementary Fig. 1**). These observations
confirmed that the five new variants identified by saturation prime editing conferred dominant
cellular resistance to rapamycin. Furthermore, mTORC1 signaling remained fully functional with
the F2108L mutation but was completely blocked with the silent F2108F mutation
(**Supplementary Fig. 1 and 2**). However, intermediate levels of signaling in the presence of
rapamycin were observed with F2108I, F2108M, and F2108G while weak activity observed with
F2108V and F2108K (**Supplementary Fig. 1 and 2**). These observations suggest that robust K562
cell growth can be maintained with partial mTORC1 signaling. These *MTOR*-F2108 mutations
could potentially be used to pharmacologically fine-tune mTORC1 signaling while maintaining
robust cellular growth. Overall, mTORC1 signaling remains completely functional in the presence
of rapamycin with F2108L, and this mutation was selected to develop a marker-free selection
strategy to engineer rapamycin-resistant CAR-T cells.
146
Concomitant transgene integration and induction of rapamycin resistance by nuclease-
driven *MTOR* editing in primary T cells.
While creating F2108L by itself via PE could be used to potentiate CAR-T cells, we aimed to
determine if we could couple rapamycin resistance to transgene integration in a single step by
editing the *MTOR* locus via homology-directed repair (HDR). As we previously showed for the
essential *ATP1A1* gene[32], this strategy can work by nesting a transgene within an intron without
disrupting the recipient gene[32,33]. We first designed two sgRNAs targeting *MTOR* intron 45 in
proximity to F2108 using CRISPOR[41] (see G4 and G5 in **Fig. 1a**). These two chemically modified
sgRNAs generated 86-90% indels in primary T cells when delivered as RNPs with both SpCas9

and the high-fidelity variant HiFiCas9[42] (**Fig. 2a**). To assess their genome-wide specificity, we

performed GUIDE-Seq[43,44] in primary CD3[+] T cells from two healthy donors using wild-type

SpCas9. While 9 potential off-target sites were identified with *MTOR*-G5, only one was identified

with *MTOR*-G4 in CD3[+] T cells (**Fig. 2b** and **Supplementary Fig. 3**). Similar results were

obtained in K562 cells via plasmid overexpression (**Supplementary Fig. 3**). Since *MTOR*-G4 is

highly active with the HiFiCas9 variant, we prioritized this sgRNA for our studies.

Next, using Cas9 RNPs and PCR-generated linear dsDNA donors[45], we confirmed that rapamycin

resistance could be achieved in primary CD3[+] T cells and observed a marked increase to up to

62% of alleles harboring the *MTOR*-F2108L mutation after selection (**Supplementary Fig. 4**).

This increase in HDR in the population upon selection was concomitant with a decrease of indels

detected at the cut site, suggesting efficient elimination of cells not having undergone HDR

(**Supplementary Fig. 4**). No enrichment was observed when using a donor that harbored the

wildtype *MTOR*-F2108 codon, confirming that robust growth in the presence of rapamycin

depends on the *MTOR*-F2108L mutation (**Supplementary Fig. 4**). In these experiments, a dose of

25 nM rapamycin was chosen based on growth curves indicating that maximal inhibition occurred

in the 16-32 nM range under our conditions (data not shown). A similar dose was used in previous

studies[27] and it falls between the immunosuppressive window of 3-22 nM and the maximal steady-

state plasma drug concentrations observed in clinical settings[46,47]. Thus, the installation of the well-

characterized *MTOR*-F2108L mutation[34,36] via HDR confers dominant cellular resistance to

rapamycin in primary T cells.

We then devised the strategy to simultaneously target a therapeutic transgene and introduce the

F2108L mutation to the *MTOR* locus (**Fig. 3a**). After targeted integration, *MTOR*-F2108L is

expressed on the forward DNA strand while the moderately active human *PGK1* promoter drives

the expression of a therapeutic transgene of interest on the reverse DNA strand (**Fig. 3a**). As a first

test, we targeted an expression cassette encoding the fluorescent protein mScarlet-I to the *MTOR*

locus in CD3[+] T cells using linear dsDNA donors[45]. Rapamycin selection reproducibly doubled

targeted integration frequency and we observed up to 31% of cells expressing the nested reporter

transgene (**Supplementary Fig. 5**). Similarly, we achieved 11% knock-in efficacy of a CD19-

CAR-2A-EGFP gene cassette (4.6 kb) following treatment for 8 days with rapamycin

(**Supplementary Fig. 6**). While encouraging, the use of large linear dsDNA donor caused toxicity,

limited proliferation, and was generally prohibitive for highly efficient gene targeting[45]. In

addition, these experiments were performed with CD3[+] T cells retained in leukoreduction system

(LRS) chambers that are typically discarded by blood banks. We found that these cells have higher

death rates and limited growth potential under our genome editing conditions in contrast to cells

isolated from whole blood donations. Nevertheless, taken together, these data illustrate that

rapamycin resistant primary T cells can be engineered to express transgenes in a single homology-

directed step using CRISPR-Cas9.

One-step generation of rapamycin resistant CAR-T cells

To improve targeting efficacy, we delivered the CD19-CAR-2A-EGFP donor using an adeno-

associated virus 6 (AAV6) vector[10,11,20] (**Fig. 3a**). Following Cas9 RNP nucleofection, CD3[+] T

cells isolated from whole blood (Donor ID: WB 1) were transduced with the AAV6 vector in a

small volume for 8 hours and then transferred to the culture medium for three days before

rapamycin treatment. Cells were reactivated four days post-transfection and expanded for an

additional 7 days before quantification of transgene-expressing cells (**Fig. 3b**). We observed a

substantial increase in editing efficacy both pre and post 8 days of selection as determined by out-

out PCR and Sanger sequencing in the bulk populations (**Fig. 3c**). Yields of CAR-2A-EGFP[+] cells

reached 82% and 85% at MOIs of 5×10^3 and 1×10^4, respectively as determined by FACS-based

analysis (**Fig. 3d**). These results were reproduced with CD3[+] T cells isolated from a second whole

blood donor (Donor ID: WB2) and two LRS donors (Donor ID: LRS1, LRS2) (**Fig. 3e**). In total,

cells from four distinct healthy donors generated similar outcomes, albeit with lower efficiency

when using cells isolated from LRS chambers (**Fig. 3e**).

To test for functionality, we performed luciferase-based cytotoxicity assays with selected CAR-T

effector cells and CD19[+] NALM6 target cells stably expressing firefly luciferase and RFP. We

also generated a rapamycin-resistant NALM6 cell line (NALM6-RapaR) by installing the *MTOR*-

F2108L mutation to assess the combinational impact of rapamycin. We observed high levels of

cytotoxicity with and without rapamycin at the highest effector to target ratios (**Fig. 3f**). At lower

effector to target ratios, higher levels of cytotoxicity were observed in the presence of rapamycin,

confirming that these CAR-T cells can be used in combination with the mTOR inhibitor to target

cancer cells (**Fig. 3f**). This combinatorial effect was not observed with NALM6-RapaR cells (**Fig.**
3f). These data indicate that immunosuppressive doses of rapamycin can be used in combination
with these CAR-T cells to increase antitumor activity *in vitro*.

Rapamycin has previously been shown to accelerate the memory T-cell differentiation program
and may impact the phenotype of treated cells[48,49]. To begin to probe the impact of rapamycin
treatment *ex vivo*, we analyzed differentiation and exhaustion markers and performed multi-color
flow cytometry panels after 8 and 15 days of treatment using CD3$^+$ T cells isolated from whole
blood of two additional healthy donors (ID: WB3, WB4) (**Supplementary Fig. 7**). In these
experiments, editing rates remained high (~80% CAR-2A-EGFP$^+$ cells) and we observed a
minimal impact of rapamycin treatment on the expression of differentiation and exhaustion
markers, as determined by CD4, CD8, CD45RO, CCR7, and PD1 staining, on both rapamycin-
sensitive and rapamycin-resistant T cells (**Supplementary Fig. 7**). Rapamycin-resistant CAR-2A-
EGFP$^+$ populations displayed a mix of central memory (CCR7$^+$/CD45RO$^+$) and effector memory
(CCR7$^-$/CD45RO$^+$) T cells after 8 days of treatment with rapamycin, while most cells displayed
an effector memory (CCR7$^-$/CD45RO$^+$) phenotype after 15 days of treatment (**Supplementary**
Fig. 7). Overall, rapamycin treatment (25 nM) had little impact on T cells' differentiation in our
ex vivo expansion protocol, and our 14 days expansion timeline (8 days of rapamycin treatment)
yields a mix of central memory and effector memory CAR-T cells.

Combinatorial CAR-T cell therapy with rapamycin slows leukemia progression *in vivo*
We then established a xenograft leukemia model using NOD/SCID/IL-2Rγ-null (NSG) male and
female mice injected with NALM6 cells stably expressing RFP and firefly luciferase. NALM6
cells were injected intravenously, and rapamycin treatment (4 mg/kg, daily) started three days
later. Rapamycin-resistant CD19-CAR-2A-EGFP$^+$-CAR T cells (Donor ID: WB2, see **Fig. 3e**),
which underwent *ex vivo* selection with 25 nM rapamycin for eight days (see **Fig. 3a,b**) , were
administered intravenously four days after tumor inoculation (**Fig. 4a)**. In total, these CD19-CAR
T cells where expanded *ex vivo* for 14 days and stimulated twice before being injected in mice (see
Fig. 3b). As a negative control, we used CD3$^+$ T cells untransduced (UT) with AAV6 donors but
having undergone nucleofection with the Cas9-*MTOR*-G4 RNP. Tumor engraftment and
progression were assessed via bioluminescence imaging twice per week for seven weeks. While

rapid tumor progression was observed with UT T cells without treatment, rapamycin slowed tumor
progression and conferred an additional week of survival (**Fig. 4b,c**). Of importance, rapamycin-
resistant CD19-CAR-T cells selected *ex vivo* could efficiently target NALM6 cells *in vivo* in the
presence or absence of rapamycin (**Fig. 4b,c**). Combinatorial rapamycin administration with
rapamycin-resistant CAR-T cells further slowed tumor progression and allowed the mice to
survive for up to seven weeks post-tumor inoculation (**Fig. 4b,c**). While we observed antitumor
control during the first weeks, tumors eventually rebounded and killed all mice (**Fig. 4b,c**). Our
results corroborate previous findings demonstrating the limited persistence of second-generation
CARs harboring a CD28 costimulation domain compared to 4-1BB[50–52]. Indeed, the second-
generation CD19-CAR architecture used during this study[53] contains a CD28 transmembrane
domain, which can form an heterodimer with CD28[54], and a CD28 costimulation domain that has
been linked to lower persistence compared to 4-1BB[6,52,55]. Nevertheless, our results confirmed that
our targeting strategy can create functional rapamycin-resistant CAR-T cells that can be used in
combination with rapamycin to slow tumor progression *in vivo*, offering an opportunity to target
cancer cells' metabolism to facilitate tumor eradication.

Pharmacological control of CAR-T cell activity using rapamycin

Using intracellular[56] or extracellular[46] rapamycin-regulated dimerization domains to
pharmacologically control chimeric antigen receptor assembly is a promising approach for
developing safer CAR-T cell therapies[57]. However, these systems rely on the non-
immunosuppressive rapamycin analog AP21967 which was primarily developed as a tool for
chemical biology and do not have suitable pharmacokinetic properties for clinical use[56], or on non-
immunosuppressive doses of rapamycin, which limit their clinical potential. Thus, we tested
whether we could target a dimerizing agent-regulated immunoreceptor complex (DARIC)[46], which
can be toggled between on and off states using rapamycin, to the *MTOR* locus.

Building on previous studies[10,46], we sought to test whether multiplexed targeting of separate
DARIC components could be performed at the *MTOR* and *TRAC* loci simultaneously (**Fig. 5a**).
Taking advantage of the dynamic regulation of CAR expression at the *TRAC* locus[10], which averts
tonic CAR signaling, we targeted a CD22 recognition domain (DARIC 2) and a 41BB-CD3z
signaling domain (DARIC 3) linked by a P2A self-cleaving peptide to the *TRAC* locus (**Fig. 5a**).

In parallel, we targeted an additional CD19 recognition domain (DARIC 1) to the *MTOR* locus to

confer dominant cellular resistance to rapamycin while providing an additional antigen specificity

to potentially overcome antigen escape[6,7,55,58,59]. Multiplexed targeted integration of both DARIC

cassettes was observed at *MTOR* and *TRAC*, as determined by out-out PCR (**Fig. 5b**). An average

of 58% of alleles harboring the *MTOR*-F2108L was observed, confirming that multiplexed genome

editing did not decrease targeted transgene integration at *MTOR* (**Fig. 5b**). We performed FACS-

based cytotoxicity assays using unselected CAR-T cells in the presence or absence of 25 nM

rapamycin with DARIC-T cells that were resistant (RapaR-CD19-CD22-DARIC) or not (CD22-

DARIC) to rapamycin. Strikingly, cytotoxicity was only observed with rapamycin-resistant CD19-

CD22-DARIC-T cells in the presence of rapamycin, confirming DARIC immunoreceptor

dimerization only in the presence of the drug (**Fig. 5c**). As observed above, rapamycin inhibited

NALM6 cell growth to some extent, but not NALM6-RapaR which harbors the *MTOR*-F2108L

mutation (**Fig. 5c**). Rapamycin-sensitive CD22-DARIC-T cells, where the *MTOR*-F2108L-

DARIC-1 AAV6 donor was omitted during transduction, could not target cancer cells in the

presence of an immunosuppressive dose of 25 nM rapamycin (**Fig. 5c**). We note that the CD22

scFv construction (DARIC 2) we engineered has never been tested for functionality by itself, thus

the absence of cytotoxicity of CD22-DARIC cells in presence of rapamycin should be interpreted

with caution. Altogether, this multiplexed strategy offers the opportunity to use

immunosuppressive doses of rapamycin in combination with DARIC-T cells to decrease tumor

cell growth and to pharmacologically fine-tune CAR-T cells' activity.

DISCUSSION

Engineering primary T cells with programmable nucleases for cell therapy applications remains

challenging. In this work, we describe a marker-free selection approach to simultaneously target a

therapeutic transgene of interest and a point mutation to the *MTOR* locus to generate dominant

cellular resistance to rapamycin. We show that mTORC1 signaling remained functional and cells

grew robustly in the presence of rapamycin after targeted transgene integration. Our results further

demonstrate that introns provide an additional non-coding location for targeted transgene

integration via intron nesting[32,33]. Taking advantage of this naturally occurring genomic

architecture allows the co-modification of an endogenous gene during the integration of a

therapeutic transgene of interest in a single gene editing event. However, this strategy is limited to

moderately active promoters, such as the human *PGK1* promoter used in this study, since transcriptional interference between the overlapping genes[33] would be detrimental for mTORC1 signaling, and consequently, CAR-T cell activation and expansion[21,30]. This could be prohibitive for applications where high levels of transgene expression are needed to reach a therapeutic threshold.

The metabolic competition between cancer and T cells has previously been shown to drive cancer progression[60], and metabolic interventions could be used in combination to improve clinical outcomes[61]. Hence, targeting cancer metabolism via mTORC1 inhibition could potentially slow tumor progression and facilitate complete tumor eradication. Taken together, our findings suggest that rapamycin could be administered to enrich CRISPR-engineered cells *in vivo* while providing a fitness advantage against tumor cells. Of note, since double editing events at two different loci are not statistically independent, rapamycin co-selection could also be performed to enrich a second modification of interest[32,62].

Our cytotoxicity assays corroborate previous findings regarding the combinatorial impact of rapamycin with *piggyBac*-modified rapamycin-resistant CAR-T cells[27]. Nonetheless, our CRISPR-based strategy offers advantages over the use of a two *piggyBac* transposon vectors system to target a CAR and a large rapamycin-resistant mTOR transgene cassette ($\approx$7.65kb)[27] to random genomic locations[27–29].

While we focused on the standard CD19[+] NALM6 acute lymphoblastic leukemia model during this study, combinatorial CAR-T cell therapy with rapamycin could potentially improve antitumor activity against solid tumors displaying hyperactive mTOR signaling[34,63,64]. Moreover, using a dimerizing agent-regulated immunoreceptor complex (DARIC) switchable between on and off states using rapamycin is a promising strategy to develop safer CAR-T cells therapies[46,57]. One limitation of this system is the use of non-immunosuppressive rapamycin dosing[46] to prevent the inhibition of CAR-T cells' activation, which could be overcome using rapamycin-resistant DARIC-T cells. This system could be used to fine-tune CAR-T cells activity with rapamycin pulsing to prevent tonic CAR signaling, and consequently, exhaustion[46,65]. In line with this, transient rest induced by a drug-regulatable CAR or the tyrosine kinase inhibitor dasatinib has

been shown to restore functionality in exhausted CAR-T cells through epigenetic remodeling[66–68].

Most importantly, rapamycin-regulatable CAR architectures could be used as safety switches to

dampen toxicity and severe adverse events associated with cytokine release syndrome[46,66–70].

During this study, we also identified new rapamycin resistance mutations via saturation prime

editing that displayed intermediate levels of mTORC1 signaling in the presence of rapamycin.

These additional mutations offer interesting opportunities to investigate the potential of

pharmacologically fine-tuning mTORC1 activity for cell therapy applications. mTOR plays key

in T cell fate decisions and differentiation[30,48]. Reducing mTORC1 signaling has been shown to

trigger the formation of stem cell-like memory T cells and also enhance CAR-T cell antitumor

activity[49,71]. In addition, Lamarthée and colleagues have shown that transient mTOR inhibition

with rapamycin rescues 4-1BB CAR-Tregs from tonic signal-induced dysfunction[72]. Considering

that mTORC1 signaling acts as a metabolic rheostat[24], fine-tuning this pathway could potentially

have a positive impact on CAR-T cells persistence, and consequently, CAR-T cells antitumor

activity. While this hypothesis warrants further investigation, our results demonstrate that

engineering the *MTOR* locus with CRISPR allows the marker-free selection of cells with different

levels of mTORC1 signaling in the presence of rapamycin.

Altogether, the strategy presented here should facilitate the *ex vivo* and *in vivo* enrichment of

CRISPR-engineered cells expressing a therapeutic transgene of interest, providing a versatile

platform for cell therapy applications.

METHODS

K562 and NALM6 cell culture and transfection

K562 cells were obtained from the ATCC (CCL-243) and NALM6 cells stably transduced to

express a FLUC-T2A-RFP-IRES-Puro reporter gene cassette (Biosettia) were gently provided by

Scott McComb (National Research Council of Canada). K562 and NALM6 cells were maintained

at 37 °C under 5% CO_2 in RPMI medium supplemented with 10% FBS, penicillin–streptomycin,

and GlutaMAX. Cells were routinely tested for the absence of mycoplasma contamination.

Rapamycin (Cayman Chemicals, Cat 53123-88-9) was dissolved at 10 mg/ml in DMSO, working

dilutions were prepared in water and stored at -20°C. AZD8055 (STEMCELL Technologies) was

dissolved at 10 mM in DMSO, working dilutions were prepared in water and stored at -20°C.

DMSO alone was diluted in water and used as vehicle control. Ouabain octahydrate (Sigma) was

dissolved at 5 mg/ml in hot water, working dilutions were prepared in water and stored at -20°C.

K562 and NALM6 cells (2×10^5 cells/transfection) were transfected using a Lonza 4D

nucleofector™ with a SF nucleofection kit (Lonza) following manufacturer's recommendations.

For prime editing transfections, K562 cells were transfected with 0.75 µg of PE vector, 0.25 µg of

epegRNA vector, and 0.1 µg of nick sgRNA vector. NALM6 cells were transfected with 50 pmols

of SpCas9 RNP (IDT) and 0.75 µg of dsDNA donor harboring *MTOR*-F2108L and a PAM

mutation to generate the NALM6-RapaR cell line. Cells were treated with the indicated

concentration of rapamycin 3 days post-nucleofection until all non-resistant cells were eliminated.

Genome editing vectors and recombinant AAV production

Guide RNAs were designed with CRISPOR[41] and their sequences are provided in the

Supplementary material section. When required, DNA sequences for the guides were modified

at position 1 to encode a G, owing to the transcription requirement of the human U6 promoter.

Guide RNAs were cloned into pX330-U6-Chimeric_BB-CBh-hSpCas9 (a gift from Feng Zhang;

Addgene 42230) and their sequences are provided in the **Supplementary material** section.

Plasmid donors were cloned into pUC19 with short homology arms (700-800 bp) and all sequences

were confirmed by Sanger sequencing. Prime editing (PE) experiments were performed with

pCMV-PEmax[73] (a gift from David Liu; Addgene 174820), and pU6-tevopreq1-GG-acceptor[39] (a

gift from David Liu; Addgene 174038). To repurpose the mTORC1 signaling reporter mVenus-

TOSI previously developed for mouse[38], the N-terminal residues (1-82) of the human *PDCD4*

gene were codon optimized using the GenSmart™ codon optimization tool, synthesized as a

gBlocks™, and cloned into ATP1A1_T804N_hPGK1_mScarlet-I_Donor (Addgene 173207)

upstream of the mScarlet-I-NLS cassette using AflII and NcoI. The CD19-CAR-28z-2A-EGFP

gene cassette from pSLCAR-CD19-28z (a gift from Scott McComb: Addgene 135991)[53] was

cloned into the *MTOR* donor vector by Gibson assembly without the 3xFLAG. The CD19-DARIC

expression vector was designed in-house based on previously described constructions[46] and US

patent 2012/0266971 A1. The CD22-DARIC construction was designed using a previously

described CD22 scFv fragment[59]. DARIC sequences were codon-optimized using GenSmart™

codon optimization tool, synthesized by GenScript, and cloned into the AAV6 vector using Gibson

assembly. Donor sequences are provided in the **Supplementary material** section. The AAV6

vectors were produced by the vector core facility at the Canadian neurophotonics platform

(molecular tools). ITR integrity was assessed following BssHII digestion of the AAV plasmid

before production. The virus was resuspended in PBS 320 mM NaCl + 5% D-sorbitol + 0.001%

pluronic acid (F-68), aliquoted, and stored at −80°C. The vector yields were $1x10^{12}$, $8.5x10^{12}$, and

$6.9x10^{12}$ GC/ml for the *MTOR*-F2108L-CD19-CAR-2A-EGFP, *MTOR*-F2108L_CD19_scFv, and

TRAC_T2A_CD22-DARIC donors, respectively.

Primary human T cells isolation and culture

Primary CD3$^+$ human T cells were isolated from anonymized healthy human donors either from

fresh whole blood (WB) or leukocyte reduction system (LRS) chambers (Héma-Québec), or from

human peripheral blood leukopaks (STEMCELL). EasySep™ Human T cell isolation kit

(STEMCELL) or CD3 microbeads (Miltenyi) were used for positive magnetic selection of CD3$^+$

T cells from peripheral blood mononuclear cells (PBMCs). Primary CD3$^+$ T cells were cultured

with Immunocult-XF T cell expansion medium supplemented with 1% penicillin-streptomycin and

300 U/ml IL-2. Cells were thawed and activated for 3 days with Immunocult™ human

CD3/CD28/CD2 T cell activator (STEMCELL) before nucleofection.

Primary human T cells transfection

dsDNA donors were produced as described previously[45]. Briefly, donor sequences were cloned

into a cloning vector and PCR amplification (25 cycles) was performed using Kapa-HiFi

polymerase (Roche). DNA purification was performed by solid-phase reversible immobilization

using AMPure XP (Beckman Coulter) magnetic beads. For ribonucleoprotein complex (RNP)

formulation, 100 pmols of sgRNAs (Alt-R, IDT) were mixed with 0.8 µl of 100 mg/ml

polyglutamic acid (Sigma) and 50 pmols of SpCas9 nuclease (IDT) as previously described[74]. The

RNP mix was incubated at 37°C for 10 minutes and the dsDNA donors were added to the mix and

incubated at room temperature for 5 minutes before transfection. Primary CD3$^+$ T cells ($1x10^6$

cells/transfection) were transfected using a Lonza 4D nucleofector™ (Pulse code EH115) with a

P3 nucleofection kit (Lonza) following manufacturer's recommendations with the indicated DNA

concentrations. CD3$^+$ T cells were resuspended in 80 µl media without IL-2 directly after

nucleofection and incubated at 37°C for 15 minutes before transfer into culture media plates at

1×10^6 cells/ml. For AAV6 transduction, CD3$^+$ T cells were resuspended in 80 µl media directly

after nucleofection and AAV6 vectors were added. Cells were incubated at 37°C for 8 hours at

high cell density before transfer into culture media plates. For rapamycin selection, CD3$^+$ T cells

were treated with or without 25 nM rapamycin on Day 3 post-nucleofection, restimulated with

Immunocult™ human CD3/CD28/CD2 T cell activator (STEMCELL) 4 days post-transfection,

and expanded for an additional 7 days.

Sanger sequencing analysis and out-out PCRs

Genomic DNA was extracted with QuickExtract DNA extraction solution (EpiCentre) following

manufacturer's recommendations. Primers used in this study and the PCR product sizes are

provided in the **Supplementary material** section. PCR amplifications were performed with 30

cycles of amplification with Phusion polymerase. Sanger sequencing was performed on PCR

amplicons to quantify the percentage of edited alleles using TIDE[75], TIDER[76], and BEAT[77]. To

quantify the percentage of alleles harboring the *MTOR*-F2108L mutation after targeted transgene

integration, we performed PCR amplifications to amplify both WT and HDR alleles with a forward

primer binding outside the region of homology of our donors, and a reverse primer binding inside

the homology arm upstream of the SpCas9 target sites (out-in PCR). Genomic DNA from WT

cells and cells transfected or transduced only with the donor were used as negative controls in all

experiments. Kapa-HiFi polymerase was used for out-out PCRs. To detect targeted integration,

out-out PCRs were performed with primers that bind outside of the homology regions of the

plasmid/dsDNA donors. Wild type K562 or CD3$^+$ T cells genomic DNA was used as a control for

all PCRs.

Flow cytometry

The percentage of fluorescent cells was quantified using a BD LSRII flow cytometer using

FACSdiva v6.1.2 software, and 1×10^5 cells were analyzed for each condition. For mTORC1

signaling assays, cells were washed with PBS and cultured at least 3 days without rapamycin. Cells

were treated with the indicated concentration of rapamycin or AZD8055 for 24 hours before

analysis. FACS sorting was performed using a BD FACS Aria Fusion flow cytometer for the

saturation prime editing experiment, and 5×10^5 cells were sorted for each condition.

Multicolor flow cytometry panels were performed using a Cytek Aurora flow cytometer. T cells

were stained with fixable viability dye eF506 (Invitrogen Cat 65-0866-18) for 15 minutes at 4°C,

then washed and resuspended in Human TruStain FcX blocking solution (BioLegend Cat

2711505) and Brilliant Stain Buffer Plus (BD Biosciences Cat 566385) for 10 minutes at room

temperature. After blocking, cells were stained with the following antibody cocktail diluted in flow

cytometry staining buffer: mouse anti-human CD3 BV750 (BioLegend Cat 344846), mouse anti-

human CD4 AF700 (BioLegend Cat 300526), mouse anti-human CD8 PerCP (BioLegend Cat

301030), mouse anti-human CD279 BV421 (BioLegend Cat 329920), mouse anti-human CCR7

APC-Fire 750 (BioLegend Cat 353246), and mouse anti-human CD45RO PE-Cy7 (Invitrogen Cat

25-0457-42). Flow cytometric data were analyzed using SpectroFlo (Cytek) and FlowJo V10 (Tree

Star).

Saturation prime editing and high-throughput DNA sequencing

The *MTOR*-F2108 saturation epegRNA (tevopreQ1)[39] vector was designed to install NNK codons

and a silent R2109R PAM mutation at *MTOR* exon 45, synthesized as a gBlock™ (IDT), and

cloned into pU6-tevopreq1-GG-acceptor[39] (a gift from David Liu; Addgene 174038). K562 cells

stably expressing the mSc-TOSI reporter were transfected with PE3max-epegRNA (tevopreQ1)

vectors and treated with 0.5 µM rapamycin starting 3 days post-transfection until all non-resistant

cells were eliminated. Following selection, cells were treated for 24 hours with 0.5 µM rapamycin

and FACS-sorted for low mSc-TOSI fluorescence intensity to enrich cells with functional

mTORC1 signaling in the presence of rapamycin. Genomic DNA was harvested after selection

and after FACS sorting for low and high mSc-TOSI fluorescence intensity. For amplicons

sequencing, primers containing Illumina forward and reverse adapters were used for a first round

of PCR using the Kapa-HiFi polymerase. PCR products were purified using AMPure XP magnetic

beads and their quality was evaluated by electrophoresis. A second round of PCR and bead

purification was performed for indexing, and amplicons were sequenced on an Illumina MiSeq

instrument. Alignment of amplicon sequences to the reference *MTOR* sequence was performed

using CRISPResso2[78], and all possible NNK codons were detected.

GUIDE-Seq

GUIDE-Seq was performed as previously described[43,44], except for improvements of the Next

Generation Sequencing (NGS) library preparation procedure reported below. GUIDE-Seq oligos

harboring 5' phosphorylation modification and two phosphorothioate linkages between the last

two nucleotides at the 3' end were synthesized by IDT. Oligos were mixed in 50 mM NaCl, 10

mM Tris-HCl (pH 8.0), 1 mM EDTA, and annealed by heating the solution to 95°C for 10 minutes,

followed by gradual cooling on a thermocycler. For K562 cells, $2x10^5$ cells were electroporated

with 750 ng of SpCas9-sgRNA vectors and 100 pmol of annealed GUIDE-Seq oligos. For primary

$CD3^+$ T cells, $1x10^6$ cells were electroporated with 50 pmol SpCas9 RNP and 100 pmol annealed

GUIDE-Seq oligos. Genomic DNA was harvested using QIAmp UCP DNA Micro Kit (QIAGEN).

GUIDE-Seq oligo tag integration was confirmed using TIDE[75] and DECODR[79] webtools from

Sanger sequencing. The library preparation method was improved starting from the original

GUIDE-Seq procedure[43,44]. After mechanical fragmentation, end-repair and A-Tailing were

directly performed using the NEBNext UltraII DNA kit (New England Biolabs). A new custom

universal Y-adapter based on the Illumina TruSeq sequences was designed to include a Unique

Molecular Index (UMI) in the sequencing read instead of the index read. This modification enables

mixing of GUIDE-Seq libraries with other Illumina library types (amplicon, shotgun, RNASeq,

etc.) in the sequencing run by using standard index read lengths. This new adapter was ligated as

described in the instructions of the NEBNext UltraII procedure using a final concentration of 133

nM. Ligated DNA was purified using 0.9X volume of AMPure XP beads (Beckman-Coulter), then

split in 2 for PCR amplifications with either Minus or Plus primers using Q5 DNA Polymerase

(New England Biolabs) with the following cycling conditions: 98°C 30 sec; 10 cycles of 98°C 10

sec, 55°C 30 sec, 72°C 30 sec; 15 cycles of 98°C 10 sec, 65°C 30 sec, 72°C 30 sec; 72°C 2 min;

4°C hold. New primers with Illumina sequence tails were specifically designed for this improved

method. The use of Illumina sequences at the Plus/Minus amplification stage eliminates the use of

custom primers for sequencing. PCR products were purified with 0.9X AMPure XP beads and

quantified using Qubit dsDNA HS Assay (Invitrogen). Then, 10 ng DNA was used for a second

PCR amplification with Illumina TruSeq dual-indexing primers (98°C 30 sec; 8 cycles of 98°C 10

sec, 55°C 30 sec, 72°C 30 sec; 72°C 2 min; 4°C hold). The barcoding at the PCR stage enables a

much simpler and higher multiplexing capacity. PCR products were purified using 0.85X AMPure

XP beads, quantified using Qubit dsDNA HS Assay and ran on a Bioanalyzer High Sensitivity

DNA chip (Agilent). Samples were pooled in equimolar amounts and sequenced on 1.4% (35M

read block) of a NovaSeq S4 PE150 lane (Illumina) at the Centre d'Expertise et de Services
Génome Québec (Montréal, QC). The remaining portion of the sequencing lane contained a
mixture of library types from multiple organisms (*M. plutonius*, *P. larvae*, *F. graminearum*, *P.*
neglectus, *Colletotrichum* spp., *Sporothrix* spp.).
531

Analysis of GUIDE-Seq sequencing
A new pipeline, named guideseq_ibis, was designed to analyze the data from the improved
GUIDE-Seq procedure (https://github.com/enormandeau/guideseq_ibis). Using this pipeline, the
raw reads were trimmed for quality using trimmomatic (v0.36, min_length 100, crop_length 200).
Then, only reads containing the expected alien sequence (maximum hamming distance of 1) and
only one copy of the ODN sequence (maximum hamming distance of 6) were retained. For these
reads, the Unique Read Identifiers (UMIs) and first eight nucleotides were used to rename the
sequence. These tagged reads were then mapped onto the latest GRCh38 human genome assembly
with bwa (v0.7.17-r1188, -T 10) and then samtools (v1.12, -S -q 1 -F 4 -F 256 -F 2048). The
alignment sam files were then sorted by chromosome name and position. The UMI and eight first
nucleotides of the reads and the starting positions of their alignments were used to detect duplicated
reads. Namely, only one copy of reads with identical UMIs, eight first nucleotides, and alignment
start position were kept. Then, Double Stranded Breaks (DSBs) were identified. For this, alignment
sites were scanned for peaks of coverage in decreasing order of depth of coverage and pairs of
ODN+ and ODN- were reported until the pairs did not meet the minimum coverage threshold
(min_length 100, min_coverage 50, window_size 10, position_error 5, bin_size 10000). To
account for small errors in sequencing and mapping, read counts are collected within
`window_size` nucleotides around each peak and pairs of ODN sequences were kept if they fell
withing `position_error` nucleotides of their relative expected positions. Identified sequences on
target and off-target were reported for each sample with their chromosome and position
localisation, the identifier and name of any gene they overlapped (using a simplifited annotation
table from GRCh38.p14), and metrics about the counts of the ODN+ and ODN- reads. Finally,
annotation of the found targets and off-targets was performed to add information about the guide's
name, the distance between the target sequence and the expected guide sequence, as well as a
representation of the modified nucleotides along the guide sequence.
557

Cytotoxicity assays

Luciferase-based cytotoxic assays were performed as previously described[10,80]. After rapamycin selection, CD19-CAR-T cells were washed with PBS, and co-cultured in 96-well plates in supplemented RPMI medium with NALM6 cells stably expressing firefly luciferase and RFP at the indicated effector to target ratio (E:T) for 18 hours. Target cells incubated with WT CD3$^+$ T cells were used to determine maximal luciferase expression (maximal relative light units; RLU$_{max}$). After coculture, an equal volume of luciferase substrate (Bright-Glo, Promega) was added to each well. Emitted light was detected with a Tecan luminescence plate reader and cell lysis was determined as $(1 - (RLU_{sample})/(RLU_{max})) \times 100$. FACS-based cytotoxicity assays were performed as previously described[46], except with a prolonged coculture time. T cells were washed with PBS, and cocultured in 12-well plates in supplemented RPMI medium with 50:50 mixtures of rapamycin-resistant K562-EBFP (CD19-negative) cells and NALM6-RFP cells at the indicated effector to target ratio (E:T) for 72 hours. Cell viability was assessed using the fixable viability dye eFluor™ 660 (eBioscience). The ratio of viable NALM6-RFP and K562-EBFP cells following coculture in the presence or absence of 25 nM rapamycin was used to calculate the percentage of cytotoxicity. Target cells incubated with WT CD3$^+$ T cells were used to determine maximal NALM6-RFP growth.

Mouse systemic tumor model.

The *in vivo* CAR-T cells experiments were performed according to the *Canadian Guide for the Care and Use of Laboratory Animals*. The Université Laval Animal Care and Use Committee approved the procedure. 8-10 weeks-old NOD/SCID/IL-2Rγ-null (NSG) male and female mice were acquired from Jackson Laboratory. 0.5×10^6 NALM6-RFP-FLUC cells were administered via tail vein injection. Three days later, daily rapamycin treatment (4 mg/kg) or vehicle (filter-sterilized solution of 5% Tween80, 5% PEG-400, 0.7% DMSO) was administered by intraperitoneal (IP) injection. Four days after tumor inoculation, 2×10^6 CAR-T cells (≈86% CAR-2A-EGFP$^+$ cells) or untransduced (UT) T cells were administered via tail vein injection. CAR-T cells expanded for 14 days, including 8 days of treatment/selection with 25 nM rapamycin, were used for the *in vivo* experiment. Tumor burden was measured twice per week by bioluminescence imaging using the IVIS LuminaIII Imaging System (PerkinElmer). Mice were anesthetized using isoflurane, and 150 mg/kg D-Luciferin (Goldbio) was administered via IP injection. Data were

589 analyzed using the Living Image software (PerkinElmer). The maximum tumor burden and
590 survival endpoint were defined by hindlimb paralysis or other clinical signs of distress. There were
591 no instances in which this maximum tumor burden was exceeded.
592

593 **DATA AVAILABILITY**

594 Source data for the figures are provided as a Source Data file. All raw Sanger sequencing data
595 generated in this study are available on request from the corresponding author [YD].
596

597 **CODE AVAILABILITY**

598 The adapted GUIDE-Seq analysis pipeline developed and used in this study is freely available at
599 https://github.com/enormandeau/guideseq_ibis.
600

REFERENCES

1. Brentjens, R. J. *et al.* CD19-targeted t cells rapidly induce molecular remissions in adults
with chemotherapy refractory acute leukemia. *Sci. Transl. Med.* **5**, 177ra38 (2013).
2. Lee, D. W. *et al.* T cells expressing CD19 chimeric antigen receptors for acute
lymphoblastic leukaemia in children and young adults: A phase 1 dose-escalation trial.
Lancet **385**, 517–528 (2015).
3. Neelapu, S. S. *et al.* Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-
Cell lymphoma. *N. Engl. J. Med.* **377**, 2531–2544 (2017).
4. Park, J. H. *et al.* Long-term follow-up of CD19 CAR therapy in acute lymphoblastic
leukemia. *N. Engl. J. Med.* **378**, 449–459 (2018).
5. Maloney, D. G. Anti-CD19 CAR T cell therapy for lymphoma — off to the races! *Nat.*
Rev. Clin. Oncol. **16**, 279–280 (2019).
6. Labanieh, L., Majzner, R. G. & Mackall, C. L. Programming CAR-T cells to kill cancer.
Nat. Biomed. Eng. **2**, 377–391 (2018).
7. Larson, R. C. & Maus, M. V. Recent advances and discoveries in the mechanisms and
functions of CAR T cells. *Nat. Rev. Cancer* **21**, 145–161 (2021).
8. Bailey, S. R. & Maus, M. V. Gene editing for immune cell therapies. *Nat. Biotechnol.* **37**,
1425–1434 (2019).
9. Ellis, G. I., Sheppard, N. C. & Riley, J. L. Genetic engineering of T cells for

immunotherapy. *Nat. Rev. Genet.* **16**, 103–107 (2021).

10.   Eyquem, J. *et al.* Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances

tumour rejection. *Nature* **543**, 113–117 (2017).

11.   Mansilla-soto, J. *et al.* HLA-independent T cell receptors for targeting tumors with low

antigen density. *Nat. Med.* **28**, 345–352 (2022).

12.   Wei, J. *et al.* Targeting REGNASE-1 programs long-lived effector T cells for cancer

therapy. *Nature* **576**, 471–476 (2019).

13.   Giuffrida, L. *et al.* CRISPR/Cas9 mediated deletion of the adenosine A2A receptor

enhances CAR T cell efficacy. *Nat. Commun.* **12**, 3236 (2021).

14.   Carnevale, J. *et al.* RASA2 ablation in T cells boosts antigen sensitivity and long-term

function. *Nature* **609**, 174–182 (2022).

15.   Depil, S., Duchateau, P., Grupp, S. A., Mufti, G. & Poirot, L. 'Off-the-shelf' allogeneic

CAR T cells: development and challenges. *Nat. Rev. Drug Discov.* **19**, 185–199 (2020).

16.   Stadtmauer, E. A. *et al.* CRISPR-engineered T cells in patients with refractory cancer.

Science **7365**, eaba7365 (2020).

17.   Ottaviano, G. *et al.* Phase 1 clinical trial of CRISPR-engineered CAR19 universal T cells

for treatment of children with refractory B cell leukemia. *Sci. Transl. Med.* **3010**,

eabq3010 (2022).

18.   Shy, B. R. *et al.* High-yield genome engineering in primary cells using a hybrid ssDNA

repair template and small-molecule cocktails. *Nat. Biotechnol.* (2022)

doi:10.1038/s41587-022-01418-8.

19.   Paul, B. *et al.* Efficient enrichment of gene-modified primary T cells via CCR5-Targeted

integration of mutant dihydrofolate reductase. *Mol. Ther. Methods Clin. Dev.* **9**, 347–357

(2018).

20.   Wiebking, V. *et al.* Metabolic engineering generates a transgene-free safety for cell

therapy. *Nat. Biotechnol.* **38**, 1441–1450 (2020).

21.   Saxton, R. A. & Sabatini, D. M. mTOR signaling in growth, metabolism, and disease. *Cell*

168, 960–976 (2017).

22.   Battaglioni, S., Benjamin, D., Wälchli, M., Maier, T. & Hall, M. N. mTOR substrate

phosphorylation in growth control. *Cell* **185**, 1814–1836 (2022).

23.   Liu, G. Y. & Sabatini, D. M. mTOR at the nexus of nutrition, growth, ageing and disease.

651   *Nat. Rev. Mol. Cell Biol.* **21**, 183–203 (2020).

652 24. Valvezan, A. J. & Manning, B. D. Molecular logic of mTORC1 signalling as a metabolic

653   rheostat. *Nat. Metab.* **1**, 321–333 (2019).

654 25. Mossmann, D., Park, S. & Hall, M. N. mTOR signalling and cellular metabolism are

655   mutual determinants in cancer. *Nat. Rev. Cancer* **18**, 744–757 (2018).

656 26. Sun, Q. *et al.* Mammalian target of rapamycin up-regulation of pyruvate kinase isoenzyme

657   type M2 is critical for aerobic glycolysis and tumor growth. *Proc. Natl. Acad. Sci. U. S. A.*

658   **108**, 4129–4134 (2011).

659 27. Huye, L. E. *et al.* Combining mTor inhibitors with rapamycin-resistant T cells: A two-

660   pronged approach to tumor elimination. *Mol. Ther.* **19**, 2239–2248 (2011).

661 28. Micklethwaite, K. P. *et al.* Investigation of product-derived lymphoma following infusion

662   of piggyBac-modified CD19 chimeric antigen receptor T cells. *Blood* **138**, 1391–1405

663   (2021).

664 29. Bishop, D. C. *et al.* Development of CAR T-cell lymphoma in 2 of 10 patients effectively

665   treated with piggyBac-modified CD19 CAR T cells. *Blood* **138**, 1504–1509 (2021).

666 30. Chi, H. Regulation and function of mTOR signalling in T cell fate decisions. *Nat. Rev.*

667   *Immunol.* **12**, 325–338 (2012).

668 31. Erwood, S. *et al.* Saturation variant interpretation using CRISPR prime editing. *Nat.*

669   *Biotechnol.* **40**, 885–895 (2022).

670 32. Levesque, S. *et al.* Marker-free co-selection for successive rounds of prime editing in

671   human cells. *Nat. Commun.* **13**, 5909 (2022).

672 33. Wright, B. W., Molloy, M. P. & Jaschke, P. R. Overlapping genes in natural and

673   engineered genomes. *Nat. Rev. Genet.* **23**, 154–168 (2021).

674 34. Wagle, N. *et al.* Response and acquired resistance to everolimus in anaplastic thyroid

675   cancer. *N. Engl. J. Med.* **371**, 1426–1433 (2014).

676 35. Lorenz, M. C. & Heitman, J. TOR mutations confer rapamycin resistance by preventing

677   interaction with FKBP12-rapamycin. *J. Biol. Chem.* **270**, 27531–27537 (1995).

678 36. Rodrik-Outmezguine, V. S. *et al.* Overcoming mTOR resistance mutations with a new-

679   generation mTOR inhibitor. *Nature* **534**, 272–276 (2016).

680 37. Choi, J., Chen, J., Schreiber, S. L. & Clardy, J. Structure of the FKBP12-rapamycin

681   complex interacting with the binding domain of human FRAP. *Science* **273**, 239–242

(1996).

38.   Oki, T. *et al.* Imaging dynamic mTORC1 pathway activity in vivo reveals marked shifts

that support time-specific inhibitor therapy in AML. *Nat. Commun.* **12**, 245 (2021).

39.   Nelson, J. W. *et al.* Engineered pegRNAs improve prime editing efficiency. *Nat.*

Biotechnol. **40**, 402–410 (2021).

40.   Chresta, C. M. *et al.* AZD8055 is a potent, selective, and orally bioavailable ATP-

competitive mammalian target of rapamycin kinase inhibitor with in vitro and in vivo

antitumor activity. *Cancer Res.* **70**, 288–298 (2010).

41.   Concordet, J. P. & Haeussler, M. CRISPOR: Intuitive guide selection for CRISPR/Cas9

genome editing experiments and screens. *Nucleic Acids Res.* **46**, 242–245 (2018).

42.   Vakulskas, C. A. *et al.* A high-fidelity Cas9 mutant delivered as a ribonucleoprotein

complex enables efficient gene editing in human hematopoietic stem and progenitor cells.

Nat. Med. **24**, 1216–1224 (2018).

43.   Tsai, S. Q. *et al.* GUIDE-seq enables genome-wide profiling of off-target cleavage by

CRISPR-Cas nucleases. *Nat. Biotechnol.* **33**, 187–198 (2015).

44.   Malinin, N. L. *et al.* Defining genome-wide CRISPR–Cas genome-editing nuclease

activity with GUIDE-seq. *Nat. Protoc.* **16**, 5592–5615 (2021).

45.   Roth, T. L. *et al.* Reprogramming human T cell function and specificity with non-viral

genome targeting. *Nature* **559**, 405–409 (2018).

46.   Leung, W.-H. *et al.* Sensitive and adaptable pharmacological control of CAR T cells

through extracellular receptor dimerization. *JCI Insight* **4**, e124430 (2019).

47.   Mahalati, K. & Kahan, B. D. Clinical pharmacokinetics of sirolimus. *Clin.*

Pharmacokinet. **40**, 573–585 (2001).

48.   Araki, K. *et al.* mTOR regulates memory CD8 T-cell differentiation. *Nature* **460**, 108–112

(2009).

49.   Dormond, O., Romero, P., Zhang, L. & Romero, P. Modulation of mTOR signalling

triggers the formation of stem cell-like memory T cells. *EBioMedicine* **4**, 50–61 (2016).

50.   Long, A. H. *et al.* 4-1BB costimulation ameliorates T cell exhaustion induced by tonic

signaling of chimeric antigen receptors. *Nat. Med.* **21**, 581–590 (2015).

51.   Philipson, B. I. *et al.* 4-1BB costimulation promotes CAR T cell survival through

noncanonical NF-kB signaling. *Sci. Signal.* **13**, eaay8248 (2020).

52.   Cappell, K. M. & Kochenderfer, J. N. A comparison of chimeric antigen receptors
containing CD28 versus 4-1BB costimulatory domains. *Nat. Rev. Clin. Oncol.* **18**, 715–
727 (2021).

53.   Bloemberg, D. *et al.* A high-throughput method for characterizing novel chimeric antigen
receptors in Jurkat cells. *Mol. Ther. - Methods Clin. Dev.* **16**, 238–254 (2020).

54.   Muller, Y. D. *et al.* The CD28-transmembrane domain mediates chimeric antigen receptor
heterodimerization with CD28. *Front. Immunol.* **12**, 639818 (2021).

55.   Majzner, R. G. & Mackall, C. L. Clinical lessons learned from the first leg of the CAR T
cell journey. *Nat. Med.* **25**, 1341–1355 (2019).

56.   Wu, C. Y., Roybal, K. T., Puchner, E. M., Onuffer, J. & Lim, W. A. Remote control of
therapeutic T cells through a small molecule-gated chimeric receptor. *Science* **350**,
aab4077 (2015).

57.   Flugel, C. L. *et al.* Overcoming on-target, off-tumour toxicity of CAR T cell therapy for
solid tumours. *Nat. Rev. Clin. Oncol.* **20**, 49–62 (2023).

58.   Shah, N. N. & Fry, T. J. Mechanisms of resistance to CAR T cell therapy. *Nat. Rev. Clin.*
Oncol. **16**, 372–385 (2019).

59.   Fry, T. J. *et al.* CD22-targeted CAR T cells induce remission in B-ALL that is naive or
resistant to CD19-targeted CAR immunotherapy. *Nat. Med.* **24**, 20–28 (2018).

60.   Chang, C. H. *et al.* Metabolic competition in the tumor microenvironment is a driver of
cancer progression. *Cell* **162**, 1229–1241 (2015).

61.   McGuirk, S., Audet-Delage, Y. & St-Pierre, J. Metabolic fitness and plasticity in cancer
progression. *Trends in Cancer* **6**, 49–61 (2020).

62.   Agudelo, D. *et al.* Marker-free coselection for CRISPR-driven genome editing in human
cells. *Nat. Methods* **14**, 615–620 (2017).

63.   Grabiner, B. C. *et al.* A diverse array of cancer-associated MTOR mutations are
hyperactivating and can predict rapamycin sensitivity. *Cancer Discov.* **4**, 554–563 (2014).

64.   Wagle, N. *et al.* Activating mTOR mutations in a patient with an extraordinary response
on a phase I trial of everolimus and pazopanib. *Cancer Discov.* **4**, 546–553 (2014).

65.   Gumber, D. & Wang, L. D. Improving CAR-T immunotherapy: Overcoming the
challenges of T cell exhaustion. *eBioMedicine* **77**, 103941 (2022).

66.   Mestermann, K. *et al.* The tyrosine kinase inhibitor dasatinib acts as a pharmacologic

on/off switch for CAR T cells. *Sci. Transl. Med.* **11**, eaau5907 (2019).

67.   Weber, E. W. *et al.* Pharmacologic control of CAR-T cell function using dasatinib. *Blood*
Adv. **3**, 711–717 (2019).

68.   Weber, E. W. *et al.* Transient rest restores functionality in exhausted CAR-T cells through
epigenetic remodeling. *Science* **372**, eaba1786 (2021).

69.   Fajgenbaum, D. C. & June, C. H. Cytokine storm. *N. Engl. J. Med.* **383**, 2255–2273
(2020).

70.   Morris, E. C., Neelapu, S. S., Giavridis, T. & Sadelain, M. Cytokine release syndrome and
associated neurotoxicity in cancer immunotherapy. *Nat. Rev. Immunol.* **22**, 85–96 (2022).

71.   Alizadeh, D. *et al.* IL15 enhances CAR-T cell antitumor activity by reducing mTORC1
activity and preserving their stem cell memory phenotype. *Cancer Immunol. Res.* **7**, 759–
772 (2019).

72.   Lamarthée, B. *et al.* Transient mTOR inhibition rescues 4-1BB CAR-Tregs from tonic
signal-induced dysfunction. *Nat. Commun.* **12**, 6446 (2021).

73.   Chen, P. J. *et al.* Enhanced prime editing systems by manipulating cellular determinants of
editing outcomes. *Cell* **184**, 5635–5652 (2021).

74.   Nguyen, D. N. *et al.* Polymer-stabilized Cas9 nanoparticles and modified repair templates
increase genome editing efficiency. *Nat. Biotechnol.* **38**, 44.49 (2019).

75.   Brinkman, E. K., Chen, T., Amendola, M. & Van Steensel, B. Easy quantitative
assessment of genome editing by sequence trace decomposition. *Nucleic Acids Res.* **42**,
e168 (2014).

76.   Brinkman, E. K. *et al.* Easy quantification of template-directed CRISPR/Cas9 editing.
Nucleic Acids Res. **46**, e58 (2018).

77.   Xu, L., Liu, Y. & Han, R. BEAT: A Python program to quantify base editing from Sanger
sequencing. *Cris. J.* **2**, 223–229 (2019).

78.   Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence
analysis. *Nat. Biotechnol.* **37**, 215–226 (2019).

79.   Bloh, K. *et al.* Deconvolution of complex DNA repair (DECODR): establishing a novel
deconvolution algorithm for comprehensive analysis of CRISPR-edited Sanger
sequencing data. *Cris. J.* **4**, 120–131 (2021).

80.   Hamieh, M. *et al.* CAR T cell trogocytosis and cooperative killing regulate tumour antigen

escape. *Nature* **568**, 112–116 (2019).

ACKNOWLEDGEMENTS

This study was supported by a grant from Mitacs and by the Fondation du CHU de Québec. JJ. L.

is supported by a CIHR Project Grant (PJT407274). S.L. was supported by a Frederick Banting

and Charles Best Canada graduate scholarship from CIHR, and a Graduate student scholarship

from the Fonds de la Recherche du Québec-Santé (FRQS). G.C. holds a Vanier Canada graduate

scholarship from CIHR. Salary support was provided by the FRQS to Y.D. B.B. and G.M. are

supported by the "Programme d'appui aux plateformes technologiques stratégiques" from the

Ministère de l'Économie, de l'Innovation et de l'Énergie Québec. We thank the staff at the flow

cytometry core facility of the CHU de Quebec Research Center for their training and assistance.

We thank Marie-Ève Paquet and the vector core facility staff at the Canadian neurophotonics

platform for AAV6 production. We also thank Scott McComb (National Research Council) for

kindly providing NALM6-RFP-FLUC cells.

COMPETING INTERESTS

791   S.L. and Y.D. are co-inventors on a patent application (WO 2023/015376 A1) related to this work.

AUTHOR'S CONTRIBUTIONS

794   Conceptualization, S.L. and Y.D.; methodology, S.L., G.C., V.D., C.G., JP. F., S.V., E.N., G.M.,

795   N.D., J.L., B.B., JJ. L., and Y.D.; investigation, S.L., G.C., V.D., C.G., JP. F., S.V., E.N., G.M.,

796   N.D., J.L., B.B., JJ. L., and Y.D.; writing original draft, S.L.; writing, review and editing, S.L.,

797   and Y.D.; supervision, J.L., B.B., JJ. L., and Y.D.; funding acquisition, J.L., JJ. L., and Y.D. All

798   authors reviewed the manuscript and approved its final version.

a

b

c

d

Figure 1. Identification of mutations conferring rapamycin resistance via saturation prime editing. (**a**) Schematic representation of the SpCas9 target sites, the reverse transcriptase (RT) template and the primer binding site (PBS) used to perform genome editing at *MTOR*. To perform saturation prime editing, G3 is used as a pegRNA, and G4 or G5 are used as a complementary nick sgRNA (PE3). The *MTOR*-F2108 codon is mutated to NNK and an additional silent PAM mutation is introduced. (**b**) Schematic of mScarlet-I mTOR signaling indicator (mSc-TOSI) degradation under mTORC1 signaling. (**c**) Histogram plot of mSc-TOSI intensity in bulk populations of cells harboring different *MTOR*-F2108 mutations. Where indicated, cells were treated for 24 hours with 50 nM rapamycin or 50 nM AZD8055 before FACS analysis. Representative images are from one of two independent biological replicates performed at different times with equivalent results. (**d**) High throughput sequencing quantification of the *MTOR*-F2108 mutations introduced via saturation prime editing following rapamycin selection and FACS sorting. The fold enrichment of each *MTOR*-F2108 mutation was calculated from the before selection sample.

a

b

Figure 2. Identification of highly active and specific sgRNAs targeting the *MTOR* locus. (a)

Small indels quantification as determined by TIDE analysis from Sanger sequencing. Primary

CD3$^+$ and CD8$^+$ T cells were transfected with the indicated nuclease/sgRNA RNPs and genomic

DNA was harvested 3 days post-transfection. $n = 2$ independent biological replicates performed at

different times with CD3$^+$ and CD8$^+$ T cells from two different healthy donors. (**b**) Candidate off-

target sites identified using GUIDE-Seq. CD3$^+$ T cells were transfected with SpCas9 RNPs

targeting *MTOR* intron 45 and GUIDE-Seq dsDNA. Four days post-nucleofection, CD3$^+$ T cells

were restimulated and expanded for an additional 7 days. Genomic DNA was harvested after

expansion (11 days post-nucleofection). Dots represent matches with the intended target sequence,

mismatches are colored, and nucleotide bulges are highlighted with a star. GUIDE-Seq read counts

are from one of two independent biological replicates performed with CD3$^+$ T cells from two

different donors (see **Supplementary Fig. 3**).

Figure 3
Levesque et al.

Figure 3. Targeting a CAR to the *MTOR* locus allows the enrichment of rapamycin-resistant CAR-T cells. (**a**) Schematic representation of CD19-CAR-2A-EGFP targeting to the reverse DNA strand of the *MTOR* locus using an AAV6 donor. The F2108L mutation is introduced via the left homology arm (HA) and transgene expression is driven by a human *PGK1* promoter. (**b**) Representative timeline for nucleofection and *ex vivo* rapamycin selection. (**c**) Out-Out PCR for CD19-CAR-2A-EGFP knock-in detection at *MTOR*. Whole blood (WB) primary CD3$^+$ T cells from WB donor 1 (WB 1) were transfected with a SpCas9-G4 RNP and transduced with an AAV6 vector with the indicated multiplicity of infection (MOI). T cells were treated (rapamycin) or not (vehicle) with 25 nM rapamycin 3 days post-transfection for 8 days. The percentage of edited alleles was determined by TIDER from Sanger sequencing. $n = 1$ experiment. (**d**) Same as in (**c**), but FACS-based quantification of targeted CD19-CAR-2A-EGFP integration. (**e**) Same as in (**c,d**) with a MOI of 5x10^3 and primary CD3$^+$ T cells isolated from WB or a leukocyte reduction system (LRS) from three additional healthy donors. $n = 3$ independent biological replicates performed in triplicate at different times with CD3$^+$ T cells from the indicated healthy donor. Each data point represents a technical replicate. (**e**) Luciferase-based cytotoxicity assay. Following rapamycin selection, CD19-CAR-T cells were incubated with the indicated effector to target (E:T) ratio with NALM6 cells stably expressing firefly luciferase (FLUC) and 25 nM rapamycin. Rapamycin-resistant NALM6 cells harboring the *MTOR*-F2108L mutation (NALM6-RapaR) were used as a control to analyze the combinatorial impact of rapamycin. Luminescence was measured after 18 hours of incubation. $n = 2$ independent biological replicates performed at different times with CD3$^+$ T cells from two different healthy donors (WB donor 1 and LRS donor 2). Each data point represents the average of three technical replicates. *hPGK1*, human phosphoglycerate kinase 1 promoter. PA, polyadenylation signal. HA, homology arm.

Figure 4
Levesque et al.

Figure 4. Combinatorial CAR-T cell therapy with rapamycin slows leukemia progression *in*

***vivo*. (a)** Timeline and experimental setup for the pre-B acute lymphoblastic leukemia xenograft

mouse model. Male and female NSG mice were challenged with 0.5 x 10^5 NALM6-RFP-FLUC

cells and daily rapamycin treatment (4 mg/kg) started three days later. Four days after tumor

inoculation, 2 x 10^6 untransduced (UT) T cells or CD19-CAR-T cells were injected (Donor ID

WB2, see **Fig. 3e**). (**b**) Bioluminescence (BLI) quantification and Kaplan-Meier survival plot.

Leukemia engraftment, bio-distribution, and tumor progression were assessed by BLI imaging two

times per week. The radiance (photons/s) of the regions of interest (ROIs) corresponds to the area

containing the whole back side of the body. $n = 8$ mice per group. (**c**) BLI and tumor bio-

distribution of all mice from (**b**) over seven weeks. The color barcode represents the radiance scale

(photons/s/cm^2/sr).

Figure 5. Multiplexed DARIC targeting to the *MTOR* and *TRAC* loci allows pharmacological control of CAR-T cells activity. (**a**) Schematic representation of DARIC-1 targeting to the reverse DNA strand of the *MTOR* locus, and DARIC-2 and 3 targeting to the *TRAC* locus using AAV6 donors. (**b**) Out-Out PCR for DARIC-1 knock-in detection at *MTOR* and DARIC-2 and 3 knock-in at *TRAC*. Whole blood (WB) primary CD3$^+$ T cells were transfected with a SpCas9-G4 RNP and transduced with AAV6 vectors with a multiplicity of infection (MOI) of 5x10^3 for each AAV6. Genomic DNA was harvested 3 days post-transfection. The percentage of edited alleles was determined by TIDER from Sanger sequencing. Representative images are from one of two independent biological replicates. (**c**) FACS-based cytotoxicity assay. CD3$^+$ T cells were electroporated with *MTOR*/*TRAC* RNPs and transduced with both *MTOR*/*TRAC* AAV6 donors to generate rapamycin-resistant (RapaR) CD19-CD22-DARIC-T cells. For rapamycin-sensitive CD22-DARIC-T cells, the *MTOR* AAV6 donor was omitted. Three days post-nucleofection, DARIC-T cells were incubated in the presence (rapamycin) or absence (vehicle) of 25 nM rapamycin with NALM6 cells stably expressing RFP and K562-RapaR stably expressing EBFP with an effector to target ratio of 1:1. Rapamycin-resistant NALM6 cells harboring the *MTOR*-F2108L mutation (NALM6-RapaR) were used as a control to analyze the combinatorial impact of rapamycin. The percentage of cytotoxicity was measured by FACS after 72 hours of incubation. *n* = 2 independent biological replicates. Each data point represents the average of three technical replicates.