Joint modelling of whole genome sequence data for human height via approximate message passing

Al Depope[1,*], Jakub Bajzik[1], Marco Mondelli[1,†,*], Matthew R. Robinson[1,†,*]

1 Institute of Science and Technology Austria, Klosterneuburg, Austria.

* corresponding authors
† indicates joint supervision

Abstract

Human height is a model for the genetic analysis of complex traits, and recent studies suggest the presence of thousands of common genetic variant associations and hundreds of low-frequency/rare variants. Here, we show that existing variable selection methods may fail to either select the correct genomic regions, or the appropriate genetic variants due to the complex and potentially long-distance correlation structure of whole genome sequence (WGS) data. To address this, we develop a new algorithmic paradigm based on approximate message passing, gVAMP, which we show accurately localizes associations to variants with the correct frequency and position in the DNA. We apply gVAMP to jointly model the relationship of tens of millions of WGS variants with human height in hundred of thousands of UK Biobank individuals. We find a power relationship of effect size and frequency of -1.1 ± 0.2, identify 59 rare variants and gene burden scores alongside many hundreds of DNA regions containing common variant associations, and show that understanding the genetic basis of complex traits will require the joint analysis of hundreds of millions of variables measured on millions of people. The polygenic risk scores obtained from gVAMP have high accuracy (including a prediction accuracy of $\sim 46\%$ for human height) and outperform current methods for downstream tasks such as mixed linear model association testing across 13 UK Biobank traits. In conclusion, gVAMP offers a scalable foundation towards a wider range of analyses in WGS data.

Keywords: whole genome regression; variable selection; polygenic risk scores; approximate message passing

Introduction

Efficient utilization of large-scale biobank data is crucial for inferring the genetic basis of disease and predicting health outcomes from DNA. By mapping associations to regions of the DNA, we seek to understand the underlying genetic architecture of phenotype. The common approach is to use single-marker or single-gene burden score regression [1–4], which gives marginal associations that do not account for linkage disequilibrium (LD) among loci. Fine-mapping methods are then employed with the aim of controlling for LD when conducting variable selection and resolving the associations to a set of likely causal variants [5].

Here, we first show that in whole genome sequence (WGS) data, even weak associations that are physically distant from causal variants will be discovered as significant at scale when using marginal association testing. Applying fine-mapping methods, which focus only on genome-wide significant loci within one region at a time in isolation from the rest of the genome, then results in severe miscalibration with a compromise of power. To address these issues we propose a new framework, gVAMP, a form of whole genome regression (WGR), capable of fitting tens of millions of WGS variants jointly for hundreds of thousands of individuals. We extensively benchmark gVAMP in simulations on WGS data and show it accurately localizes associations to variants with the correct frequency and position in the DNA, outperforming marginal testing and marginal testing plus fine-mapping. We then describe an application of gVAMP to WGS data and measurements of human height for hundreds of thousands of UK Biobank participants, where we find a stronger relationship of effect size and variant frequency than previously considered [6].

We additionally validate the application of gVAMP for polygenic risk score prediction on a number of datasets by benchmarking against the state of the art. Here, gVAMP outperforms widely used summary statistic methods such LDpred2 [7] and SBayesR [8]; its performance matches that of MCMC sampling schemes [9], but with a dramatic speed-up in time (analysing 8.4M SNPs jointly in under a day as opposed to weeks); and it provides a small improvement over the recently proposed LDAK-KVIK approach [10]. When polygenic risk scores are used for association testing, gVAMP also outperforms both the REGENIE [1] and LDAK-KVIK [10] software. In summary, our approach lays the foundations for a wider range of analyses in large WGS datasets.

Results

Overview of the approach

Our focus is on the simple idea of joint modeling of all genetic variants genome-wide controlling for local and long-range LD. To do this, we consider a general form of whole-genome Bayesian variable selection regression (BVSR), common to genome-wide association studies (GWAS) [8, 9], estimating the effects vector $\boldsymbol{\beta} \in \mathbb{R}^P$ from a vector of phenotype measurements $\boldsymbol{y} = (y_1, \ldots, y_N) \in \mathbb{R}^N$ given by

$$y_i = \langle \boldsymbol{x}_i, \boldsymbol{\beta} \rangle + \epsilon_i, \quad \text{for } i \in \{1, \ldots, N\}. \tag{1}$$

Here, $\boldsymbol{x}_i$ is the row of the normalized genotype matrix $\boldsymbol{X}$ corresponding to the i-th individual, $\langle \boldsymbol{x}_i, \boldsymbol{\beta} \rangle = \boldsymbol{x}_i^T \boldsymbol{\beta}$ denotes the inner product, and $\boldsymbol{\epsilon} = (\epsilon_1, \ldots, \epsilon_N)$ is an unknown noise vector with

multivariate normal distribution $\mathcal{N}(0, \gamma_\epsilon^{-1} \cdot \boldsymbol{I})$ and unknown noise precision γ_ϵ^{-1}. To allow for a range of genetic effects, we select the prior on $\boldsymbol{\beta}$ to be of an adaptive spike-and-slab form:

$$\beta_j \sim (1 - \lambda) \cdot \delta_0(\cdot) + \lambda \cdot \sum_{l=1}^{L} \pi_l \cdot \mathcal{N}(\cdot, 0, \sigma_l^2), \quad \text{for } j \in \{1, \ldots, P\}, \qquad (2)$$

where we learn from the data: the DNA variant inclusion rate $\lambda \in [0, 1]$, the unknown number of Gaussian mixtures L, the mixture probabilities $(\pi_l)_{l=1}^{L}$ and the variances for the slab components $(\sigma_l^2)_{l=1}^{L}$.

Current commonly applied association testing [1–4], fine-mapping [5] and polygenic risk score methods [9, 11] are all based on forms of Equations (1) and (2), with parameters estimated by restricted maximum likelihood (REML), Markov Chain Monte Carlo (MCMC), expectation maximisation (EM), or variational inference (VI). REML and MCMC are computationally intensive and slow; while EM and VI are faster, they trade speed for accuracy with few theoretical guarantees. Furthermore, current software implementations of these algorithms limit either the number of markers or individuals. Mixed linear model association (MLMA) approaches are typically restricted to using less than one million SNPs to control for the polygenic background [1, 2], resulting in a loss of power [9] and the potential for inadequate control for fine-scale confounding factors [12]. Polygenic risk score algorithms are limited to a few million SNPs, and lose power by modeling only blocks of genetic markers [7, 8]. Likewise, fine-mapping methods are generally limited to focal segments of the DNA [5], and they are unable to fit all genome-wide DNA variants together. Thus, no existing approach can apply the statistical model of Equations (1) and (2) to truly estimate the effects vector $\boldsymbol{\beta}$ genome-wide in large-scale WGS data.

We overcome this issue by developing a new approach for GWAS inference, dubbed *genomic Vector Approximate Message Passing* (gVAMP). Approximate Message Passing (AMP) [13–15] refers to a family of iterative algorithms with several attractive properties: *(i)* AMP allows the usage of a wide range of Bayesian priors; *(ii)* the AMP performance for high-dimensional data can be precisely characterized by a simple recursion called state evolution [16]; *(iii)* using state evolution, joint association test statistics can be obtained [17]; and *(iv)* AMP achieves Bayes-optimal performance in several settings [17–19]. However, we find that existing AMP algorithms proposed for various applications [20–23] cannot be transferred to biobank analyses as: *(i)* they are entirely infeasible at scale, requiring expensive singular value decompositions; and *(ii)* they give diverging estimates of the signal in either simulated genomic data or the UK Biobank data. To address the problem, we combine a number of principled approaches to produce an algorithm tailored to whole genome regression as described in the Methods (Algorithm 1).

gVAMP provides both *(i)* a point estimate of the posterior mean $\mathbb{E}[\boldsymbol{\beta} \mid \boldsymbol{X}, \boldsymbol{y}]$ which can be directly used to create a polygenic risk score, and *(ii)* a statistical framework, via state evolution, to test whether each marker makes a non-trivial contribution to the outcome. Specifically, state evolution yields both p-values [17] and posterior inclusion probabilities (PIP) for each SNP (see "gVAMP state evolution association testing" in the Methods).

BVSR approaches can, in principle, assess uncertainty in which variables to select, even when the variables are highly correlated. However, typically a complex posterior distribution is obtained (or

samples approximating it), which can be difficult to interpret. Indeed, highly correlated variables pose an obvious challenge as they are difficult to distinguish from one another, resulting in a loss of power to select them. One approach to this is to construct credible sets of correlated variables (given a pre-defined correlation or 'purity') that capture relevant variables from an approximation of the posterior [5]. However, this is computationally challenging at scale and, thus, limited in practice to small data subsets, requiring ad hoc decisions about how large a given DNA region should be. Additionally, our simulation study below shows that this approach can considerably overstate the number of significantly associated genomic regions when applied to WGS data one-region-at-a-time. Another approach, that may be more applicable to large-scale applications, is to focus on selecting groups of variables, rather than individual ones. Here, we use a simple version of this idea, which is to first cluster into groups only commonly occurring very highly correlated variables, selecting the most common 'tagging' variant, leaving all rare variation unchanged. As gVAMP is a BVSR model with a sparsity inducing prior that is adaptively learned from the data, we can then determine the probability that either single variants, or groups of highly correlated common variants make a contribution to the phenotypic variance. The only choice is to define the correlation to create the groups, and we note that other variable selection methods applied to genomic studies typically use ≥ 0.5 to pre-define the correlation of the selected variables [5].

Here, we select an absolute correlation of ≥ 0.6, which enabled us to analyze a large data set of nearly 20 million WGS variants jointly, in a way that keeps the majority of the structure of the data unchanged. We validate below the choice of an absolute correlation ≥ 0.6 in WGS data: a lower correlation leads to worse variable selection performance; a higher correlation does not guarantee a statistically significant improvement, while leading to a larger data set which is computationally more expensive to analyze.

Finally, we learn all unknown parameters in an adaptive Bayes expectation-maximization (EM) framework [24, 25], which avoids expensive cross-validation and yields biologically informative inference of the phenotypic variance attributable to the genomic data (SNP heritability, h^2_{SNP}) allowing for a full genome-wide characterization of the genetic architecture of human complex traits in WGS data.

Variable selection in sequencing data

We first assess the ability of the gVAMP framework to estimate genetic effects, by taking UK Biobank WGS data from chromosome 12 (1,922,505 SNPs and 411,536 unrelated individuals of European ancestry) and simulating a wide range of phenotypes that differ in *(i)* the variance attributable to the SNPs, *(ii)* the number of causal variants (163 or 400), *(iii)* the frequency of the causal variants, and *(iv)* the relationship of marker frequency and effect size (see Methods and Supplementary Table S1 for an overview of the scenarios). We allocate causal variants within a window (20Mb width for 400 causal variants, or 8Mb width for 163 causal variants) of chromosome 12, selecting one SNP as causal within a block of 50kb, leaving the remainder of the chromosome as null (Figure 1a), see Methods for more details. Following standard practice, we adjust the simulated phenotype by the leading 20 principal components (PCs) calculated from the WGS data of chromosome 12 and analyze the data with either: *(i)* standard GWAS marginal testing one-SNP-at-a-time; *(ii)* our proposed gVAMP; *(iii)* the SuSiE variable selection model [5] run on 50kb regions for which a

significant variant was discovered in the GWAS.

We evaluate the localization accuracy of our variable selection with respect to *(i)* the correct position in the DNA (within 50kb), *(ii)* a variant with a given correlation to a true underlying causal variant, and *(iii)* a variant with the correct frequency with respect to the correlated causal variant. Across all 12 simulation scenarios, we find that marginal GWAS testing in WGS data discovers associations that are correlated to, but potentially physically distant from, the simulated causal variants (Figure 1a gives an example of this finding, which occurs across all settings). When the marginal results are then used to identify DNA regions upon which to conduct variable selection with SuSiE, this in turn leads to SuSiE having a large number of discoveries across the DNA (Figure 1a), which are often caused by significant correlations with very distant causal variants (Figure 1d). In contrast, gVAMP outperforms SuSiE across all 12 different simulation scenarios (Figure 1b, Figure S1). For gVAMP, we use the per-SNP PIP to directly compare to SuSiE, and we find that gVAMP localizes associations to the correct position in the DNA (within 50kb) with a well-calibrated error rate that is far lower than marginal testing, or marginal testing plus variable selection with SuSiE (Figure 1b, Figure S1, Figure S2). Specifically, when assessing performance, we define *(i)* the true positive rate (TPR) as the number of identified 50kb windows divided by the total number of windows containing a causal variant, and *(ii)* the false discovery rate (FDR) as the number of identified 50kb windows that do not contain a causal variant divided by the total number of windows identified. We say that a window is identified if the algorithm returns a variant in it.

This way of assessing performance consists of localizing an association to the correct 50kb window in the DNA, and it is akin to treating associated variants that are distant as independent, which corresponds to the underlying assumption for running SuSiE fine-mapping in blocks of the DNA. By selecting PIP ≥ 0.95, our simulations suggest that, across most settings, we have approximately probability 95% of detecting the correct genomic region (Figure S2). We note that SuSiE is not designed to differentiate null genomic regions from those containing an effect, rather it is designed so that marginal estimates are used to first select the regions, and then SuSiE determines the PIP of variants within the window.

We also assess the ability of gVAMP to localize the signal, by creating a list of variants correlated to a given threshold to each identified variant within a larger DNA region (1Mb). In this context, we define a false positive if a causal variant is not present within the list. We find that gVAMP generally detects associations that have absolute correlation ≥ 0.4 to a causal variant, with again a calibrated error rate (Figure S3). Furthermore, we find that the frequency of the discovered SNPs from gVAMP matches that of the correlated true causal variants (Figure 1c and Figure S4). Thus, across our WGS simulation settings, gVAMP localizes associations to within 50kb of a causal variant, to variants of similar frequency to the causal variants, and generally with a correlation of at least 0.4 to a causal variant (Figure S3). We note that the receiver operator curves (ROC, plotting TPR as a function of FDR) results do not depend upon whether we do p-value or PIP association testing with gVAMP (Figure S5).

The advantage of the gVAMP framework is confirmed in a large set of additional simulation comparisons. In particular, we find that giving SuSiE the true prior as initialization, running SuSiE in windows up to 500kb, or running SuSiE-inf combining inference across blocks only marginally improves performance (Figure S6). We also find that the results do not depend upon the choice of

the variable selection approach, as multiple different implementations of FINEMAP all yield results ₁₅₈ that are similar to equivalent SuSiE implementations (Figure S6). ₁₅₉

We note that, prior to all analyses presented in this work, we make a design choice in the selection of WGS data, where: *(i)* we pick variants of MAF $> 10^{-4}$ in order to allow a reasonable signal recovery power, as given a sample size of roughly 400,000 individuals, only about 80 observations are non-zero at this cut-off; and *(ii)* we create sets of highly correlated variants with absolute correlation > 0.6 within a 1Mb radius and then select the single most common variant from each set, which is motivated in our discussion above (Figure S7). We find that these analysis design choices do not bias the results of gVAMP: in all of our simulation settings, we select causal variants from the full WGS data of chromosome 12 and, when we analyze the processed data, we find that *(i)* the variance attributable to SNPs of different frequency is generally unbiased (Figure 1e and Figure S8), and *(ii)* the frequency of the discovered SNPs matches that of the correlated true causal variants (Figure 1c and Figure S4). A slight bias exists for common variants, where the frequency of the variants selected by gVAMP can around 1-2% higher than the correlated causal variant, which is due to picking the most common variant from each set when grouping common highly correlated variants (Figure S4). Finally, we show that the direction of the simulated power relationship between effect size and minor allele frequency is also recovered by gVAMP (Figure S9). Taken all together, these findings suggest that existing methods fall short of the performance required for variable selection in WGS data, while gVAMP both reliably localizes the signal to the correct genomic regions and it selects a variant of appropriate frequency.

Finally, we examine whether the genetic effects recovered in imputed data are the same as those inferred in WGS data across our 12 scenarios. For the same set of UK Biobank individuals, we use the phenotypes simulated from the full WGS data, but we swap the WGS data for the latest imputed SNP data release. We use the same data selection as for the WGS, where we filter for variants of MAF > 0.0001, create sets of highly correlated variants with absolute correlation > 0.6 within a 1Mb radius and also select imputation info score 0.8. We find that the frequency of the SNPs discovered at ≥ 0.95 PIP is generally closer to that of the correlated true causal variant for WGS data, as compared to imputed data (Figure 1c and Figure S4). Furthermore, while the analysis of WGS data consistently recovers the variance attributable to SNPs of different frequency, analyzing the imputed data leads to a significant misestimation in several of the simulation scenarios we consider (Figure S8).

The genetic architecture of human height in WGS data ₁₈₉

We describe the genetic architecture of human height inferred from 16,854,878 WGS variants observed for 415,000 UK Biobank individuals. We find significant amounts of common genetic variation detected and a considerable contribution of rare genetic variants (Figure 2). Generally, the regions identified are replicated in the most recent GWAS study of human height [26] within 100-500kb (Figure 2a). We find that for rare variants identified in the WGS analysis, variants discovered by Yengo *et al.* [26] of strongest correlation and most similar MAF significant at $p \leq 5 \cdot 10^{-8}$ within a 500kb region, are all more common in frequency (Figure 2b). There is a drop in association testing power for rare variants, with lower average marginal Chi-squared test statistic values for the rarest WGS variants identified (Figure 2c) and this, coupled with few rare variants being present in the

Figure 1. Simulation study of WGS UK Biobank data chromosome 12 where 6.25% of the phenotypic variance is attributable to 163 SNPs, with a power relationship between the effect size and frequency (setting d in Supplementary Table S1). *(a)* The y-axis shows -log10 of 1 minus the posterior inclusion probability (PIP) of gVAMP (blue), -log10 of 1 minus the PIP of the fine-mapping approach SuSiE (orange), and -log10 of the p-values obtained from marginal one-SNP-at-a-time regression testing (gray). Simulated causal variants are all allocated to the left of the black line, leaving the rest of the chromosome as "null". While marginal estimates and SuSiE PIPs consistently discover variants in the null region, in the same region the PIP values of gVAMP are generally well controlled below 10^{-10}. *(b)* The true positive rate and false discovery rate of marginal testing plus SuSiE (orange) and gVAMP (blue) calculated as associations in the same 50kb window as a true causal variant. *(c)* The average frequency difference of a gVAMP selected variant from the most correlated true causal variant for rare and common variants. The frequency of variants selected by gVAMP in either WGS or imputed data is unbiased for either rare or common SNPs. *(d)* The maximum squared correlation of the discoveries made by SuSiE that were not in the same 50kb window as a true causal variant. Large correlations with the causal variants localized in the first 8-20Mb are rather common across the whole chromosome, especially for rare WGS variants. *(e)* The proportion of the sum of the squared regression coefficients attributable to variants of different frequency as estimated from gVAMP in WGS or imputed SNP data. The ground truth was calculated as the sum of squared true effects. The variance attributable to SNPs of different frequency matches the ground truth for gVAMP in WGS and imputed data. Error bars (in *(c)* and *(e)*) or bands (in *(b)*) give the standard deviation across 5 simulation replicates.

study by Yengo *et al.* [26] explains the discrepancy. Taken together, this suggests that when we discover a region in the WGS analysis, similar DNA regions have previously been identified, but the effects may be localized to different sets of variants.

We can directly determine the power relationship between effect size and minor allele frequency, which is close to -1, a value significantly larger than that previously reported within the empirical literature [6] (Figure 3a and Table S3). We note that under a population genetic model of stabilizing selection α should be -1 for variants with large enough effect and closer to zero for weaker variants [27]. We estimate the effects of 17,852 WES gene burden scores when fit alongside the WGS data (Figure 3b) and sum up the joint effects to determine the variation in height attributable to each gene (Figure 3c). We see little relationship between the variance attributable to the burden score of a given gene and the SNPs annotated to the gene (Figure S10), with some notable exceptions of: ACAN, ADAMTS17, ADAMTS10, and LCORL. Additionally, across annotations, we find that the phenotypic variance attributable to different DNA regions is higher for intergenic and intronic

Figure 2. The genetic architecture of human height inferred by gVAMP from 16,854,878 whole genome sequence variants in the UK Biobank. *(a)* For each whole genome sequence (WGS) variant discovered at PIP $\geq 95\%$, we determine whether we also identify a variant within a given number of base pairs (distance, x-axis) from: *(i)* the latest height GWAS study by Yengo *et al.* [26], selecting variants at significance $p \leq 5 \cdot 10^{-8}$ from the marginal summary statistics for European individuals excluding the UK Biobank; or *(ii)* an analysis of imputed SNP data with gVAMP, selecting variants with PIP $\geq 95\%$. The proportion overlap (y-axis) is calculated as the proportion of the identified WGS variants that are discovered within a certain base pair distance. *(b)* For each WGS variant discovered at PIP $\geq 95\%$, we determine: *(i)* the variant discovered by Yengo *et al.* of most similar MAF significant at $p \leq 5 \cdot 10^{-8}$ within a 500kb region; or *(ii)* the imputed variant of most similar minor allele frequency (MAF) discovered at PIP $\geq 95\%$ within a 500kb region. We then calculate the mean difference in MAF of the identified WGS variant and the imputed/GWAS variant, with error bars showing twice the standard error. *(c)* For each WGS variant discovered at PIP $\geq 95\%$, we determine the maximum marginal association test statistic of any variant with absolute correlation ≥ 0.6. Boxplots of these maximum Chi-squared test statistic values are then given for different MAF groups, showing a drop in average power to detect rare variants. *(d)* Density plot of the MAF distribution of all discovered variants at PIP $\geq 95\%$ for 2,174,071 imputed SNPs ("2M imputed SNPs"), 8,430,446 imputed SNPs fit jointly alongside 17,852 whole exome sequencing (WES) gene burden scores ("8M imputed SNPs + WES burden scores"), and 16,854,878 WGS SNPs fit jointly alongside 17,852 WES gene burden scores ("17M imputed SNPs + WES burden scores"). Irrespective of the data analyzed the frequency distributions are similar, with the exception that the WGS analysis identifies a greater number of variants (both SNPs and gene burden scores) at lower frequencies. *(e)* We calculate the proportional contribution to phenotypic variance across different MAF groups for 8,430,446 imputed SNPs fit jointly alongside 17,852 whole exome sequencing (WES) gene burden scores ("8M imputed SNPs + WES burden scores"), and 16,854,878 WGS SNPs fit jointly alongside 17,852 WES gene burden scores ("17M imputed SNPs + WES burden scores"). A larger proportion of variance is attributable to rare variation at the expense of more common variants in WGS data.

variants (Figure 3d). However, when adjusting for the number of SNPs contained by each category ₂₁₂

Figure 3. The relationship between effect size and minor allele frequency, burden score effects and the genetic architecture of human height inferred from 16,854,878 whole genome sequence (WGS) variants in the UK Biobank. *(a)* For different posterior inclusion probability thresholds (0.5 to 0.95 and ≥ 0.95), we plot the relationship between joint effect size and minor allele frequency (MAF) for 16,854,878 WGS variants ("17M WGS SNPs"), 8,430,446 imputed SNP variants ("8M imputed SNPs") and 17,852 whole exome sequence (WES) gene burden scores fit alongside either the WGS or imputed variants. *(b)* Across 17,852 WES gene burden scores, we find general concordance of the estimated effect sizes when fit alongside either imputed or WGS data, for most but not all genes, with squared correlation 0.532. *(c)* Likewise, we also find general concordance of the phenotypic variance attributable to markers annotated to each of the 17,852 genes, when fit conditional on either imputed or WGS variants, with squared correlation 0.494. *(d)* Finally, we show similar patterns of enrichment when annotating markers to functional annotations in either the average effect size relative to the genome-wide average effect size (labeled "effect size") or the proportion of variance attributable to each group (labeled "variance"), from joint estimation of either imputed or WGS data.

by using the average effect size of the group relative to the average effect size genome-wide, we find that exonic variants that are nonsynonymous, splicing and stop-gain have the largest average effects of the WGS variants included within our model (Figure 3d).

19 out of the 21 genome-wide significant gene burden scores are for genes that have previous GWAS height associations linked to them in Open Targets, but our analysis suggests that the effect is attributable to a rare protein coding variant rather than the common variants suggested by current genome-wide association studies. This includes insulin-like growth factor 2 genes IGF1R and IGF2BP2, and known height genes ACAN, ADAMTS10 and ADAMTS17, CRISPLD1, DDR2, FLNB, HMG20B, LTBP2, LCORL, NF1, and NPR2. Novel genes are COL27A1 which encodes a collagen type XXVII alpha 1 chain, and HECTD1. The top genes, where gVAMP attributes $\geq 0.04\%$ of height variation in addition to those listed above are EFEMP1, ZBTB38, and ZFAT. We find that many genome-wide significant height-associated rare variants discovered in the WGS data were not found in previous UK Biobank analyses in Open Targets including: rs116467226, an intronic variant by TPRG1; rs766919361, an intronic variant by FGF18; rs141168133, an intergenic variant 19kb from ID4; rs150556786, an upstream gene variant for GRM4; rs574843917, a non-coding transcript exon variant in GPR21; rs532230290, an intergenic variant 42kb from SCYL1; rs543038394, intronic in OVOL1; rs1247942912, a non-coding transcript exon variant in AC024257.3; rs577630729, a regulatory region variant for ISG20; and rs140846043, a non-coding transcript exon variant of MIRLET7BHG. 10 out of our top 38 height-associated WGS variants of $\leq 1\%$ MAF were not previously discovered, and only become height-associated when conditioning on the entire polygenic background captured by WGS data within our analysis. Generally though, marginal estimates made in WGS data for height have similar joint effect sizes when included within the gVAMP model (Figure S11).

We recall that rare variants are directly observed in WGS data as opposed to being largely inferred within imputed data. Rare variants are also more prone to imputation errors and the subset of variants that can be inferred accurately must, by definition, have some correlation to the variants measured on the genotyping array platform [28]. This is confirmed by our finding that using the same MAF threshold of 0.0001 as for the WGS data results in estimates with small out-of-sample prediction accuracy from either gVAMP (6% correlation), ridge regression (4% correlation), or an elastic net (3% correlation). All of these models appear to fit the training data well, but they do not generalise well out-of-sample, and this is likely due to the highly correlated nature of the dataset creating difficulties in separating the effects of correlated rare and common variants. Thus, we can only make comparisons between WGS and imputed data by increasing the MAF threshold used to 0.002, creating a set with 8,430,446 imputed SNPs and a subset of 2,174,071 SNPs (see Methods). For WGS variants discovered at posterior inclusion probability PIP $\geq 95\%$, we determine whether the same variant, or a variant within a given number of base pairs (distance, x-axis), is identified in imputed SNP data at the same PIP threshold. We find moderate overlap in the locations of the SNPs that we detect across the two datasets within 10kb (a small proportion maps to the same location) and substantial overlap at over 100kb either side of the WGS findings (Figure 2a). Generally, we find that the variants found are similar in distribution across the WGS and imputed SNP analyses, with significant amounts of common genetic variation detected and a slight increase in the frequency of the common variants selected by gVAMP in WGS data (Figure 2d). The exception

are rare variants, where the WGS analysis identifies a greater proportion of variants (both SNPs ₂₅₅
and gene burden scores) at lower frequencies (Figure 2d) and finds a larger proportion of variance ₂₅₆
attributable to rare variation at the expense of slightly less variation attributable to common variants ₂₅₇
(Figure 2e). We find a general concordance of the estimated effect sizes for most but not all WES ₂₅₈
gene burden scores (Figure 3b), and a general concordance across markers annotated to each of ₂₅₉
the 17,852 genes, conditional on all other markers, across the imputed SNP and WGS analysis ₂₆₀
(Figure 3c). WES gene burden score effects can differ in the WGS analysis by being either: *(i)* more ₂₆₁
strongly associated because of improved conditioning on the genetic background, or *(ii)* attenuated ₂₆₂
because the variants that comprise the burden score are also included as variants in the analysis. ₂₆₃
Finally, across annotations, we find similar patterns across both WGS and imputed SNP analyses ₂₆₄
(Figure 3d). ₂₆₅

gVAMP estimates the proportion of phenotypic variance in human height attributable to WGS ₂₆₆
data as 0.652, a value that is comparable to 0.63 for the imputed SNP data, to the previously ₂₆₇
published REML estimate in a different WGS dataset [29] and to family-based estimates [30]. This ₂₆₈
is expected as the imputed data is a subset of the WGS data and very low frequency variants ₂₆₉
are expected to make little contribution to the population-level variance, so that the addition of ₂₇₀
large numbers of rare variants will have little impact, especially if they capture signal that was ₂₇₁
previously weakly tagged by more common variation (Figure 2). gVAMP identifies 2,070 sets of ₂₇₂
variants in the 2,174,071 imputed SNP data at PIP ≥ 95%, which is consistent with findings from ₂₇₃
mixed-linear model association testing described in the next section within imputed data (Table S4). ₂₇₄
We note that in the WGS data gVAMP identifies 498 genome-wide significant effects at PIP ≥ 95%. ₂₇₅
There are two potential explanations for this. The first is that some of the height associations are ₂₇₆
re-allocated to DNA variants of lower minor allele frequency, which then have lower power to be ₂₇₇
detected above a PIP ≥ 95% threshold. The second explanation is that two variants can have very ₂₇₈
low correlation with each other in the imputed data, while still both being moderately correlated to ₂₇₉
a third SNP. If the third SNP is not in the imputed panel but is height associated, this would result ₂₈₀
in two quasi-independent associations in the imputed data, but only one in the WGS data. We ₂₈₁
find that our 498 WGS findings are within 50kb of 1,042 of the 2,070 sets of variants identified by ₂₈₂
gVAMP in the 2,174,071 imputed SNP data and are within 500kb of 1989 of the 2,070 variant sets. ₂₈₃
Thus, the WGS gVAMP findings are often neighbored by multiple variants discovered by gVAMP ₂₈₄
in the imputed SNP data. ₂₈₅

Validation and benchmarking of gVAMP ₂₈₆

As gVAMP is the only algorithm that scales to tens of millions of WGS variants, we can only ₂₈₇
validate and benchmark against state-of-the-art polygenic risk score prediction approaches in a ₂₈₈
number of alternative datasets. Applying our model to 13 traits in the UK Biobank imputed data ₂₈₉
(training data sample size and trait codes given in Table S1), we compare the prediction accuracy of ₂₉₀
gVAMP to the widely used summary statistic methods LDpred2 [7] and SBayesR [8], and to the ₂₉₁
individual-level methods GMRM [9] and LDAK [10]. gVAMP matches or outperforms all methods ₂₉₂
for most phenotypes (Figure 4 and Table 1 with the best prediction accuracy highlighted in bold). ₂₉₃
Specifically, for human height (HT), our prediction accuracy is the highest and individual-level ₂₉₄
approaches clearly outperform summary statistic ones: we obtain an accuracy of 45.7%, which is a ₂₉₅

Figure 4. Validating gVAMP through polygenic risk score accuracy and association testing benchmarks in the UK Biobank within imputed SNP data. *(a)* Relative prediction accuracy of gVAMP in a hold-out set of the UK Biobank across 13 traits as compared to other approaches. *(b)* Relative number of leave-one-chromosome-out (LOCO) associations of gVAMP across 13 UK Biobank traits as compared to other approaches at 8,430,446 markers.

97.8% relative increase over LDpred2 and a 26.2% relative increase over SBayesR (Figure 4a and Table 1). We also note that our polygenic risk score has higher accuracy than that obtained from SBayesR using the latest height GWAS study of 3.5M people of 44.7% [26], despite our sample size of only 414,055. Thus, the challenge now is to use efficiently the increased sample size coming from multiple human cohorts, see "Discussion" for additional details.

The polygenic risk score accuracy presented may represent the limit of what is achievable for these traits at the UK Biobank sample size. While modeling 2.17M SNPs improves accuracy over modeling 880,000 variants, we find rather small differences when going from 2.17M to 8M. This is because the 2.17M data set is an LD-grouped subset of the 8M data, and the addition of highly correlated variants captures little additional phenotypic variance. gVAMP estimates the h_{SNP}^2 of each of the 13 traits at an average of 3.4% less than GMRM when using 2,174,071 SNP markers, but at an average of only 3.9% greater than GMRM when using 8,430,446 SNP markers. This implies that only slightly more of the phenotypic variance is captured by the SNPs when the full imputed SNP data are used (Figure S12). We also note that moving from the imputed data to the WGS data, leads to a reduction in out-of-sample prediction for human height. This is likely because of the small increase in variance explained and the reduction in the MAF of the markers included within the WGS model, which are then largely absent from the test set. In summary, gVAMP (an adaptive mixture) performs similarly to LDAK (an elastic net) and GMRM (an MCMC sampling algorithm), improving over both when analyzing the full set of 8,430,446 imputed SNPs (Table 1 and Figure 4).

Next, we compare gVAMP to REGENIE, GMRM and LDAK-KVIK for association testing of the 13 traits in the UK Biobank imputed data. gVAMP performs similarly to mixed-linear model

Table 1. Polygenic risk score prediction accuracy R^2 for 13 different traits from statistical models trained in the UK Biobank data and tested in a UK Biobank hold-out set. Training data sample size and trait codes are given in Table S1 for each trait. The sample size of the hold-out test set is $15,000$ for all phenotypes. LDpred2 and SBayesR give estimates obtained from the LDpred2 and SBayesR software respectively, using summary statistic data of 8,430,446 SNPs obtained from the REGENIE software. LDAK gives the estimates obtained by the LDAK software using the elastic net predictor and individual-level data of 2,174,071 SNP markers. GMRM denotes estimates obtained from a Bayesian mixture model at 2,174,071 SNP markers ("GMRM 2M"). gVAMP denotes estimates obtained from the adaptive EM Bayesian mixture model within a vector approximate message passing (VAMP) framework presented in this work, using either 887,060 ("gVAMP 880k"), 2,174,071 ("gVAMP 2M"), or 8,430,446 SNP markers ("gVAMP 8M").

Phen	LDpred	SBayesR	LDAK 2.17M	GMRM 2.17M	gVAMP 880k	gVAMP 2.17M	gVAMP 8M
CHOL	0.147	0.149	0.152	**0.153**	0.140	0.152	**0.153**
EOSI	0.107	0.112	0.120	0.122	0.114	0.120	**0.124**
HbA1c	0.087	0.090	0.092	0.092	0.085	0.092	**0.095**
HDL	0.199	0.208	0.210	0.213	0.192	0.209	**0.219**
MCH	0.178	0.215	0.214	0.221	0.203	0.218	**0.223**
MCV	0.196	0.234	0.235	**0.244**	0.222	0.240	**0.244**
RBC	0.186	0.191	0.192	**0.199**	0.182	0.195	*0.198*
BMI	0.100	0.118	0.140	0.133	0.107	0.132	**0.141**
DBP	0.065	0.067	0.070	**0.071**	0.058	0.065	**0.071**
FVC	0.098	0.103	**0.112**	0.111	0.097	0.109	**0.112**
BMD	0.188	0.194	0.195	0.201	0.183	0.198	**0.204**
HT	0.231	0.362	0.449	0.450	0.419	0.449	**0.457**
SBP	0.068	0.071	0.072	**0.073**	0.061	0.072	**0.073**

association leave-one-chromosome-out (MLMA-LOCO) testing conducted using a predictor from GMRM, with the use of the full 8,430,446 imputed SNP markers improving performance (Table 2 and Figure 4b). REGENIE yields far fewer associations than either GMRM or gVAMP for all traits (Table 2 and Figure 4b), with LDAK-KVIK being equivalent to REGENIE. We additionally verify that placing the predictors obtained by gVAMP into the second step of the LDAK-KVIK algorithm, also yields improved performance as compared to the LDAK model (Table S4). Finally, using gVAMP to conduct variable selection, we find that hundreds of the LOCO marker associations for each trait can be localized (Table 2), with the obvious caveat that these results are for imputed SNP data and re-analysis of WGS data may yield different results as we show above for height. Using height as an example, we find that 3367 of the 5242 MLMA-LOCO associations can be localized directly to the 2070 imputed variant sets discovered by gVAMP via the PIP testing within 50kb. Thus, the gVAMP findings obtained by performing variable selection via the state evolution framework are often neighbored by multiple variants discovered by MLMA-LOCO in the imputed SNP data.

The increase in MLMA-LOCO findings for gVAMP over LDAK and REGENIE within the UK Biobank is consistent with the extensive simulation study results presented in Supplementary Note 1.

There, we show that MLMA-LOCO testing power scales with the out-of-sample prediction accuracy only for gVAMP and not for the other methods (Figure S13). The other conclusions obtained for the 13 UK Biobank traits are also corroborated by the simulation study of Supplementary Note 1. First, we show that gVAMP outperforms summary statistic [7, 8] and individual-level approaches [10] for polygenic risk score prediction across a range of settings (Figure S13). We then find that when the predictors obtained from gVAMP are used to conduct downstream tasks, such as MLMA-LOCO testing, they outperform both REGENIE [1] and LDAK-KVIK [10] (Figures S14 and S15). Contrary to previous work, we show the importance of including as large a number of predictors as possible when generating the polygenic risk scores used in the first stage of MLMA-LOCO for both the power and control of type-I error (Figures S14 and S15). Additional ablations varying the number of markers included in the model lead to similar findings (Figures S16, S17, S18 and S19). We also find that in practice our variational inference framework has similar performance to an MCMC approach, GMRM, that has a similar mixture prior [9] (Figure S20).

In terms of compute time and resource use, gVAMP completes in less than 1/2 of the time given the same data and compute resources as compared to REGENIE (Figure S21a), it is dramatically faster than GMRM (12.5× speed-up, Figure S20c), and it is marginally faster than LDAK when run on our simulated data (Figure S21a). Across the 13 traits, gVAMP is faster than LDAK-KVIK given the same SNPs and compute resources, but it requires more RAM (roughly the plink file size on disk, Figure S21b).

Discussion

We provide evidence that existing variable selection methods can fail to either select the correct genomic regions, or the appropriate genetic variants due to the complex and potentially long-distance correlation structure of whole genome sequence (WGS) data. As a result, applying fine-mapping methods one region at a time in isolation from the rest of the genome, may result in miscalibration and incorrect localisation of the underlying genetic effects. Motivated by this, we develop an adaptive form of Bayesian variable selection regression capable of fitting tens of millions of WGS variants jointly for hundreds of thousands of individuals. We conclusively demonstrate that gVAMP outperforms marginal testing plus fine-mapping approaches for localizing a signal to variants with the correct frequency and position in the DNA. We then apply the model to a set of 17M WGS variants, revealing a stronger relationship between effect size and allele frequency than previously considered, and localizing associations to numerous rare variants and gene burden scores as well as hundreds of regions containing common variation. To the best of our knowledge, this provides the highest-dimensional genomic analysis to date.

gVAMP provides a point estimate from the posterior distribution and, as such, its performance is expected to match that of standard MCMC Gibbs sampling algorithms. We show that this is the case in imputed SNP data: gVAMP and GMRM have comparable out-of-sample prediction accuracy, comparable estimates of the proportion of variance attributable to the DNA, and comparable number of MLMA-LOCO findings. However, unlike Gibbs sampling algorithms which cannot scale to current data sizes, gVAMP exhibits a remarkable speed-up which allows high-dimensional biobank data with full WGS observations to be analyzed. Taken together, gVAMP offers a flexible framework

Table 2. Genome-wide significant associations for 13 **UK Biobank traits from the software GMRM, gVAMP, LDAK-KVIK and REGENIE at 8,430,446 genetic variants from either mixed linear model association leave-one-chromosome-out (MLMA-LOCO) testing or gVAMP variable selection.** "REGENIE 880k LOCO" denotes results obtained from MLMA-LOCO testing using the REGENIE software, with 882,727 SNP markers used for step 1 and 8,430,446 markers used for the LOCO testing of step 2. "GMRM 2M LOCO" refers to MLMA-LOCO testing at 8,430,446 SNPs, using a Bayesian MCMC mixture model in step 1, with 2,174,071 SNP markers. "LDAK 2M LOCO" refers to MLMA-LOCO testing at 8,430,446 SNPs, using the LDAK-KVIK software in step 1, with 2,174,071 SNP markers. "gVAMP 2M LOCO" refers to MLMA-LOCO testing at 8,430,446 SNPs, using the framework presented here, where in step 1 2,174,071 markers are used and the second step identical to that of REGENIE. We then extend that to using all 8,430,446 SNP markers ("gVAMP 8M LOCO") in step 1. We also present gVAMP posterior inclusion probability (PIP) testing when applying a correlation threshold on the markers prior to analysis of $\rho = 0.9$ to the SNP markers (2,174,071 markers). For LOCO testing, the values give the number of genome-wide significant linkage disequilibrium independent associations selected based upon a standard p-value threshold of less than $5 \cdot 10^{-8}$ and R^2 between SNPs in a 1 Mb genomic segment of less than 1%. For the PIP testing, values give the number of genome-wide significant associations selected based upon a ≥ 0.95 threshold.

Trait	REGENIE 880k LOCO	GMRM 2M LOCO	LDAK 2M LOCO	gVAMP 2M LOCO	gVAMP 8M LOCO	gVAMP PIP
CHOL	571	627	482	603	**632**	381
EOSI	572	**693**	549	630	**693**	527
HbA1c	337	**429**	321	385	*413*	227
HDL	692	1009	639	812	**1092**	536
MCH	746	889	690	810	**916**	691
MCV	970	**1190**	903	1062	*1185*	898
RBC	897	1229	856	1079	**1325**	651
BMI	688	**1376**	726	1175	*1266*	261
DBP	291	425	303	419	**468**	315
FVC	549	**994**	577	721	*986*	318
BMD	522	**874**	549	668	*867*	491
HT	2712	5070	2703	4452	**5242**	2070
SBP	311	499	323	388	**529**	245

suitable to *(i)* conduct variable selection in WGS data (something that SuSiE, FINEMAP and their variants simply cannot do), and *(ii)* create polygenic risk scores with high accuracy that can also be used for downstream tasks such as MLMA-LOCO testing, with improved performance over LDAK-KVIK and REGENIE.

There are a number of remaining limitations and possible ways to expand and improve the analyses presented here. Our results suggest that the prioritization of relevant regions, genes and gene sets is feasible at current sample sizes, but the detection of specific height-associated mutations is expected to be a challenging task. There are a very large number of rare variants within the human population that are missing from our analysis of 16,854,878 WGS variants. While conducting variable selection genome-wide for these will require millions of WGS samples, it may be beneficial to include as many of these rare variants as possible. Our algorithm requires random-access memory (RAM) that is the same as the plink data file on disk, and although we evidence that this is not prohibitive to running the algorithm on the UK Biobank cloud computing, two alternative

solutions could be to: *(i)* apply gVAMP region-by-region, or *(ii)* make slight modifications in the implementation, so that gVAMP streams data, at the cost of slightly increased run time. Our ongoing work focuses on exploring these options.

A further limitation is that we have exclusively tested and analyzed individuals of European ancestry. Combining data of different ancestries, or utilising admixed populations is of great importance [31], and previous work suggests that better modeling within a single large biobank can facilitate improved association testing in other global biobanks [32]. However, tackling this problem requires the development of a modeling framework that explicitly allows for differences in MAF, LD and effect size across the human population, which we leave to future work.

Additionally, we also removed first-degree relatives from all of our analyses to avoid confounding of shared environmental effects. While some relatedness still exists even after removing first-degree relatives, we show in simulations that gVAMP performs well in both WGS and imputed data when planting a signal on real observed genotype data. Our motivation is that genotype-phenotype data of close relatives is often utilized for follow-up studies and there are unique models (to which our framework could be adapted) employed for such tasks. Additionally, removing first-degree relatives loses only a few thousand individuals from the UK Biobank, which does not alter the sample size in a significant way. We note that methods such as fastGWA [4] or the equivalent implementation in LDAK, could be used where a first step calculates the parameters needed to control for environmental confounding before the model is then run.

Finally, we also now seek to extend the model presented here to different outcome distributions (binary outcomes, time-to-event, count data, etc.), to have a grouped prior for annotations, to model multiple outcomes jointly, and to do all of this using summary statistic as well as individual-level data across different biobanks. Progress in these areas has been made in the AMP literature, see e.g. [33, 34]. Combining data sets together in a principled way without a loss of accuracy, whilst preserving privacy, is key to obtaining the sample sizes that are likely required to fully explore the genetic basis of complex traits.

In summary, gVAMP is a different way to create genetic predictors and to conduct variable selection. With increasing sample sizes reducing standard errors, a vast number of genomic regions are being identified as significantly associated with trait outcomes by one-SNP-at-a-time association testing. Such large numbers of findings will make it increasingly difficult to determine the relative importance of a given mutation, especially in whole genome sequence data with dense, highly correlated variants. Thus, it is crucial to develop statistical approaches fitting all variants jointly and asking whether, given the LD structure of the data, there is evidence for an effect at each locus.

Methods

gVAMP algorithm

Approximate message passing (AMP) was originally proposed for linear regression [13, 16, 35] assuming a Gaussian design matrix $\boldsymbol{X}$. To accommodate a wider class of structured design matrices, vector approximate message passing (VAMP) was introduced in [14]. The performance of VAMP can be precisely characterized via a deterministic, low-dimensional *state evolution* recursion, for any right-orthogonally invariant design matrix. We recall that a matrix is right-orthogonally invariant

---

Algorithm 1 gVAMP

1: **Input:** preprocessed normalized genotype matrix $\boldsymbol{X} \in \mathbb{R}^{N \times P}$, max number of iterations N_{it}, initial estimate of effect sizes $\boldsymbol{r}_{1,0} = \boldsymbol{0}_P \in \mathbb{R}^P$, initial estimate of effect sizes precision $\gamma_{1,0} = 10^{-6} > 0$, initial estimate of noise precision $\gamma_{\epsilon,0} = 2$, initial set of parameters defining the prior distribution $\Theta_0 = \{\lambda, (\pi_i^{(0)})_{i=1}^L, (\sigma_i^{(0)})_{i=1}^L\}$, max number of variance auto-tuning steps $N_{\text{var\_tune}} = 5 \in \mathbb{N}$, threshold for stopping criterion $\varepsilon = 10^{-4} > 0$, damping factor $\rho \in (0,1)$.

2: **for** $t = 0, 1, \ldots, N_{\text{it}}$ **do**

3:     **Denoising step**

4:     **for** $k = 0, 1, \ldots, N_{\text{var\_tune}}$ **do**

5:         $\hat{\boldsymbol{\beta}}_{1,t} = \mathbb{E}[\boldsymbol{\beta} | \boldsymbol{r}_{1,t} = \boldsymbol{\beta} + \mathcal{N}(0, \gamma_{1,t}^{-1}\boldsymbol{I}), \gamma_{1,t}, \Theta_t]$

6:         **if** $t > 0$ **then**

7:             Variance auto-tuning step of estimation error for $\boldsymbol{\beta}$ in the denoising step, called $\gamma_{1,t}$

8:             EM update of the prior distribution parameters Θ, called Θ_t

9:             **if** $|\gamma_{1,t} - \gamma_{1,t}^{(\text{previous})}| < 10^{-3}$ **then**

10:                **break**

11:             **end if**

12:         **end if**

13:     **end for**

14:     **if** $t \geq 0$ **then**

15:         $\hat{\boldsymbol{\beta}}_{1,t} = \rho \cdot \hat{\boldsymbol{\beta}}_{1,t} + (1 - \rho) \cdot \hat{\boldsymbol{\beta}}_{1,t-1}$

16:     **end if**

17:     $\alpha_{1,t} = \gamma_{1,t} \cdot \langle \text{Var}[\boldsymbol{\beta} | \boldsymbol{r}_{1,t} = \boldsymbol{\beta} + \mathcal{N}(0, \gamma_{1,t}^{-1}\boldsymbol{I}), \gamma_{1,t}, \Theta_t] \rangle$

18:     $\gamma_{2,t} = \gamma_{1,t} \cdot (1 - \alpha_{1,t})/\alpha_{1,t}$

19:     $\boldsymbol{r}_{2,t} = (\hat{\boldsymbol{\beta}}_{1,t} - \alpha_{1,t}\boldsymbol{r}_{1,t})/(1 - \alpha_{1,t})$

20:     **LMMSE step**

21:     $\hat{\boldsymbol{\beta}}_{2,t} = (\gamma_{\epsilon,t}\boldsymbol{X}^T\boldsymbol{X} + \gamma_{2,t}\boldsymbol{I})^{-1}(\gamma_{\epsilon,t}\boldsymbol{X}^T\boldsymbol{y} + \gamma_{2,t}\boldsymbol{r}_{2,t})$

22:     $\alpha_{2,t} = \gamma_{2,t} \cdot \text{Tr}[(\gamma_{\epsilon,t}\boldsymbol{X}^T\boldsymbol{X} + \gamma_{2,t}\boldsymbol{I})^{-1}]/P$

23:     $\gamma_{1,t+1} = \gamma_{2,t} \cdot (1 - \alpha_{2,t})/\alpha_{2,t}$

24:     **if** $t > 1$ **then**

25:         Variance auto-tuning step of estimation error for $\boldsymbol{\beta}$ in the LMMSE step, called $\gamma_{2,t}$

26:     **end if**

27:     $\boldsymbol{r}_{1,t+1} = (\hat{\boldsymbol{\beta}}_{2,t} - \alpha_{2,t}\boldsymbol{r}_{2,t})/(1 - \alpha_{2,t})$

28:     EM update of the estimate of γ_ϵ, called $\gamma_{\epsilon,t}$

29:     **if** $t \geq 1$ and $||\hat{\boldsymbol{\beta}}_{1,t} - \hat{\boldsymbol{\beta}}_{1,t-1}||_2/||\hat{\boldsymbol{\beta}}_{1,t-1}||_2 < \varepsilon$ **then**

30:         **break**

31:     **end if**

32: **end for**

33: **return** $\hat{\boldsymbol{\beta}}_{1,t}$

---

if its right singular vectors are distributed according to the Haar measure, i.e., they are uniform in the group of orthogonal matrices. Empirical evidence [36,37] suggests that the predictions for rotationally invariant models hold for a wide class of scenarios, and a line of work has aimed at showing that this is case for AMP algorithms [38,39].

gVAMP extends EM-VAMP, introduced in [14,24,25], in which the prior parameters are adaptively learnt from the data via EM, and it is an iterative procedure consisting of two steps: *(i)* denoising, and *(ii)* linear minimum mean square error estimation (LMMSE). The denoising step accounts for

the prior structure given a noisy estimate of the signal $\boldsymbol{\beta}$, while the LMMSE step utilizes phenotype values to further refine the estimate by accounting for the LD structure of the data.

A key feature of the algorithm is the so called *Onsager correction*: this is added to ensure the asymptotic normality of the noise corrupting the estimates of $\boldsymbol{\beta}$ at every iteration. Here, in contrast to MCMC or other iterative approaches, the normality is guaranteed under mild assumptions on the normalized genotype matrix. This property allows a precise performance analysis via state evolution and, consequently, the optimization of the method.

In particular, the quantity $\gamma_{1,t}$ in line 7 of Algorithm 1 is the state evolution parameter tracking the error incurred by $\boldsymbol{r}_{1,t}$ in estimating $\boldsymbol{\beta}$ at iteration t. The state evolution result gives that $\boldsymbol{r}_{1,t}$ is asymptotically Gaussian, i.e., for sufficiently large N and P, $\boldsymbol{r}_{1,t}$ is approximately distributed as $\mathcal{N}(\boldsymbol{\beta}, \gamma_{1,t}^{-1}\boldsymbol{I})$. Here, $\boldsymbol{\beta}$ represents the signal to be estimated, with the prior learned via EM steps at iteration t:

$$\beta_i \sim (1 - \lambda_t) \cdot \delta_0(\cdot) + \lambda_t \cdot \sum_{l=1}^{L} \pi_{t,l} \cdot \mathcal{N}(\cdot, 0, \sigma_{t,l}^2), \quad \forall\, i = 1, \ldots, P.$$

Compared to Equation (2), the subscript t in $\lambda_t, \pi_{t,l}, \sigma_{t,l}$ indicates that these parameters change through iterations, as they are adaptively learned by the algorithm. Similarly, $\boldsymbol{r}_{2,t}$ is approximately distributed as $\mathcal{N}(\boldsymbol{\beta}, \gamma_{2,t}^{-1}\boldsymbol{I})$. The Gaussianity of $\boldsymbol{r}_{1,t}, \boldsymbol{r}_{2,t}$ is enforced by the presence of the Onsager coefficients $\alpha_{1,t}$ and $\alpha_{2,t}$, see lines 17 and 22 of Algorithm 1, respectively. We also note that $\alpha_{1,t}$ (resp. $\alpha_{2,t}$) is the state evolution parameter linked to the error incurred by $\hat{\boldsymbol{\beta}}_{1,t}$ (resp. $\hat{\boldsymbol{\beta}}_{2,t}$).

The vectors $\boldsymbol{r}_{1,t}, \boldsymbol{r}_{2,t}$ are obtained after the LMMSE step, and they are further improved via the denoising step, which respectively gives $\hat{\boldsymbol{\beta}}_{1,t}, \hat{\boldsymbol{\beta}}_{2,t}$. In the denoising step, we exploit our estimate of the approximated posterior by computing the conditional expectation of $\boldsymbol{\beta}$ with respect to $\boldsymbol{r}_{1,t}, \boldsymbol{r}_{2,t}$ in order to minimize the mean square error of the estimated effects. For example, let us focus on the pair $(\boldsymbol{r}_{1,t}, \hat{\boldsymbol{\beta}}_{1,t})$ (analogous considerations hold for $(\boldsymbol{r}_{2,t}, \hat{\boldsymbol{\beta}}_{2,t})$). Then, we have that

$$\hat{\boldsymbol{\beta}}_{1,t} = f_t(\boldsymbol{r}_{1,t}) = \mathbb{E}[\boldsymbol{\beta} | \boldsymbol{r}_{1,t} = \boldsymbol{\beta} + \mathcal{N}(0, \gamma_{1,t}^{-1}\boldsymbol{I}), \lambda_t, \{\pi_{t,l}\}_{l=1}^{L}, \{\sigma_{t,l}^2\}_{l=1}^{L}]. \tag{3}$$

Here, $f_t : \mathbb{R} \rightarrow \mathbb{R}$ denotes the denoiser at iteration t and the notation $f_t(\boldsymbol{r}_{1,t})$ assumes that the denoiser f_t is applied component-wise to elements of $\boldsymbol{r}_{1,t}$. Note that, in line 15 of Algorithm 1, we take this approach one step further by performing an additional step of damping, see "Algorithm stability" below.

From Bayes theorem, one can calculate the posterior distribution (which here has the form of a spike-and-slab mixture of Gaussians) and obtain its expectation. Hence, by denoting a generic component of $\boldsymbol{r}_{1,t}$ as r_1, it follows that

$$
\begin{aligned}
f_t(r_1) &= \frac{\lambda_t \cdot \sum_{l=1}^{L} \pi_{t,l} \cdot \frac{r_1 \cdot \sigma_{t,l}^2}{\gamma_{1,t}^{-1} + \sigma_{t,l}^2} \cdot \mathcal{N}(r_1; 0, \gamma_{1,t}^{-1} + \sigma_{t,l}^2)}{(1 - \lambda_t) \cdot \mathcal{N}(r_1; 0, \gamma_{1,t}^{-1}) + \lambda_t \sum_{l=1}^{L} \pi_{t,l} \cdot \mathcal{N}(r_1; 0, \gamma_{1,t}^{-1} + \sigma_{t,l}^2)} \\[2mm]
&= \frac{\lambda_t \cdot \sum_{l=1}^{L} \pi_{t,l} \cdot \frac{r_1 \cdot \sigma_{t,l}^2}{(\gamma_{1,t}^{-1} + \sigma_{t,l}^2)^{3/2}} \cdot \mathrm{EXP}(\sigma_{t,l}^2)}{(1 - \lambda_t) \cdot \gamma_1^{1/2} \cdot \mathrm{EXP}(0) + \lambda_t \cdot \sum_{l=1}^{L} \pi_{t,l} \cdot \frac{1}{(\gamma_{1,t}^{-1} + \sigma_{t,l}^2)^{1/2}} \cdot \mathrm{EXP}(\sigma_{t,l}^2)},
\end{aligned}
\tag{4}
$$

where $\mathcal{N}(r_1; 0, \gamma_{1,t}^{-1} + \sigma_{t,l}^2)$ denotes the probability density function of a Gaussian with mean 0 and variance $\gamma_{1,t}^{-1} + \sigma_{t,l}^2$ evaluated at r_1. Furthermore, we set

$$\mathrm{EXP}(\sigma^2) = \exp\left(-\frac{r_1^2}{2} \cdot \frac{\sigma_{t,*}^2 - \sigma^2}{(\gamma_{1,t}^{-1} + \sigma^2)(\gamma_{1,t}^{-1} + \sigma_{t,*}^2)} \right),$$

with $\sigma_{t,*}^2 := \max_l(\sigma_{t,l}^2)$. This form of the denoiser is particularly convenient, as we typically deal with very sparse distributions when estimating genetic associations. We also note that the calculation of the Onsager coefficient in line 17 of Algorithm 1 requires the evaluation of a conditional variance, which is computed as the ratio of the derivative of the denoiser over the error in the estimation of the signal, i.e.,

$$\mathrm{Var}[\beta_i | (\boldsymbol{r}_{1,t})_i = \beta_i + \mathcal{N}(0, \gamma_{1,t}^{-1} \boldsymbol{I}), \lambda_t, \{\pi_{t,l}\}_{l=1}^L, \{\sigma_{t,l}^2\}_{l=1}^L] = f_t'((\boldsymbol{r}_{1,t})_i)/\gamma_{1,t}. \tag{5}$$

The calculation of the derivative of f_t is detailed in Supplementary Note 2.

If one has access to the singular value decomposition (SVD) of the data matrix $\boldsymbol{X}$, the per-iteration complexity is of order $\mathcal{O}(NP)$. However, at biobank scales, performing the SVD is computationally infeasible. Thus, the linear system $(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I})^{-1}(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{y} + \gamma_{2,t} \boldsymbol{r}_{2,t})$ (see line 21 of Algorithm 1) needs to be solved using an iterative method, in contrast to having an analytic solution in terms of the elements of the singular value decomposition of $\boldsymbol{X}$. In the next section, we provide details on how we overcome this issue.

Scaling up using warm-start conjugate gradients

We approximate the solution of the linear system $(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I})^{-1}(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{y} + \gamma_{2,t} r_{2,t})$ with a symmetric and positive-definite matrix via the *conjugate gradient method* (CG), see Algorithm 2 in Supplementary Note 2, which is included for completeness. If κ is the condition number of $\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I}$, the method requires $\mathcal{O}(\sqrt{\kappa})$ iterations to return a reliable approximation.

Additionally, inspired by [40], we initialize the CG iteration with an estimate of the signal from the previous iteration of gVAMP. This warm-starting technique leads to a reduced number of CG steps that need to be performed and, therefore, to a computational speed-up. However, this comes at the expense of potentially introducing spurious correlations between the signal estimate and the Gaussian error from the state evolution. Such spurious correlations may lead to algorithm instability when run for a large number of iterations (also extensively discussed below). This effect is prevented by simply stopping the algorithm as soon as the R^2 measure on the training data starts decreasing.

In order to calculate the Onsager correction in the LMMSE step of gVAMP (see line 22 of Algorithm 1), we use the Hutchinson estimator [41] to estimate the quantity $\mathrm{Tr}[(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I})^{-1}]/P$. We recall that this estimator is unbiased, in the sense that, if $\boldsymbol{u}$ has i.i.d. entries equal to -1 and $+1$ with the same probability, then

$$\mathbb{E}[\boldsymbol{u}^T(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I})^{-1} \boldsymbol{u}/P] = \mathrm{Tr}[(\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I})^{-1}]/P.$$

Furthermore, in order to perform an EM update for the noise precision γ_ϵ one has to calculate the trace of a matrix which is closely connected to the one we have seen in the previous paragraph.

In order to do so efficiently, i.e., to avoiding solving another large-dimensional linear system, we store the inverted vector $(\gamma_{\epsilon,t}\boldsymbol{X}^T\boldsymbol{X} + \gamma_{2,t}\boldsymbol{I})^{-1}\boldsymbol{u}$ and reuse it again in the EM update step (see the subparagraph on EM updates).

Algorithm stability

We find that the application of existing EM-VAMP algorithms to the UK Biobank dataset leads to diverging estimates of the signal. This is due to the fact that the data matrix (the SNP data) might not conform to the properties required in [14], especially that of right-rotational invariance. Furthermore, incorrect estimation of the noise precision in line 28 of Algorithm 1 may also lead to instability of the algorithm, as previous applications of EM-VAMP generally do not leave many hyperparameters to estimate.

To mitigate these issues, different approaches have been proposed including row or/and column normalization, damping (i.e., doing convex combinations of new and previous estimates) [42], and variance auto-tuning [25]. In particular, to prevent EM-VAMP from diverging and ensure it follows its state evolution, we empirically observe that the combination of the following techniques is crucial.

1. We perform *damping* in the space of denoised signals. Thus, line 15 of Algorithm 1 reads as

$$\hat{\boldsymbol{\beta}}_{1,t} = \rho \cdot \mathbb{E}[\boldsymbol{\beta}|\boldsymbol{r}_{1,t}, \Theta_t] + (1-\rho) \cdot \hat{\boldsymbol{\beta}}_{1,t-1},$$

    in place of $\hat{\boldsymbol{\beta}}_{1,t} = \mathbb{E}[\boldsymbol{\beta}|\boldsymbol{r}_{1,t}, \Theta_t]$. Here, $\rho \in (0,1)$ denotes the damping factor. This ensures that the algorithm is making smaller steps when updating a signal estimate.

2. We perform *auto-tuning* of $\gamma_{1,t}$ via the approach from [25]. Namely, in the auto-tuning step, one refines the estimate of $\gamma_{1,t}$ and the prior distribution of the effect size vector $\boldsymbol{\beta}$ by jointly re-estimating them. If we denote the previous estimates of $\gamma_{1,t}$ and Θ with $\gamma_{1,t}^{(\text{previous})}$ and $\Theta^{(\text{previous})}$, then this is achieved by setting up an expectation-maximization procedure whose aim is to maximize

$$\mathbb{E}\big[\log p(\boldsymbol{\beta}, \boldsymbol{r}_{1,t}|\gamma_{1,t}, \Theta)|\boldsymbol{r}_{1,t}, \gamma_{1,t}^{(\text{previous})}, \Theta^{(\text{previous})}\big]$$

    with respect to $\gamma_{1,t}$ and Θ.

3. We *filter* the design matrix for first-degree relatives to reduce the correlation between rows, which has the additional advantage of avoiding potential confounding of shared-environmental effects among relatives.

Estimation of the prior and noise precision via EM

The VAMP approach in [14] assumes exact knowledge of the prior on the signal $\boldsymbol{\beta}$, which deviates from the setting in which genome-wide association studies are performed. Hence, we adaptively learn the signal prior from the data using expectation-maximization (EM) steps, see lines 8 and 28 of Algorithm 1. This leverages the variational characterization of EM-VAMP [24], and its rigorous theoretical analysis presented in [25]. In this subsection, we summarize the hyperparameter

estimation results derived based upon [43] in the context of our model. We find that the final
update formulas for our hyperparameter estimates are as follows.

- Sparsity rate λ: We define $\{\zeta_j\}_{j=1}^P$ as:

$$\zeta_j := \frac{\lambda_t \cdot \sum_{i=1}^L \pi_{i,t} \cdot \mathcal{N}((\boldsymbol{r}_{1,t})_j; 0, \sigma_{i,t}^2 + \gamma_{1,t}^{-1})}{\lambda_t \cdot \sum_{i=1}^L \pi_{i,t} \cdot \mathcal{N}((\boldsymbol{r}_{1,t})_j; 0, \sigma_{i,t}^2 + \gamma_{1,t}^{-1}) + (1 - \lambda_t) \cdot \mathcal{N}((\boldsymbol{r}_{1,t})_j; 0, \gamma_{1,t}^{-1})}, \quad \forall j = 1, \ldots, P.$$

(6)

The intuition behind $\{\zeta_j\}_{j=1}^P$ is that each ζ_j tells what fraction of posterior probability mass
was assigned to the event that it has a non-zero effect. Then, the update formula for the
sparsity rate λ_{t+1} reads as

$$\lambda_{t+1} = \frac{1}{P} \sum_{j=1}^P \zeta_j.$$

- Probabilities of mixture components in the slab part $\{\pi_i\}_{i=1}^L$: We define $\{\xi_{j,i}\}_{i=1,j=1}^{L,P}$ as

$$\xi_{j,i} = \frac{\pi_{i,t} \mathcal{N}((\boldsymbol{r}_{1,t})_j; 0, \sigma_i^2 + \gamma_{1,t}^{-1})}{\sum_{l=1}^L \pi_{l,t} \cdot \mathcal{N}((\boldsymbol{r}_{1,t})_j; 0, \sigma_l^2 + \gamma_{1,t}^{-1})}, \quad \forall i = 1, \ldots, L, \ \forall j = 1, \ldots, P.$$

The intuition behind $\{\xi_{j,i}\}_{i=1,j=1}^{L,P}$ is that each $\xi_{j,i}$ approximates the posterior probability that
a marker j belongs to a mixture i conditional on the fact that it has non-zero effect. Thus,
the update formula for $\pi_{i,t+1}$ reads as

$$\pi_{i,t+1} = \frac{\sum_{j=1}^P \zeta_j \xi_{j,i}}{\sum_{j=1}^P \zeta_j}, \quad \forall i = 1, \ldots, L.$$

- Variances of mixture components in the slab part $\{\sigma_i^2\}_{i=1}^L$: Using the same notation, the up-
date formula reads as

$$\sigma_{i,t+1}^2 = \frac{\sum_{j=1}^P \zeta_j \cdot \xi_{j,i} \cdot \left[\left(\frac{(\boldsymbol{r}_{1,t})_j \cdot \gamma_{1,t}}{\gamma_{1,t} + \sigma_{i,t}^{-2}} \right)^2 + \frac{1}{\gamma_{1,t} + \sigma_{i,t}^{-2}} \right]}{\sum_{j=1}^P \zeta_j \cdot \xi_{j,i}}, \quad \forall i = 1, \ldots, L.$$

Here we also introduce a mixture merging step, i.e., if the two mixtures are represented by
variances that are close to each other in relative terms, then we merge those mixtures together.
Thus, we adaptively learn the mixture number.

- Precision of the error γ_ϵ : We define $\boldsymbol{\Sigma}_t := (\gamma_{\epsilon,t} \boldsymbol{X}^T \boldsymbol{X} + \gamma_{2,t} \boldsymbol{I})^{-1}$. Then, the update formula
for the estimator of γ_ϵ reads as

$$\gamma_{\epsilon,t+1} = \frac{1}{\frac{||\boldsymbol{y} - \boldsymbol{X}\hat{\beta}_{2,t}||^2}{N} + \frac{\text{Tr}(\boldsymbol{X}\boldsymbol{\Sigma}_t \boldsymbol{X}^T)}{N}}.$$

In the formula above, the term $||\boldsymbol{y} - \boldsymbol{X}\hat{\beta}_{2,t}||^2/N$ takes into account the quality of the fit

of the model, while the term $\text{Tr}(\boldsymbol{X}\boldsymbol{\Sigma}_t\boldsymbol{X}^T)/N$ prevents overfitting by accounting for the structure of the prior distribution of the effect sizes via the regularization term $\gamma_{2,t}$. We note that the naive evaluation of this term would require an inversion of a matrix of size $P \times P$. We again use the Hutchinson estimator for the trace to approximate this object, i.e., $\text{Tr}(\boldsymbol{X}\boldsymbol{\Sigma}_t\boldsymbol{X}^T) = \text{Tr}(\boldsymbol{X}^T\boldsymbol{X}\boldsymbol{\Sigma}_t) \approx \boldsymbol{u}^T(\boldsymbol{X}^T\boldsymbol{X}\boldsymbol{\Sigma}_t)\boldsymbol{u}$, where $\boldsymbol{u}$ has i.i.d. entries equal to -1 and $+1$ with the same probability. Furthermore, instead of solving a linear system $\boldsymbol{\Sigma}_t\boldsymbol{u}$ with a newly generated $\boldsymbol{u}$, we re-use the $\boldsymbol{u}$ sampled when constructing the Onsager coefficient, thus saving the time needed to construct the object $\boldsymbol{\Sigma}_t\boldsymbol{u}$.

C++ code optimization

Our open-source gVAMP software (`https://github.com/medical-genomics-group/gVAMP`) is implemented in C++, and it incorporates parallelization using the OpenMP and MPI libraries. MPI parallelization is implemented in a way that the columns of the normalized genotype matrix are approximately equally split between the workers. OpenMP parallelization is done on top of that and used to further boost performance within each worker by simultaneously performing operations such as summations within matrix vector product calculations. Moreover, data streaming is employed using a lookup table, enabling byte-by-byte processing of the genotype matrix stored in PLINK format with entries encoded to a set $\{0, 1, 2\}$:

$$\left(\boxed{0 \;\; 1}\;\boxed{0 \;\; 0}\;\boxed{1 \;\; 1}\;\boxed{1 \;\; 0}\right) \mapsto \left(\boxed{\text{NaN}}\boxed{2}\boxed{0}\boxed{1}\right)$$

The lookup table enables streaming in the data in bytes, where every byte (8 bits) encodes the information of 4 individuals. This reduces the amount of memory needed to load the genotype matrix. In addition, given a suitable computer architecture, our implementation supports SIMD instructions which allow handling four consecutive entries of the genotype matrix simultaneously. To make the comparisons between different methods fair, the results presented in the paper do not assume usage of SIMD instructions. Additionally, we emphasize that all calculations take un-standardized values of the genotype matrix in the form of standard PLINK binary files, but are conducted in a manner that yields the parameter estimates one would obtain if each column of the genotype matrix was standardized.

UK Biobank data

Participant inclusion

UK Biobank has approval from the North-West Multicenter Research Ethics Committee (MREC) to obtain and disseminate data and samples from the participants (`https://www.ukbiobank.ac.uk/ethics/`), and these ethical regulations cover the work in this study. Written informed consent was obtained from all participants.

Our objective is to use the UK Biobank to provide proof of principle of our approach and to compare to state-of-the-art methods in applications to biobank data. We first restrict our analysis to a sample of European-ancestry UK Biobank individuals to provide a large sample size and more uniform genetic background with which to compare methods. To infer ancestry, we use both self-reported ethnic background (UK Biobank field 21000-0), selecting coding 1, and genetic ethnicity

(UK Biobank field 22006-0), selecting coding 1. We project the 488,377 genotyped participants 582 onto the first two genotypic principal components (PC) calculated from 2,504 individuals of the 583 1,000 Genomes project. Using the obtained PC loadings, we then assign each participant to the 584 closest 1,000 Genomes project population, selecting individuals with PC1 projection $\leq$ absolute 585 value 4 and PC2 projection $\leq$ absolute value 3. We apply this ancestry restriction as we wish to 586 provide the first application of our approach, and to replicate our results, within a sample that 587 is as genetically homogeneous as possible. Our approach can be applied within different human 588 groups (by age, genetic sex, ethnicity, etc.). However, combining inference across different human 589 groups requires a model that is capable of accounting for differences in minor allele frequency and 590 linkage disequilibrium patterns across human populations. Here, the focus is to first demonstrate 591 that our approach provides an optimal choice for biobank analyses, and ongoing work focuses on 592 exploring differences in inference across a diverse range of human populations. Secondly, samples 593 are also excluded based on UK Biobank quality control procedures with individuals removed of *(i)* 594 extreme heterozygosity and missing genotype outliers; *(ii)* a genetically inferred gender that did 595 not match the self-reported gender; *(iii)* putative sex chromosome aneuploidy; *(iv)* exclusion from 596 kinship inference; *(v)* withdrawn consent. 597

Whole genome sequence data 598

We process the population-level WGS variants, recently released on the UK Biobank DNAnexus 599 platform. We use BCF tools to process thousands of pVCF files storing the chunks of DNA sequences, 600 applying elementary filters on genotype quality (GQ $\leq$ 10), local allele depth (smpl_sum LAD < 8), 601 missing genotype (F_MISSING > 0.1), and minor allele frequency (MAF < 0.0001). We select this 602 MAF threshold as it means that on average about 80 people will have a genotype that is non-zero, 603 which was the lowest frequency for which we felt that there was adequate power in the data to 604 detect the variants. 605

Simultaneously, we normalize the indels to the most recent reference, removing redundant data 606 fields to reduce the size of the files. For all chromosomes separately, we then concatenate all 607 the pre-processed VCF files and convert them into PLINK format. The compute nodes on the 608 DNAnexus system are quite RAM limited, and it is not possible to analyse the WGS data outside 609 of this system, which restricts the number of variants that can be analysed jointly. To reduce the 610 number of variants to the scale which can be fit in the largest computational instance available on 611 the DNAnexus platform, we rank variants by minor allele frequency and remove the variants in 612 high LD with the most common variants using the PLINK clumping approach, setting a 1000 kb 613 radius, and R^2 threshold to 0.36. This selects a focal common variant from a group of other common 614 variants with correlation ≥ 0.6, which serves to capture the common variant signal into groups, 615 whilst keeping all rare variation within the data. Finally, we remove the variants sharing the same 616 base pair position, not keeping any of these duplicates, and merge all the chromosomes into a large 617 data instance, including the final 16,854,878 WGS variants. 618

Imputed SNP data

We use genotype probabilities from version 3 of the imputed autosomal genotype data provided by the UK Biobank to hard-call the single nucleotide polymorphism (SNP) genotypes for variants with an imputation quality score above 0.3. The hard-call-threshold is 0.1, setting the genotypes with probability ≤ 0.9 as missing. From the good quality markers (with missingness less than 5% and p-value for the Hardy-Weinberg test larger than 10^{-6}, as determined in the set of unrelated Europeans) we select those with rs identifier, in the set of European-ancestry participants, providing a dataset of 23,609,048 SNPs. For comparison in the WGS simulation, we select chromosome 12 and process the data in the same way as for the WGS data above, selecting MAF ≥ 0.0001 and selecting one marker from a set of variants with LD $R^2 \geq 0.6$.

We further subset the 23,609,048 SNP data by selecting markers with MAF ≥ 0.002 to give 8,430,446 autosomal markers. We then select an additional two subsets of these data by selecting one marker from a set of variants with LD $R^2 \geq 0.8$ within a 1MB window to obtain 2,174,071 markers, and selecting with LD $R^2 \geq 0.5$ to obtain 882,727 SNP markers. This results in the selection of two subsets of "tagging variants", with only variants in very high LD with the tag SNPs removed. This allows us to compare analysis methods that are restricted in the number of SNPs that can be analyzed, but still provide them a set of markers that are all correlated with the full set of imputed SNP variants, limiting the loss of association power by ensuring that the subset is correlated to those SNPs that are removed.

Whole exome sequence data burden scores

We then combine this data with the UK Biobank whole exome sequence data. The UK Biobank final release dataset of population level exome variant calls files is used (`https://doi.org/10.1101/572347`). Genomic data preparation and aggregation is conducted with custom pipeline (repo) on the UK Biobank Research Analysis Platform (RAP) with DXJupyterLab Spark Cluster App (v. 2.1.1). Only biallelic sites and high quality variants are retained according to the following criteria: individual and variant missingness $< 10\%$, Hardy-Weinberg Equilibrium p-value $> 10^{-15}$, minimum read coverage depth of 7, at least one sample per site passing the allele balance threshold > 0.15. Genomic variants in canonical, protein coding transcripts (Ensembl VERSION) are annotated with the Ensembl Variant Effect Predictor (VEP) tool (docker image ensemblorg/ensembl-vep:release_110.1). High-confidence (HC) loss-of-function (LoF) variants are identified with the LOFTEE plugin (v1.0.4_GRCh38). For each gene, homozygous or multiple heterozygous individuals for LoF variants have received a score of 2, those with a single heterozygous LoF variant 1, and the rest 0. We chose to use the WES data to create the burden scores rather than the WGS data as existing well-tested pipelines were available.

Phenotypic records

Finally, we link these DNA data to the measurements, tests, and electronic health record data available in the UK Biobank [44] and, for the imputed SNP data, we select 7 blood based biomarkers and 6 quantitative measures which show $\geq 15\%$ SNP heritability and $\geq 5\%$ out-of-sample prediction accuracy [9]. Our focus is on selecting a group of phenotypes for which there is sufficient power to

observe differences among approaches. We split the sample into training and testing sets for each phenotype, selecting 15,000 individuals that are unrelated (SNP marker relatedness < 0.05) to the training individuals to use as a testing set. This provides an independent sample of data with which to access prediction accuracy. We restrict our prediction analyses to this subset as predicting across other biobank data introduces issues of phenotypic concordance, minor allele frequency and linkage disequilibrium differences. In fact, our objective is to simply benchmark methods on as uniform a dataset as we can. As stated, combining inference across different human groups, requires a model that is capable of accounting for differences in minor allele frequency and linkage disequilibrium patterns across human populations and, while our algorithmic framework can provide the basis of new methods for this problem, the focus here is on benchmarking in the simpler linear model setting. Samples sizes and traits used in our analyses are given in Table S2. Prior to analysis all phenotypic values were adjusted by the participant's recruitment age, their genetic sex, north-south and east-west location coordinates, measurement center, genotype batch, and the leading 20 genetic PCs.

Statistical analysis in the UK Biobank

Simulation study for WGS data

To demonstrate the usefulness of gVAMP as a signal localization tool, we investigate a series of 12 simulated scenarios in which the signal is planted on top of raw WGS chromosome 12 genotype data, which consists of $1,922,505$ variants. The data is further split into train set with $411,536$ individuals present and test set which contains $14,926$ individuals. As in the real data analysis, we rank variants by minor allele frequency and remove the variants in high LD with the most common variants using the PLINK clumping approach, setting a 1000kb radius, and R^2 threshold to 0.36 to obtain a subset of $800,674$ variants belonging to chromosome 12 which we further use for benchmarking.

Finally, the signal itself is simulated from a Bernoulli-Gaussian prior in which the variance of the slab part is determined as the ratio between desired heritability and the expected number of causal variants. We examine the effect that simulating a non-zero α-power law has on the inference process and on the quality of estimation of α (see Supplementary Note 3 for more details on the simulation of effects under α-power law and estimation pipeline). In the simulation scenarios labeled as rare, we only allow variants to be causal if their MAF belongs to an interval $[0.001, 0.01]$. The simulation details for each of the 12 scenarios we considered are reported in Table S1. When simulating effect sizes we group variants into windows of size $50,000$ bp and simulate exactly one causal variant in a number of consecutive valid windows at the beginning of the chromosome. A window is not valid if there is at most one variant in a clumped dataset belonging to it, as the benchmark methods cannot run on singletons. All simulated traits are PCA corrected for the first 20 principal components following standard practice.

We run the SuSiE method either separately for each window or by using a 10 times larger window size, comparing the difference. For each window block, we first load the data using the *plinkFile* library, standardize and fill the NA values by means, and finally run SuSiE using the *susieR* library. SuSiE-inf, FINEMAP and FINEMAP-inf are ran over windows using window-specific

summary statistics: marginal effects and LD matrices generated by PLINK. In the pre-processing stage, we tailor the data format for a specific framework and remove NA values. We run SuSiE-inf and FINEMAP-inf in one framework allowing to initialize FINEMAP-inf using tau-squared and sigma-squared parameters generated by SuSiE-inf to improve accuracy. When running FINEMAP, we use a shotgun stochastic search subprogram. When evaluating quality of signal localization, we define the true positive as a discovery within a $50\,\mathrm{kb}$ window where the causal signal was planted.

We run the gVAMP method using as prior initialization a spike at zero and 22 slab mixtures. We let the variance of those mixtures follow a geometric progression in the interval $[10^{-7}, 1]$, and we let the probabilities follow a geometric progression with factor $1/2$. The prior probability $1 - \lambda$ of variants being assigned to the 0 mixture is initialized to 99%. Variant effect sizes are initialized with 0. Based on the experiments in the UK Biobank imputed dataset, in WGS we use damping factor $\rho = 0.05$ and run the method until iteration 30 when the hyper-parameters stabilize.

gVAMP model parameters for the analysis of human height in WGS data

We apply gVAMP to the WGS data to analyse human height using the largest computational instance currently available on the DNAnexus platform, employing 128 cores and $1921.4\,\mathrm{GB}$ total memory. Efficient C++ gVAMP implementation allows for parallel computing, utilizing OpenMP and MPI libraries. Here, we split the memory requirements and computational workload between 2 OpenMP threads and 64 MPI workers. The prior initialization of gVAMP is similar to that used in the simulation study, except that we opt for a sparser prior with probability 99.5% of variants being assigned to the 0 mixture. As before, the SNP marker effect sizes are initialised with 0. Based on the experiments in the UK Biobank imputed dataset, in WGS we set the initial damping factor ρ to 0.1, and adjust it to 0.05 from iteration 4 onward, which stabilizes the algorithm and leads us to stop it at iteration 18. We also note that, after the first few iterations, the marker inclusion probabilities are stable.

gVAMP model parameters for imputed SNP data

We run gVAMP on the 13 UK Biobank phenotypes on the full 8,430,446 SNP set, and on the 2,174,071 and 882,727 LD clumped SNP set. We find that setting the damping factor ρ to 0.1 performs well for all the 13 outcomes in the UK Biobank that we have considered. For the prior initialization, we use a spike at zero and 22 slab mixtures following a geometric progression in the interval $[10^{-6}, 10]$. The probabilities of the 22 slab mixtures follow a geometric progression with factor $1/2$, and the probability of being assigned to the 0 mixture is 97%. This configuration works well for all phenotypes. We also note that our inference of the number of mixtures, their probabilities, their variances and the SNP marker effects is not dependent upon specific starting parameters for the analyses of the 2,174,071 and 882,727 SNP datasets, and the algorithm is rather stable for a range of initialization choices. Similarly, the algorithm is stable for different choices of the damping ρ, as long as said value is not too large.

Generally, appropriate starting parameters are not known in advance and this is why we learn them from the data within the EM steps of our algorithm. However, it is known that EM can be sensitive to the starting values given and, thus, we recommend initialising a series of models at

different values to check that this is not the case (similar to starting multiple Monte Carlo Markov chains in standard Bayesian methods). The feasibility of this recommendation is guaranteed by the significant speed-up of our algorithm compared to existing approaches, see Supplementary Note 1, Figure S21.

Given the same data, gVAMP yields estimates that more closely match GMRM when convergence in the training R^2, SNP heritability, residual variance, and out-of-sample test R^2 are smoothly monotonic within around 10-40 iterations. Following this, training R^2, SNP heritability, residual variance, and out-of-sample test R^2 may then begin to slightly decrease as the number of iterations becomes large. Thus, as a stopping criterion for the 2,174,071 and 882,727 SNP datasets, we choose the iteration that maximizes the training R^2.

We highlight the iterative nature of our method. Thus, improved computational speed and more rapid convergence is achieved by providing better starting values for the SNP marker effects. Specifically, when moving from 2,174,071 to 8,430,446 SNPs, only columns with correlation $R^2 \geq 0.8$ are being added back into the data. Thus, for the 8,430,446 SNP set, we initialise the model with the converged SNP marker and prior estimates obtained from the 2,174,071 SNP runs, setting to 0 the missing markers. Furthermore, we lower the value of the damping factor ρ, with typical values being 0.05 and 0.01. We experiment both with using the noise precision from the initial 2,174,071 SNP runs and with setting it to 2. We then choose the model that leads to a smoothly monotonic curve in the training R^2. We observe that SNP heritability, residual variance, and out-of-sample test R^2 are also smoothly monotonic within 25 iterations. Thus, as a stopping criterion for the 8,430,446 SNP dataset, we choose the estimates obtained after 25 iterations for all the 13 traits. We follow the same process when extending the analyses to include the WES rare burden gene scores.

Polygenic risk scores and SNP heritability

gVAMP produces SNP effect estimates that can be directly used to create polygenic risk scores. The estimated effect sizes are on the scale of normalised SNP values, i.e., $(\boldsymbol{X}_j - \mu_{X_j})/SD(\boldsymbol{X}_j)$, with μ_{X_j} the column mean and $SD(\boldsymbol{X}_j)$ the standard deviation, and thus SNPs in the out-of-sample prediction data must also be normalized. We provide an option within the gVAMP software to do phenotypic prediction, returning the adjusted prediction R^2 value when given input data of a PLINK file and a corresponding file of phenotypic values. gVAMP estimates the SNP heritability as the phenotypic variance (equal to 1 due to normalization) minus 1 divided by the estimate of the noise precision, i.e., $h^2_{SNP} = 1 - 1/\gamma_\epsilon$.

We compare gVAMP to a MCMC sampler approach (GMRM) with a similar prior (the same number of starting mixtures) as presented in [9]. We select this comparison as the MCMC sampler was demonstrated to exhibit the highest genomic prediction accuracy up to date [9]. We run GMRM for 2000 iterations, taking the last 1800 iterations as the posterior. We calculate the posterior means for the SNP effects and the posterior inclusion probabilities of the SNPs belonging to the non-zero mixture group. GMRM estimates the SNP heritability in each iteration by sampling from an inverse χ^2 distribution using the sum of the squared regression coefficient estimates. We also compare to the LDAK-KVIK elastic net software [10], run using their suggested parameters and default options.

Finally, we also compare gVAMP to the summary statistics prediction methods LDpred2 [7] and SBayesR [8] using summary statistic data of 8,430,446 SNPs obtained from the REGE-

NIE software. For SBayesR, following the recommendation on the software webpage (`https://cnsgenomics.com/software/gctb/#SummaryBayesianAlphabet`), after splitting the genomic data per chromosomes, we calculate the so-called *shrunk* LD matrix, which use the method proposed by [45] to shrink the off-diagonal entries of the sample LD matrix toward zero based on a provided genetic map. We make use of all the default values: `--genmap-n` 183, `--ne` 11400 and `--shrunk-cutoff` 10^{-5}. Following that, we run the SBayesR software using summary statistics generated via the REGENIE software (see "Mixed linear association testing" below) by grouping several chromosomes in one run. Namely, we run the inference jointly on the following groups of chromosomes: $\{1\}, \{2\}, \{3\}, \{4\}, \{5, 6\}, \{7, 8\}, \{9, 10, 11\}, \{12, 13, 14\}$ and $\{15, 16, 17, 18, 19, 20, 21, 22\}$. This allows to have locally joint inference, while keeping the memory requirements reasonable. All the traits except for Blood cholesterol (CHOL) and Heel bone mineral density T-score (BMD) give non-negative R^2; CHOL and BMD are then re-run using the option to remove SNPs based on their GWAS p-values (threshold set to 0.4) and the option to filter SNPs based on LD R-Squared (threshold set to 0.64). For more details on why one would take such an approach, one can check `https://cnsgenomics.com/software/gctb/#FAQ`. As the obtained test R^2 values are still similar, as a final remedy, we run standard linear regression over the per-group predictors obtained from SBayesR on the training dataset. Following that, using the learned parameters, we make a linear combination of the per-group predictors in the test dataset to obtain the prediction accuracy given in the table.

For LDpred2, following the software recommendations, we create per-chromosome banded LD matrices with the window size of 3cM. After the analysis of the genome-wide run of LDpred2, we establish that the chains do not converge even after tuning the shrinkage factor, disabling the sign jump option and disabling the usage of MLE (`use_MLE=FALSE` option). For this reason, we opt to run LDpred2 per chromosome, in which case the chains converge successfully. Twenty chains with different proportion of causal markers are run in the LDpred2 method, for each of the chromosomes independently. Then, a standard linear regression involving predictors from different chromosomes is performed to account for correlations between SNPs on different chromosomes, which achieved better test R^2 than the predictors obtained by stacking chromosomal predictors. In summary, for both LDpred2 and SBayesR we have tried to find the optimal solution to produce the highest possible out-of-sample prediction accuracy, contacting the study authors, if required, for guidance.

Mixed linear model association testing

We conduct mixed linear model association testing using a leave-one-chromosome-out (LOCO) estimation approach on the 8,430,446 and 2,174,071 imputed SNP markers. LOCO association testing approaches have become the field standard and they are two-stage: a subset of markers is selected for the first stage to create genetic predictors; then, statistical testing is conducted in the second stage for all markers one-at-a-time. We consider REGENIE [1], as it is a recent commonly applied approach and LDAK-KVIK [10]. We also compare to GMRM [9], a Bayesian linear mixture of regressions model that has been shown to outperform REGENIE for LOCO testing. For the first stage of LOCO, REGENIE is given 887,060 markers to create the LOCO genetic predictors, even if it is recommended to use 0.5 million genetic markers. We compare the number of significant loci obtained from REGENIE to those obtained if one were to replace the LOCO predictors with:

(i) those obtained from GMRM using the LD pruned sets of 2,174,071 and 887,060 markers; and *(ii)* those obtained from gVAMP at all 8,430,446 markers and the LD pruned sets of 2,174,071 and 887,060 markers. We note that obtaining predictors from GMRM at all 8,430,446 markers is computationally infeasible, as using the LD pruned set of 2,174,071 markers already takes GMRM several days. In contrast, gVAMP is able to use all 8,430,446 markers and still be faster than GMRM with the LD pruned set of 2,174,071 markers. LDAK-KVIK is given 2,174,071 markers to facilitate a direct comparison with gVAMP and we compare *(i)* placing the predictors of gVAMP directly within the LDAK-KVIK step 2; *(ii)* using gVAMP step 2 procedure which is identical to that of REGENIE.

LOCO testing does not control for linkage disequilibrium within a chromosome. Thus, to facilitate a simple, fair comparison across methods, we clump the LOCO results obtained with the following PLINK commands: `--clump-kb` 1000 `--clump-r2` 0.01 `--clump-p1` 0.00000005. Therefore, within 1Mb windows of the DNA, we calculate the number of independent associations (squared correlation ≤ 0.01) identified by each approach that pass the genome-wide significance testing threshold of $5 \cdot 10^{-8}$. As LOCO can only detect regions of the DNA associated with the phenotype and not specific SNPs, given that it does not control for the surrounding linkage disequilibrium, a comparison of the number of uncorrelated genome-wide significance findings is conservative. A full benchmark of approaches in a simulation study is also provided in Supplementary Note 1, where we consider a wider range of metrics, scenarios and comparisons.

gVAMP association testing

In line with the standard outputs of MCMC methods, the AMP framework additionally offers the possibility of calculating posterior inclusion probabilities (PIPs) jointly for each of the SNPs. Once the prior information in a particular iteration is updated, we use Equation 6 to approximate the probability of each SNP belonging to each of the slab mixture components. Finally, by summing up all the mentioned probabilities for a particular SNP we obtain its posterior inclusion probability. To conduct variable selection, we apply a threshold of $PIP \geq 0.95$.

We can additionally exchange PIP values for p-values. Relying on the properties of the EM-VAMP estimator, whose noise is asymptotically Gaussian due to the Onsager correction [14], we have that $\boldsymbol{r}_{1,t} \approx \boldsymbol{\beta} + \mathcal{N}(0, \gamma_{1,t}^{-1}\boldsymbol{I})$, where $\boldsymbol{\beta}$ is the ground-truth value of the underlying genetic effects vector. More precisely, one can show that $\frac{1}{N}\|\boldsymbol{r}_{1,t} - \boldsymbol{\beta} - \mathcal{N}(0, \gamma_{1,t}^{-1}\boldsymbol{I})\| \to 0$, as $N, P \to \infty$, with the ratio N/P being fixed. Therefore, for each marker with index j, a one-sided p-value for the hypothesis test H0 : $\beta_j = 0$ is given by $\Phi(-|(\boldsymbol{r}_{1,t})_i| \cdot \gamma_{1,t}^{1/2})$, where Φ is the CDF of a standard normal distribution and $(\boldsymbol{r}_{1,t})_i$ denotes the i-th component of the vector $\boldsymbol{r}_{1,t}$.

Data availability

This project uses the UK Biobank data under project number 35520. UK Biobank genotypic and phenotypic data is available through a formal request at (`http://www.ukbiobank.ac.uk`). All summary statistic estimates are released publicly on Dryad: `https://doi.org/xx.xxxx/dryad.xxxxxxxxx`.

Code availability

The gVAMP code `https://github.com/medical-genomics-group/gVAMP` is fully open source. The scripts used to execute the model are available at `https://github.com/medical-genomics-group/gVAMP`. R version 4.2.1 is available at `https://www.r-project.org/`. PLINK version 1.9 is available at `https://www.cog-genomics.org/plink/1.9/`. REGENIE is available at `https://github.com/rgcgithub/regenie`. bignspr 1.12.4 package that contains LDpred2 is available at `https://privefl.github.io/bigsnpr/index.html`. SBayesR is available at `https://cnsgenomics.com/software/gctb/#Overview`.

Acknowledgments

We would like to thank our three reviewers whose comments improved the presentation of this work. We thank Malgorzata Borczyk for creating the gene burden scores. We thank Robin Beaumont, Amedeo Roberto Esposito, Gareth Hawkes, Philip Schniter, Matthew Stephens, Pragya Sur, Peter Visscher, Michael Weedon and Harry Wright for providing valuable suggestions and comments on earlier versions of the work. We would like to acknowledge the participants and investigators of the UK Biobank study. This project was funded by a Lopez-Loreta Prize to MM, by an SNSF Eccellenza Grant to MRR (PCEGP3-181181), and by core funding from the Institute of Science and Technology Austria. High-performance computing was supported by the Scientific Service Units (SSU) of IST Austria through resources provided by Scientific Computing (SciComp).

Author contributions

MM and MRR conceived the study. AD, MM and MRR designed the study. AD derived the model and the algorithm, with input from MM and MRR. AD wrote the software, with input from MM and MRR. AD, JB, MM, and MRR conducted the analysis and wrote the paper. All authors approved the final manuscript prior to submission.

Author competing interests

MRR receives research funding from Boehringer Ingelheim for work that is unrelated to that presented here. All other authors declare no competing interests.

Ethical approval declaration

This project uses UK Biobank data under project 35520. UK Biobank genotypic and phenotypic data is available through a formal request at `http://www.ukbiobank.ac.uk`. The UK Biobank has ethics approval from the North West Multi-centre Research Ethics Committee (MREC). Methods were carried out in accordance with the relevant guidelines and regulations, with informed consent obtained from all participants.

References

1. Mbatchou, J. *et al.* Computationally efficient whole-genome regression for quantitative and binary traits. *Nature Genetics* **53**, 1097–1103 (2021).

2. Loh, P.-R. *et al.* Efficient bayesian mixed-model analysis increases association power in large cohorts. *Nature Genetics* **47**, 284–290 (2015).

3. Zhou, W. *et al.* Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nature Genetics* **50**, 1335–1341 (2018).

4. Jiang, L., Zheng, Z., Fang, H. & Yang, J. A generalized linear mixed model association tool for biobank-scale data. *Nature Genetics* **53**, 1616–1621 (2021).

5. Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A Simple New Approach to Variable Selection in Regression, with Application to Genetic Fine Mapping. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **82**, 1273–1300 (2020).

6. Speed, D. *et al.* Reevaluation of snp heritability in complex human traits. *Nature genetics* **49**, 986–992 (2017).

7. Privé, F., Arbel, J. & Vilhjálmsson, B. J. LDpred2: better, faster, stronger. *Bioinformatics* **36**, 5424–5431 (2020).

8. Lloyd-Jones, L. R. *et al.* Improved polygenic prediction by bayesian multiple regression on summary statistics. *Nature Communications* **10**, 5086 (2019).

9. Orliac, E. J. *et al.* Improving gwas discovery and genomic prediction accuracy in biobank data. *Proceedings of the National Academy of Sciences* **119**, e2121279119 (2022).

10. Hof, J. & Speed, D. Ldak-kvik performs fast and powerful mixed-model association analysis of quantitative and binary phenotypes. *medRxiv* 2024–07 (2024).

11. Spence, J. P., Sinnott-Armstrong, N., Assimes, T. L. & Pritchard, J. K. A flexible modeling and inference framework for estimating variant effect sizes from gwas summary statistics. *bioRxiv* (2022).

12. Lawson, D. J. *et al.* Is population structure in the genetic biobank era irrelevant, a challenge, or an opportunity? *Human Genetics* **139**, 23–41 (2020).

13. Donoho, D. L., Maleki, A. & Montanari, A. Message Passing Algorithms for Compressed Sensing. *Proceedings of the National Academy of Sciences* **106**, 18914–18919 (2009).

14. Rangan, S., Schniter, P. & Fletcher, A. K. Vector approximate message passing. *IEEE Transactions on Information Theory* **65**, 6664–6684 (2019).

15. Feng, O. Y., Venkataramanan, R., Rush, C., Samworth, R. J. *et al.* A unifying tutorial on approximate message passing. *Foundations and Trends® in Machine Learning* **15**, 335–536 (2022).

16. Bayati, M. & Montanari, A. The dynamics of message passing on dense graphs, with applications to compressed sensing. *IEEE Transactions on Information Theory* **57**, 764–785 (2011).

17. Montanari, A. & Venkataramanan, R. Estimation of low-rank matrices via approximate message passing. *Annals of Statistics* **45**, 321–345 (2021).

18. Barbier, J., Krzakala, F., Macris, N., Miolane, L. & Zdeborová, L. Optimal errors and phase transitions in high-dimensional generalized linear models. *Proceedings of the National Academy of Sciences* **116**, 5451–5460 (2019).

19. Barbier, J., Camilli, F., Mondelli, M. & Sáenz, M. Fundamental limits in structured principal component analysis and how to reach them. *Proceedings of the National Academy of Sciences* **120** (2023).

20. Jeon, C., Ghods, R., Maleki, A. & Studer, C. Optimality of large MIMO detection via approximate message passing. In *IEEE International Symposium on Information Theory*, 1227–1231 (2015).

21. Metzler, C. A., Maleki, A. & Baraniuk, R. G. From denoising to compressed sensing. *IEEE Trans. Information Theory* **62**, 5117–5144 (2016).

22. Eksioglu, E. M. & Tanc, A. K. Denoising AMP for MRI reconstruction: BM3D-AMP-MRI. *SIAM Journal on Imaging Sciences* **11**, 2090–2109 (2018).

23. Zhong, X., Su, C. & Fan, Z. Empirical bayes pca in high dimensions. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **84**, 853–878 (2022).

24. Fletcher, A. K. & Schniter, P. Learning and free energies for vector approximate message passing. In *2017 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, 4247–4251 (2017).

25. Fletcher, A. K., Sahraee-Ardakan, M., Rangan, S. & Schniter, P. Rigorous dynamics and consistent estimation in arbitrarily conditioned linear systems. In *Advances in Neural Information Processing Systems*, vol. 30 (2017).

26. Yengo, L. *et al.* A saturated map of common genetic variants associated with human height. *Nature* **610**, 704–712 (2022).

27. Simons, Y. B., Bullaughey, K., Hudson, R. R. & Sella, G. A population genetic interpretation of gwas findings for human quantitative traits. *PLoS biology* **16**, e2002985 (2018).

28. Si, Y., Vanderwerff, B. & Zöllner, S. Why are rare variants hard to impute? coalescent models reveal theoretical limits in existing algorithms. *Genetics* **217** (2021).

29. Wainschtein, P. *et al.* Assessing the contribution of rare variants to complex trait heritability from whole-genome sequence data. *Nature Genetics* **54**, 263–273 (2022).

30. Robinson, M. R. *et al.* Genotype–covariate interaction effects and the heritability of adult body mass index. *Nature Genetics* **49**, 1174–1181 (2017).

31. Hindorff, L. A. *et al.* Prioritizing diversity in human genomics research. *Nature Reviews Genetics* **19**, 175–185 (2018).

32. Campos, A. I. *et al.* Boosting the power of genome-wide association studies within and across ancestries by using polygenic scores. *Nature Genetics* **55**, 1769–1776 (2023).

33. Schniter, P., Rangan, S. & Fletcher, A. K. Vector approximate message passing for the generalized linear model. In *2016 50th Asilomar conference on signals, systems and computers*, 1525–1529 (IEEE, 2016).

34. Fletcher, A. K., Pandit, P., Rangan, S., Sarkar, S. & Schniter, P. Plug in estimation in high dimensional linear inverse problems: A rigorous analysis. *Journal of Statistical Mechanics: Theory and Experiment* **2019**, 124021 (2019).

35. Krzakala, F., Mézard, M., Sausset, F., Sun, Y. & Zdeborová, L. Probabilistic reconstruction in compressed sensing: algorithms, phase diagrams, and threshold achieving matrices. *Journal of Statistical Mechanics: Theory and Experiment* **2012**, P08009 (2012).

36. Donoho, D. & Tanner, J. Observed universality of phase transitions in high-dimensional geometry, with implications for modern data analysis and signal processing. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **367**, 4273–4293 (2009).

37. Monajemi, H. *et al.* Deterministic matrices matching the compressed sensing phase transitions of gaussian random matrices. *Proceedings of the National Academy of Sciences* **110**, 1181–1186 (2013).

38. Wang, T., Zhong, X. & Fan, Z. Universality of approximate message passing algorithms and tensor networks. *The Annals of Applied Probability* **34**, 3943–3994 (2024).

39. Dudeja, R., Lu, Y. M. & Sen, S. Universality of approximate message passing with semi-random matrices. *Annals of Probability* (2023).

40. Skuratovs, N. & Davies, M. E. Warm-starting in message passing algorithms. In *IEEE International Symposium on Information Theory (ISIT)*, 1187–1192 (2022).

41. Hutchinson, M. A stochastic estimator of the trace of the influence matrix for laplacian smoothing splines. *Communications in Statistics - Simulation and Computation* **19**, 433–450 (1990).

42. Vila, J., Schniter, P., Rangan, S., Krzakala, F. & Zdeborová, L. Adaptive damping and mean removal for the generalized approximate message passing algorithm. In *IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, 2021–2025 (2015).

43. Vila, J. & Schniter, P. Expectation-maximization gaussian-mixture approximate message passing. *IEEE Transactions on Signal Processing* **61** (2012).

44. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

45. Wen, X. & Stephens, M. Using linear predictors to impute allele frequencies from summary or pooled genotype data. *The Annals of Applied Statistics* **4**, 1158 – 1182 (2010).

46. Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M. & Price, A. L. Advantages and pitfalls in the application of mixed-model association methods. *Nature Genetics* **46**, 100–106 (2014).

47. Daetwyler, H. D., Villanueva, B. & Woolliams, J. A. Accuracy of predicting the genetic risk of disease using a genome-wide approach. *PLOS ONE* **3**, 1–8 (2008).

Supplementary information

Joint modeling of whole genome sequence data for human height via approximate message passing

Al Depope, Jakub Bajzik, Marco Mondelli, Matthew R. Robinson

Supplementary Tables

Table S1. The 12 simulation settings of a whole genome sequence (WGS) data simulation study using chromosome 12 of the UK Biobank. Each line gives the code used in the study (from *a* to *l*), the number of causal variants that were simulated one per 50kb block (163 or 400) starting from the beginning of the full WGS data of chromosome 12, the proportion of phenotypic variance attributable to the causal variants (heritability, 6.25% or 12.5%), whether variants were randomly allocated with respect to the frequency of the SNP or whether variants of frequency between 1% and 0.1% were chosen (rare, yes or no), and finally the power relationship (α) between the causal variant effect size and the frequency. Note that α is on the scale of standardized genetic variants, which translates as -1 for 0 values and -1.5 for values of -0.5 for genetic variants on the unstandardized scale.

ID	Number of causal variants	Heritability	Rare	α
a	163	6.25%	Yes	0.0
b	163	6.25%	No	0.0
c	163	6.25%	Yes	-0.5
d	163	6.25%	No	-0.5
e	400	12.5%	No	-0.5
f	400	12.5%	Yes	-0.5
g	400	12.5%	No	0.0
h	400	12.5%	Yes	0.0
i	400	6.25%	No	-0.5
j	400	6.25%	Yes	-0.5
k	400	6.25%	No	0.0
l	400	6.25%	Yes	0.0

Table S2. The 13 **UK Biobank traits used within the study**. Phenotypic names and their codes used in the study. The sample size, N, gives the number of individuals with training data measures.

Phenotype	Code	Sample size, N
Blood: cholesterol	CHOL	395,025
Blood: eosinophill count	EOSI	401,452
Blood: glycated haemoglobin	HbA1c	394,912
Blood: High density lipoprotein	HDL	360,286
Blood: mean corpuscular haemoglobin	MCH	402,201
Blood mean corpuscular volume	MCV	402,202
Red blood cell count	RBC	402,204
Body mass index	BMI	413,595
Diastolic blood pressure	DBP	377,358
Forced vital capacity	FVC	376,724
Heel bone mineral density	BMD	231,693
Standing height	HT	414,055
Systolic blood pressure	SBP	377,347

Table S3. The relationship between minor allele frequency and effect size for human height in 16,854,878 whole genome sequence (WGS) variants and 8,430,446 imputed SNP variants in the UK Biobank. We estimate the power law α parameter value with one standard deviation confidence interval from a whole-genome sequence gVAMP analysis of standing height (*WGS SNPs*) compared to a gVAMP analysis of imputed SNPs (*Imputed SNPs*). The standard deviation is calculated by treating each chromosome specific α estimate as one realization. Columns named *WES (WGS + WES)* and *WES (Imputed + WES)* give point estimate of α parameter considering only 17,852 gene burden score effects obtained in joint runs with either WGS data or imputed data. Rows represent different PIP thresholds used to select variants based on which the α parameter is estimated. α values are presented for genetic variants that are unscaled (0,1,2 scale).

PIP threshold	WGS SNPs	Imputed SNPs	WES	
			WGS + WES	Imputed + WES
0.95	-1.1135 $\pm$ 0.2351	-1.0046 $\pm$ 0.0588	-1.0546	-1.0662
0.5	-0.9661 $\pm$ 0.1010	-0.7570 $\pm$ 0.1332	-1.0168	-0.9276

Table S4. Genome-wide significant associations for 13 **UK Biobank traits from the software gVAMP and LDAK-KVIK at 8,430,446 genetic variants from mixed linear model association leave-one-chromosome-out (MLMA-LOCO) testing.** "LDAK 2M LOCO" refers to MLMA-LOCO testing at 8,430,446 SNPs, using the LDAK-KVIK software in step 1, with 2,174,071 SNP markers. "gVAMP 2M LOCO 1" refers to MLMA-LOCO testing at 8,430,446 SNPs, using the framework presented here, where in step 1 2,174,071 markers are used and the second step is identical to that of LDAK-KVIK. "gVAMP 2M LOCO 2" is the same as "gVAMP 2M LOCO 1" but with the second step identical to REGENIE. The values are the number of genome-wide significant linkage disequilibrium independent associations selected based upon a standard p-value threshold of less than $5 \cdot 10^{-8}$ and R^2 between SNPs in a 1 Mb genomic segment of less than 1%.

Trait	LDAK 2M LOCO	gVAMP 2M LOCO 1	gVAMP 2M LOCO 2
CHOL	482	625	603
EOSI	549	666	630
HbA1c	321	400	385
HDL	639	855	812
MCH	690	858	810
MCV	903	1128	1062
RBC	856	1140	1079
BMI	726	1202	1175
DBP	303	357	419
FVC	577	759	721
BMD	549	681	668
HT	2703	4747	4452
SBP	323	410	388

Supplementary Figures

Figure S1. Comparison of the true positive rate (TPR, x-axis) versus the false discovery rate (FDR, y-axis) between gVAMP and marginal GWAS association testing plus fine-mapping implemented in the SuSiE algorithm across 12 simulation scenarios using whole genome sequence (WGS) data of chromosome 12 in the UK Biobank. For the detailed description of different simulation settings, see Supplementary Table S1. The TPR and FDR of marginals plus SuSiE (orange) and gVAMP (blue) are calculated as associations that were within 50 kb of a true causal variant. For each 50kb DNA segment, we use a Bonferroni corrected p-value significance threshold for marginal GWAS testing and then having identified associated regions we examine the ability of SuSiE to conduct variable selection across PIP thresholds (orange). Error bands give the standard deviation across simulation replicates.

Figure S2. Performance in terms of the false discovery rate of marginal GWAS testing plus fine-mapping with SuSiE and gVAMP at different posterior inclusion probability thresholds in a whole genome sequence (WGS) data simulation study of chromosome 12 in the UK Biobank. For the detailed description of different simulation settings, see Supplementary Table S1. Error bands give the standard deviation across simulation replicates. As the posterior inclusion probability (PIP) threshold increases to 95%, the false discovery rate declines to around 5% in most settings for gVAMP as expected. For comparison, at each 50kb DNA segment we use a Bonferroni corrected p-value significance threshold for marginal GWAS testing and then having identified associated regions we examine the ability of SuSiE to conduct variable selection across PIP thresholds (orange). Due to its reliance on the marginal testing significance threshold, the FDR of SuSiE is basically flat. Note that the performance of SuSiE remains poor even without the marginal testing.

Figure S3. Performance in terms of the true positive rate and the false discovery rate of marginal GWAS regression association testing, marginal GWAS testing plus fine-mapping with SuSiE, and gVAMP at posterior inclusion probability ≥ 0.95 in a whole genome sequence (WGS) data simulation study of chromosome 12 in the UK Biobank. We use a Bonferroni corrected p-value significance threshold for marginal GWAS (gray) and $PIP > 0.95$ for SuSiE applied to 50kb regions identified as significant in the marginal analysis (orange). Marginal GWAS and SuSiE do not control the FDR in these WGS data simulation settings (see Supplementary Table S1). In contrast, using a $PIP > 0.95$ threshold for the gVAMP testing (blue) controls FDR, whilst yielding high TPR. Error bars give the standard deviation across simulation replicates. "gVAMP $\rho \geq 0.4$" indicates the FDR of gVAMP to select a variable with absolute correlation ≥ 0.4 to a causal variant.

Figure S4. The difference in frequency (y-axis) of variants selected by gVAMP at posterior inclusion probability ≥ 0.95 to the most correlated causal variant in a whole genome sequence (WGS) data simulation study using chromosome 12 of the UK Biobank. The frequency of variants selected by gVAMP in an anlysis of either WGS (blue) or imputed (orange) SNP data is unbiased for either rare (left part of each sub-plot) or common (right part of each sub-lot) SNPs. Error bars give the standard deviation across simulation replicates.

Figure S5. gVAMP provides both p-values and posterior inclusion probabilities for each SNP. The TPR and FDR of gVAMP in a whole genome sequence (WGS) data simulation study of chromosome 12 in the UK Biobank. Using p-values (gray dashed line) or posterior inclusion probability values (PIP, blue solid line) gives indistinguishable curves. TPR and FDR are calculated as associations that are within 50kb of a true causal variant. Error bands give the standard deviation across simulation replicates.

Figure S6. Benchmarking of gVAMP to a wide range of fine-mapping methods applied to marginal GWAS summary data in a whole genome sequence (WGS) data simulation study of chromosome 12 in the UK Biobank. We compare the true positive rate (TPR) and false discovery rate (FDR) of gVAMP (blue) versus using a Bonferroni corrected p-value significance threshold for marginal GWAS testing, and then having identified regions, we examine the ability of SuSiE to conduct variable selection across PIP thresholds (orange). Giving SuSiE the true prior as initialization (SuSiE true prior), running SuSiE in windows up to 500kb (SuSiE 500kb window), or running SuSiE-inf combining inference across blocks only marginally improves performance. Both FINEMAP and FINEMAP-inf also yield results that are similar to the equivalent SuSiE implementations. TPR and FDR are calculated as associations that are within 50kb of a true causal variant. Error bands give the standard deviation across simulation replicates.

Figure S7. Comparison of the true positive rate (TPR, x-axis) versus the false discovery rate (FDR, y-axis) of gVAMP when grouping common variants into sets of different minimum correlation (ρ) in a whole genome sequence (WGS) data simulation study of chromosome 12 in the UK Biobank. The TPR and FDR of are calculated as associations that were within 50 kb of a true causal variant. Error bands give the standard deviation across simulation replicates.

Figure S8. The proportion of variance attributable to DNA loci of different frequency in a whole genome sequence (WGS) data simulation study using chromosome 12 of the UK Biobank. We plot the proportion of the sum of the squared regression coefficients attributable to variants of different frequency as estimated from gVAMP in WGS (blue) or imputed SNP (orange) data. The ground truth (green) is calculated as the sum of squared true effects. The variance attributable to SNPs of different frequency matches the ground truth for gVAMP in WGS data and is more likely to be mis-estimated when gVAMP is applied to imputed SNP data. Error bars give the standard deviation across simulation replicates.

Figure S9. Estimation of power law exponent α of the relationship between effect size and minor allele frequency (MAF) in a whole genome sequencing (WGS) data simulation study using chromosome 12 of the UK Biobank. Across different simulation settings (see Supplementary Table S1) where the power relationship of effect size and MAF is simulated to be either 0 or −0.5, gVAMP infers the correct direction of the power law on either using standardized variants with zero mean and unit variance, i.e. on the variance scale in (a), or on the 'effect size' scale of non-standardized genetic variants in (b), but can moderately underestimate the true value. The underestimation is likely due to the limited number of causal variants used within the simulation study, and we expect performance to improve in a genome-wide setting. We estimate the power law exponent following the derivation given in Supplementary Note 3 genome-wide across all variants identified at posterior inclusion probability ≥ 0.95. Error bars give the standard deviation across simulation replicates.

Figure S10. The variance attributable to gene burden scores calculated from whole exome sequence (WES) data (y-axis) shows no relationship to the variance attributable to DNA variants within and around the gene (x-axis) estimated from a gVAMP analysis of either 8,430,446 imputed SNP markers or 16,854,878 whole genome sequence (WGS) variants.

Figure S11. Marginal regression coefficient estimates of the relationship of 16,854,878 whole genome sequence (WGS) variants with human height, plotted against the regression coefficient values obtained from gVAMP. "WGS" refers to the gVAMP run on the WGS SNPs, "Imputed" refers to the gVAMP run on 8,430,446 imputed SNP markers, and "Sim" gives the same comparison of marginal and gVAMP estimates in setting d. The comparison is plotted for common (minor allele frequency, MAF $\geq 1\%$) and rare variants (MAF $< 1\%$).

Figure S12. SNP heritability estimation of GMRM versus gVAMP with different numbers of SNP markers across 13 traits in the UK Biobank. Comparison of the proportion of phenotypic variation attributable to 2,174,071 autosomal SNP genetic markers (SNP heritability) estimated by GMRM (x-axis) to the SNP heritability estimated by gVAMP (y-axis) at either the same 2,174,071 SNPs (red) or 8,430,446 SNP markers (blue). The slope of the lines shows a 1-to-1 relationship of gVAMP to GMRM, but with an average of 3.4% lower estimate for gVAMP at 2.17M SNPs. Analysing 8.4M SNPs with gVAMP increases the heritability estimate over GMRM by 3.9%, which is consistent with an increase in phenotypic variance captured by the full imputed sequence data, as opposed to a selected subset of SNP markers. The dashed grey line gives $y = x$.

Supplementary Note 1

Figure S13. Simulation study of out-of-sample prediction accuracy in UK Biobank imputed genotype data. We compare out-of-sample prediction accuracy (y-axis) for polygenic risk scores (PRS) created at different sets of markers ("880k" = 887,060, "1m" = 1,410,525, "2m" = 2,174,071 SNPs) from gVAMP (blue) and an elastic net model implemented in the LDAK software (red), as compared to the prediction accuracy obtained at the same sets of SNPs from a ridge regression model implemented in the LDAK software (x-axis). Grey lines give the 1:1 line. Comparisons are made across three settings: "V1" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers and effects are drawn from a Gaussian, "V2" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects that have a power relationship to the minor allele frequency of -0.5, and "V3" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects drawn from an elastic net distribution.

Simulation study in imputed SNP data

To support our empirical analyses we also conduct a simulation study in imputed SNP data using the 8,430,446 UK Biobank genetic marker data with 414,055 individuals. We first randomly sample 40,000 causal variants genome-wide to give a highly polygenic genetic basis. We then simulate three scenarios. First, in *V1*, we allocate effect sizes from a Gaussian with mean zero and variance 0.5/40000, where 0.5 is the proportion of variance attributable to the SNP markers. Multiplying the simulated SNP effects by normalized values of the 40,000 causal markers, gives a vector of genetic values of length $N = 414055$ with variance 0.5. To this we add a vector of noise, drawn from a

Figure S14. Simulation study of power for mixed linear model leave-one-chromosome out association testing in UK Biobank imputed genotype data. We compare the average chi-squared value at the causal variants (y-axis) for leave-one-chromosome-out association testing (LOCO) when the first step LOCO predictors are created at different sets of markers ("880k" = 887,060, "1m" = 1,410,525, "2m" = 2,174,071 SNPs) from gVAMP (blue), an elastic net model implemented in the LDAK software (red), and Regenie (green). Comparisons are made across three settings: "V1" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers and effects are drawn from a Gaussian, "V2" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects that have a power relationship to the minor allele frequency of -0.5, and "V3" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects drawn from an elastic net distribution.

Gaussian with mean zero and variance 0.5, to produce a response variable of length N, with zero mean and unit variance. Second, in *V2*, we simulate an α-power relationship between the causal variant effect size and the minor allele frequency (see Supplementary Note 3), with $\alpha = -0.5$. Third, in *V3*, we simulate under an elastic net prior, where with probability 0.8 we sample effect sizes from a standard Gaussian, with probability 0.2 we sample from a double exponential distribution, and we then scale the effects vector so that multiplying them with normalized values of the 40,000 causal markers, gives a vector of genetic values of length $N = 414055$ with variance 0.5.

We analyze the simulated response variable with gVAMP, using either 8,430,446, 2,174,071, 1,410,525 or 887,060 SNP markers with identical initialization to that described in the Methods for the empirical UK Biobank study. We also analyze the data with REGENIE using 2,174,071, 1,410,525 or 887,060 SNP markers for the first stage and 8,430,446 SNP markers for the second stage

Figure S15. Simulation study of true positive rate (TPR), false discovery rate (FDR) and type-I error for mixed linear model leave-one-chromosome out association testing in UK Biobank imputed genotype data. We compare performance of leave-one-chromosome-out association testing (LOCO) when the first step LOCO predictors are created at different sets of markers ("880k" = 887,060, "1m" = 1,410,525, "2m" = 2,174,071 SNPs) from gVAMP (blue), an elastic net model implemented in the LDAK software (red), and Regenie (green). Comparisons are made across three settings: "V1" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers and effects are drawn from a Gaussian, "V2" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects that have a power relationship to the minor allele frequency of -0.5, and "V3" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects drawn from an elastic net distribution. For each 1Mb window that contains an identified genome-wide significant association, we then ask if the identified variants are correlated (squared correlation ≥ 0.01) to a causal variant: if so, we classify the window as a true positive; otherwise, we classify it as a false positive. The true positive rate is calculated as the number of true positive 1Mb windows divided by the total number of simulated causal variants, and it is also known as the recall, or sensitivity, reflecting the power of a statistical test. We provide two measures of the false discovery rate: FDR1 is calculated as the number of false positive 1Mb windows divided by the number of 1Mb windows that contain genome-wide significant associations, and it is a measure of the proportion of independent discoveries that are false. FDR2 is calculated as the total number of false positive SNPs (squared correlation ≤ 0.01 with a causal variant) identified divided by the total number of identified genome-wide significant associations. We calculate the type-I error rate (TIER) as the number of SNPs with no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant) that pass the Bonferroni genome-wide significance testing threshold divided by the total number of SNPs that have no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant).

LOCO testing. We then repeat the analysis again with LDAK-KVIK using 2,174,071, 1,410,525 or 887,060 SNP markers for the first stage and 8,430,446 SNP markers for the second stage LOCO testing. This creates three simulation scenarios where the heritability and number of causal variants is fixed, but the effect size distributions vary. We note that because the 2,174,071, 1,410,525 or 887,060 SNP marker sets are LD reduced subsets of the larger 8,430,446 SNP markers, the simulated causal variants are in LD with markers used in step 1, but they may not be present within the

Figure S16. Simulation study of λ_{GC} value of mixed linear model leave-one-chromosome out association testing in UK Biobank imputed genotype data. We compare performance of leave-one-chromosome-out association testing (LOCO) when the first step LOCO predictors are created at different sets of markers, from gVAMP (blue), an elastic net model implemented in the LDAK software (red), and Regenie (green). Comparisons are made across six settings: "V1" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers and effects are drawn from a Gaussian, "V2" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects that have a power relationship to the minor allele frequency of -0.5, "V3" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects drawn from an elastic net distribution, "V4" where 40,000 causal variants are randomly selected from 887,060 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, "V5" where 40,000 causal variants are randomly selected from 2,174,071 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, and "V6" where 5,000 causal variant effects are randomly selected from 2,174,071 SNPs, with effects that contribute 25% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5. For each 1Mb window that contains an identified genome-wide significant association, we then ask if the identified variants are correlated (squared correlation ≥ 0.01) to a causal variant: if so, we classify the window as a true positive; otherwise, we classify it as a false positive. The true positive rate is calculated as the number of true positive 1Mb windows divided by the total number of simulated causal variants, and it is also known as the recall, or sensitivity, reflecting the power of a statistical test. We provide two measures of the false discovery rate: FDR1 is calculated as the number of false positive 1Mb windows divided by the number of 1Mb windows that contain genome-wide significant associations, and it is a measure of the proportion of independent discoveries that are false. FDR2 is calculated as the total number of false positive SNPs (squared correlation ≤ 0.01 with a causal variant) identified divided by the total number of identified genome-wide significant associations. We calculate the type-I error rate (TIER) as the number of SNPs with no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant) that pass the Bonferroni genome-wide significance testing threshold divided by the total number of SNPs that have no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant).

data. This is in contrast to the simulations presented in the LDAK-KVIK [10] and REGENIE [1] papers, where the causal variants are always within the subset of SNPs used in the first stage of the

Figure S17. Simulation study of out-of-sample prediction accuracy in UK Biobank imputed genotype data. We compare out-of-sample prediction accuracy (y-axis) for polygenic risk scores (PRS) created at different sets of markers from gVAMP and an elastic net model implemented in the LDAK software. Grey lines give the 1:1 line. Comparisons are made across three settings: "V4" where 40,000 causal variants are randomly selected from 887,060 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, "V5" where 40,000 causal variants are randomly selected from 2,174,071 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, and "V6" where 5,000 causal variant effects are randomly selected from 2,174,071 SNPs, with effects that contribute 25% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5.

analysis.

We begin by comparing the out-of-sample prediction accuracy obtained by each approach for the different marker sets, following the same procedures outlined in the Methods for the empirical UK Biobank analysis. Note that it is not possible to obtain SNP marker effects for the model used in REGENIE step 1 and so we approximate this with a ridge-regression model implemented in the LDAK software. For the out-of-sample prediction, we use a hold-out set of 15,000 individuals that are unrelated (SNP marker relatedness < 0.05) to the training individuals. We present these results in Figure S13.

We then compare the LOCO association testing results obtained by REGENIE, LDAK-KVIK, and gVAMP. We estimate the average chi-squared value at the causal variants and present these results in Figure S14. We give a simple, fair comparison of the true positive rate (TPR) and false discovery rate (FDR) across methods controlling for LD, where we calculate the number of independent associations (squared correlation ≤ 0.01) identified by each approach that pass a Bonferroni genome-wide significance testing threshold of 0.05/8430446 within 1Mb windows of the DNA. Specifically, for each 1Mb window that contains an identified genome-wide significant

Figure S18. Simulation study of power for mixed linear model leave-one-chromosome out association testing in UK Biobank imputed genotype data. We compare the average chi-squared value at the causal variants (y-axis) for leave-one-chromosome-out association testing (LOCO) when the first step LOCO predictors are created at different sets of markers, from gVAMP (blue), an elastic net model implemented in the LDAK software (red), and Regenie (green). Comparisons are made across three settings: "V4" where 40,000 causal variants are randomly selected from 887,060 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, "V5" where 40,000 causal variants are randomly selected from 2,174,071 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, and "V6" where 5,000 causal variant effects are randomly selected from 2,174,071 SNPs, with effects that contribute 25% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5.

association, we then ask if it is correlated (squared correlation ≥ 0.01) to a causal variant: if so, we classify it as a true positive; otherwise, we classify it as a false positive. The true positive rate is calculated as the number of true positive 1Mb windows divided by the total number of simulated causal variants, and it is also known as the recall, or sensitivity, reflecting the power of a statistical test. We provide two measures of the false discovery rate: FDR1 is calculated as the number of false positive 1Mb windows divided by the number of 1Mb windows that contain genome-wide significant associations, and it is a measure of the proportion of independent discoveries that are false. FDR2 is calculated as the total number of false positive SNPs (squared correlation ≤ 0.01 with a causal variant) identified divided by the total number of identified genome-wide significant associations. As genome-wide association studies aim to detect regions of the DNA associated with the phenotype, the definition of a false discovery as the detection of a variant at genome-wide significance when that variant has squared correlation ≤ 0.01 with a causal variant within 1Mb is a very conservative one. Additionally, we calculate the type-I error rate as the number of SNPs with no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant) that pass

Figure S19. Simulation study of true positive rate (TPR), false discovery rate (FDR) and type-I error for mixed linear model leave-one-chromosome out association testing in UK Biobank imputed genotype data. We compare performance of leave-one-chromosome-out association testing (LOCO) when the first step LOCO predictors are created at different sets of markers, from gVAMP (blue), an elastic net model implemented in the LDAK software (red), and Regenie (green). Comparisons are made across three settings: "V4" where 40,000 causal variants are randomly selected from 887,060 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, "V5" where 40,000 causal variants are randomly selected from 2,174,071 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, and "V6" where 5,000 causal variant effects are randomly selected from 2,174,071 SNPs, with effects that contribute 25% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5. For each 1Mb window that contains an identified genome-wide significant association, we then ask if the identified variants are correlated (squared correlation ≥ 0.01) to a causal variant: if so, we classify the window as a true positive; otherwise, we classify it as a false positive. The true positive rate is calculated as the number of true positive 1Mb windows divided by the total number of simulated causal variants, and it is also known as the recall, or sensitivity, reflecting the power of a statistical test. We provide two measures of the false discovery rate: FDR1 is calculated as the number of false positive 1Mb windows divided by the number of 1Mb windows that contain genome-wide significant associations, and it is a measure of the proportion of independent discoveries that are false. FDR2 is calculated as the total number of false positive SNPs (squared correlation ≤ 0.01 with a causal variant) identified divided by the total number of identified genome-wide significant associations. We calculate the type-I error rate (TIER) as the number of SNPs with no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant) that pass the Bonferroni genome-wide significance testing threshold divided by the total number of SNPs that have no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant).

the Bonferroni genome-wide significance testing threshold divided by the total number of SNPs that have no correlation to a causal variant (squared correlation ≤ 0.01 with a causal variant). We present these results in Figure S15. Finally, we also calculate the ratio of the median chi-square to the expected median chi-square values (λ_{GC}) and we present these results in Figure S16. We conduct five simulation replicates for each of the three simulation scenarios, as we find that this is

Figure S20. Comparison of MCMC Gibbs sampling algorithm GMRM to gVAMP in an imputed UK Biobank data simulation study. For a simulation setting where 40,000 causal variants are randomly selected from 8,430,446 SNP markers and effects are drawn from a Gaussian ("V1"), in *(a)*, we show results from standard leave-one-chromosome-out (LOCO) association testing where we select either all markers, 2,174,071 markers, or 887,060 markers for the first stage and then use all markers for the second stage LOCO testing. GMRM utilises 2,174,071 markers, whilst gVAMP can utilise the full range. We calculate the true positive rate (TPR) and the false discovery rate (FDR) and find that the FDR is well controlled at 5% or less for both gVAMP and GMRM. For *(b)*, we compare out-of-sample prediction accuracy for polygenic risk scores created at different sets of markers from gVAMP (8,430,446 and 2,174,071) and GMRM (2,174,071). *(c)* gives the run time in hours for the first stage analysis of gVAMP and GMRM, across different marker sets using 50 CPU from a single compute node. *(d)* gives a comparison of the proportion of phenotypic variation attributable to either 8,430,446 or a subset of 2,174,071 autosomal single nucleotide polymorphism (SNP) genetic markers (SNP heritability) estimated by GMRM and gVAMP. We consider three simulation scenarios: "V1" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers and effects are drawn from a Gaussian, "V4" where 40,000 causal variants are randomly selected from 887,060 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5, "V5" where 40,000 causal variants are randomly selected from 2,174,071 SNPs, with effects that contribute 50% to the phenotypic variance and have a power relationship to the minor allele frequency of -0.5. Points give the posterior means for GMRM and the convergence of gVAMP from five simulation replicates. Analysing 8,430,446 SNPs with gVAMP increases the heritability estimate over GMRM. This is consistent with an increase in phenotypic variance captured by the full imputed sequence data, as opposed to analyzing a selected subset of SNP markers, in which case gVAMP estimates are lower than those obtained from GMRM. Given the same data containing all the causal variants, the algorithms perform similarly irrespective of the underlying effect size distributions ("V4" and "V5").

more than sufficient to contrast methods, as the results were very consistent across replicates.

We then further conduct another three simulation scenarios: *V4* where the 40,000 causal variants are drawn from the 2,174,071 SNP subset and we simulate a α-power relationship between the causal variant effect size and the minor allele frequency (see Supplementary Note 3), with $\alpha = -0.5$; *V5* where the 40,000 causal variants are drawn from the 887,060 SNP subset and we simulate a α-power relationship between the causal variant effect size and the minor allele frequency (see Supplementary Note 3), with $\alpha = -0.5$; and *V6* where the 5,000 causal variants are drawn from the 2,174,071 SNP subset with SNP-heritability = 0.25 and we simulate a α-power relationship between the causal variant effect size and the minor allele frequency (see Supplementary Note 3),

Figure S21. Timing comparison of LDAK elastic net prediction model and gVAMP in (a) an imputed UK Biobank data simulation study and (b) across 13 UK Biobank traits. *(a)* For simulation setting "V3" where 40,000 causal variants are randomly selected from 8,430,446 SNP markers, with effects drawn from an elastic net distribution, we show the total run time of the algorithms when run on different numbers of imputed SNP markers and given different numbers of CPU resources from a single AMD EPYC 7763 (64 cores, 2.45 GHz base-frequency) compute machine. *(b)* A boxplot of the run time of the algorithms when run on 13 different UK Biobank traits and 2,174,071 SNPs, when given 60 CPU. Training data sample size is given in Table S1 for each trait.

with $\alpha = -0.5$. Here, our objective in *V4*, *V5*, and *V6* is to compare REGENIE, LDAK-KVIK and gVAMP in the scenario where the causal variants are present in the data used to create the predictors for the first step of LOCO as to date simulations for these methods have assumed this. This ensures that our findings are not just driven by only having SNPs correlated with the causal variants in step 1. We present these results, comparing prediction accuracy of LDAK to gVAMP in Figure S17, comparing the chi-squared values at the causal variants from REGENIE, LDAK, and gVAMP in Figure S18, and comparing the TPR and FDR of REGENIE, LDAK, and gVAMP in Figure S19.

Additionally, we compare to GMRM using 2,174,071 SNP markers (as this completes within reasonable compute time and resource use), running for 2,500 iterations with 500 iteration burn in. Our objective is to compare GMRM and gVAMP to empirically assess the Bayes optimality of gVAMP when applied to genomic data. We compare the LOCO association testing, the out-of-sample prediction accuracy, the run time, and the SNP heritability of the two methods under the scenarios "V1", "V4" and "V5". We present these results in Figure S20. Finally, we compare the run time of REGENIE, LDAK and gVAMP both in a controlled simulation setting "V3" and across the 13 UK Biobank traits. We present these results in Figure S21.

Imputed data simulation study results

For mixed linear model association testing (MLMA), association testing power (TPR) depends upon the sample size and the out-of-sample prediction accuracy of the predictors obtained from

the first step [46]. Polygenic risk score prediction accuracy depends upon h^2_{SNP}, the number of true underlying causal variants and the sample size [47], which are fixed in our simulation allowing a comparison of prediction accuracy obtained by different approaches across different genetic architectures. We find that prediction accuracy increases as marker number increases for all approaches (Figure S13). The prediction accuracy obtained by gVAMP is always slightly higher than that obtained by the LDAK model in every scenario and this difference becomes larger as more markers are used (Figure S13).

When these predictors are used for leave-one-chromosome-out MLMA testing, we find that the average chi-square values of the causal variants increases as marker number increases only for gVAMP (Figure S14). The values obtained by gVAMP are always higher than those obtained by LDAK or REGENIE and this difference becomes larger as more markers are used (Figure S14). We find that gVAMP has better LOCO association testing performance as compared to REGENIE or LDAK in terms of the TPR and FDR (Figure S15). In the simulation settings presented here, REGENIE does not control the FDR unless larger numbers of markers are used in step 1 (Figure S15). LDAK controls the FDR and type-I error well, comparative to gVAMP, but has lower TPR (Figure S15). gVAMP with larger numbers of SNPs used for step 1 gives higher TPR, whilst controlling the FDR below 5% (Figure S15).

Similar patterns are also observed in the scenarios where the causal variants are present in the data used to create the predictors for the first step of LOCO, where gVAMP shows improved prediction accuracy (Figure S17), higher average-chi square values of the causal variants (Figure S18), higher TPR and controlled FDR (Figure S19). Across all simulation settings, we find that the median chi-square value obtained from an MLMA-LOCO analysis with gVAMP is well controlled, showing the moderate elevation expected under polygenicity (Figure S16). This is in sharp contrast to REGENIE, which is elevated in all settings (Figure S16). The values obtained by LDAK rely on their proposal for adjusting the chi-square statistics and were generally lower than those obtained by MLMA-LOCO with gVAMP (Figure S16), even if we use the gVAMP LOCO predictors in place of the LDAK-KVIK ones and using LDAK-KVIK's proposed step 2 (Table S4).

Finally, we also show that gVAMP has comparative performance to a similar MCMC implementation of a Bayesian mixture of Gaussians variable selection regression model (Figure S20). Analysing 8,430,446 SNPs with gVAMP increases the SNP-heritability estimate over GMRM. This is consistent with an increase in phenotypic variance captured by the full imputed sequence data, as opposed to analyzing a selected subset of SNP markers, in which case gVAMP estimates are lower than those obtained from GMRM (Figure S20). Given the same data containing all the causal variants, the algorithms perform similarly irrespective of the scenario (Figure S20).

Supplementary Note 2

Onsager correction calculation

In order to ensure Gaussianity of residuals, gVAMP calculates the so-called Onsager correction based on (5). For such calculation, the derivative of the denoising function f_t defined in (3) is required. Let us denote the numerator and denominator of (4) with $\text{Num}(r_1)$ and $\text{Den}(r_1)$, respectively. Then,

$$
\frac{\partial \text{Num}(r_1)}{\partial r_1} = \lambda_t \cdot \sum_{l=1}^{L} \pi_{t,l} \cdot \frac{\sigma_{t,l}^2}{(\gamma_{1,t}^{-1} + \sigma_{t,l}^2)^{3/2}} \cdot \text{EXP}(\sigma_{t,l}^2) \cdot \left[1 - r_1^2 \cdot \frac{(\sigma_{t,*}^2 - \sigma_{t,l}^2)}{(\gamma_{1,t}^{-1} + \sigma_{t,l}^2)(\gamma_{1,t}^{-1} + \sigma_{t,*}^2)} \right],
$$

$$
\frac{\partial \text{Den}(r_1)}{\partial r_1} = -r_1 \cdot \left[\lambda_t \cdot \sum_{l=1}^{L} \frac{\pi_{t,l}}{(\gamma_{1,t}^{-1} + \sigma_{t,l}^2)} \cdot \frac{\sigma_{t,*}^2 - \sigma_{t,l}^2}{(\gamma_{1,t}^{-1} + \sigma_{t,l}^2)(\gamma_{1,t}^{-1} + \sigma_{t,*}^2)} \cdot \text{EXP}(\sigma_{t,l}^2) \right.
$$
$$
\left. + (1 - \lambda_t) \cdot \frac{\gamma_{1,t}^{3/2} \cdot \sigma_{t,*}^2}{(\gamma_{1,t}^{-1} + \sigma_{t,*}^2)} \cdot \text{EXP}(0) \right].
$$

Thus, the Onsager correction reads

$$
\frac{\partial}{\partial r_1} \left(\frac{\text{Num}(r_1)}{\text{Den}(r_1)} \right) = \frac{\frac{\partial}{\partial r_1} \text{Num}(r_1)}{\text{Den}(r_1)} - \frac{\text{Num}(r_1) \cdot \frac{\partial}{\partial r_1} \text{Den}(r_1)}{(\text{Den}(r_1))^2}.
$$

Conjugate gradient algorithm for solving linear systems

Algorithm 2 Conjugate gradient method for solving a symmetric linear system $\boldsymbol{Ax} = \boldsymbol{b}$.

1: **Input:** Initial estimate of the solution $\boldsymbol{x}_0$, initial residual $\boldsymbol{r}_0 = \boldsymbol{b}$, initial search direction $\boldsymbol{p}_0 = \boldsymbol{r}_0$, linear system matrix $\boldsymbol{A}$, right-hand side vector $\boldsymbol{b}$, stopping error threshold $\varepsilon > 0$.
2: **for** $n = 1, 2, 3, \ldots$ **do**
3: $\quad \alpha_n = \frac{\boldsymbol{r}_{n-1}^T \boldsymbol{r}_{n-1}}{\boldsymbol{p}_{n-1}^T \boldsymbol{A} \boldsymbol{p}_{n-1}}$
4: $\quad \boldsymbol{x}_n = \boldsymbol{x}_{n-1} + \alpha_n \boldsymbol{p}_{n-1}$
5: $\quad \boldsymbol{r}_n = \boldsymbol{r}_{n-1} - \alpha_n \boldsymbol{A} \boldsymbol{p}_{n-1}$
6: $\quad \beta_n = \frac{\boldsymbol{r}_n^T \boldsymbol{r}_n}{\boldsymbol{r}_{n-1}^T \boldsymbol{r}_{n-1}}$
7: $\quad \boldsymbol{p}_n = \boldsymbol{r}_n + \beta_n \boldsymbol{p}_{n-1}$
8: $\quad$ **If** $n \geq 1$ and $||\boldsymbol{x}_n - \boldsymbol{x}_{n-1}||_2 / ||\boldsymbol{x}_n||_2 < \varepsilon$, then **break**
9: **end for**
10: **return** $\boldsymbol{x}_n$

Supplementary Note 3

Simulation of effects under α-power law

In expectation, we randomly select $\lambda \cdot P$ causal SNP positions. For each SNP position j, upon randomly choosing a mixture group l it belongs to, according to the true effect probability distribution, we draw its effect size from $\mathcal{N}(0, \sigma_l^2 \cdot (2 \cdot \mathrm{MAF}_j \cdot [1 - \mathrm{MAF}_j])^\alpha)$. Given a desired heritability h^2 of the simulated trait, we normalize the estimates so that the sum of squares of all simulated effects is equal to h^2.

Estimation of α

Let us denote a set of causal SNPs with $\mathcal{C}$. Then, under the assumption that the effect of the causal SNP with index j satisfies the following relationship with their minor allele frequency (MAF_j):

$$\beta_j \sim c_l \cdot \mathcal{N}(0, (2 \cdot \mathrm{MAF}_j \cdot [1 - \mathrm{MAF}_j])^\alpha), \quad \forall j \in \mathcal{C},$$

where c_l is a constant depending on the mixture group l and not on MAF_j, one can rewrite this relationship as

$$\beta_j = c_l \cdot (2 \cdot \mathrm{MAF}_j \cdot [1 - \mathrm{MAF}_j])^{\alpha/2} \cdot Z_j, \quad \forall j \in \mathcal{C}$$

where $Z_j \sim \mathcal{N}(0, 1)$ and mutually independent. This implies that

$$\log\left(\beta_j^2\right) = \text{constant} + \log\left(Z_j^2\right) + \alpha \cdot \log(2 \cdot \mathrm{MAF}_j \cdot [1 - \mathrm{MAF}_j]).$$

Hence, by regressing $\log\left(\beta_j^2\right)$ onto $\log(2 \cdot \mathrm{MAF}_j \cdot [1 - \mathrm{MAF}_j])$ with intercept term, for a subset of SNPs that pass the PIP threshold of 0.95, one can read out an estimate of α as the slope parameter in the regression.